PERTURBED LATTICE CROSSES AND HEISENBERG UNIQUENESS PAIRS

DANYLO RADCHENKO AND JOÃO P.G. RAMOS

ABSTRACT. This work focuses on two questions raised by H. Hedenmalm and A. Montes-Rodríguez [\[27\]](#page-35-0) on Heisenberg Uniqueness Pairs for perturbed lattice crosses.

The first of them deals with a *complete* characterization of $\beta > 0$ for which, for a fixed $\theta \in \mathbb{R}$, the translated lattice cross $\Lambda_{\beta}^{\theta} = ((\mathbb{Z} + {\theta}) \times {\theta}) \cup ({0} \times {\beta} \mathbb{Z})$ satisfies that $(\Gamma, \Lambda_{\beta}^{\theta})$ is a Heisenberg Uniqueness Pair, where Γ is the hyperbola in \mathbb{R}^2 with axes as asymptotes. As a matter of fact, we show that $(\Gamma, \Lambda^{\theta}_{\beta})$ is an H.U.P. *if and only if* $\beta \leq 1$, confirming a prediction made in [\[27\]](#page-35-0).

Furthermore, under modified decay conditions on the measures under consideration, we are able to prove *sharp* results for when a perturbed lattice cross $\Lambda_{A,B}$ is such that $(\Gamma, \Lambda_{A,B})$ is an H.U.P. In particular, under such decay conditions, this solves another question posed by H. Hedenmalm and A. Montes-Rodríguez.

Our techniques run through the analysis of the action of the operator which maps the Fourier transform of an L^1 function ψ to the Fourier transform of $t^{-2}\psi(1/t)$. In other words, we analyze the operator taking the restriction to the x-axis of a solution u to the Klein-Gordon equation to its restriction to the y-axis. This operator turns out to be related to the action of the Four-dimensional Fourier transform on radial functions, which enables us to use the philosophical framework and techniques of discrete uncertainty principles for the Fourier transform.

1. INTRODUCTION

1.1. **Historical Background.** Let $f : \mathbb{R} \to \mathbb{C}$ be a measurable, square-integrable function. A crucial question in Fourier analysis, with consequences in fields ranging from physics to signal processing is: how well can one recover the information of such a function f and its Fourier transform \hat{f} , given one only has partial access to such information?

As a first partial answer to such a question, one may invoke the classical *Heisenberg uncertainty principle*. It guarantees that any function $f \in L^2$ cannot be too concentrated in space, unless it is the case that its Fourier transform \hat{f} is 'spread out' in frequency: in particular, the mass of any such f cannot be 'overwhelmingly concentrated' in a space ball of small radius around any $x_0 \in \mathbb{R}$. while at the same time the mass of \hat{f} being concentrated in another ball of small radius around $\xi_0 \in \mathbb{R}$.

As striking and elegant as such a principle may seem, it represents only the beginning of a rich literature on the subject of uncertainty principles. Indeed, several results in the literature have been of instrumental nature in further understanding the relationship between space-frequency concentration and recovery of functions.

We highlight, in particular: (i) the Shannon-Whittaker formula [\[40,](#page-36-0) [48\]](#page-36-1), which states that, for a function $f \in L^2$ whose Fourier transform is compactly supported on a compact interval, we may recover it by its values from a certain rescaling of the set of integers. This implies that, if a Fourier transform of a compactly supported function vanishes on certain rescaling of the integers, then it must vanish identically; (ii) Hardy's Uncertainty Principle [\[25\]](#page-35-1), which shows that, if a function and its Fourier transform are pointwise bounded by a multiple of a certain low-variance Gaussian, then it must vanish identically; (iii) and finally, the Amrein-Berthier-Benedicks theorem $[6, 1, 5]$ $[6, 1, 5]$ $[6, 1, 5]$ $[6, 1, 5]$ $[6, 1, 5]$, which states that, for any two measurable sets $A, B \subset \mathbb{R}$ of *finite* measure, a function cannot have support in \tilde{A} with its Fourier transform supported in \tilde{B} , unless it is *identically zero*.

In this manuscript, we shall be interested in uncertainty principles of a similar flavour, but which are related to a certain given *partial differential equation*. In fact, all of the results mentioned above may be interpreted as suitable uniqueness results for the *harmonic oscillator*: if we define, for a fixed $f \in L^2$, the function $\Phi_f(x,t)$ as being a solution to

$$
i \partial_t \Phi_f = (-\Delta + \pi^2 |x|^2) \Phi_f, \text{ for } (x, t) \in \mathbb{R} \times \mathbb{R},
$$

with $\Phi_f(x,0) = f(x)$ for $x \in \mathbb{R}$, then one has that $\Phi_f(x,1/8)$ is equal to (a constant multiple of) the Fourier transform of f . Hence, the uncertainty-flavoured results mentioned above translate naturally as results about *uniqueness of solutions* to the time-dependent harmonic oscillator above.

In line with these results, we highlight two main lines of related work. The first of them is dedicated to extending the philosophical view of uncertainty principles as uniqueness results for certain partial differential equations as a more concrete device. Here, we mention, for instance, the celebrated works by L. Escauriaza, C. Kenig, C. Ponce and L. Vega [\[12,](#page-35-2) [13,](#page-35-3) [14,](#page-35-4) [10,](#page-34-3) [15\]](#page-35-5), where the authors prove a sharp version of Hardy's uncertainty principle for general Schrödinger equations with a potential. In that same line of work, we also highlight the following articles [\[31,](#page-35-6) [23,](#page-35-7) [33\]](#page-35-8) and the references therein.

The second line of work mentioned above is the main focus of this manuscript. In order to describe it, let first μ denote a Borel measure on \mathbb{R}^2 . Suppose, moreover, that $\text{supp}(\mu) \subset \Gamma_0$, where Γ_0 is some smooth curve in \mathbb{R}^2 . A set $\Lambda \subseteq \mathbb{R}^2$ is said to form a *Heisenberg Uniqueness Pair* (or H.U.P., for shortness) with Γ_0 if, whenever μ above satisfies $\hat{\mu}|_{\Lambda} = 0$, then we must have $\mu \equiv 0$.

This definition as such may seem unrelated to the topic at hand of uniqueness results for partial differential equations, but, as a matter of fact, in most cases of curves Γ_0 explored in the literature so far this relationship is not only evident but also crucial. Indeed, we highlight here three main cases:

I. The parabola $\Gamma = \{(x, y): y = x^2\}$. In this case, any measure μ supported on Γ yields through Fourier transform a solution to the free Schrödinger equation $\partial_t \hat{\mu} + \frac{i}{\pi}$ $\frac{i}{\pi}\partial_x^2\hat{\mu} = 0;$

II. The circle $\Gamma = \{(x, y): x^2 + y^2 = 1\}$. Here, one readily obtains that $\hat{\mu}$ satisfies the *Helmholtz* equation $\Delta \hat{\mu} = \pi^2 \hat{\mu};$

III. The hyperbola $\Gamma = \{(x, y): xy = 1\}$. Finally, here one has that, if $\text{supp}(\mu) \subset \Gamma$, then $\hat{\mu}$ satisfies the Klein-Gordon equation $\partial_x \partial_y \hat{\mu} + \pi^2 \hat{\mu} = 0$.

Note that, in light of these considerations, the first line of work described above [\[12,](#page-35-2) [14,](#page-35-4) [13,](#page-35-3) [10,](#page-34-3) [15\]](#page-35-5) is directly related to the concept of Heisenberg Uniqueness Pairs for the Schrödinger equation. In this work, on the other hand, we shall focus on the third case, in which Heisenberg Uniqueness Pairs are directly related to uniqueness sets for the Klein-Gordon equation. We refer the reader to

 $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ $[14, 24, 16, 30, 34, 41, 42, 46, 17]$ for references in the directions of the first two other instances described above.

In spite of the fact that the definition of Heisenberg Uniqueness Pairs becomes natural in the context introduced above, it seems, to the best of our knowledge, that it has first been explicitly introduced only recently in the seminal work of H. Hedenmalm and A. Montes-Rodriguez [\[27\]](#page-35-0). There, besides introducing the concept and discussing some basic examples similar to the ones above, they completely resolve the question of determining all $\alpha, \beta > 0$ such that the pair $(\Gamma, \Lambda_{\alpha,\beta})$ is a Heisenberg Uniqueness Pair, where Γ denotes the hyperbola $\Gamma = \{(x, y) \in \mathbb{R}^2 : xy = 1\}$, and $\Lambda_{\alpha,\beta} = ((\alpha \mathbb{Z}) \times \{0\}) \cup (\{0\} \times (\beta \mathbb{Z}))$ denotes the *lattice cross* with dilation parameters α,β .

More specifically, they showed that $(\Gamma, \Lambda_{\alpha,\beta})$ is an H.U.P. if and only if $\alpha\beta \leq 1$. Their result, in addition to possessing a beautiful proof using elements from dynamical systems in novel ways, drew new connections between fields in analysis and raised several important questions in the direction of Fourier uncertainty principles. Effectively, several subsequent works further exploited the connections between such problems and the Klein-Gordon equation such as [\[28,](#page-35-13) [7,](#page-34-4) [29,](#page-35-14) [3,](#page-34-5) [2,](#page-34-6) [18\]](#page-35-15), and, as recently highlighted [\[38,](#page-36-6) [3,](#page-34-5) [2\]](#page-34-6), Heisenberg Uniqueness Pairs for the hyperbola turn out to have a connection to recent *Fourier interpolation results*, such as the interpolation formula due to the first author and M. Viazovska [\[37\]](#page-36-7), which possesses a particularly important connection to the solution of the sphere packing problem in dimensions 8 and 24 [\[8,](#page-34-7) [45,](#page-36-8) [9\]](#page-34-8).

1.2. Main results. In this manuscript, we shall focus on a question initially raised by H. Hedenmalm and A. Montes-Rodríguez:

Question 1 (Open problem (a) in [\[27\]](#page-35-0)). If one defines the translated lattice crosses

$$
\Lambda_{\beta}^{\theta} = ((\mathbb{Z} + \{\theta\}) \times \{0\}) \cup (\{0\} \times \beta \mathbb{Z}), \tag{1.1}
$$

for which $\beta > 0$ is $(\Gamma, \Lambda_{\beta}^{\theta})$ a Heisenberg Uniqueness Pair?

Moreover, if one considers suitable perturbations $\tilde{\Lambda} \subset (\mathbb{R} \times \{0\}) \cup (\{0\} \times \mathbb{R})$ of lattice crosses, when is $(\Gamma, \tilde{\Lambda})$ a H.U.P.?

The authors conjecture in [\[27\]](#page-35-0) that the methods developed there could be adapted to prove results for the first of those questions. On the other hand, the perturbative regime suggested by them seemed to require new ideas outside of the scope of their techniques, and it was left open. Regarding the latter question, we note that the only result that we are aware of in this direction was by the second author and M. Stoller [\[38\]](#page-36-6), which showed that, at least for certain smooth classes of measures, it is indeed the case that one is able to perturb $\Lambda_{\alpha,\beta}$ slightly and still obtain positive results in the case when $\alpha \cdot \beta = 1$.

Regarding the first question on lattice crosses translated by a fixed quantity $\theta \in \mathbb{R}$, D. Giri and R. Rawat [\[20\]](#page-35-16) were the first ones to consider it explicitly after the original comment in [\[28\]](#page-35-13). In their manuscript, it was first claimed that $(\Gamma, \Lambda^{\theta}_{\beta})$ is a Heisenberg uniqueness pair if and only if $\beta \leq p$.

Although the examples constructed in [\[20\]](#page-35-16) show that $\beta \leq p$ is a necessary condition, the "if" part of this result turned out to be false, as noted by the authors themselves in [\[21\]](#page-35-17), due to a mistake in a computation undertaken on [\[20,](#page-35-16) page 8], while defining their main operators in analogy to the original work of Hedenmalm and Montes-Rodríguez.

The main results in this manuscript are inspired by and provide answers to Question [1.](#page-2-0) Indeed, the first result we highlight here deals with the context of translated lattice crosses. Effectively, we are able to show that, independently of how $\theta \in \mathbb{R}$ is chosen, $(\Gamma, \Lambda^{\theta}_{\beta})$ is a Heisenberg Uniqueness Pair (H.U.P.) if and only if $\beta \leq 1$, settling hence the first part of Question [1.](#page-2-0)

Theorem 1. Let $\Gamma = \{(x, y) \in \mathbb{R}^2 : xy = 1\}$ denote the hyperbola, and consider the translated lattice crosses Λ_{β}^{θ} as defined in [\(1.1\)](#page-2-1), for some arbitrary $\theta \in \mathbb{R}$. Then $(\Gamma, \Lambda_{\beta}^{\theta})$ is a Heisenberg Uniqueness Pair if and only if $\beta \leq 1$.

Before describing the proof of Theorem [1,](#page-3-0) we remark on its surprisingly challenging nature by comparing our result with the ones obtained by using the techniques available in the literature of H.U.P.'s for the hyperbola so far.

Indeed, in spite of the fact that the original prediction in [\[27\]](#page-35-0) is, after all, correct, the techniques needed in order to fully accomplish it differ significantly from the original ones. This is especially highlighted by the previously available results in this direction in the literature: in order to prove the fact that $\beta \leq 1$ implies that $(\Gamma, \Lambda^{\theta}_{\beta})$ is indeed a H.U.P., one needs to resort to the subsequent work [\[28\]](#page-35-13), which in spite of being close in spirit to the results in [\[27\]](#page-35-0), was only developed several years later. For the proof that $\beta \leq 1$ is also *necessary*, the differences are even more drastic: as a matter of fact, the counterexamples in [\[27\]](#page-35-0) are functions of the form

$$
\psi(t) = P_{z_1}(t) - P_{z_2}(t),\tag{1.2}
$$

where $P_z(t) = \frac{y}{(x-t)^2+y^2}$ is the Poisson kernel at $z = x + iy$, and $z_1, z_2 \in \mathbb{C}_+$. In [\[20\]](#page-35-16), the authors, while employing the same Poisson extension techniques, obtain that $\beta \leq p$ is a necessary condition, whenever $\theta = \frac{1}{n}$ $\frac{1}{p}, p \in \mathbb{N}$. Since this lower bound on β blows up as $p \to \infty$, the gap between the necessary and sufficient conditions converges to infinity as the parameter $\frac{1}{p} \to 0$, which is diametrically opposite to what happens when $\theta = 0$.

In order to circumvent these issues, we need two main ingredients for the proof of Theorem [1:](#page-3-0) the first is a corrected proof of the "if" part of the result from [\[20\]](#page-35-16), which, as mentioned above, can be obtained by employing the techniques from [\[28\]](#page-35-13), as noted in the corrigendum [\[21\]](#page-35-17).

The second main ingredient, and the main novel feature of this work, is a new set of counterexamples for the "only if" part, going beyond the ones given by [\(1.2\)](#page-3-1). These new counterexamples are inspired by the correspondence between Heisenberg Uniqueness Pairs and Fourier uncertainty principles highlighted in [\[3\]](#page-34-5), together with recent constructions of A. Kulikov, F. Nazarov and M. Sodin [\[32\]](#page-35-18) of Fourier non-uniqueness sets in dimension 1.

In order to lay the foundation for an iterative process, as done in [\[32\]](#page-35-18), we need to consider a different setup, arising from the operator T which takes as input a function f and returns the restriction of the solution u to the Klein-Gordon equation $\partial_x \partial_y u + \pi^2 u = 0$ with $u|_{\{y=0\}} = f$ to the y-axis. An equivalent way to define this operator is by letting it be the operator which takes as input the Fourier transform of a function ψ , and returns as an output the Fourier transform of the function $t^{-2}\psi(1/t)$. As we shall see, this operator is, as in [\[3\]](#page-34-5), related to certain four-dimensional Fourier transforms of radial functions.

The crucial difference here is that its definition allows us to bypass radiality, making it possible for one to consider asymmetric sets in the counterexample construction - a feature not achiavable

through a purely Fourier-based approach. Moreover, the properties of the operator T (see Proposition [2.4\)](#page-10-0) show that the iteration process has to be done in two separate parts, according to the positive and negative parts of the zero set under consideration. This allows us to maintain the crucial fact that the operator T takes functions supported on a half-line to functions supported on the opposite half-line, allowing one to deduce results also in the case of one-sided H.U.P.'s – as highlighted in Section [5.2.](#page-29-0)

Our next result deals, on the other hand, with the topic of H.U.P.'s for perturbed lattice crosses, as stated in the second part of Question [1.](#page-2-0) Even in the case dealt with in [\[38\]](#page-36-6), the perturbations considered are taken to be origin-symmetric and considerably close to the original lattice cross points, with the additional constraint that the functions considered there are taken to be realvalued, which excludes a large variety of cases. In that context, we give below the first instance of sharp conditions in order for such a perturbed lattice cross to form a H.U.P. with the hyperbola, under different decay assumptions when compared to the ones in, for instance, [\[27,](#page-35-0) [28\]](#page-35-13).

In order to state such a result, we let first

$$
\mathcal{H}_{\ell}(\mathbb{R}) = \{g : \mathbb{R} \to \mathbb{C} : g \in L^{2}((1+|y|^{2\ell}) dy), \,\widehat{g} \in C_{0}(\mathbb{R})\},\,
$$

where $\ell \in \mathbb{Z}_{\geq 0}$. We remark that these spaces are somewhat reminiscent of the ones defined in [\[23\]](#page-35-7) in the context of H.U.P.'s for the parabola.

Definition 1.1. Let μ be a measure on \mathbb{R}^2 supported on Γ which can be represented as $d\mu(t, 1/t) =$ $\psi(t) dt$, for some $\psi \in \mathcal{H}_{\ell}(\mathbb{R})$, and let Λ be a subset of $(\mathbb{R} \times \{0\}) \cup (\{0\} \times \mathbb{R})$. We say that the pair (Γ, Λ) is an \mathcal{H}_{ℓ} -Heisenberg Uniqueness Pair (or \mathcal{H}_{ℓ} -H.U.P.) if for any μ as above with $\hat{\mu}|_{\Lambda} = 0$, we have $\mu \equiv 0$.

Note that, if $\ell \geq 1$, then $\mathcal{H}_{\ell} \subset L^1(\mathbb{R})$, and hence the concept of \mathcal{H}_{ℓ} -H.U.P.'s is well-defined in that case, since $\hat{\mu}(x, 0) = \hat{\psi}(x)$, and $\hat{\mu}(0, y) = \hat{\varphi}(y)$, where $\varphi(t) = \frac{1}{t^2} \psi\left(\frac{1}{t}\right)$ $(\frac{1}{t}) \in L^1$. For the $\ell = 0$ case, note that $\psi \in \mathcal{H}_0$ is equivalent to $\varphi \in L^2(t^2 dt)$, which means that the Fourier transform of $t \cdot \varphi(t)$ is an L^2 L^2 function. By standard distribution theory arguments (see Section 2 below), this implies that the Fourier transform of φ as a distribution may be identified with a *continuous* function of polynomial growth, and hence $\hat{\mu}|_{(\mathbb{R}\times{0})\cup({0}\times\mathbb{R})}$ is defined pointwise and continuous, which shows that the concept of \mathcal{H}_0 -H.U.P.'s is also well-defined.

With that in mind, we shall use the following notation for the *generalized lattice cross*

$$
\Lambda_{\mathbf{A},\mathbf{B}} = (A \times \{0\}) \cup (\{0\} \times B),
$$

where $A = \{a_n\}_{n \in \mathbb{Z}}$, $B = \{b_n\}_{n \in \mathbb{Z}}$ are two separated sequences of points on \mathbb{R} , in the sense that

$$
\min(|a_{n+1}-a_n|,|b_{n+1}-b_n|) > \delta, \qquad \forall n \in \mathbb{Z},
$$

for some $\delta > 0$. We will sometimes also call such sets uniformly discrete.

Theorem 2. Let $\Gamma = \{(x, y) \in \mathbb{R}^2 \colon xy = 1\}$ denote the hyperbola. The following assertions hold:

(I) Suppose that

$$
\sup_{n} |a_{n+1} - a_n| = \alpha, \ \sup_{n} |b_{n+1} - b_n| = \beta.
$$

Then the pair $(\Gamma, \Lambda_{A,B})$ is an \mathcal{H}_0 -Heisenberg Uniqueness Pair if $\alpha\beta < 1$. Furthermore, $(\Gamma, \Lambda_{\mathbf{A}, \mathbf{B}})$ is an \mathcal{H}_2 -H.U.P. if $\alpha \beta = 1$.

(II) Suppose that

$$
\inf_{n} |a_{n+1} - a_n| = \alpha, \ \inf_{n} |b_{n+1} - b_n| = \beta,
$$

with $\alpha\beta > 0$. Then $(\Gamma, \Lambda_{A,B})$ is not an \mathcal{H}_1 -H.U.P. (and hence also neither a \mathcal{H}_0 -H.U.P., nor a regular H.U.P.) if $\alpha\beta > 1$.

In order to prove the positive part of this result, we use a *Poincaré-Wirtinger inequality* argument: by using such estimates, one can essentially control the L^2 -norm of $x \cdot \psi(x)$, where ψ is the density of a measure μ as above with $\hat{\mu}|_{\Lambda_{A,B}} = 0$, strictly from below by the L^2 norm of ψ itself. On the other hand, the same argument can be used for $\varphi(t) = t^{-2}\psi(1/t)$, which implies the reverse *strict* inequality, yielding a contradiction.

The sketch of proof above is inspired by the positive Fourier Uniqueness results from [\[32\]](#page-35-18), but again with differences induced by the context of the result. It is striking to note that, by considering the definition of a \mathcal{H}_0 -H.U.P., then the argument carried out here allows for a considerable simplification of the subcritical results from [\[27\]](#page-35-0). It is further important to notice that the class of functions \mathcal{H}_0 is inherently *different* from the ones considered in [\[27,](#page-35-0) [28,](#page-35-13) [7\]](#page-34-4), which implies that, in the case when $A = \alpha \mathbb{Z}, B = \beta \mathbb{Z}, \alpha \beta \leq 1$, our results actually *extend* the ones in those references to different classes. We refer the reader to Sections [4.2](#page-18-0) and [5.3](#page-31-0) below for more details.

Additionally, we remark that Theorem [2](#page-4-0) may be interpreted alternatively as a first step towards an answer to *another* question in [\[27\]](#page-35-0): in [\[27,](#page-35-0) Open problem (d)], the authors ask whether it can be proved that, for a general distributional solution u to $\partial_x \partial_y u + \pi^2 u = 0$, such that u is a continuous function vanishing at infinity, then $u|_{\Lambda_{\alpha,\beta}} = 0 \Rightarrow u \equiv 0$ if $\alpha\beta \leq 1$. In that regard, Theorem [2](#page-4-0) does not fully remove the Fourier analysis aspect of the matter, but it accomplishes the goal of having a condition merely dependent on the function u , which is that its restriction to the real axis belongs to $L^2(\mathbb{R})$, and that it vanishes at infinity. We believe that a similar approach could lead to a solution to this problem, which we wish to investigate in future work.

For Part (II) of Theorem [2,](#page-4-0) we will use essentially the same techniques used in order to construct the counterexamples in the "only if" part of Theorem [1.](#page-3-0) For that reason, and for the inherent interest in such a construction in its own right, we state an unified result concerning the counterexamples constructed.

Theorem 3. Let $A = \{a_n\}_{n \in \mathbb{Z}}$ and $B = \{b_n\}_{n \in \mathbb{Z}}$ be two sequences of real numbers such that

$$
\min\left(\liminf_{n\to\pm\infty}|a_{n+1}-a_n|,\liminf_{n\to\pm\infty}|b_{n+1}-b_n|\right)>1.
$$

Then there exists a function $\psi \in L^1(\mathbb{R})$ such that $\psi \not\equiv 0$ and

$$
\int_{\mathbb{R}} e^{-\pi i \tau a_n} \psi(\tau) d\tau = \int_{\mathbb{R}} e^{-\pi i b_n/\tau} \psi(\tau) d\tau = 0, \quad \forall n \in \mathbb{Z}.
$$
\n(1.3)

We further note that the results in Theorem [2](#page-4-0) can also be strengthened so that the density condition in its positive part are only taken to hold in a limit sense. The proof of such a result, as expected, runs through a generalization of the results in [\[32\]](#page-35-18) to higher dimensions. We defer the discussion of this to Section [5.1.](#page-20-0)

Lastly, we briefly point out that our methods allow for a *complete* solution of the original problems studied by H. Hedenmalm and A. Montes-Rodríguez in [\[27\]](#page-35-0), under the \mathcal{H}_0 -condition, even in the critical $\alpha\beta = 1$ case. It also yields a solution of the \mathcal{H}_0 -modification of Theorem [1.](#page-3-0) We briefly highlight these facts in the following result (see Section [5.3](#page-31-0) for further details):

Theorem 4. Let Γ be the hyperbola as defined in Theorem [1,](#page-3-0) and $\theta \in \mathbb{R}$. If $\Lambda^{\theta}_{\alpha,\beta}$ = $((\alpha \mathbb{Z} + {\theta}) \times {\theta}) \cup ({0} \times {\beta} \mathbb{Z}), \text{ then } (\Gamma, \Lambda^{\theta}_{\alpha, \beta}) \text{ is a } \mathcal{H}_0\text{-H.U.P. if, and only if, } \alpha \beta \leq 1.$

The proof of that result employs similar techniques as that of Theorems [1](#page-3-0) and [2](#page-4-0) in order to deal with the subcritical and supercritical cases. In the critical case, however, a more subtle analysis is needed, and the equality case in the Poincaré–Wirtinger inequality yields that any function yielding a candidate to fail the H.U.P. property satisfies two different kinds of periodicity conditions. One of them is a usual 2-periodicity condition, and the other is that the function composed with $1/t$ is still 2-periodic. By using classical results on the geometry of discrete groups, we are able to conclude that the only functions satisfying those conditions are constants, and a direct analysis shows that the only alternative is that the functions we deal with are identically zero.

The manuscript is organized as follows: we first set, in Section [2,](#page-6-0) the groundwork for our arguments, defining the operator T and stating its most important properties. We then prove in Section [3](#page-11-0) Theorem [3,](#page-5-0) since it will be of universal use throughout the manuscript. In Section [4,](#page-17-0) we finally prove Theorems [1](#page-3-0) and Theorem [2.](#page-4-0) Finally, in Section [5,](#page-20-1) we discuss applications of our main results and techniques: in particular, we comment on an asymptotic generalization of Theorem [2](#page-4-0) in Section [5.1;](#page-20-0) we prove a one-sided lattice cross result, as studied in [\[28\]](#page-35-13), in Section [5.2;](#page-29-0) and we comment further on the endpoint case of Theorem [2](#page-4-0) and its relationship to summation formulas in Section [5.3,](#page-31-0) providing there also the proof of Theorem [4.](#page-6-1)

2. Preliminaries

2.1. Notation and definitions. We shall use the following notation throughout the manuscript: whenever not further specified, $a \leq b$ means that $a \leq C \cdot b$, where $C > 0$ is a finite numerical constant whose exact value is not important for the respective context.

We define the Fourier transform of $f \in L^1(\mathbb{R}^d)$ in dimension $d \in \mathbb{N}$ as

$$
\mathcal{F}_d(f)(\xi) = \int_{\mathbb{R}^d} f(x)e^{-2\pi ix\xi} dx.
$$
\n(2.1)

In some cases, however, it will be convenient to consider a different normalization:

$$
\widehat{f}(\xi) = \frac{1}{\sqrt{2}^d} \int_{\mathbb{R}^d} f(x) e^{-\pi i x \cdot \xi} dx.
$$
\n(2.2)

This distinction will spare us from some unnecessary calculations with changes of variables. The inverse of the operator defined in [\(2.2\)](#page-6-2) will be denoted by

$$
\breve{f}(\xi) = \frac{1}{\sqrt{2}^d} \int_{\mathbb{R}^d} f(x) e^{\pi i x \xi} dx.
$$

Finally, for a given class of measurable functions C, we will usually write that (Γ, Λ) is a C-Heisenberg Uniqueness Pair (or C−H.U.P.) if, given any μ supported on Γ absolutely continuous with density

 $\psi \in \mathcal{C}$ and such that $\hat{\mu}|_{\Lambda} = 0$, we have $\mu \equiv 0$. Here, note that the normalization employed is crucial in order to state an accurate version of Theorems [1](#page-3-0) and [2.](#page-4-0)

2.2. Function spaces and embeddings. We shall use the following notation for *homogeneous* Sobolev spaces:

$$
H^{s}(\mathbb{R}) = \left\{ f \in \mathcal{S}'(\mathbb{R}) : \ \hat{f} \text{ is a function in } L^{1}_{loc}(\mathbb{R} \setminus \{0\}) \text{ and } \int_{\mathbb{R}} |\hat{f}(\xi)|^{2} |\xi|^{2s} d\xi < +\infty \right\},\tag{2.3}
$$

where $\mathcal{S}(\mathbb{R})$ denotes the Schwartz class, and $\mathcal{S}'(\mathbb{R})$ denotes the space of tempered distributions. In the case of a domain $\Omega \subset \mathbb{R}^d$, we will use the following notation for the *inhomogeneous* Sobolev space:

 $W^{1,p}(\Omega) = \{f : \Omega \to \mathbb{C} \colon \nabla f \text{ exists in the weak sense, and } ||f||_p + ||\nabla f||_p < +\infty \}.$

With regard to Sobolev spaces on intervals of the real line, we highlight the classical $Poincaré$ -Wirtinger inequality, which will be crucial in the proof of Theorem [2,](#page-4-0) as well as in Section [5.1.](#page-20-0) We refer the reader to [\[26,](#page-35-19) Section 7.7] for a proof of this result.

Theorem (Poincaré-Wirtinger). Let $(a, b) = I \subset \mathbb{R}$ be an open interval. Suppose $f \in W^{1,2}(I)$ is such that $f(a) = f(b) = 0$. Then we have that

$$
\int_{I} |f(t)|^2 dt \le \left(\frac{|I|}{\pi}\right)^2 \int_{I} |f'(t)|^2 dt,
$$
\n(2.4)

with equality if, and only if, $f(x) = c \sin \left(\frac{\pi(x-a)}{|I|} \right)$ $|I|$ for some $c \in \mathbb{C}$.

We also note the following embedding estimate, which will be particularly useful in Section [5.1:](#page-20-0)

Proposition 2.1. There exists $C > 0$ such that, for any $\theta \in L^2(\mathbb{R}) \cap H^1(\mathbb{R})$ with $\theta(0) = 0$, we have

$$
\|\theta\|_{L^2(dt/|t|)} \leq C \|\theta\|_{L^2}^{1/2} \|\theta'\|_{L^2}^{1/2}.
$$

Proof. We begin by noting that $\theta \in H^1(\mathbb{R})$ implies readily that $\theta \in AC(\mathbb{R})$. Moreover, we have the estimate

$$
|\theta(s)| \leq \int_0^s |\theta'(t)| dt \leq s^{1/2} \left(\int_0^s |\theta'(t)|^2 dt \right)^{1/2} \leq ||\theta'||_2 \cdot s^{1/2}.
$$

Hence, we take $\delta > 0$ and split

$$
\|\theta\|_{L^2(dt/|t|)}^2 = \int_{[-\delta,\delta]} \frac{|\theta(t)|^2}{|t|} dt + \int_{\mathbb{R}\setminus[-\delta,\delta]} \frac{|\theta(t)|^2}{|t|} dt
$$

$$
\leq 2\delta \|\theta'\|_{L^2}^2 + \frac{1}{\delta} \|\theta\|_2^2.
$$

Choosing $\delta > 0$ so as to minimize the right-hand side above directly implies the claim.

Finally, we prove the fact mentioned in the introduction that $\psi \in \mathcal{H}_0$ implies that $\hat{\varphi} \in C(\mathbb{R})$. We remark first that, as previously noted, $\psi \in L^2$ implies $\varphi \in L^2(x^2 dx)$. This shows that the Fourier transform of $x\varphi(x)$ is L^2 -integrable, which is equivalent to the assertion that $\hat{\varphi} \in H^1(\mathbb{R})$. The next result then proves the claim that $\hat{\varphi} \in C(\mathbb{R})$.

Proposition 2.2. Let $f \in H^1(\mathbb{R})$. Then f may be represented as a continuous function.

Proof. Let f' denote the derivative of f as a tempered distribution. Then we have that, for $g \in \mathcal{S}(\mathbb{R})$,

$$
\langle f', g \rangle = -\langle f, g' \rangle = -\langle \hat{f}, \hat{g'} \rangle.
$$

Since $\hat{f} \in L^1_{loc}(\mathbb{R} \setminus \{0\})$, we may write the pairing $\langle \hat{f}, \hat{g'} \rangle = -\pi i \cdot \int_{\mathbb{R}} \hat{f}(\xi) \xi \overline{\hat{g}(\xi)} d\xi$, whenever \hat{g} has compact support not containing 0. Since the latter integral is bounded by $\pi ||f||_{H_1} ||g||_2$, we may actually extend this identification to all $q \in \mathcal{S}(\mathbb{R})$. Hence,

$$
\langle f', g \rangle = \int_{\mathbb{R}} (\pi i \xi) \widehat{f}(\xi) \overline{\widehat{g}(\xi)} d\xi = \langle \widehat{\pi i \xi \cdot \widehat{f}, g} \rangle.
$$

This shows that \hat{f}' is a locally integrable function, and $\hat{f}'(\xi) = \pi i \xi \hat{f}(\xi) \in L^2$, which shows that $f' \in L^2(\mathbb{R})$. Now, let $h \in \mathcal{S}'(\mathbb{R})$ denote another distribution such that $h' = f'$. We then have

$$
\langle h - f, g' \rangle = -\langle h' - f', g \rangle = 0
$$

for all $g \in \mathcal{S}(\mathbb{R})$. On the other hand,

$$
\langle h-f,g'\rangle=\langle \widehat{h-f},(\pi i\xi)\cdot \widehat{g}\rangle.
$$

Let then η be compactly supported, and $0 \notin \text{supp}(\eta)$. Then $\eta(\xi) = \pi i \xi \hat{g}(\xi)$ for some $g \in \mathcal{S}(\mathbb{R})$, and hence $\langle \widehat{h - f}, \eta \rangle = 0$. Thus, we conclude that $\widehat{h - f}$ is supported at 0. Thus, $\widehat{h - f} = \sum_{k=0}^{N} a_k \partial_x^k$ for some $N \in \mathbb{N}$. Since this implies that $h - f$ is a polynomial, and since $\langle h - f, g' \rangle = 0$ for all $q \in \mathcal{S}(\mathbb{R})$, we conclude that $h - f$ is a constant.

In order to conclude the proof, note that $f_0(x) = \int_0^x f'(t) dt$ satisfies that $f'_0 = f'$ in the sense of distributions. Since f_0 is (Hölder) continuous, and since by the considerations above f_0 and f differ by a constant, f is continuous, as desired.

2.3. Main equivalences and the operator T . In this section, we shall begin to analyze the operator described in the Introduction, which maps a "reasonable" function f to the restriction to the y-axis of the solution u to the Klein-Gordon equation

$$
\partial_x \partial_y u + \pi^2 u = 0,
$$

with $u(x, 0) = f(x)$. Indeed, we shall define the operator T by its action on functions $\theta \in C_c^{\infty}(\mathbb{R})$ such that $\theta(0) = 0$ as

$$
T\theta(\xi) = -\pi\sqrt{|\xi|} \int_{\{y:\ \xi y < 0\}} \frac{\theta(y)}{\sqrt{|y|}} J_1(2\pi\sqrt{|\xi y|}) dy,\tag{2.5}
$$

where J_{ν} denotes the Bessel function of the first kind and order $\nu \in \mathbb{R}$. Clearly, for $\theta \in C_c^{\infty}(\mathbb{R})$, the operator T above is well-defined and finite for each $\xi \in \mathbb{R}$.

The definition of this operator seems cumbersome and unrelated to the Klein-Gordon equation at first, but, as we shall see below, a reformulation of it shows that T is the most "natural" operator to consider when analyzing Heisenberg uniqueness pairs involving the hyperbola Γ , and in fact matches the previously stated relationship. For that purpose, we start with the following result.

Lemma 2.1. For $\theta \in C_c^{\infty}(\mathbb{R})$ with $\theta(0) = 0$, define $\Theta_{\pm} : \mathbb{R}^4 \to \mathbb{C}$ by

$$
\Theta_{\pm}(u) = \frac{\theta(\pm |u|^2)}{|u|^2}.
$$
\n(2.6)

Then

$$
T\theta(\xi) = -|\xi|\mathcal{F}_4(\Theta_{-\text{sgn}(\xi)})(\sqrt{|\xi|}),\tag{2.7}
$$

where \mathcal{F}_4 denotes the Fourier transform in dimension 4.

Proof. We suppose that $\xi < 0$, with the other case being entirely analogous. We recall the formula for the Fourier transform of a radial function $g : \mathbb{R}^4 \to \mathbb{C}$ in $L^1(\mathbb{R}^4)$:

$$
\mathcal{F}_4(g)(\eta) = 2\pi \int_0^\infty g(s) \left(\frac{s}{|\eta|}\right) J_1(2\pi |\eta|s) \, s \, ds.
$$

Changing variables $y = s^2$ in the definition [\(2.5\)](#page-8-0), we get

$$
T\theta(\xi) = -2\pi\sqrt{|\xi|} \int_0^\infty \theta(s^2) J_1(2\pi\sqrt{|\xi|}s) ds
$$

=
$$
-2\pi |\xi| \int_0^\infty \frac{\theta(s^2)}{s^2} \left(\frac{s}{\sqrt{|\xi|}}\right) J_1(2\pi\sqrt{|\xi|}s) s ds,
$$

which concludes the proof.

We are now able to prove our main equivalence for the definition of T , which enables us to connect it to Heisenberg Uniqueness pairs:

Proposition 2.3. Under the conditions above, if $\psi \in L^1$ is such that $\hat{\psi} = \theta$, then

$$
T\theta(\xi) = \frac{1}{\sqrt{2}} \int_{\mathbb{R}} \psi(t) e^{-\pi i \xi/t} dt.
$$
\n(2.8)

Proof. Let us suppose that $\xi < 0$, since the other case is entirely analogous. Then the considerations in [\[3,](#page-34-5) Section 2.1] together with Lemma [2.1](#page-8-1) imply the following: if

$$
\theta(t) = \frac{1}{\sqrt{2}} \int_{\mathbb{R}} e^{-\pi i t s} \psi(s) \, ds,
$$

then the four-dimensional Fourier transform of Θ at η is given by

$$
-\frac{1}{\sqrt{2}|\eta|^2}\int_{\mathbb{R}}e^{\pi i|\eta|^2/t}\psi(t)\,dt.
$$

Here, we made crucial use of the fact that $\theta(0) = \hat{\psi}(0) = 0$, in order to use the results from [\[3\]](#page-34-5). Replacing then η by $\sqrt{|\xi|}$ and using that $|\xi| = -\xi$ yields the claim.

Remark 2.1. Proposition [2.3](#page-9-0) is similar in spirit to Proposition 5.2.1 in [\[28\]](#page-35-13): indeed, by simply using a formal Fubini argument in [\(2.8\)](#page-9-1) with that result, one may (formally) recover Proposition [2.3.](#page-9-0) As a matter of fact, the formal argument outlined here may be made formal through multiplication with a decay-inducing factor and then taking limits. For brevity of exposition, however, we decided not to include such computations.

A crucial feature of Proposition [2.3](#page-9-0) is that it allows us to extend the definition of T to general continuous, bounded functions θ , satisfying $\theta = \hat{\psi}$, where $\psi \in L^1$ and $\hat{\psi}(0) = 0$. Indeed, we may always write $T\theta = \hat{\varphi}$ whenever $\theta \in C_c^{\infty}(\mathbb{R})$, where $\varphi(t) = t^{-2}\psi(1/t)$. Since $\varphi \in L^1$ whenever $\psi \in L^1$, we may use this equivalent definition to extend T to the more general class of $\theta \in (\widehat{L}^1)_0 = {\theta =$ $\widehat{\psi}, \psi \in L^1, \theta(0) = 0$.

Finally, note that if u is a solution to $\partial_x \partial_y u + \pi^2 u = 0$, with the additional constraint that $u(x, 0) = \theta(x)$, then if we may write $\hat{\psi} = \hat{\theta}$, it follows from a direct Fourier characterization argument that $u(x, y) = \frac{1}{\sqrt{2}}$ $\frac{1}{2} \int_{\mathbb{R}} \psi(t) e^{-\pi i (x \cdot t + y/t)} dt$, and Proposition [2.3](#page-9-0) shows that $u(0, y) = T\theta(y)$, concluding hence the claim about the relationship between T and the Klein-Gordon equation.

In the next result we highlight some other important quantitative facts about the operator T in terms of decay and regularity:

Proposition 2.4. Under the same hypotheses of Proposition [2.3,](#page-9-0) the following assertions hold:

(I) For all admissible θ , we have that

$$
||T\theta||_{L^2(dt/|t|)} = ||\theta||_{L^2(dt/|t|)}
$$

$$
||T\theta||_{L^2(\mathbb{R})} = \frac{1}{\sqrt{2\pi}} ||\theta'||_{L^2(\mathbb{R})}.
$$

,

- (II) $(T \circ T)\theta = \theta$ holds for any θ admissible;
- (III) $\text{supp}(\theta) \subset \mathbb{R}_+$ if and only if $\text{supp}(T\theta) \subset \mathbb{R}_-$.

Proof. We start with the proof of the first equality in (I) . We have, by (2.6) , (2.7) and Plancherel,

$$
\int_{\mathbb{R}_-} \frac{|T\theta(\xi)|^2}{|\xi|} d\xi = 2 \int_{\mathbb{R}_+} |\mathcal{F}_4(\Theta)(t)|^2 t^3 dt = 2 \int_{\mathbb{R}_+} |\Theta(t)|^2 t^3 dt = \int_{\mathbb{R}_+} \frac{|\theta(s)|^2}{|s|} ds.
$$

By doing the same for the integral over \mathbb{R}_+ , we conclude that part. For the second equality, we may write, with the aid of Proposition [2.3,](#page-9-0)

$$
||T\theta||_2^2 = \int_{\mathbb{R}} |t^{-2}\psi(1/t)|^2 dt = \int_{\mathbb{R}} t^2 |\psi(t)|^2 dt = \frac{1}{2\pi^2} ||\theta'||_2^2,
$$

as desired. For (II), note that $T\theta$ is admissible if θ is: indeed, Part (I) shows that $T\theta \in L^2(dt/|t|)$. Moreover, since T is given by the Fourier transform of the function $t^{-2}\psi(1/t) \in L^1$, it is continuous. Finally, an analogous argument to the one in Part (I) with the definition of T shows that $(T\theta)' \in L^2$. Hence, the operator $(T \circ T)\theta$ is well-defined.

For the desired identity, we first note that

$$
\widetilde{T\theta}(y) = y^{-2}\psi(1/y),
$$

and thus

$$
\theta(t) = \widehat{\psi}(t) = \frac{1}{\sqrt{2}} \int_{\mathbb{R}} y^{-2} \widetilde{T} \theta(1/y) e^{-\pi i y t} dy = \frac{1}{\sqrt{2}} \int_{\mathbb{R}} \widetilde{T} \theta(y) e^{-\pi i t/y} dy = (T \circ T) \theta(t).
$$

Finally, Part (III) follows directly from (2.5) .

Note the following direct consequence of Proposition [2.4:](#page-10-0) $\theta \in L^2(\mathbb{R}) \cap H^1(\mathbb{R})$ if, and only if, $T\theta \in L^2(\mathbb{R}) \cap H^1(\mathbb{R})$. We use this fact to state the following main equivalence result, which will be crucial throughout the rest of the paper:

Proposition 2.5. The following assertions are equivalent:

- (I) There exists $\psi \in L^2((1+t^2)dt)$ with $\hat{\psi}(0) = 0$ and $\int_{\mathbb{R}} \psi(t) e^{-\pi i a_n t} dt = 0 = \int_{\mathbb{R}} \psi(t) e^{-\pi i b_n/t} dt =$ $0, \forall n \in \mathbb{Z}$;
- (II) There exists $\theta \in L^2(dt/|t|) \cap L^2(\mathbb{R}) \cap H^1(\mathbb{R})$, such that $\theta(a_n) = T\theta(b_n) = 0$, $\forall n \in \mathbb{Z}$.

The proof of this result is a direct consequence of Proposition [2.1,](#page-7-0) Proposition [2.3](#page-9-0) and Proposition [2.4,](#page-10-0) and its statement will provide us with the right framework for both our positive and negative results.

3. Proof of Theorem [3](#page-5-0)

3.1. Preliminary Reductions. In order to begin our process of building a function ψ satisfying the hypotheses of Theorem [3,](#page-5-0) we start by recalling some concepts and reductions from [\[32\]](#page-35-18).

Indeed, first and foremost, we define the *density of order p* (if it exists) of a sequence $Z = \{z_j\}_{j \in \mathbb{Z}}$ of points in the complex plane as

$$
\mathcal{D}_p(Z) = \lim_{r \to \infty} \frac{\#(Z \cap D_r(0))}{r^p}.
$$

We also need the concept of *smooth sequences*, as defined in [\[32\]](#page-35-18): a sequence Z with p-density $\mathcal{D}_p(Z)$ is called *p*-smooth if $z_j = r_j e^{i\theta}$, for some fixed $\theta \in [0, 2\pi)$, and the following conditions are met:

- (I) The function $r \mapsto |\#(Z \cap D_r(0)) \mathcal{D}_p(Z)r^p|$ is bounded for $r > 0$;
- (II) There is $d > 0$ such that $r_{j+1} r_j \ge d(1+r_j)^{1-p}$.

Our first reduction will show that, if A and B are sequences as in the statement of Theorem [3,](#page-5-0) then we may 'expand' them a bit so that they become more regular, while still satisfying the same asymptotic properties in terms of density:

Lemma 3.1. Let $C = \{c_n\}_{n\geq 0}$ be an increasing sequence in \mathbb{R}_+ , such that, for some $\sigma > 0$,

$$
\liminf_{n \to \infty} (c_{n+1} - c_n) > \sigma.
$$

Then, for any $D > 1/\sigma$, there exists a 1-smooth sequence $C' \supset C$ with $\mathcal{D}_1(C') = D$ and such that $|C' \setminus C| = +\infty.$

We refer the reader to [\[32,](#page-35-18) Claim 7] for a proof of this fact. Using Lemma [3.1,](#page-11-1) we then replace A and B by A' and B', where $A' \cap (0, +\infty)$ is a sequence given by Lemma [3.1](#page-11-1) for $A \cap (0, +\infty)$, with its definition for negative values being analogous. The set B' is built in an entirely analogous way from B. By the Lemma again, we may still assume that $\mathcal{D}_1(A')$, $\mathcal{D}_1(B') < 1$. We shall then work with A' , B' throughout the steps of the proof below.

3.2. Entire functions with prescribed zeros. Let $k : [-\pi, \pi] \to \mathbb{R}$ be defined by

$$
k(\theta) = (\pi \beta) \sin(2\theta) - \gamma \cos(2\theta)
$$
\n(3.1)

whenever $\theta \in [0, \pi/2]$, and extended to $\theta \in [-\pi, \pi]$ such that it becomes even and symmetric with respect to $\pi/2$. For such k, we call a discrete set $\mathcal{Z} = \mathcal{Z}_1^+ \cup \mathcal{Z}_1^- \cup \mathcal{Z}_2^+ \cup \mathcal{Z}_2^- \subset \mathbb{C}$, where $\mathcal{Z}^{\pm}_{j}\subset$ $\left\{\arg(z)=\frac{(j\pm 1)\pi}{2}\right\}$ $\Big\}$, k–regular if:

(I) \mathcal{Z}_j^{\pm} has density $m_j/4\pi$ with respect to exponent 2. That is, it holds that

$$
|\mathcal{Z}_j^{\pm} \cap (0, r e^{i\theta_j^{\pm}})| \sim r^2 m_j / 4\pi
$$

as $r \to \infty$, where $m_1 = 2\pi\beta$, $m_2 = 2\gamma$, with $\theta_j^{\pm} = \frac{(j\pm 1)\pi}{2}$;

(II) the disks $\{D_z\}_{z\in\mathcal{Z}} = \{B(z, c\cdot(1+|z|)^{-1})\}_{z\in\mathcal{Z}}$ are all *disjoint* for some $c > 0$.

We now state the following result by Levin, which will be crucial in our construction:

Theorem 5 (Theorem 5, Ch. II, [\[35\]](#page-36-9)). Let k be as in [\(3.1\)](#page-11-2). Then, for any k-regular set \mathcal{Z} , there exists an entire function S, whose zeroes are simple and coincide with Z, such that, for every $\varepsilon > 0$,

$$
|S(re^{i\theta})| \le C_{\varepsilon}e^{(k(\theta)+\varepsilon)r^2}, \,\, everywhere \,\, in \,\, \mathbb{C},\tag{I}
$$

$$
|S(re^{i\theta})| \ge c_{\varepsilon}e^{(k(\theta)-\varepsilon)r^2}, \text{ whenever } w = re^{i\theta} \notin \bigcup_{z \in \mathcal{Z}} D_z.
$$
 (II)

We refer the reader to either the original work of Levin [\[35\]](#page-36-9) or the recent manuscript [\[32\]](#page-35-18), where a self-contained proof of this fact, which works in our context, is available in the appendix.

As consequence of (I) and (II) in Theorem [5,](#page-12-0) we have that, for $z \in \mathcal{Z}_i$,

$$
|S'(z)| \ge c_{\varepsilon} e^{(k(\arg z) - \varepsilon)|z|^2}.
$$
\n(3.2)

This can be seen, for instance, by applying the mean value property to the harmonic function $\log(|S(w)/(w-z)|)$ in the disk D_z .

3.3. Almost-interpolation bases. Our next step is to construct functions by using Theorem [5](#page-12-0) which will give rise to our iteration scheme. We relabel the sequences A and B so that

$$
A \cap (0, +\infty) = \{a_k\}_{k \ge 1}, \qquad B \cap (0, +\infty) = \{b_k\}_{k \ge 1}.
$$

In what follows, we will mainly deal with the set A , but the same construction can be repeated to B by a verbatim adaptation.

We define \tilde{A}_1 to be the sequence of elements of the form $\sqrt{a_n}$ whenever $n > 0$, and we let \tilde{A}_2 analogously be the sequence of elements of the form $\sqrt{-a_n}$ whenever $n \leq 0$.

We start by taking $\mathcal{Z}_1^+(1) = \tilde{A}_1$, and setting $\mathcal{Z}_1^-(1) = -\mathcal{Z}_1^+(1)$. We then add two other sets $\mathcal{Z}_2^{\pm}(1)$ so that the properties in the definition of k-regularity are fulfilled, and $-\mathcal{Z}_2^+(1) = \mathcal{Z}_2^-(1)$. The exact choice of these sets is not important, as long as the set $\mathcal{Z}(1) := \cup_{\pm} \cup_{j=1,2} \mathcal{Z}_j^{\pm}(1)$ obtained in the end is origin-symmetric and is k_1 -regular, for some function k_1 as above. By repeating the same construction for $\mathcal{Z}_1^+(2) = \tilde{A}_2$, we obtain in a similar manner a k_2 -regular set $\mathcal{Z}(2)$ associated to a (possibly) different function k_2 . Note, however, that the functions k_1, k_2 are (crucially) only tied to $\mathcal{Z}(1)$, $\mathcal{Z}(2)$ through the densities of these sets.

We then use Theorem [5](#page-12-0) for $\mathcal{Z}(i), i \in \{1,2\}$. This yields an entire function S_i in each case with the desired zero properties at the sets $\mathcal{Z}(i)$, respectively. Moreover, we may assume that such a function is always *even*. This follows directly from the construction of S_i , as given either in [\[35\]](#page-36-9) or [\[32\]](#page-35-18): if we let $E_2(z) = (1-z)e^{z+z^2/2}$, then S_i is obtained as a limit as $R \to \infty$ of the functions

$$
S_{i,R}(z) = e^{\alpha z + \beta z^2} \prod_{\lambda \in \mathcal{Z}_i \cap D_R(0)} E_2(z/\lambda),
$$

where $\alpha, \beta \in \mathbb{C}$. We claim that, if S_i satisfies the conditions of Theorem [5,](#page-12-0) then so does $R_i(z) :=$ $e^{-\alpha z} \cdot S_i(z)$. Indeed, the zero assumption does not change by multiplying by complex exponentials. Moreover, since $|e^{\alpha z}| \leq C_{\varepsilon} e^{\varepsilon |z|^2}$ for all $\varepsilon > 0$, the bounds (I) and (II) in Theorem [5](#page-12-0) are fulfilled. Therefore, we may suppose without loss of generality that $\alpha = 0$ in the definition of $S_{i,R}$.

Since we have shown that we may take $S_{i,R}$ to be even by construction for each R, then we may also take S_i to be even, as desired. Consider then, for a fixed $a \in \tilde{A}_i$, the function

$$
\mathcal{S}_{i,a}(z) = \frac{S_i(z)}{z^2 - a^2}.
$$

Since S_i is even, this new function is again an entire function. By applying the maximum modulus principle to this function on a small disk around a, and using the bounds we have on S_i outside it, we can easily conclude that it satisfies

$$
|\mathcal{S}_{i,a}(z)| \lesssim C_{\varepsilon} e^{(k(\theta) + \varepsilon)|z|^2}
$$
\n(3.3)

in all of \mathbb{C} , uniformly on $a \in \tilde{A}_i$.

Now, for any $n \in \mathbb{Z}$, let $a(n)$ be the positive real number for which $a(n)^2 = \pm a_n$, $a(n) \in \tilde{A}_1 \cup \tilde{A}_2$. We then define a function $\theta_n : \mathbb{R} \to \mathbb{C}$ in the following way: if $a(n) \in \tilde{A}_1$, we define

$$
\theta_n(t) = \begin{cases} t \cdot \mathcal{S}_{1,a(n)}(\sqrt{t}), & \text{if } t > 0, \\ 0, & \text{if } t < 0. \end{cases}
$$

The definition of θ_n for $a(n) \in \tilde{A}_2$ is analogous. These functions will serve as "approximate" interpolation functions: θ_n vanishes on A, except for the point a_n . Although no analogous interpolation property holds for $T(\theta_n)$, we have a (weak) surrogate of that through the following strong decay bounds:

Lemma 3.2. There exist absolute constants $\alpha' > \alpha'' > 0$ such that the following holds. For each $x \in \mathbb{R}$, each $\xi \in \mathbb{R}$ and each $n \in \mathbb{Z}$, we have

$$
|\theta_n(x)| \le Ce^{-\alpha''|x|}, \quad |T(\theta_n)(\xi)| \le Ce^{-\alpha'|\xi|}.
$$

Proof. Without loss of generality, let us assume that $a_n > 0$. By Proposition [2.4,](#page-10-0) we have that $\text{supp}(\mathcal{T}\theta_n) \subset \mathbb{R}_+$, and hence it is enough to prove the bound for $\xi < 0$. We shall write $a := a(n)$ throughout the proof.

We then recall [\(2.7\)](#page-9-2), which, for $\Theta_n(u) = \frac{\theta_n(|u|^2)}{|u|^2}$ $\frac{(|u|^2)}{|u|^2}$ for $u \in \mathbb{R}^4$, allows us to write

$$
T(\theta_n)(\xi) = -|\xi| \cdot \mathcal{F}_4(\Theta_n)(\sqrt{|\xi|}). \tag{3.4}
$$

By recalling the definition of θ_n , we have that

$$
\mathcal{F}_4(\Theta_n)(\eta) = \int_{\mathbb{R}^4} \mathcal{S}_{1,a}(|x|) e^{-2\pi ix \cdot \eta} dx.
$$

On the other hand, $S_{1,a}$ is an *even* function of z, which allows it to be written as $S_{1,a}(z) = H_{1,a}(z^2)$, for some entire function $H_{1,a}$. Note that $|H_{1,a}(re^{i\theta})| \leq e^{-cr}$ if θ is close to either 0 or π , by the properties of $\mathcal{S}_{i,a}$.

Hence, taking these considerations into account, we may write the integral defining $\mathcal{F}_4(\Theta_n)$ through Fubini's theorem as a four-fold integral, as follows:

$$
\mathcal{F}_4(\Theta_n)(\eta) = \int_{\mathbb{R}^4} H_{1,a}(x_1^2 + x_2^2 + x_3^2 + x_4^2) e^{-2\pi ix \cdot \eta} dx.
$$

By the previously mentioned decay properties of $H_{1,a}$, we are able to change contours in each one of those one-dimensional integrals. This allows us to write

$$
\mathcal{F}_4(\Theta_n)(\eta) = \int_{\mathbb{R}^4} H_{1,a}(\|x+iy\|^2) e^{-2\pi i (x+iy)\cdot \eta} dx.
$$
 (3.5)

Here, we use the convention that, for $x, y \in \mathbb{R}^4$, $||x + iy||^2 := \sum_{j=1}^4 (x_j + iy_j)^2$. From [\(3.5\)](#page-14-0), we may bound $|\mathcal{F}_4(\Theta_n)(\eta)|$ as follows. First, let $y = -t\eta$, where $t > 0$ will be chosen later. Then

$$
|\mathcal{F}_4(\Theta_n)(\eta)| \le e^{-2\pi|y||\eta|} \left| \int_{\mathbb{R}^4} |H_{1,a}(\|x+iy\|^2)|e^{-\varepsilon r} e^{\varepsilon r} dx \right|
$$

$$
\le e^{-2\pi|y||\eta|} \sup_{x \in \mathbb{R}^4} \left(|H_{1,a}(\|x+iy\|^2) e^{\varepsilon r}| \times \left(\int_{\mathbb{R}^4} e^{-\varepsilon r} dx \right) \right). \tag{3.6}
$$

where we write $||x + iy||^2 = |x|^2 - |y|^2 + 2i\langle x, y \rangle =: re^{i\theta}$. Here, $\varepsilon > 0$ will be chosen later. We now note that

$$
r = ((|x|^2 - |y|^2)^2 + 4\langle x, y \rangle^2)^{1/2} \ge ||x|^2 - |y|^2|,
$$

which implies

$$
\int_{\mathbb{R}^4} e^{-\varepsilon r} \, dx \le C \int_0^\infty e^{-\varepsilon |s^2 - |y|^2} |s^3 \, ds = \frac{C}{2} \int_0^\infty e^{-\varepsilon |s' - |y|^2} |s' \, ds'.
$$

On the other hand, it is not hard to see that the expression on the right-hand side above is bounded by $C_{\varepsilon}(1+|y|^2)$. Thus, from [\(3.6\)](#page-14-1), we obtain

$$
|\mathcal{F}_4(\Theta_n)(\eta)| \le C_{\varepsilon} (1+|y|^2) e^{-2\pi |y||\eta|} \sup_{x \in \mathbb{R}^4} (|H_{1,a}(\|x+iy\|^2)|e^{\varepsilon r}).
$$

From the bounds available on $S_{1,a}$, we conclude that $|H_{1,a}(re^{i\theta})| \leq C_{\varepsilon}e^{(k(\theta/2)+\varepsilon)r}$. Thus,

$$
|H_{1,a}(\|x+iy\|^2)|e^{\varepsilon r}\leq C_\varepsilon e^{(k(\theta/2)+2\varepsilon)r}.
$$

We now note that, if $s < 1$ is sufficiently close to 1, and $\gamma = \pi s \omega^{-2}$, with ω sufficiently large, then any function k defined as in (3.1) satisfies

$$
k(\theta) < \pi \omega^2 \sin^2(\theta), \quad \text{whenever } \theta \in [0, \pi/2]. \tag{3.7}
$$

For a proof of this fact, we refer the reader to $[32, \text{Claim 8}]$. Since the inequality in (3.7) is *strict* for the whole compact interval $[0, \pi/2]$, if we choose $\varepsilon > 0$ small enough, we will have $k(\theta/2) + 2\varepsilon \leq$ $\pi\omega^2 \sin^2(\theta/2)$, whenever $\theta \in [-\pi, \pi]$. Thus, for such $\varepsilon > 0$, we obtain the following bound:

$$
|\mathcal{F}_4(\Theta_n)(\eta)| \le C'(1+|y|^2) \sup_{x \in \mathbb{R}^4} e^{2\pi \left(\frac{\omega^2}{2}\sin^2(\theta/2)r - |y||\eta|\right)}.
$$

On the other hand, $r \leq |x|^2 + |y|^2$ and $\sin^2(\theta/2) = \frac{1}{2}(1 - \cos(\theta))$, which implies that $\sin^2(\theta/2)r =$ 1 $\frac{1}{2}(r - r\cos(\theta)) \le \frac{1}{2}(|x|^2 + |y|^2 - (|x|^2 - |y|^2)) = |y|^2$. This shows finally that

$$
|\mathcal{F}_4(\Theta_n)(\eta)| \le C'(1+|y|^2) \sup_{x \in \mathbb{R}^4} e^{2\pi \left(\frac{\omega^2}{2}|y|^2 - |y||\eta|\right)}.
$$
 (3.8)

Recalling that $y = -t\eta$, $t > 0$, and optimizing on the parameter t, we get from [\(3.8\)](#page-14-3) that

$$
|\mathcal{F}_4(\Theta_n)(\eta)| \lesssim (1+|\eta|^2)e^{-\frac{\pi}{\omega^2}|\eta|^2} = (1+|\eta|^2)e^{-\frac{\gamma}{s}|\eta|^2}.
$$
 (3.9)

Gathering [\(3.9\)](#page-14-4) with [\(3.4\)](#page-13-0), the definition of θ_n , [\(3.3\)](#page-13-1) and [\(3.1\)](#page-11-2), one is able to finish the proof. \Box

With Lemma [3.2](#page-13-2) at hand, we define

$$
\varrho_n(t) := \frac{\theta_n(t)}{a(n)^2 \cdot \mathcal{S}_{i,a(n)}(a(n))}.
$$

By [\(3.2\)](#page-12-1), it follows that $|\mathcal{S}_{i,a(n)}(a(n))| \ge c_{\varepsilon}e^{-(\gamma+\varepsilon)|a_n|^2}$ holds for each n, which implies, together with Lemma [3.2,](#page-13-2) that there are $\alpha' > \alpha > \alpha'' > 0$ such that

$$
|\varrho_n(t)| \le Ce^{-\alpha''|t| + \alpha|a_n|},
$$

$$
|T(\varrho_n)(\xi)| \le Ce^{-\alpha'|\xi| + \alpha|a_n|}.
$$

Moreover, we have that $\text{supp}(\varrho_n) \subset \mathbb{R}_{\text{sgn}(n)}$, where we use the convention that $\text{sgn}(0) = -1$. Finally, note that $\varrho_n(a_j) = \delta_{n,j}$ whenever $sgn(n) = sgn(j)$, and that, since T is an isometry between H^1 and L^2 , we have that

$$
\|\varrho_n'\|_2 = c \cdot \|T(\varrho_n)\|_2 \le Ce^{\alpha |a_n|}.
$$

By running the same argument in the exact same fashion as we did above, we are also able to find a sequence of functions σ_n with the property that

$$
|\sigma_n(t)| \le Ce^{-\alpha''|t| + \alpha|b_n|},
$$

$$
|T(\sigma_n)(\xi)| \le Ce^{-\alpha'|\xi| + \alpha|b_n|},
$$

with $\text{supp}(\sigma_n) \subset \mathbb{R}_{\text{sgn}(n)}$, $\sigma_n(b_j) = \delta_{n,j}$ whenever $\text{sgn}(n) = \text{sgn}(j)$, and $\|\sigma'_n\|_2 \le Ce^{\alpha|b_n|}$.

3.4. Iteration Scheme and Conclusion. We are now ready to set up the main iteration process. Fix $\delta > 0$ sufficiently small (to be chosen later), and take $L > 0$ such that

$$
\sum_{k \in A_L} e^{(\alpha - \alpha')|a_k|} + \sum_{j \in B_L} e^{(\alpha - \alpha')|b_j|} < \delta,
$$

where $A_L = \{k \in \mathbb{Z} : |a_k| > L\}, B_L = \{j \in \mathbb{Z} : |b_j| > L\}.$ We further divide these sets into

$$
A_L^+ = A_L \cap (0, +\infty), \ A_L^- = A_L \setminus A_L^+, \ B_L^+ = B_L \cap (0, +\infty), \ B_L^- = B_L \setminus B_L^+.
$$

With that in mind, we define the Banach spaces \mathfrak{B}^{\pm} of pairs of sequences $\kappa = (\{s_k\}_{k \in A_L^{\pm}}, \{r_j\}_{j \in B_L^{\mp}})$ such that the norm

$$
\|\kappa\|_{\mathfrak{B}^{\pm}} := \sum_{k \in A_L^{\pm}} |s_k| e^{\alpha |a_k|} + \sum_{j \in B_L^{\mp}} |r_j| e^{\alpha |b_j|} < +\infty.
$$
 (3.10)

We are then able to show the following:

Proposition 3.1. For each sequence $\kappa \in \mathfrak{B}^{\pm}$, there exists a continuous function $f_{\pm} : \mathbb{R} \to \mathbb{C}$ with $f_{\pm} \in L^2 \cap H^1$ such that $\text{supp}(f_{\pm}) \subset \mathbb{R}_{\pm}$ and

$$
f_{\pm}(a_k) = s_k, \forall k \in A_L^{\pm},
$$

$$
Tf_{\pm}(b_j) = r_j, \forall j \in B_L^{\mp}.
$$

Proof. Suppose without loss of generality that we are dealing with \mathfrak{B}^+ . We start by taking

$$
f_0(t) = \sum_{k \in A_L^+} s_k \varrho_k(t) + \sum_{j \in B_L^-} r_j T(\sigma_j)(t).
$$
 (3.11)

Since supp $(\sigma_i) \subset \mathbb{R}_+$ if $j < 0$, we have that supp $(f_0) \subset \mathbb{R}_+$. Moreover, it follows directly from the definition that we have the following pointwise bound on f_0 and $T(f_0)$:

$$
|T(f_0)(t)| + |f_0(t)| \le C ||\kappa||_{\mathfrak{B}^+} e^{-\alpha''|t|}.
$$
\n(3.12)

Furthermore, we may bound the H^1 norm of f_0 as follows:

$$
||f_0'||_2 \le \sum_{k \in A_L^+} |s_k||\varrho_k'||_2 + \sum_{j \in B_L^-} |r_j|||T(\sigma_j)'||_2
$$

$$
\le C \left(\sum_{k \in A_L^+} |s_k|e^{\alpha|a_k|} + \sum_{j \in B_L^-} |r_j|e^{\alpha|r_j|} \right) \le C ||\kappa||_{\mathfrak{B}^+}.
$$
 (3.13)

In a completely analogous manner, by arguing with $(Tf_0)'$ instead and using that $T(Tf_0) = f_0$, we are able to conclude that $||f_0||_2 \leq C ||\kappa||_{\mathfrak{B}^+}$ as well.

We now define a linear map $\mathcal T$ from \mathfrak{B}^+ to itself, by $\mathcal T \kappa = (\{(\mathcal T^1\kappa)_k\}_{k \in A_L^+}, \{(\mathcal T^2\kappa)_j\}_{j \in B_L^-}),$ where

$$
(\mathcal{T}^1 \kappa)_k = f_0(a_k) - s_k,
$$

$$
(\mathcal{T}^2 \kappa)_j = T(f_0)(b_j) - r_j.
$$

Now, we simply notice that

$$
\sum_{k \in A_L^+} |(\mathcal{T}^1 \kappa)_k|e^{\alpha|a_k|} \le C \sum_{k \in A_L^+} \sum_{j \in B_L^-} |r_j|e^{(\alpha - \alpha')|a_k| + \alpha|b_j|}
$$

$$
\le C \|\kappa\|_{\mathfrak{B}^+} \sum_{k \in A_L^+} e^{(\alpha - \alpha')|a_k|} < C\delta \|\kappa\|_{\mathfrak{B}^+}.
$$

The same argument applied to $(\mathcal{T}^2\kappa)_j$ shows that

$$
\|\mathcal{T}\kappa\|_{\mathfrak{B}^+}\leq 2C\delta\|\kappa\|_{\mathfrak{B}^+}.
$$

Take thus $\delta < \frac{1}{4C}$ above. We now define f_1 analogously as in [\(3.11\)](#page-15-0), but with $\mathcal{T}\kappa$ in place of κ , and in general we define f_n recursively to be the function interpolating $\mathcal{T}^{(n)}\kappa$ as coefficients, as in [\(3.11\)](#page-15-0). Here, $\mathcal{T}^{(n)}$ denotes the *n*−th iterate of the map $\mathcal T$ defined above. We claim that

$$
f_+ := -\sum_{n\geq 0} f_n
$$

satisfies the hypotheses of the proposition. Indeed, since we showed that $\mathcal T$ is a contraction in \mathfrak{B}^+ , [\(3.12\)](#page-16-0) implies that the function f_+ , as well as $T f_+$, is pointwise bounded by $e^{-\alpha''|t|}$. The telescoping nature of the definition of $\mathcal T$ then implies that $f_+(a_k) = s_k$ for each $k \in A_L^+$, as well as $Tf_+(b_j) = r_j$, for all $j \in B_L^-$. Moreover, since f_+ is an uniform limit of continuous functions, it is itself continuous, and since supp $(f_n) \subset \mathbb{R}_+$, the same holds for f_+ . Finally, from [\(3.13\)](#page-16-1) we obtain that $f_+ \in L^2 \cap H^1$ as desired, with the bound

$$
||f_+||_2 + ||f'_+||_2 \leq C ||\kappa||_{\mathfrak{B}^+}
$$

being additionally fulfilled. \square

Recall that, in the beginning of the proof, we replaced A and B by A' and B', with $|A' \setminus A|$ = $|B' \setminus B| = +\infty$. It follows thus that the arguments above for such modified sets results, in the same

way, in a solution to the respective interpolation problem. We let then s_k, r_j be sequences such that $s_k = 0 = b_j$ if k and j are indices of an element of the original sets A and B, respectively, and set them as arbitrary values in the complement $A' \setminus A$, $B' \setminus B$, with the only constraint that they do not violate [\(3.10\)](#page-15-1). This shows that the space of functions which vanish on $A_{\pm} \setminus [-L, L]$ with their image under T vanishing on $B_{\pm} \setminus [-L, L]$ is infinite-dimensional.

On the other hand, since there are only finitely many points of A_{\pm} and B_{\mp} inside $[-L, L]$, we conclude that there must be an infinite-dimensional space of functions F_{\pm} such that

$$
F_{\pm}(a_k) = 0
$$
, $\forall k$ such that $sgn(k) = \pm$,
 $TF_{\pm}(b_j) = 0$, $\forall j$ such that $sgn(j) = \mp$.

Since the support of F_{\pm} is in \mathbb{R}_{\pm} , and the support of TF_{\pm} is in \mathbb{R}_{\mp} , the vanishing statement [\(1.3\)](#page-5-1) ψ_{\pm} such that $\widehat{\psi_{\pm}} = F_{\pm}$. All that is left is to show that ψ_{\pm} are integrable: this can be achieved by noticing that, by our construction, we have that

$$
||F_{\pm}||_2 + ||F'_{\pm}||_2 < +\infty,
$$
\n(3.14)

and hence, since $\psi_{\pm} = \widetilde{F_{\pm}}$, then

$$
\|\psi_{\pm}\|_{L^{1}(\mathbb{R})} \leq C \left(\int_{\mathbb{R}} |\psi_{\pm}(t)|^{2} (1+t^{2}) dt \right)^{1/2} \leq C' \left(\|F_{\pm}\|_{2} + \|F'_{\pm}\|_{2} \right) < +\infty,
$$

as desired, concluding the proof of Theorem [3.](#page-5-0)

4. Proof of Theorems [1](#page-3-0) and [2](#page-4-0)

4.1. Proof of Theorem [1.](#page-3-0) We start with the 'if' part. The proof below follows essentially the same arguments as in [\[28\]](#page-35-13); it has been brought to our attention that the authors of [\[20\]](#page-35-16) have recently provided in [\[21\]](#page-35-17) an analogous argument. We decided, however, to include it here for completeness.

Indeed, suppose we have $\psi \in L^1(\mathbb{R})$ such that

$$
\int_{\mathbb{R}} e^{-\pi i \tau (n+\theta)} \psi(\tau) d\tau = 0, \ \forall n \in \mathbb{Z},
$$
\n(4.1)

$$
\int_{\mathbb{R}} e^{-\pi i(\beta n)/\tau} \psi(\tau) d\tau = 0, \ \forall \, n \in \mathbb{Z}.
$$
\n(4.2)

Let $M_{\theta}\psi(\tau) := e^{-\pi i \theta \tau} \psi(\tau)$. Equation [\(4.1\)](#page-17-1) is equivalent to $\sum_{j\in\mathbb{Z}} (M_{\theta}\psi)(2j+t) = 0$ for almost every $t \in \mathbb{R}$. Analogously, changing variables $\tau \mapsto \beta/\tau$ in [\(4.2\)](#page-17-2), we obtain that

$$
\sum_{j\in\mathbb{Z}}\frac{1}{(\tau+2j)^2}\psi\left(\frac{\beta}{\tau+2j}\right)=0,
$$

for almost all $t \in \mathbb{R}$. Isolating the terms associated with $j = 0$ in both of these relations, we obtain

$$
\psi(t) = -\sum_{j \in \mathbb{Z}^*} e^{-2\pi i j \theta} \psi(2j + t),\tag{4.3}
$$

$$
\frac{1}{t^2}\psi(\beta/t) = -\sum_{j\in\mathbb{Z}^*} \frac{1}{(t+2j)^2}\psi\left(\frac{\beta}{t+2j}\right). \tag{4.4}
$$

Equivalently, (4.4) rewrites as

$$
\psi(s) = -\sum_{j \in \mathbb{Z}^*} \frac{\beta^2}{(\beta + 2sj)^2} \psi\left(\frac{\beta s}{\beta + 2sj}\right). \tag{4.5}
$$

Combining (4.3) and (4.5) yields

$$
\psi(s) = \sum_{k,j \in \mathbb{Z}^*} e^{-2\pi i j \theta} \frac{\beta^2}{(\beta + 2(s + 2k)j)^2} \psi \left(\frac{\beta(s + 2k)}{\beta + 2(s + 2k)j} \right).
$$
(4.6)

Define the operator $T_\beta: L^1[-1,1] \to L^1[-1,1]$ as

$$
T_{\beta}f(t) = \sum_{j\in\mathbb{Z}^*} \frac{\beta}{(t+2j)^2} f\left(\frac{\beta}{t+2j}\right).
$$

With this notation, [\(4.6\)](#page-18-2) implies that

$$
|\psi(t)| \le (T_{\beta}^2 |\psi|)(t)
$$
, for almost all $t \in (-1, 1)$. (4.7)

From this point on, we have to distinguish between cases: first of all, if $\beta \in (0,1)$, it suffices to use Proposition 3.13.1 in [\[28\]](#page-35-13), which says that $T_{\beta}^{2l}|\psi| \to 0$ in $L^1[-1,1]$. On the other hand, if $\beta = 1$, we only need to use [\[28,](#page-35-13) Proposition 3.13.13], which also guarantees to us that $\int_{-1+\delta}^{1-\delta} T_1^{2l} |\psi| \to 0$ as $l \to \infty$, for any $\delta > 0$. Thus, in either case, we obtain $\psi \equiv 0$ in (-1, 1). By [\(4.5\)](#page-18-1), $\psi \equiv 0$ on R.

For the 'only if' part, fix $\beta > 1$. We simply take a $\psi_0 \in L^1, \psi_0 \neq 0$, given by Theorem [3](#page-5-0) such that

$$
\widehat{\psi_0}(\sqrt{\beta}(n+\theta)) = 0, \forall n \in \mathbb{Z},
$$

$$
T\widehat{\psi_0}(\sqrt{\beta}n) = 0, \forall n \in \mathbb{Z}.
$$

By taking $\psi(t) = \psi_0(t/\sqrt{\beta})$, this shows that Λ_{β}^{θ} does not form a H.U.P. with the hyperbola, whenever $\beta > 1$, finishing the proof of that result.

4.2. Proof of Theorem [2.](#page-4-0) Since Theorem [3](#page-5-0) provides us, upon scaling, with the examples for the $\alpha\beta > 1$ case, we focus on the other two cases.

Applying a scaling if necessary we may assume $\alpha = \beta \leq 1$. Let ψ be a function satisfying [\(1.3\)](#page-5-1) for such sequences, and define $\varphi(t) = t^{-2}\psi(1/t)$. We use the definition [\(2.2\)](#page-6-2) in what follows.

Step 1: We first show that the conditions in Theorem [2](#page-4-0) imply that $\psi \in L^2((1+x^2) dx)$. Indeed, by the Poincaré-Wirtinger inequality [\(2.4\)](#page-7-1) applied to each interval (b_k, b_{k+1}) , we have

$$
\frac{1}{\pi^2} \int_{b_k}^{b_{k+1}} |\widehat{\varphi}'(t)|^2 dt \ge (b_{k+1} - b_k)^2 \frac{1}{\pi^2} \int_{b_k}^{b_{k+1}} |\widehat{\varphi}'(t)|^2 dt \ge \int_{b_k}^{b_{k+1}} |\widehat{\varphi}(t)|^2 dt.
$$

Summing the inequality above over $k \in [-N, N) \cap \mathbb{Z}$ yields

$$
\frac{1}{\pi^2} \int_{\mathbb{R}} |\hat{\varphi}'(t)|^2 dt \ge \frac{1}{\pi^2} \int_{b_{-N}}^{b_N} |\hat{\varphi}'(t)|^2 dt \ge \int_{b_{-N}}^{b_N} |\hat{\varphi}(t)|^2 dt.
$$
 (4.8)

Using Fatou's lemma on the right-hand side of [\(4.8\)](#page-18-3) then yields that $\hat{\varphi} \in L^2$, which is equivalent after a change of variables to $x \cdot \psi(x) \in L^2$, concluding this step.

Step 2: Suppose first $\alpha\beta < 1$, and that $\psi \neq 0$ satisfies the conditions of Theorem [2.](#page-4-0) After rescaling we may assume $\alpha = \beta < 1$. Then condition [\(1.3\)](#page-5-1) translates to $\hat{\psi}(a_n) = 0 = T\hat{\psi}(b_n)$. We have proved in step 1 that $\hat{\varphi} = T\hat{\psi}$ and $\hat{\psi}$ belong to $H^1(\mathbb{R})$. Then by the Poincaré-Wirtinger inequality (2.4) , we have

$$
\int_{a_n}^{a_{n+1}} \left| \frac{\widehat{\psi}'(x)}{\pi} \right|^2 dx \ge (a_{n+1} - a_n)^2 \int_{a_n}^{a_{n+1}} \left| \frac{\widehat{\psi}'(x)}{\pi} \right|^2 dx \ge \int_{a_n}^{a_{n+1}} |\widehat{\psi}(x)|^2 dx, \tag{4.9}
$$

with equality if and only if $\hat{\psi} = 0$ on (a_n, a_{n+1}) . Summing up all these inequalities and using Plancherel twice, we obtain:

$$
\int_{\mathbb{R}} t^2 |\psi(t)|^2 dt = \int_{\mathbb{R}} \left| \frac{\widehat{\psi}'(x)}{\pi} \right|^2 dx > \int_{\mathbb{R}} |\widehat{\psi}(x)|^2 dx
$$

=
$$
\int_{\mathbb{R}} |\psi(t)|^2 dt = \int_{\mathbb{R}} t^{-2} |\psi(1/t)|^2 dt = \int_{\mathbb{R}} t^2 |\varphi(t)|^2 dt.
$$

On the other hand, running the same argument as in step 1, we obtain that

$$
\int_{\mathbb{R}} t^2 |\psi(t)|^2 dt < \int_{\mathbb{R}} t^2 |\varphi(t)|^2 dt,
$$

an obvious contradiction, which stems from the fact that we supposed that $\psi \neq 0$, concluding the proof under the conditions of Theorem [2,](#page-4-0) since $\hat{\psi}, T\hat{\psi} \in H^1$ if, and only if, $\psi \in L^2((1+t^2) dt)$.

Step 3: Now, suppose we are in the $\alpha\beta = 1$ case. Through a dilation argument again, we may take $\alpha = \beta = 1$. By applying the proof above, we get that we must have equality in the Poincaré-Wirtinger inequality

$$
(a_{n+1} - a_n)^2 \int_{a_n}^{a_{n+1}} \left| \frac{\hat{\psi}'(x)}{\pi} \right|^2 dx \ge \int_{a_n}^{a_{n+1}} |\hat{\psi}(x)|^2 dx, \tag{4.10}
$$

for each $n \in \mathbb{Z}$. This promptly implies that there is $t_n \in \mathbb{C}$ such that $\widehat{\psi}(x) = t_n \cdot \sin\left(\pi \frac{x-a_n}{a_{n+1}-a_n}\right)$ $a_{n+1}-a_n$ \int for $x \in [a_n, a_{n+1}].$

Suppose thus that $\hat{\psi} \neq 0$. Then there is $n_0 \in \mathbb{Z}$ such that $t_{n_0} \neq 0$. We now focus on the next endpoint a_{n_0+1} . Since $\psi \in \mathcal{H}_2$, we have that $\hat{\psi} \in C^1(\mathbb{R})$, and hence $\hat{\psi}'(a_{n_0+1})$ is well-defined. But, since $\hat{\psi}(x) = t_{n_0+1} \sin\left(\pi \frac{x-a_{n_0+1}}{a_{n_0+2}-a_{n_0+1}}\right)$ for $x \in [a_{n_0+1}, a_{n_0+2}]$, by comparing the right and left limits

$$
\lim_{r \to 0} \frac{\widehat{\psi}(a_{n_0+1}+r) - \widehat{\psi}(a_{n_0+1})}{r} = \lim_{r \to 0} \frac{\widehat{\psi}(a_{n_0+1}-r) - \widehat{\psi}(a_{n_0+1})}{-r} = \widehat{\psi}'(a_{n_0+1}),
$$

we obtain that

$$
-\frac{t_{n_0}}{a_{n_0+1}-a_{n_0}}=\frac{t_{n_0+1}}{a_{n_0+2}-a_{n_0+1}}.
$$

This shows that $t_{n_0+1} \neq 0$. Since the same argument may be used for the endpoint a_{n_0} , we have $t_{n_0-1} \neq 0$, and hence $t_n \neq 0$ for all $n \in \mathbb{Z}$ by induction. Hence, $\hat{\psi} \neq 0$ on each of the intervals $[a_n, a_{n+1}]$. Looking back to the proof, we see that the inequality

$$
\int_{a_n}^{a_{n+1}} \left| \frac{\widehat{\psi}'(x)}{\pi} \right|^2 dx \ge (a_{n+1} - a_n)^2 \int_{a_n}^{a_{n+1}} \left| \frac{\widehat{\psi}'(x)}{\pi} \right|^2 dx
$$

must actually become an *equality* for each $n \in \mathbb{Z}$, and hence $a_{n+1} - a_n = 1$ whenever $\hat{\psi}' \neq 0$ in (a_n, a_{n+1}) . Repeating the endpoint derivative argument we used above for each $n \in \mathbb{Z}$ implies readily that

$$
-t_n=t_{n+1},\,\forall\,n\in\mathbb{Z},
$$

which, on the other hand, shows that $\hat{\psi}(x) = c \cdot \sin(\pi(x - \theta))$ for all $x \in \mathbb{R}$, for some $c \in \mathbb{C} \setminus \{0\}$ and some $\theta \in \mathbb{R}$. Since this last function is *not* a Fourier transform of an L^1 function by Riemann-Lebesgue, we arrive at a contradiction, which stems from supposing that $\psi \neq 0$, as desired. \square

Remark 4.1. It is important to note that same endpoint results may be obtained without making use of continuity of $\hat{\psi'}$, only using square-summability of ψ and a slightly different *regularity* assumption on ψ . Moreover, in the regular case where A is a translated copy of Z and $B = \mathbb{Z}$, we may obtain the same result once more solely under the assumption that $\psi \in \mathcal{H}_0$. We refer the reader to Section [5.3](#page-31-0) for a more detailed discussion on that topic.

5. Applications

5.1. An asymptotic version of Theorem [2.](#page-4-0) As promised in the introduction, the main goal of this section will be to prove an asymptotic version of Theorem [2.](#page-4-0) In order to do so, we define the following auxiliary class of functions:

$$
\mathcal{C}_{1/2} = \left\{ \psi \in \mathcal{H}_0 \colon \widehat{\psi} \in L^2(dt/|t|) \right\}.
$$
\n
$$
(5.1)
$$

The notation employed for this space is inspired by Proposition [2.1,](#page-7-0) which shows that the $L^2(dt/|t|)$ norm is bounded by an interpolation of L^2 and H^1 norms. Our theorem then reads as follows:

Theorem 6. Let $A = \{a_n\}_n$ and $B = \{b_n\}_n$ be two separated sequences. Then the following assertions hold:

(I) Suppose $0 \in A \cup B$. If we let

$$
\limsup_{n \to \pm \infty} |a_{n+1} - a_n| = \alpha, \limsup_{n \to \pm \infty} |b_{n+1} - b_n| = \beta,
$$

then $(\Gamma, \Lambda_{\mathbf{A}, \mathbf{B}})$ is a $\mathcal{C}_{1/2}$ -H.U.P. if $\alpha\beta < 1$.

(II) If we let

$$
\liminf_{n \to \pm \infty} |a_{n+1} - a_n| = \alpha, \liminf_{n \to \pm \infty} |b_{n+1} - b_n| = \beta,
$$

then $(\Gamma, \Lambda_{\mathbf{A}, \mathbf{B}})$ is not a $C_{1/2}$ -H.U.P. if $\alpha\beta > 1$.

Part (II) of the result above is a direct consequence of Theorem [3,](#page-5-0) after a suitable dilation. Hence, we focus on proving Part (I) in what follows.

5.1.1. A higher-dimensional Fourier uniqueness result. In order to prove Theorem [6,](#page-20-2) we need to use the techniques developed in [\[32\]](#page-35-18) in the context of Fourier uniqueness pairs. In particular, we will need a higher-dimensional version of their main Fourier uniqueness result.

Here, it is important to note in which sense we define vanishing: indeed, for $d = 1$, all functions in $H¹ \cap L²$ are automatically *continuous*, and hence pointwise evaluation is well-defined. On the other hand, in higher dimensions this is not the case; for that reason we will say that a function $f \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$ vanishes on a sphere of radius $r > 0$ if we have that

$$
\mathbb{T}_r(f) = 0,
$$

where $\mathbb{T}_r : W^{1,2}(B_r(0)) \to L^2(r\mathbb{S})$ denotes the Sobolev-trace operator for the ball of center 0 and radius $r > 0$.

Theorem 7. Let $f : \mathbb{R}^d \to \mathbb{R}$ be a radial function with $f \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$ vanishing on a set of centered spheres of radii $\{\lambda_i\}_i$, with $\mathcal{F}_d(f)$ being continuous and vanishing on centered spheres with $\text{radii } \{\gamma_i\}_i$, where the aforementioned sequences satisfy

$$
\limsup_{i \to \infty} \lambda_i^{p-1} |\lambda_{i+1} - \lambda_i| < \alpha,
$$
\n
$$
\limsup_{i \to \infty} \gamma_i^{q-1} |\gamma_{i+1} - \gamma_i| < \beta,
$$

with $1/p + 1/q = 1$, $p \ge 2$ and $\alpha^{1/p} \beta^{1/q} < 1/2$. Then $f \equiv 0$.

Since the $d = 1$ case has been treated in [\[32\]](#page-35-18), we suppose that $d \geq 2$ throughout the proof. We begin our discussion of the proof of this result with the following simple lemma:

Lemma 5.1. Let $t > 0$ and let $\varepsilon \geq 1 - \left(1 + \frac{1}{2t}\right)^{-\frac{d-1}{2}}$. Let $f : \mathbb{R}^d \to \mathbb{R}$ be a function such that $f \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$, supported outside of a centered ball of radius 1, which vanishes on a set of (centered) spheres with $(1 - \varepsilon)(2t)^{-1}$ -dense set of radii. Then, for all convex increasing C^1 -functions $\Phi : \mathbb{R}_+ \to \mathbb{R}_+$, we have

$$
\Phi(t^2) \int_{\mathbb{R}^d} |f(x)|^2 dx \le \int_{\mathbb{R}^d} \Phi(|\xi|^2) |\mathcal{F}_d(f)(\xi)|^2 d\xi.
$$

Proof. The proof of this result is similar to that of $[32, \text{Lemma 3}]$. We begin with the following:

Claim 5.1. Let $\mathbf{C}(a, b)$ be the least of all $C > 0$ such that, for the annulus $A(a, b) = \{x \in \mathbb{R}^d : a <$ $|x| < b$, we have

$$
||f||_{L^{2}(A(a,b))} \leq C||\nabla f||_{L^{2}(A(a,b))},
$$
\n(5.2)

for each $f \in W^{1,2}(A(a,b))$ with $f|_{\partial A(a,b)} = 0$, in the sense of Sobolev traces. Then there exists a function u_0 such that

$$
||u_0||_{L^2(A(a,b))} = \mathbf{C}(a,b)||\nabla u_0||_{L^2(A(a,b))}.
$$
\n(5.3)

Furthermore, we have that any such function u_0 is *radially symmetric*.

Proof of the Claim [5.2.](#page-21-0) The existence of such a function follows directly from the Rellich-Kondrachov theorem. For the radiality, note first that any function satisfying [\(5.3\)](#page-21-1) must satisfy the following Dirichlet-Laplace eigenvalue problem:

$$
\begin{cases}\n-\Delta u_0 = \lambda_1 u_0 & \text{in } A(a, b), \\
u_0 = 0 & \text{on } \partial A(a, b).\n\end{cases}
$$
\n(5.4)

Suppose now that u_0 is not radial, and consider, for a fixed rotation $R \in SO(d)$, the function

$$
u_0^R(x) = u_0(Rx).
$$

This new function is, by direct computation, still a solution of [\(5.4\)](#page-21-2). Since the first eigenfunction is unique, and since, by polar coordinates,

$$
\int_{A(a,b)} u_0 = \int_{A(a,b)} u_0^R
$$

,

it follows by uniqueness of the first eigenfunction that $u_0^R = u_0$. But this is the same as saying that any function u_0 satisfying [\(5.3\)](#page-21-1) must be *radial*, concluding the proof of the claim.

Our next claim deals with a relationship between Poincaré constants on an annulus and the real line interval associated with it:

Claim 5.2. Let $C(a, b)$ be as in Claim [5.1.](#page-21-3) Let also $C(a, b)$ denote the best constant in the inequality

$$
||f||_{L^{2}((a,b))} \leq C||f'||_{L^{2}((a,b))}.
$$
\n(5.5)

Then we have

$$
\mathbf{C}(a,b) \leq \left(\frac{b}{a}\right)^{\frac{d-1}{2}} \mathfrak{C}(a,b).
$$

Proof of Claim [5.2.](#page-22-0) By Claim [5.1,](#page-21-3) we have that (5.3) holds for u_0 . On the other hand, since u_0 is radial, we identify it with its restriction to the positive real line. Then, writing $\omega_d = \frac{\pi^{d/2}}{\Gamma(d/2+1)}$ to denote the volume of the unit ball in \mathbb{R}^d ,

$$
||u_0||_{L^2(A(a,b))}^2 = d \cdot \omega_d \cdot \int_a^b |u_0(r)|^2 r^{d-1} dr, \quad ||\nabla u_0||_{L^2(A(a,b))}^2 = d \cdot \omega_d \cdot \int_a^b |u'_0(r)|^2 r^{d-1} dr.
$$

Since $a^{d-1} \le r^{d-1} \le b^{d-1}$ holds in this case, then we have

$$
||u_0||_{L^2(A(a,b))}^2 \le d \cdot \omega_d b^{d-1} ||u_0||_{L^2(a,b)}^2
$$

$$
\le d \cdot \omega_d b^{d-1} \mathfrak{C}(a,b)^2 ||u'_0||_{L^2((a,b))}^2 \le (b/a)^{d-1} \mathfrak{C}(a,b)^2 ||\nabla u_0||_{L^2(A(a,b))}^2.
$$

By the definition of $\mathbf{C}(a, b)$, we conclude the desired assertion.

We now conclude in a similar way as in $[32, \text{ Lemma } 3]$: indeed, let f be as in the statement of Lemma [5.1.](#page-21-4) By the Poincaré inequality for annuli $A(r_i, r_{i+1})$, together with Claim [5.2,](#page-22-0) we obtain

$$
\int_{\mathbb{R}^d} |f(x)|^2 dx \le \left(\frac{r_{i+1}}{r_i}\right)^{d-1} \cdot \frac{(1-\varepsilon)^2}{(2\pi t)^2} \int_{\mathbb{R}^d} |\nabla f(x)|^2 dx.
$$

By Plancherel, we obtain

$$
\int_{\mathbb{R}^d} t^2 |f(x)|^2 dx \le (1 - \varepsilon)^2 \left(1 + \frac{1}{2t} \right)^{d-1} \int_{\mathbb{R}^d} |\xi|^2 |\mathcal{F}_d(f)(\xi)|^2 d\xi,
$$

since $r_i > 1$ and $r_{i+1} - r_i \leq (2t)^{-1}$. By the condition on ε , we prove the desired inequality with $\Phi(t) = t$. In order to prove for general Φ , one repeats the last step in the proof of Lemma 3 in [\[32\]](#page-35-18) verbatim. This finishes the proof.

The next ingredient is a version of the main result in [\[32,](#page-35-18) Section 5.2]. More specifically, we will show the following:

Proposition 5.1. Let $f : \mathbb{R}^d \to \mathbb{R}$ be a function with $f \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$ such that f vanishes on centered spheres with radii $\{\lambda_i\}_i$, and $\mathcal{F}_d(f)$ vanishes on centered spheres with radii $\{\gamma_i\}_i$, where

 $\{\lambda_i\}_i, \{\gamma_i\}_i$ are as in the statement of Theorem [7.](#page-21-5) Then we have that f is a Schwartz function. More specifically, f belongs to the Gelfand-Shilov class of functions satisfying

$$
\int_{\mathbb{R}^d} |f(x)|^2 e^{c|x|^p} dx, \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 e^{c|\xi|^q} d\xi < +\infty,
$$
\n(5.6)

for some $c > 0$.

Proof. Since the proof follows the same lines as that of [\[32,](#page-35-18) Section 5.2], we only indicate the parts where changes are needed. We let then $a = p - 1$, $b = q - 1$. By a rescaling argument we may also assume that $\alpha = \beta < 1/2$ from the conditions on λ_i and γ_i in Theorem [7.](#page-21-5)

We start by proving that, under the conditions of Proposition [5.1,](#page-22-1) we have $\mathcal{F}_d(f) \in L^2(\mathbb{R}^d) \cap$ $H^1(\mathbb{R}^d)$. First, let $F_0 \in C_c^{\infty}(\mathbb{R}^d)$ be a smooth function, with $\text{supp}(F_0) \subset B_1(0)$. Take $v \in \mathbb{R}^d$ sufficiently large and consider $f \cdot F_{0,v}$, where $F_{0,v}(x) = F_0(x - v)$. In order to apply Lemma [5.1](#page-21-4) to $f \cdot F_{0,v}$, we need to show that $f \cdot F_{0,v} \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$. Since we have that $\nabla (f \cdot F_{0,v}) =$ $(\nabla f) \cdot F_{0,v} + f \cdot (\nabla F_{0,v})$, and since $\nabla F_{0,v}$ is a bounded function with compact support, this last claim readily follows.

We then apply Lemma [5.1](#page-21-4) to $f \cdot F_{0,v}$ with $\Phi(t) = t$. For |v| sufficiently large, $f \cdot F_{0,v}$ vanishes on a set of spheres with radii are at least $(c(|v| - 1)^a)^{-1}$ -dense. This shows that

$$
\int_{\mathbb{R}^d} |f(x)F_{0,v}(x)|^2 |x|^{2a} dx \lesssim (|v| - 1)^{2a} \int_{\mathbb{R}^d} |f(x)F_{0,v}(x)|^2 dx
$$

$$
\lesssim \int_{\mathbb{R}^d} |\mathcal{F}_d(f) * \mathcal{F}_d(F_{0,v})(\xi)|^2 |\xi|^2 d\xi.
$$
 (5.7)

We then integrate both the left-hand and right-hand sides of (5.7) with respect to v such that $R > |v| > X_0$, with X_0 a fixed large constant. On the one hand, the integral of the left-hand side over such v is at least

$$
\int_{B_{R-1}(0)\setminus B_{X_0+1}(0)} |f(x)|^2 |x|^{2a} dx.
$$

On the other hand, the same integral applied to the right-hand side of [\(5.7\)](#page-23-0) is bounded, by Plancherel, by

$$
\int_{\mathbb{R}^d \times \mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\mathcal{F}_d(F_0)(\eta)|^2 |\xi + \eta|^2 d\xi d\eta.
$$
 (5.8)

Since

$$
\int_{\mathbb{R}^d} |\mathcal{F}_d(F_0)(\eta)|^2 \left(1 + \frac{|\eta|}{|\xi|}\right)^2 d\eta \le C
$$

for $|\xi|$ sufficiently large, we may conclude that (5.8) is bounded by

$$
C\int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 (1+|\xi|^2) dx < +\infty,
$$

for some $C > 0$. Hence, gathering all this information, we have that

$$
\int_{B_{R-1}(0)\setminus B_{X_0+1}(0)} |f(x)|^2 |x|^{2a} dx \le C(\|f\|_{L^2}^2 + \|f\|_{H^1}^2). \tag{5.9}
$$

We may then take $R \to \infty$ on the left-hand side of [\(5.9\)](#page-23-2) to obtain that

$$
\int_{\mathbb{R}^d} |f(x)|^2 |x|^{2a} dx < +\infty.
$$

Since $a = p - 1 \ge 1$, this shows that $\mathcal{F}_d(f) \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$, as desired.

We then note that the assertion that each f satisfying the statement of Theorem [7](#page-21-5) must belong to the Schwartz class can be proved by essentially simply following the proof above, now employing the same argument on both spatial and frequency sides, and replacing the weight used by a sequence of weights $\Phi_t^p(s)$ which agree with $|s|^p$ on a large set, are convex and grow linearly at infinity. We shall skip the details in that proof, and refer the reader to [\[32,](#page-35-18) Section 5.1] for further details.

For the proof of [\(5.6\)](#page-23-3), a slightly more precise argument is needed. With that in mind, let

$$
F = 1_{B_u} * \underbrace{1_{B_{u/2k}} * \cdots * 1_{B_{u/2k}}}_{k-\text{fold convolution}}
$$

where we define $\underline{1}_B(x) := \frac{1}{|B|} 1_B(x)$, and B_r denotes the Euclidean ball with center 0 and radius r. Note that, for a function f as in the statement of the proposition, there exists $X_0 > 1$ such that, if $|x| > X_0$, then $|x|$ belongs to an interval of length at most $(2\sigma|x|^{p-1})^{-1}$, where $\sigma > 1$ and the endpoint of such interval are radii of spheres where f vanishes. With that in mind, take v to be a vector in \mathbb{R}^d such that $|v| - \frac{3}{2}u > X_0$. Then the function $f \cdot F_v$, where $F_v(x) = F(x - v)$, satisfies the hypotheses of Lemma [5.1](#page-21-4) with $t = \frac{1+\sigma}{2}$ $\frac{\pm \sigma}{2}(|v| - \frac{3}{2}u)^{2a} = \tilde{\sigma}(|v| - \frac{3}{2}u)^{2a}$, since we can take $\varepsilon \gtrsim \sigma - 1$ there. Hence, using $\Phi(t) = t^{\theta}$, we obtain that

$$
\left(|v|-\frac{3}{2}u\right)^{2a\theta}\int_{\mathbb{R}^d}|f(x)|^2|F_v(x)|^2\,dx\leq \tilde{\sigma}^{-2\theta}\int_{\mathbb{R}^d}|\xi|^{2\theta}|(\mathcal{F}_d(f)*\mathcal{F}_d(F_v)(\xi)|^2\,d\xi.
$$

Again, we integrate both sides over $v \in \mathbb{R}^d$ such that $|v| > \frac{3}{2}$ $\frac{3}{2}u + X_0$. On the left-hand side, we bound it from below, in analogy to the one-dimensional case, by

$$
\left(\frac{K-4}{K}\right)^{2a\theta} \int_{|x| \ge K u} |x|^{2a\theta} |f(x)|^2 dx \cdot \left(\int_{\mathbb{R}^d} |F(y)|^2 dy\right),\tag{5.10}
$$

,

where $K > 4$ and $u \geq X_0$. For the right-hand side, we bound it from above by

$$
\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |(\mathcal{F}_d(f) * \mathcal{F}_d(F_v)(\xi)|^2 |\xi|^{2\theta} d\xi \right) dv = \int_{\mathbb{R}^d \times \mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\mathcal{F}_d(F)(\eta)|^2 |\xi + \eta|^{2\theta} d\xi d\eta. \tag{5.11}
$$

We now need to estimate the contribution on the integral above stemming from the Fourier transform of *F*. Effectively, the Fourier transform of $\frac{1}{B_1}$ is \mathcal{F}_d $\sqrt{ }$ $\frac{1}{B_1}$ $\Bigg) (\xi) = \frac{1}{\omega_d} \cdot |\xi|^{-d/2} J_{d/2}(2\pi |\xi|),$ where J_{ν} denotes the Bessel function of order ν . Hence, since $\frac{1}{B_r}(x) = r^{-d} \frac{1}{B_1}(x/r)$ for all $r > 0$,

$$
\mathcal{F}_d(F)(\eta) = u^{d/2} \frac{J_{d/2}(2\pi \cdot u|\eta|)}{|\eta|^{d/2}} \cdot \left(\frac{J_{d/2}(2\pi \cdot (u/2k) \cdot |\eta|)}{\omega_d \cdot |(u/2k) \cdot \eta|^{d/2}} \right)^k.
$$

We now claim that

$$
(|\xi| + |\eta|) \cdot \frac{|J_{d/2}(\pi \frac{u}{k} |\eta|)|}{\omega_d \cdot |\frac{u}{2k} \eta|^{d/2}} \leq |\xi| + \sqrt{\frac{2d+4}{\pi}} \frac{k}{u \pi}.
$$

Indeed, since \mathcal{F}_d $\sqrt{ }$ $\frac{1}{B_1}$ $\left|\int_0^{\infty}$ 1, the inequality is trivially true if $|\eta| \leq \sqrt{\frac{2d+4}{\pi}}$ π k $\frac{k}{u\pi}$. If, on the other hand, $|\eta| > \sqrt{\frac{2d+4}{\pi}}$ π k $\frac{\kappa}{u\pi}$, we write the Bessel function $J_{d/2}(t)$ as

$$
J_{d/2}(t) = \frac{(d-1) \cdot (\frac{1}{2}t)^{\frac{d}{2}-1}}{\sqrt{\pi} \Gamma(\frac{d+1}{2})} \int_0^{\frac{1}{2}\pi} \sin(t \cos \theta) \sin^{d-2} \theta \cos \theta d\theta;
$$

see, for instance, [\[47,](#page-36-10) (7), §III.3.3, p. 48]. From this formula, it follows that

$$
2^{d/2}\Gamma\left(\frac{d}{2}+1\right)\frac{|J_{d/2}(t)|}{t^{d/2-1}} \le \sqrt{\frac{2d+4}{\pi}},\tag{5.12}
$$

upon using Gautschi's inequality [\[11,](#page-35-20) (5.6.4)], which asserts that $\Gamma(x+1) \leq (x+1)^{1-s} \cdot \Gamma(x+s)$ for all $x \geq 0$, $s \in (0,1)$, and the fact that

$$
(d-1)\left|\int_0^{\frac{\pi}{2}} \sin(t\cos\theta)\sin^{d-2}\theta\cos\theta\,d\theta\right| \leq \int_0^{\frac{\pi}{2}} |\sin^{d-1}|'(\theta)|\,d\theta = 1.
$$

It follows hence from [\(5.12\)](#page-25-0) that

$$
(|\xi| + |\eta|) \cdot \frac{|J_{d/2}(\pi \frac{u}{k} |\eta|)|}{\omega_d \cdot |\frac{u}{2k} \eta|^{d/2}} \le |\xi| + |\eta| \cdot \left(2^{d/2} \Gamma\left(\frac{d}{2} + 1\right) \frac{|J_{d/2}(\pi \frac{u}{k} |\eta|)|}{|\pi \frac{u}{k} \eta|^{d/2}} \right)
$$

$$
\le |\xi| + \sqrt{\frac{2d + 4}{\pi}} \frac{k}{u\pi}.
$$

Therefore, using that the Fourier transform of the (normalized) unit ball in dimension d is uniformly bounded by 1, for $k \ge \theta$, we conclude that the right-hand side of (5.11) is bounded by

$$
\int_{\mathbb{R}^d \times \mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 \left(|\xi| + \sqrt{\frac{2d+4}{\pi}} \frac{k}{u\pi} \right)^{2\theta} |\mathcal{F}_d(\mathbf{1}_{B_u})(\eta)|^2 d\xi d\eta
$$

$$
= \omega_d \cdot u^d \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 \left(|\xi| + \sqrt{\frac{2d+4}{\pi}} \frac{k}{u\pi} \right)^{2\theta} d\xi.
$$

On the other hand, taking into account that $\int_{\mathbb{R}^d} |F(y)|^2 dy \geq \omega_d \cdot 2^{-d} u^d$ – since $F \equiv 1$ in $B_{u/2}$ – in [\(5.10\)](#page-24-1), we obtain the improved lower bound

$$
\omega_d \cdot 2^{-d} u^d \left(\frac{K-4}{K}\right)^{2a\theta} \int_{|x| \ge K u} |x|^{2a\theta} |f(x)|^2 dx.
$$
 (5.13)

Putting all these considerations together, we obtain

$$
\left(\frac{K-4}{K}\right)^{2a\theta} \int_{|x|\geq Ku} |x|^{2a\theta} |f(x)|^2 dx \leq 2^d \tilde{\sigma}^{-2\theta} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 \left(|\xi| + \sqrt{\frac{2d+4}{\pi}} \frac{k}{u\pi}\right)^{2\theta} d\xi. \tag{5.14}
$$

At this point, we can just repeat the last part of the proof of the one-dimensional case verbatim. Effectively, we rewrite (5.14) as

$$
\int_{|x|\geq K u} |x|^{2a\theta} |f(x)|^2 dx \leq 2^d \left(\frac{1}{\tilde{\sigma}} \left(\frac{K}{K-4}\right)^a\right)^{2\theta} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 \left(|\xi| + \sqrt{\frac{2d+4}{\pi}} \frac{k}{u\pi}\right)^{2\theta} d\xi. \tag{5.15}
$$

We choose K large enough so that $\frac{1}{\tilde{\sigma}}$ $\left(K \right)$ $K-4$ a^{a} < 1, take $k \in [\theta, \theta \pi]$, and bound the integral on the right-hand side of [\(5.15\)](#page-25-2) by

$$
\left(\frac{K+1}{K}\right)^{2\theta} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\xi|^{2\theta} d\xi + \int_{|\xi| \le K\sqrt{\frac{2d+4}{\pi} \frac{k}{u\pi}}} |\mathcal{F}_d(f)(\xi)|^2 \left(|\xi| + \sqrt{\frac{2d+4}{\pi} \frac{k}{u\pi}}\right)^{2\theta} d\xi
$$

$$
\le \left(\frac{K+1}{K}\right)^{2\theta} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\xi|^{2\theta} d\xi + \left(\sqrt{\frac{2d+4}{\pi}}(K+1)\right)^{2\theta} \left(\frac{\theta}{u}\right)^{2\theta} \int_{\mathbb{R}^d} |f(x)|^2 dx.
$$

Since

$$
\int_{|x| \le K u} |f(x)|^2 |x|^{2a\theta} dx \le K^{2a\theta} u^{2a\theta} \int_{\mathbb{R}^d} |f(x)|^2 dx,
$$

inserting these considerations into [\(5.15\)](#page-25-2) we obtain that

$$
\int_{\mathbb{R}^d} |x|^{2a\theta} |f(x)|^2 dx \le 2^d \left(\frac{1}{\tilde{\sigma}} \left(\frac{K}{K-4}\right)^a \cdot \frac{K+1}{K}\right)^{2\theta} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\xi|^{2\theta} d\xi
$$

$$
+ \left(K^{2a\theta} u^{2a\theta} + 2^{d+1} \left(\sqrt{\frac{2d+4}{\pi}} (K+1)\right)^{2\theta} \left(\frac{\theta}{u}\right)^{2\theta}\right) \int_{\mathbb{R}^d} |f(x)|^2 dx. (5.16)
$$

Using now that the factor in front of the first integral on the right-hand side above converges to zero as $\theta \to \infty$, and taking $u = \theta^{1/(a+1)}$, which is allowed since X_0 is fixed, we get

$$
\int_{\mathbb{R}^d} |f(x)|^2 |x|^{2a\theta} \, dx \le \frac{1}{2} \int_{\mathbb{R}^d} |\mathcal{F}_d(f)(\xi)|^2 |\xi|^{2\theta} \, d\xi + C_d^{2\theta} \theta^{\frac{2a}{a+1}\theta} \int_{\mathbb{R}^d} |f(x)|^2 \, dx. \tag{5.17}
$$

Now, notice that the exact same can be done if we switch the roles of f and \hat{f} . By doing so, if we let $\kappa = a\theta$, we obtain

$$
\int_{\mathbb{R}^d} |x|^{2a\theta} |f(x)|^2 dx + \int_{\mathbb{R}^d} |\xi|^{2b\kappa} |\mathcal{F}_d(f)(\xi)|^2 d\xi \leq C_d^{\theta} \theta^{\frac{2a}{a+1}\theta} \int_{\mathbb{R}^d} |f(x)|^2 dx.
$$

Calling $\theta = \frac{a+1}{2a}$ $\frac{p+1}{2a}\ell$, where $\ell \in \mathbb{Z}_+$, and reverting back $a = p - 1$, $b = q - 1$, we obtain

$$
\int_{\mathbb{R}^d} |x|^{p\ell} |f(x)|^2 dx + \int_{\mathbb{R}^d} |\xi|^{q\ell} |\mathcal{F}_d(f)(\xi)|^2 d\xi \leq C_d^{\ell} \ell! \int_{\mathbb{R}^d} |f(x)|^2 dx.
$$

Dividing both sides by $(2C_d)^{\ell} \ell!$ and summing over all $\ell \geq 0$ then yields our claim.

Proof of Theorem [7.](#page-21-5) From Proposition [5.1,](#page-22-1) we wish to conclude, in analogy to [\[32,](#page-35-18) Section 5.3] and using the methods from [\[39,](#page-36-11) Section 5.2.3, Step 2], that f and $\mathcal{F}_d(f)$ may be extended—when identified with their radial representation—as entire functions of order p and q , respectively.

Indeed, it readily follows from the techniques from [\[32\]](#page-35-18) that for some $c' > 0$, one has $|f(x)| \lesssim$ $e^{-c'|x|^p}, |\mathcal{F}_d(f)(\xi)| \lesssim e^{-c'|\xi|^q}$, for each $x, \xi \in \mathbb{R}^d$. In order to conclude the assertion on analyticity, we follow the overall structure already present in [\[32,](#page-35-18) Section 5.3]. As a matter of fact, our first observation is that, since we can represent

$$
\mathcal{F}_d(f)(\xi) = (2\pi)^{d/2} \int_0^\infty f(r) \cdot r^{d-1} \mathcal{A}_{\frac{d-2}{2}}(|\xi|r) \, dr,\tag{5.18}
$$

where $A_{\nu}(s) = (2\pi s)^{\nu} J_{\nu}(2\pi s)$, and since A_{ν} is an even analytic function of its argument, which satisfies, when extended to the whole complex plane,

$$
|\mathcal{A}_{\nu}(\xi + i\eta)| \le C_d e^{2\pi |\eta|},\tag{5.19}
$$

the decay conditions from $f, \mathcal{F}_d(f)$ above imply that $\mathcal{F}_d(f)$ and f can be extended (when regarded as one-variable functions) to the whole complex plane as entire functions of orders p and q , respectively. With this in mind, we let

$$
\kappa_1(\theta) = \frac{1}{2\pi} \limsup \frac{\log |f(re^{i\theta})|}{r^p},
$$

$$
\kappa_2(\theta) = \frac{1}{2\pi} \limsup \frac{\log |\mathcal{F}_d(f)(\rho e^{i\theta})|}{\rho^q}
$$

denote the Phragmén-Lindelöff indicators of f and $\mathcal{F}_d(f)$, respectively. We then denote $K_i = |\kappa_i(0)|$. For these indicators, the following Claim—which is a version of [\[32,](#page-35-18) Claim 2]—holds:

Claim 5.3. We have

$$
\kappa_1(\theta) \le \frac{1}{p(qK_2)^{p/q}} |\sin \theta|^p,
$$

\n
$$
\kappa_2(\varphi) \le \frac{1}{q(pK_1)^{q/p}} |\sin \varphi|^q,
$$
\n(5.20)

for all $\theta, \varphi \in [0, 2\pi)$.

Proof. The only difference between the proof of this result and that of [\[32,](#page-35-18) Claim 2] is that the current one must resort to [\(5.18\)](#page-26-0) and [\(5.19\)](#page-27-0). Once these two formulas are applied, the rest of the proof in [\[32,](#page-35-18) Claim 2] can be repeated verbatim, and hence we omit that part. \square

We then state the following crucial claim:

Claim 5.4. With the definitions above, we have that, for $\theta \in \left(0, \frac{\pi}{2} \right)$ $_{2p}$ $\big)$,

$$
K_1 \ge \frac{1}{p} \tan(p\theta) - \frac{1}{p(qK_2)^{p/q}} \frac{(\sin \theta)^p}{\cos(p\theta)}
$$
(5.21)

and, for $\varphi \in \left(0, \frac{\pi}{2\epsilon}\right)$ $\overline{2q}$ $\big)$,

$$
K_2 \ge \frac{1}{q} \tan(q\varphi) - \frac{1}{q(pK_1)^{q/p}} \frac{(\sin \varphi)^q}{\cos(q\varphi)}.
$$
\n
$$
(5.22)
$$

Proof of Claim [5.4.](#page-27-1) The proof is a verbatim adaptation of [\[32,](#page-35-18) Equation $(5.3.1)$], and hence we omit it.

In order to conclude the proof of Theorem [7,](#page-21-5) we use the inequality $\frac{\tan p\theta}{p} > \frac{\sin \theta}{(\cos p\theta)}$ $\frac{\sin \theta}{(\cos p\theta)^{1/p}}, \theta \in (0, \pi/2p),$ in the statement of Claim [5.4.](#page-27-1) Letting $X = \frac{\sin \theta}{(\cos \theta)}$ $\frac{\sin \theta}{(\cos p\theta)^{1/p}}$, we get from these considerations that

$$
K_1 > X - \frac{1}{(qK_2)^{p/q}} \frac{X^p}{p}, \, \forall X > 0.
$$

A direct optimization argument concludes that the maximum of the right-hand side of the expression above is at $X = qK_2$, for which we obtain $K_1 > K_2$. On the other hand, we may exchange the roles of f and $\mathcal{F}_d(f)$ in the proof above, which allows us to obtain $K_2 > K_1$, which is an obvious contradiction. This contradiction arises when we suppose that $f \neq 0$, which finishes the proof. \Box Remark 5.1. As it turns out, Theorem [7](#page-21-5) has consequences for arbitrary functions vanishing on spherical shells, rather than only radial ones, as explicitly stated in the next result.

Corollary 5.1. Let $f, \mathcal{F}_d(f) \in L^2(\mathbb{R}^d) \cap H^1(\mathbb{R}^d)$. Suppose, moreover, that

$$
f|_{\lambda_j \cdot \mathbb{S}^{d-1}} = \mathcal{F}_d(f)|_{\gamma_j \cdot \mathbb{S}^{d-1}} \equiv 0, \ \forall j \ge 1,
$$

where $\{\lambda_j\}_j, \{\gamma_j\}_j$ are as in Theorem [7.](#page-21-5) Then $f \equiv 0$.

The proof of that result follows from using Proposition [5.1,](#page-22-1) which ensures that $f \in \mathcal{S}(\mathbb{R}^d)$ under the hypotheses of Corollary [5.1,](#page-28-0) together with Theorem [7](#page-21-5) in conjunction with the following result, originally proved by M. Stoller [\[43\]](#page-36-12):

Theorem (Corollary 2.2 in [\[43\]](#page-36-12)). Fix a dimension $d > 2$ and fix two subsets R, $\hat{R} \subset (0,\infty)$. Suppose that for all $p \in \{d + 2m : m \in \mathbb{N}_0\}$ an all radial Schwartz functions f defined on \mathbb{R}^p , the following implication holds:

$$
\left(f|_{\bigcup_{r\in R}rS^{p-1}}=0 \text{ and } \mathcal{F}_d(f)|_{\bigcup_{\rho\in \hat{R}}\rho S^{p-1}}=0\right)\Longrightarrow f=0.
$$

Then the same implication holds for arbitrary $f \in \mathcal{S}(\mathbb{R}^d)$.

5.1.2. Proof of Theorem [6.](#page-20-2) We are now able to move on to the proof of Part (I) of Theorem [6:](#page-20-2)

Proof of Theorem [2,](#page-4-0) Part (I). Suppose we have a $\psi \in C_{1/2}$ satisfying the conditions in Part (I) of Theorem [2.](#page-4-0) Let $\Theta_+(u) = \frac{\theta(|u|^2)}{|u|^2}$ $\frac{(|u|)}{|u|^2}$. By the reductions in Section [2.3,](#page-8-3) and by [\(2.7\)](#page-9-2), the following estimate holds:

$$
\int_{\mathbb{R}^4} |\Theta_+(u)|^2 (1+|u|^2) du = c \cdot \int_0^\infty \frac{|\theta(s^2)|^2}{s^4} (1+s^2)s^3 ds = \frac{c}{2} \int_{\mathbb{R}_+} |\theta(t)|^2 \frac{1+t}{t} dt
$$

$$
\lesssim \|\theta\|_{L^2}^2 + \|\theta\|_{L^2(dt/|t|)}^2.
$$
 (5.23)

Let $A_+ = A \cap (0, +\infty)$, $B_- = B \cap (-\infty, 0)$. Since $\Theta_+(\sqrt{a}) = \mathcal{F}_4(\Theta_+)(\sqrt{-b}) = 0$, for all $a \in A_+$, $b \in$ B−, and since

$$
\limsup_{n \to \infty} \sqrt{a_{n+1}} |\sqrt{a_{n+1}} - \sqrt{a_n}| \le \frac{1}{2} \limsup_{n \to \infty} |a_{n+1} - a_n| < \frac{\alpha}{2},
$$

and, by the same token,

$$
\limsup_{n \to -\infty} \sqrt{-b_n} |\sqrt{-b_{n+1}} - \sqrt{-b_n}| < \frac{\beta}{2},
$$

the sequences of radii $\gamma_i = \sqrt{a_i}$, for $a_i > 0$, $\lambda_i = \sqrt{-b_i}$ for $b_i < 0$ satisfy the hypotheses of Theorem [7](#page-21-5) with $p = q = 2$. Moreover, since Θ_+ is a continuous function by hypothesis, we are in a perfect position in order to apply Theorem [7.](#page-21-5) It then plainly implies that $\Theta_+ \equiv 0$. Since the same can be done for the negative part of θ , we have $\theta \equiv 0$, finishing the proof.

Remark 5.2. As mentioned after the statement of Theorem [6,](#page-20-2) the definition of the class $\mathcal{C}_{1/2}$ is inspired by Proposition [2.1:](#page-7-0) indeed, if one assumes the stronger property that $\theta \in L^2 \cap H^1$, then it follows directly that $T\theta \in L^2 \cap H^1$ by Proposition [2.4.](#page-10-0) By that same result and Proposition [2.1,](#page-7-0) we have promptly that $T\theta, \theta \in L^2 \cap L^2(dt/|t|)$, which is what we end up using in the proof of Part (I) of Theorem [6.](#page-20-2)

This highlights the fact that the main results in this manuscript deal with many classes of functions that are different in comparison to the ones in the previous literature. Indeed, the pioneering article $[27]$ and subsequent works such as $[28]$ all deal with $\psi \in L^1(\mathbb{R})$, which automatically implies that $\theta = \hat{\psi}$ and $T\theta$ are both continuous functions. Our results here, however, assume only $\psi \in L^2$ in terms of integrability, and thus the condition $\hat{\psi} \in C(\mathbb{R})$ is needed in order to make sense of pointwise values of θ .

To that extent, we remark that there is a vast array of functions $\psi \notin L^1(\mathbb{R})$ satisfying the conditions above: fix for instance any $f_0 \in C(\mathbb{R})$ 2-periodic such that its sequence of Fourier series coefficients is *not* in $\ell^1(\mathbb{Z})$. A simple example of such a function is given by

$$
f_0(x) = \sum_{n=1}^{\infty} \frac{1}{n^2} \sin \left((2^{n^3} + 1)\pi |x| \right), \qquad x \in [-1, 1],
$$

extended 2-periodically, as first observed by F \acute{e} jer [\[36,](#page-36-13) [44\]](#page-36-14). Then consider

$$
\psi(x) = \sum_{n \in \mathbb{Z}} \hat{f}_0(n) \cdot e^{\pi i (x-n)} 1_{[n-1,n+1]}(x).
$$

By Plancherel's theorem, it follows easily that $\psi \in L^2(\mathbb{R})$, but $\psi \notin L^1(\mathbb{R})$. Moreover, we have that

$$
\widehat{\psi}(\xi) = 2\left(\sum_{n \in \mathbb{Z}} \widehat{f}_0(n) e^{\pi i n \xi}\right) \cdot \frac{\sin(\pi(\xi - 1))}{\pi(\xi - 1)} = 2 \cdot f_0(\xi) \cdot \frac{\sin(\pi(\xi - 1))}{\pi(\xi - 1)}.
$$

By construction, we have that $\hat{\psi}$ is continuous and vanishes at infinity, as desired. Moreover, this construction incidentally yields a function in the class $C_{1/2}$ which is not in $L¹$. Indeed, we have $|\hat{\psi}(\xi)| \leq |\xi|$ for small ξ , and hence $\hat{\psi} \in L^2(d\xi/|\xi|)$ as a consequence, which further emphasizes the fundamental difference between the results presented here and the ones in [\[27,](#page-35-0) [28,](#page-35-13) [29,](#page-35-14) [7\]](#page-34-4) and other related work.

5.2. One-sided information on space and frequency. Although we stated Theorem [2](#page-4-0) as a recovery result for perturbed lattice crosses, we can actually obtain perturbed versions of the *one*sided results from [\[28\]](#page-35-13). In order to state those, we will need a bit of additional notation.

Definition 5.1. For a set $\Lambda \subset \mathbb{R}^2$, we say that $\psi \in \mathcal{H}_{\ell}$ belongs to the class $\mathcal{H}_{\ell}(\Lambda)$ if, for any $\xi = (\xi_1, \xi_2) \in \Lambda$, we have

$$
E\psi(\xi_1, \xi_2) = \int_{\mathbb{R}} \psi(t)e^{\pi i(t\xi_1 + \xi_2/t)} dt = 0.
$$
 (5.24)

For that class, we define in analogy to [\[28,](#page-35-13) Definition 1.2.1] the \mathcal{H}_{ℓ} -Zariski closure of Λ as

$$
\mathrm{z clos}^{\ell}_{2}(\Lambda) = \left\{ \xi \in \mathbb{R}^{2} \colon \forall \, \psi \in \mathcal{H}_{\ell}(\Lambda), E\psi(\xi) = 0 \right\}.
$$

We are now able to state our next result. Recall that, for $A, B \subset \mathbb{R}$, we define $\Lambda_{A,B} = (A \times \{0\}) \cup$ $({0} \times B)$.

Theorem 8. Let $A \subset \mathbb{R}_{\geq 0}, B \subset \mathbb{R}_{\leq 0}$ be two uniformly discrete sets. Then:

(I) Suppose that

$$
\sup_{n} |a_{n+1} - a_n| = \alpha, \ \sup_{n} |b_{n+1} - b_n| = \beta,
$$

with $\alpha\beta$ < 1. Then $z\cos_2^0(\Lambda_{A,B}) = \mathbb{R}_{\geq 0} \times \mathbb{R}_{\leq 0}$. Moreover, if $\alpha\beta = 1$, $z\cos_2^2(\Lambda_{A,B}) =$ $\mathbb{R}_{\geq 0} \times \mathbb{R}_{\leq 0}$ as well.

(II) Suppose that

$$
\inf_{n} |a_{n+1} - a_n| = \alpha, \inf_{n} |b_{n+1} - b_n| = \beta,
$$

with $\alpha\beta > 1$. Then $\mathbb{R}_{\geq 0} \times \mathbb{R}_{\leq 0} \setminus z \text{clos}_2^2(\Lambda_{\mathbf{A}, \mathbf{B}})$ contains infinitely many points.

Proof. We start with the case $\alpha\beta < 1$ in Part (I). Without loss of generality, $\alpha = \beta < 1$. Let $\psi \in \mathcal{H}_0$ be a function such that

$$
E\psi(a_n, 0) = E\psi(0, b_k) = 0, \forall n, k \in \mathbb{N}.
$$
\n(5.25)

The crucial observation we shall use throughout is that we may replace ψ by $\tilde{\psi}(t) = \frac{\psi - i \mathbf{H} \psi}{2}$ so that [\(5.25\)](#page-30-0) still holds for $\tilde{\psi}$, where $\mathbf{H}\psi(t)$ denotes the *Hilbert transform* of ψ at t, defined as

$$
\mathbf{H}\psi(t) = \frac{1}{\pi} \mathrm{p.v.} \int_{\mathbb{R}} \frac{\psi(x)}{t - x} dx.
$$
 (5.26)

Indeed, note first of all that $\tilde{\psi} \in L^2$. Moreover, if we simply use the property that

$$
\widehat{\mathbf{H}\psi}(\xi) = -i\operatorname{sgn}(\xi)\widehat{\psi}(\xi),
$$

we obtain that

$$
E\tilde{\psi}(a_n,0) = \frac{1}{2} \left(\int_{\mathbb{R}} \psi(t) e^{-\pi ita_n} - i \int_{\mathbb{R}} \mathbf{H} \psi(t) e^{-\pi ita_n} dt \right) = \theta(a_n) = 0, \forall n \ge 0.
$$

For the second equality, we need the following observation:

$$
\mathbf{H}\psi(1/t) = \frac{1}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{\psi(x)}{\frac{1}{t} - x} dx = \frac{t}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{\psi(x)}{1 - t \cdot x} dx \n= \frac{t^2}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{\psi(x) \cdot x}{1 - t \cdot x} dx + \frac{t}{\pi} \int \psi = \frac{t^2}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{\psi(1/s) \cdot (1/s)}{1 - \frac{t}{s}} \frac{ds}{s^2} \n= -\frac{t^2}{\pi} \text{p.v.} \int_{\mathbb{R}} \left(\frac{1}{s^2} \psi\left(\frac{1}{s}\right)\right) \frac{ds}{t - s} = -t^2 \mathbf{H}\varphi(t),
$$
\n(5.27)

where $\varphi(s) = \psi(1/s) \cdot (1/s^2)$, and where we used that $\int \psi = 0$. Hence,

$$
\int_{\mathbb{R}} \frac{1}{t^2} \mathbf{H} \psi(1/t) e^{-\pi i b_n t} dt = -\int_{\mathbb{R}} \mathbf{H} \varphi(t) e^{-\pi i b_n t} dt,
$$

and thus we obtain $\widehat{H\varphi} \in C(\mathbb{R} \setminus \{0\})$, and also

$$
E\tilde{\psi}(0,b_n) = \frac{1}{2} \left(\int_{\mathbb{R}} \varphi(t) e^{-\pi i b_n t} dt + i \int_{\mathbb{R}} \mathbf{H} \varphi(t) e^{-\pi i b_n t} dt \right) = T\theta(b_n) = 0, \forall n \ge 0,
$$

again by the properties of the Hilbert transform, as $b_n < 0$ for all n. Hence, we may suppose that $\tilde{\psi}$ is of the form above in what follows, and as a consequence we may always assume that $\hat{\psi}$ is supported on $\mathbb{R}_{\geq 0}$.

The rest of the proof is similar to that of Theorem [2:](#page-4-0) we start by once more observing that the Poincaré-Wirtinger argument of [\(4.8\)](#page-18-3) yields again that $\hat{\varphi} \in L^2(\mathbb{R})$, and hence $\psi \in L^2(x^2 dx)$. Applying now the same argument as in [\(4.9\)](#page-19-0), we have

$$
\int_{\mathbb{R}} |x|^2 |\psi(x)|^2 \, dx = \int_{\mathbb{R}} \left| \frac{\hat{\psi}'(x)}{\pi} \right|^2 \, dx = \int_0^\infty \left| \frac{\hat{\psi}'(x)}{\pi} \right|^2 \, dx \ge \int_0^\infty |\hat{\psi}(x)|^2 \, dx = \int_{\mathbb{R}} |\psi(x)|^2 \, dx. \tag{5.28}
$$

On the other hand, the same argument can be applied to φ by [\(5.27\)](#page-30-1): indeed, we have that $\hat{\varphi} = T\theta$. Since if supp $(\theta) \subset \mathbb{R}_{\geq 0}$, we have by Proposition [2.4](#page-10-0) that supp $(T\theta) \subset \mathbb{R}_{\leq 0}$, the same Poincaré-Wirtinger argument works to show that

$$
\int_{\mathbb{R}} |x|^2 |\varphi(x)|^2 dx \ge \int_{\mathbb{R}} |\varphi(x)|^2 dx.
$$
\n(5.29)

We then must have equality in the inequalities (5.28) and (5.29) , which implies equality in (4.9) . If $\alpha = \beta < 1$, this is only possible if $\psi \equiv 0$ whenever ψ has Fourier support on $\mathbb{R}_{\geq 0}$. Equivalently, undoing the change $\psi \mapsto \tilde{\psi}$, we have that

$$
\int_{\mathbb{R}} \psi(t) e^{\pi i \eta/t} dt = \int_{\mathbb{R}} \psi(t) e^{-\pi i t \xi} dt = 0, \forall \xi, \eta \ge 0.
$$

By [\[28,](#page-35-13) Proposition 1.7.2], we conclude that

$$
\int_{\mathbb{R}} \psi(t) e^{\pi i (\eta/t - \xi t)} dt = 0, \forall \xi, \eta \ge 0,
$$

which is the desired claim in that case. For the $\alpha\beta = 1$ case, we repeat the analysis of Section [4.2](#page-18-0) for the $\alpha\beta = 1$ case verbatim for $\tilde{\psi}$, which again allows us to conclude that $\psi - iH\psi \equiv 0 \equiv \varphi + iH\varphi$. By the considerations above, this implies the desired result also in that result.

Now, for Part (II) of Theorem [8,](#page-29-1) we simply note that the functions ψ_{\pm} built in Section [3.4](#page-15-2) satisfy exactly our assumptions, where we extend the sets A and B to the whole real line so that they satisfy the same uniform discreteness hypotheses. Hence, since $\psi_{\pm} \neq 0$, it follows that the inclusion zclos₂($\Lambda_{\mathbf{A},\mathbf{B}}$) $\subset \mathbb{R}_{\geq 0} \times \mathbb{R}_{\leq 0}$ is strict, and since the functions ψ_{\pm} are not identically zero, the complement of $z\text{clos}_2(\Lambda_{\mathbf{A},\mathbf{B}})$ in $\mathbb{R}_{\geq 0} \times \mathbb{R}_{\leq 0}$ must necessarily contain infinitely many points, as desired. \Box

Remark 5.3. The argument above shows that the exact same result as Theorem [8](#page-29-1) holds if $A \subset$ $\mathbb{R}_{\leq 0}, B \subset \mathbb{R}_{\geq 0}$, under the same hypotheses on the distribution of the difference of consecutive terms.

5.3. The critical case of Theorem [2.](#page-4-0) We now explain a different way to obtain the critical case $\alpha\beta = 1$ of Part (I) of Theorem [2,](#page-4-0) under several different sets of assumptions. We formulate it as a result of its own, which includes in its statement Theorem [4.](#page-6-1)

Theorem 9. Let A and B be as in Theorem [2.](#page-4-0) The following assertions hold:

(I) Let

$$
\mathcal{C} = \left\{ \psi \in \mathcal{H}_0(\mathbb{R}) \colon \int_{\mathbb{R}} |\widehat{\psi}(x)| |x|^{1/4} dx < +\infty \right\}.
$$

Then, if $\alpha\beta = 1$, ($\Gamma, \Lambda_{\mathbf{A}, \mathbf{B}}$) is a C-H.U.P.

(II) If $A = {\alpha n + \theta}_{n \in \mathbb{Z}}$ and $B = {\beta n}_{n \in \mathbb{Z}}$, then $(\Gamma, \Lambda_{A,B})$ is a \mathcal{H}_0 -H.U.P. if, and only if, $\alpha \beta \leq 1$.

Proof. Effectively, by the same argument as in the proof in Section [4,](#page-17-0) we are able to conclude first that, in both cases, $\psi \in L^2((1+x^2)dx)$. Furthermore, we also have that $a_{n+1} - a_n = 1$ whenever $\hat{\psi} \neq 0$ on $[a_n, a_{n+1}]$. In that case, we may conclude that

$$
\widehat{\psi}(x) = t_n \sin\left(\pi \frac{x - a_n}{a_{n+1} - a_n}\right) = t_n \sin(\pi(x - a_n)) \text{ for } x \in [a_n, a_{n+1}).
$$

That is,

$$
\hat{\psi}(x) = \sum_{n \in \mathbb{Z}} t_n \sin(\pi(x - a_n)) 1_{[a_n, a_{n+1}]}. \tag{5.30}
$$

Taking inverse Fourier transforms on both sides, we obtain

$$
\psi(t) = \sum_{n \in \mathbb{Z}} t_n e^{\pi i t (a_n + 1/2)} \left(\cos(\pi(\cdot)) \widetilde{1}_{(-1/2, 1/2)} \right)(t).
$$

Since we may compute

$$
\int_{-1/2}^{1/2} \cos(\pi x) e^{\pi i x t} dx = \frac{1}{2i\pi} \left(\frac{e^{\frac{\pi}{2}i(t+1)} - e^{-\frac{\pi}{2}i(t+1)}}{t+1} + \frac{e^{\frac{\pi}{2}i(t-1)} - e^{-\frac{\pi}{2}i(t-1)}}{t-1} \right)
$$

= $\frac{1}{\pi} \left(\frac{\sin(\frac{\pi}{2}(t+1))}{t+1} + \frac{\sin(\frac{\pi}{2}(t-1))}{t-1} \right)$
= $\frac{1}{\pi} \cos(\pi t/2) \left(\frac{1}{t+1} - \frac{1}{t-1} \right) = -\frac{2}{\pi} \frac{\cos(\pi t/2)}{t^2 - 1},$

we obtain that

$$
\psi(t) = -\frac{1}{\sqrt{2}\pi(t^2 - 1)} \sum_{n \in \mathbb{Z}} t_n \left(e^{\pi i t a_{n+1}} + e^{\pi i t a_n} \right). \tag{5.31}
$$

Since $\psi \in L^2(\mathbb{R})$, [\(5.30\)](#page-32-0) shows that $\{t_n\}_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$. By the exact same argument applied to $\varphi(t) = t^{-2}\psi(1/t)$, we obtain that also

$$
t^{-2}\psi(1/t) = -\frac{1}{\sqrt{2}\pi(t^2 - 1)}\sum_{n \in \mathbb{Z}} r_n \left(e^{\pi i t b_{n+1}} + e^{\pi i t b_n}\right),\tag{5.32}
$$

with $\{r_n\}_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$. By combining [\(5.31\)](#page-32-1) and [\(5.32\)](#page-32-2), we get that

$$
\sum_{n\in\mathbb{Z}} t_n \left(e^{\pi ita_{n+1}} + e^{\pi ita_n}\right) = -\sum_{n\in\mathbb{Z}} r_n \left(e^{\pi ib_{n+1}/t} + e^{\pi ib_n/t}\right), \forall t \in \mathbb{R}.\tag{5.33}
$$

This last equation will be the main setup for our proofs.

Part (I): We first deal with the case of general C -H.U.P.'s. Note that both sides in (5.33) define a tempered distribution. Hence, taking Fourier transforms on both sides implies that

$$
\sum_{n\in\mathbb{Z}} t_n \left(\delta_{a_{n+1}} + \delta_{a_n} \right) = -\sum_{n\in\mathbb{Z}} r_n \left(\int_{\mathbb{R}} e^{-\pi i t \xi} e^{\pi i b_{n+1}/t} dt + \int_{\mathbb{R}} e^{-\pi i t \xi} e^{\pi i b_n/t} dt \right). \tag{5.34}
$$

Here, the integrals on the right-hand side of [\(5.34\)](#page-32-4) have to be interpreted in the principal value sense. Now, using the results from [\[28,](#page-35-13) Proposition 5.2.1], such integrals may be evaluated as follows:

$$
\lim_{\varepsilon \to 0} \int_{\mathbb{R}} e^{-\pi i (t\xi - \eta/t)} e^{-\varepsilon|t|} dt = 2\pi \cdot \delta_0(|\eta|/|\xi|) - 2\pi \sqrt{|\eta|/|\xi|} J_1\left(2\pi \sqrt{|\xi\eta|}\right) 1_{\{\xi\eta < 0\}}.\tag{5.35}
$$

Hence, rewriting (5.34) in light of (5.35) ,

$$
\sum_{n\in\mathbb{Z}} t_n \left(\delta_{a_{n+1}} + \delta_{a_n} \right)
$$
\n
$$
= \frac{2\pi}{\sqrt{|\xi|}} \sum_{n\in\mathbb{Z}} r_n \left(\sqrt{|b_n|} J_1 \left(2\pi \sqrt{|\xi b_n|} \right) 1_{\{b_n \xi < 0\}} + \sqrt{|b_{n+1}|} J_1 \left(2\pi \sqrt{|\xi b_{n+1}|} \right) 1_{\{b_{n+1}\xi < 0\}} \right). \tag{5.36}
$$

Now, since $|J_1(s)| \leq C/s^{1/2}$ holds for any $s > 0$, we conclude that, if we know that

$$
\sum_{n \in \mathbb{Z}} |r_n||b_n|^{1/4} < +\infty,\tag{5.37}
$$

then the right-hand side of [\(5.36\)](#page-32-6) converges absolutely for any $\xi: |\xi| > 0$. This shows that $t_{n+1} =$ $-t_n$ for all but at most one $n \in \mathbb{Z}$, which is a contradiction to the fact that $\{t_n\}_{n\in\mathbb{Z}} \in \ell^2(\mathbb{Z})$, unless $\psi \equiv 0$, the desired conclusion which we desired to reach. Since [\(5.37\)](#page-33-0) follows if, for instance,

$$
\int_{\mathbb{R}} |\widehat{\psi}(x)||x|^{1/4} dx < +\infty,
$$

we finish this part accordingly.

Part (II): We focus, in the remaining part, on the original case of translated lattice crosses con-sidered in [\[27,](#page-35-0) [28,](#page-35-13) [20,](#page-35-16) [21\]](#page-35-17). We rewrite [\(5.33\)](#page-32-3) with $a_n = n + \theta, b_n = n$ as follows:

$$
e^{i\theta t} \sum_{n \in \mathbb{Z}} t_n \left(e^{\pi i nt} + e^{\pi i (n+1)t} \right) = - \sum_{n \in \mathbb{Z}} r_n \left(e^{\pi i (n+1)/t} + e^{\pi i n/t} \right).
$$
 (5.38)

Let the absolute value of the left-hand side of [\(5.38\)](#page-33-1) be denoted by $\Psi(t)$. This equation tells us thus that Ψ satisfies

$$
\begin{cases}\n\Psi(t+2) = \Psi(t) & \text{for all } t \in \mathbb{R}, \\
\Psi\left(\frac{1}{t}\right) = \Psi\left(\frac{1}{t+2}\right) & \text{for all } t \in \mathbb{R} \setminus \{0, -2\}.\n\end{cases}
$$
\n(5.39)

We further note that, by the properties of ψ , we must have necessarily $\Psi \in L^2(-1,1)$. We claim that these conditions imply that Ψ is constant.

Indeed, if not, then fix an arbitrary $\alpha > 0$ such that the set

$$
E_\alpha = \{t \in \mathbb{R} \colon \Psi(t) > \alpha\}
$$

satisfies $0 < |E_\alpha \cap (-1, 1)| < 2$. Consider now the function 1_{E_α} . By (5.39) , $t \in E_\alpha$ holds if and only if $t + 2 \in E_\alpha$, and also $t \in E_\alpha$ if and only if $\frac{t}{1-2t} \in E_\alpha$. Hence, 1_{E_α} satisfies the relationships in [\(5.39\)](#page-33-2) as well. We now wish to contradict the fact that $0 < |E_\alpha \cap (-1, 1)| < 2$, implying that either $|E_{\alpha}| = 0$ or $|\mathbb{R} \setminus E_{\alpha}| = 0$. Since $\alpha > 0$ is arbitrary, we may conclude that Ψ itself is constant. Hence, replacing Ψ by $1_{E_{\alpha}}$ in the argument below if needed, we may suppose that $\Psi \in L^{\infty}(\mathbb{R})$.

The rest of the argument follows the same footsteps as in [\[27,](#page-35-0) Section 5]: indeed, while one could simply invoke Proposition 5.1 in [\[27\]](#page-35-0), we chose to include the argument used for it below for completeness.

Let then $\tilde{\Psi}: \mathbb{C}_+ \to \mathbb{C}$ denote the Poisson extension of Ψ to the upper half plane. By definition $\tilde{\Psi}$ is a bounded harmonic function on the upper half plane, and a direct computation shows that

$$
\begin{cases} \tilde{\Psi}(z+2) = \tilde{\Psi}(z) & \text{for all } z \in \mathbb{C}_{+}, \\ \tilde{\Psi}(z) = \tilde{\Psi}\left(\frac{z}{1-2z}\right) & \text{for all } z \in \mathbb{C}_{+}. \end{cases}
$$
\n(5.40)

Let G denote the group of transformations preserving \mathbb{C}_+ generated by $z \mapsto z + 2$ and $z \mapsto \frac{z}{1-2z}$. By for instance the results in [\[19\]](#page-35-21) and [\[4\]](#page-34-9), we deduce that G is a *discrete and free group*, and a fundamental domain for \mathbb{C}_+/\mathcal{G} is given by

$$
\mathcal{D} = \left\{ z \in \mathbb{C}_+ : |\text{Re}(z)| < 1, \left| z - \frac{1}{2} \right| > \frac{1}{2} \text{ and } \left| z + \frac{1}{2} \right| > \frac{1}{2} \right\}.
$$

It follows that $\tilde{\Psi}$ defines a bounded harmonic function on \mathcal{D} , and since the domain \mathcal{D} only has cusps at $0, \infty$ and ± 1 , these are removable singularities of $\tilde{\Psi}$ when seen as a harmonic function on the quotient \mathbb{C}_+/\mathcal{G} . Thus, $\tilde{\Psi}$ extends to a harmonic function on the compact Riemann surface induced by \mathbb{C}_+/\mathcal{G} , which, by Liouville's theorem, implies $\tilde{\Psi}$ is constant, which yields in turn that Ψ is constant.

Finally, note that, given that Ψ is constant, the only viable option is that $\Psi \equiv 0$, since turning to [\(5.31\)](#page-32-1) shows that $\psi(t) = \frac{c}{t^2 - 1} \in L^2(\mathbb{R})$, which can only hold if $c = 0$, as desired.

Remark 5.4. Note that [\(5.37\)](#page-33-0) is fundamentally different from the \mathcal{H}_2 condition of Theorem [2.](#page-4-0) Indeed, the condition of belonging to the class $\mathcal C$ may be suitably translated into an assertion on regularity of ψ . On the other hand, $\psi \in \mathcal{H}_2$ is an assertion on the *decay* of ψ . We believe, in light of these comments, that a more careful analysis could lead to a sharper result on uniqueness of distributions solving [\(5.33\)](#page-32-3) with square-summable coefficients, possibly by employing techniques from almost periodic functions, as done in [\[22\]](#page-35-22).

ACKNOWLEDGEMENTS

We thank Mateus Sousa and Felipe Gonçalves for several discussions that led to the investigation of the main results in this manuscript.

D.R. acknowledges funding by the European Union (ERC, FourIntExP, 101078782). Views and opinions expressed are those of the author(s) only and do not necessarily reflect those of the European Union or European Research Council (ERC). Neither the European Union nor ERC can be held responsible for them.

REFERENCES

- [1] W. O. Amrein and A. M. Berthier, *On support properties of* L p *-functions and their Fourier transforms*, J. Functional Analysis 24 (1977), no. 3, 258–267. MR 0461025
- [2] Andrew Bakan, Haakan Hedenmalm, and Alfonso Montes-Rodriguez, *Hyperbolic Fourier series and the Klein-Gordon equation*, arXiv preprint arXiv:2401.06871 (2024).
- [3] Andrew Bakan, Haakan Hedenmalm, Alfonso Montes-Rodríguez, Danylo Radchenko, and Maryna Viazovska, *Fourier uniqueness in even dimensions*, Proceedings of the National Academy of Sciences 118 (2021), no. 15, e2023227118.
- [4] Alan F. Beardon, *The geometry of discrete groups*, Graduate Texts in Mathematics, vol. 91, Springer-Verlag, New York, 1995, Corrected reprint of the 1983 original. MR 1393195
- [5] Michael Benedicks, *On Fourier transforms of functions supported on sets of finite Lebesgue measure*, J. Math. Anal. Appl. 106 (1985), no. 1, 180–183. MR 780328
- [6] A. M. Berthier and J. M. Jauch, A theorem on the support of functions in $L^2(\mathbf{R})$ and of their Fourier transforms, Lett. Math. Phys. 1 (1975/76), no. 2, 93–97. MR 412727
- [7] Francisco Canto-Martín, Håkan Hedenmalm, and Alfonso Montes-Rodriguez, *Perron-Frobenius operators and the Klein-Gordon equation.*, Journal of the European Mathematical Society (EMS Publishing) 16 (2014), no. 1, 31–66.
- [8] Henry Cohn, Abhinav Kumar, Stephen Miller, Danylo Radchenko, and Maryna Viazovska, *The sphere packing problem in dimension* 24, Annals of Mathematics 185 (2017), no. 3, 1017–1033.
- [9] $____\$ contrast *Dniversal optimality of the* E₈ *and Leech lattices and interpolation formulas*, Annals of Mathematics 196 (2022), no. 3, 983–1082.
- [10] M. Cowling, L. Escauriaza, C. E. Kenig, G. Ponce, and L. Vega, *The Hardy uncertainty principle revisited*, Indiana Univ. Math. J. 59 (2010), no. 6, 2007–2025. MR 2919746
- [11] *NIST Digital Library of Mathematical Functions*, <https://dlmf.nist.gov/>, Release 1.1.12 of 2023-12-15, F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [12] L. Escauriaza, C. E. Kenig, G. Ponce, and L. Vega, *On uniqueness properties of solutions of Schrödinger equations*, Comm. Partial Differential Equations 31 (2006), no. 10-12, 1811–1823. MR 2273975
- [13] , *Convexity properties of solutions to the free Schrödinger equation with Gaussian decay*, Math. Res. Lett. 15 (2008), no. 5, 957–971. MR 2443994
- [14] , *Hardy's uncertainty principle, convexity and Schrödinger evolutions*, J. Eur. Math. Soc. (JEMS) 10 (2008), no. 4, 883–907. MR 2443923
- [15] Luis Escauriaza, Carlos E Kenig, Gustavo Ponce, and Luis Vega, *The sharp Hardy uncertainty principle for Schrödinger evolutions*, Duke Mathematical Journal 155 (2010), no. 1, 163–187.
- [16] Aingeru Fernandez-Bertolin, Karlheinz Gröchenig, and Philippe Jaming, *From Heisenberg uniqueness pairs to properties of the Helmholtz and Laplace equations*, Journal of Mathematical Analysis and Applications 469 (2019), no. 1, 202–219.
- [17] Saifallah Ghobber and Philippe Jaming, *Uncertainty principles for integral operators*, arXiv preprint arXiv:1206.1195 (2012).
- [18] Somnath Ghosh and Debkumar Giri, *Density of exponentials and perron-frobenius operators*, Advances in Mathematics 457 (2024), 109932.
- [19] J. Gilman and B. Maskit, *An algorithm for* 2*-generator Fuchsian groups*, Michigan Math. J. 38 (1991), no. 1, 13–32. MR 1091506
- [20] Deb Kumar Giri and Rama Rawat, *Heisenberg uniqueness pairs for the hyperbola*, Bulletin of the London Mathematical Society 53 (2021), no. 1, 16–25.
- [21] , *Corrigendum: Heisenberg uniqueness pairs for the hyperbola*, Bulletin of the London Mathematical Society 54 (2022), no. 5, 2041–2043.
- [22] Felipe Gonçalves, *A classification of Fourier summation formulas and crystalline measures*, arXiv preprint arXiv:2312.11185 (2023).
- [23] Felipe Gonçalves and João P. G. Ramos, *A note on discrete Heisenberg uniqueness pairs for the parabola*, Bull. Sci. Math. 174 (2022), Paper No. 103095. MR 4362794
- [24] Karlheinz Gröchenig and Philippe Jaming, *The Cramér–Wold theorem on quadratic surfaces and Heisenberg uniqueness pairs*, Journal of the Institute of Mathematics of Jussieu 19 (2020), no. 1, 117–135.
- [25] G. H. Hardy, *A Theorem Concerning Fourier Transforms*, J. London Math. Soc. 8 (1933), no. 3, 227–231. MR 1574130
- [26] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge, at the University Press,, 1952, 2d ed. MR 46395
- [27] Haakan Hedenmalm and Alfonso Montes-Rodríguez, *Heisenberg uniqueness pairs and the Klein-Gordon equation*, Annals of Mathematics (2011), 1507–1527.
- [28] , *The Klein-Gordon equation, the Hilbert transform, and dynamics of Gauss-type maps*, Journal of the European Mathematical Society 22 (2020), no. 6, 1703–1757.
- [29] , *The Klein-Gordon equation, the Hilbert transform and Gauss-type maps:* H [∞] *approximation*, Journal d'Analyse Mathématique 144 (2021), 119-190.
- [30] Philippe Jaming and Karim Kellay, *A dynamical system approach to Heisenberg uniqueness pairs*, Journal d'Analyse Mathématique 134 (2018), no. 1, 273-301.
- [31] Christoph Kehle and João PG Ramos, *Uniqueness of solutions to nonlinear Schrödinger equations from their zeros*, Annals of PDE 8 (2022), no. 2, 21.
- [32] Aleksei Kulikov, Fedor Nazarov, and Mikhail Sodin, *Fourier uniqueness and non-uniqueness pairs*, arXiv preprint arXiv:2306.14013 (2023).
- [33] Aleksei Kulikov, Lucas Oliveira, and João PG Ramos, *On Gaussian decay rates of harmonic oscillators and equivalences of related Fourier uncertainty principles*, Revista matemática iberoamericana 40 (2023), no. 2, 481– 502.
- [34] Nir Lev, *Uniqueness theorems for Fourier transforms*, Bulletin des sciences mathematiques 135 (2011), no. 2, 134–140.
- [35] B Ya Levin, *Lectures on entire functions*, vol. 150, American Mathematical Soc., 1996.
- [36] Jean-Pierre Merx, *A continuous function with divergent Fourier series*, Available at https://www.mathcounterexamples.net/continuous-function-with-divergent-fourier-series/ (2016).
- [37] Danylo Radchenko and Maryna Viazovska, *Fourier interpolation on the real line*, Publ. Math. Inst. Hautes Etudes ´ Sci. 129 (2019), 51–81. MR 3949027
- [38] Jo˜ao P.G. Ramos and Martin Stoller, *Perturbed Fourier uniqueness and interpolation results in higher dimensions*, Journal of Functional Analysis 282 (2022), no. 12, 109448.
- [39] Jo˜ao Pedro Ramos and Mateus Sousa, *Fourier uniqueness pairs of powers of integers*, Journal of the European Mathematical Society 24 (2022), no. 12, 4327–4351.
- [40] Claude E. Shannon, *Communication in the presence of noise*, Proc. I.R.E. 37 (1949), 10–21. MR 28549
- [41] Per Sjölin, *Heisenberg uniqueness pairs and a theorem of Beurling and Malliavin*, Bulletin des sciences mathematiques 135 (2011), no. 2, 125–133.
- [42] , *Heisenberg uniqueness pairs for the parabola*, Journal of Fourier Analysis and Applications 19 (2013), 410–416.
- [43] Martin Stoller, *Fourier interpolation from spheres*, Trans. Amer. Math. Soc. 374 (2021), no. 11, 8045–8079. MR 4328691
- [44] Pál Turán, *Beispiele stetiger Funktionen mit divergenter Fourierreihe*, pp. 538–541, Birkhäuser Basel, Basel, 1970.
- [45] Maryna S. Viazovska, *The sphere packing problem in dimension 8*, Ann. of Math. (2) 185 (2017), no. 3, 991–1015. MR 3664816
- [46] Francisco Javier Gonzalez Vieli, *A uniqueness result for the Fourier transform of measures on the sphere*, Bulletin of the Australian Mathematical Society 86 (2012), no. 1, 78–82.
- [47] George Neville Watson, *A treatise on the theory of bessel functions*, Cambridge Mathematical Library, Cambridge University Press, 1995.
- [48] Edmund Taylor Whittaker, *XVIII.—on the functions which are represented by the expansions of the interpolationtheory*, Proceedings of the Royal Society of Edinburgh 35 (1915), 181–194.

LABORATOIRE PAUL PAINLEVÉ, UNIVERSITY OF LILLE, FRANCE. *Email address*: danradchenko@gmail.com

Institute of Mathematics, EPF Lausanne, Switzerland. *Email address*: joaopgramos95@gmail.com