UNVEILING f −BIHARMONIC θ_{α} −SLANT CURVES IN S− SPACE FORMS

ŞABAN GÜVENÇ

ABSTRACT. In this paper, we firstly provide a concise overview of S −manifolds, f-biharmonicity and θ_{α} -slant curves. We then derive a key equation and analyze it in detail to establish the necessary and sufficient conditions for θ_{α} -slant curves to be f-biharmonic. Finally, we present an example to support our findings.

1. INTRODUCTION

In 1964, J. Eells and L. Maire explored key aspects of harmonic maps, later extending their ideas to k-harmonic maps [\[7\]](#page-12-0). Building on this, G. Y. Jiang studied the case for $k = 2$, deriving foundational variational formulas for 2-harmonic maps [\[15\]](#page-12-1). B. Y. Chen, in a comprehensive survey, addressed biharmonic submanifolds of Euclidean space, particularly focusing on the condition $\Delta H = 0$, with Δ being the Laplace operator and H representing the mean curvature vector field [\[5\]](#page-12-2). Notably, if the ambient space is Euclidean, the works of Jiang and Chen converge.

Expanding this field, J. T. Cho, J. Inoguchi, and J. E. Lee introduced the concept of slant curves in Sasakian manifolds, drawing a parallel to Lancret's theorem for Euclidean space curves [\[6\]](#page-12-3). Their findings highlighted that non-geodesic curves in Sasakian 3-manifolds are slant curves when the ratio of $(\tau \pm 1)$ to the geodesic curvature k remains constant.

Further advancements were made by D. Fetcu and C. Oniciuc, who proposed a method to generate biharmonic submanifolds in Sasakian space forms by utilizing the characteristic vector field flow ξ [\[9\]](#page-12-4). They demonstrated that this flow transforms biharmonic integral submanifolds into biharmonic anti-invariant submanifolds. Building on this work, the author and C . Ozgür examined biharmonic slant curves in S -space forms, aiming to generalize these findings [\[10\]](#page-12-5).

This paper seeks to extend the results for biharmonic slant curves by exploring f–biharmonic θ_{α} –slant curves in S–space forms. The subsequent sections provide definitions, examine the necessary conditions for these curves to be proper f−biharmonic and conclude with an example that demonstrate these concepts.

2. Background and Definitions

Let (M, q) be a $(2m + s)$ -dimensional Riemannian manifold. M is called a *framed metric manifold* with a *framed metric structure* $(\phi, \xi_{\alpha}, \eta_{\alpha}, g), \alpha \in \{1, ..., s\},$

²⁰²⁰ Mathematics Subject Classification. Primary 53C25; Secondary 53C40, 53A04.

Key words and phrases. θ_{α} –slant curve, f-biharmonic curve, Frenet curve, S-space form.

if it satisfies:

$$
\phi^2 X = -X + \sum_{\alpha=1}^s \eta_\alpha(X)\xi_\alpha, \quad \eta_\alpha(\phi(X)) = 0, \quad \eta_\alpha(\xi_\beta) = \delta_{\alpha\beta}, \quad \phi(\xi_\alpha) = 0,
$$

$$
g(X, Y) = g(\phi X, \phi Y) + \sum_{\alpha=1}^s \eta_\alpha(X)\eta_\alpha(Y), \tag{2.1}
$$

$$
\eta_\alpha(X) = g(X, \xi), \quad d\eta_\alpha(X, Y) = -d\eta_\alpha(Y, X) = g(X, \phi Y),
$$

where ϕ is a (1, 1)-type tensor field of rank $2m; \xi_1, ..., \xi_s$ are vector fields; $\eta_1, ..., \eta_s$ are 1-forms, and g is a Riemannian metric on $M; X, Y \in TM$ and $\alpha, \beta \in \{1, ..., s\}$ (see [\[17\]](#page-13-0), [\[20\]](#page-13-1)). The structure $(\phi, \xi_{\alpha}, \eta_{\alpha}, g)$ is said to be an *S*-structure if the Nijenhuis tensor of ϕ is equal to $-2d\eta_{\alpha} \otimes \xi_{\alpha}$ for all $\alpha \in \{1, ..., s\}$ [\[2\]](#page-12-6).

If $s = 1$, a framed metric structure is equivalent to an almost contact metric structure, and an S-structure is equivalent to a Sasakian structure. For an Sstructure, we have the following equations [\[2\]](#page-12-6):

$$
(\nabla_X \phi) Y = \sum_{\alpha=1}^s \left\{ g(\phi X, \phi Y) \xi_\alpha + \eta_\alpha(Y) \phi^2 X \right\},\tag{2.2}
$$

and

$$
\nabla \xi_{\alpha} = -\phi,\tag{2.3}
$$

for all $\alpha = 1, ..., s$. In the case of $s = 1, (2.3)$ $s = 1, (2.3)$ can be derived from [\(2.2\)](#page-1-1).

Let $X \in T_pM$ be orthogonal to $\xi_1, ..., \xi_s$. The plane section spanned by $\{X, \phi X\}$ is called a ϕ -section in T_pM , and its sectional curvature is referred to as the ϕ *sectional curvature.* Let $(M, \phi, \xi_\alpha, \eta_\alpha, g)$ be an S-manifold. If M has constant ϕ -sectional curvature, its curvature tensor R is given by

$$
R(X,Y)Z = \sum_{\alpha,\beta} \left\{ \eta_{\alpha}(X)\eta^{\beta}(Z)\phi^{2}Y - \eta_{\alpha}(Y)\eta^{\beta}(Z)\phi^{2}X \right. \\ \left. - g(\phi X, \phi Z)\eta_{\alpha}(Y)\xi_{\beta} + g(\phi Y, \phi Z)\eta_{\alpha}(X)\xi_{\beta} \right\} \\ + \frac{c+3s}{4} \left\{ -g(\phi Y, \phi Z)\phi^{2}X + g(\phi X, \phi Z)\phi^{2}Y \right\} \\ + \frac{c-s}{4} \left\{ g(X,\phi Z)\phi Y - g(Y,\phi Z)\phi X + 2g(X,\phi Y)\phi Z \right\}, \tag{2.4}
$$

for $X, Y, Z \in TM$ [\[4\]](#page-12-7). In this case, M is called an S-space form and is denoted by $M(c)$. If $s = 1$, an S-space form is equivalent to a Sasakian space form [\[1\]](#page-12-8).

Let (M, g) and (N, h) be Riemannian manifolds, and $\phi : M \to N$ a differentiable map. A *harmonic map* is a critical point of the energy functional of ϕ , defined as

$$
E(\phi) = \frac{1}{2} \int_M \left| d\phi \right|^2 v_g,
$$

(see [\[8\]](#page-12-9)). Furthermore, a *biharmonic map* is a critical point of the bienergy functional

$$
E_2(\phi) = \frac{1}{2} \int_M |\tau(\phi)|^2 \, v_g,
$$

where $\tau(\phi) = \text{trace} \nabla d\phi$ and is called *the first tension field* of ϕ . Jiang derived the biharmonic map equation [\[15\]](#page-12-1):

$$
\tau_2(\phi) = -J^{\phi}(\tau(\phi)) = -\Delta \tau(\phi) - \text{trace} R^N(d\phi, \tau(\phi))d\phi = 0,
$$

where J^{ϕ} denotes the Jacobi operator of ϕ . It is evident that harmonic maps are biharmonic; thus, non-harmonic biharmonic maps are referred to as *proper biharmonic*.

In [\[18\]](#page-13-2), Y.L. Ou proved the following lemma:

Lemma 1. [\[18\]](#page-13-2) *A curve* γ : $(a, b) \rightarrow (M, g)$ *parametrized by arclength is an fbiharmonic curve with a function* $f:(a, b) \rightarrow (0, \infty)$ *if and only if*

$$
f(\nabla_T \nabla_T \nabla_T T - R(T, \nabla_T T)T) + 2f'\nabla_T \nabla_T T + f''\nabla_T T = 0.
$$

As a result, γ is a proper f-biharmonic curve if and only if

$$
\tau_3 = \nabla_T \nabla_T \nabla_T T - R(T, \nabla_T T)T + 2\frac{f'}{f} \nabla_T \nabla_T T + \frac{f''}{f} \nabla_T T = 0,
$$

where f is a non-constant function.

Let $\gamma: I \to M$ be a unit-speed curve in an m-dimensional Riemannian manifold (M, g) . The curve γ is called a *Frenet curve of osculating order* r ($1 \leq r \leq m$), if there exist g-orthonormal vector fields $V_1, V_2, ..., V_r$ along the curve satisfying the Frenet equations:

$$
T = V_1 = \gamma',
$$

\n
$$
\nabla_T V_1 = k_1 V_2,
$$

\n
$$
\nabla_T V_j = -k_{j-1} V_{j-1} + k_j V_{j+1}, \quad 1 < j < r,
$$

\n
$$
\nabla_T V_r = -k_{r-1} V_{r-1}.
$$
\n(2.5)

Here, $k_1, ..., k_{r-1}$ are positive functions known as the curvatures of γ . If $k_1 = 0$, then γ is referred to as a *geodesic*. If k_1 is a non-zero positive constant and $r = 2, \gamma$ is called a *circle*. If $k_1, ..., k_{r-1}$ are all non-zero positive constants, then γ is called a *helix of order* $r (r \geq 3)$. When $r = 3$, it is simply referred to as a *helix*.

A submanifold of an S-manifold is said to be an *integral submanifold* if $\eta_{\alpha}(X) =$ $0, \alpha \in \{1, ..., s\}$, where X is tangent to the submanifold [\[16\]](#page-13-3). A *Legendre curve* is a 1-dimensional integral submanifold of an S-manifold $(M^{2m+s}, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$. More precisely, a unit-speed curve $\gamma : I \to M$ is a Legendre curve if T is g-orthogonal to all ξ_{α} $(\alpha = 1, ..., s)$ [\[19\]](#page-13-4).

3. θ_{α} –SLANT CURVES IN S–MANIFOLDS

In this section, we review the concept of θ_{α} -slant curves in S-manifolds:

Definition 1. [\[13\]](#page-12-10) Let $M = (M^{2m+s}, \phi, \xi_\alpha, \eta_\alpha, g)$ be an *S*-manifold, and let γ : I → *M be a unit-speed curve.* γ *is called a* $θ_α$ -*slant curve if there are constant angles* θ_{α} ($\alpha = 1, \ldots, s$) *such that* $\eta_{\alpha}(T) = \cos \theta_{\alpha}$. These angles θ_{α} are referred to *as the contact angles of* γ *.*

It is easy to observe that Definition [1](#page-2-0) generalizes the family of slant curves to θ_{α} −slant curves. In particular, a θ_{α} −slant curve is called *slant* if all its contact angles are equal (see [\[10\]](#page-12-5)).

For a θ_{α} -slant curve, it is known that $\eta_{\alpha}(V_2) = 0$ for all $\alpha = 1, \ldots, s$. The following notations are used similarly to [\[13\]](#page-12-10):

$$
a = \sum_{\alpha=1}^{s} \cos^{2} \theta_{\alpha}, \quad b = \sum_{\alpha=1}^{s} \cos \theta_{\alpha}, \quad V = \sum_{\alpha=1}^{s} \cos \theta_{\alpha} \xi_{\alpha}.
$$

Corollary 1. [\[13\]](#page-12-10) *If* γ *is a slant curve, then*

$$
a = s \cos^2 \theta
$$
, $b = s \cos \theta$, $V = \cos \theta \sum_{\alpha=1}^{s} \xi_{\alpha}$,

where θ *denotes the common contact angle of* γ *.*

\mathbf{q} saban güvenç

Let γ be a non-geodesic unit-speed θ_{α} -slant curve. Then, $g(\phi T, \phi T) = 1 - a \ge 0$. If $a = 1$, we have $\phi T = 0$, which implies $T = \mathcal{V}$. Hence, $\nabla_T T = \nabla_{\mathcal{V}} \mathcal{V} = 0$, meaning that γ is a geodesic as an integral curve of $\mathcal V$.

Proposition 1. [\[13\]](#page-12-10) *For a non-geodesic unit-speed* θ_{α} *-slant curve in an* S-manifold,

$$
a = \sum_{\alpha=1}^{s} \cos^2 \theta_{\alpha} < 1.
$$

Proposition 2. [\[13\]](#page-12-10) *For a non-geodesic unit-speed* θ_{α} *-slant curve in an S*-manifold $(M, \phi, \xi_\alpha, \eta_\alpha, g)$ *, we have*

$$
\nabla_T \phi T = (1 - a) \sum_{\alpha=1}^{s} \xi_{\alpha} + b(-T + V) + k_1 \phi V_2.
$$
 (3.1)

4. f-BIHARMONIC θ_{α} -SLANT CURVES IN S-SPACE FORMS

In this section, we consider proper f-biharmonic θ_{α} -slant curves in S-space forms. Let γ be a unit-speed θ_{α} -slant curve in an S-space form $(M, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$. In [\[13\]](#page-12-10), it is shown that

$$
R(T, \nabla_T T) T = -k_1 \left[b^2 + \frac{c+3s}{4} (1-a) \right] V_2 - 3k_1 \frac{c-s}{4} g(\phi T, V_2) \phi T,
$$

$$
\tau_2(\gamma) = \nabla_T \nabla_T \nabla_T T - R(T, \nabla_T T) T.
$$

Thus, we can calculate

$$
\tau_3 = \nabla_T \nabla_T \nabla_T T - R(T, \nabla_T T) T + 2 \frac{f'}{f} \nabla_T \nabla_T T + \frac{f''}{f} \nabla_T T
$$
\n
$$
= \left(-3k_1 k_1' - 2k_1^2 \frac{f'}{f} \right) T
$$
\n
$$
+ \left[k_1'' - k_1^3 - k_1 k_2^2 + k_1 \left[b^2 + \frac{c + 3s}{4} (1 - a) \right] + 2 \frac{f'}{f} k_1' + \frac{f''}{f} k_1 \right] V_2
$$
\n
$$
+ \left(2k_1' k_2 + k_1 k_2' + 2 \frac{f'}{f} k_1 k_2 \right) V_3 + k_1 k_2 k_3 V_4 + 3k_1 \frac{c - s}{4} g(\phi T, V_2) \phi T.
$$
\n(4.1)

As a result, we can state the following theorem:

Theorem 1. γ *is a proper f-biharmonic* θ_{α} -slant curve in an S-space form $(M, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$ *if and only if*

$$
3\frac{k_1'}{k_1} + 2\frac{f'}{f} = 0,\t\t(4.2)
$$

$$
k_1^2 + k_2^2 = \frac{k_1''}{k_1} + \frac{f''}{f} + 2\frac{f'}{f}\frac{k_1'}{k_1} + b^2 + \frac{c+3s}{4}(1-a) + 3\frac{c-s}{4}g(\phi T, V_2)^2, \tag{4.3}
$$

$$
k'_2 + 2k_2 \frac{k'_1}{k_1} + 2k_2 \frac{f'}{f} + 3 \frac{c-s}{4} g(\phi T, V_2) g(\phi T, V_3) = 0,
$$
\n(4.4)

$$
k_2 k_3 + 3\frac{c-s}{4} g(\phi T, V_2)g(\phi T, V_4) = 0
$$
\n(4.5)

and $g(\tau_3, \phi T) = 0$ *.*

Proof. Let γ be a proper f-biharmonic θ_{α} -slant curve. Then $\tau_3 = 0$. From equation [4.1,](#page-3-0) if we apply T, V_2 , V_3 , V_4 , and ϕ T, we obtain equations [\(4.2\)](#page-3-1), [\(4.3\)](#page-3-2), [\(4.4\)](#page-3-3), [\(4.5\)](#page-3-4), and $g(\tau_3, \phi T) = 0$. Conversely, if the given equations are satisfied for a θ_{α} -slant curve, it can be easily shown that $\tau_3 = 0$. Therefore, γ is a proper f-biharmonic curve. \square

The following Lemma will be crucial for the reader to understand the results of next cases:

Lemma 2. Let $y = y(x)$ be a real valued function. Furthermore, let $c_2 \geq 0$, $\lambda \geq 0$ *be real constants,* $\varepsilon \in \{-1, 0, +1\}$ *and* $u = u(x) = 2\lambda x + c_4$ *for an arbitrary constant* c4*. Then the autonomous ODE*

$$
3 (y')^{2} - 2yy'' = 4y^{2} [(1 + c_{2}^{2}) y^{2} - \varepsilon \lambda^{2}]
$$

has the general solution of the form

$$
y = \frac{\pm\sqrt{N} + M}{D},\tag{4.6}
$$

where the functions N*,* M *and* D *denotes*

(*i*) *for* $\varepsilon = +1$ *and* $\lambda > 0$:

$$
N = \lambda^2 \sec^2 u. \left[-\left(1 + c_2^2 + c_3^2\right) \sec^2 u + \left(1 + c_2^2 - c_3^2\right) \right],\tag{4.7}
$$

$$
M = \lambda c_3 \sec^2 u, \ D = (1 + c_2^2) \sec^2 u - (1 + c_2^2 - c_3^2); \tag{4.8}
$$

(ii) for
$$
\varepsilon = -1
$$
 and $\lambda > 0$:

$$
N = \lambda^2 \operatorname{sech}^2 u. \left[\left(1 + c_2^2 + c_3^2 \right) \operatorname{sech}^2 u - \left(1 + c_2^2 + c_3^2 \right) \right],\tag{4.9}
$$

$$
M = \lambda c_3 \operatorname{sech}^2 u, \ D = \left(1 + c_2^2\right) \operatorname{sech}^2 u - \left(1 + c_2^2 + c_3^2\right); \tag{4.10}
$$

(*iii*) *for* $\varepsilon = 0$ *or* $\lambda = 0$:

$$
N = 0, \ M = 4c_3,\tag{4.11}
$$

$$
D = c_3^2 x^2 + 2c_3^2 c_4 x + c_3^2 c_4^2 + 16c_2^2 + 16.
$$
 (4.12)

Note that the arbitrary constants of the general solutions in Lemma [2](#page-4-0) are denoted by c_3 and c_4 to distinguish from c_1 and c_2 , which will be used in some other equations later.

Now, we are all set to consider the equation $g(\tau_3, \phi T) = 0$ from all points of view. Notice that if the coefficient of ϕT vanishes, then $g(\tau_3, \phi T) = 0$ is satisfied directly. The cases are:

Case I. $c = s$.

Case II. $c \neq s$ and $q(\phi T, V_2) = 0$.

When the coefficient of ϕT does not vanish, we will investigate two more cases: **Case III.** $c \neq s$ and $\phi T \parallel V_2$.

Case IV. $c \neq s$ and $g(\phi T, V_2) \neq 0$ or $\pm \sqrt{1-a}$.

One might ask what happens if ϕT itself vanishes as a vector field. In this case, as shown in [\[13\]](#page-12-10), we have $T = V = \sum_{\alpha=1}^{s} \cos \theta_{\alpha} \xi_{\alpha}$ and $k_1 = 0$. Thus, equation [\(4.2\)](#page-3-1) implies that f is a constant. Consequently, γ cannot be a proper f -biharmonic curve. Note that being proper f -biharmonic means f -biharmonic but not biharmonic. Likewise, being proper biharmonic means biharmonic but not harmonic.

Case I. $c = s$.

In this case, equations [\(4.2-](#page-3-1)[4.5\)](#page-3-4) become

$$
3\frac{k_1'}{k_1} + 2\frac{f'}{f} = 0,\t\t(4.13)
$$

$$
k_1^2 + k_2^2 = \frac{k_1''}{k_1} + \frac{f''}{f} + 2\frac{f'}{f}\frac{k_1'}{k_1} + b^2 + s(1 - a),\tag{4.14}
$$

$$
k'_2 + 2k_2 \frac{k'_1}{k_1} + 2k_2 \frac{f'}{f} = 0,\t\t(4.15)
$$

$$
k_2 k_3 = 0,\t\t(4.16)
$$

By solving these equations, we can state the following theorem:

Theorem 2. *Under the assumption* $c = s$, γ *is a proper f-biharmonic* θ_{α} -slant *curve in* $(M, \phi, \xi_\alpha, \eta_\alpha, g)$ *if and only if:*

(*i*) γ *is of osculating order* $r = 3$ *with* $f = c_1 k_1^{-3/2}$, $\frac{k_2}{k_1} = c_2$, and $k_1 = k_1$ (*t*) *is of the form*

$$
k_1(t) = \frac{\pm\sqrt{N} + M}{D},
$$

where the functions N*,* M *and* D *are denoted by equations* (4.[7\)](#page-4-1) *and* [\(4](#page-4-2).8) *and*

$$
\lambda_1 = \sqrt{b^2 + s(1 - a)} > 0, \ u = u(t) = 2\lambda_1 t + c_4,
$$

 $c_1 > 0$, $c_2 > 0$, c_3 *and* c_4 *are arbitrary constants, t is the arc-length parameter; or* (ii) γ *is of osculating order* $r = 2$ *and satisfies the above equations and inequalities, except for the inequality* $c_2 > 0$ *, where instead* $c_2 = 0$ *.*

Proof. Let $k_1 = k_1(t)$, where t denotes the arc-length parameter. From equation (4.[13\)](#page-5-0), it is easy to see that $f = c_1 k_1^{-3/2}$ for an arbitrary constant $c_1 > 0$. Thus, we obtain

$$
\frac{f'}{f} = \frac{-3 k_1'}{2 k_1}, \quad \frac{f''}{f} = \frac{15}{4} \left(\frac{k_1'}{k_1}\right)^2 - \frac{3 k_1''}{2 k_1}.
$$
\n(4.17)

If $k_2 = 0$, then γ is of osculating order $r = 2$, and equations (4.[13\)](#page-5-0) and (4.[14\)](#page-5-1) must be satisfied. Therefore, the second equation combined with (4.[17\)](#page-5-2) yields the ODE

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2 \left[k_1^2 - \left(b^2 + s(1-a) \right) \right]. \tag{4.18}
$$

This ODE can be solved using Lemma [2.](#page-4-0) Firstly, $1 + c_2^2 = 1$ gives us $c_2 = 0$. Since $s \geq 1, 1 - a > 0$ and $b^2 \geq 0$, we have $b^2 + s(1 - a) > 0$. We can write

$$
b^{2} + s(1 - a) = sgn (b^{2} + s(1 - a)) \cdot (\sqrt{b^{2} + s(1 - a)})^{2}
$$

= $\varepsilon \lambda_{1}^{2}$,

where we denote $\lambda_1 = \sqrt{b^2 + s(1 - a)}$. Now that $\varepsilon = +1$ and $\lambda_1 > 0$, using Lemma [2](#page-4-0) (i), we find that $k_1(t)$ is of the form [\(4](#page-4-3).6) where N, M and D are as in equations (4.[7\)](#page-4-1) and (4.[8\)](#page-4-2) (with $c_2 = 0$). Notice that $c_2 = 0$ is equivalent to $k_2 = 0 = c_2 k_1$ when $r = 2$.

If k_2 = constant $\neq 0$, we find that f is a constant, indicating that γ is not proper f-biharmonic in this case. For $k_2 \neq$ constant, from equation [\(4.16\)](#page-5-3), we have $k_3 = 0$. Thus, γ is of osculating order $r = 3$. Using equation [\(4.17\)](#page-5-2), equation [\(4.15\)](#page-5-4) gives us $\frac{k_2}{k_1} = c_2$, where $c_2 > 0$ is a constant. Substituting these results into equation (4.14) , we obtain the ODE

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2[(1+c_2^2)k_1^2 - (b^2 + s(1-a))]
$$

which gives the general solution using Lemma [2](#page-4-0) and the proof is completed. \Box

Case II. $c \neq s$ and $\phi T \perp V_2$.

In this case, $g(\phi T, V_2) = 0$. From Theorem [1,](#page-3-5) we obtain

$$
\begin{array}{c} 3\frac{k_1'}{k_1}+2\frac{f'}{f}=0,\\ k_1^2+k_2^2=\frac{k_1''}{k_1}+\frac{f''}{f}+2\frac{k_1'}{k_1}\frac{f'}{f}+b^2+\frac{c+3s}{4}(1-a),\\ k_2'+2k_2\frac{f'}{f}+2k_2\frac{k_1'}{k_1}=0,\\ k_2k_3=0.\end{array}
$$

Firstly, we need the following Lemma from [\[13\]](#page-12-10):

Lemma 3. [\[13\]](#page-12-10) Let γ be a θ_{α} -*slant curve of order* $r = 3$ *in an* S-*space form* $(M, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$ and $\phi T \perp V_2$. Then, $\{T, V_2, V_3, \phi T, \nabla_T \phi T, \xi_1, \ldots, \xi_s\}$ is linearly in*dependent.* So $dim M \geq 5 + s$.

Now we have the following theorem:

Theorem 3. Let γ be a θ_{α} -*slant curve in an S*-*space form* $(M^{2m+s}, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$ *,* $\alpha \in \{1, ..., s\}, c \neq s \text{ and } \phi T \perp V_2$. Then γ is proper f-biharmonic if and only if (1) γ *is of osculating order* $r = 3$ *with* $f = c_1 k_1^{-3/2}$, $\frac{k_2}{k_1} = c_2$, $m \ge 3$,

$$
\{T=V_1, V_2, V_3, \phi T, \nabla_T \phi T, \xi_1, ..., \xi_s\}
$$

is linearly independent and $k_1 = k_1(t)$ *is of the form*

$$
k_1 = \frac{\pm\sqrt{N} + M}{D}
$$

where the functions N*,* M *and* D *are denoted by*

(a) *equations* [\(4](#page-4-2).7) *and* (4.8), *for* $b^2 + [(c+3s)/4](1-a) > 0;$

(b) *equations* [\(4](#page-4-4).9) *and* (4.[10\)](#page-4-5), *for* $b^2 + [(c+3s)/4](1-a) < 0$;

(c) *equations* (4.[11\)](#page-4-6) *and* (4.[12\)](#page-4-7), *for* $b^2 + [(c+3s)/4](1-a) = 0;$

with

$$
\lambda_2 = \sqrt{\left|b^2 + \frac{(c+3s)}{4}(1-a)\right|}, \ u = u(t) = 2\lambda_2 t + c_4,
$$

for arbitrary constants $c_1 > 0$, $c_2 > 0$, c_3 *and* c_4 *; or*

(2) γ *is of osculating order* $r = 2$ *and satisfies the above equations and inequalities, except for* $m \geq 3$, $c_2 > 0$,

$$
\{T = V_1, V_2, V_3, \phi T, \nabla_T \phi T, \xi_1, ..., \xi_s\}
$$

is linearly independent, where instead $m \geq 2$, $c_2 = 0$,

$$
\{T = V_1, V_2, \phi T, \nabla_T \phi T, \xi_1, ..., \xi_s\}
$$

is linearly independent.

Proof. The proof is similar to the proof of Theorem [2.](#page-5-5) In this case, we have the ODE

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2[(1+c_2^2)k_1^2 - \left[b^2 + \frac{(c+3s)}{4}(1-a)\right].
$$

We use Lemma [2](#page-4-0) using the fact that

$$
b^{2} + \frac{c+3s}{4}(1-a) = sgn\left(b^{2} + \frac{c+3s}{4}(1-a)\right)\left(\sqrt{\left|b^{2} + \frac{c+3s}{4}(1-a)\right|}\right)^{2}
$$

$$
= \varepsilon\lambda_{2}^{2}.
$$

For $b^2 + [(c+3s)/4] (1-a) > 0$, we have $\varepsilon = +1$ and $\lambda_2 > 0$. For $b^2 + [(c+3s)/4] (1-a)$ a) < 0, we find $\varepsilon = -1$ and $\lambda_2 > 0$. Finally $b^2 + [(c + 3s)/4] (1 - a) = 0$ gives $\varepsilon = 0$ and $\lambda_2 = 0$.

Case III. $c \neq s$, $\phi T \parallel V_2$.

As a result of the assumptions of this case, we have $\phi T = \epsilon \sqrt{1 - a} V_2$, $g(\phi T, V_2) =$ $\epsilon\sqrt{1-a}$, $g(\phi T, V_3) = 0$ and $g(\phi T, V_4) = 0$, where $\epsilon = sgn(g(\phi T, V_2)) = \pm 1$. From equations (4.4) (4.4) and (4.17) (4.17) , we can write

$$
k'_2 + 2k_2 \left(\frac{-3}{2} \frac{k'_1}{k_1}\right) + 2k_2 \frac{k'_1}{k_1} = 0.
$$
 (4.19)

Integrating (4.[19\)](#page-7-0), we have

$$
\frac{k_2}{k_1} = c_2,
$$

for some constant $c_2 > 0$. Likewise in [\[13\]](#page-12-10), for non-constant k_1 , one can show that if $\phi T = \epsilon \sqrt{1 - aV_2}$, then

$$
k_2 = \sqrt{ad^2 - as + b^2 + 2\epsilon bd + s},
$$

where $d = k_1/\sqrt{1 - a}$. So we get

$$
\frac{\sqrt{ad^2 - as + b^2 + 2ebd + s}}{k_1} = c_2,
$$

which is equivalent to

$$
\left(c_2^2 + \frac{a}{a-1}\right)k_1^2 + \frac{2\epsilon b}{a-1}k_1 + (as - b^2 - s) = 0.
$$

If this equation is quadratic or linear in terms of k_1 , then k_1 becomes a constant and γ cannot be proper f-biharmonic. Let us assume

$$
c_2^2 + \frac{a}{a-1} = 0,
$$

$$
\frac{2eb}{a-1} = 0,
$$

$$
as - b^2 - s = 0.
$$

The second equation above gives that $b = 0$. Then the last equation reduces to $as - s = 0$, that is, $a = 1$. But in this case γ becomes a geodesic as an integral curve of V and cannot be proper f-biharmonic. Hence, we give the following result:

Theorem 4. *There does not exist any proper* f-biharmonic θ_{α} −*slant curve in an* S-space form $(M^{2m+s}, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$ with $c \neq s$ and $\phi T \parallel V_2$.

Case IV. $c \neq s$ and $g(\phi T, V_2)$ is not 0 or $\pm \sqrt{1-a}$.

In this final case, let $(M^{2m+s}, \phi, \xi_\alpha, \eta_\alpha, g)$ be an S-space form and $\gamma : I \to M$ a θ_{α} -slant curve of osculating order r. Note that $r = 2$ corresponds to $\phi T \in \text{span}\{V_2\},$ which was investigated in Case III. So let $3 \leq r \leq 2m + s$. If γ is f-biharmonic, then $\phi T \in \text{span}\{V_2, V_3, V_4\}$. Let $\beta(t)$ denote the angle function between ϕT and V_2 ,

that is, $g(\phi T, V_2) = \sqrt{1 - a} \cos \beta(t)$. If we differentiate $g(\phi T, V_2)$ along γ and use equations (2.5) (2.5) and (3.1) , we get

$$
-\sqrt{1-a}\beta'(t)\sin\beta(t) = \nabla_T g(\phi T, V_2) = g(\nabla_T \phi T, V_2) + g(\phi T, \nabla_T V_2)
$$

= $g\left((1-a)\sum_{\alpha=1}^s \xi_\alpha + b(-T+V) + k_1 \phi V_2, V_2\right) + g(\phi T, -k_1 T + k_2 V_3)$
= $k_2 g(\phi T, V_3).$ (4.20)

If we write $\phi T = g(\phi T, V_2)V_2 + g(\phi T, V_3)V_3 + g(\phi T, V_4)V_4$, Theorem [1](#page-3-5) gives us

$$
3\frac{k_1'}{k_1} + 2\frac{f'}{f} = 0,\t\t(4.21)
$$

$$
k_1^2 + k_2^2 = \frac{k_1''}{k_1} + \frac{f''}{f} + 2\frac{f'}{f}\frac{k_1'}{k_1} + \left[b^2 + \frac{c+3s}{4}(1-a) + 3\frac{c-s}{4}(1-a)\cos^2\beta\right],
$$
 (4.22)

$$
k_2' + 2k_2 \frac{k_1'}{k_1} + 2k_2 \frac{f'}{f} + 3 \frac{c - s}{4} \sqrt{1 - a} \cos \beta g(\phi T, V_3) = 0, \tag{4.23}
$$

$$
k_2 k_3 + 3 \frac{c - s}{4} \sqrt{1 - a} \cos \beta g(\phi T, V_4) = 0.
$$
 (4.24)

If we substitute (4.17) (4.17) into (4.22) (4.22) and (4.23) (4.23) , we find

$$
k_1^2 + k_2^2 = b^2 + \frac{c+3s}{4}(1-a) + 3\frac{c-s}{4}(1-a)\cos^2\beta - \frac{k_1''}{2k_1} + \frac{3}{4}\left(\frac{k_1'}{k_1}\right)^2, \quad (4.25)
$$

$$
k_2' - \frac{k_1'}{k_1}k_2 + 3\frac{c-s}{4}\sqrt{1-a}\cos\beta g(\phi T, V_3) = 0.
$$
 (4.26)

If we multiply (4.26) by $2k_2$ and use (4.20) , we obtain

$$
2k_2k'_2 - 2\frac{k'_1}{k_1}k_2^2 + \frac{3(c-s)}{4}(1-a)(-2\beta'\cos\beta\sin\beta) = 0.
$$
 (4.27)

Let us denote $v(t) = k_2^2(t)$, where t is the arc-length parameter. Then [\(4.27\)](#page-8-4) becomes

$$
v' - 2\frac{k'_1}{k_1}v = -\frac{3(c-s)}{4}(1-a)(-2\beta'\cos\beta\sin\beta),\tag{4.28}
$$

which is a linear ODE. If we solve (4.28) , we get the following results: i) If β is a constant, then

$$
\frac{k_2}{k_1} = c_2,\t\t(4.29)
$$

where $c_2 > 0$ is an arbitrary constant. From (4.[20\)](#page-8-3) and (4.[30\)](#page-8-6), we find $g(\phi T, V_3)$ = 0. Since $\|\phi T\| = \sqrt{1-a}$ and $\phi T = \sqrt{1-a} \cos \beta V_2 + g(\phi T, V_4)V_4$, we obtain $g(\phi T, V_4) = \pm \sqrt{1 - a} \sin \beta$. Using (4.[22\)](#page-8-0) and (4.[29\)](#page-8-7), we have

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2 \left[(1+c_2^2)k_1^2 - b^2 - \frac{c+3s+3(c-s)\cos^2\beta}{4}(1-a) \right].
$$

ii) If $\beta = \beta(t)$ is a non-constant function, then

$$
k_2^2 = -\frac{3(c-s)}{4}(1-a)\cos^2\beta + \mu(t)k_1^2,
$$
\n(4.30)

where

$$
\mu(t) = -\frac{3(c-s)}{2}(1-a) \int \frac{\cos^2 \beta k'_1}{k_1^3} dt.
$$
\n(4.31)

If we substitute (4.30) (4.30) into (4.25) (4.25) , we find

$$
\left[1+\mu(t)\right]k_1^2 = b^2 + \frac{c+3s}{4}(1-a) + \frac{3(c-s)}{2}(1-a)\cos^2\beta - \frac{k_1''}{2k_1} + \frac{3}{4}\left(\frac{k_1'}{k_1}\right)^2.
$$

Hence, we can state the following final theorem:

Theorem 5. Let $\gamma: I \to M$ be a θ_{α} -*slant curve of osculating order* r *in an* S*space form* $(M^{2m+s}, \phi, \xi_{\alpha}, \eta_{\alpha}, g)$ *, where* $r \geq 3$ *,* $c \neq s$ *,* $g(\phi T, V_2) = \sqrt{1 - a} \cos \beta(t)$ *is not* 0 *or* $\pm \sqrt{1-a}$. *Then* γ *is proper f*-biharmonic *if and only if* $f = c_1 k_1^{-3/2}$ *and* (i) *if* β *is a constant,*

$$
\frac{k_2}{k_1} = c_2,
$$

\n
$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2 \left\{ (1+c_2^2)k_1^2 - \left[b^2 + \frac{c+3s+3(c-s)\cos^2\beta}{4}(1-a) \right] \right\}
$$

\n
$$
k_2k_3 = \pm \frac{3(c-s)\sin 2\beta}{8}(1-a),
$$

(*ii*) *if* β *is a non-constant function,*

$$
k_2^2 = -\frac{3(c-s)}{4}(1-a)\cos^2\beta + \mu(t) \cdot k_1^2,
$$

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2 \left\{ (1+\mu(t))k_1^2 - \left[b^2 + \frac{c+3s+3(c-s)\cos^2\beta}{4}(1-a) \right] \right\},
$$

$$
k_2k_3 = \pm \frac{3(c-s)\sin 2\beta \sin w}{8}(1-a),
$$

where c_1 *and* c_2 *are positive constants,*

$$
\phi T = \sqrt{1 - a} \left(\cos \beta V_2 \pm \sin \beta \cos w V_3 \pm \sin \beta \sin w V_4 \right), \tag{4.32}
$$

 $w = w(t)$ *is the angle function between* V_3 *and the orthogonal projection of* ϕT *onto* $span\{V_3, V_4\}$. w *is related to* β *by* $cos w = \frac{\pi}{3}/k_2$ *and* $\mu(t)$ *is given by*

$$
\mu(t) = -\frac{3(c-s)}{2} (1-a) \int \frac{\cos^2 \beta k_1'}{k_1^3} dt.
$$

Proof. The proof follows directly from the preceding calculations. The equation $\cos w = \pm \frac{\beta'}{k_2}$ is derived using equations [\(4.20\)](#page-8-3) and [\(4.32\)](#page-9-0).

In case β is a constant, we can give the following direct corollary of Theorem [5:](#page-9-1)

Corollary 2. Let $\gamma : I \to M$ be a θ_{α} -*slant curve of osculating order* $r \geq 3$ *in* an S-space form $(M^{2m+s}, \phi, \xi_\alpha, \eta_\alpha, g)$, where $c \neq s$, $g(\phi T, V_2) = \sqrt{1 - a} \cos \beta$ is a *constant and* $\beta \in (0, 2\pi) \setminus \{\frac{\pi}{2}, \pi, \frac{3\pi}{2}\}.$ Then γ *is proper* f-biharmonic if and only if $f = c_1 k_1^{-3/2}$, $\frac{k_2}{k_1} = c_2$ and $k_1 = k_1(t)$ *is of the form*

$$
k_1 = \frac{\pm\sqrt{N} + M}{D}
$$

where the functions N*,* M *and* D *are denoted by*

(a) *equations* [\(4](#page-4-1).7) *and* (4.[8\)](#page-4-2), *for* $b^2 + \{ [c + 3s + 3(c - s) \cos^2 \beta] / 4 \} (1 - a) > 0$; (b) *equations* [\(4](#page-4-4).9) *and* (4.[10\)](#page-4-5), *for* $b^2 + \{ [c + 3s + 3(c - s) \cos^2 \beta] / 4 \} (1 - a) < 0;$ (c) *equations* (4.[11\)](#page-4-6) *and* (4.[12\)](#page-4-7), *for* $b^2 + \{ [c + 3s + 3(c - s) \cos^2 \beta] / 4 \}$ (1-a) = 0; *with*

$$
\lambda_4 = \sqrt{\left| b^2 + \frac{c + 3s + 3(c - s)\cos^2\beta}{4} (1 - a) \right|}, u = u(t) = 2\lambda_4 t + c_4,
$$

for arbitrary constants $c_1 > 0$, $c_2 > 0$, c_3 *and* c_4 *;*

$$
k_2 k_3 = \pm \frac{3(c-s)\sin 2\beta}{8} (1-a)
$$

and $\phi T = \sqrt{1 - a} (\cos \beta V_2 \pm \sin \beta V_4)$ *.*

Proof. In this case, we have the ODE

$$
3(k'_1)^2 - 2k_1k''_1 = 4k_1^2[(1+c_2^2)k_1^2 - \left[b^2 + \frac{c+3s+3(c-s)\cos^2\beta}{4}(1-a)\right].
$$

Using Lemma [2](#page-4-0) and Theorem [5,](#page-9-1) the proof is clear. \Box

5. CONSTRUCTION OF AN EXAMPLE IN $\mathbb{R}^4(-6)$

Firstly, let us recall the structures defined on a special S-manifold. Consider $M = \mathbb{R}^{2m+s}$ with the coordinate functions $\{x_1, ..., x_m, y_1, ..., y_m, z_1, ..., z_s\}$ and the following structures:

$$
\xi_{\alpha} = 2 \frac{\partial}{\partial z_{\alpha}}, \quad \alpha = 1, ..., s,
$$

$$
\eta^{\alpha} = \frac{1}{2} \left(dz_{\alpha} - \sum_{i=1}^{m} y_{i} dx_{i} \right), \quad \alpha = 1, ..., s,
$$

$$
\phi X = \sum_{i=1}^{m} Y_{i} \frac{\partial}{\partial x_{i}} - \sum_{i=1}^{m} X_{i} \frac{\partial}{\partial y_{i}} + \left(\sum_{i=1}^{m} Y_{i} y_{i} \right) \left(\sum_{\alpha=1}^{s} \frac{\partial}{\partial z_{\alpha}} \right),
$$

$$
g = \sum_{\alpha=1}^{s} \eta^{\alpha} \otimes \eta^{\alpha} + \frac{1}{4} \sum_{i=1}^{m} (dx_{i} \otimes dx_{i} + dy_{i} \otimes dy_{i}).
$$

Here,

$$
X = \sum_{i=1}^{m} \left(X_i \frac{\partial}{\partial x_i} + Y_i \frac{\partial}{\partial y_i} \right) + \sum_{\alpha=1}^{s} \left(Z_{\alpha} \frac{\partial}{\partial z_{\alpha}} \right) \in \chi(M).
$$

Then, $(\mathbb{R}^{2m+s}, \phi, \xi_{\alpha}, \eta^{\alpha}, g)$ becomes an S-space form with constant ϕ -sectional curvature $-3s$. This manifold is denoted by $\mathbb{R}^{2n+s}(-3s)$ [\[14\]](#page-12-11). The following vector fields

$$
X_i = 2\frac{\partial}{\partial y_i}, \quad X_{m+i} = \phi X_i = 2\left(\frac{\partial}{\partial x_i} + y_i \sum_{\alpha=1}^s \frac{\partial}{\partial z_\alpha}\right), \quad \xi_\alpha = 2\frac{\partial}{\partial z_\alpha}
$$

form a g-orthonormal basis of $\chi(M)$, and the Riemannian connection is given by

$$
\nabla_{X_i} X_j = \nabla_{X_{m+i}} X_{m+j} = 0, \quad \nabla_{X_i} X_{m+j} = \delta_{ij} \sum_{\alpha=1}^s \xi_\alpha, \quad \nabla_{X_{m+i}} X_j = -\delta_{ij} \sum_{\alpha=1}^s \xi_\alpha,
$$

$$
\nabla_{X_i} \xi_\alpha = \nabla_{\xi_\alpha} X_i = -X_{m+i}, \quad \nabla_{X_{m+i}} \xi_\alpha = \nabla_{\xi_\alpha} X_{m+i} = X_i,
$$

[\[14\]](#page-12-11). Let us choose $m = 2$ and $s = 2$. Now, let $\gamma : I \to \mathbb{R}^6(-6)$, $\gamma = (\gamma_1, ..., \gamma_6)$ be a unit-speed θ_{α} -slant curve. We can calculate

$$
T = \frac{1}{2} \left[\gamma_3' X_1 + \gamma_4' X_2 + \gamma_1' X_3 + \gamma_2' X_4 + \xi_2 \right],
$$

where we take $\theta_1 = \frac{\pi}{2}$ and $\theta_2 = \frac{\pi}{3}$. Then, $g(T, T) = 1$ gives us

$$
(\gamma_1')^2 + (\gamma_2')^2 + (\gamma_3')^2 + (\gamma_4')^2 = 3.
$$
 (5.1)

We also have

$$
\phi T = \frac{1}{2} \left[-\gamma_1' X_1 - \gamma_2' X_2 + \gamma_3' X_3 + \gamma_4' X_4 \right]. \tag{5.2}
$$

After calculations, we obtain

$$
\nabla_T T = \frac{1}{2} \left[\begin{array}{c} (\gamma_3'' + \gamma_1') X_1 + (\gamma_4'' + \gamma_2') X_2 \\ + (\gamma_1'' - \gamma_3') X_3 + (\gamma_2'' - \gamma_4') X_4 \end{array} \right]. \tag{5.3}
$$

Let us select

$$
k_1 = \frac{1}{2+t^2},
$$

\n
$$
k_2 = \frac{1}{2+t^2},
$$
\n(5.4)

$$
k_3 = \frac{\sqrt{17}}{4} (2 + t^2), \tag{5.5}
$$

$$
g(\phi T, V_2) = \frac{\sqrt{3}}{2} \cos \beta = \text{constant},
$$

$$
\cos \beta = \pm \frac{\sqrt{2}}{6} (\beta \approx 1.3329 \text{ or } 1.8087),
$$

$$
f = (2 + t^2)^{3/2},
$$

From (5.3) and (5.4) , we get

$$
\begin{bmatrix} (\gamma_3'' + \gamma_1')^2 + (\gamma_4'' + \gamma_2')^2 \\ + (\gamma_1'' - \gamma_3')^2 + (\gamma_2'' - \gamma_4')^2 \end{bmatrix} = \frac{4}{(2 + t^2)^2}.
$$

We also have $\eta^1(T) = \cos \theta_1 = 0$ and $\eta^2(T) = \cos \theta_2 = 1/2$, which leads to

$$
\gamma_5' = \gamma_1' \gamma_3 + \gamma_2' \gamma_4,
$$

$$
\gamma_6' = 1 + \gamma_1' \gamma_3 + \gamma_2' \gamma_4.
$$

Using $\nabla_T T = k_1 V_2$, we get

$$
g(\phi T, V_2) = \frac{1}{k_1} g(\phi T, \nabla_T T)
$$

=
$$
\frac{2 + t^2}{4} (-3 + \gamma_1'' \gamma_3' - \gamma_1' \gamma_3'' + \gamma_2'' \gamma_4' - \gamma_2' \gamma_4'').
$$

Notice that for $c_1 = 1$, $c_2 = 1$, $c_3 = 4$, $c_4 = 0$, Corollary [2](#page-9-2) (c) is satisfied. In fact, since $b = 1/2$, $c = -6$, $s = 2$, $\sin 2\beta = \pm \sqrt{17}/9$ and $a = 1/4$, we have

$$
b^{2} + \frac{c + 3s + 3(c - s) \cos^{2} \beta}{4} (1 - a) = 0,
$$

$$
k_{1} = \frac{1}{2 + t^{2}} = \frac{4c_{3}}{c_{3}^{2}x^{2} + 2c_{3}^{2}c_{4}x + c_{3}^{2}c_{4}^{2} + 16c_{2}^{2} + 16},
$$

$$
k_{2} = \frac{1}{2 + t^{2}} = c_{2}k_{1},
$$

$$
k_2 k_3 = \pm \frac{\sqrt{17}}{4} = \pm \frac{3(c-s)\sin 2\beta}{8} (1-a).
$$

As a result, under these circumstances, γ becomes a proper f-biharmonic θ_{α} -slant curve for $f = (2 + t^2)^{3/2}$ in $\mathbb{R}^6(-6)$.

6. Conclusions and Future Work

In this study, the properties of f-biharmonic θ_{α} -slant curves defined on Smanifolds have been investigated. In the future, it will be important to explore broader classes of these structures and conduct studies on applications of θ_{α} –slant curves and their implications in various systems. Recall that θ_{α} –slant curves are defined as all the contact angles are constant seperately. As an idea, two possible generalizations can be given as

$$
a = \sum_{i=1}^{s} \cos^2 \theta_{\alpha} = constant
$$

or

$$
b = \sum_{i=1}^{s} \cos \theta_{\alpha} = constant,
$$

where the contact angles do not need to be constant but their sum or squared sum to be constant. Then θ_{α} –slant curves will be a subclass and the results on these curves will be corollaries of those future studies.

REFERENCES

- [1] Baikoussis, C., Blair, D. E.: On Legendre curves in contact 3-manifolds. Geom. Dedicata. 49, 135-142 (1994).
- [2] Blair, D. E.: Geometry of manifolds with structural group $U(n) \times O(s)$. J. Differential Geometry. 4, 155-167 (1970).
- [3] Blair, D. E.: Riemannian geometry of contact and symplectic manifolds, Second edition. Progress in Mathematics, 203. Birkhauser Boston, Inc., Boston, MA, (2010).
- [4] Cabrerizo, J. L., Fernandez, L. M., Fernandez M.: The curvature of submanifolds of an S-space form. Acta Math. Hungar. 62, 373-383 (1993).
- [5] Chen, B.Y.: A report on submanifolds of finite type. Soochow J. Math. 22, 117-337 (1996).
- [6] Cho, J. T., Inoguchi, J., Lee, J. E.: On slant curves in Sasakian 3-manifolds. Bull. Austral. Math. Soc. **74**, 359-367 (2006).
- [7] Eells, Jr. J., Lemaire, L.: Selected topics in harmonic maps. Amer. Math. Soc. Providence, R.I. (1983).
- [8] Eells, Jr. J., Sampson, J. H.: Harmonic mappings of Riemannian manifolds. Amer. J. Math. 86, 109-160 (1964).
- [9] Fetcu, D., Oniciuc, C.: Explicit formulas for biharmonic submanifolds in Sasakian space forms. Pacific J. Math. 240, 85-107 (2009).
- [10] Güvenç, S., Ozgür, C.: On slant curves in S-manifolds. Commun. Korean Math. Soc. 33 (1), 293-303 (2018).
- [11] Güvenç, Ş. and Özgür, C.: C−parallel and C−proper Slant Curves of S−manifolds. Filomat 33 (19), 6305-6313 (2019).
- [12] Güvenç, S.: A Note on f-biharmonic Legendre Curves in S-space forms. International Electronic Journal of Geometry, 12 (2), 260-267 (2019).
- [13] Güvenç, S.: An Extended Family of Slant Curves in S-manifolds. Mathematical Sciences and Applications E-Notes. 8 (1) , 69-77 (2020).
- [14] Hasegawa, I., Okuyama, Y., Abe, T.: On p-th Sasakian manifolds. J. Hokkaido Univ. Ed. Sect. II A. **37** (1), 1-16 (1986).
- [15] Jiang, G. Y.: 2-harmonic maps and their first and second variational formulas. Chinese Ann. Math. Ser. A. 7, 389-402 (1986).

$\begin{array}{c} {\bf \text{SABAN GÜVENQ}} \end{array}$

- [16] Kim, J. S., Dwivedi, M. K., Tripathi, M. M.: Ricci curvature of integral submanifolds of an S-space form. Bull. Korean Math. Soc. 44, 395-406 (2007).
- [17] Nakagawa, H.: On framed f-manifolds. Kodai Math. Sem. Rep. 18, 293-306 (1966).
- [18] Ou, Y.L.: On f−biharmonic maps and f−biharmonic submanifolds. Pacific Journal of Mathematics **271** (2), 461-477.
- [19] Özgür, C., Güvenç, Ş.: On biharmonic Legendre curves in S-space forms. Turkish J. Math. 38 (3), 454-461 (2014).
- [20] Yano, K., Kon, M.: Structures on Manifolds. Series in Pure Mathematics, 3. World Scientific Publishing Co. Singapore (1984).

(Ş. Güvenç) BALIKESIR UNIVERSITY, DEPARTMENT OF MATHEMATICS, 10145, ÇAĞIŞ, BALIKEsir, TURKEY

 $Email~address, \, \S.$ Güvenç: sguvenc@balikesir.edu.tr

