

ASYMPTOTIC BEHAVIORS FOR VOLTERRA TYPE MCKEAN-VLASOV STOCHASTIC INTEGRAL EQUATIONS WITH SMALL NOISE

SHANQI LIU, YAOZHONG HU, AND HONGJUN GAO

ABSTRACT. This work is devoted to studying asymptotic behaviors for Volterra type McKean-Vlasov stochastic differential equations with small noise. By applying the weak convergence approach, we establish the large and moderate deviation principles. In addition, we obtain the central limit theorem and find the Volterra integral equation satisfied by the limiting process, which involves the Lions derivative of the drift coefficient.

Keywords. Stochastic Volterra integral equations, McKean-Vlasov SDEs, Large deviation principles, Central limit Theorem, Moderate deviation principles.

AMS 2020 Subject Classifications. 60H20; 60F10; 60F05.

1. INTRODUCTION

Consider the following Volterra type McKean-Vlasov stochastic differential equations (SDEs) on \mathbb{R}^d :

$$X_{t,\xi}^\varepsilon = \xi + \int_0^t K_1(t,s)b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})ds + \sqrt{\varepsilon} \int_0^t K_2(t,s)\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})dW_s, \quad 0 \leq t \leq T, \quad (1.1)$$

where $\mathcal{L}_{X_{t,\xi}^\varepsilon} \in \mathcal{P}_2(\mathbb{R}^d)$ denotes the probability law of $X_{t,\xi}^\varepsilon$, $\varepsilon \in (0, 1]$ is a small parameter, $K_i(t, s)$, $i = 1, 2$ are two positive functions defined on the simplex $\Delta_T = \{0 \leq s < t \leq T\}$ which may be singular, W_t is an m -dimensional Brownian motion defined on the classical Wiener space $(\Omega, \mathcal{F}, \mathbb{P})$, ξ is a \mathbb{R}^d valued random variable independent of the Brownian motion B , and the coefficients $\sigma : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}^d \times \mathbb{R}^m$, $b : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}^d$ are Borel measurable. Here we use $\mathcal{P}_2(\mathbb{R}^d)$ to denote the set of all square integrable probability measures on \mathbb{R}^d .

This paper aims to investigate the asymptotic behavior for (1.1). More precisely, we will study the asymptotic behaviors of

$$\frac{X_{t,\xi}^\varepsilon - X_{t,\xi}^0}{\sqrt{\varepsilon} h(\varepsilon)}, \quad t \in [0, T],$$

as $\varepsilon \rightarrow 0$, where X^0 satisfies the following deterministic Volterra integral equation (formally setting $\varepsilon = 0$ in (1.1)):

$$X_{t,\xi}^0 = \xi + \int_0^t K_1(t,s)b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})ds, \quad (1.2)$$

where δ is the Dirac point measure. We focus particularly on the following three tasks.

- (i) The choice $h(\varepsilon) = \frac{1}{\sqrt{\varepsilon}}$ provides the large deviation principle (LDP) with speed ε^{-1} (e.g. Theorem 3.1 in Section 3).
- (ii) The choice $h(\varepsilon) = 1$ yields the central limit theorem (CLT), i.e. $\frac{X_{t,\xi}^\varepsilon - X_{t,\xi}^0}{\sqrt{\varepsilon}}$ converges to a stochastic process which solves a Volterra integral equation involving the Lions derivative of the drift coefficient (e.g. Theorems 4.1 in Section 4).

(iii) The choice $h(\varepsilon)$ satisfying

$$h(\varepsilon) \rightarrow \infty, \quad \sqrt{\varepsilon}h(\varepsilon) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0,$$

fills the gap between the CLT and LDP scales, that is, it provides the moderate deviation estimate with speed $h^2(\varepsilon)$ (e.g. Theorem 5.1 in Section 5).

To complete the first two tasks we shall use the weak convergence criteria presented in [13] and the key point is to identify the corresponding controlled Volterra equations. The main difficulty to complete the third task is to obtain estimate of the form $\mathbb{E}\left(\sup_{t \in [0, T]} |Z_{t, \xi}^\varepsilon - Z_{t, \xi}|^p\right)$ (see Section 4 for the definition of $Z_{t, \xi}^\varepsilon$ and $Z_{t, \xi}$) in particular due to the singularity of kernels. We shall apply extended Kolmogorov's continuity criteria (see e.g. [35]) to overcome this difficulty. This approach extends the approach of Qiao [28] to distribution dependent case.

In the absence of the Volterra kernels, that is the kernels $K_1 = K_2 = 1$, the equation (1.1) is reduced to classical McKean-Vlasov SDEs (also called distribution dependent SDEs [29] or mean-field SDEs). In this case the equation was first studied by Kac in [15, 16], which describe limiting behaviors of individual particles in an interacting particle system of mean-field type when the number of particles goes to infinity (propagation of chaos) and has widely applied to various fields. We refer to [12, 22] and the references therein for more details. For this type of classical McKean-Vlasov SDEs, the small noise asymptotic behaviors have been extensively investigated in recent years. Herrmann et al. [11] obtained the LDP in path space equipped with the uniform norm, when the drift is superlinear but coercive, and the diffusion coefficient is a constant. Dos Reis et al. [8] obtained the LDP in both uniform and Hölder topologies under the assumption that the drift and diffusion coefficients satisfy some extra Hölder continuity with respect to time.

When the coefficients b and σ do not depend on the distribution $\mathcal{L}_{X_{t, \xi}^\varepsilon}$ of the solution process (1.1) is reduced to stochastic Volterra integral equations or Volterra type SDEs. There are now a large amount of work dedicated to this type of equations. They can be driven by general semimartingales in addition to Brownian motion. When the kernel is singular, usually the coefficients needs to be nicer and when the kernel is more regular, the coefficients can be allowed to be more general. Among the large body of literature let us mention [1, 2, 3, 4, 18, 24, 26, 27]. Let us also point out that the Euler and Milstein schemes for these equations with singular kernels was studied in [19]. For these Volterra-type SDEs with regular kernel, a Freidlin–Wentzell's type LDP was established in [23] in continuous functions space and was later extended in [17] to the Besov-Orlicz space. The case of singular kernels was studied in [35] and then in [36]. The MDP for one dimensional stochastic Volterra equations with regular kernels the MDP was studied in [20]. More recently, Jacquier and Pannier [13] established LDP and MDP for stochastic Volterra systems with singular kernels under weaker conditions. Qiao [28] first proved CLT for stochastic Volterra-type SDEs with singular kernels. To the best of our knowledge, there are no LDP, CLT, and MDP results for Volterra-type McKean-Vlasov stochastic integral equations so far.

Compared with classical SDEs, Volterra type SDEs are more complex. On the one hand, these equations are in general neither Markovian nor semi-martingales, and include fractional Brownian motion with Hurst index $H \in (0, 1)$ as a special case when $b = 0$, $\sigma = 1$, and

$$K_2(t, s) = K^H(t, s) = c_H(t-s)^{H-\frac{1}{2}} + c_H\left(\frac{1}{2}-H\right)\int_s^t (\theta-s)^{H-\frac{3}{2}}\left(1-(\frac{s}{\theta})^{\frac{1}{2}-H}\right)d\theta,$$

with

$$c_H = \left(\frac{2H\Gamma(\frac{3}{2}-H)}{\Gamma(H+\frac{1}{2})\Gamma(2-2H)}\right)^{\frac{1}{2}}.$$

Volterra-type McKean-Vlasov SDEs first appeared in [32], in which the authors studied the well-posedness for a class of mean-field backward stochastic Volterra integral equations associated with the Pontryagin's type maximum principle for a controlled mean field Volterra system. This result is then extended in [34] to study the well-posedness and Pontryagin's type maximum principle of mean-field backward doubly stochastic Volterra integral equation. Very recently, the well-posedness for Volterra type McKean-Vlasov SDEs is considered in ([25, Theorem 2.3]) when the kernel is singular but the coefficients b and σ are Lipschitzian, or when kernel is regular with Hölder continuous coefficients ([14, Theorem 2.10 or Theorem 1.3]). The result [25, Theorem 2.3] is extended in [21] to a more general case by applying Volterra-type Gronwall's inequality.

The rest of the paper is organized as follows. In Section 3, we establish the large deviation principle for Volterra type McKean-Vlasov SDEs. The CLT and MDP are presented in Section 4 and Section 5 respectively. To facilitate the reading of the paper, we postpone some auxiliary lemmas to Appendix A.

Throughout this paper C_p denotes a positive constant which may changes from line to line, where the subscript p is used to emphasize that the constant depends on certain parameter(s) p .

2. PRELIMINARIES

2.1. Notations. Throughout this paper, we denote by \mathbb{R}^d the n -dimensional Euclidean space. Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ to be a given complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$, where \mathcal{F}_t is a nondecreasing family of sub- σ fields of \mathcal{F} satisfying the usual conditions. Let \mathcal{P} be the space of all probability measures μ on \mathbb{R}^d , and let

$$\mathcal{P}_2(\mathbb{R}^d) = \left\{ \mu \in \mathcal{P}(\mathbb{R}^d) : \mu(|\cdot|^2) := \int_{\mathbb{R}^d} |x|^2 \mu(dx) < \infty \right\}.$$

It is well known that \mathcal{P}_2 is a Polish space under the Wasserstein distance

$$\mathbb{W}_2(\mu, \nu) := \inf_{\pi \in \mathcal{C}(\mu, \nu)} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |x - y|^2 \pi(dx, dy) \right)^{\frac{1}{2}}, \quad \mu, \nu \in \mathcal{P}_2(\mathbb{R}^d),$$

where $\mathcal{C}(\mu, \nu)$ is the set of couplings for μ and ν ; that is, $\pi \in \mathcal{C}(\mu, \nu)$ is a probability measure on $\mathbb{R}^d \times \mathbb{R}^d$ such that $\pi(\cdot \times \mathbb{R}^d) = \mu$ and $\pi(\mathbb{R}^d \times \cdot) = \nu$.

For measurable functions K and L on $\mathbb{R}_+ \times \mathbb{R}_+$ the convolutions $K * L$ is defined by

$$(K * L)(t, s) = \int_s^t K(t, u) L(u, s) du,$$

for almost all $(t, s) \in \Delta$, where $\Delta := \{(t, s) \in \mathbb{R}_+ \times \mathbb{R}_+ : s \leq t\}$.

2.2. L -derivatives for the functions on $\mathcal{P}_2(\mathbb{R}^d)$. For any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$, the tangent space at μ is given by

$$T_{\mu, 2} := L^2(\mathbb{R}^d \mapsto \mathbb{R}^d; \mu) := \left\{ \phi : \mathbb{R}^d \mapsto \mathbb{R}^d; \phi \text{ is measurable with } \mu(|\phi|^2) := \int_{\mathbb{R}^d} |\phi(x)|^2 \mu(dx) < \infty \right\}.$$

For $\phi \in T_{\mu, 2}$, we set $\|\phi\|_{T_{\mu, 2}} := \int_{\mathbb{R}^d} |\phi(x)|^2 \mu(dx)$.

Definition 2.1. Let $f : \mathcal{P}_2(\mathbb{R}^d) \mapsto \mathbb{R}$ be a continuous function, and let I be the identity map on \mathbb{R}^d .

(i) If for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$

$$T_{\mu, 2} \ni \phi \mapsto D_\phi^L f(\mu) := \lim_{\varepsilon \rightarrow 0} \frac{f(\mu \circ (I + \varepsilon \phi)^{-1}) - f(\mu)}{\varepsilon} \in \mathbb{R}$$

is a well-defined bounded linear functional, then we say that f is intrinsically differentiable at μ and we denote the intrinsic derivative of f at μ by $D_\phi^L f(\mu)$.

(ii) If for any $\mu \in \mathcal{P}_2(\mathbb{R}^d)$

$$\lim_{\|\phi\|_{T_{\mu,2}} \rightarrow 0} \frac{|f(\mu \circ (\text{Id} + \phi)^{-1}) - f(\mu) - D_\phi^L f(\mu)|}{\|\phi\|_{T_{\mu,2}}} = 0,$$

f is called L -differentiable at μ with the L -derivative (i.e. Lions derivative) $D^L f(\mu)$. By the Riesz representation theorem, there exists a unique element $D^L f(\mu) \in T_{\mu,2}$ satisfying

$$\langle D^L f(\mu), \phi \rangle_{T_{\mu,2}} := \int_{\mathbb{R}^d} \langle D^L f(\mu)(x), \phi(x) \rangle \mu(dx) = D_\phi^L f(\mu), \quad \forall \phi \in T_{\mu,2}.$$

(iii) Denote by $C^1(\mathcal{P}_2(\mathbb{R}^d))$ the set of all continuous functionals f on $\mathcal{P}_2(\mathbb{R}^d)$ satisfying that f is L -differentiable at any point $\mu \in \mathcal{P}_2(\mathbb{R}^d)$, and the L -derivative $D^L f(\mu)(x)$ has a version jointly continuous in $(\mu, x) \in \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R}^d$. Moreover, let $C_b^1(\mathcal{P}_2(\mathbb{R}^d))$ be the subset of $C^1(\mathcal{P}_2(\mathbb{R}^d))$ such that $D^L f(\mu)(x)$ is bounded: there is a C_f , depending only on f so that $|D^L f(\mu)(x)| \leq C_f$ for all $\mu \in \mathcal{P}_2$ and $x \in \mathbb{R}^d$.

For a vector-valued function $f = (f_i)$, or a matrix-valued function $f = (f_{i,j})$ with L -differentiable components, we write

$$D_\phi^L f(\mu) = (D_\phi^L f_i(\mu)), \quad \text{or} \quad D_\phi^L f(\mu) = (D_\phi^L f_{i,j}(\mu)), \quad \mu \in \mathcal{P}_2(\mathbb{R}^d).$$

2.3. Volterra type Growall's inequality. We first recall the following lemma due to Girpenberg [10] (Theorem 1).

Lemma 2.2. Let $K : \Delta \rightarrow \mathbb{R}_+$ be a measurable function. Assume that for any $T > 0$,

$$t \mapsto \int_0^t K(t, s) ds \in L^\infty(0, T)$$

and

$$\limsup_{\varepsilon \downarrow 0} \left\| \int_{\cdot + \varepsilon}^{\cdot} K(\cdot + \varepsilon, s) ds \right\|_{L^\infty(0, T)} < 1.$$

Define

$$R_1(t, s) := K(t, s), \quad R_{n+1}(t, s) := \int_s^t K(t, u) R_n(u, s) du, \quad n \in \mathbb{N}. \quad (2.3)$$

Then for any $T > 0$, there exist constants $C_T > 0$ and $\gamma \in (0, 1)$ such that

$$\left\| \int_0^{\cdot} R_n(\cdot, s) ds \right\|_{L^\infty(0, T)} \leq C_T n \gamma^n, \quad \forall n \in \mathbb{N}.$$

In particular, the series

$$R(t, s) := \sum_{n=1}^{\infty} R_n(t, s) \quad (2.4)$$

converges for almost all $(t, s) \in \Delta$. Moreover, we have

$$R(t, s) - K(t, s) = (K * R)(t, s) - (R * K)(t, s), \quad (2.5)$$

and for any $T > 0$,

$$t \mapsto \int_0^t R(t, s) ds \in L^\infty(0, T). \quad (2.6)$$

The function R defined by (2.4) is called the resolvent of the kernel K . The set of all functions K in Lemma 2.2 will be denoted by \mathcal{K} . We now present Gronwall's lemma of Volterra type due to Zhang [36] (Lemma 2.2).

Lemma 2.3. *Let $K \in \mathcal{K}$ and R_n and R be defined by (2.3) and (2.4), respectively. Let $f, g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be two measurable functions satisfying that for any $T > 0$ and some $n \in \mathbb{N}$,*

$$t \mapsto \int_0^t R_n(t, s)f(s)ds \in L^\infty(0, T)$$

and for almost all $t \in (0, \infty)$,

$$\int_0^t R(t, s)g(s)ds < \infty.$$

If for almost all $t \in (0, \infty)$,

$$f(t) \leq g(t) + \int_0^t K(t, s)f(s)ds,$$

then for almost all $t \in (0, \infty)$,

$$f(t) \leq g(t) + \int_0^t R(t, s)g(s)ds.$$

2.4. A criterion of large deviation principles. Let E be a Polish space. For each $\varepsilon > 0$, let X^ε be an E -valued random variable given on $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$.

Definition 2.4. (rate function) *A function $I : E \rightarrow [0, +\infty)$ is called a rate function if I is lower semicontinuous. Moreover, a rate function I is called a good rate function if the level set $\{x \in E : I(x) \leq N\}$ is compact for each constant $N < \infty$.*

Definition 2.5. (Large deviation principle) *A sequence of random variables $\{X^\varepsilon\}$ is said to satisfy the LDP on E with rate function I if the following lower and upper bound conditions hold:*

(i) (Lower bound) For any open set $G \subset E$,

$$\liminf_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(X^\varepsilon \in G) \geq - \inf_{x \in G} I(x).$$

(ii) (Upper bound) For any closed set $F \subset E$,

$$\limsup_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{P}(X^\varepsilon \in F) \leq - \sup_{x \in F} I(x).$$

Definition 2.6. (Laplace principle) *We say that X^ε satisfies the Laplace principle with the rate function I , if for any real bounded continuous function g on E*

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \log \mathbb{E} \left\{ \exp \left[- \frac{g(X^\varepsilon)}{\varepsilon} \right] \right\} = - \inf_{x \in E} (g(x) + I(x)).$$

Lemma 2.7. (Varadhan's lemma [33]) *Let the random sequence $\{X^\varepsilon\}$ with values in a Polish space E fulfill the LDP with a good rate function I . Then $\{X^\varepsilon\}$ fulfills the Laplace principle on E with the same rate function I .*

Lemma 2.8. (Bryc's converse [7]) *The Laplace principle implies the LDP with the same good rate function.*

Combination of Lemmas 2.7 and 2.8 yields that if E is a Polish space and I is a good rate function, then the LDP and Laplace principle are equivalent.

Denote $C_T^d := C([0, T], \mathbb{R}^d)$ the space of continuous functions from $[0, T]$ to \mathbb{R}^d equipped with the supremum norm $\|\phi\|_T := \sup_{t \in [0, T]} |\phi_t|$ and let

$$\mathcal{A} := \left\{ v : \Omega \times [0, T] \rightarrow \mathbb{R}^m \text{ progressively measurable, } E \left[\int_0^T |v_t|^2 dt \right] < +\infty \right\},$$

$$\mathcal{S}_N := \left\{ v \in L^2 : \int_0^T |v_s|^2 ds \leq N \right\},$$

and

$$\mathcal{A}_N := \left\{ v \in \mathcal{A} : v \in \mathcal{S}_N \right\}.$$

Definition 2.9. [13] For all $N \in \mathbb{N}$ and $v \in \mathcal{A}_N$ we define

$$\begin{aligned} \mathcal{G}_{v,N}^0 := & \left\{ \phi : \Omega \rightarrow C_T^d ; \text{ so that there exist } \{\varepsilon_n\}_{n \in \mathbb{N}} \subset [0, \infty) \text{ with } \lim_{n \rightarrow \infty} \varepsilon_n = 0 \right. \\ & \text{and a sequence } \{v^{\varepsilon_n}\}_{n \in \mathbb{N}} \subset \mathcal{A}_N \text{ with } \lim_{n \rightarrow \infty} v^{\varepsilon_n} = v \text{ in distribution such that} \\ & \left. \phi = \lim_{n \rightarrow \infty} \mathcal{G}^{\varepsilon_n} \left(W + \frac{1}{\sqrt{\varepsilon_n}} \int_0^\cdot v_s^{\varepsilon_n} ds \right) \text{ in distribution} \right\}, \end{aligned}$$

and $\mathcal{G}_{v,N}^0$ is empty if v is not in \mathcal{A}_N . For all $v \in \mathcal{A}$, we also denote $\mathcal{G}_v^0 := \bigcup_{n \in \mathbb{N}} \mathcal{G}_{v,N}^0$.

Then we define the functional $I : C_T^d \rightarrow [0, +\infty]$ given by

$$I(\phi) := \inf \left\{ \frac{1}{2} \int_0^T |v_s|^2 ds : v \in L^2 \text{ such that } \phi \in \mathcal{G}_v^0 \right\}. \quad (2.7)$$

Definition 2.10. [13] We say that $\phi \in C_T^d$ is uniquely characterized if there exists a sequence $\{v^n\}_{n \in \mathbb{N}} \subset L^2$ such that

$$\mathcal{G}_{v^n}^0 = \{\phi\} \quad \text{and} \quad \frac{1}{2} \int_0^T |v_s^n|^2 ds \leq I(\phi) + \frac{1}{n},$$

for all $n \in \mathbb{N}$. In particular, if there exists $\tilde{v} \in L^2$ which attains the infimum in (2.7) and $\mathcal{G}_{v^n}^0 = \{\phi\}$ then ϕ is uniquely characterized, because one can choose $v^n = \tilde{v}$ for all $n \in \mathbb{N}$.

The above functions ϕ, v, v^n, \tilde{v} are deterministic.

In order to formulate the sufficient condition for Laplace principle (equivalently, Large Deviations Principle). We also need the following basic assumption from [13].

(U) For any $\delta > 0$ and any $\phi \in C_T^d$ such that $I(\phi) < \infty$, there exists ϕ^δ uniquely characterized such that $\|\phi - \phi^\delta\|_T \leq \delta$ and $\|I(\phi) - I(\phi^\delta)\|_T \leq \delta$.

Lemma 2.11. [13] Assume that

Condition (i): For all $N > 0$, all $v \in \mathcal{A}_N$ and all families $\{v^\varepsilon\}_{\varepsilon > 0}$ in \mathcal{A}_N converging in distribution to v as ε tends to zero, $\{\mathcal{G}^\varepsilon(W + \frac{1}{\sqrt{\varepsilon}} \int_0^\cdot v_s^\varepsilon ds)\}$ is tight.

Condition (ii): The functional I defined by (2.7) has compact level sets, i.e. I is a good rate function.

Condition (iii): Assumption (U) holds.

Then the family $\{\mathcal{G}^\varepsilon(W)\}_{\varepsilon > 0}$ satisfies the Laplace principle and, by equivalence, satisfies the Large Deviations Principle with rate function I and speed ε^{-1} .

2.5. Technical lemmas. In order to prove our main results, we first state some assumptions used later.

(H1) There exist positive constant $\gamma > 0$ and $C_T > 0$ such that for any $0 \leq t < t' \leq T$

$$\begin{aligned} \int_0^t |K_i(t', s) - K_i(t, s)|^2 ds &\leq C_T |t' - t|^{2\gamma}, \quad t \in [0, T], \quad i = 1, 2, \\ \int_t^{t'} K_i^2(t', s) ds &\leq C_T |t - t'|^{2\gamma}, \quad t \in [0, T], \quad i = 1, 2. \end{aligned}$$

(H2) For any $x, y \in \mathbb{R}^d$, $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ and $t \in [0, T]$

$$|b(t, x, \mu) - b(t, y, \nu)| + \|\sigma(t, x, \mu) - \sigma(t, y, \nu)\| \leq L_1(|x - y| + \mathbb{W}_2(\mu, \nu)),$$

where $L_1 > 0$ is a positive constant.

(H3) For any $x \in \mathbb{R}^d$, $\mu \in \mathcal{P}_2(\mathbb{R}^d)$ and $t \in [0, T]$

$$|b(t, x, \mu)| + \|\sigma(t, x, \mu)\| \leq L_2(1 + |x| + \mathbb{W}_2(\mu, \delta_0)),$$

where $L_2 > 0$ is a positive constant.

Remark 2.12. (i) Note that by (H1) and Hölder's inequality, the kernel $K(t, s) := C_1 K_1(t, s) + C_2 K_2(t, s) + C_3 K_1^2(t, s) + C_4 K_2^2(t, s)$ with positive constants $C_i \geq 0, i = 1, 2, 3, 4$ satisfies the following properties:

$$t \mapsto \int_0^t K(t, s) ds \in L^\infty(0, T)$$

and

$$\limsup_{\varepsilon \downarrow 0} \left\| \int_{\cdot}^{\cdot + \varepsilon} K(\cdot + \varepsilon, s) ds \right\|_{L^\infty(0, T)} < 1.$$

Let R_n^K and R^K be defined by (2.3) and (2.4) in terms of K respectively. By (2.3), (2.5), (2.6) and (H1), when $n = 1$ we have

$$\int_0^T R_1^K(t, s) ds = \int_0^T K(t, s) ds < \infty,$$

and

$$\int_0^t R^K(t, s) ds < \infty.$$

This enables us to use Lemma 2.3.

(ii) It is clear that under the assumptions (H1)-(H3), there exists a unique solution $X_{t,\xi}^\varepsilon$ of (1.1) ($X_{t,\xi}^0$ of (1.2)) (Theorem 3.1 of [21]).

Lemma 2.13. Under the assumption (H1), the following inequalities hold for any adapted \mathbb{R}^d -valued process Y and $\mathbb{R}^{d \times m}$ -valued process Z :

(i) for $p \geq 2$ and $t \in [0, T]$,

$$\mathbb{E} \left[\left| \int_0^t K_1(t, s) Y_s ds \right|^p \right] \leq C_{p,T} \int_0^t K_1(t, s) \cdot \mathbb{E} |Y_s|^p ds,$$

(ii) for $p \geq 2$ and $t \in [0, T]$,

$$\mathbb{E} \left[\left| \int_0^t K_2(t, s) Z_s dW_s \right|^p \right] \leq C_{p,T} \int_0^t K_2^2(t, s) \cdot \mathbb{E} \|Z_s\|^p ds,$$

(iii) for $p \geq 2, v \in \mathcal{A}_N$ and $t \in [0, T]$,

$$\mathbb{E}\left[\left|\int_0^t K_1(t, s) Z_s v_s ds\right|^p\right] \leq C_{p,T,N} \int_0^t K_1^2(t, s) \cdot \mathbb{E}\|Z_s\|^p ds,$$

(iv) for $p \geq 1, t \in [0, T]$ and $h \geq 0$ with $t + h \leq T$,

$$\mathbb{E}\left[\left|\int_0^t (K_1(t+h, s) - K_1(t, s)) Y_s ds\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_1(t+h, s) Y_s ds\right|^p\right] \leq C_{p,T} h^{\gamma p} \sup_{t \in [0, T]} \mathbb{E}[|Y_t|^p],$$

(v) for $p \geq 2, t \in [0, T]$ and $h \geq 0$ with $t + h \leq T$,

$$\mathbb{E}\left[\left|\int_0^t (K_2(t+h, s) - K_2(t, s)) Z_s dW_s\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_2(t+h, s) Z_s dW_s\right|^p\right] \leq C_{p,T} h^{\gamma p} \sup_{t \in [0, T]} \mathbb{E}\|Z_t\|^p,$$

(vi) for $p \geq 2, v \in \mathcal{A}_N, t \in [0, T]$ and $h \geq 0$ with $t + h \leq T$,

$$\mathbb{E}\left[\left|\int_0^t (K_1(t+h, s) - K_1(t, s)) Z_s v_s ds\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_1(t+h, s) Z_s v_s ds\right|^p\right] \leq C_{T,N} h^{\gamma p} \sup_{t \in [0, T]} \mathbb{E}\|Z_t\|^p.$$

Proof. Let us show (i), (iii) and (iv) first. Take $p \geq 2$. Hölder's inequality show that

$$\begin{aligned} \mathbb{E}\left[\left|\int_0^t K_1(t, s) Y_s ds\right|^p\right] &\leq \int_0^t K_1(t, s) \cdot \mathbb{E}[|Y_s|^p] ds \cdot \left|\int_0^t K_1(t, s) ds\right|^{p-1} \\ &\leq C_{p,T} \int_0^t K_1(t, s) \cdot \mathbb{E}[|Y_s|^p] ds. \end{aligned}$$

For (iii), by Hölder's inequality and recall $v \in \mathcal{A}_N$ we have

$$\begin{aligned} \mathbb{E}\left[\left|\int_0^t K_1(t, s) Z_s v_s ds\right|^p\right] &\leq \left(\int_0^t v_s^2 ds\right)^{\frac{p}{2}} \cdot \mathbb{E}\left[\left|\int_0^t K_1^2(t, s) \cdot \|Z_s\|^2 ds\right|^{\frac{p}{2}}\right] \\ &\leq N^{\frac{p}{2}} \int_0^t K_1^2(t, s) \cdot \mathbb{E}\|\|Z_s\|^p\| ds \cdot \left|\int_0^t K_1^2(t, s) ds\right|^{\frac{p}{2}-1} \\ &\leq C_{p,T,N} \int_0^t K_1^2(t, s) \cdot \mathbb{E}\|\|Z_s\|^p\| ds. \end{aligned}$$

For (iv), by Minkowski's integral inequality we observe

$$\begin{aligned} &\mathbb{E}\left[\left|\int_0^t (K_1(t+h, s) - K_1(t, s)) Y_s ds\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_1(t+h, s) Y_s ds\right|^p\right] \\ &\leq \left(\int_0^t |K_1(t+h, s) - K_1(t, s)| \cdot [\mathbb{E}|Y_s|^p]^{\frac{1}{p}} ds\right)^p + \left(\int_t^{t+h} K_1(t+h, s) \cdot [\mathbb{E}|Y_s|^p]^{\frac{1}{p}} ds\right)^p \\ &\leq \sup_{t \in [0, T]} \mathbb{E}[|Y_s|^p] \left(\left(\int_0^t |K_1(t+h, s) - K_1(t, s)| ds\right)^p + \left(\int_t^{t+h} K_1(t+h, s) ds\right)^p \right) \\ &\leq C_{p,T} h^{\gamma p} \sup_{t \in [0, T]} \mathbb{E}[|Y_t|^p]. \end{aligned}$$

We next show (ii) and (v). For (ii), let $p \geq 2$, using BDG's inequality, Minkowski's integral inequality and Hölder's inequality we have

$$\begin{aligned} \mathbb{E}\left[\left|\int_0^t K_2(t, s) Z_s dW_s\right|^p\right] &\leq C_p \mathbb{E}\left[\left(\int_0^t K_2^2(t, s) \|Z_s\|^2 ds\right)^{\frac{p}{2}}\right] \\ &\leq C_p \int_0^t K_2^2(t, s) \cdot \mathbb{E}\|\|Z_s\|^p\| ds \cdot \left|\int_0^t K_2^2(t, s) ds\right|^{\frac{p}{2}-1} \end{aligned}$$

$$\leq C_{p,T} \int_0^t K_2^2(t,s) \cdot \mathbb{E}[\|Z_s\|^p] ds.$$

For (v), take $p \geq 2$. By the BDG's inequality and Minkowski's inequality, we obtain

$$\begin{aligned} & \mathbb{E}\left[\left|\int_0^t (K_2(t+h,s) - K_2(t,s)) Z_s dW_s\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_2(t+h,s) Z_s dW_s\right|^p\right] \\ & \leq C_p \mathbb{E}\left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 \cdot \|Z_s\|^2 ds\right)^{\frac{p}{2}} + C_p \mathbb{E}\left(\int_t^{t+h} K_2^2(t+h,s) \cdot \|Z_s\|^2 ds\right)^{\frac{p}{2}} \\ & \leq C_p \left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 \cdot \mathbb{E}[\|Z_s\|^p]^{\frac{2}{p}} ds\right)^{\frac{p}{2}} + C_p \left(\int_t^{t+h} K_2^2(t+h,s) \mathbb{E}[\|Z_s\|^p]^{\frac{2}{p}} ds\right)^{\frac{p}{2}} \\ & \leq C_p \sup_{t \in [0,T]} \mathbb{E}[\|Z_t\|^p] \left[\left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 ds\right)^{\frac{p}{2}} + \left(\int_t^{t+h} K_2^2(t+h,s) ds\right)^{\frac{p}{2}}\right] \\ & \leq C_{p,T} h^{\gamma p} \sup_{t \in [0,T]} \mathbb{E}[\|Z_t\|^p]. \end{aligned}$$

Finally, we show (vi). By Hölder's inequality and recall $v \in \mathcal{A}_N$ we have

$$\begin{aligned} & \mathbb{E}\left[\left|\int_0^t (K_2(t+h,s) - K_2(t,s)) Z_s v_s ds\right|^p\right] + \mathbb{E}\left[\left|\int_t^{t+h} K_2(t+h,s) Z_s v_s ds\right|^p\right] \\ & \leq \left(\int_0^t v_s^2 ds\right)^{\frac{p}{2}} \cdot \mathbb{E}\left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 \cdot \|Z_s\|^2 ds\right)^{\frac{p}{2}} \\ & \quad + \left(\int_0^t v_s^2 ds\right)^{\frac{p}{2}} \cdot \mathbb{E}\left(\int_t^{t+h} K_2^2(t+h,s) \cdot \|Z_s\|^2 ds\right)^{\frac{p}{2}} \\ & \leq C_N \left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 \cdot \mathbb{E}[\|Z_s\|^p]^{\frac{2}{p}} ds\right)^{\frac{p}{2}} + \left(\int_t^{t+h} K_2^2(t+h,s) \mathbb{E}[\|Z_s\|^p]^{\frac{2}{p}} ds\right)^{\frac{p}{2}} \\ & \leq C_N \sup_{t \in [0,T]} \mathbb{E}[\|Z_t\|^p] \left(\left(\int_0^t |K_2(t+h,s) - K_2(t,s)|^2 ds\right)^{\frac{p}{2}} + \left(\int_t^{t+h} K_2^2(t+h,s) ds\right)^{\frac{p}{2}}\right) \\ & \leq C_{T,N} h^{\gamma p} \sup_{t \in [0,T]} \mathbb{E}[\|Z_t\|^p], \end{aligned}$$

which completes the proof. \square

Lemma 2.14. *Under the assumptions **(H1)**-**(H3)**, it follows that for $p \geq 1, \xi, \eta \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E}|X_{t,\xi}^\varepsilon|^p \leq C_{p,T,L_2}(1 + |\xi|^p),$$

and

$$\sup_{t \in [0,T]} \mathbb{E}|X_{t,\xi}^\varepsilon - X_{t,\eta}^\varepsilon|^p \leq C_{p,T,L_1}|\xi - \eta|^p,$$

where the constants C_{p,T,L_1} and C_{p,T,L_2} are independent of ε .

Proof. We can assume $p \geq 2$, which is sufficient for $p \geq 1$ by Lyapunov inequality. By Lemma 2.13 (i)-(ii), **(H1)**, **(H3)** and Lemma 2.6 in [8], for $0 \leq \varepsilon < 1$ we have

$$\begin{aligned} & \mathbb{E}|X_{t,\xi}^\varepsilon|^p \\ & = \mathbb{E}\left|\xi + \int_0^t K_1(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) ds + \sqrt{\varepsilon} \int_0^t K_2(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s\right|^p \\ & \leq 3^{p-1}|\xi|^p + 3^{p-1}\mathbb{E}\left|\int_0^t K_1(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) ds\right|^p + 3^{p-1}\mathbb{E}\left|\sqrt{\varepsilon} \int_0^t K_2(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s\right|^p \end{aligned}$$

$$\begin{aligned}
&\leq 3^{p-1}|\xi|^p + C_{p,T} \int_0^t K_1(t,s) \cdot \mathbb{E}|b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})|^p ds + C_{p,T} \int_0^t K_2^2(t,s) \cdot \mathbb{E}\|\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})\|^p ds \\
&\leq 3^{p-1}|\xi|^p + C_{p,T,L_2} \int_0^t K_1(t,s) \cdot \mathbb{E}\left|1 + |X_{s,\xi}^\varepsilon| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0)\right|^p ds \\
&\quad + C_{p,T,L_2} \int_0^t K_2^2(t,s) \cdot \mathbb{E}\left|1 + |X_{s,\xi}^\varepsilon| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0)\right|^p ds \\
&\leq 3^{p-1}|\xi|^p + C_{p,T,L_2} \int_0^t K_1(t,s)(1 + \mathbb{E}|X_{s,\xi}^\varepsilon|^p) ds + C_{p,T,L_2} \int_0^t K_2^2(t,s)(1 + \mathbb{E}|X_{s,\xi}^\varepsilon|^p) ds \\
&\leq 3^{p-1}|\xi|^p + C_{p,T,L_2} + C_{p,T,L_2} \int_0^t K_{1,2}(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon|^p ds,
\end{aligned}$$

where $K_{1,2}^*(t,s) := C_{p,T,L_2} \cdot (K_1(t,s) + K_2^2(t,s))$ and the constant C_{p,T,L_2} is independent of ε .

Let $R^{K_{1,2}^*}$ be defined by (2.4) in terms of $K_{1,2}$. By Remark 2.12, Lemma 2.3 and (2.6), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}
\mathbb{E}|X_{t,\xi}^\varepsilon|^p &\leq 3^{p-1}|\xi|^p + C_{p,T,L_2} + \int_0^t R^{K_{1,2}^*}(s) \cdot (3^{p-1}|\xi|^p + C_{p,T,L_2}) ds \\
&\leq C_{p,T,L_2}(1 + |\xi|^p).
\end{aligned}$$

Similarly, we can prove

$$\mathbb{E}|X_{t,\xi}^\varepsilon - X_{t,\eta}^\varepsilon|^p \leq C_{p,T,L_1}|\xi - \eta|^p.$$

□

Proceeding as in the proof of Lemma 2.14, we have the following Lemma associated with $X_{t,\xi}^0$.

Lemma 2.15. *Under the assumptions **(H1)-(H3)**, it follows that for $p \geq 1, \xi, \eta \in \mathbb{R}^d$*

$$\sup_{t \in [0, T]} |X_{t,\xi}^0|^p \leq C_{p,T,L_2}(1 + |\xi|^p),$$

and

$$\sup_{t \in [0, T]} |X_{t,\xi}^0 - X_{t,\eta}^0|^p \leq C_{p,T,L_1}|\xi - \eta|^p,$$

where the constants C_{p,T,L_1} and C_{p,T,L_2} are independent of ε .

Lemma 2.16. *Denote $Q_t^\varepsilon := \frac{X_{t,\xi}^\varepsilon - X_{t,\xi}^0}{\sqrt{\varepsilon h(\varepsilon)}}$. Assume that **(H1)-(H3)** hold. Then for $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0, T]} \mathbb{E}|Q_t^\varepsilon|^p \leq \frac{C_{p,T,L_1,L_2}}{|h(\varepsilon)|^p}(1 + |\xi|^p),$$

where the constant C_{p,T,L_1,L_2} is independent of ε .

Proof. Write

$$Q_t^\varepsilon = \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon h(\varepsilon)}} ds + \frac{1}{h(\varepsilon)} \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s. \quad (2.8)$$

By Lemma 2.13 (i)-(ii) and **(H1)-(H3)**, we have

$$\mathbb{E}|Q_t^\varepsilon|^p$$

$$= \mathbb{E}\left|\int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon h(\varepsilon)}} ds + \frac{1}{h(\varepsilon)} \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s\right|^p$$

$$\begin{aligned}
&\leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)} ds \right|^p + 2^{p-1} \mathbb{E} \left| \frac{1}{h(\varepsilon)} \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right|^p \\
&\leq C_{p,T} \int_0^t K_1(t,s) \cdot \mathbb{E} \left| \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)} \right|^p ds \\
&\quad + \frac{C_{p,T}}{|h(\varepsilon)|^p} \int_0^t K_2^2(t,s) \cdot \mathbb{E} \|\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})\|^p ds \\
&\leq C_{p,T,L_1} \int_0^t K_1(t,s) \cdot \mathbb{E} \left| \frac{|X_{s,\xi}^\varepsilon - X_{s,\xi}^0| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)} \right|^p ds \\
&\quad + \frac{C_{p,T,L_2}}{|h(\varepsilon)|^p} \int_0^t K_2^2(t,s) \cdot \mathbb{E} \left| (1 + |X_{s,\xi}^\varepsilon| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0)) \right|^p ds \\
&\leq C_{p,T,L_1} \int_0^t K_1(t,s) \cdot \mathbb{E} |Q_{s,\xi}^\varepsilon|^p ds + \frac{C_{p,T,L_2}}{|h(\varepsilon)|^p} \int_0^t K_2^2(t,s) \cdot (1 + \mathbb{E} |X_{s,\xi}^\varepsilon|^p) ds \\
&\leq \frac{C_{p,T,L_2}}{|h(\varepsilon)|^p} (1 + |\xi|^p) + C_{p,T,L_1} \int_0^t K_1(t,s) \cdot \mathbb{E} |Q_{s,\xi}^\varepsilon|^p ds,
\end{aligned}$$

where in the last step Lemma 2.14 is used.

Let $R^{\hat{K}_1}$ be defined by (2.4) in terms of $\hat{K}_1 := C_{p,T,L_1} K_1(t,s)$. By Remark 2.12, Lemma 2.3 and (2.6), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}
\mathbb{E} |Q_t^\varepsilon|^p &\leq \frac{C_{p,T,L_2}}{|h(\varepsilon)|^p} (1 + |\xi|^p) + \int_0^t R^{\hat{K}_1}(t,s) \cdot \frac{C_{p,T,L_2}}{|h(\varepsilon)|^p} (1 + |\xi|^p) ds \\
&\leq \frac{C_{p,T,L_1,L_2}}{|h(\varepsilon)|^p} (1 + |\xi|^p).
\end{aligned}$$

This proves the lemma. \square

Lemma 2.17. *Let $M : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R} \rightarrow \mathbb{R}^d$, $N, G : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{P}_2(\mathbb{R}^d) \times \mathbb{R} \rightarrow \mathbb{R}^d$ satisfy*

$$M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \leq C(1 + |\Lambda_{s,x}^{\varepsilon,v}| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0) + \frac{1}{\sqrt{\varepsilon} h(\varepsilon)} \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0})),$$

$$N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \vee G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \leq C(1 + |\Lambda_{s,x}^{\varepsilon,v}| + |X_{s,\xi}^0| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0))$$

for some finite constant C . Let $\Lambda_{t,x}^{\varepsilon,v}$ satisfy

$$\begin{aligned}
\Lambda_{t,x}^{\varepsilon,v} &= x + \int_0^t K_1(t,s) M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) ds \\
&\quad + \int_0^t K_1(t,s) N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) v_s ds + \int_0^t K_2(t,s) G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) dW_s,
\end{aligned} \tag{2.9}$$

where $X_{s,\xi}^0$ and $X_{s,\xi}^\varepsilon$ are defined as before. Then, for $p > 2, N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$

$$\sup_{t \in [0, T]} \mathbb{E} |\Lambda_{t,x}^{\varepsilon,v}|^p \leq C_{p,T,N,L_2} (1 + |\xi|^p),$$

where the constant C_{p,T,N,L_2} is independent of $\varepsilon \in (0, 1]$, and $v \in \mathcal{A}_N$ (but it may depend on \mathcal{A}_N or N).

Proof. Since $\Lambda_{t,\xi}^{\varepsilon,v}$ satisfies (2.9), we have

$$\mathbb{E}[|\Lambda_{t,x}^{\varepsilon,v}|^p]$$

$$\begin{aligned}
&= \mathbb{E} \left[\left| x + \int_0^t K_1(t, s) M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) ds \right. \right. \\
&\quad \left. \left. + \int_0^t K_1(t, s) N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) v_s ds + \int_0^t K_2(t, s) G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) dW_s \right|^p \right] \\
&\leq 4^{p-1} |x|^p + 4^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) ds \right|^p \\
&\quad + 4^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) v_s ds \right|^p \\
&\quad + 4^{p-1} \mathbb{E} \left| \int_0^t K_2(t, s) G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) dW_s \right|^p \\
&:= 4^{p-1} |x|^p + I_1 + I_2 + I_3.
\end{aligned}$$

For the term I_1 , by Lemma 2.13-(i), **(H1)-(H3)**, Lemmas 2.14 and 2.16 we have

$$\begin{aligned}
I_1 &\leq C_{p,T} \int_0^t K_1(t, s) \cdot \mathbb{E} \left| M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \right|^p ds \\
&\leq C_{p,T} \int_0^t K_1(t, s) \cdot \mathbb{E} \left| 1 + |\Lambda_{s,x}^{\varepsilon,v}| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0) + \frac{1}{\sqrt{\varepsilon} h(\varepsilon)} \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}) \right|^p ds \\
&\leq C_{p,T} \int_0^t K_1(t, s) \cdot (1 + \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p + \mathbb{E}|X_{s,\xi}^\varepsilon|^p + \mathbb{E}|Q_{t,\xi}^\varepsilon|^p) ds \\
&\leq C_{p,T,L_1,L_2} (1 + |\xi|^p) + C_{p,T} \int_0^t K_1(t, s) \cdot \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p ds.
\end{aligned}$$

For the term I_2 , by Lemma 2.13-(iii), **(H1)-(H3)**, Lemmas 2.14 and 2.15 we have

$$\begin{aligned}
I_2 &\leq C_{p,T,N} \int_0^t K_1^2(t, s) \cdot \mathbb{E} \left\| N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \right\|^p ds \\
&\leq C_{p,T,N} \int_0^t K_1^2(t, s) \cdot \mathbb{E} \left| 1 + |\Lambda_{s,x}^{\varepsilon,v}| + |X_{s,\xi}^0| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0) \right|^p ds \\
&\leq C_{p,T,N} \int_0^t K_1^2(t, s) \cdot (1 + \mathbb{E}|X_{s,\xi}^{\varepsilon,v}|^p + \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p + \mathbb{E}|X_{s,\xi}^0|^p) ds \\
&\leq C_{p,T,N,L_2} (1 + |\xi|^p) + C_{p,T,N} \int_0^t K_1^2(t, s) \cdot \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p ds.
\end{aligned}$$

For the term I_3 , applying 2.13-(ii), **(H1)-(H3)**, Lemmas 2.14 and 2.15 yields

$$\begin{aligned}
I_3 &\leq C_{p,T} \int_0^t K_2^2(t, s) \cdot \mathbb{E} \left| G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \right|^p ds \\
&\leq C_{p,T} \int_0^t K_2^2(t, s) \cdot \mathbb{E} \left| 1 + |\Lambda_{s,x}^{\varepsilon,v}| + |X_{s,\xi}^0| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0) \right|^p ds \\
&\leq C_{p,T} \int_0^t K_2^2(t, s) \cdot (1 + \mathbb{E}|X_{s,\xi}^{\varepsilon,v}|^p + \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p + \mathbb{E}|X_{s,\xi}^0|^p) ds \\
&\leq C_{p,T,L_2} (1 + |\xi|^p) + C_{p,T} \int_0^t K_2^2(t, s) \cdot \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p ds.
\end{aligned}$$

Therefore

$$\mathbb{E}|\Lambda_{t,x}^{\varepsilon,v}|^p \leq C_{p,T,N,L_1,L_2} (1 + |\xi|^p) + \int_0^t K_{1,2}^*(t, s) \cdot \mathbb{E}|\Lambda_{s,x}^{\varepsilon,v}|^p ds,$$

where $K_{1,2}^*(t, s) := C_{p,T} \cdot (K_1(t, s) + K_2^2(t, s)) + C_{p,T,N} K_2^2(t, s)$ and the constants $C_{p,T}, C_{p,T,N}$ are independent of ε .

Let $R^{K_{1,2}}$ be defined by (2.4) in terms of $K_{1,2}^*$. By Remark 2.12, Lemma 2.3 and (2.6), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}\mathbb{E}|\Lambda_{t,x}^{\varepsilon,v}|^p &\leq C_{p,T,N,L_1,L_2}(1 + |\xi|^p) + \int_0^t R^{K_{1,2}}(t,s) \cdot C_{p,T,N,L_1,L_2}(1 + |\xi|^p) ds \\ &\leq C_{p,T,N,L_1,L_2}(1 + |\xi|^p).\end{aligned}$$

The proof of the lemma is then completed. \square

Lemma 2.18. *Assume that $\Lambda_{t,x}^{\varepsilon,v}$ satisfies (2.9) and (H1)-(H3) hold, if $p > 2 \vee \frac{2}{\gamma}$, $t, s \in [0, T]$, $N > 0$, $\xi \in \mathbb{R}^d$ and $\{v^\varepsilon\}_{\varepsilon>0}$ is a family in \mathcal{A}_N , in addition, assume there exist a constant C such that*

$$M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \leq C(1 + |\Lambda_{s,x}^{\varepsilon,v}| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0) + \frac{1}{\sqrt{\varepsilon}h(\varepsilon)} \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0})),$$

$$N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \vee G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) \leq C(1 + |\Lambda_{s,x}^{\varepsilon,v}| + |X_{s,\xi}^0| + \mathbb{W}_2(\mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_0)).$$

Then $\Lambda^{\varepsilon,v^\varepsilon}$ admits a version which is Hölder continuous on $[0, T]$ of any order $\alpha < \frac{\gamma}{2} - \frac{1}{p}$, uniformly for all $\varepsilon > 0$. Denoting again this version by $\Lambda^{\varepsilon,v^\varepsilon}$, one can find a $\varepsilon_0 > 0$ such that

$$\sup_{0 < \varepsilon < \varepsilon_0} \mathbb{E} \left[\left(\sup_{0 \leq s < t \leq T} \frac{|\Lambda_{t,x}^{\varepsilon,v^\varepsilon} - \Lambda_{s,x}^{\varepsilon,v^\varepsilon}|}{|t-s|^\alpha} \right)^p \right] \leq C_{\alpha,p,T,N,L_2,\xi},$$

for all $\alpha \in [0, \frac{\gamma}{2} - \frac{1}{p}]$. Moreover, the family of random processes $\{\Lambda^{\varepsilon,v^\varepsilon}\}_{\varepsilon>0}$ is tight in C_T .

Proof. For any $s < t$, we have

$$\begin{aligned}&\Lambda_{t,x}^{\varepsilon,v^\varepsilon} - \Lambda_{s,x}^{\varepsilon,v^\varepsilon} \\ &= \int_s^t K_1(t,u) M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v^\varepsilon}, h(\varepsilon)) du + \int_s^t K_1(t,u) N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) v_u du \\ &\quad + \int_s^t K_2(t,u) G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) dW_u \\ &\quad + \int_0^s (K_1(t,u) - K_1(s,u)) \cdot M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v^\varepsilon}, h(\varepsilon)) du \\ &\quad + \int_0^s (K_1(t,u) - K_1(s,u)) \cdot N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) v_u du \\ &\quad + \int_0^s (K_2(t,u) - K_2(s,u)) \cdot G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) dW_u \\ &:= J_1 + J_2 + J_3 + J_4 + J_5 + J_6.\end{aligned}$$

Note that under the assumptions on M, N, G , Lemmas 2.14, 2.15, 2.16 and 2.17 give us

$$\sup_{s \in [0,T]} \mathbb{E}|M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v^\varepsilon}, h(\varepsilon))|^p \leq C_{p,T,N,L_1,L_2}(1 + |\xi|^p),$$

$$\sup_{s \in [0,T]} \mathbb{E}\|N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon))\|^p \leq C_{p,T,N,L_2}(1 + |\xi|^p),$$

$$\sup_{s \in [0,T]} \mathbb{E}\|G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon))\|^p \leq C_{p,T,N,L_2}(1 + |\xi|^p).$$

For the terms J_1 and J_4 , combining Lemma 2.13-(iv), Lemmas 2.14-2.17 we have

$$\mathbb{E}|J_1|^p + \mathbb{E}|J_4|^p \leq C_{p,T}|t-s|^{\gamma p} \sup_{s \in [0,T]} \mathbb{E}|M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v^\varepsilon}, h(\varepsilon))|^p$$

$$\leq C_{p,T,N,L_1,L_2}(1+|\xi|^p)|t-s|^{\gamma p}.$$

For the terms J_2 and J_5 , by Lemma 2.13-(vi), Lemmas 2.14-2.17, we have

$$\begin{aligned}\mathbb{E}|J_2|^p + \mathbb{E}|J_5|^p &\leq C_{T,N}|t-s|^{\gamma p} \sup_{s \in [0,T]} \mathbb{E}\|N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon))\|^p \\ &\leq C_{p,T,N,L_2}(1+|\xi|^p)|t-s|^{\gamma p}.\end{aligned}$$

For the terms J_3 and J_6 , by Lemma 2.13-(v), Lemmas 2.14-2.17, we have

$$\begin{aligned}\mathbb{E}|J_3|^p + \mathbb{E}|J_6|^p &\leq C_{p,T}|t-s|^{\gamma p} \sup_{t \in [0,T]} \mathbb{E}\|G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon))\|^p \\ &\leq C_{p,T,N,L_2}(1+|\xi|^p)|t-s|^{\gamma p}.\end{aligned}$$

Combining the above estimations, we have

$$\mathbb{E}|\Lambda_{t,x}^{\varepsilon,v} - \Lambda_{s,x}^{\varepsilon,v}|^p \leq C_{p,T,N,L_1,L_2}(1+|\xi|^p)|t-s|^{\gamma p}.$$

This gives the bound in the lemma. The existence of a continuous version now follows from the Kolmogorov's continuity theorem (e.g. Theorem I. 2.1 in [30]). Furthermore, Aldous theorem ([5, Theorem 16.10]) can be used to conclude that the sequence $\{X_{\cdot,\xi}^{\varepsilon,v}\}_{\varepsilon>0}$ is tight. \square

3. LARGE DEVIATION PRINCIPLE FOR VOLTERRA TYPE MCKEAN-VLASOV SDES

In this section, we will use the weak convergence approach to obtain the LDP for Volterra type McKean-Vlasov SDEs.

It is clear that under the assumptions **(H1)-(H3)**, there exists a unique strong solution $X_{t,\xi}^\varepsilon$ of (1.1) (Theorem 3.1 of [21]). As ε tends to zero, $X_{\cdot,\xi}^\varepsilon$ will converge to $X_{\cdot,\xi}^0$, where $X_{\cdot,\xi}^0$ satisfies (1.2). In order to prove Laplace principle for (1.1), we take

$$E = C_T^d := C([0, T], \mathbb{R}^d), \quad \mathcal{G}^\varepsilon(W) := X_{\cdot,\xi}^\varepsilon,$$

where $C([0, T], \mathbb{R}^d)$ denotes the space of continuous functions from $[0, T]$ to \mathbb{R}^d equipped with the supremum norm $\|\phi\|_T := \sup_{t \in [0, T]} |\phi_t|$. Then by Girsanov's theorem, the shifted version $X_{\cdot,\xi}^{\varepsilon,v} := \mathcal{G}^\varepsilon(W + \frac{1}{\sqrt{\varepsilon}} \int_0^\cdot v_s ds)$ is the unique strong solution of the controlled equation under \mathbb{P} :

$$\begin{aligned}X_{t,\xi}^{\varepsilon,v} &= \xi + \int_0^t K_1(t, s)b(s, X_{s,\xi}^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon})ds + \int_0^t K_1(t, s)\sigma(s, X_{s,\xi}^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon})v_s ds \\ &\quad + \sqrt{\varepsilon} \int_0^t K_2(t, s)\sigma(s, X_{s,\xi}^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon})dW_s.\end{aligned}\tag{3.10}$$

Under some appropriate conditions, we will prove that $X_{t,\xi}^{\varepsilon,v}$ will converge to ϕ_t as $\varepsilon \rightarrow 0$, where ϕ_t is the unique solution of the following deterministic Volterra equation

$$\phi_t = \xi + \int_0^t K_1(t, s)b(s, \phi_s, \delta_{X_{s,\xi}^0})ds + \int_0^t K_1(t, s)\sigma(s, \phi_s, \delta_{X_{s,\xi}^0})v_s ds.\tag{3.11}$$

We now state the results of LDP.

Theorem 3.1. *Under the assumptions **(H1)-(H3)**, the family $\{X_{\cdot,\xi}^\varepsilon\}_{\varepsilon>0}$, the unique solution of (1.1), satisfies the Large Deviation Principle with rate function (2.7) and speed ε^{-1} , where \mathcal{G}_v^0 corresponds to the unique solution of the limiting equation (3.11).*

We shall use Lemma 2.11 to prove this theorem. Note that the third condition in Lemma 2.11 is automatically satisfied if the limiting equation (3.11) has a unique solution. We shall verify the first two conditions in Lemma 2.11, which will be given by Lemmas 3.3 and 3.8. But before this we need some preparation.

Lemma 3.2. *Assume that **(H1)-(H3)** hold, then there exists a unique solution ϕ_t such that for almost all $t \geq 0$, $N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$, some constant C_{T,N,L_2}*

$$\sup_{t \in [0,T]} |\phi_t| \leq C_{T,N,L_2}(1 + |\xi|), \quad (3.12)$$

for almost all $t \in [0, T]$.

Lemma 3.2 implies that the solution of (3.11) is unique. By Remark 3.12 in [13], we only need to check Conditions (i) and (ii) in Theorem 2.11.

Let

$$x := \xi, \Lambda_{t,\xi}^{\varepsilon,v} := X_{t,\xi}^{\varepsilon,v}, M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := b(s, X_{s,\xi}^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon})$$

and

$$N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := \sigma(s, X_{s,\xi}^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}), G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := \sqrt{\varepsilon}\sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)).$$

Then Lemmas 2.17 and 2.18 give us that

Lemma 3.3. *Under the assumptions **(H1)-(H3)**, it follows that for $p > 2, N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$. Hence $\{X^{n,v}, v^n\}_{n \geq 0}$ converges almost surely in the product topology on $C_T^d \times \mathcal{S}_N$, and the limit is the $C_T^d \times \mathcal{S}_N$ -valued random variable (ϕ, v) in a possibly different probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ as n tends to $+\infty$.*

$$\sup_{t \in [0,T]} \mathbb{E}|X_{t,\xi}^{\varepsilon,v}|^p \leq C_{p,T,N,L_2}(1 + |\xi|^p),$$

where the constant C_{p,T,N,L_2} is independent of $\varepsilon \in (0, 1], v \in \mathcal{A}_N$ (but it depends on \mathcal{A}_N or N).

Lemma 3.4. *Assume that **(H1)-(H3)** hold, if $p > 2 \vee \frac{2}{\gamma}$, $t, s \in [0, T]$, $N > 0, \xi \in \mathbb{R}^d$ and $\{v^\varepsilon\}_{\varepsilon > 0}$ is a family in \mathcal{A}_N , then $X^{\varepsilon,v^\varepsilon}$ admits a version which is Hölder continuous on $[0, T]$ of any order $\alpha < \frac{\gamma}{2} - \frac{1}{p}$, uniformly for all $\varepsilon > 0$. Denoting again this version by $X^{\varepsilon,v^\varepsilon}$, one has for all $\varepsilon > 0$ small enough*

$$\mathbb{E} \left[\left(\sup_{0 \leq s < t \leq T} \frac{|X_{t,\xi}^{\varepsilon,v^\varepsilon} - X_{s,\xi}^{\varepsilon,v^\varepsilon}|}{|t-s|^\alpha} \right)^p \right] \leq C_{p,T,N,L_2,\xi},$$

for all $\alpha \in [0, \frac{\gamma}{2} - \frac{1}{p}]$. Moreover, the family of random processes $\{X_{\cdot,\xi}^{\varepsilon,v^\varepsilon}\}_{\varepsilon > 0}$ is tight in C_T .

By a similar procedure, we can get that the solution ϕ_t of (3.11) also have Hölder continuous path, that is

Lemma 3.5. *Assume that **(H1)-(H3)** hold, for $t, s \in [0, T]$, $N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$, then the solution ϕ_t of (3.11) has for all $\varepsilon > 0$ small enough*

$$|\phi_t - \phi_s| \leq C_{T,N,L_2}(1 + |\xi|)|t - s|^\gamma.$$

Since $X_{t,\xi}^\varepsilon - X_{t,\xi}^0 = Q_{t,\xi}^\varepsilon|_{h(\varepsilon)=\frac{1}{\sqrt{\varepsilon}}}$, so Lemma 2.16 is simplified as

Lemma 3.6. *Under the assumptions **(H1)-(H3)**, it follows that for $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E}|X_{t,\xi}^\varepsilon - X_{t,\xi}^0|^p \leq \varepsilon^{\frac{p}{2}} C_{p,T,L_1,L_2}(1 + |\xi|^p),$$

where the constant C_{p,T,L_1,L_2} is independent of ε .

Lemma 3.7. *The set \mathcal{G}_v^0 from Definition 2.9 is characterized by*

$$\mathcal{G}_v^0 = \left\{ \phi : [0, T] \rightarrow \mathbb{R}^d \mid \phi_t = \xi + \int_0^t K_1(t, s) b(s, \phi_s, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, \phi_s, \delta_{X_{s,\xi}^0}) v_s ds \right\}.$$

Proof. The proof of this lemma follows the technique in Theorem 1 [6] or Lemma 3.17 of [13]. Before we characterize \mathcal{G}_v^0 , we first identify $\mathcal{G}_{v,N}^0$. Recall the definition 2.9 of $\mathcal{G}_{v,N}^0$. For $N \in \mathbb{N}$ and $v \in \mathcal{A}_N$, consider a subsequence $\{\varepsilon_n\}_{n \in \mathbb{N}} \subset [0, \infty)$ with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and a sequence $\{v^{\varepsilon_n}\}_{n \in \mathbb{N}} \subset \mathcal{A}_N$ such that $\lim_{n \rightarrow \infty} v^{\varepsilon_n} = v$ in distribution, and assume that $X^{n,v} := X^{\varepsilon_n, v^{\varepsilon_n}}$ converges in distribution to some random variable ϕ with values in C_T^d . To simplify the representation, we define $v^n := v^{\varepsilon_n}$, $X^{\varepsilon_n} := X^n$.

By Skorohod representation theorem ([9, Theorem 3.15]) we can work with almost sure convergence for the purpose of identifying the limit. Hence $\{X^{n,v}, v^n\}_{n \geq 0}$ converges almost surely in the product topology on $C_T^d \times \mathcal{S}_N$, and the limit is a $C_T^d \times \mathcal{S}_N$ -valued random variable (ϕ, v) in a possibly different probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ as n tends to $+\infty$. The convergence of the couple also takes place in distribution. For $t \in [0, T]$, define $\Phi_t : C_T^d \times \mathcal{S}_N \rightarrow \mathbb{R}$ as

$$\Phi_t(\omega, f) := \left| \omega_t - \xi - \int_0^t K_1(t, s) \cdot (b(s, \omega_s, \delta_{X_{s,\xi}^0}) + \sigma(s, \omega_s, \delta_{X_{s,\xi}^0}) f_s) ds \right| \wedge 1.$$

It is clear that Φ_t is bounded, and we also can show Φ_t is continuous. Let $\omega^n \rightarrow \omega$ in C_T^d and $f^n \rightarrow f$ in \mathcal{S}_N with respect the weak topology.

For $\omega, \omega^n \in C_T^d$, $f, f^n \in \mathcal{S}_N$, by Lemma 2.15 and (H1)-(H3) we have that

$$\begin{aligned} |\Phi_t(\omega, f) - \Phi_t(\omega^n, f^n)| &\leq |\omega_t - \omega_t^n| + \int_0^t K_1(t, s) \cdot |b(s, \omega_s, \delta_{X_{s,\xi}^0}) - b(s, \omega_s^n, \delta_{X_{s,\xi}^0})| ds \\ &\quad + \int_0^t K_1(t, s) \cdot |(\sigma(s, \omega_s, \delta_{X_{s,\xi}^0}) - \sigma(s, \omega_s^n, \delta_{X_{s,\xi}^0})) f_s^n + (f_s - f_s^n) \sigma(s, \omega_s, \delta_{X_{s,\xi}^0})| ds \\ &\leq \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + \left(\int_0^t K_1^2(t, s) ds \right)^{\frac{1}{2}} \cdot \left(\int_0^t |b(s, \omega_s, \delta_{X_{s,\xi}^0}) - b(s, \omega_s^n, \delta_{X_{s,\xi}^0})|^2 ds \right)^{\frac{1}{2}} \\ &\quad + \left(\int_0^t |f_s^n|^2 ds \right)^{\frac{1}{2}} \cdot \left(\int_0^t K_1^2(t, s) \cdot \|\sigma(s, \omega_s, \mathcal{L}_{X_{s,\xi}^0}) - \sigma(s, \omega_s^n, \mathcal{L}_{X_{s,\xi}^0})\|^2 ds \right)^{\frac{1}{2}} \\ &\quad + \sup_{t \in [0, T]} \|\sigma(t, \omega_t, \mathcal{L}_{X_{t,\xi}^0})\| \int_0^t K_1(t, s) \cdot |f_s - f_s^n| ds \\ &\leq \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{T,L_1} \left(\int_0^t |\omega_s - \omega_s^n|^2 ds \right)^{\frac{1}{2}} + C_{N,L_1} \left(\int_0^t K_1^2(t, s) \cdot |\omega_s - \omega_s^n|^2 ds \right)^{\frac{1}{2}} \\ &\quad + C_{L_2} \sup_{t \in [0, T]} (1 + |\omega_t| + |X_t^0|) \int_0^t K_1(t, s) \cdot |f_s - f_s^n| ds \\ &\leq \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{T,L_1} \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{N,T,L_1} \sup_{t \in [0, T]} |\omega_t - \omega_t^n| \\ &\quad + C_{T,L_2} (1 + \sup_{t \in [0, T]} \omega_t) \int_0^t K_1(t, s) \cdot |f_s - f_s^n| ds \\ &\leq C_{N,T,L_1} \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{L_2} (1 + \sup_{t \in [0, T]} \omega_t) \int_0^t K_1(t, s) \cdot |f_s - f_s^n| ds. \end{aligned}$$

Since f^n tends to f weakly in L^2 and $\lim_{n \rightarrow \infty} |\omega_t - \omega_t^n| = 0$, so we get $\Phi_t(f^n, \omega^n)$ tends to $\Phi_t(f, \omega)$ as n tends to ∞ , which implies Φ_t is continuous and therefore

$$\lim_{n \rightarrow \infty} \mathbb{E}[\Phi_t(X^{n,v}, v^n)] = \mathbb{E}[\Phi_t(\phi, v)].$$

It remains to prove that the right hand of above equality is actually equal to zero. By BDG's inequality, Minkowski's inequality, Lemmas 2.14, 3.3 and 3.6 we have

$$\begin{aligned} \mathbb{E}[\Phi_t(X^{n,v}, v^n)] &\leq \mathbb{E}\left|\int_0^t K_1(t, s) \cdot (b(s, X_s^{n,v}, \mathcal{L}_{X_s^n}) - b(s, X_s^{n,v}, \delta_{X_{s,\xi}^0})) ds\right| \\ &\quad + \mathbb{E}\left|\int_0^t K_1(t, s) \cdot (\sigma(s, X_s^{n,v}, \mathcal{L}_{X_s^n}) - \sigma(s, X_s^{n,v}, \delta_{X_{s,\xi}^0})) v_s^n ds\right| \\ &\quad + \sqrt{\varepsilon_n} \mathbb{E}\left|\int_0^t K_2(t, s) \cdot \sigma(s, X_s^{n,v}, \mathcal{L}_{X_s^n}) dW_s\right| \\ &\leq L_1 \mathbb{E}\int_0^t K_1(t, s) \cdot \mathbb{W}_2(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0}) ds \\ &\quad + \left(\int_0^t |v_s^n|^2 ds\right)^{\frac{1}{2}} \cdot \mathbb{E}\left(\int_0^t K_1^2(t, s) \cdot \|\sigma(s, X_s^{n,v}, \mathcal{L}_{X_s^n}) - \sigma(s, X_s^{n,v}, \delta_{X_{s,\xi}^0})\|^2 ds\right)^{\frac{1}{2}} \\ &\quad + \sqrt{\varepsilon_n} \mathbb{E}\left(\int_0^t K_2^2(t, s) \cdot \|\sigma(s, X_s^{n,v}, \mathcal{L}_{X_s^n})\|^2 ds\right)^{\frac{1}{2}} \\ &\leq L_1 \int_0^t K_1(t, s) \cdot (\mathbb{E}|X_s^n - X_{s,\xi}^0|^p)^{\frac{1}{p}} ds \\ &\quad + \sqrt{NL_2} \mathbb{E}\left(\int_0^t K_1^2(t, s) \cdot \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0}) ds\right)^{\frac{1}{2}} \\ &\quad + \sqrt{\varepsilon_n L_2} \mathbb{E}\left(\int_0^t K_2^2(t, s) \cdot (1 + |X_s^{n,v}|^2 + \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_0)) ds\right)^{\frac{1}{2}} \\ &\leq \sqrt{\varepsilon_n} C_{T,L_1,L_2,\xi} + \sqrt{NL_2} \left[\mathbb{E}\left(\int_0^t K_1^2(t, s) \cdot \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0}) ds\right)^{\frac{p}{2}}\right]^{\frac{1}{p}} \\ &\quad + \sqrt{\varepsilon_n L_2} \left[\mathbb{E}\left(\int_0^t K_2^2(t, s) \cdot (1 + |X_s^{n,v}|^2 + \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_0)) ds\right)^{\frac{p}{2}}\right]^{\frac{1}{p}} \\ &\leq \sqrt{\varepsilon_n} C_{T,L_1,L_2,\xi} + \sqrt{NL_2} \left[\left(\int_0^t K_1^2(t, s) \cdot \mathbb{E}(\mathbb{W}_2^p(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0}))^{\frac{2}{p}} ds\right)^{\frac{p}{2}}\right]^{\frac{1}{p}} \\ &\quad + \sqrt{\varepsilon_n L_2} \left[\left(\int_0^t K_2^2(t, s) \cdot \mathbb{E}(1 + |X_s^{n,v}|^p + \mathbb{W}_2^p(\mathcal{L}_{X_s^n}, \delta_0))^{\frac{2}{p}} ds\right)^{\frac{p}{2}}\right]^{\frac{1}{p}} \\ &\leq \sqrt{\varepsilon_n} C_{T,N,L_1,L_2,\xi} + \sqrt{\varepsilon_n} C_{T,N,L_2,\xi} \\ &\leq C_{T,N,L_1,L_2,\xi} \sqrt{\varepsilon_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Hence $\mathbb{E}[\Phi_t(\phi, v)] = 0$, which implies that ϕ satisfies (3.11) almost surely, for all $t \in T$. By Lemma 3.5, we have that ϕ has continuous paths, so it satisfies (3.11) for all $t \in T$, almost surely, which means $\mathcal{G}_{v,N}^0$ consists of the unique solution of (3.11). Since this definition is independent of N , it extends to \mathcal{G}_v^0 . The proof is complete. \square

This Lemma shows that the set \mathcal{G}_v^0 corresponds to the unique solution ϕ_t of (3.11).

Lemma 3.8. *Assume that (H1)-(H3) hold, the functional I in (2.7) has compact level sets.*

Proof. By Lemma 3.7, we have that

$$I(\phi) := \inf \left\{ \frac{1}{2} \int_0^T |v_s|^2 ds : v \in L^2, \phi_t = \xi + \int_0^t K_1(t, s) b(s, \phi_s, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, \phi_s, \delta_{X_{s,\xi}^0}) v_s ds \right\}.$$

We are going to prove that the functional I defined in (2.7) has compact level sets, that is for all $N > 0$, the level sets

$$L_N := \{\phi \in C_T^d : I(\phi) \leq N\}$$

are compact. Fix $N > 0$ and consider an arbitrary sequence $\{\phi^n\}_{n \in \mathbb{N}} \subset L_N$, we will show that there exists a converging subsequence such that the limit belongs to L_N .

We first prove that $\{\phi^n\}_{n \in \mathbb{N}}$ is relative compact. According to classical Arzelà-Ascoli's Theorem, we only need to show that $\{\phi^n\}_{n \in \mathbb{N}}$ is uniformly bounded and equicontinuous. For all $n \in \mathbb{N}$ and $t \in [0, T]$, there exists $v^n \in L^2$ such that $\frac{1}{2} \int_0^T |v_t^n|^2 dt \leq N$ and $\phi^n \in \mathcal{G}_{v^n}^0$, which implies that $v^n \in \mathcal{S}_{2N}$ and ϕ^n is the solution of following Volterra type McKean-Vlasov integral equation:

$$\phi_t^n = \xi + \int_0^t K_1(t, s) b(s, \phi_s^n, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, \phi_s^n, \delta_{X_{s,\xi}^0}) v_s^n ds.$$

By Lemmas 3.2 and 3.5, we obtain $\{\phi^n\}_{n \in \mathbb{N}}$ is relatively compact which ensures that L_N is relatively compact for any $N > 0$.

Next we show that $\{\phi^n\}_{n \in \mathbb{N}}$ is closed, that is, let $\{\phi^n\}_{n \in \mathbb{N}}$ be a converging sequence of L_N and denote its limit by $\phi \in C_T^d$, we want to prove that the limit $\phi \in L_N$. Since $v^n \in \mathcal{S}_{2N}$ which is a compact space with respect to the weak topology. Hence there exists a subsequence $\{n_k\}_{k \in \mathbb{N}}$ such that v^{n_k} converges weakly in L^2 to the limit $v \in \mathcal{S}_{2N}$ and $\lim_{k \rightarrow \infty} \phi^{n_k} = \phi$.

Notice that for $v, v^{n_k} \in \mathcal{S}_{2N}$, by (H1)-(H3), Hölder's inequality, Lemmas 2.15 and 3.2 we have that

$$\begin{aligned} & \left| \phi_t^{n_k} - \left(\xi + \int_0^t K_1(t, s) b(s, \phi_s, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, \phi_s, \delta_{X_{s,\xi}^0}) v_s ds \right) \right| \\ & \leq \left| \int_0^t K_1(t, s) \cdot (b(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) - b(s, \phi_s, \delta_{X_{s,\xi}^0})) ds \right| \\ & \quad + \left| \int_0^t K_1(t, s) \cdot (\sigma(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) v_s^{n_k} - \sigma(s, \phi_s, \delta_{X_{s,\xi}^0}) v_s) ds \right| \\ & \leq L_1 \int_0^t K_1(t, s) \cdot |\phi_s^{n_k} - \phi_s| ds + \int_0^t K_1(t, s) \cdot \|\sigma(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) - \sigma(s, \phi_s, \delta_{X_{s,\xi}^0})\| \cdot |v_s^{n_k}| ds \\ & \quad + \int_0^t K_1(t, s) \cdot \|\sigma(s, \phi_s, \delta_{X_{s,\xi}^0})\| \cdot |v_s^{n_k} - v_s| ds \\ & \leq L_1 \int_0^t K_1(t, s) \cdot |\phi_s^{n_k} - \phi_s| ds + \left(\int_0^t |v_s^{n_k}|^2 ds \right)^{\frac{1}{2}} \cdot \left(\int_0^t K_1^2(t, s) \cdot \|\sigma(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) - \sigma(s, \phi_s, \delta_{X_{s,\xi}^0})\|^2 ds \right)^{\frac{1}{2}} \\ & \quad + C_{L_2} \int_0^t K_1(t, s) \cdot (1 + |\phi_s| + \mathbb{W}_2(\delta_{X_{s,\xi}^0}, \delta_0)) \cdot |v_s^{n_k} - v_s| ds \\ & \leq C_{L_1, T} \sup_{t \in [0, T]} |\phi_t^{n_k} - \phi_t| + C_{N, L_1} \left(\int_0^t K_1^2(t, s) \cdot |\phi_s^{n_k} - \phi_s|^2 ds \right)^{\frac{1}{2}} \\ & \quad + C_{L_2} (1 + \sup_{t \in [0, T]} |\phi_t| + \sup_{t \in [0, T]} |X_t^0|) \cdot \int_0^t K_1(t, s) \cdot |v_s^{n_k} - v_s| ds \\ & \leq C_{L_1, T} \sup_{t \in [0, T]} |\phi_t^{n_k} - \phi_t| + C_{N, L_1, T} \sup_{t \in [0, T]} |\phi_t^{n_k} - \phi_t| + C_{T, N, L_2} \int_0^t K_1(t, s) \cdot |v_s^{n_k} - v_s| ds \end{aligned}$$

$$\begin{aligned} &\leq C_{N,L_1,T} \sup_{t \in [0,T]} |\phi_t^{n_k} - \phi_t| \rightarrow 0 \quad \text{as } n_k \rightarrow \infty \quad (\lim_{k \rightarrow \infty} \phi^{n_k} = \phi) \\ &+ C_{T,N,L_2} \int_0^t K_1(t,s) \cdot |v_s^{n_k} - v_s| ds \rightarrow 0 \quad \text{as } n_k \rightarrow \infty \quad (v^{n_k} \text{ converges weakly in } L^2 \text{ to } v). \end{aligned}$$

So we easily obtain that

$$\begin{aligned} \phi_t &= \lim_{k \rightarrow \infty} \phi^{n_k} \\ &= \lim_{k \rightarrow \infty} \left(\xi + \int_0^t K_1(t,s) b(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t,s) \sigma(s, \phi_s^{n_k}, \delta_{X_{s,\xi}^0}) v_s^{n_k} ds \right) \\ &= \xi + \int_0^t K_1(t,s) b(s, \phi_s, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t,s) \sigma(s, \phi_s, \delta_{X_{s,\xi}^0}) v_s ds. \end{aligned}$$

Since $v \in \mathcal{S}_{2N}$ so that $\phi \in L_N$, which concludes the proof of the closure and therefore of the compactness of L_N . \square

Thus, we have verified the two conditions in Lemma 2.11 (by Lemma 3.4 and Lemma 3.8) and Theorem 3.1 is proved.

4. THE CENTRAL LIMIT THEOREM FOR VOLTERRA TYPE MCKEAN-VLASOV SDEs

In this section, we will study the asymptotic behavior of

$$\frac{X_{t,\xi}^\varepsilon - X_{t,\xi}^0}{\sqrt{\varepsilon}}, \quad 0 \leq t \leq T,$$

where X^0 satisfies the deterministic Volterra integral equation (1.2). Now we impose some assumptions on the coefficient functions that will enable us to study central limit theorem:

(H4) For any $x, y \in \mathbb{R}^d$, $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ and $t \in [0, T]$

$$\|\nabla b(t, x, \mu)\| \leq L_3, \quad \|D^L b(t, x, \mu)\|_{T_{\mu,2}} \leq L_3,$$

where $\nabla b(t, x, \mu)$ denotes the derivative of $b(t, x, \mu)$ with respect to x , $D^L b(t, x, \mu)$ is the Lions derivative with respect to μ , and $L_3 > 0$ is a positive constant.

(H5) For $t \in [0, T]$, $x, y \in \mathbb{R}^d$ and $X, Y, \phi \in L^2(\Omega \mapsto \mathbb{R}^d, \mathbb{P})$

$$\begin{aligned} &\left| \mathbb{E} \langle D^L b(t, x, \mathcal{L}_X)(X), \phi \rangle - \mathbb{E} \langle D^L b(t, y, \mathcal{L}_Y)(Y), \phi \rangle \right| \\ &\leq L_4 \left(|x - y| + \mathbb{W}_2(\mathcal{L}_X, \mathcal{L}_Y) + (\mathbb{E}|X - Y|^2)^{\frac{1}{2}} \right) \cdot (\mathbb{E}|\phi|^2)^{\frac{1}{2}} \end{aligned}$$

for some positive constant $L_4 > 0$.

(H6) The derivative of $\nabla b(t, \cdot, \mu)$ exists and for any $x, y \in \mathbb{R}^d$, $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ and $t \in [0, T]$

$$\|\nabla b(t, x, \mu) - \nabla b(t, y, \nu)\| \leq L_5 (|x - y| + \mathbb{W}_2(\mu, \nu)),$$

for some positive constant $L_5 > 0$.

Define $Z_{t,\xi}^\varepsilon = Q_{t,\xi}^\varepsilon|_{h(\varepsilon)=1} = \frac{X_{t,\xi}^\varepsilon - X_{t,\xi}^0}{\sqrt{\varepsilon}}$, and we can write

$$Z_{t,\xi}^\varepsilon = \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s, \quad (4.13)$$

Then we construct another Volterra type McKean-Vlasov SDE of $Z_{t,\xi}$:

$$\begin{aligned} Z_{t,\xi} &= \int_0^t K_1(t,s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right] ds \\ &\quad + \int_0^t K_2(t,s) \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s. \end{aligned} \quad (4.14)$$

Assume **(H1)-(H4)**. There exists a unique solution $Z_{t,\xi}$ of (4.14) (Theorem 3.1 of [21]). We now state main results in this section.

Theorem 4.1. *Suppose that the assumptions **(H1)-(H6)** are satisfied, then it holds that for any $R > 0$ and $p > 2$*

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} \left(\sup_{t \in [0,T], |\xi| \leq R} |Z_{t,\xi}^\varepsilon - Z_{t,\xi}|^p \right) = 0.$$

We first prepare some preliminary lemmas needed for the proof of the main theorem.

Lemma 4.2. [29] *Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be an atomless space, and let $X, Y \in L^2(\Omega \mapsto \mathbb{R}^d, \mathbb{P})$ with $\mathcal{L}_X = \mu$. If either X and Y are bounded and f is L -differentiable at μ , or $f \in C_b^1(\mathcal{P}_2(\mathbb{R}^d))$, then*

$$\lim_{\varepsilon \rightarrow 0} \frac{f(\mathcal{L}_{X+\varepsilon Y}) - f(\mu)}{\varepsilon} = \mathbb{E} \langle D^L f(\mu)(X), Y \rangle.$$

Consequently,

$$\left| \lim_{\varepsilon \rightarrow 0} \frac{f(\mathcal{L}_{X+\varepsilon Y}) - f(\mu)}{\varepsilon} \right| = |\mathbb{E} \langle D^L f(\mu)(X), Y \rangle| \leq \|D^L f(\mu)\|_{T_{\mu,2}} \sqrt{\mathbb{E}|Y|^2}.$$

First we estimate $Z_{t,\xi}^\varepsilon = Q_{t,\xi}^\varepsilon|_{h(\varepsilon)=1}$. Lemma 2.16 implies that Lemma 4.3 (i).

Lemma 4.3. (i) *Under the assumptions **(H1)-(H3)**, it follows that for $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E} |Z_{t,\xi}^\varepsilon|^p \leq C_{p,T,L_1,L_2}(1 + |\xi|^p).$$

(ii) *Under the assumptions **(H1)-(H6)**, it follows that for $p > 2, \xi, \eta \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E} |Z_{t,\xi}^\varepsilon - Z_{t,\eta}^\varepsilon|^p \leq C_{p,T,L_1,L_2,L_3,L_4,L_5}(1 + |\xi|^p)|\xi - \eta|^p.$$

(iii) *Under the assumptions **(H1)-(H3)**, it follows that for any $t, t' \in [0, T]$, $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E} |Z_{t,\xi}^\varepsilon - Z_{t',\xi}^\varepsilon|^p \leq C_{p,T,L_1,L_2}(1 + |\xi|^p)|t - t'|^{\gamma p},$$

where all the constants on the above right hand sides are independent of ε .

Proof. We are going to prove (iii) and the case (ii) is similar. For any $t > t'$ we have

$$\begin{aligned} &\mathbb{E} |Z_{t,\xi}^\varepsilon - Z_{t',\xi}^\varepsilon|^p \\ &= \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right. \\ &\quad \left. - \int_0^{t'} K_1(t',s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds - \int_0^{t'} K_2(t',s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right|^p \\ &= \mathbb{E} \left| \int_0^{t'} (K_1(t,s) - K_1(t',s)) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds \right|^p \end{aligned}$$

$$\begin{aligned}
& + \int_0^{t'} (K_2(t, s) - K_2(t', s)) \cdot \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s + \int_{t'}^t K_1(t, s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds \\
& + \left| \int_{t'}^t K_2(t, s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right|^p \\
\leq & 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_1(t, s) - K_1(t', s)) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds \right|^p \\
& + 4^{p-1} \mathbb{E} \left| \int_{t'}^t K_1(t, s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds \right|^p \\
& + 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_2(t, s) - K_2(t', s)) \cdot \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right|^p \\
& + 4^{p-1} \mathbb{E} \left| \int_{t'}^t K_2(t, s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right|^p := \Xi_1 + \Xi_2.
\end{aligned}$$

For the term Ξ_1 , by Lemma 2.13 (iv), Lemma 4.3 (i), (H1)-(H3), we have

$$\begin{aligned}
\Xi_1 & \leq C_{p,T} |t - t'|^{\gamma p} \sup_{t \in [0, T]} \mathbb{E} \left| \frac{b(t, X_{t,\xi}^\varepsilon, \mathcal{L}_{X_{t,\xi}^\varepsilon}) - b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})}{\sqrt{\varepsilon}} \right|^p \\
& \leq C_{p,T,L_1} |t - t'|^{\gamma p} \sup_{t \in [0, T]} \mathbb{E} |Z_{t,\xi}^\varepsilon|^p \\
& \leq C_{p,T,L_1,L_2} (1 + |\xi|^p) |t - t'|^{\gamma p}.
\end{aligned}$$

For the term Ξ_2 , applying 2.13 (v), Lemma 2.14, (H1)-(H3), we have

$$\begin{aligned}
\Xi_2 & \leq C_{p,T} |t' - t|^{\gamma p} \sup_{t \in [0, T]} \mathbb{E} \|\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon})\|^p \\
& \leq C_{p,T,L_2} |t' - t|^{\gamma p} (1 + \sup_{t \in [0, T]} \mathbb{E} |X_{t,\xi}^\varepsilon|^p) \\
& \leq C_{p,T,L_2} (1 + |\xi|^p) |t' - t|^{\gamma p}.
\end{aligned}$$

Hence, for all $0 \leq t < t' \leq T$,

$$\mathbb{E} |Z_{t,\xi}^\varepsilon - Z_{t',\xi}^\varepsilon|^p \leq C_{p,T,L_1,L_2} (1 + |\xi|^p) |t' - t|^{\gamma p}.$$

□

Next, we are going to take care of bound for $Z_{t,\xi}$.

Lemma 4.4. *Assume (H1)-(H4). Then for $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0, T]} \mathbb{E} |Z_{t,\xi}|^p \leq C_{p,T,L_2,L_3} (1 + |\xi|^p),$$

where the constant C_{p,T,L_2,L_3} is independent of ε .

Proof. By (4.14), BDG's inequality, Hölder's inequality, (H1), (H3) and (H4) we have

$$\begin{aligned}
\mathbb{E} |Z_{t,\xi}|^p & = \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right] ds \right. \\
& \quad \left. + \int_0^t K_2(t, s) \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s \right|^p \\
& \leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right] ds \right|^p
\end{aligned}$$

$$\begin{aligned}
& + 2^{p-1} \mathbb{E} \left| \int_0^t K_2(t, s) \sigma(s, X_s^0, \mathcal{L}_{X_s^0}) dW_s \right|^p \\
& \leq 2^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right. \\
& \quad \left. + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right|^p ds \cdot \left| \int_0^t K_1(t, s) ds \right|^{p-1} \\
& \quad + C_p \mathbb{E} \left(\int_0^t K_2^2(t, s) \cdot \left\| \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) \right\|^2 ds \right)^{\frac{p}{2}} \\
& \leq C_{p,L_3} \mathbb{E} \int_0^t K_1(t, s) \cdot (|Z_{s,\xi}|^p + \mathbb{E}|Z_{s,\xi}|^p) ds \cdot \left| \int_0^t K_1(t, s) ds \right|^{p-1} \\
& \quad + C_p \mathbb{E} \int_0^t K_2^2(t, s) \cdot \left\| \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) \right\|^p ds \cdot \left| \int_0^t K_2^2(t, s) ds \right|^{\frac{p}{2}-1} \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E}|Z_{s,\xi}|^p ds + C_{p,T,L_2} \int_0^t K_2^2(t, s) \cdot (1 + \mathbb{E}|X_{s,\xi}^0|^p) ds \\
& \leq C_{p,T,L_2}(1 + |\xi|^p) + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E}|Z_{s,\xi}|^p ds,
\end{aligned}$$

where in the last inequality Lemma 2.15 is used.

By Remark 2.12, Lemma 2.3 and (2.6), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}
\mathbb{E}|Z_{t,\xi}|^{2p} & \leq C_{p,T,L_2}(1 + |\xi|^p) + \int_0^t R^{K_1^*}(t, s) \cdot C_{p,T,L_2}(1 + |\xi|^p) ds \\
& \leq C_{p,T,L_2,L_3}(1 + |\xi|^p).
\end{aligned}$$

□

Lemma 4.5. *Under the assumptions **(H1)**-**(H5)**, we have that for $p > 2, \xi, \eta \in \mathbb{R}^d$*

$$\sup_{t \in [0, T]} \mathbb{E}|Z_{t,\xi} - Z_{t,\eta}|^p \leq C_{p,T,L_2,L_3,L_4,L_5}(1 + |\xi|^p)|\xi - \eta|^p,$$

where the constant C_{p,T,L_2,L_3,L_4,L_5} is independent of ε .

Proof. By (4.14), BDG's inequality, Hölder's inequality, **(H1)** and **(H3)-(H5)** we have

$$\begin{aligned}
& \mathbb{E}|Z_{t,\xi} - Z_{t,\eta}|^p \\
& = \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right] ds \right. \\
& \quad \left. + \int_0^t K_2(t, s) \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s - \int_0^t K_2(t, s) \sigma(t, X_{s,\eta}^0, \delta_{X_{s,\eta}^0}) dW_s \right. \\
& \quad \left. - \int_0^t K_1(t, s) \cdot \left[\nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\eta} \rangle \right] ds \right|^p \\
& \leq 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})) ds \right|^p \\
& \quad + 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle - \mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\eta} \rangle) ds \right|^p \\
& \quad + 3^{p-1} \mathbb{E} \left| \int_0^t K_2(t, s) \cdot (\sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \sigma(t, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})) dW_s \right|^p \\
& \leq 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})) ds \right|^p
\end{aligned}$$

$$\begin{aligned}
& + 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\nabla_{Z_{s,\xi}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0}) - \nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})) ds \right|^p \\
& + 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle - \mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\xi} \rangle) ds \right|^p \\
& + 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot (\mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\xi} \rangle - \mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\eta} \rangle) ds \right|^p \\
& + C_p \mathbb{E} \left(\int_0^t K_2^2(t, s) \cdot \|\sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \sigma(t, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})\|^2 ds \right)^{\frac{p}{2}} \\
& \leq 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot |\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})|^p ds \cdot \left(\int_0^t K_1(t, s) ds \right)^{p-1} \\
& + 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot |\nabla_{Z_{s,\xi}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0}) - \nabla_{Z_{s,\eta}} b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})|^p ds \cdot \left(\int_0^t K_1(t, s) ds \right)^{p-1} \\
& + 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot |\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0) - D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\xi} \rangle|^p ds \cdot \left(\int_0^t K_1(t, s) ds \right)^{p-1} \\
& + 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot |\mathbb{E} \langle D^L b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})(X_{s,\eta}^0), Z_{s,\xi} - Z_{s,\eta} \rangle|^p ds \cdot \left(\int_0^t K_1(t, s) ds \right)^{p-1} \\
& + C_p \mathbb{E} \int_0^t K_2^2(t, s) \cdot \|\sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \sigma(t, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})\|^p \cdot ds \left(\int_0^t K_2^2(t, s) ds \right)^{\frac{p}{2}-1} \\
& \leq C_{p,T,L_5} \mathbb{E} \int_0^t K_1(t, s) \cdot (|X_{s,\xi}^0 - X_{s,\eta}^0|^p + \mathbb{W}_2^p(\delta_{X_{s,\xi}^0}, \delta_{X_{s,\eta}^0})) \cdot |Z_{s,\xi}|^p ds + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi} - Z_{s,\eta}|^p ds \\
& + C_{p,T,L_4} \mathbb{E} \int_0^t K_1(t, s) \cdot (|X_{s,\xi}^0 - X_{s,\eta}^0|^p + \mathbb{W}_2^p(\delta_{X_{s,\xi}^0}, \delta_{X_{s,\eta}^0}) + (\mathbb{E} |X_{s,\xi}^0 - X_{s,\eta}^0|^2)^{\frac{p}{2}}) \cdot (\mathbb{E} |Z_{s,\xi}|^2)^{\frac{p}{2}} ds \\
& + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi} - Z_{s,\eta}|^p ds + C_{p,T,L_2} \mathbb{E} \int_0^t K_2^2(t, s) \cdot (|X_{s,\xi}^0 - X_{s,\eta}^0|^p + \mathbb{W}_2^p(\delta_{X_{s,\xi}^0}, \delta_{X_{s,\eta}^0})) ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi} - Z_{s,\eta}|^p ds + C_{p,T,L_4} \mathbb{E} \int_0^t K_1(t, s) \cdot \mathbb{E} |X_{s,\xi}^0 - X_{s,\eta}^0|^p ds \\
& + C_{p,T,L_2} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^0 - X_{s,\eta}^0|^p ds + C_{p,T,L_4} \int_0^t K_1(t, s) \cdot \mathbb{E} |X_{s,\xi}^0 - X_{s,\eta}^0|^p \cdot \mathbb{E} |Z_{s,\xi}|^p ds \\
& + C_{p,T,L_5} \int_0^t K_1(t, s) \cdot (\mathbb{E} |X_{s,\xi}^0 - X_{s,\eta}^0|^{2p})^{\frac{1}{2}} \cdot (\mathbb{E} |Z_{s,\xi}|^{2p})^{\frac{1}{2}} ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi} - Z_{s,\eta}|^p ds + C_{p,T,L_2,L_3,L_4} (1 + |\xi|^p) |\xi - \eta|^p,
\end{aligned}$$

where in the last inequality Lemmas 2.15 and 4.4 are used.

Finally, by Remark 2.12 and Lemma 2.3, we have

$$\begin{aligned}
\mathbb{E} |Z_{t,\xi} - Z_{t,\eta}|^p & \leq C_{p,T,L_2,L_3,L_4,L_5} (1 + |\xi|^p) |\xi - \eta|^p + \int_0^t R^{K_1^*}(t, s) \cdot (C_{p,T,L_2,L_3,L_4,L_5} (1 + |\xi|^p) |\xi - \eta|^p) ds \\
& \leq C_{p,T,L_2,L_3,L_4,L_5} (1 + |\xi|^p) |\xi - \eta|^p.
\end{aligned}$$

This completes the proof of the lemma. \square

Lemma 4.6. *Under the assumptions (H1)-(H4), we have that for any $t, t' \in [0, T]$, $p > 2$, $\xi \in \mathbb{R}^d$*

$$\sup_{t \in [0, T]} \mathbb{E} |Z_{t,\xi} - Z_{t',\xi}|^p \leq C_{p,T,L_2,L_3} (1 + |\xi|^p) |t - t'|^{\gamma p},$$

where the constant C_{p,T,L_2,L_3} is independent of ε .

Proof. For any $t > t'$ we have

$$\begin{aligned}
& \mathbb{E}|Z_{t,\xi} - Z_{t',\xi}|^p \\
&= \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right. \\
&\quad - \int_0^{t'} K_1(t',s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \\
&\quad + \left. \int_0^t K_2(t,s) \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s - \int_0^{t'} K_2(t',s) \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s \right|^p \\
&= \mathbb{E} \left| \int_0^{t'} (K_1(t,s) - K_1(t',s)) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right. \\
&\quad + \left. \int_0^{t'} (K_2(t,s) - K_2(t',s)) \cdot \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s + \int_{t'}^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s \right. \\
&\quad + \left. \int_{t'}^t K_1(t,s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right|^p \\
&\leq 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_1(t,s) - K_1(t',s)) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right|^p \\
&\quad + 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_2(t,s) - K_2(t',s)) \cdot \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s \right|^p \\
&\quad + 4^{p-1} \mathbb{E} \left| \int_{t'}^t K_1(t,s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right|^p \\
&\quad + 4^{p-1} \mathbb{E} \left| \int_{t'}^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s \right|^p := \Theta_1 + \Theta_2 + \Theta_3 + \Theta_4.
\end{aligned}$$

For the term Θ_1 , by Hölder's inequality, Lemma 4.4, (H1), (H3) and (H4), we have

$$\begin{aligned}
\Theta_1 &:= 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_1(t,s) - K_1(t',s)) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E}\langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right|^p \\
&\leq C_{p,L_3} \mathbb{E} \int_0^{t'} |K_1(t,s) - K_1(t',s)| \cdot (\mathbb{E}|Z_{s,\xi}|^p + |Z_{s,\xi}|^p) ds \cdot \left(\int_0^{t'} |K_1(t,s) - K_1(t',s)| ds \right)^{p-1} \\
&\leq C_{p,T,L_3} \int_0^{t'} |K_1(t,s) - K_1(t',s)| \cdot \mathbb{E}|Z_{s,\xi}|^p ds |t - t'|^{\gamma(p-1)} \\
&\leq C_{p,T,L_2,L_3} (1 + |\xi|^p) |t - t'|^{\gamma p}.
\end{aligned}$$

For the term Θ_2 , applying BDG's inequality, Hölder's inequality, extended Minkowski's inequality, Lemma 2.14, (H1) and (H3), we have

$$\begin{aligned}
\Theta_2 &:= 4^{p-1} \mathbb{E} \left| \int_0^{t'} (K_2(t,s) - K_2(t',s)) \cdot \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s \right|^p \\
&\leq C_p \mathbb{E} \left(\int_0^{t'} |K_2(t,s) - K_2(t',s)|^2 \cdot \|\sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0})\|^2 ds \right)^{\frac{p}{2}} \\
&\leq C_{p,L_2} \mathbb{E} \left(\int_0^{t'} |K_2(t,s) - K_2(t',s)|^2 \cdot (1 + |X_{s,\xi}^0|^2 + \mathbb{W}_2^2(\mathcal{L}_{X_{s,\xi}^0}, \delta_0)) ds \right)^{\frac{p}{2}} \\
&\leq C_{p,L_2} \left(\int_0^{t'} |K_2(t,s) - K_2(t',s)|^2 \cdot (\mathbb{E}[1 + |X_{s,\xi}^0|^p + \mathbb{W}_2^p(\mathcal{L}_{X_{s,\xi}^0}, \delta_0)])^{\frac{2}{p}} ds \right)^{\frac{p}{2}}
\end{aligned}$$

$$\begin{aligned} &\leq C_{p,L_2} \left(\int_0^{t'} |K_2(t,s) - K_2(t',s)|^2 \cdot (1 + (\mathbb{E}|X_{s,\xi}^0|^p)^{\frac{2}{p}}) ds \right)^{\frac{p}{2}} \\ &\leq C_{p,T,L_2} (1 + |\xi|^p) \left(\int_0^t |K_2(t,s) - K_2(t',s)|^2 ds \right)^{\frac{p}{2}} \leq C_{p,T,L_2} (1 + |\xi|^p) |t' - t|^{\gamma p}. \end{aligned}$$

For the term Θ_3 , by Hölder's inequality, Lemma 4.4, (H1), (H3) and (H4), we have

$$\begin{aligned} \Theta_3 &:= 4^{p-1} E \left| \int_{t'}^t K_1(t,s) \cdot \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(t, X_{t,\xi}^0, \delta_{X_{t,\xi}^0})(X_{t,\xi}^0), Z_{t,\xi} \rangle \right] ds \right|^p \\ &\leq C_{p,L_3} \mathbb{E} \int_{t'}^t K_1(t,s) \cdot (\mathbb{E}|Z_{s,\xi}|^p + |Z_{s,\xi}|^p) ds \cdot \left(\int_t^{t'} K_1(t,s) ds \right)^{p-1} \\ &\leq C_{p,T,L_3} \int_{t'}^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}|^p ds |t - t'|^{\gamma(p-1)} \\ &\leq C_{p,T,L_2,L_3} (1 + |\xi|^p) |t - t'|^{\gamma p}. \end{aligned}$$

For the term Θ_4 , by BDG's inequality, Hölder's inequality, Lemma 2.14, (H1) and (H3), we have

$$\begin{aligned} \Theta_4 &:= 4^{p-1} \mathbb{E} \left| \int_{t'}^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0}) dW_s \right|^p \\ &\leq C_p \mathbb{E} \left(\int_{t'}^t K_2^2(t,s) \|\sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0})\|^2 ds \right)^{\frac{p}{2}} \\ &\leq C_p \mathbb{E} \int_{t'}^t K_2^2(t,s) \cdot \|\sigma(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^0})\|^p ds \cdot \left(\int_{t'}^t K_2^2(t,s) ds \right)^{\frac{p}{2}-1} \\ &\leq C_{p,T} \mathbb{E} \int_{t'}^t K_2^2(t,s) \cdot (1 + |X_{s,\xi}^0|^p + \mathbb{W}_2^p(\mathcal{L}_{X_{s,\xi}^0}, \delta_0)) ds |t - t'|^{\gamma(p-2)} \\ &\leq C_{p,T,L_2} (1 + |\xi|^p) |t - t'|^{\gamma p}. \end{aligned}$$

Hence, for all $0 \leq t < t' \leq T$,

$$\mathbb{E}|Z_{t,\xi} - Z_{t',\xi}|^p \leq C_{p,T,L_2,L_3} (1 + |\xi|^p) |t' - t|^{\gamma p}.$$

This proves the lemma. \square

Now we bound $Z_{t,\xi}^\varepsilon - Z_{t,\xi}$.

Lemma 4.7. Suppose that the assumptions (H1)-(H6) are satisfied, then it holds that for $p > 2, \xi \in \mathbb{R}^d$

$$\mathbb{E}|Z_{t,\xi}^\varepsilon - Z_{t,\xi}|^p \leq C_{p,T,L_3,L_4} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}),$$

where the constant C_{p,T,L_3,L_4} is independent of ε .

Proof.

$$\begin{aligned} &\mathbb{E}|Z_{t,\xi}^\varepsilon - Z_{t,\xi}|^p \\ &= \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right. \\ &\quad \left. - \int_0^t K_1(t,s) \left[\nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) + \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right] ds - \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) dW_s \right|^p \\ &\leq 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \left(\frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon})}{\sqrt{\varepsilon}} - \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right) ds \right|^p \end{aligned}$$

$$\begin{aligned}
& + 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\frac{b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right) \right|^p ds \\
& + 3^{p-1} \mathbb{E} \left| \int_0^t K_2(t, s) \cdot (\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})) dW_s \right|^p \\
& := \Pi_1 + \Pi_2 + \Pi_3.
\end{aligned}$$

For the term Π_1 , by Hölder's inequality, Lemmas 3.6, 4.3, (H1)-(H4) and (H6) we have

$$\begin{aligned}
\Pi_1 & = 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon})}{\sqrt{\varepsilon}} - \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right) ds \right|^p \\
& \leq 3^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon})}{\sqrt{\varepsilon}} - \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right|^p ds \cdot \left(\int_0^t K_1(t, s) ds \right)^{p-1} \\
& \leq C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) du \right|^p ds \\
& \leq C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| \nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \nabla_{Z_{s,\xi}^\varepsilon} b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) \right|^p du ds \\
& \quad + C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| \nabla_{Z_{s,\xi}^\varepsilon} b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \nabla_{Z_{s,\xi}} b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) \right|^p du ds \\
& \leq C_{p,T,L_5} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 |M_s^\xi(u) - X_{s,\xi}^0|^p du \cdot |Z_{s,\xi}^\varepsilon|^p ds + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds + C_{p,T,L_5} \int_0^t K_1(t, s) \cdot \mathbb{E} (|X_{s,\xi}^\varepsilon - X_{s,\xi}^0|^p \cdot |Z_{s,\xi}^\varepsilon|^p) ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds + C_{p,T,L_5} \int_0^t K_1(t, s) \cdot (\mathbb{E} |Z_{s,\xi}^\varepsilon|^{2p})^{\frac{1}{2}} \cdot (\mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\xi}^0|^{2p})^{\frac{1}{2}} ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds + C_{p,T,L_1, L_2} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}),
\end{aligned}$$

where $M_s^\xi(u) = X_{s,\xi}^0 + u(X_{s,\xi}^\varepsilon - X_{s,\xi}^0)$, $u \in [0, T]$.

For the term Π_2 , by Hölder's inequality, Lemmas 3.6, 4.3, and (H1)-(H5) we have

$$\begin{aligned}
\Pi_2 & = 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\frac{b(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right) ds \right|^p \\
& \leq 3^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\int_0^1 (\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r)), Z_{s,\xi}^\varepsilon \rangle \right. \right. \\
& \quad \left. \left. - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right) dr ds \right|^p \\
& \leq 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\int_0^1 (\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r)), Z_{s,\xi}^\varepsilon \rangle \right. \right. \\
& \quad \left. \left. - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle \right) dr ds \right|^p + 6^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \right. \\
& \quad \cdot \left(\int_0^1 (\mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi}^\varepsilon \rangle - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \rangle) dr ds \right|^p \\
& \leq 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r)), Z_{s,\xi}^\varepsilon \rangle - \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi}^\varepsilon \rangle dr \right|^p ds \\
& \quad \cdot \left| \int_0^t K_1(t, s) ds \right|^{p-1} + 6^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \mathbb{E} \langle D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi}^\varepsilon \rangle, Z_{s,\xi} \rangle \right|^p ds
\end{aligned}$$

$$\begin{aligned}
& - \mathbb{E} \left| D^L b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})(X_{s,\xi}^0), Z_{s,\xi} \right|^p ds \cdot \left| \int_0^t K_1(t,s) ds \right|^{p-1} \\
& \leq C_{p,T,L_4} \mathbb{E} \int_0^t K_1(t,s) \cdot \int_0^1 \left| \mathbb{W}_2(\mathcal{L}_{R_s^\xi(r)}, \delta_{X_{s,\xi}^0}) + (\mathbb{E}|R_s^\xi(r) - X_{s,\xi}^0|^2)^{\frac{1}{2}} \right|^p dr \cdot (\mathbb{E}|Z_{s,\xi}^\varepsilon|^2)^{\frac{p}{2}} ds \\
& \quad + C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds \\
& \leq C_{p,T,L_4} \int_0^t K_1(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\xi}^0|^p \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon|^p ds + C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds \\
& \leq C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds + C_{p,T,L_1,L_2,L_4} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}),
\end{aligned}$$

where $R_s^\xi(r) := X_{s,\xi}^0 + r(X_{s,\xi}^\varepsilon - X_{s,\xi}^0)$, $t \in [0, T]$.

For the term Π_3 , by BDG's inequality, Hölder's inequality, Lemma 3.6 and (H1)-(H3), we have

$$\begin{aligned}
\Pi_3 &= 3^{p-1} \mathbb{E} \left| \int_0^t K_2(t,s) \cdot (\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})) dW_s \right|^p \\
&\leq C_p \mathbb{E} \left(\int_0^t K_2^2(t,s) \cdot \| \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \|^2 ds \right)^{\frac{p}{2}} \\
&\leq C_p \mathbb{E} \int_0^t K_2^2(t,s) \cdot \| \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(t, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \|^p ds \cdot \left(\int_0^t K_2^2(t,s) ds \right)^{\frac{p}{2}-1} \\
&\leq C_{p,T,L_2} \int_0^t K_2^2(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\xi}^0|^p ds \\
&\leq C_{p,T,L_1,L_2} \varepsilon^{\frac{p}{2}} (1 + |\xi|^p) \leq C_{p,T,L_1,L_2} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}).
\end{aligned}$$

So by Remark 2.12 and Lemma 2.3, we have

$$\begin{aligned}
\mathbb{E}|Z_{t,\xi}^\varepsilon - Z_{t,\xi}|^p &\leq C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\xi}|^p ds + C_{p,T,L_1,L_2,L_4} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}) \\
&\leq C_{p,T,L_1,L_2,L_3,L_4} \varepsilon^{\frac{p}{2}} (1 + |\xi|^{2p}).
\end{aligned}$$

□

We are now ready to prove Theorem 4.1.

Proof of Theorem 4.1. We define

$$Z(r, t, \xi) = \begin{cases} Z_{t,\xi}, & r = 0, \\ Z_{t,\xi}^{\varepsilon_n} + \frac{r-\varepsilon_n T}{\varepsilon_{n+1}} (Z_{t,\xi}^{\varepsilon_{n+1}} - Z_{t,\xi}^{\varepsilon_n}), & \varepsilon_{n+1} T < r \leq \varepsilon_n T, n \in \mathbb{N}. \end{cases}$$

Fix $R > 0$, by Lemmas 4.3[(ii)], 4.3[(iii)], 4.5, 4.6, 4.7, there exists a constant $\beta := \beta(\gamma) > 0$ such that for all $p > 2$, $t, t' \in [0, T]$, $\xi, \eta \in D_R := \{\xi \in \mathbb{R}^d, |\xi| \leq R\}$, we have

$$\mathbb{E} \left| Z(r, t, \xi) - Z(r', t', \eta) \right|^p \leq C_{p,T,R,L_3,L_4} (|r - r'|^{\beta p} + |t - t'|^{\beta p} + |\xi - \eta|^{\beta p}).$$

Thus, by Kolmogorov's continuity criterion, there is a p -order integrable random variable $A(\omega)$ such that

$$\sup_{t \in [0, T], |\xi| \leq R} \left| Z(r, t, \xi) - Z(r', t, \xi) \right| \leq A(\omega) |r - r'|^\lambda, \quad a.s.,$$

where $\lambda \in (0, \beta - \frac{d+2}{p})$. In particular, taking $r = 0, r' = \varepsilon_n T$ we have

$$\mathbb{E} \left(\sup_{t \in [0, T], |\xi| \leq R} |Z_{t,\xi}^{\varepsilon_n} - Z_{t,\xi}| \right)^p \leq |\varepsilon_n T|^{p\lambda} \mathbb{E}|A(\omega)|^p,$$

which yields the desired convergence. \square

5. THE MODERATE DEVIATION PRINCIPLE FOR VOLTERRA TYPE MCKEAN-VLASOV SDES

In this section, we shall study moderate deviation principle for our Volterra type McKean-Vlasov SDEs.

For $\varepsilon > 0$, define

$$Y_t^\varepsilon := \frac{X_{\cdot,\xi}^\varepsilon - X_{\cdot,\xi}^0}{\sqrt{\varepsilon} h(\varepsilon)} = \frac{\mathcal{G}^\varepsilon(W) - X_{\cdot,\xi}^0}{\sqrt{\varepsilon} h(\varepsilon)} := \mathcal{J}^\varepsilon(W). \quad (5.15)$$

Assume that $h(\varepsilon)$ satisfies

$$h(\varepsilon) \rightarrow \infty, \quad \sqrt{\varepsilon} h(\varepsilon) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

This $\mathcal{J}^\varepsilon : C_T^m \rightarrow C_T^d$ is a Borel measurable map for each $\varepsilon > 0$, and the Y_t^ε satisfies the following Volterra type McKean-Vlasov SDEs:

$$\begin{aligned} Y_t^\varepsilon &= \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)} ds \\ &\quad + \frac{1}{h(\varepsilon)} \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s. \end{aligned} \quad (5.16)$$

It is clear that the existence and unique of Y_t^ε hold by Theorem 3.1 of [21], then the moderate deviation principle for $\{X_t^\varepsilon\}_{\varepsilon>0}$ is equivalent to the large deviation principle for $\{Y_t^\varepsilon\}_{\varepsilon>0}$.

Applying Girsanov's theorem, the shifted version $Y_t^{\varepsilon,v} := \mathcal{J}^\varepsilon(W + h(\varepsilon) \int_0^t v_s ds)$ is the unique strong solution of the controlled equation under \mathbb{P} :

$$\begin{aligned} Y_t^{\varepsilon,v} &= \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)} ds \\ &\quad + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}) v_s ds \\ &\quad + \frac{1}{h(\varepsilon)} \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s. \end{aligned} \quad (5.17)$$

For all $v \in \mathcal{A}$, we define \mathcal{J}_v^0 to be the solution of the limiting Volterra integral equation

$$\psi_t = \int_0^t K_1(t,s) \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds. \quad (5.18)$$

To obtain the MDP, in addition to **(H1)-(H3)** and **(H6)**, we also need the following assumption.

(H7) There exists a positive constant L_6 such that for $t \in [0, T]$

$$\nabla b(t, 0, \delta_0) \leq L_6.$$

Define the new rate function as

$$\Lambda(\psi) := \inf \left\{ \frac{1}{2} \int_0^T |v_s|^2 ds : v \in L^2 \text{ such that } \psi \in \mathcal{J}_v^0 \right\}. \quad (5.19)$$

We now state our results of MDP.

Theorem 5.1. *Under **(H1)-(H3)** and **(H6)-(H7)**, the family $\{X_t^\varepsilon\}_{\varepsilon>0}$, the unique solution of (1.1), satisfies the Moderate Deviations Principle with rate function (5.19) and speed $h^2(\varepsilon)$, where \mathcal{J}_v^0 corresponds to the unique solution of the limiting equation (5.18).*

We are going to use Lemma 2.11 and will then verify the two conditions there.

Step 1 Verification of the Condition (i) of Lemma 2.11.

Lemma 5.2. *Assume that **(H1)-(H3)** and **(H6)-(H7)** hold. Then there exists a unique solution ψ_t such that for almost all $t \geq 0$, $N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$, some constant C_{T,N,L_2,L_5,L_6}*

$$\sup_{t \in [0,T]} |\psi_t| \leq C_{T,N,L_2,L_5,L_6} (1 + |\xi|), \quad (5.20)$$

for almost all $t \in [0, T]$.

From Lemma 2.16, we easily get the following estimate associated with $Y^\varepsilon := Q_t^\varepsilon = \frac{X^\varepsilon - X^0}{\sqrt{\varepsilon h(\varepsilon)}}$.

Lemma 5.3. *Under the assumptions **(H1)-(H3)**, it follows that for $p > 2, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E}|Y_t^\varepsilon|^p \leq \frac{1}{|h(\varepsilon)|^p} C_{p,T,L_1,L_2} (1 + |\xi|^p),$$

where the constant C_{p,T,L_1,L_2} is independent of ε .

Let

$$\begin{aligned} x := 0, \Lambda_{t,\xi}^{\varepsilon,v} := Y_t^{\varepsilon,v}, N(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}), \\ M(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \delta_{X_{s,\xi}^0}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon} h(\varepsilon)}, \\ G(s, X_{s,\xi}^0, \mathcal{L}_{X_{s,\xi}^\varepsilon}, \Lambda_{s,x}^{\varepsilon,v}, h(\varepsilon)) := \frac{1}{h(\varepsilon)} \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon} h(\varepsilon) Y_s^{\varepsilon,v}, \mathcal{L}_{X_{s,\xi}^\varepsilon}). \end{aligned}$$

Then Lemma 2.17 and Lemma 2.18 give us that

Lemma 5.4. *Under the assumptions **(H1)-(H3)**, for $p > 2, N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$*

$$\sup_{t \in [0,T]} \mathbb{E}|Y_t^{\varepsilon,v}|^p \leq C_{p,T,N,L_1,L_2} (1 + |\xi|^p),$$

where the constant C_{p,T,N,L_1,L_2} is independent of ε, v .

Lemma 5.5. *Assume that **(H1)-(H3)** hold. If $p > 2 \vee \frac{2}{\gamma}$, $t, s \in [0, T]$, $N > 0, \xi \in \mathbb{R}^d$ and $\{v^\varepsilon\}_{\varepsilon > 0}$ is a family in \mathcal{A}_N , then $Y^{\varepsilon,v^\varepsilon}$ admits a version which is Hölder continuous on $[0, T]$ of any order $\alpha < \frac{\gamma}{2} - \frac{1}{p}$, uniformly for all $\varepsilon > 0$. Denoting again this version by $Y^{\varepsilon,v^\varepsilon}$, one has for all $\varepsilon > 0$ small enough*

$$\mathbb{E} \left[\left(\sup_{0 \leq s < t \leq T} \frac{|Y_t^{\varepsilon,v^\varepsilon} - Y_s^{\varepsilon,v^\varepsilon}|}{|t-s|^\alpha} \right)^p \right] \leq C_{p,T,N,L_1,L_2,\xi},$$

for all $\alpha \in [0, \frac{\gamma}{2} - \frac{1}{p}]$. Moreover, the family of random variables $\{Y^{\varepsilon,v^\varepsilon}\}_{\varepsilon > 0}$ is tight in C_T .

By almost same procedures, we can get that the solution ψ_t of (5.18) also have Hölder continuous path, that is

Lemma 5.6. *Assume that **(H1)-(H3)** and **(H6)** hold, for $t, s \in [0, T]$, $N > 0, v \in \mathcal{A}_N, \xi \in \mathbb{R}^d$, then the solution ψ_t of (5.18) has for all $\varepsilon > 0$ small enough*

$$|\psi_t - \psi_s| \leq C_{T,N,L_2} (1 + |\xi|^p) |t - s|^\gamma.$$

This verifies Condition (i) of Lemma 2.11.

Step 2 Verification of the Condition (ii) of Lemma 2.11.

Lemma 5.7. *The set \mathcal{J}_v^0 from Definition 2.9 (\mathcal{G} is replaced by \mathcal{J} and $\varepsilon_n = \frac{1}{h(\varepsilon_n)}$) is characterized by*

$$\mathcal{J}_v^0 = \left\{ \psi : [0, T] \rightarrow \mathbb{R}^d \mid \psi_t = \int_0^t K_1(t, s) \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds \right\}.$$

Proof. Recall the definition 2.9 of $\mathcal{J}_{v,N}^0$. For $N \in \mathbb{N}$ and $v \in \mathcal{A}_N$, consider a subsequence $\{\varepsilon_n\}_{n \in \mathbb{N}} \subset [0, \infty)$ with $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ and a sequence $\{v^{\varepsilon_n}\}_{n \in \mathbb{N}} \subset \mathcal{A}_N$ such that $\lim_{n \rightarrow \infty} v^{\varepsilon_n} = v$ in distribution, and assume that $Y^{n,v} := Y^{\varepsilon_n, v^{\varepsilon_n}}$ converges in distribution to some random variable ψ with values in C_T^d . To simplify the representation, we denote $v^n := v^{\varepsilon_n}$.

By Skorohod representation theorem we can work with almost sure convergence for the purpose of identifying the limit. Hence, we may assume that $\{Y^{n,v}, v^n\}_{n \geq 0}$ converges almost surely in the product topology on $C_T^d \times \mathcal{S}_N$, and the limit is a $C_T^d \times \mathcal{S}_N$ -valued random variable (ψ, v) in a possibly different probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ as n tends to $+\infty$. For $t \in [0, T]$, define $\Psi_t : C_T^d \times \mathcal{S}_N \rightarrow \mathbb{R}$ as

$$\Psi_t(\omega, f) := \left| \omega_t - \int_0^t K_1(t, s) \nabla_{\omega_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds - \int_0^t K_2(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) f_s ds \right| \wedge 1.$$

It is clear that Ψ_t is bounded, and we are going to show that Ψ_t is continuous. To this end, let $\omega^n \rightarrow \omega$ in C_T^d and $f^n \rightarrow f$ in \mathcal{S}_N with respect the weak topology.

For $\omega, \omega^n \in C_T^d, f, f^n \in \mathcal{S}_N$, by **(H1)-(H3), (H6)-(H7)** and Lemma 2.15 we have that

$$\begin{aligned} & |\Psi_t(\omega, f) - \Psi_t(\omega^n, f^n)| \\ & \leq |\omega_t - \omega_t^n| + \int_0^t K_1(t, s) \cdot \|\nabla_{\omega_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{\omega_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\| ds \\ & \quad + \int_0^t K_2(t, s) \cdot |\sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) f_s - \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) f_s^n| ds \\ & \leq \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + \int_0^t K_1(t, s) \cdot |\omega_s - \omega_s^n| \cdot \|\nabla b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla b(s, 0, \delta_0)\| ds \\ & \quad + \int_0^t K_1(t, s) \cdot |\omega_s - \omega_s^n| \cdot \|\nabla b(s, 0, \delta_0)\| ds + \int_0^t K_2(t, s) \cdot |f_s - f_s^n| \cdot \|\sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\| ds \\ & \leq \sup_{t \in [0, T]} |\omega_t - \omega_t^n| \cdot \left[1 + L_5 \int_0^t K_1(t, s) \cdot (|X_{s,\xi}^0| + \mathbb{W}_2(\delta_{X_{s,\xi}^0}, \delta_0)) ds \right] \\ & \quad + C_{T,L_6} \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{L_2} \int_0^t K_2(t, s) \cdot |f_s - f_s^n| \cdot (1 + |X_{s,\xi}^0| + \mathbb{W}_2(\delta_{X_{s,\xi}^0}, \delta_0)) ds \\ & \leq C_{T,L_2,L_5,L_6} \sup_{t \in [0, T]} |\omega_t - \omega_t^n| + C_{L_2} \int_0^t K_2(t, s) \cdot |f_s - f_s^n| ds. \end{aligned}$$

Since f^n tends to f weakly in L^2 and $\lim_{n \rightarrow \infty} |\omega_t - \omega_t^n| = 0$, we see that $\Psi_t(f^n, \omega^n)$ tends to $\Psi_t(f, \omega)$ as n tends to ∞ , which implies Ψ_t is continuous and therefore

$$\lim_{n \rightarrow \infty} \mathbb{E}[\Psi_t(Y^{n,v}, v^n)] = \mathbb{E}[\Psi_t(\psi, v)].$$

Now it suffices for us to prove that the right hand of above equality is actually equal to zero.

Since

$$\mathbb{E}[\Psi_t(Y^{n,v}, v^n)] = \mathbb{E}\left[\left| \int_0^t K_1(t, s) \cdot \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} ds \right| \right]$$

$$\begin{aligned}
& + \int_0^t K_2(t, s) \cdot \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) v_s^n ds \\
& + \frac{1}{h(\varepsilon_n)} \int_0^t K_2(t, s) \cdot \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) dW_s \\
& - \int_0^t K_1(t, s) \nabla_{Y_s^{n,v}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds - \int_0^t K_2(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s^n ds \Big| \wedge 1 \Big] \\
& \leq \mathcal{V}_1 + \mathcal{V}_2 + \mathcal{V}_3,
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{V}_1 & := \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left[\frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} - \nabla_{Y_s^{n,v}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right] ds \right|, \\
\mathcal{V}_2 & := \mathbb{E} \left| \int_0^t K_2(t, s) \cdot \left[\sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right] \cdot v_s^n ds \right|, \\
\mathcal{V}_3 & := \mathbb{E} \left| \frac{1}{h(\varepsilon_n)} \int_0^t K_2(t, s) \cdot \sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) dW_s \right|.
\end{aligned}$$

For the term \mathcal{V}_1 , by Lemmas 5.3, 5.4, Hölder's inequality, (H1)-(H3) and (H6), we have

$$\begin{aligned}
\mathcal{V}_1 & \leq \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} - \nabla_{Y_s^{n,v}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right| ds \\
& \leq \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} \right| ds \\
& \quad + \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \frac{b(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_{s,\xi}^0}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} - \nabla_{Y_s^{n,v}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right| ds \\
& \leq L_1 \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \frac{\mathbb{W}_2(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon_n} h(\varepsilon_n)} \right| ds \\
& \quad + \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \nabla_{Y_s^{n,v}} b(s, P_s^n(u), \mathcal{L}_{X_{s,\xi}^0}) - \nabla_{Y_s^{n,v}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) du \right| ds \\
& \leq L_1 \int_0^t K_1(t, s) \cdot (\mathbb{E}|Y_s^n|^p)^{\frac{1}{p}} ds + L_5 \sqrt{\varepsilon_n} h(\varepsilon_n) \int_0^t K_1(t, s) \cdot (\mathbb{E}|Y_s^{n,v}|^p)^{\frac{1}{p}} ds \\
& \leq C_{T,L_1,L_2,\xi} \frac{1}{h(\varepsilon_n)} + C_{T,L_1,L_2,L_5,\xi} \sqrt{\varepsilon_n} h(\varepsilon_n),
\end{aligned}$$

where $P_s^n(u) := P_s^{\varepsilon_n}(u) := X_{s,\xi}^0 + u \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, u \in [0, T]$.

For the term \mathcal{V}_2 , by Lemma 3.6, Lemma 5.4, Hölder's inequality, Minkowski's inequality and (H1)-(H3), we have

$$\begin{aligned}
\mathcal{V}_2 & \leq \sqrt{N} \mathbb{E} \left(\int_0^t K_2^2(t, s) \cdot \|\sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n}) - \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\|^2 ds \right)^{\frac{1}{2}} \\
& \leq \sqrt{NL_1} \left[\mathbb{E} \left(\int_0^t K_2^2(t, s) \cdot (\varepsilon_n |h(\varepsilon_n)|^2 |Y_s^{n,v}|^2 + \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0})) ds \right)^{\frac{p}{2}} \right]^{\frac{1}{p}} \\
& \leq \sqrt{NL_1} \left(\int_0^t K_2^2(t, s) \cdot [\mathbb{E}(|\sqrt{\varepsilon_n} h(\varepsilon_n)|^p |Y_s^{n,v}|^p + \mathbb{W}_2^p(\mathcal{L}_{X_s^n}, \delta_{X_{s,\xi}^0}))]^{\frac{2}{p}} ds \right)^{\frac{1}{2}} \\
& \leq \sqrt{NL_1} \left(\int_0^t K_2^2(t, s) \cdot [\sqrt{\varepsilon_n} h(\varepsilon_n)|^p \mathbb{E}|Y_s^{n,v}|^p + \mathbb{E}|X_s^n - X_{s,\xi}^0|^p]^{\frac{2}{p}} ds \right)^{\frac{1}{2}}
\end{aligned}$$

$$\leq C_{T,N,L_1,L_2,\xi} \sqrt{\varepsilon_n} h(\varepsilon_n) + C_{T,N,L_1,L_2,\xi} \sqrt{\varepsilon_n}.$$

For the term \mathcal{V}_3 , by Lemmas 2.14, 2.15, and 5.4, Hölder's inequality, Minkowski's inequality and (H1)-(H3), we have

$$\begin{aligned} \mathcal{V}_3 &\leq \frac{1}{h(\varepsilon_n)} \mathbb{E} \left(\int_0^t K_2^2(t,s) \cdot \|\sigma(s, X_{s,\xi}^0 + \sqrt{\varepsilon_n} h(\varepsilon_n) Y_s^{n,v}, \mathcal{L}_{X_s^n})\|^2 ds \right)^{\frac{1}{2}} \\ &\leq \frac{C_{L_1}}{h(\varepsilon_n)} \left[\mathbb{E} \left(\int_0^t K_2^2(t,s) \cdot (1 + |X_{s,\xi}^0|^2 + \varepsilon_n h(\varepsilon_n) |Y_s^{n,v}|^2 + \mathbb{W}_2^2(\mathcal{L}_{X_s^n}, \delta_0)) ds \right)^{\frac{p}{2}} \right]^{\frac{1}{p}} \\ &\leq \frac{C_{L_1}}{h(\varepsilon_n)} \left(\int_0^t K_2^2(t,s) \cdot (1 + |X_{s,\xi}^0|^p + |\sqrt{\varepsilon_n} h(\varepsilon_n)|^p \mathbb{E} |Y_s^{n,v}|^p + \mathbb{E} \mathbb{W}_2^p(\mathcal{L}_{X_s^n}, \delta_0))^{\frac{2}{p}} ds \right)^{\frac{1}{2}} \\ &\leq C_{T,L_1,L_2,\xi} \frac{1}{h(\varepsilon_n)} + C_{T,N,L_1,L_2,\xi} \sqrt{\varepsilon_n} h(\varepsilon_n). \end{aligned}$$

Therefore

$$\mathbb{E}[\Psi_t(Y^{n,v}, v^n)] \leq C_{T,N,L_1,L_2,\xi} \frac{1}{h(\varepsilon_n)} + C_{T,N,L_1,L_2,\xi} \sqrt{\varepsilon_n} h(\varepsilon_n) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Hence $\mathbb{E}[\Psi_t(\psi, v)] = 0$, which implies that ψ satisfies (5.18) almost surely, for all $t \in T$. By Lemma 5.6, we have that ψ has continuous paths, so it satisfies (5.18) for all $t \in T$, almost surely,. Thus, $\mathcal{J}_{v,N}^0$ consists of unique solution of (3.11). Since this definition is independent of N , it extends to \mathcal{J}_v^0 . The proof is complete. \square

This Lemma shows that the set \mathcal{J}_v^0 corresponds to the unique solution ψ_t of (5.18).

Lemma 5.8. *Assume that (H1)-(H3) and (H6)-(H7) hold, the functional Λ in (5.19) has compact level sets.*

Proof. By Lemma 5.7, we have that

$$\Lambda(\psi) := \inf \left\{ \frac{1}{2} \int_0^T |v_s|^2 ds : v \in L^2, \psi_t = \int_0^t K_1(t,s) \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds \right\}.$$

To prove the functional Λ defined in (5.19) has compact level sets, that is for all $N > 0$, the level sets

$$L_N := \{\psi \in C_T^d : I(\phi) \leq N\}$$

are compact. Fix $N > 0$ and consider an arbitrary sequence $\{\psi^n\}_{n \in \mathbb{N}} \subset L_N$, we will show that there exists a converging subsequence such that the limit belongs to L_N .

We first prove that $\{\psi^n\}_{n \in \mathbb{N}}$ is relatively compact. According to classical Arzelà-Ascoli's Theorem, we only need to show that $\{\psi^n\}_{n \in \mathbb{N}}$ is uniformly bounded and equicontinuous. For all $n \in \mathbb{N}$ and $t \in [0, T]$, there exists $v^n \in L^2$ such that $\frac{1}{2} \int_0^T |v_t^n|^2 dt \leq N$ and $\psi^n \in \mathcal{G}_{v^n}^0$, which implies that $v^n \in \mathcal{S}_{2N}$ and ψ^n is the solution of following Volterra type McKean-Vlasov integral equation:

$$\psi_t^n = \int_0^t K_1(t,s) \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s^n ds.$$

By Lemmas 5.2 and 5.6, we obtain $\{\psi^n\}_{n \in \mathbb{N}}$ is relatively compact which ensures that L_N is relatively compact for any $N > 0$.

Next we show that $\{\psi^n\}_{n \in \mathbb{N}}$ is closed. Namely, we want to prove that if $\{\psi^n\}_{n \in \mathbb{N}}$ be a converging sequence of L_N with limit $\psi \in C_T^d$, then $\phi \in L_N$. Since $v^n \in \mathcal{S}_{2N}$ which is a compact space with respect to the weak topology. Hence there exists a subsequence $\{\psi^{n_k}\}_{k \in \mathbb{N}}$ such that v^{n_k} converges weakly in \mathcal{S}_{2N} such that v^{n_k} converges weakly in L^2 to the limit $v \in \mathcal{S}_{2N}$ and $\lim_{k \rightarrow \infty} \psi^{n_k} = \psi$.

Notice that for $v, v^{n_k} \in \mathcal{S}_{2N}$, by **(H1)-(H3)**, Hölder's inequality and Lemma 2.15 we have that

$$\begin{aligned}
& \left| \phi_t^{n_k} - \left(\int_0^t K_1(t, s) \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds \right) \right| \\
& \leq \left| \int_0^t K_1(t, s) \cdot (\nabla_{\psi_s^{n_k}} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})) ds \right| \\
& \quad + \left| \int_0^t K_1(t, s) \cdot \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \cdot (v^{n_k} - v) ds \right| \\
& \leq \int_0^t K_1(t, s) \cdot \|\nabla b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla b(s, 0, \delta_0) + \nabla b(s, 0, \delta_0)\| \cdot |\psi_s^{n_k} - \psi_s| ds \\
& \quad + \int_0^t K_1(t, s) \cdot \|\sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\| \cdot |v^{n_k} - v| ds \\
& \leq L_6 \int_0^t K_1(t, s) \cdot (|X_{s,\xi}^0| + \mathbb{W}_2(\delta_{X_{s,\xi}^0}, \delta_0)) \cdot |\psi_s^{n_k} - \psi_s| ds + L_5 \int_0^t K_1(t, s) \cdot |\psi_s^{n_k} - \psi_s| ds \\
& \quad + \sqrt{L_2} \int_0^t K_1(t, s) \cdot (1 + |X_{s,\xi}^0|^2 + \mathbb{W}_2^2(\delta_{X_{s,\xi}^0}, \delta_0))^{\frac{1}{2}} \cdot |v^{n_k} - v| ds \\
& \leq C_{T,L_2,L_5,L_6,\xi} \sup_{t \in [0,T]} |\psi_t^{n_k} - \psi_t| \quad \text{as } n_k \rightarrow \infty \quad (\lim_{k \rightarrow \infty} \phi^{n_k} = \phi) \\
& \quad + C_{T,L_2,\xi} \int_0^t K_1(t, s) \cdot |v_s^{n_k} - v_s| ds \rightarrow 0 \quad \text{as } n_k \rightarrow \infty \quad (v^{n_k} \text{ converges weakly in } L^2 \text{ to } v).
\end{aligned}$$

So we obtain easily that

$$\begin{aligned}
\psi_t &= \lim_{k \rightarrow \infty} \psi_t^{n_k} \\
&= \lim_{k \rightarrow \infty} \left(\int_0^t K_1(t, s) \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s^n ds \right) \\
&= \int_0^t K_1(t, s) \nabla_{\psi_s} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_1(t, s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds.
\end{aligned}$$

Since $v \in \mathcal{S}_{2N}$ we see $\psi \in L_N$, which concludes the proof of the closeness and therefore of the compactness of L_N . \square

Combining Lemma 5.5 and Lemma 5.8, Theorem 5.1 is a disect result of Lemma 2.11.

APPENDIX A. APPENDIX

A.1. Proof of Lemma 3.2.

Proof. We utilize the successive approximation scheme. Define recursively $(\phi^n)_{n \geq 1}$ as follows: $\phi_t^1 := \xi$, $t \in [0, T]$ and for $n = 0, 1, 2, \dots$

$$\phi_t^{n+1} := \xi + \int_0^t K_1(t, s) b(s, \phi_s^n, \delta_{X_s^0}) ds + \int_0^t K_1(t, s) \sigma(s, \phi_s^n, \delta_{X_s^0}) v_s ds, t \in [0, T]. \quad (1.21)$$

By Hölder's inequality, Lemma 2.15, **(H1)** and **(H3)** we have

$$\begin{aligned}
|\phi_t^{n+1}|^2 &\leq 3|\xi|^2 + 3 \left| \int_0^t K_1(t, s) b(s, \phi_s^n, \delta_{X_s^0}) ds \right|^2 + 3 \left| \int_0^t K_1(t, s) \sigma(s, \phi_s^n, \delta_{X_s^0}) v_s ds \right|^2 \\
&\leq 3|\xi|^2 + 3 \int_0^t K_1(t, s) \cdot |b(s, \phi_s^n, \delta_{X_s^0})|^2 ds \cdot \left| \int_0^t K_1(t, s) ds \right|^2 \\
&\quad + 3N \int_0^t K_1^2(t, s) \cdot \|\sigma(s, \phi_s^n, \delta_{X_s^0})\|^2 ds
\end{aligned}$$

$$\begin{aligned}
&\leq 3|\xi|^2 + C_{T,N,L_2} + C_{T,L_2} \int_0^t K_1(t,s) \cdot |\phi_s^n|^2 ds + C_{p,T,N,L_2} \int_0^t K_1^2(t,s) \cdot |\phi_s^n|^2 ds \\
&\quad + C_{T,N,L_2} \int_0^t |X_{s,\xi}^0|^2 ds \\
&\leq C_{T,N,L_2}(1 + |\xi|^2) + \int_0^t (C_{T,L_2} K_1(t,s) + C_{T,N,L_2} K_1^2(t,s)) \cdot |\phi_s^n|^2 ds. \tag{1.22}
\end{aligned}$$

This means

$$f_m(t) \leq C_{T,N,L_2}(1 + |\xi|^2) + \int_0^t K_{1,1}(t,s) \cdot f_m(s) ds,$$

where

$$f_m(t) := \sup_{n=1,\dots,m} |\phi_s^n|^2,$$

and $K_{1,1} := C_{T,L_2} K_1(t,s) + C_{T,N,L_2} K_1^2(t,s)$ is independent of m .

Let $R^{K_{1,1}}$ be defined by (2.4) in terms of $K_{1,2}$. By Remark 2.12, Lemma 2.3, (2.5), (2.6) and (H1), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}
\sup_{n \in \mathbb{N}} |\phi_t^n|^2 &= \lim_{m \rightarrow \infty} f_m(t) \leq C_{p,T,N,L_2} \left(|\xi|^2 + 1 + \int_0^t R^{K_{1,1}}(t,s) \cdot (|\xi|^2 + 1) ds \right) \\
&\leq C_{T,N,L_2}(1 + |\xi|^2). \tag{1.23}
\end{aligned}$$

On the other hand, we set

$$Z_{n,m}(t) := \phi_t^n - \phi_t^m$$

and

$$f(t) := \limsup_{n,m \rightarrow \infty} |Z_{n,m}(t)|^2.$$

First, by (1.23) we see

$$f(t) \leq \sup_{n,m} |Z_{n,m}(t)|^2 \leq 2 \sup_n |\phi_t^n|^p < \infty.$$

By Hölder's inequality, (H1) and (H2) we have

$$\begin{aligned}
|Z_{n+1,m+1}(t)|^2 &= |\phi_t^{n+1} - \phi_t^{m+1}|^2 \\
&\leq 2 \left| \int_0^t |b(t,s, \phi_s^n, \delta_{X_{s,\xi}^0}) - b(t,s, \phi_s^m, \delta_{X_{s,\xi}^0})| ds \right|^2 \\
&\quad + 2 \left| \int_0^t (\sigma(t,s, \phi_s^n, \delta_{X_{s,\xi}^0}) - \sigma(t,s, \phi_s^m, \delta_{X_{s,\xi}^0})) v_s ds \right|^2 \\
&\leq C_{L_1} \left| \int_0^t K_1(t,s) \cdot |\phi_s^n - \phi_s^m| ds \right|^2 \\
&\quad + 2N \left| \int_0^t K_1^2(t,s) \cdot \|\sigma(t,s, \phi_s^n, \delta_{X_{s,\xi}^0}) - \sigma(t,s, \phi_s^m, \delta_{X_{s,\xi}^0})\|^2 ds \right| \\
&\leq C_{L_1} \int_0^t K_1(t,s) \cdot |\phi_s^n - \phi_s^m|^2 ds \cdot \int_0^t K_1(t,s) ds \\
&\quad + C_{N,L_1} \int_0^t K_1^2(t,s) \cdot |\phi_s^n - \phi_s^m|^2 ds \\
&\leq C_{T,L_1} \int_0^t K_1(t,s) \cdot |\phi_s^n - \phi_s^m|^2 ds + C_{T,N,L_1} \int_0^t K_1^2(t,s) \cdot |\phi_s^n - \phi_s^m|^2 ds \\
&\leq C_{T,N,L_1} \int_0^t (K_1(t,s) + K_1^2(t,s)) \cdot |Z_{n,m}(t)|^2 ds.
\end{aligned}$$

By (1.23), **(H1)** and applying Fatou's lemma, we have

$$f(t) \leq C_{T,N,L_1} \int_0^t (K_1(t,s) + K_1^2(t,s)) \cdot f(s) ds.$$

By Remark 2.12 and Lemma 2.3, we get for almost all $t \in [0, T]$,

$$f(t) = \limsup_{n,m \rightarrow \infty} |Z_{n,m}(t)|^p = 0.$$

Hence, there exists a ϕ_t such that for almost all $t \in [0, T]$,

$$\lim_{m \rightarrow \infty} |\phi_t^n - \phi_t|^2 = 0.$$

Now taking limits in (1.21) gives the existence. The uniqueness follows from a similar calculation and Lyapunov inequality. \square

A.2. Proof of Lemma 4.3[(ii)].

Proof. By BDG's inequality, Hölder's inequality and **(H1)-(H6)** we have

$$\begin{aligned} & \mathbb{E}|Z_{t,\xi}^\varepsilon - Z_{t,\eta}^\varepsilon|^p \\ &= \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) dW_s \right. \\ &\quad \left. - \int_0^t K_1(t,s) \cdot \frac{b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon}) - b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})}{\sqrt{\varepsilon}} ds - \int_0^t K_2(t,s) \sigma(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon}) dW_s \right|^p \\ &\leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \left(\frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} - \frac{b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon}) - b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})}{\sqrt{\varepsilon}} \right) ds \right|^p \\ &\quad + 2^{p-1} \mathbb{E} \left| \int_0^t K_2(t,s) \cdot (\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon})) dW_s \right|^p \\ &\leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \left(\frac{b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} + \frac{b(s, X_{s,\xi}^\varepsilon, \delta_{X_{s,\xi}^0}) - b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})}{\sqrt{\varepsilon}} \right) ds \right. \\ &\quad \left. - \int_0^t K_1(t,s) \cdot \left(\frac{b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon}) - b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})}{\sqrt{\varepsilon}} + \frac{b(s, X_{s,\eta}^\varepsilon, \delta_{X_{s,\eta}^0}) - b(s, X_{s,\eta}^0, \delta_{X_{s,\eta}^0})}{\sqrt{\varepsilon}} \right) ds \right|^p \\ &\quad + C_p \mathbb{E} \left(\int_0^t K_2^2(t,s) \cdot \|\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon})\|^2 ds \right)^{\frac{p}{2}} \\ &\leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t,s) \cdot \left(\int_0^1 \mathbb{E} \langle D^L b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{R_s^\xi(r)}) (R_s^\xi(r), Z_{s,\xi}^\varepsilon) \rangle dr \right. \right. \\ &\quad \left. \left. + \int_0^1 \nabla_{Z_{s,\xi}^\varepsilon} b(s, X_{s,\xi}^0 + u(X_{s,\xi}^\varepsilon - X_{s,\xi}^0), \delta_{X_{s,\xi}^0}) du \right) ds \right. \\ &\quad \left. - \int_0^t K_1(t,s) \cdot \left(\int_0^1 \mathbb{E} \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)}) (R_s^\eta(r), Z_{s,\eta}^\varepsilon) \rangle dr \right. \right. \\ &\quad \left. \left. + \int_0^1 \nabla_{Z_{s,\eta}^\varepsilon} b(s, X_{s,\eta}^0 + u(X_{s,\eta}^\varepsilon - X_{s,\eta}^0), \delta_{X_{s,\eta}^0}) du \right) ds \right|^p \\ &\quad + C_p \mathbb{E} \int_0^t K_2^2(t,s) \cdot \|\sigma(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{X_{s,\xi}^\varepsilon}) - \sigma(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{X_{s,\eta}^\varepsilon})\|^p ds \cdot \left| \int_0^t K_2^2(t,s) ds \right|^{\frac{p}{2}-1} \end{aligned}$$

$$\begin{aligned}
&\leq 2^{p-1} \mathbb{E} \left| \int_0^t K_1(t, s) \cdot \left(\int_0^1 \mathbb{E} \langle D^L b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r)), Z_{s,\xi}^\varepsilon \rangle - \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r)), Z_{s,\eta}^\varepsilon \rangle \mathbf{d}r \right. \right. \\
&\quad \left. \left. + \int_0^1 (\nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \delta_{X_{t,\xi}^0}) - \nabla_{Z_{s,\eta}^\varepsilon} b(s, M_s^\eta(u), \delta_{X_{t,\eta}^0})) \mathbf{d}u \right) \mathbf{d}s \right|^p \\
&\quad + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s \\
&\leq 2^{p-1} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \mathbb{E} (\langle D^L b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r), Z_{s,\xi}^\varepsilon) - \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r)), Z_{s,\eta}^\varepsilon \rangle) \mathbf{d}r \right. \\
&\quad \left. + \int_0^1 (\nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \delta_{X_{t,\xi}^0}) - \nabla_{Z_{s,\eta}^\varepsilon} b(s, M_s^\eta(u), \delta_{X_{t,\eta}^0})) \mathbf{d}u \right|^p \mathbf{d}s \cdot \left| \int_0^t K_1(t, s) \mathbf{d}s \right|^{p-1} \\
&\quad + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s \\
&\leq C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \left| \langle D^L b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r), Z_{s,\xi}^\varepsilon) - \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r)), Z_{s,\eta}^\varepsilon \rangle \right|^p \mathbf{d}r \mathbf{d}s \right. \\
&\quad \left. + C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| \nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \delta_{X_{t,\xi}^0}) - \nabla_{Z_{s,\eta}^\varepsilon} b(s, M_s^\eta(u), \delta_{X_{t,\eta}^0}) \right|^p \mathbf{d}u \mathbf{d}s \right. \\
&\quad \left. + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s \right. \\
&\leq C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \left| \int_0^1 \mathbb{E} (\langle D^L b(s, X_{s,\xi}^\varepsilon, \mathcal{L}_{R_s^\xi(r)})(R_s^\xi(r), Z_{s,\xi}^\varepsilon) - \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r)), Z_{s,\xi}^\varepsilon \rangle) \mathbf{d}r \mathbf{d}s \right|^p \\
&\quad + C_{p,T} E \int_0^t K_1(t, s) \cdot \left| \int_0^1 \mathbb{E} (\langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r), Z_{s,\xi}^\varepsilon) - \langle D^L b(s, X_{s,\eta}^\varepsilon, \mathcal{L}_{R_s^\eta(r)})(R_s^\eta(r)), Z_{s,\eta}^\varepsilon \rangle) \mathbf{d}r \mathbf{d}s \right|^p \\
&\quad + C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| \nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \delta_{X_{t,\xi}^0}) - \nabla_{Z_{s,\eta}^\varepsilon} b(s, M_s^\eta(u), \delta_{X_{t,\eta}^0}) \right|^p \mathbf{d}u \mathbf{d}s \\
&\quad + C_{p,T} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| \nabla_{Z_{s,\eta}^\varepsilon} b(s, M_s^\eta(u), \delta_{X_{t,\eta}^0}) - \nabla_{Z_{s,\xi}^\varepsilon} b(s, M_s^\xi(u), \delta_{X_{t,\xi}^0}) \right|^p \mathbf{d}u \mathbf{d}s \\
&\quad + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s \\
&\leq C_{p,T,L_4} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 \left| |\mathcal{L}_{R_s^\xi(r)} - \mathcal{L}_{R_s^\eta(r)}| + \mathbb{W}_2(\mathcal{L}_{R_s^\xi(r)}, \mathcal{L}_{R_s^\eta(r)}) + (\mathbb{E} |R_s^\xi(r) - R_s^\eta(r)|^2)^{\frac{1}{2}} \right|^p \mathbf{d}r \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon|^p \mathbf{d}s \\
&\quad + C_{p,T,L_3} \mathbb{E} \int_0^t K_1(t, s) \cdot |Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p \mathbf{d}s + C_{p,T,L_5} \mathbb{E} \int_0^t K_1(t, s) \cdot \int_0^1 |M_s^\xi(u) - M_s^\eta(u)|^p \mathbf{d}u \cdot |Z_{s,\xi}^\varepsilon|^p \mathbf{d}s \\
&\quad + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p \mathbf{d}s + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s \\
&\leq C_{p,T,L_4} \int_0^t K_1(t, s) \cdot \mathbb{E} (|X_{s,\xi}^0 - X_{s,\eta}^0|^p + |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon|^p \mathbf{d}s \\
&\quad + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p \mathbf{d}s \\
&\quad + C_{p,T,L_5} \mathbb{E} \int_0^t K_1(t, s) \cdot (|X_{s,\xi}^0 - X_{s,\eta}^0|^p + |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p) \cdot |Z_{s,\xi}^\varepsilon|^p \mathbf{d}s \\
&\quad + C_{p,T,L_3} \int_0^t K_1(t, s) \cdot \mathbb{E} |Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p \mathbf{d}s + C_{p,T,L_1} \int_0^t K_2^2(t, s) \cdot \mathbb{E} |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \mathbf{d}s
\end{aligned}$$

$$\begin{aligned}
&\leq C_{p,T,L_4} \int_0^t K_1(t,s) \cdot \mathbb{E}|X_{s,\xi}^0 - X_{s,\eta}^0|^p \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon|^p ds + C_{p,T,L_4} \int_0^t K_1(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon|^p ds \\
&\quad + C_{p,T,L_5} \int_0^t K_1(t,s) \cdot \mathbb{E}((|X_{s,\xi}^0 - X_{s,\eta}^0|^p + |X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p) \cdot |Z_{s,\xi}^\varepsilon|^p) ds \\
&\quad + C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p ds + C_{p,T,L_1} \int_0^t K_2^2(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p ds \\
&\leq C_{p,T,L_4} \int_0^t K_1(t,s) \cdot \mathbb{E}|X_{s,\xi}^0 - X_{s,\eta}^0|^p \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon|^p ds + C_{p,T,L_4} \int_0^t K_1(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon|^p ds \\
&\quad + C_{p,T,L_5} \int_0^t K_1(t,s) \cdot (\mathbb{E}|X_{s,\xi}^0 - X_{s,\eta}^0|^{2p})^{\frac{1}{2}} \cdot (\mathbb{E}|Z_{s,\xi}^\varepsilon|^{2p})^{\frac{1}{2}} ds \\
&\quad + C_{p,T,L_5} \int_0^t K_1(t,s) \cdot (\mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^{2p})^{\frac{1}{2}} \cdot (\mathbb{E}|Z_{s,\xi}^\varepsilon|^{2p})^{\frac{1}{2}} ds \\
&\quad + C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p ds + C_{p,T,L_1} \int_0^t K_2^2(t,s) \cdot \mathbb{E}|X_{s,\xi}^\varepsilon - X_{s,\eta}^\varepsilon|^p ds \\
&\leq C_{p,T,L_1,L_2,L_4,L_5} (1 + |\xi|^p)|\xi - \eta|^p + C_{p,T,L_3} \int_0^t K_1(t,s) \cdot \mathbb{E}|Z_{s,\xi}^\varepsilon - Z_{s,\eta}^\varepsilon|^p ds,
\end{aligned}$$

where $R_s^\xi(r) := X_{s,\xi}^0 + r(X_{s,\xi}^\varepsilon - X_{s,\xi}^0)$, $R_s^\eta(r) := X_{s,\eta}^0 + r(X_{s,\eta}^\varepsilon - X_{s,\eta}^0)$, $M_s^\xi(u) := X_{s,\xi}^0 + u(X_{s,\xi}^\varepsilon - X_{s,\xi}^0)$, $M_s^\eta(u) := X_{s,\eta}^0 + u(X_{s,\eta}^\varepsilon - X_{s,\eta}^0)$, $r, u \in [0, T]$, and in the last inequality Lemma 2.14, Lemma 2.15 and Lemma 4.3 are used.

Finally, let $R^{K_1^*}$ be defined by (2.4) with $K_1^* := C_{p,T,L_3} K_1(t,s)$. From Remark 2.12 and Lemma 2.3, it follows that

$$\begin{aligned}
\mathbb{E}|Z_{t,\xi}^\varepsilon - Z_{t,\eta}^\varepsilon|^p &\leq C_{p,T,L_1,L_2,L_4,L_5} (1 + |\xi|^p)|\xi - \eta|^p \\
&\quad + \int_0^t R^{K_1^*}(s) \cdot (C_{p,T,L_1,L_2,L_4,L_5} (1 + |\xi|^p)|\xi - \eta|^p) ds \\
&\leq C_{p,T,L_1,L_2,L_3,L_4,L_5} (1 + |\xi|^p)|\xi - \eta|^p.
\end{aligned}$$

□

A.3. Proof of Theorem 5.2.

Proof. We utilize the successive approximation scheme. Define recursively $(\psi^n)_{n \geq 1}$ as follows: $\psi_t^1 := 0$, $t \in [0, T]$ and for each $n \in \mathbb{N}_+$,

$$\psi_t^{n+1} := \int_0^t K_1(t,s) \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds + \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds, t \in [0, T]. \quad (1.24)$$

By Hölder's inequality, (H1), (H3), (H6)-(H7) and Lemma 2.15 we have

$$\begin{aligned}
|\psi_t^{n+1}|^2 &\leq 2 \left| \int_0^t K_1(t,s) \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) ds \right|^2 + 2 \left| \int_0^t K_2(t,s) \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) v_s ds \right|^2 \\
&\leq 2 \int_0^t K_1(t,s) \left\| \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right\|^2 ds \cdot \left| \int_0^t K_1(t,s) ds \right| \\
&\quad + 2N \int_0^t K_1^2(t,s) \left\| \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right\|^2 ds \\
&\leq C_T \int_0^t K_1(t,s) \cdot \left\| \nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{\psi_s^n} b(s, 0, \delta_0) + \nabla_{\psi_s^n} b(s, 0, \delta_0) \right\|^2 ds \\
&\quad + C_{T,N} \int_0^t K_2^2(t,s) \cdot \left\| \sigma(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) \right\|^2 ds
\end{aligned}$$

$$\begin{aligned}
&\leq C_{T,L_5} \int_0^t K_1(t,s) \cdot |X_{s,\xi}^0|^2 \cdot |\psi_s^n|^2 ds + C_{T,L_6} \int_0^t K_1(t,s) \cdot |\psi_s^n|^2 ds \\
&\quad + C_{T,N,L_2} \int_0^t K_2^2(t,s) \cdot (1 + |X_{s,\xi}^0|^2) ds \\
&\leq C_{T,L_2,L_5,L_6} (1 + |\xi|^2) \int_0^t K_1(t,s) \cdot |\psi_s^n|^2 ds + C_{T,N,L_2} (1 + |\xi|^2) \\
&:= C_{T,N,L_2} (1 + |\xi|^2) + \int_0^t \tilde{K}_1(t,s) \cdot |\psi_s^n|^2 ds,
\end{aligned} \tag{1.25}$$

where $\tilde{K}_1(t,s) := C_{T,L_2,L_5,L_6} (1 + |\xi|^2) K_1(t,s)$. If we define

$$g_m(t) := \sup_{n=1,\dots,m} |\psi_s^n|^2,$$

then

$$g_m(t) \leq C_{T,N,L_2} (1 + |\xi|^2) + \int_0^t \tilde{K}_1(t,s) \cdot g_m(s) ds,$$

where the constant C_{T,N,L_2} is independent of m .

Let $R^{\tilde{K}_1}$ be defined by (2.4) in terms of \tilde{K}_1 . Note that by (2.3), (2.5), (2.6) and (H1), when $n = 1$ we have

$$\int_0^T R^{\tilde{K}_1}(t,s) g_m(s) ds = \int_0^T \tilde{K}_1(t,s) g_m(s) ds < \infty,$$

and

$$\begin{aligned}
&\int_0^t R^{\tilde{K}_1}(t,s) \cdot |\xi|^2 ds \\
&= \int_0^t \tilde{K}_1(t,s) \cdot |\xi|^2 ds + \int_0^t \left(\int_s^t R^{\tilde{K}_1}(t,u) \tilde{K}_1(u,s) du \right) \cdot |\xi|^2 ds \\
&= \int_0^t \tilde{K}_1(t,s) ds \cdot |\xi|^2 + \int_0^t R^{\tilde{K}_1}(t,u) \left(\int_0^u \tilde{K}_1(u,s) ds \right) du \cdot |\xi|^2 < \infty.
\end{aligned}$$

Hence, by Lemma 2.3, (2.5), (2.6) and (H1), we obtain that for almost all $t \in [0, T]$,

$$\begin{aligned}
\sup_{n \in \mathbb{N}} |\psi_t^n|^2 &= \lim_{m \rightarrow \infty} g_m(t) \leq C_{T,N,L_2} (1 + |\xi|^2) + \int_0^t R^{\tilde{K}_1}(t,s) \cdot (1 + |\xi|^2) ds \\
&\leq C_{T,N,L_2,L_5,L_6} (1 + |\xi|^2).
\end{aligned} \tag{1.26}$$

Denote

$$W_{n,m}(t) := \psi_t^n - \psi_t^m$$

and

$$g(t) := \limsup_{n,m \rightarrow \infty} |W_{n,m}(t)|^2.$$

By Hölder's inequality, (H1) and (H6)-(H7) we have

$$\begin{aligned}
|W_{n+1,m+1}(t)|^2 &= |\psi_t^{n+1} - \psi_t^{m+1}|^2 \\
&\leq \left| \int_0^t K_1(t,s) \cdot \|\nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{\psi_s^m} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\| ds \right|^2 \\
&\leq \int_0^t K_1(t,s) \cdot \|\nabla_{\psi_s^n} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla_{\psi_s^m} b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0})\|^2 ds \cdot \int_0^t K_1(t,s) ds \\
&\leq C_T \int_0^t K_1(t,s) \cdot |\psi_s^n - \psi_s^m|^2 \cdot \|\nabla b(s, X_{s,\xi}^0, \delta_{X_{s,\xi}^0}) - \nabla b(s, 0, \delta_0) + \nabla b(s, 0, \delta_0)\|^2 ds
\end{aligned}$$

$$\leq C_{T,L_2,L_5,L_6} (1 + |\xi|^2) \int_0^t K_1(t,s) \cdot W_{n,m}(s) ds.$$

By (1.26), **(H1)** and applying Fatou's lemma, we have

$$g(t) \leq C_{T,N,L_2,L_5,L_6} \int_0^t K_1(t,s) \cdot g(s) ds.$$

By using Lemma 2.3, we get for almost all $t \in [0, T]$,

$$g(t) = \limsup_{n,m \rightarrow \infty} |W_{n,m}(t)|^2 = 0.$$

Hence, there exists a ϕ_t such that for almost all $t \in [0, T]$,

$$\lim_{m \rightarrow \infty} |\psi_t^n - \psi_t|^2 = 0.$$

Now taking limit in (1.24) gives the existence. The uniqueness follows from a similar calculation. Lyapunov inequality gives desired estimate. \square

CONFLICTS OF INTERESTS

The authors declare no conflict of interests.

Acknowledgement S. Liu was supported by China Scholarship Council. Y. Hu was supported by the NSERC discovery fund and a Centennial fund of University of Alberta. H. Gao was supported in part by the NSFC Grant Nos. 12171084, the Jiangsu Provincial Scientific Research Center of Applied Mathematics under Grant No. BK20233002 and the fundamental Research Funds for the Central Universities No. RF1028623037.

REFERENCES

1. E. Abi Jaber, M. Larsson, and S. Pulido, *Affine Volterra processes*, Ann. Appl. Probab. **29** (2019), no. 5, 3155–3200.
2. J. Ackermann, T. Kruse, and L. Overbeck, *Inhomogeneous affine Volterra processes*, Stochastic Process. Appl. **150** (2022), 250–279.
3. M. A. Berger and V. J. Mizel, *Volterra equations with Itô integrals. I*, J. Integral Equations **2** (1980), no. 3, 187–245.
4. ———, *Volterra equations with Itô integrals. II*, J. Integral Equations **2** (1980), no. 4, 319–337.
5. P. Billingsley, *Convergence of probability measures*, second ed., Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons, Inc., New York, 1999, A Wiley-Interscience Publication.
6. A. Chiarini and M. Fischer, *On large deviations for small noise Itô processes*, Adv. in Appl. Probab. **46** (2014), no. 4, 1126–1147.
7. A. Dembo and O. Zeitouni, *Large deviations techniques and applications*, second ed., Applications of Mathematics (New York), vol. 38, Springer-Verlag, New York, 1998.
8. G. dos Reis, W. Salkeld, and J. Tugaut, *Freidlin-Wentzell LDP in path space for McKean-Vlasov equations and the functional iterated logarithm law*, Ann. Appl. Probab. **29** (2019), no. 3, 1487–1540.
9. J. Duan and W. Wang, *Effective dynamics of stochastic partial differential equations*, Elsevier Insights, Elsevier, Amsterdam, 2014.
10. G. Gripenberg, *On the resolvents of nonconvolution Volterra kernels*, Funkcial. Ekvac. **23** (1980), no. 1, 83–95.
11. S. Herrmann, P. Imkeller, and D. Peithmann, *Large deviations and a Kramers' type law for self-stabilizing diffusions*, Ann. Appl. Probab. **18** (2008), no. 4, 1379–1423.
12. W. Hong, S. Li, and W. Liu, *Large deviation principle for McKean-Vlasov quasilinear stochastic evolution equations*, Appl. Math. Optim. **84** (2021), no. suppl. 1, S1119–S1147.
13. A. Jacquier and A. Pannier, *Large and moderate deviations for stochastic Volterra systems*, Stochastic Process. Appl. **149** (2022), 142–187.
14. L. Jie, L. Luo, and H. Zhang, *One-dimensional McKean-Vlasov stochastic Volterra equations with Hölder diffusion coefficients*, Statist. Probab. Lett. **205** (2024), Paper No. 109970, 11.

15. M. Kac, *Foundations of kinetic theory*, Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954–1955, vol. III, Univ. California Press, Berkeley-Los Angeles, Calif., 1956, pp. 171–197.
16. ———, *Probability and related topics in physical sciences*, Lectures in Applied Mathematics, Interscience Publishers, London-New York, 1959, With special lectures by G. E. Uhlenbeck, A. R. Hibbs, and B. van der Pol.
17. E. H. Laknel, *Large deviation for stochastic Volterra equation in the Besov-Orlicz space and application*, Random Oper. Stochastic Equations **11** (2003), no. 4, 333–350.
18. M. Li, C. Huang, and Y. Hu, *Asymptotic separation for stochastic Volterra integral equations with doubly singular kernels*, Appl. Math. Lett. **113** (2021), Paper No. 106880, 7.
19. ———, *Numerical methods for stochastic Volterra integral equations with weakly singular kernels*, IMA J. Numer. Anal. **42** (2022), no. 3, 2656–2683.
20. Y. Li, R. Wang, N. Yao, and S. Zhang, *A moderate deviation principle for stochastic Volterra equation*, Statist. Probab. Lett. **122** (2017), 79–85.
21. S. Liu and H. Gao, *Volterra type McKean-Vlasov SDEs with singular kernels: Well-posedness, Propagation of Chaos and Euler schemes*, arXiv preprint arXiv:2311.06863 (2023).
22. W. Liu, Y. Song, J. Zhai, and T. Zhang, *Large and moderate deviation principles for McKean-Vlasov SDEs with jumps*, Potential Anal. **59** (2023), no. 3, 1141–1190.
23. D. Nualart and C. Rovira, *Large deviations for stochastic Volterra equations*, Bernoulli **6** (2000), no. 2, 339–355.
24. É. Pardoux and P. Protter, *Stochastic Volterra equations with anticipating coefficients*, Ann. Probab. **18** (1990), no. 4, 1635–1655.
25. D. J. Prömel and D. Scheffels, *Mean-field stochastic Volterra equations*, arXiv preprint arXiv:2307.13775 (2023).
26. ———, *Stochastic Volterra equations with Hölder diffusion coefficients*, Stochastic Process. Appl. **161** (2023), 291–315.
27. P. Protter, *Volterra equations driven by semimartingales*, Ann. Probab. **13** (1985), no. 2, 519–530.
28. H. Qiao, *The central limit theorem for stochastic Volterra equations with singular kernels*, arXiv preprint arXiv:2303.01715 (2023).
29. P. Ren and F-Y. Wang, *Bismut formula for Lions derivative of distribution dependent SDEs and applications*, J. Differential Equations **267** (2019), no. 8, 4745–4777.
30. D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, third ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 293, Springer-Verlag, Berlin, 1999.
31. A. Richard, X. Tan, and F. Yang, *Discrete-time simulation of stochastic Volterra equations*, Stochastic Process. Appl. **141** (2021), 109–138.
32. Y. Shi, T. Wang, and J. Yong, *Mean-field backward stochastic Volterra integral equations*, Discrete Contin. Dyn. Syst. Ser. B **18** (2013), no. 7, 1929–1967.
33. S. R. S. Varadhan, *Large deviations and applications*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 46, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1984.
34. H. Wu and J. Hu, *Mean field backward doubly stochastic Volterra integral equations and their applications*, Discrete Contin. Dyn. Syst. Ser. S **16** (2023), no. 5, 937–966.
35. X. Zhang, *Euler schemes and large deviations for stochastic Volterra equations with singular kernels*, J. Differential Equations **244** (2008), no. 9, 2226–2250.
36. ———, *Stochastic Volterra equations in Banach spaces and stochastic partial differential equation*, J. Funct. Anal. **258** (2010), no. 4, 1361–1425.

SCHOOL OF MATHEMATICAL SCIENCE, NANJING NORMAL UNIVERSITY, NANJING 210023, CHINA

Email address: shanqiliumath@126.com

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON T6G 2G1, CANADA

Email address: yaozhong@ualberta.ca

SCHOOL OF MATHEMATICS, SOUTHEAST UNIVERSITY, NANJING 211189, CHINA

Email address: hjgao@seu.edu.cn (Corresponding author)