On the Amplitude of Vortex Entropy: A Semiclassical Treatment

Yehao Guo,^{1,†} Dong Qiu,¹ Haiwen Liu,² and Jie Xiong^{1,*}

¹School of Physics and State Key Laboratory of Electronic Thin Films and Integrated Devices,

University of Electronic Science and Technology of China, Chengdu 611731, China

²Department of Physics and Center for Advanced Quantum Studies, Beijing Normal University,

Beijing 100875, China

Despite a long history of Nernst effect in superconductors, a satisfactory theory on its amplitude in vortex liquid phase is still absent. The central quantity of vortex Nernst signals is the entropy s_{ϕ} carried by each vortex. Here we show a semiclassical treatment based on London equation and Pippard nonlocal generalization. The derived s_{ϕ} is a function of both temperatures and magnetic fields. Its magnitude s_{ϕ}^{amp} scales with normal-state conductivity σ_n . Estimations based on our formula show good consistency with experimentally determined values. In dirty limit, the relation is further simplified into a Wiedemann-Franzlike ratio $s_{\phi}^{amp}/\sigma_n \sim k_B \ln 2/\sigma_Q$ if taking parameter values deduced from Homes' law, where $\sigma_Q = 4e^2/h$ is two-dimensional quantum conductivity. We also address related issues, including bounds to s_{ϕ} , the Nernst signal and viscosity-entropy density ratio, which are all expressed in fundamental physical constants.

[†]Contact author: <u>yehao.guo@std.uestc.edu.cn</u> ^{*}Contact author: <u>jiexiong@uestc.edu.cn</u> Introduction. In the presence of magnetic fields, the Nernst effect is the transverse electric field under longitudinal thermal gradient. The Nernst signals can be generated either by quasiparticles, or by shortlived Cooper pairs or by vortices [1-3]. While the amplitude of the former two has been understood [1,2], there is no satisfactory account of the vortex Nernst effect. Vortices, each carrying a flux quantum $\Phi_0 =$ h/2e, enter type-II superconductors under magnetic fields above the lower critical field H_{c1} . Its Nernst signal, experimentally defined as the ratio of transverse electric field to longitudinal temperature gradient, can be calculated through the relation $N = s_{\phi} \rho_{xx} / \Phi_0$, where ρ_{xx} is the longitudinal resistivity and s_{ϕ} is the entropy carried by each vortex [4]. Compared with the more familiar ρ_{xx} , the central quantity is s_{ϕ} . Stephen first investigated s_{ϕ} near H_{c1} [5]. Sergeev *et al* obtained a smaller s_{ϕ} by considering vortex cores only [6]. However, the predicted s_{ϕ} exceeded the experimental value in Nb-doped strontium titanate (Nb: SrTiO₃) by fifty times [7]. The application of Uemura's law [8] to Bi₂Sr₂CaCu₂O_{8+x} (BSCCO) predicted a constant s_{ϕ} and failed to capture the evolution of its magnitude upon doping [9]. The critical clue was pointed out by Behnia that vortex entropy may depend on the normal-state resistivity ρ_n [3]. Historically, similar situation was encountered in 1950s by Pippard, who found the penetration depth λ varied significantly with the addition of impurities, although T_C was almost the same [10]. By generalizing London equation to a nonlocal form, confirmed later by Bardeen-Cooper-Schrieffer theory microscopically [11], Pippard connected λ with the mean free path l thus the effect of impurities, obtaining successful fitting to experimental data.

Inspired by Behnia and Pippard, we present a semiclassical theory on s_{ϕ} here. The free energy density of vortices F_v is formulated based on London equation. The entropy density S can then be derived through $S = -\partial F_v/\partial T$. Divided by areal density $n_v = B/\Phi_0$, we obtain the average entropy s_{ϕ} per vortex. The normal-state conductivity σ_n is introduced by Pippard nonlocal generalization [10]. After the product of two dimensionless parameters is deduced empirically from Homes' law [12,13], we estimate the amplitude s_{ϕ}^{amp} of various superconductors, showing good consistency with experiments. In dirty limit, our result is simplified into $s_{\phi}^{amp}/\sigma_n \sim k_B \ln 2/\sigma_Q$ with two-dimensional (2D) quantum conductivity $\sigma_Q = 4e^2/h$, reminiscent of Widemann-Franz law [14]. We also addressed bounds to s_{ϕ}^{amp} , N and viscosity-entropy density ratio η/S , which can all be expressed in fundamental physical constants. The minimum s_{ϕ}^{amp} to keep superconductivity in dirty limit is $k_B \ln 2$, and the upper bound to N is $k_B \ln 2/e$. The lower bound $\eta/S \ge \pi\hbar/k_B \ln 2$ resembles the holographic bound [15].

Vortex entropy and its amplitude. The free energy density F_v of vortices consists of F_{v1} , gained outside vortex cores due to induction fields and kinetic energy of supercurrents, and F_{v2} inside cores due to the loss of condensation. F_{v1} is calculated by an integral in real space, or equivalently, summation in reciprocal space using Plancherel's theorem:

$$F_{\mathrm{v1}} = \frac{1}{2\mu_0} \int d\boldsymbol{r} \left[\boldsymbol{h}(\boldsymbol{r})^2 + \lambda^2 \left(\nabla \times \boldsymbol{h}(\boldsymbol{r}) \right)^2 \right] = \frac{1}{2\mu_0} \sum_{\boldsymbol{k}} (1 + \lambda^2 k^2) \boldsymbol{h}(\boldsymbol{k})^2 \tag{1}$$

h(k) is solved by Fourier transform of London equation $\nabla \times \nabla \times \lambda^2 h(r) + h(r) = \sum_i \Phi_0 \delta(r - R_i) \hat{z}$ in a unit area:

$$h(\mathbf{k}) = \frac{n_{\rm v} \Phi_0}{1 + \lambda^2 k^2} = \frac{B}{1 + \lambda^2 k^2}$$
(2)

 $h(\mathbf{k})$ is nonzero only when \mathbf{k} equals multiples of reciprocal lattice vector. For the sake of simplicity, we will treat it as a continuous variable, replacing the summation in Eq. (1) by an integral from $2\pi/a$ to $2\pi/2\xi$, where a and ξ are vortex lattice constant and coherence length, respectively. The constant 1 in the dominator of Eq. (2) can also be omitted, which is justified under fields high enough so that

 $\lambda/\xi \ge \lambda k \ge \lambda/a \gg 1$. This procedure is equivalent to neglect the spatial variation of $h(\mathbf{r})$ except for a homogeneous background ($\mathbf{k} = 0$). The distribution of supercurrent, carrying kinetic energy around vortices, leads to a logarithmic dependence on the ratio a/ξ or $\mu_0 H_{c2}/B$. The free energy density in Eq. (1) is thus expressed as:

$$F_{\rm v1} = \frac{B^2}{2\mu_0} + \frac{n_{\rm v}\Phi_0^2}{8\pi\mu_0\lambda^2} \ln\frac{\mu_0H_{\rm c2}}{B}$$
(1')

The loss of condensation inside the vortex core amounts to $\pi\xi^2 \mu_0 H_c^2/2$ per vortex, where H_c is the thermodynamic critical field with the relation $2\sqrt{2}\mu_0\pi\xi H_c\lambda = \Phi_0$ [16]. Therefore, the free energy density F_{v2} is:

$$F_{v2} = \frac{n_v \Phi_0^2}{8\pi\mu_0 \lambda^2 2}$$
(3)

Adding F_{v1} and F_{v2} , we obtain the total free energy density of vortices:

$$F_{\rm v} = \frac{B^2}{2\mu_0} + \frac{n_{\rm v}\Phi_0^2}{8\pi\mu_0\lambda^2} (\frac{1}{2} + \ln\frac{\mu_0 H_{\rm c2}}{B})$$
(4)

The temperature dependence is introduced through λ and H_{c2} , by assuming:

$$\frac{1}{\lambda^2(T)} = \frac{1}{\lambda^2(0)} f(\varepsilon)$$

$$H_{c2}(T) = H_{c2}(0) f(\varepsilon)$$
(5)

where $\varepsilon = 1 - T/T_c$, and $f(\varepsilon)$ is a non-decreasing function of ε with f(0) = 0. The simplest form is Ginzburg-Landau (G-L) type $f(\varepsilon) = \varepsilon$ near T_c . The vortex entropy density is calculated through $S = -\partial F_v/\partial T$. Divided by n_v , we obtain the average entropy per vortex:

$$s_{\phi}(T,B) = \frac{\Phi_0^2}{8\pi\mu_0\lambda^2(0)T_{\rm C}}\frac{{\rm d}f}{{\rm d}\varepsilon} \left(\frac{3}{2} + \ln\frac{\mu_0 H_{\rm c2}}{B}\right) \tag{6}$$

 s_{ϕ} is the function of both temperatures and magnetic fields, with its amplitude characterized by $s_{\phi}^{amp} = \Phi_0^2/8\pi\mu_0\lambda^2(0)T_c$. According to G-L theory, $df/d\varepsilon = 1$ near T_c . As the temperature is lowered, $df/d\varepsilon$ will be smaller since H_{c2} deviates from the G-L form and eventually saturates. The conclusion can also be arrived by analyzing the limit at T = 0. Starting from a superconducting ground state (S = 0) under zero field, the third law of thermodynamics requires s_{ϕ} is zero as the field is turned on, leading to $df/d\varepsilon = 0$ in Eq. (6). This implies H_{c2} should saturate as T approaching zero, consistent with the usual tendency of $H_{c2}(T)$ in experiments.

We utilize Pippard's nonlocal generalization [10] to $\lambda(0)$. The London penetration depth λ_L is defined by $\lambda_L^2 = m/n_s e^2 \mu_0$, where *m* and *e* are electron mass and charge respectively, and n_s is the density of superconducting electrons. When these quantities are those of Cooper pairs, λ_L^2 remain the same. We define two dimensionless parameters $\gamma = \hbar v_F / \xi k_B T_C$ and $\delta = n_s/n$, where v_F is Fermi velocity and *n* is the total electron density. Combining $l = v_F \tau$ and $\sigma_n = 1/\rho_n = ne^2 \tau/m$, Pippard's λ is expressed as:

$$\frac{1}{\lambda^2(0)} = \frac{1}{\lambda_L^2(0)} \frac{l}{\xi + l} = \delta \gamma \mu_0 \frac{k_{\rm B}}{\hbar} \frac{T_{\rm C}}{\rho_{\rm n}} \frac{1}{1 + l/\xi}$$
(7)

In dirty limit $l \ll \xi$, Eq. (7) is simplified into the same form as Homes' law [12,13] which gives $\delta \gamma \sim 2.8 \pm 0.6$ empirically:

$$\frac{1}{\lambda^2(0)} = \delta \gamma \mu_0 \frac{k_{\rm B}}{\hbar} \frac{T_{\rm C}}{\rho_{\rm n}} = \delta \gamma \mu_0 \frac{k_{\rm B}}{\hbar} \sigma_{\rm n} T_{\rm C} \tag{7'}$$

The only difference is that ρ_n here is taken at T = 0, while that of Homes' law is taken around T_c . With Eq. (7), the amplitude s_{ϕ}^{amp} can be rewritten as:

$$s_{\phi}^{\rm amp} = \frac{\delta\gamma\Phi_0^2}{8\pi\rho_{\rm n}}\frac{k_{\rm B}}{\hbar}\frac{1}{1+l/\xi} = \frac{\delta\gamma\,k_{\rm B}}{4}\frac{\sigma_{\rm n}}{\sigma_{\rm 0}}\frac{1}{1+l/\xi} \tag{8}$$

where $\sigma_Q = 4e^2/h$. Eq. (8) is our main result, connecting vortex entropy to normal conductivity directly.

Estimations on s_{ϕ}^{amp} . We take ρ_n as $\rho_n(T_c)$ for simplicity and use the typical value $\delta \gamma = 2.8$ given by Homes' law [12,13]. The unit of ρ_n is taken to be $\mu \Omega \cdot \text{cm}$ as commonly used in experiments, Eq. (8) can be rewritten numerically as:

$$s_{\phi}^{\text{amp}} = \frac{1}{1 + l/\xi} \frac{6.2 \times 10^{-12}}{\rho_{\text{n}}} \,\mu\Omega \cdot \text{cm} \cdot \text{J}/(\text{K} \cdot \text{m})$$
 (8')

For dirty $(l/\xi \ll 1)$ superconductors with $\rho_n \sim 100 \,\mu\Omega \cdot \text{cm}$ and c-axis constant of a few nanometers, s_{ϕ}^{amp} per sheet is around 10^{-23} J/K, on the same order of Boltzmann constant k_B [3,7,17]. For general cases, we collect parameters including ρ_n , ξ and l, calculating s_{ϕ}^{amp} according to Eq. (8'). The predicted s_{ϕ}^{amp} are shown in TABLE I, together with the experimentally determined values $s_{\phi}^{\text{exp}} = N\Phi_0/\rho_{xx}$ at Nernst peaks.

TABLE I. Related parameters, theoretically predicted s_{ϕ}^{amp} based on Eq. (8'), and experimentally determined s_{ϕ}^{exp} at Nernst peaks (the corresponding magnetic field and temperature are listed in the brackets) in FeTe_{0.6}Se_{0.4} [18], MoGe [7,19], Nb: SrTiO₃ [7], epitaxial Bi₂Sr₂CaCu₂O_{8+x} films (epi-BSCCO) [20,21], exfoliated Bi₂Sr₂CaCu₂O_{8+x} flakes (exf-BSCCO) with different doping levels [9], YBa₂Cu₃O_{7-\delta} [20,22], bulk and thin NbSe₂ flakes [23] and Nb₈₀Mo₂₀ [24,25].

$ ho_{ m n}$	l	ξ	S_{ϕ}^{amp}	s_{ϕ}^{\exp}
$[\mu\Omega \cdot cm]$	[nm]	[nm]	[10 ⁻¹⁴ J K ⁻¹ m ⁻¹]	$[10^{-14} \mathrm{J}\mathrm{K}^{-1}\mathrm{m}^{-1}]$
400	3.4	1.6	0.50	1.6 (24 T, 10 K)
150	0.3	7.7	3.9	3.1 (4 T, 3.2 K)
140	36	62	2.8	2.3 (0.06 T, 0.16 K)
140	3.4	1.42	1.3	1.6 (12 T, 77 K)
180	6	1.5	0.69	0.95 (9 T, 75 K)
250	4	1.4	0.64	0.44 (12 T, 65 K)
1000	1.8	1.6	0.29	0.053 (12 T, 50 K)
4200	0.3	3.3	0.13	0.011 (12 T, 35 K)
100	4.8	1.45	1.4	1.1 (12 T, 82 K)
110	11.7	9.1	2.5	3.5 (0.8 T, 2.82 K)
7.2	106	8.1	6.1	17 (1.2 T, 4.12 K)
5.5	5.3	39	99	100 (0.03 T, 3.5 K)
	$[\mu\Omega \cdot cm]$ 400 150 140 140 140 180 250 1000 4200 100 110 7.2	$\begin{array}{c c} [\mu\Omega\cdot cm] & [nm] \\ \hline 400 & 3.4 \\ 150 & 0.3 \\ 140 & 36 \\ 140 & 3.4 \\ 180 & 6 \\ 250 & 4 \\ 1000 & 1.8 \\ 4200 & 0.3 \\ 100 & 4.8 \\ 110 & 11.7 \\ 7.2 & 106 \\ \end{array}$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	$[\mu\Omega \cdot cm]$ $[nm]$ $[nm]$ $[10^{-14} J K^{-1}m^{-1}]$ 4003.41.60.501500.37.73.914036622.81403.41.421.318061.50.6925041.40.6410001.81.60.2942000.33.30.131004.81.451.411011.79.12.57.21068.16.1

Despite the wide range of ρ_n , the difference between s_{ϕ}^{amp} and s_{ϕ}^{exp} in each material is within one order. The exceptions are underdoped BSCCO flakes with $T_C \leq 56$ K. One possible reason is that when temperatures are close to T_C , the thermal activation of Bogoliubov quasiparticles is significant. While in dirty materials these quasiparticles contribute few Nernst signals due to short mean free path [2], they could produce remarkable ρ_{xx} . Therefore, $s_{\phi}^{exp} = N\Phi_0/\rho_{xx}$ would be smaller than the mere value of vortex entropy. In fact, in Ref. [9] the temperature of the Nernst peak gradually approaches and exceeds T_C eventually as the doping is decreased.

Bounds of s_{ϕ}^{amp} and N. In dirty limit $(l/\xi \ll 1)$, Eq. (7') leads to a ratio $s_{\phi}^{\text{amp}}/\sigma_{n} = 0.7k_{\text{B}}/\sigma_{\text{Q}} \approx k_{\text{B}} \ln 2/\sigma_{\text{Q}}$ reminiscent of Wiedemann-Franz law [14], implying a particle-transport picture of 2e

charges carrying entropy and conductivity quanta. It is interesting that $k_B \ln 2$ is the quantum of entropy cost by losing one bit of information, while σ_Q is the scale of dissipation per transverse mode in the conductor (differed by a factor 2) according to Landauer's transport formula [26]. The combination of these two scales reminds of Landauer's principle [27] that *erasing information requires energy to be dissipated as heat*. In the present case, the heat should be produced through the quantum dissipator σ_Q as each bit of information is erased.

According to boson-vortex duality in 2D disordered superconductors [28,29], superconductorinsulator transition takes place at the critical conductivity $\sigma_c = \sigma_Q$ (the length disappears in units of σ in 2D). Therefore, the lower bound to keep superconductivity ($\sigma_n \ge \sigma_0$) would be:

$$s_{\phi}^{\rm amp} \ge \frac{\delta \gamma}{4} k_{\rm B} \tag{9}$$

If taking $\delta \gamma = 2.8$, then Eq. (9) coincides with the value proposed in Ref. [3], which is reasonable since $k_{\rm B} \ln 2$ should be the minimum entropy to discern a vortex structure, viz., regions inside or outside the core. If the entropy was smaller, a vortex cannot be defined due to lack of information, and the system would become normal. If the critical conductivity $\sigma_{\rm c} \neq \sigma_{\rm Q}$, which, for example, can be tuned in Josephson junction arrays by the ratio of Josephson coupling to Coulomb blockade energy [30], the bound can also be larger or smaller than $k_{\rm B} \ln 2$. For cleaner materials with $l \geq \xi$, the entropy can also be smaller due to the factor $1/(1 + l/\xi)$ in Eq. (8).

The vortex Nernst signal can be expressed as $N = s_{\phi}\rho_{xx}/\Phi_0$ [4]. For ideally free vortex liquid, the flux flow resistance [31–33] is $\rho_f = \rho_n H/H_{c2}$, while for pinned liquid, there is also an exponential factor $\exp(-\varepsilon/T)$ characterizing the thermal activation of mobile vortices [34], where the energy barrier $\varepsilon > 0$ depends on pinning details. Therefore, $\rho_{xx} = \rho_f \exp(-\varepsilon/T) < \rho_f$. Considering $1/(1 + l/\xi) < 1$ and $df/d\varepsilon \le 1$, N follows the inequation:

$$N < \frac{s_{\phi}\rho_{\rm f}}{\Phi_0} < \frac{\delta\gamma}{8} \frac{k_B}{e} \left(\frac{3}{2} + \ln\frac{\mu_0 H_{\rm c2}}{B}\right) \frac{B}{\mu_0 H_{\rm c2}}$$
(10)

The right-hand side is an increasing function on $B/\mu_0 H_{c2}$, reaching its maximum at $\mu_0 H_{c2}$. However, our result based on London equation neglects the depletion of n_s outside cores due to finite momentum. The inaccuracy could become more significant as the field approaching H_{c2} , so we resort to G-L theory at such limit, which takes account of this nonlinear effect. The G-L free energy density [35] is:

$$F_{\rm GL} = F_{\rm n} + \frac{B^2}{2\mu_0} - \frac{1}{2\mu_0} \frac{(\mu_0 H_{\rm c2} - B)^2}{1 + (2\kappa^2 - 1)\beta_{\rm A}}$$

$$\sim F_{\rm n} + \frac{B^2}{2\mu_0} - \frac{1}{2\mu_0} \frac{(\mu_0 H_{\rm c2} - B)^2}{2\kappa^2}$$
(11)

where ~ is associated with G-L parameter $\kappa \sim \lambda/\xi \gg 1$ and the lattice parameter $\beta_A \sim 1$ [35]. However, it should be noted that F_{GL} contains contributions from both thermally activated Bogoliubov quasiparticles and vortices. The former corresponds to $F_n - \mu_0 H_c^2/2$ under zero field. F_v is extracted from F_{GL} by subtracting $F_n - \mu_0 H_c^2/2$, as recently used to calculate chemical penitential of vortices [36]. Combined with the relation $H_{c2} = \sqrt{2}\kappa H_c = \Phi_0/2\pi\xi^2$, we obtain F_v of vortices and the corresponding s_{ϕ} :

$$F_{\rm v} = \frac{B^2}{2\mu_0} \left(1 - \frac{1}{2\kappa^2}\right) + \frac{n_{\rm v}\Phi_0^2}{4\pi\lambda^2}$$
$$s_{\phi} = \frac{\Phi_0^2}{4\pi\mu_0\lambda^2(0)T_{\rm c}}\frac{\mathrm{d}f}{\mathrm{d}\varepsilon}$$
(12)

It turns out that Eq. (12) can be extrapolated from Eq. (6) by taking $B = \mu_0 H_{c2}$ and replacing 3/2 with

2. Therefore, the upper bound of N is:

$$N < \frac{\delta \gamma}{4} \frac{k_B}{e} \tag{10'}$$

For $\delta\gamma$ of order unity, Eq. (10') is of order 10 μ V/K. If taking $\delta\gamma = 2.8$, then $N < k_{\rm B} \ln 2/e$.

The lower bound of viscosity-entropy density ratio. The viscosity-entropy density ratio η/S of vortex liquid was proposed to have a minimum similar with those of classical liquids [3,37]. Here we address this problem. The Bardeen-Stephen (BS) viscosity [31] is $\eta = \mu_0 H_{c2} \Phi_0 / \rho_n$. The entropy density is obtained through $S = n_v s_{\phi}$ after inserting Eq. (8) into Eq. (6). Considering $df/d\varepsilon \leq 1$, the viscosity-entropy density ratio is:

$$\frac{\eta}{S} \ge \frac{8\pi}{\delta\gamma} \frac{\hbar}{k_{\rm B}} \left[\frac{B}{\mu_0 H_{\rm c2}} \left(\frac{3}{2} + \ln \frac{\mu_0 H_{\rm c2}}{B} \right) \right]^{-1}$$
(13)

The right-hand side is a decreasing function on $B/\mu_0 H_{c2}$, reaching its minimum as $B = \mu_0 H_{c2}$. Similar with Eq. (10), the constant 3/2 is replaced by 2 at H_{c2} . Therefore, the minimum is, if taking $\delta \gamma = 2.8$:

$$\frac{\eta}{S} \ge \frac{4\pi}{\delta\gamma} \frac{\hbar}{k_{\rm B}} \sim \frac{\pi\hbar}{k_{\rm B} \ln 2} \tag{13'}$$

which suggests that if $k_{\rm B} \ln 2$ sets the scale of entropy, then the viscosity would be set by $\pi\hbar = h/2$, half of the Planck constant. Since BS viscosity is applicable to the free vortex liquid, the general viscosity could be enhanced due to vortex-vortex and vortex-pinning interaction, and Eq. (13) still holds. The lower bound resembles the holographic one in relativistic hydrodynamics [15] where the coefficient of $\hbar/k_{\rm B}$ is $1/4\pi$, and that of classical liquid [37] which has an additional square root of proton-electron mass ratio.

Discussion. The low-field s_{ϕ} by Stephen [5] can be reproduced by replacing the integral cutoff at lattice constant a in Eq. (1) with penetration depth λ , giving the factor $2\ln\kappa$ rather than $\ln(\mu_0 H_{c2}/B)$. Sergeev's result [6] is the same as our expression for amplitude s_{ϕ}^{amp} . Both results are independent of temperatures, fields, and resistivities. The finite s_{ϕ} close to H_{c2} in Eq. (12) contrasts with that obtained by Maki and Caroli [38–40], where s_{ϕ} should approach zero at $B = \mu_0 H_{c2}$. Their result can be reproduced if calculating $S = -\partial F_{GL}/\partial T$ directly from Eq. (11), then we would obtain $S \propto (\mu_0 H_{c2} - B)$. The discrepancy can be resolved by noting that S calculated from F_{GL} also contains the contribution from thermally activated quasiparticles. To estimate vortex entropy, we have utilized Homes' law to deduce $\delta\gamma$. The procedure can also be reversed, for example, one obtains s_{ϕ} , the parameter γ and characteristic length scales like ξ and l through thermoelectric transport and scanning-tunneling microscopy, the superconducting portion δ can then be estimated.

In summary, we have derived the average entropy s_{ϕ} of each vortex through a thermodynamic method. We further relate its amplitude s_{ϕ}^{amp} with the normal-state resistivity. The consistency with experimental values suggests our consideration is reasonable. The inclusion of resistivity also enables us to estimate bounds to vortex entropy, Nernst signals and viscosity-entropy density ratio. The factor ln 2 is incorporated from $\delta \gamma = 2.8$ given empirically by Homes' law [12,13]. Therefore, the last piece of the whole picture might reside in the mystery of Homes' law.

Acknowledgment. We are grateful to discussions with K. Behnia, G. E. Volovik, Ding Zhang, and Yang Wang. The work was supported by the National Key Research and Development Program of China (2021YFA0718800).

References:

- K. Behnia, The Nernst effect and the boundaries of the Fermi liquid picture, J. Phys.: Condens. Matter 21, 113101 (2009).
- [2] K. Behnia and H. Aubin, Nernst effect in metals and superconductors: a review of concepts and experiments, Rep. Prog. Phys. 79, 046502 (2016).
- [3] K. Behnia, Nernst response, viscosity and mobile entropy in vortex liquids, J. Phys.: Condens. Matter 35, 074003 (2023).
- [4] R. P. Huebener, Magnetic Flux Structures in Superconductors (Springer-Berlin, Heidelberg, 2013).
- [5] M. J. Stephen, Galvanomagnetic and Related Effects in Type-II Superconductors, Phys. Rev. Lett. 16, 801 (1966).
- [6] A. Sergeev, M. Reizer, and V. Mitin, Thermomagnetic vortex transport: Transport entropy revisited, Europhys. Lett. 92, 27003 (2010).
- [7] C. W. Rischau, Y. Li, B. Fauqué, H. Inoue, M. Kim, C. Bell, H. Y. Hwang, A. Kapitulnik, and K. Behnia, Universal Bound to the Amplitude of the Vortex Nernst Signal in Superconductors, Phys. Rev. Lett. 126, 077001 (2021).
- [8] Y. J. Uemura et al., Universal Correlations between $T_{\rm C}$ and $n_{\rm s}/m^*$ (Carrier Density over Effective Mass) in High- $T_{\rm C}$ Cuprate Superconductors, Phys. Rev. Lett. **62**, 2317 (1989).
- [9] S. Hu, J. Qiao, G. Gu, Q.-K. Xue, and D. Zhang, Vortex entropy and superconducting fluctuations in ultrathin underdoped Bi₂Sr₂CaCu₂O_{8+x} superconductor, Nat. Commun. 15, 4818 (2024).
- [10] A. B. Pippard and W. L. Bragg, An experimental and theoretical study of the relation between magnetic field and current in a superconductor, Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 216, 547 (1953).
- [11] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of Superconductivity, Phys. Rev. 108, 1175 (1957).
- [12] C. C. Homes *et al.*, A universal scaling relation in high- temperature superconductors, Nature 430, 539 (2004).
- [13] D. N. Basov, R. D. Averitt, D. Van Der Marel, M. Dressel, and K. Haule, Electrodynamics of correlated electron materials, Rev. Mod. Phys. 83, 471 (2011).
- [14] N. W. Ashcroft and N. D. Mermin, Solid State Physics (Cengage, 1976).
- [15] P. K. Kovtun, D. T. Son, and A. O. Starinets, Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics, Phys. Rev. Lett. 94, 111601 (2005).
- [16] M. Tinkham, Introduction to Superconductivity (Courier Corporation, 2004).
- [17] R. P. Huebener and H.-C. Ri, Vortex transport entropy in cuprate superconductors and Boltzmann constant, Physica C: Superconductivity and Its Applications 591, 1353975 (2021).
- [18] A. Pourret, L. Malone, A. B. Antunes, C. S. Yadav, P. L. Paulose, B. Fauque, and K. Behnia, Strong correlation and low carrier density in Fe_{1+y}Te_{0.6}Se_{0.4} as seen from its thermoelectric response, Phys. Rev. B 83, 020504 (2011).
- [19] A. Rogachev, A. T. Bollinger, and A. Bezryadin, Influence of High Magnetic Fields on the Superconducting Transition of One-Dimensional Nb and MoGe Nanowires, Phys. Rev. Lett. 94, 017004 (2005).
- [20] H.-C. Ri, R. Gross, F. Gollnik, A. Beck, R. P. Huebener, P. Wagner, and H. Adrian, Nernst, Seebeck, and Hall effects in the mixed state of YBa₂Cu₃O_{7-δ} and Bi₂Sr₂CaCu₂O_{8+x} thin films: A comparative study, Phys. Rev. B 50, 3312 (1994).
- [21] T. Honma, K. Yamaya, F. Minami, and S. Takekawa, Temperature dependences of anisotropic

resistivity in the a-b plane and Hall coefficient in a single-crystal $Bi_2Sr_2CaCu_2O_y$, Physica C: Superconductivity **176**, 209 (1991).

- [22] A. L. Solovjov, H. U. Habermeier, and T. Haage, Fluctuation conductivity in YBa₂Cu₃O_{7-y} films with different oxygen content. I. Optimally and lightly doped YBCO films, Low Temp. Phys. 28, 17 (2002).
- [23] X.-Q. Li, Z.-L. Li, J.-J. Zhao, and X.-S. Wu, Electrical and thermoelectric study of twodimensional crystal of NbSe₂, Chinese Phys. B 29, 087402 (2020).
- [24] O. L. De Lange and F. A. Otter, Flux flow effects in a nearly reversible type II superconductor, J. Low Temp. Phys. 18, 31 (1975).
- [25] R. A. French and J. Lowell, Effect of Temperature and of Substitutional Mo on the Magnetic Behavior of Superconducting Nb, Phys. Rev. 173, 504 (1968).
- [26] S. Datta, Electronic Transport in Mesoscopic Systems (Cambridge Univ. Press, 2009).
- [27] C. H. Bennett, Notes on Landauer's principle, reversible computation, and Maxwell's Demon, Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics 34, 501 (2003).
- [28] M. P. A. Fisher, Quantum phase transitions in disordered two-dimensional superconductors, Phys. Rev. Lett. 65, 923 (1990).
- [29] M. P. A. Fisher and D. H. Lee, Correspondence between two-dimensional bosons and a bulk superconductor in a magnetic field, Phys. Rev. B 39, 2756 (1989).
- [30] R. Fazio, Quantum phase transitions and vortex dynamics in superconducting networks, Physics Reports 355, 235 (2001).
- [31] J. Bardeen and M. J. Stephen, Theory of the Motion of Vortices in Superconductors, Phys. Rev. 140, A1197 (1965).
- [32] P. Nozières and W. F. Vinen, The motion of flux lines in type II superconductors, The Philosophical Magazine: A Journal of Theoretical Experimental and Applied Physics 14, 667 (1966).
- [33] Y. B. Kim, C. F. Hempstead, and A. R. Strnad, Flux-Flow Resistance in Type-II Superconductors, Phys. Rev. 139, A1163 (1965).
- [34] M. V. Feigel'man, V. B. Geshkenbein, and A. I. Larkin, Pinning and creep in layered superconductors, Physica C: Superconductivity 167, 177 (1990).
- [35] P.-G. De Gennes, Superconductivity of Metals and Alloys (CRC press, Boca Raton, 2018).
- [36] A. Auerbach and D. Arovas, Hall anomaly and moving vortex charge in layered superconductors, SciPost Phys. 8, 061 (2020).
- [37] K. Trachenko and V. V. Brazhkin, Minimal quantum viscosity from fundamental physical constants, Sci. Adv. 6, eaba3747 (2020).
- [38] K. Maki, Motion of the vortex lattice in a dirty type II superconductor, J. Low Temp. Phys. 1, 45 (1969).
- [39] C. Caroli and K. Maki, Motion of the Vortex Structure in Type-II Superconductors in High Magnetic Field, Phys. Rev. 164, 591 (1967).
- [40] K. Maki, Vortex motion in superconductors, Physica 55, 124 (1971).