Model-assisted and Knowledge-guided Transfer Regression for the Underrepresented Population

Doudou Zhou¹, Mengyan Li^{*2}, Tianxi Cai³, and Molei Liu^{†4}

¹Department of Statistics and Data Science, National University of Singapore

²Department of Mathematical Sciences, Bentley University

³Department of Biostatistics, Harvard T.H. Chan School of Public Health. ⁴Department of Biostatistics, Columbia Mailman School of Public Health.

Abstract

Covariate shift and outcome model heterogeneity are two prominent challenges in leveraging external sources to improve risk modeling for underrepresented cohorts in paucity of accurate labels. We consider the transfer learning problem targeting some unlabeled minority sample encountering (i) covariate shift to the labeled source sample collected on a different cohort; and (ii) outcome model heterogeneity with some majority sample informative to the targeted minority model. In this scenario, we develop a novel model-assisted and knowledge-guided transfer learning targeting underrepresented population (MAKEUP) approach for high-dimensional regression models. Our MAKEUP approach includes a model-assisted debiasing step in response to the covariate shift, accompanied by a knowledge-guided sparsifying procedure leveraging the majority data to enhance learning on the minority group. We also develop a model selection method to avoid negative knowledge transfer that can work in the absence of gold standard labels on the target sample. Theoretical analyses show that MAKEUP provides efficient estimation for the target model on the minority group. It maintains robustness to the high complexity and misspecification of the nuisance models used for covariate shift correction, as well as adaptivity to the model heterogeneity and potential negative transfer between the majority and minority groups. Numerical studies demonstrate similar advantages in finite sample settings over existing approaches. We also illustrate our approach through a real-world application about the transfer learning of Type II diabetes genetic risk models on some underrepresented ancestry group.

[∗]Zhou and Li are equal contributors.

[†]Email: ml4890@cumc.columbia.edu

Keywords: Covariate shift; Sampling disparity; Model heterogeneity; Double robustness; Model calibration; Transfer Lasso.

1 Introduction

1.1 Background

Genetic disease risk modeling plays a critical role in decision-making and knowledge discovery, with wide-ranging applications, particularly in precision medicine [\(Moons et al., 2012;](#page-27-0) [Amarasingham et al., 2014;](#page-24-0) [Matheny et al., 2010\)](#page-27-1). The extensive adoption of electronic health records (EHRs) provides abundant data for developing real-world predictive models that incorporate a broad spectrum of demographic and clinical information. When EHRs are linked with specimen bio-repositories, they form biobank datasets containing detailed genomic and phenotypic information at the individual level. This integration facilitates the development and implementation of personalized risk prediction models, such as polygenic risk scores, which significantly enhance prediction accuracy [\(Steyerberg et al., 2013;](#page-27-2) [Parikh](#page-27-3) [et al., 2016;](#page-27-3) [Li et al., 2020\)](#page-26-0).

Despite their potential, fully leveraging EHR biobank data for precision medicine research remains challenging due to several key obstacles. The high dimensionality of biobank data, which includes codified diagnosis, medication, procedure information, free-text clinical notes, lab results, and large-scale genomic data, complicates analysis. The lack of gold standard labels and the presence of noisy outcome data further hinder accurate disease risk modeling and prediction. Establishing these gold standard labels often requires manual chart review, as diagnostic codes are frequently insufficient or unavailable. Another significant challenge is ensuring fairness in prediction algorithms for minority populations, whose smaller sample sizes compared to the majority can lead to biased models and concerns about generalizability for algorithms trained to optimize an overall performance. For instance, in the MGB biobank data, around 90% subjects are European descent and only 5% are African American. Such disparity in data availability naturally leads to the unfair results that the risk prediction accuracy on the majority European population is much higher than the minority ancestry groups [\(West et al., 2017;](#page-28-0) [Martin et al., 2019\)](#page-26-1). These challenges motivate the exploration of transfer learning methods to develop risk prediction models for underrepresented populations by borrowing information from both larger majority populations and labeled source populations. For instance, in our real-world study using the Mass General Brigham (MGB) biobank [\(Castro et al., 2022\)](#page-24-1), we aim to leverage labeled source data from biobank participants enrolled in 2014, across both white and non-white subgroups, to build a genetic risk prediction model for type 2 diabetes in the unlabeled non-white biobank population of 2021.

In such transfer studies, **covariate shift** between the two cohorts arises as a key due to various reasons. For instance, EHR cohorts can vary in their coding systems and versions [\(Guo et al., 2021\)](#page-25-0) and notably, MGB underwent a change on its coding system to EPIC around 2015, which is likely to result in the distributional shift of EHR features. Also, covariate shifts can be caused by the demographic variation of the cohort at different institutions or years, the emergence of new diseases, and the evolving practice patterns of healthcare professionals [\(Braithwaite, 2018\)](#page-24-2). For example, the deterioration of model performance was observed in acute kidney injury prediction at Veterans Affairs (VA) hospitals from 2003 to 2012 due to temporal shift of its observational cohort [\(Davis et al., 2017\)](#page-25-1). As a consequence, capturing covariate shift and correcting its potential bias play an essential role in effectively porting and generalizing data resources and knowledge across multiple EHR and biobank cohorts.

To improve fairness, one important strategy is to leverage knowledge of the model obtained from the large majority sample to assist learning on the minority groups. Recent studies like [Verma et al. \(2023\)](#page-28-1) established that a large proportion of the genetic effects on a wide range of phenotypes tend to be consistent and transferable across different ancestry groups. This reveals the potential of knowledge transfer to improve the prediction accuracy on underrepresented ancestry groups and achieve better fairness in personalized medicine. Nevertheless, the presence of outcome model heterogeneity across different ancestry groups and phenotypes is universal and non-negligible [\(De Lillo et al., 2021;](#page-25-2) [Wang et al.,](#page-28-2) [2024,](#page-28-2) e.g.), which could induce bias and negative knowledge transfer. This calls for knowledge transfer approaches efficient in leveraging informative majority sources while staying robust to potential model shifts between the sub-populations.

1.2 Related literature

Both covariate shift and outcome model shift have been studied as the main challenges in transfer learning literature under different scenarios. To address covariate shift, [Huang](#page-25-3) [et al. \(2007\)](#page-25-3) developed an importance weighting approach based on kernel mean matching to characterize the density ratio between the source and target and use it to reweight the source sample. [Reddi et al. \(2015\)](#page-27-4) proposed a robust importance weighting strategy to achieve good bias and variance trade-off in the presence of strong covariate shift. [Wang \(2023\)](#page-28-3) developed a novel pseudo-labeling approach to optimize the performance of kernel ridge regression. Their method is adaptive to unknown and potentially severe covariate shift. More close to our work, recent studies like [Chakrabortty et al. \(2019\)](#page-24-3), [Liu et al. \(2023\)](#page-26-2), [Qiu et al.](#page-27-5) [\(2023\)](#page-27-5), [Zhou et al. \(2024\)](#page-28-4), and [Tian et al. \(2024\)](#page-28-5) considered a doubly robust framework that augments the importance weighted estimating equations with an imputation model for the outcome on target, to achieve robustness to model misspecification and excessive estimation errors. Nevertheless, these existing methods address neither high-dimensional outcome models nor the disparity sampling issue highlighted in the proposed framework.

The proposed method also relates to the knowledge-guided transfer learning literature

addressing outcome model shift that has been explored recently. In specific, [Bastani \(2021\)](#page-24-4) considered the scenario that the source (majority) and target (minority) model coefficients have a relatively small difference and proposed a source-coefficient-guided shrinkage estimation approach for the target model. [Li et al. \(2022\)](#page-26-3) focused on the optimal transfer learning of linear models under a similar assumption that the difference between source and target models is smaller or sparser than the outcome model itself. They also proposed a data-driven procedure to ensure effective and non-negative transfer in the presence of non-informative sources. [Tian and Feng \(2023\)](#page-27-6) extended this so-called Trans-Lasso framework to accommodate high-dimensional sparse generalized linear models (GLMs) and [Lin and Reimherr](#page-26-4) [\(2024\)](#page-26-4) studied the transfer smooth regression setting using kernel ridge methods with a similar spirit. In addition, [Li et al. \(2023\)](#page-26-5) considered a federated learning setup targeting the underrepresented group and leveraged the idea of Trans-Lasso to improve the estimation efficiency. Moreover, [Cai et al. \(2024\)](#page-24-5) and [He et al. \(2024\)](#page-25-4) developed knowledge transferring methods addressing shifts of both the covariates and outcome models. Different from all these works, we consider a more challenging setup in which neither the majority nor the minority data in the target sample includes any observations of the outcome Y . In this setup, existing approaches are not applicable, as they rely on labeled samples to conduct the knowledge transfer regression with shrinkage to source models.

Our theoretical framework relates to the literature on model-assisted semiparametric estimation with high-dimensional or highly complex nuisance models. In this direction, the doubly robust estimation of conditional models or heterogeneous treatment effects (HTE) shares more closely related objectives with our approach. For example, [Fan et al. \(2022\)](#page-25-5) and [Semenova and Chernozhukov \(2021\)](#page-27-7) extended the double machine learning (DML) framework [\(Chernozhukov et al., 2018\)](#page-24-6) to infer the HTE on low-dimensional effect modifiers with high-dimensional adjustment covariates. [Kennedy \(2023\)](#page-26-6) studied the DML estimation of HTE models with high-complexity and nontrivial structures such as smoothness or sparsity. [Kato \(2024\)](#page-25-6) extended this approach for debiased inference of high-dimensional parametric HTE functions. [Baybutt and Navjeevan \(2023\)](#page-24-7) incorporated the idea of model calibration [\(Tan, 2020,](#page-27-8) e.g.) for low-dimensional smooth estimation of HTE with potentially misspecified high-dimensional parametric nuisance models. To our best knowledge, none of these existing strategies can be adopted to our setup with high-dimensional sparse target model under covariate shift while simultaneously fulfilling the following two desirable properties: (i) double robustness to the misspecification of nuisance models; and (ii) double robustness to the slow convergence rates (i.e., poor sparsity level) of the nuisance models. This is due to the technical difficulty of realizing asymptotic linearity on the sparse estimators with excessive regularization errors.

1.3 Contribution

Our problem setup to be introduced in Section [2](#page-4-0) presents a unique combination of three main challenges: (1) covariate shift between the source and target cohorts, (2) sampling disparity on the minority subgroup accompanied by model heterogeneity between the majority and minority, and (3) high-dimensionality. To address these challenges simultaneously, we propose a novel approach called Model-Assisted and KnowledgE-guided transfer regression targeting Underrepresented Population (MAKEUP). It includes a calibrated model-assisted estimation procedure to achieve robust and efficient correction of covariate shift, followed by an adaptive and safe knowledge transfer approach to assist risk modeling for the targeted underrepresented sub-population with the majority sample. The main methodological contributions of our developments are summarized as follows.

First, for covariate shift correction when both the predictors and adjustment features are of high dimensionality, our approach ensures simultaneous model-double-robustness and rate-double robustness through a novel strategy consisting of debiasing, calibration, and sparsifying steps. Specifically, our method is proven to be consistent when either the importance weighting or imputation models with some parametric forms of the high-dimensional adjustment covariates are correctly specified. Additionally, these two nuisance estimators have only a "second-order" impact on the convergence rate of the resulting sparse estimator for the outcome model, in a similar sense as the rate-doubly-robust property [\(Kennedy,](#page-26-6) [2023;](#page-26-6) [Smucler et al., 2019\)](#page-27-9). Such a desirable robustness property cannot be achieved through existing doubly robust methods due to the excessive regularization bias and asymptotic nonlinearity of both the nuisance and target estimators.

Second, our approach achieves adaptive knowledge transfer from the majority subgroup to the targeted underrepresented subgroup without any collected labels. This is accomplished by incorporating the idea of source-guided shrinkage with our doubly robust covariate shift correction procedure. Previous methods like Trans-Lasso [\(Li et al., 2022\)](#page-26-3) cannot be applied in our setup since the regression problem on the unlabeled target cannot be formed in a straightforward way preserving robustness to the misspecification and excessive errors of the nuisance models. We address this challenge by introducing a debiased-coefficient construction for sparse estimation of the target model. It inherits the model- and rate-robust properties from our covariate shift correction and allows the natural incorporation of source-guided shrinkage and negative transfer protection, which achieve adaptive and effective knowledge transfer.

2 Problem Setup

Denote $Y \in \mathbb{R}$ as the outcome of interest, $R \in \{0,1\}$ as the subgroup indicator with $R = 1$ indicating the majority group and $R = 0$ being the underrepresented minority group, and

 $\mathbf{Z} = (\mathbf{X}^{\mathsf{T}}, \mathbf{W}^{\mathsf{T}})^{\mathsf{T}} \in \mathbb{R}^{q+p}$ as the covariate vector, where $\mathbf{X} = (X_1, \dots, X_q)^{\mathsf{T}}$ is a vector of risk factors to predict Y, with the first element $X_1 = 1$, and $\mathbf{W} = (W_1, \ldots, W_p)^\intercal$ denotes the auxiliary covariates informative to Y but not as part of the risk factors of interest. Both X and W can be of high dimension. Our goal is to construct a prediction model of $Y \sim X$ for the underrepresented group $(R = 0)$ on some target population \mathcal{T} :

$$
\mathbb{E}_{\tau}[Y \mid \mathbf{X}, R = 0] = g(\mathbf{X}^{\mathsf{T}} \bar{\boldsymbol{\beta}}^{[0]}), \tag{1}
$$

where $\mathbb{E}_{\tau}[\cdot]$ denotes the expectation operator on $\mathcal{T}, g(\cdot)$ is a known link function, and $\bar{\boldsymbol{\beta}}^{[0]}$ is the high-dimensional sparse coefficient vector. We do not require the working model [\(1\)](#page-5-0) to hold and define the population parameter $\bar{\beta}^{[0]}$ as the solution to

$$
\mathbb{E}_{\tau}[\mathbf{X}\{Y - g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\} \mid R = 0] = \mathbf{0}.
$$
 (2)

This solution corresponds to the ordinary least square regression when $g(x) = x$ and logistic regression when $g(x) = 1/(1 + e^{-x})$.

We assume that the outcome Y is only labeled or collected on a source cohort S but not observed on the target \mathcal{T} , yielding an observed dataset with a total size n and in the form

$$
\mathscr{D} = \{ \mathbf{D}_i = (S_i Y_i, \mathbf{Z}_i^{\mathsf{T}}, S_i, R_i)^{\mathsf{T}} : i = 1, 2, \dots, n \},
$$

where $R_i \in \{0,1\}$ indicates the majority and minority groups and S_i is the source and target indicator that $S_i = 1$ for subjects i from the source cohort S and $S_i = 0$ for the target \mathcal{T} . Denote the sample size of each stratum as $n_{s,r} := \sum_{i=1}^n \mathbb{I}(S_i = 1, R_i = r)$ and $n_{\tau,r} := \sum_{i=1}^n \mathbb{I}(S_i = 0, R_i = r)$ for $r \in \{0,1\}$, where $\mathbb{I}(\cdot)$ is the indicator function. Also, we use $n_s := n_{s,0} + n_{s,1}$ and $n_\tau := n_{\tau,0} + n_{\tau,1}$ to represent the sample size on S and T so we have $n = n_s + n_\tau$. We consider an imbalanced sampling scenario on both S and T where $n_{\rm S,1} \gg n_{\rm S,0}$ and $n_{\rm T,1} \gg n_{\rm T,0}$. A common example in biobank studies is that $R=1$ stands for the majority ancestry group European and $R = 0$ for the underrepresented African group.

2.1 Covariate shift

We assume that for $(s, r) \in \{0, 1\}^2$, the samples of (Y, Z) are generated by

$$
Y, \mathbf{Z} \mid S = s, R = r \sim p_{\mathbf{Z}|S=s, R=r}(\mathbf{z}) \cdot p_{Y|\mathbf{Z}, R=r}(y), \tag{3}
$$

where $p_{\mathbf{Z}|S=s,R=r}(\cdot)$ and $p_{Y|\mathbf{Z},R=r}(\cdot)$ represent the distribution function of \mathbf{Z} given $S=s, R=s$ r, and that of Y conditional on **Z** and $R = r$ respectively. Importantly, given $R = r$, the distribution of covariates $\mathbf{Z} = (\mathbf{X}^\text{\tiny T}, \mathbf{W}^\text{\tiny T})^\text{\tiny T}$ is assumed to be different between S and T while the conditional distribution of $Y \mid \mathbf{Z}, R = r$ is assumed to be the same. This is a typical covariate shift scenario frequently considered when Y is not observed on \mathcal{T} .

We introduce the density ratio $h_r^*(\mathbf{z}) := p_{\mathbf{Z}|S=0,R=r}(\mathbf{z})/p_{\mathbf{Z}|S=1,R=r}(\mathbf{z})$ to characterize the covariate shift between the source and target within subgroup $r = 0, 1$. Under such an assumption in [\(3\)](#page-5-1), the conditional mean function $m_r^{\star}(\mathbf{z}) := \mathbb{E}[Y | \mathbf{Z} = \mathbf{z}, R = r]$ remains to be the same between S and T. Nevertheless, we have $\mathbb{E}_{\tau}[Y | \mathbf{X}, R = 0] \neq \mathbb{E}_{\mathcal{S}}[Y | \mathbf{X}, R = 0]$ due to the joint distributional shift of X and the auxiliary W. Consequently, directly regressing Y against **X** on S will typically produce a biased estimator for the target $\bar{\beta}^{[0]}$, no matter the model [\(1\)](#page-5-0) is correct or not. Our first main challenge is to correct this bias when using labeled observations from $\mathcal S$ to estimate the model parameters defined on $\mathcal T$.

Remark 1. In EHR-driven genetic studies such as Section [6,](#page-21-0) \bf{X} includes genetic variants and demographic variables being used to model the risk of some disease Y , while W consists of EHR proxies to Y including relevant diagnostic codes, medication prescriptions, laboratory tests, etc. In this scenario, the focus is on predicting Y using the risk factors in X rather than all covariates in **Z** because the auxiliary **W** does not appear together with **X** at the baseline, on which biomedical discoveries and decisions are supposed to be made. However, the shift between the source and target populations, such as temporal shift in the clinical profile of biobank participants, cannot be fully captured by the difference in X which mainly contains genomic information. It is thus important to include auxiliary variables W for proper population shift adjustment and transfer the knowledge of $Y \mid \mathbf{X}, \mathbf{W}$ from source to target since $Y \mid \mathbf{X}, R = 0$ can be inherently different between the source and target populations.

2.2 Model heterogeneity

Analogous to [\(2\)](#page-5-2), we define the model coefficient on the majority group from $\mathcal{T}, \bar{\beta}^{[1]}$ as the solution to the equations

$$
\mathbb{E}_{\tau}[\mathbf{X}\{Y - g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\} \mid R = 1] = \mathbf{0}.
$$

The larger $n_{s,1}$ and $n_{\tau,1}$ can lead to a more precise estimate of $\bar{\beta}^{[1]}$ compared with $\bar{\beta}^{[0]}$ learned with small underrepresented samples. It is important to mitigate this sampling disparity issue and a tentative strategy is to leverage the estimate of $\bar{\beta}^{[1]}$ as an external knowledge to guide the learning of $\bar{\beta}^{[0]}$. This is motivated by the observation that the genome-wide conditional associations or models of a wide range of diseases and traits tend to have similar patterns across different ancestry groups, as revealed in recent studies [\(Lam et al., 2019;](#page-26-7) [Verma et al.,](#page-28-1) [2023,](#page-28-1) e.g.). Meanwhile, the heterogeneity between $\bar{\beta}^{[1]}$ and $\bar{\beta}^{[0]}$ needs to be considered to avoid potential bias. For these purposes, we introduce the assumption

$$
(\bar{\beta}^{[0]}, \bar{\beta}^{[1]}) \in \left\{ \bar{\beta}^{[0]}, \bar{\beta}^{[1]} : \|\bar{\beta}^{[0]}\|_{0} \leq s_{\beta}^{[0]}, \ \|\bar{\beta}^{[0]} - \bar{\beta}^{[1]}\|_{v} \leq R_{\delta, v} \right\},\tag{4}
$$

where $\|\boldsymbol{\beta}\|_0$ represent the number of non-zero entries of the vector $\boldsymbol{\beta}$ and $\|\boldsymbol{\beta}\|_v$ is the ℓ_v -norm of β for some fixed $v \in [0, 1]$. Here, $\|\cdot\|_v$ corresponds to the exact $(v = 0)$ or approximate

 $(v \in (0, 1])$ sparsity norm, and $s_{\beta}^{[0]}$ $\bar{\beta}^{[0]}$ and $R_{\delta, v}$ represent the sparsity levels of the coefficients $\bar{\boldsymbol{\beta}}^{[0]}$ and the difference between the two subgroups $\bar{\boldsymbol{\beta}}^{[0]} - \bar{\boldsymbol{\beta}}^{[1]}$. Since the outcome models across sub-populations are presumed to be similar for most risk factors, we expect $R_{\delta,v} \ll s_{\beta}^{[0]}$ when taking $v = 0$ as a special case. This promises an efficiency gain by leveraging $\bar{\beta}^{[1]}$ to assist the learning of $\bar{\beta}^{[0]}$ in the underrepresented group, as their difference is sparser and easier to estimate compared to $\bar{\beta}^{[0]}$ itself. On the other hand, given that $s_{\beta}^{[0]}$ $\beta^{\text{[0]}}$ and $R_{\delta,v}$ are unknown in practice, we aim to maintain adaptivity and robustness to excessive model heterogeneity, i.e., the case where $R_{\delta,v}$ is large compared to $s_{\beta}^{[0]}$ $\bar{\boldsymbol{\beta}}^{[0]}$. In this scenario, the knowledge from $\bar{\boldsymbol{\beta}}^{[1]}$ can be non-informative or misleading to the target parameters $\bar{\beta}^{[0]}$, and it is desirable to avoid potential negative transfer caused by this. The problem setup described above is illustrated in Figure [1.](#page-7-0) B , and B and B and B and B and B caused by this. The problem setup description

Figure 1: The transfer learning setup considered by MAKEUP with our primary goal being risk modeling for the minority subgroup from some target cohort without labeled samples (marked in the red block).

3 Method

3.1 Model-assisted covariate shift correction

Define the empirical mean operators on the source samples, target samples, and their union for group $r \in \{0,1\}$ as: $\widehat{\mathbb{E}}_{\mathcal{S}}^{[r]} f(\mathbf{D}) = n_{\mathcal{S},r}^{-1} \sum_{i=1}^{n} \mathbb{I}(S_i = 1, R_i = r) f(\mathbf{D}_i)$, $\widehat{\mathbb{E}}_{\mathcal{T}}^{[r]} f(\mathbf{D}) =$ $n_{\tau,r}^{-1} \sum_{i=1}^{n} \mathbb{I}(S_i = 0, R_i = r) f(\mathbf{D}_i)$, and $\widehat{\mathbb{E}}_{S \cup \tau}^{[r]} f(\mathbf{D}) = (n_{S,r} + n_{\tau,r})^{-1} \sum_{i=1}^{n} \mathbb{I}(R_i = r) f(\mathbf{D}_i)$.

We begin with specifying the nuisance models for density ratio and conditional mean imputation as follows:

$$
h_r(\mathbf{z}) = \exp\left(\phi(\mathbf{z})^{\mathsf{T}} \boldsymbol{\alpha}^{[r]}\right), \text{ and } m_r(\mathbf{z}) = b(\phi(\mathbf{z})^{\mathsf{T}} \boldsymbol{\gamma}^{[r]}),
$$

where $\phi : \mathbb{R}^{p+q} \to \mathbb{R}^d$ is a basis function of **Z**, which can be chosen flexibly. For example,

one could simply take $\phi(\mathbf{z}) = \mathbf{z}$ or set $\phi(\mathbf{z})$ as a concatenation of non-linear bases of the components of **z**. The vectors $\alpha^{[r]}$ and $\gamma^{[r]}$ are the coefficients for the nuisance models that need to be estimated. The function $b(\cdot)$ is a pre-specified, monotonically increasing link function, which may or may not be the same as the link function $q(\cdot)$. Denote by $\mathbf{\Phi} = \phi(\mathbf{Z})$. When the imputation model $m_r(\mathbf{z}) = m_r^{\star}(\mathbf{z})$, the target parameter $\bar{\beta}^{[r]}$ is the solution to the following imputation-based (IM) estimating equation:

$$
\mathbb{E}_{\tau}[\mathbf{X}\{m_r(\mathbf{Z}) - g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\} \mid R = r] = \mathbf{0}.
$$

Also, when $h_r(\mathbf{z}) = h_r^*(\mathbf{z}), \bar{\boldsymbol{\beta}}^{[r]}$ solves the importance weighted (IW) estimating equation:

$$
\mathbb{E}_{S}[h_r(\mathbf{Z})\mathbf{X}\{Y-g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\}\mid R=r]=\mathbf{0}.
$$

Empirically, $\bar{\beta}^{[r]}$ can be estimated using the sample versions of these equations. However, these simple strategies are prone to potential model misspecification. Specifically, the IW method leads to inconsistency if the density ratio model h_r is misspecified, and the IM method fails when the imputation model m_r is misspecified.

To overcome this challenge, we combine the two nuisance models to construct a doubly robust estimating equation for $\bar{\beta}^{[r]}$ as

$$
\mathbb{E}_{\mathcal{S}}[h_r(\mathbf{Z})\mathbf{X}\{m_r(\mathbf{Z}) - Y\} \mid R = r] - \mathbb{E}_{\mathcal{T}}[\mathbf{X}\{m_r(\mathbf{Z}) - g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\} \mid R = r] = \mathbf{0}.
$$
 (5)

We show in Proposition [1](#page-9-0) that [\(5\)](#page-8-0) provides a consistent solution for $\bar{\beta}^{[r]}$ when either the density ratio or the imputation model is correctly specified. Motivated by this, we propose an empirical doubly robust loss function in [\(6\)](#page-8-1). Specifically, given the two nuisance parameters $\alpha^{[r]}$ and $\gamma^{[r]}$ whose estimation will be discussed later, we define the doubly robust covariate shift-corrected sparse regression for $\bar{\beta}^{[r]}$ as:

$$
\widehat{\boldsymbol{\beta}}^{[r]} = \arg\min_{\boldsymbol{\beta}\in\mathbb{R}^q} \mathcal{L}_r(\boldsymbol{\beta};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]}) + \lambda^{[r]}\|\boldsymbol{\beta}\|_1,\tag{6}
$$

where the loss function is defined as

$$
\mathcal{L}_r(\boldsymbol{\beta};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]}) = \left[\widehat{\mathbb{E}}_S^{[r]} \mathbf{X}^{\mathsf{T}} \exp(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\alpha}^{[r]}) \{b(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma}^{[r]}) - Y\} - \widehat{\mathbb{E}}_T^{[r]} \mathbf{X}^{\mathsf{T}} b(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma}^{[r]})\right] \boldsymbol{\beta} + \widehat{\mathbb{E}}_T^{[r]} G(\mathbf{X}^{\mathsf{T}} \boldsymbol{\beta}),
$$

where $G(a) = \int_0^a g(u) \mathrm{d}u$, and $\lambda^{[r]}$ is a penalization parameter. For all tuning parameters including $\lambda^{[r]}$ and others to be introduced later, we will present their theoretically optimal rate in Section [4](#page-13-0) and their empirical tuning strategies in Supplementary B.3. By our construction, $\mathbb{E}\partial_{\beta}\mathcal{L}_r(\beta;\alpha^{[r]},\gamma^{[r]})$ is equivalent to the left-hand side of [\(5\)](#page-8-0). Thus, by Proposition [1,](#page-9-0) $\mathcal{L}_r(\beta;\alpha^{[r]},\gamma^{[r]})$ is doubly robust to the misspecification of the two nuisance models in the sense that either a correct density ratio or a correct conditional mean imputation model can lead to the target solution $\bar{\boldsymbol{\beta}}^{[r]}$.

Proposition 1. Let $\boldsymbol{\beta}^{[r]}$ be the solution of the estimating equation $\mathbb{E}\partial_{\boldsymbol{\beta}}\mathcal{L}_r(\boldsymbol{\beta};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]})=0$, which is equivalent with equation [\(5\)](#page-8-0). We have $\boldsymbol{\beta}^{[r]} = \bar{\boldsymbol{\beta}}^{[r]}$ when either $h_r^{\star}(\mathbf{z}) = \exp \left(\phi(\mathbf{z})^{\mathsf{T}} \boldsymbol{\alpha}^{[r]} \right)$ or $m_r^{\star}(\mathbf{z}) = b(\phi(\mathbf{z})^{\tau} \boldsymbol{\gamma}^{[r]})$.

Motivated by the above-introduced construction, a natural strategy for the estimation of $\bar{\boldsymbol{\beta}}^{[r]}$ is to first estimate $\boldsymbol{\alpha}^{[r]}$ and $\boldsymbol{\gamma}^{[r]}$ through regularized regression

$$
\widetilde{\boldsymbol{\alpha}}^{[r]} = \underset{\boldsymbol{\alpha} \in \mathbb{R}^{q+p}}{\arg \min} \widehat{\mathbb{E}}_{\mathcal{S} \cup \mathcal{T}}^{[r]} \{ \rho_{\mathcal{T},r} S \exp(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\alpha}) - \rho_{\mathcal{S},r} (1-S) \boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\alpha} \} + \lambda_{\alpha}^{[r]} ||\boldsymbol{\alpha}||_{1};
$$
\n
$$
\widetilde{\boldsymbol{\gamma}}^{[r]} = \underset{\boldsymbol{\gamma} \in \mathbb{R}^{q+p}}{\arg \min} \widehat{\mathbb{E}}_{\mathcal{S}}^{[r]} \{ -Y \boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma} + B(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma}) \} + \lambda_{\gamma}^{[r]} ||\boldsymbol{\gamma}||_{1},
$$
\n(7)

where $\rho_{\tau,r} = (n_{\mathcal{S},r} + n_{\tau,r})/n_{\tau,r}, \ \rho_{\mathcal{S},r} = (n_{\mathcal{S},r} + n_{\tau,r})/n_{\mathcal{S},r}, \ B(a) = \int_0^a b(u) \mathrm{d}u$, and $\lambda_\alpha^{[r]}$, $\lambda_\gamma^{[r]}$ are two tuning parameters, then obtain the preliminary estimator $\widetilde{\beta}^{[r]}$ as

$$
\widetilde{\boldsymbol{\beta}}^{[r]} = \underset{\boldsymbol{\beta} \in \mathbb{R}^q}{\arg \min} \mathcal{L}_r(\boldsymbol{\beta}, \widetilde{\boldsymbol{\alpha}}^{[r]}, \widetilde{\boldsymbol{\gamma}}^{[r]}) + \lambda^{[r]} \|\boldsymbol{\beta}\|_1.
$$
\n(8)

According to our above discussion, $\tilde{\boldsymbol{\beta}}^{[r]}$ tends to be consistent when either the density ratio or the conditional mean model is correctly specified. However, it is unlikely to achieve a desirable convergence rate due to the excessive biases in $\tilde{\alpha}^{[r]}$ and $\tilde{\gamma}^{[r]}$ introduced by regularization; see Remark [5](#page-15-0) for more discussion. To address this issue, we propose a bifold bias correction approach. Denote by $\bar{\mathbf{\Sigma}}_{\boldsymbol{\beta}}^{[r]} = \mathbb{E}_{\tau}[g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\mathbf{X}\mathbf{X}^{\mathsf{T}} \mid R = r]$ and $\widehat{\mathbf{\Sigma}}_{\boldsymbol{\beta}}^{[r]} = \widehat{\mathbb{E}}_{\tau}^{[r]}[g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\mathbf{X}\mathbf{X}^{\mathsf{T}}]$ where $\dot{g}(\cdot)$ is the derivative of $g(\cdot)$. As the first fold of bias correction, we address the bias caused by the regularization term in [\(8\)](#page-9-1) through an one-step debiased construction for each $\beta_j^{[r]} = \mathbf{e}_j^{\intercal}\boldsymbol{\beta}^{[r]} \,\, (j \in \{1,\ldots,q\})$:

$$
\widehat{\beta}_{\mathrm{Deb},j}^{[r]}(\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]}) = \widetilde{\beta}_j^{[r]} + \widehat{\boldsymbol{\Omega}}_j^{[r]} \partial_{\boldsymbol{\beta}} \mathcal{L}_r(\widetilde{\boldsymbol{\beta}}^{[r]};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]}),
$$
\n(9)

where \mathbf{e}_j is the j-th unit vector in \mathbb{R}^q , $\widetilde{\beta}_j^{[r]}$ is the j-th element of $\widetilde{\beta}^{[r]}$, $\widehat{\Omega}_j^{[r]}$ j is a regularized estimation of $\bar{\mathbf{\Omega}}_i^{[r]}$ $[j^r]$, the *j*-th row of $\bar{\mathbf{\Omega}}^{[r]} = \left[\bar{\mathbf{\Sigma}}_{\bar{\mathsf{A}}^{[r]}}^{[r]}\right]$ $[\bar{\rho}^{[r]}_b]^{-1}$, and ∂_{β} is the partial derivative operator with respect to $\boldsymbol{\beta}$. To construct $\widehat{\boldsymbol{\Omega}}_{j}^{[r]}$ j^j , we use the node-wise lasso method proposed by [Van de](#page-28-6) [Geer et al. \(2014\)](#page-28-6). See Supplementary B.1 for the detailed expression of $\widehat{\Omega}_{j}^{[r]}$.,
j .

This step only reduces the biases arising from the regularization on β in [\(8\)](#page-9-1). In the second fold of bias correction, we further mitigate the biases due to the excessive regularization errors in $\tilde{\alpha}^{[r]}$ and $\tilde{\gamma}^{[r]}$. Our key idea is to further calibrate the two nuisance models and make them satisfying the moment conditions:

$$
\mathbb{E}\big[\partial_{\alpha}\partial_{\beta}\mathcal{L}_r(\bar{\boldsymbol{\beta}}^{[r]};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]})\big] = \mathbf{0}; \quad \mathbb{E}\big[\partial_{\gamma}\partial_{\beta}\mathcal{L}_r(\bar{\boldsymbol{\beta}}^{[r]};\boldsymbol{\alpha}^{[r]},\boldsymbol{\gamma}^{[r]})\big] = \mathbf{0}.\tag{10}
$$

Under these two conditions, the first-order errors in $\alpha^{[r]}$ and $\gamma^{[r]}$ can be removed through concentration in a similar spirit with the idea of Neyman orthogonality [\(Chernozhukov et al.,](#page-24-6) [2018\)](#page-24-6). This motivates us to obtain the calibrated nuisance estimators as $\hat{\alpha}_j^{[r]} = \tilde{\alpha}^{[r]} + \hat{\xi}_j^{[r]}$ j and $\widehat{\boldsymbol{\gamma}}_j^{[r]} = \widetilde{\boldsymbol{\gamma}}^{[r]} + \widehat{\boldsymbol{\zeta}}_j^{[r]}$ with

$$
\widehat{\boldsymbol{\xi}}_{j}^{[r]} = \underset{\boldsymbol{\xi} \in \mathbb{R}^{q+p}}{\operatorname{argmin}} \widehat{\mathbb{E}}_{\mathcal{S} \cup \mathcal{T}}^{[r]} \widehat{w}_{j}^{[r]} \dot{b} (\boldsymbol{\Phi}^{\mathsf{T}} \widetilde{\boldsymbol{\gamma}}^{[r]}) \mathcal{F}^{[r]}(\boldsymbol{\xi}; \widetilde{\boldsymbol{\alpha}}^{[r]}) + \lambda_{\alpha_{j}}^{[r]} \|\boldsymbol{\xi}\|_{1} ;
$$
\n
$$
\widehat{\boldsymbol{\zeta}}_{j}^{[r]} = \underset{\boldsymbol{\zeta} \in \mathbb{R}^{q+p}}{\operatorname{argmin}} \widehat{\mathbb{E}}_{\mathcal{S}}^{[r]} \widehat{w}_{j}^{[r]} \exp(\boldsymbol{\Phi}^{\mathsf{T}} \widetilde{\boldsymbol{\alpha}}^{[r]}) \mathcal{G}(\boldsymbol{\zeta}; \widetilde{\boldsymbol{\gamma}}^{[r]}) + \lambda_{\gamma_{j}}^{[r]} \|\boldsymbol{\zeta}\|_{1} ,
$$
\n(11)

where $\widehat{w}_j^{[r]} = \widehat{\Omega}_j^{[r]} \mathbf{X}$ and

$$
\mathcal{F}^{[r]}(\boldsymbol{\xi};\boldsymbol{\alpha}) = \rho_{\tau,r} S \exp{\{\boldsymbol{\Phi}^{\tau}(\boldsymbol{\alpha}+\boldsymbol{\xi})\}} - \rho_{s,r}(1-S)\boldsymbol{\Phi}^{\tau}(\boldsymbol{\alpha}+\boldsymbol{\xi}) ;
$$

$$
\mathcal{G}(\boldsymbol{\zeta};\boldsymbol{\gamma}) = -Y\boldsymbol{\Phi}^{\tau}(\boldsymbol{\gamma}+\boldsymbol{\zeta}) + B\{\boldsymbol{\Phi}^{\tau}(\boldsymbol{\gamma}+\boldsymbol{\zeta})\} .
$$

The LASSO problems in [\(11\)](#page-10-0) are designed such that their Karush–Kuh–Tucker (gradient) conditions empirically match our desirable moment conditions in [\(10\)](#page-9-2). We then plug the calibrated $\hat{\boldsymbol{\alpha}}_{j}^{[r]}$ $\begin{bmatrix} [r]_j \end{bmatrix}$ and $\begin{bmatrix} \widehat{\boldsymbol{\gamma}}_j^{[r]} \end{bmatrix}$ $j^{[r]}$ into [\(9\)](#page-9-3) for a bifold bias-corrected estimator of each $\beta_j^{[r]}$ $j^{[r]}$, denoted $\text{as}\ \widehat{\beta}^{[r]}_{\text{Deb},j} \ = \ \widehat{\beta}^{[r]}_{\text{Deb},j} (\widehat{\boldsymbol{\alpha}}^{[r]}_{j})$ $\begin{bmatrix} [r]\,,\,\widehat{\boldsymbol{\gamma}}_j^{[r]} \end{bmatrix}$ ^[r]). We also denote by $\widehat{\beta}_{\text{Deb}}^{[r]} = (\widehat{\beta}_{\text{Deb},1}^{[r]}, \ldots, \widehat{\beta}_{\text{Deb},q}^{[r]})^{\mathsf{T}}$. The aboveintroduced covariate shift correction approach is summarized in Algorithm [1.](#page-10-1)

Remark 2. Note that the weight $\hat{w}_j^{[r]}$ used in [\(11\)](#page-10-0) may not be positive definite, which could make the loss function in [\(11\)](#page-10-0) irregular and ill-posed. To handle this, one can divide the samples into two sets with positive and negative only \hat{w}_{ij} 's respectively and solve [\(11\)](#page-10-0) on them separately. Details of this stratification strategy are presented in Supplementary B.2.

Although the bias-corrected $\widehat{\beta}_{\text{Deb}}^{[r]}$ is element-wise consistent to $\bar{\beta}^{[r]}$ with a desirable convergence rate, its overall ℓ_2 -error $\|\widehat{\beta}_{\text{Deb}}^{[r]} - \bar{\beta}^{[r]}\|_2$ does not converge to zero since $\widehat{\beta}_{\text{Deb}}^{[r]}$ is a dense estimator in contrast to the sparse $\bar{\beta}^{[r]}$, with its ℓ_2 -error growing fast with the large dimension q. For the purpose of risk prediction, we further construct an ℓ_2 -consistent estimator for $\bar{\beta}^{[r]}$ through a thresholding and sparsifying approach:

$$
\widehat{\boldsymbol{\beta}}_{\text{Thr}}^{[r]} = \left\{ \text{Thr} \big(\widehat{\beta}_{\text{Deb},j}^{[r]}, \tau^{[r]} \big) \right\}_{j=1,\dots,q},\tag{12}
$$

where $\text{Three}(z, c) = z$ if $|z| \geq c$; $\text{Three}(z, c) = 0$ if $|z| < c$, and $\tau^{[r]}$ is a tuning parameter. In Theorem [1,](#page-15-1) we justify the ℓ_2 -consistency of $\widehat{\beta}_{\text{Thr}}^{[r]}$ and derive its rate of convergence. Related to our above discussion, we also comment in Remarks [4](#page-15-2) and [5](#page-15-0) that $\hat{\beta}_{\text{Thr}}^{[r]}$ is substantially more robust to regularization errors in the nuisance estimators compared with the preliminary estimator $\widetilde{\boldsymbol{\beta}}^{[r]}$.

3.2 Adaptive knowledge transfer from majority to minority

The construction of $\hat{\beta}_{\text{Thr}}^{[0]}$ enables us to borrow information from the source minority population, which may still suffer from high variability due to small sample size. Thus, we propose a knowledge-guided transfer learning procedure using the majority group estimator $\hat{\beta}_{\text{Thr}}^{[1]}$ to aid the estimation of $\bar{\beta}^{[0]}$. Similar to recent work like [Li et al. \(2022\)](#page-26-3), our approach relies on the presumption that $\bar{\pmb{\delta}} = \bar{\pmb{\beta}}^{[1]} - \bar{\pmb{\beta}}^{[0]}$ is sparser and easier to estimate well than $\bar{\pmb{\beta}}^{[0]}$, e.g., $R_{\delta,v} \ll s_{\beta}^{[0]}$ when $v = 0$ in [\(4\)](#page-6-0). The idea is to take $\widehat{\boldsymbol{\beta}}_{\text{Thr}}^{[1]}$ as a baseline (offset) and estimate the difference term $\bar{\beta}^{[0]} - \hat{\beta}^{[1]}_{\text{Thr}}$ instead of $\bar{\beta}^{[0]}$ through thresholding and sparsifying. The details are described in Algorithm [2.](#page-11-0)

Algorithm 2 Majority-knowledge-guided thresholding estimation

Input: The debiased coefficient vector $\hat{\beta}_{\text{Deb}}^{[0]}$ obtained by implementing Algorithm [1](#page-10-1) on the minority group, and the sparsified estimator $\hat{\beta}_{\text{Thr}}^{[1]}$ obtained on the majority group using (12) . **Output:** Knowledge transfer estimator $\hat{\boldsymbol{\beta}}_{\text{KTr}}^{[0]} = \hat{\boldsymbol{\beta}}_{\text{Thr}}^{[1]} + \hat{\boldsymbol{\delta}}$ where $\hat{\boldsymbol{\delta}} = (\hat{\delta}_1, \dots, \hat{\delta}_q)^{\text{T}}, \ \hat{\delta}_j = (\hat{\delta}_1, \dots, \hat{\delta}_q)^{\text{T}}$ Thre $(\hat{\beta}_{\text{Deb},j}^{[0]} - \hat{\beta}_{\text{Thr},j}^{[1]}, \tau_{\text{KTr}})$ and τ_{KTr} is a tuning parameter.

As shown in Section [4.3,](#page-16-0) when $\bar{\delta}$ is sparser than $\bar{\beta}^{[0]}$, Algorithm [2](#page-11-0) can effectively leverage knowledge from the majority group to improve the estimation efficiency on $\bar{\beta}^{[0]}$. However, when $\bar{\boldsymbol{\delta}}$ is denser than $\bar{\boldsymbol{\beta}}^{[0]}$, i.e., $\bar{\boldsymbol{\beta}}^{[1]}$ differs substantially from $\bar{\boldsymbol{\beta}}^{[0]}$, $\hat{\boldsymbol{\beta}}^{[0]}_{\text{KTr}}$ given by Algo-rithm [2](#page-11-0) could be less efficient than the minority-only estimator $\hat{\beta}_{\text{Thr}}^{[0]}$, suffering from negative knowledge transfer. To address this issue, we propose a novel negative transfer protection approach that ensures the final estimator for $\bar{\beta}^{[0]}$ to be no worse than $\hat{\beta}_{\text{Thr}}^{[0]}$ and is adaptive to the unknown transferability level between the two sub-populations (i.e., sparsity level of δ). For similar purposes, recent work like [Tian and Feng \(2023\)](#page-27-6) uses hold-out samples from the minority group to select between or ensemble the minority-only and knowledge transfer estimators. However, such a strategy is not directly applicable to our case due to the absence of labeled data on the target site $\mathcal T$. To handle this challenge, we introduce Algorithm [3](#page-12-0) that relies on a surrogate loss

$$
\widehat{\mathcal{Q}}(\boldsymbol{\beta}^{[0]};\widehat{\boldsymbol{\beta}}_{\mathrm{Deb}}^{[0]}):=\big\|\boldsymbol{\beta}^{[0]}-\widehat{\boldsymbol{\beta}}_{\mathrm{Deb}}^{[0]}\big\|_2^2,
$$

to evaluate the accuracy of $\boldsymbol{\beta}^{[0]}$. In this formulation, the dense vector $\widehat{\boldsymbol{\beta}}_{\text{Deb}}^{[0]}$ can be essentially viewed as a Gaussian vector with mean $\bar{\beta}^{[0]}$ (neglecting higher-order error terms) and, thus, serving as an appropriate criterion for model selection. We demonstrate this point more rigorously in Lemma [3.](#page-17-0)

In Algorithm [3,](#page-12-0) we adopt data-splitting on the minority samples and derive two independent bias-corrected dense vectors $\widehat{\boldsymbol{\beta}}_{\mathrm{D}\epsilon}^{[0]}$ $_{\rm Deb,1}^{[0]}$ and $\widehat{\boldsymbol{\beta}}^{[0]}_{\rm D6}$ ^[0]_{Deb,2}. One of them, says $\widehat{\boldsymbol{\beta}}_{\mathrm{D6}}^{[0]}$ $_{\text{Deb},1}^{\text{D}}$, is used to derive the sparse minority-only estimator $\widehat{\boldsymbol{\beta}}_{\text{Th}}^{[0]}$ $\widehat{\mathsf{f}}_{\rm{Tr},1}^{[0]}$ and the knowledge transfer estimator $\widehat{\boldsymbol{\beta}}_{\rm{K}^{\prime}}^{[0]}$ KTr,1 . Then, the other one $\widehat{\beta}_{\mathrm{De}}^{[0]}$ $\mathbb{D}_{\text{eb},2}$ is used to ensemble these two estimators through the surrogate loss function $\widehat{\mathcal{Q}}(\cdot; \widehat{\boldsymbol{\beta}}_{\mathrm{D}\epsilon}^{[0]}$ $\mathcal{D}_{\text{e}b,2}$). We use cross-fitting to leverage the two folds more effectively. For ensembling, we use the exponential weighting strategy [\(Dalalyan and Tsybakov, 2007,](#page-25-7) e.g.) with some pre-specified temperature parameter $a > 0$ in [\(13\)](#page-12-1). When $a = \infty$, $\hat{\beta}_{MU,k}^{[0]}$ will be the one chosen from $\widehat{\beta}_{\text{Thr},k}^{[0]}$ and $\widehat{\beta}_{\text{KTr},k}^{[0]}$ that has smaller loss $\widehat{\mathcal{Q}}$ contrasted with the debiased vector from the other fold of data. As will be shown in Theorem [3,](#page-17-1) the resulted estimator named as MAKEUP is ensured to be no worse than both $\hat{\beta}_{\text{Thr},k}^{[0]}$ and $\hat{\beta}_{\text{KTr},k}^{[0]}$, with a slight price to pay for the model ensemble or selection in Algorithm [3.](#page-12-0)

Algorithm 3 Protect against negative knowledge transfer

 $\textbf{Input:}~~ \mathscr{D}^{[r]} = \{ \mathbf{D}_i = (S_i Y_i, \mathbf{X}_i^{\scriptscriptstyle \sf T}, \mathbf{W}_i^{\scriptscriptstyle \sf T}, S_i, R_i)^{\scriptscriptstyle \sf T} : R_i = r, \,\, i \in [n] \}, r = 0, 1;$

1: Randomly split the minority data $\mathscr{D}^{[0]}$ (including both the source and target samples) into two disjoint sets $\mathscr{D}_1^{[0]}$ $\mathcal{D}_1^{[0]}$ and $\mathscr{D}_2^{[0]}$ $2^{[0]}$ of equal size.

- 2: Implement Algorithm [1](#page-10-1) on $\mathscr{D}_k^{[0]}$ ^[0] to derive $\widehat{\boldsymbol{\beta}}_{\text{Deb},k}^{[0]}$ for $k = 1, 2$ and on $\mathscr{D}^{[1]}$ for $\widehat{\boldsymbol{\beta}}_{\text{Deb}}^{[1]}$.
- **3:** Implement thresholding on the dense vectors to obtain $\hat{\beta}_{\text{Thr},k}^{[0]}$ for $k = 1, 2$ and $\hat{\beta}_{\text{Thr}}^{[1]}$.
- **4:** Implement Algorithm [2](#page-11-0) with $\widehat{\boldsymbol{\beta}}_{\text{Thr}}^{[1]}$ and $\widehat{\boldsymbol{\beta}}_{\text{Deb},k}^{[0]}$ to obtain $\widehat{\boldsymbol{\beta}}_{\text{KTr},k}^{[0]}$ for $k = 1, 2$.
-
- 5: For $k = 1, 2$, aggregate the estimators as

$$
\widehat{\boldsymbol{\beta}}_{\mathrm{MU},k}^{[0]} = w_k \widehat{\boldsymbol{\beta}}_{\mathrm{Thr},k}^{[0]} + (1-w_k) \widehat{\boldsymbol{\beta}}_{\mathrm{KTr},k}^{[0]}, \ w_k = \frac{e^{-a\widehat{\mathcal{Q}}(\widehat{\boldsymbol{\beta}}_{\mathrm{Thr},k}^{[0]},\widehat{\boldsymbol{\beta}}_{\mathrm{Deb},3-k}^{[0]})}}{e^{-a\widehat{\mathcal{Q}}(\widehat{\boldsymbol{\beta}}_{\mathrm{Thr},k}^{[0]},\widehat{\boldsymbol{\beta}}_{\mathrm{Deb},3-k}^{[0]})} + e^{-a\widehat{\mathcal{Q}}(\widehat{\boldsymbol{\beta}}_{\mathrm{KT},k}^{[0]},\widehat{\boldsymbol{\beta}}_{\mathrm{Deb},3-k}^{[0]})}}.
$$
(13)

Output: The final MAKEUP estimator $\widehat{\boldsymbol{\beta}}_{MU}^{[0]} = (\widehat{\boldsymbol{\beta}}_{MU,1}^{[0]} + \widehat{\boldsymbol{\beta}}_{MU}^{[0]}$ $_{\rm MU,2}^{^{[0]}}\big)/2.$

4 Theoretical Justification

4.1 Notations

To introduce the key sparsity assumptions and the convergence results of the nuisance estimators, we first define the population-level model parameters as follows. Let

$$
\bar{\boldsymbol{\alpha}}^{[r]} = \argmin_{\boldsymbol{\alpha} \in \mathbb{R}^{q+p}} \mathbb{E}_{\mathcal{S} \cup \mathcal{T}}^{[r]} \{ \rho_{\mathcal{T},r} S \exp(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\alpha}) - \rho_{\mathcal{S},r} (1-S) \boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\alpha} \},
$$
\n
$$
\bar{\boldsymbol{\gamma}}^{[r]} = \argmin_{\boldsymbol{\gamma} \in \mathbb{R}^{q+p}} \mathbb{E}_{\mathcal{S}}^{[r]} \{ -Y \boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma} + B(\boldsymbol{\Phi}^{\mathsf{T}} \boldsymbol{\gamma}) \},
$$

be the population-level preliminary nuisance model coefficients. For $j = 1, \ldots, q$, we let $\bar{\boldsymbol{\alpha}}_{j}^{[r]}=\bar{\boldsymbol{\alpha}}^{[r]}+\bar{\boldsymbol{\xi}}_{j}^{[r]}$ $[\bar{y}]$ and $\bar{\gamma}_j^{[r]} = \bar{\gamma}^{[r]} + \bar{\zeta}_j^{[r]}$ be the population-level calibrated coefficients corresponding to equation (11) , where

$$
\bar{\xi}_{j}^{[r]} = \underset{\xi \in \mathbb{R}^{q+p}}{\operatorname{argmin}} \mathbb{E}_{s \cup \tau}^{[r]} \bar{w}_{j}^{[r]} \dot{b}(\Phi^{\tau} \bar{\gamma}^{[r]}) \mathcal{F}^{[r]}(\xi; \bar{\alpha}^{[r]}),
$$
\n
$$
\bar{\zeta}_{j}^{[r]} = \underset{\xi \in \mathbb{R}^{q+p}}{\operatorname{argmin}} \mathbb{E}_{s}^{[r]} \bar{w}_{j}^{[r]} \exp(\Phi^{\tau} \bar{\alpha}^{[r]}) \mathcal{G}(\zeta; \bar{\gamma}^{[r]}),
$$
\n(14)

and $\bar{w}_j^{[r]} = \bar{\mathbf{\Omega}}_j^{[r]} \mathbf{X}$. When the density ratio model is correctly specified, we can show that $\bar{\boldsymbol{\xi}}_{j}^{[r]}=\boldsymbol{0} \text{ and } \bar{\boldsymbol{\alpha}}^{[r]}=\bar{\boldsymbol{\alpha}}_{j}^{[r]}$ ^[r] for all $j \in \{1, \ldots, p\}$. Similarly, when the imputation model is correct, we have $\bar{\boldsymbol{\zeta}}_j^{[r]} = \mathbf{0}$ and $\bar{\boldsymbol{\gamma}}_i^{[r]} = \bar{\boldsymbol{\gamma}}_j^{[r]}$ $j^{[r]}$. For $r \in \{0,1\}$, we introduce

$$
s_{\beta}^{[r]} = ||\bar{\boldsymbol{\beta}}^{[r]}||_0 \text{ and } s_{\text{nu}}^{[r]} = \max \left\{ ||\bar{\boldsymbol{\alpha}}^{[r]}||_0, ||\bar{\boldsymbol{\gamma}}^{[r]}||_0, \max_{j \in \{1, \dots, p\}} ||\bar{\boldsymbol{\Omega}}_j^{[r]}||_0 \right\}
$$

to respectively denote the (exact) sparsity level of the target model coefficients and the nuisance model coefficients. Considering that nuisance parameters $\bar{\xi}_i^{[r]}$ $_{j}^{[r]}$ and $\bar{\zeta}_{j}^{[r]}$ $j^{\left(r\right)}$ are defined through the reweighted regression in [\(14\)](#page-13-1), we introduce $R_{\text{nu},v}^{[r]} = \max \left\{ {\|\bar{\boldsymbol{\xi}}_j^{[r]}\|_v, \|\bar{\boldsymbol{\zeta}}_j^{[r]}\|_v \right\}$ for an arbitrary $v \in [0, 1]$ to accommodate both the exact (i.e., $v = 0$) and approximate sparsity (i.e., $v \in (0,1]$) regimes. In addition, we define $R_{\delta,v} = \|\bar{\beta}^{[1]} - \bar{\beta}^{[0]}\|_v = \|\bar{\delta}\|_v$ for any $v \in [0,1]$ to depict the model heterogeneity between the majority and minority sub-populations on the target. Again, our analyses cover both the exact and approximate sparse $\bar{\delta}$ scenarios.

We use $a_n = o(b_n)$ if $\lim_{n\to\infty} a_n/b_n = 0$, $a_n = O(b_n)$ if $\lim_{n\to\infty} |a_n/b_n| \leq C$ for some constant C, and $a_n \n\t\le b_n$ if $a_n = O(b_n)$ and $b_n = O(a_n)$. We use $\stackrel{\mathbb{P}}{\rightarrow}$ to denote convergence in probability and $\stackrel{\mathcal{D}}{\rightarrow}$ for convergence in distribution. For a sequence random variables Z_n , we use $Z_n = o_p(a_n)$ if $|Z_n|/a_n \stackrel{\mathbb{P}}{\rightarrow} 0$ and $Z_n = O_p(a_n)$ if $\lim_{C \to \infty} \lim_{n \to \infty} \mathbb{P}(|Z_n/a_n| > C) = 0$. Denote by $n_r = n_{s,r} \wedge n_{\tau,r}$ for $r \in \{0,1\}$, where $x \wedge y = \min\{x, y\}$, and assume $n_{\mathcal{S},0} = o(n_{\mathcal{S},1})$ and $n_{\mathcal{T},0} = o(n_{\mathcal{T},1})$, i.e., the minority sample sizes are much smaller than the majority on both source and target.

4.2 Covariate shift correction

We first justify the robustness and effectiveness of the model-assisted covariate shift correc-tion step introduced in Algorithm [1](#page-10-1) as well as its subsequent thresholding estimator $\hat{\beta}_{\text{Thr}}^{[r]}$. First, we introduce the key Assumption [1](#page-14-0) that at least one nuisance model is correctly specified, referred to as the model-doubly-robust assumption in semiparametric literature [\(Smucler et al., 2019\)](#page-27-9).

Assumption 1. For $r \in \{0, 1\}$, either $h_r^{\star}(\mathbf{z}) := p_{\mathbf{Z}|S=0, R=r}(\mathbf{z})/p_{\mathbf{Z}|S=1, R=r}(\mathbf{z}) = \exp(\phi(\mathbf{z})^{\intercal} \bar{\mathbf{\alpha}}^{[r]})$ or $m_r^{\star}(\mathbf{z}) \coloneqq \mathbb{E}[Y \mid \mathbf{Z} = \mathbf{z}, R = r] = b(\phi(\mathbf{z})^{\top} \bar{\pmb{\gamma}}^{[r]})$ holds.

All of our technical assumptions presented in Supplementary A.1 have been commonly used in existing literature of high-dimensional inference [\(Van de Geer et al., 2014,](#page-28-6) e.g.) and doubly robust inference [\(Tan, 2020,](#page-27-8) e.g.). As a summary, Assumption A1 rules out the heavy tail and singularity of the predictors X as well as the nuisance model covariates Φ . In this assumption, we introduce $K > 0$ such that $||\mathbf{X}_i||_{\infty} \leq K$ for all $i \in \{1, ..., n_{\mathcal{S}} + n_{\mathcal{T}}\}$ with probability approaching 1, where $||(x_1,\ldots,x_d)||_{\infty} := \max_{j\in\{1,\ldots,d\}} |x_j|$. For bounded design, we have $K = O(1)$ and for gaussian or sub-gaussian design, we can show $K = O({\log(q)}^{1/2})$.

Assumption A2 includes sparsity assumptions that $s_{\beta}^{[r]}$ $\left[r \right]_{{\beta}}, \, s_{\text{nu}}^{[r]} , \, R_{\text{nu},v}^{[r]}$ and $R_{\delta, v}$ are not excessively large given the sample sizes, which are crucial for the consistency of the nuisance and target model estimators. Similar requirements can be found in existing literature for sparse GLM [\(Wainwright, 2019\)](#page-28-7). Assumption A3 is a regularity condition on the link functions $q(\cdot)$ and $b(\cdot)$ satisfied in broad GLM settings. Assumption A4 includes the optimal rate of all tuning parameters. We first introduce Lemma [1](#page-14-1) about the convergence property of each bias-corrected $\widehat{\beta}_{\text{Deb},j}^{[r]}$ obtained in Algorithm [1.](#page-10-1) This key lemma is justified based on the convergence results of $\widehat{\boldsymbol{\Omega}}_{j}^{[r]}$ $\left[\begin{matrix}[r]\end{matrix}\right], \widetilde{\boldsymbol{\beta}}^{[r]},$ and $\{\widehat{\boldsymbol{\alpha}}_j^{[r]}\}$ $\left[r\right] _{j},\widehat{\boldsymbol{\gamma}}_{j}^{\left[r\right] }$ $\{f_j^{[r]}\}\$ established in Lemmas A1, A2, and A3; see Supplementary A.1.

Lemma [1](#page-14-0). Under Assumption 1 and Assumptions $A1 - A4$, we have

$$
\widehat{\beta}_{\mathrm{Deb},j}^{[r]} - \bar{\beta}_j^{[r]} = \bar{\mathbf{\Omega}}_j^{[r]} \partial_{\beta} \mathcal{L}_r(\bar{\boldsymbol{\beta}}^{[r]}; \bar{\boldsymbol{\alpha}}_j^{[r]}, \bar{\boldsymbol{\gamma}}_j^{[r]}) + O_p(\Delta_r),
$$

where $n_r^{1/2} \bar{\mathbf{\Omega}}_{j}^{[r]} \partial_{\boldsymbol{\beta}} \mathcal{L}_r (\bar{\boldsymbol{\beta}}^{[r]}; \bar{\boldsymbol{\alpha}}_{j}^{[r]})$ $_{j}^{[r]}, \bar{\boldsymbol{\gamma}}_{j}^{[r]}$ $j^{[r]}_j \overset{\mathcal{D}}{\rightarrow} N(0, \sigma_j^2)$ with some $\sigma_j^2 = O(1)$ and

$$
\Delta_r = \frac{K^7(s_{\text{nui}}^{[r]} + s_{\beta}^{[r]})\log(q)}{n_r} + KR_{\text{nui},v}^{[r]} \left\{ \frac{\log(q)}{n_r} \right\}^{1-v/2}.
$$
 (15)

In Lemma [1,](#page-14-1) we derive the asymptotic expansion that each $\hat{\beta}_{\text{Deb},j}^{[r]} = \bar{\beta}_j^{[r]} + \epsilon_j + \text{bias}_j$, where ϵ_j is a variance term with mean 0 and standard error of the order $n_r^{-1/2}$ and bias_j stands for a bias term free of any first-order errors from the nuisance estimators $\hat{\alpha}_j^{[r]}$ $j^{[r]}$ and $\widehat{\bm{\gamma}}^{[r]}_j$. ^[r]. Thus, $\hat{\beta}_{\text{Deb},j}^{[r]}$ is insensitive to the nuisance errors. We shall elaborate more on this point in Remark [4.](#page-15-2)

Remark 3. Note that the sparsity Assumption A2 is not sufficient to achieve the $n_r^{-1/2}$ convergence and asymptotic normality of $\widehat{\beta}_{\mathrm{Deb},j}^{[r]}$ in Lemma [1.](#page-14-1) Neither do we necessarily require this to build up our following convergence theorems. Nonetheless, we can further show that under moderately stronger sparsity conditions $s_{\text{nu}}^{[r]}$, $s_{\beta}^{[r]} = o(n_r^{1/2}/\{K^7 \log(q)\})$ and $R_v^{[r]} = o(n_r^{(1-v)/2}/[K\{\log(q)\}^{1-v/2}])$, $n_r^{1/2}(\widehat{\beta}_{\text{Deb},j}^{[r]} - \bar{\beta}_j^{[r]}$ $j^{[r]}$) will converge to some zero-mean normal distribution, enabling the interval estimation of $\bar{\beta}_i^{[r]}$ $\frac{[r]}{j}$.

Although the element-wise convergence of $\hat{\beta}_{\text{Deb}}^{[r]} = (\hat{\beta}_{\text{Deb},1}^{[r]}, \dots, \hat{\beta}_{\text{Deb},q}^{[r]})^{\top}$ is justified in Lemma [1,](#page-14-1) $\hat{\beta}_{\text{Deb}}^{[r]}$ can still fail to achieve ℓ_2 -consistency that $\|\hat{\beta}_{\text{Deb}}^{[r]} - \bar{\beta}^{[r]}\|_2 = o_p(1)$. As was pointed out in the end of Section [3.1,](#page-7-1) this is because $\hat{\beta}_{\text{Deb}}^{[r]}$ is a dense vector having its estimator error accumulated with an increasing q , which motivates us to employ the thresholding procedure to obtain $\hat{\beta}_{\text{Thr}}^{[r]}$. The ℓ_1 - and ℓ_2 - convergence rates of $\hat{\beta}_{\text{Thr}}^{[r]}$ is given in Theorem [1.](#page-15-1)

Theorem [1](#page-14-0). Under Assumption 1 and Assumptions $AI - A4$ with

$$
\tau^{[r]} = C_0 \sqrt{\log(q)} (n_r^{-1/2} + \Delta_r),
$$

for some constant $C_0 > 0$, we have

$$
\|\widehat{\boldsymbol{\beta}}^{[r]}_{\text{Thr}} - \bar{\boldsymbol{\beta}}^{[r]}\|_1 = O_p(s_{\beta}^{[r]}\tau^{[r]}), \quad \|\widehat{\boldsymbol{\beta}}^{[r]}_{\text{Thr}} - \bar{\boldsymbol{\beta}}^{[r]}\|_2 = O_p(\{s_{\beta}^{[r]}\}^{1/2}\tau^{[r]}).
$$

Remark 4. The error term Δ_r defined in [\(15\)](#page-14-2) encodes the impact of the nuisance estimators on the convergence rate of $\widehat{\beta}_{\text{Deb},j}^{[r]}$ and $\widehat{\beta}_{\text{Thr}}^{[r]}$. With the knowledge of the population nuisance models, i.e., having $\Delta_r = 0$, the error rate given in Theorem [1](#page-15-1) will match the oracle (minimax optimal) rate in the sparse estimation of $\bar{\beta}^{[r]}$. Importantly, when $v = 0$, Δ_r is proportional to an n_r^{-1} -dominated rate $(s_{\text{nu}}^{[r]} + R_{\text{nu}}^{[r]}$ $\lim_{n\to\infty} \log(q)/n_r$, which has a higher order in n_r compared to the oracle part from $\bar{\mathbf{\Omega}}_j^{[r]} \partial_{\boldsymbol{\beta}} \mathcal{L}_r (\bar{\boldsymbol{\beta}}^{[r]}; \bar{\boldsymbol{\alpha}}_j^{[r]})$ $_{j}^{[r]},\bar{\boldsymbol{\gamma}}_{j}^{[r]}$ $j^{[r]}$) = $O_p(n_r^{-1/2})$.

Remark 5. Recall that our preliminary estimator $\widetilde{\boldsymbol{\beta}}^{[r]}$ is also based on a doubly robust formulation but does not include the calibration Steps $2 - 4$ in Algorithm A2. Different $from$ $\widehat{\bm{\beta}}^{[r]}_{\text{Thr}},$ it does not get rid of the nuisance models' first-order error $\{s_{\text{nu}}^{[r]}\log(q)/n_r\}^{1/2}$; see Lemma A2. This implies that our proposed calibration steps can effectively reduce the sensitivity to errors in the nuisance estimators, which is in a similar spirit to the rate-doublyrobust property studied in [Kennedy \(2023\)](#page-26-6). Nevertheless, [Kennedy \(2023\)](#page-26-6) does not establish any model-double-robustness similar to our Assumption [1.](#page-14-0)

4.3 Knowledge transfer learning

In this section, we derive the convergence rates of the knowledge transfer estimator $\hat{\beta}_{\text{K}^{\text{r}}}^{[0]}$ KTr as well as the final version $\hat{\beta}_{MU}^{[0]}$ and demonstrate their improvement over the minority-only estimator $\widehat{\boldsymbol{\beta}}_{\text{Thr}}^{[0]}$.

Lemma 2. Suppose that Assumption [1](#page-14-0) and Assumptions $A1 - A4$ hold with

$$
\tau_{\text{KTr}} = C_1 \sqrt{\log(q)} (n_0^{-1/2} + \Delta),
$$

where $\Delta = \max_{r \in \{0,1\}} \Delta_r$, and $C_1 > 0$ is a constant. Then we have

$$
\|\widehat{\boldsymbol{\delta}}-\bar{\boldsymbol{\delta}}\|_1=O_p\Big(\Big\{\frac{\log(q)}{n_0}\Big\}^{-v/2}R_{\delta,v}\tau_{\text{\tiny KT}}\Big),\quad \|\widehat{\boldsymbol{\delta}}-\bar{\boldsymbol{\delta}}\|_2=O_p\Big(\Big\{\frac{\log(q)}{n_0}\Big\}^{-v/4}R_{\delta,v}^{1/2}\tau_{\text{\tiny KT}}\Big).
$$

Lemma [2](#page-16-1) is an important mediating result about the estimation error on $\bar{\pmb{\delta}} = \bar{\pmb{\beta}}^{[1]} - \bar{\pmb{\beta}}^{[0]}$. Based on this lemma, we derive in Theorem [2](#page-16-2) the ℓ_2 -convergence rate of the knowledge transfer estimator in Algorithm [2.](#page-11-0)

Theorem 2. Under Assumption [1](#page-14-0) and Assumptions $A1 - A4$, we have

$$
\|\widehat{\beta}_{\mathrm{KTr}}^{[0]} - \bar{\beta}^{[0]}\|_2 = O_p\left(\left\{s_{\beta}^{[1]}\log(q)\right\}^{1/2}\left(n_1^{-1/2} + \Delta_1\right) + \frac{\left\{\log(q)\right\}^{1/2 - v/4}}{n_0^{-v/4}}R_{\delta,v}^{1/2}\left(n_0^{-1/2} + \Delta\right)\right). \tag{16}
$$

The first error term on the right hand side of (16) is contributed by the majority estimator $\hat{\beta}_{\text{Thr}}^{[1]}$ and the second term corresponds to the error of $\hat{\delta}$ given in Lemma [2.](#page-16-1) We shall explain the potential efficiency gain of the knowledge transfer Algorithm [2](#page-11-0) in Remark [6.](#page-16-4)

Remark 6. For simplicity, we reasonably assume $s_{\text{nu}}^{[1]} \approx s_{\text{nu}}^{[0]}$ and $R_{\text{nui},v}^{[1]} \approx R_{\text{nui},v}^{[0]}$, which implies that $\Delta_1 = o(\Delta_0) = o(\Delta)$. Also, we take $v = 0$ and consider a typical transferable scenario with $R_{\delta,0} = o(s_{\beta}^{[0]})$ $\stackrel{[0]}{\beta})\,=\,o(s_{\beta}^{[1]}%)\,\delta(s_{\beta}^{[2]})\,\delta(s_{\beta}^{[3]})\,\delta(s_{\beta}^{[3]})$ $\bar{\delta}^{[1]}_{\beta}$), i.e., the difference $\bar{\boldsymbol{\delta}}$ is sparser than the target model coefficients as introduced in Section [2.2.](#page-6-1) Then by Theorems [1](#page-15-1) and [2,](#page-16-2) the ratio of the convergence rates between $\|\widehat{\beta}_{\mathrm{KTr}}^{[0]} - \bar{\beta}^{[0]}\|_2$ and $\|\widehat{\beta}_{\mathrm{Thr}}^{[0]} - \bar{\beta}^{[0]}\|_2$ equals to $(R_{\delta,0}/s_{\beta}^{[0]})^{1/2} = o_p(1)$, $indicating$ a substantial efficiency improvement of the knowledge transfer estimator $\widehat{\boldsymbol \beta}_{\rm KTr}^{[0]}$ over the minority-data-only estimator $\widehat{\boldsymbol{\beta}}^{[0]}_{\text{Thr}}$.

Nevertheless, in a slightly simplified scenario where $\Delta_r = O(n_r^{-1/2})$, $r \in \{0, 1\}$, we can see from Theorem [2](#page-16-2) that whenever $s_{\beta}^{[0]}$ $\delta_{\beta}^{[0]}/s_{\beta}^{[1]} = o(n_0/n_1) \text{ or } s_{\beta}^{[0]} = o(R_{\delta,\nu}\{\log(q)/n_0\}^{-\nu/2}), \widehat{\boldsymbol{\beta}}_{K'}^{[0]}$ KTr will have a larger ℓ_2 -error rate than $\hat{\boldsymbol{\beta}}_{\text{Thr}}^{[0]}$, which is referred to as negative knowledge transfer. To avoid this issue, we propose the model selection Algorithm [3](#page-12-0) to obtain the final estimator $\widehat{\beta}_{\mathrm{MU}}^{[0]}$ and will justify its effectiveness as follows. We begin with an important Lemma [3](#page-17-0) that the $\hat{\boldsymbol{\beta}}_{\rm Deb}^{[0]}$ -based loss function $\hat{\mathcal{Q}}(\boldsymbol{\beta};\hat{\boldsymbol{\beta}}_{\rm Deb}^{[0]})$ offers a good approximation of the ideal model selection criterion $\|\boldsymbol{\beta}-\bar{\boldsymbol{\beta}}^{[0]}\|$ 2 $\frac{2}{2}$.

Lemma 3. Under Assumption [1](#page-14-0) and Assumptions $A1 - A4$, we have

$$
\widehat{\mathcal{Q}}(\boldsymbol{\beta};\widehat{\boldsymbol{\beta}}_{\mathrm{Deb}}^{[0]}) = ||\boldsymbol{\beta} - \bar{\boldsymbol{\beta}}^{[0]}||_2^2 + 2(\boldsymbol{\beta} - \bar{\boldsymbol{\beta}}^{[0]})^{\mathsf{T}}(\boldsymbol{\mathcal{E}} + \mathrm{Rem}) + C_2,
$$

where $\mathcal{E}=\big\{ \bar{\mathbf{\Omega}}_1^{[0]} \partial_{\boldsymbol{\beta}} \mathcal{L}_r (\bar{\boldsymbol{\beta}}_1^{[0]}$ $_{1}^{[0]};\bar{\boldsymbol{\alpha}}_{1}^{[0]}$ $_{1}^{\left[0\right] },\bar{\boldsymbol{\gamma}}_{1}^{\left[0\right] }$ $\{\bar{\Omega}_1^{[0]}) , \ldots , \bar{\mathbf{\Omega}}_q^{[0]} \partial_{\boldsymbol{\beta}} \mathcal{L}_r (\bar{\boldsymbol{\beta}}_q^{[0]})$ ${[0] \over q}; \bar{\boldsymbol{\alpha}}_q^{[0]}, \bar{\boldsymbol{\gamma}}_q^{[0]})\}^\intercal$ is a mean-zero and concentrated empirical process satisfying $\|\mathcal{E}\|_{\infty} = O_p[\{\log(q)/n_0\}^{1/2}]$, and Rem stands for the reminder approximation-error term satisfying $\|\text{Rem}\|_{\infty} = O_p[\{\log(q)\}^{1/2} \Delta_0]$, and C_2 is a value free of the parameter β .

As discussed in Remark [4,](#page-15-2) Rem, with its maximum norm controlled by $\{\log(q)\}^{1/2}\Delta_0$, is a reminder error term encoding the second-order influence of the nuisance estimators. Meanwhile, the term $2(\beta - \bar{\beta}^{[0]})^T \mathcal{E}$ can be properly controlled through the concentration of $\mathcal E$. Leveraging this lemma as well as Theorems [1](#page-15-1) and [2,](#page-16-2) we derive the convergence rate of our final estimator $\hat{\beta}_{MU}^{[0]}$ in Theorem [3.](#page-17-1)

Theorem 3. Under Assumption [1](#page-14-0) and Assumptions $A1 - A4$, we have

$$
\|\widehat{\beta}_{\mathrm{MU}}^{[0]} - \bar{\beta}^{[0]}\|_2 = O_p\Big\{\Big(\|\widehat{\beta}_{\mathrm{Thr}}^{[0]} - \bar{\beta}^{[0]}\|_2 + \mathrm{DetectErr}\Big) \wedge \|\widehat{\beta}_{\mathrm{KTr}}^{[0]} - \bar{\beta}^{[0]}\|_2\Big\},\
$$

with the detection error term

$$
\text{DetectErr} = \left\{ \left(n_0^{-1/2} + \Delta_0 \right) \left(\left\| \widehat{\boldsymbol{\beta}}_{\text{KTr}}^{[0]} - \bar{\boldsymbol{\beta}}^{[0]} \right\|_2 + \left\| \widehat{\boldsymbol{\beta}}_{\text{Thr}}^{[0]} - \bar{\boldsymbol{\beta}}^{[0]} \right\|_2 \right) \right\}^{1/2}.
$$

When $\Delta_r = o(n_r^{-1/2})$, we have

$$
\text{DetectErr} = \left(n_0^{-1/2} \max_{r=0,1} \left\{ \frac{s_\beta^{[r]} \log(q)}{n_r} \right\}^{1/2} + n_0^{-1/2} \left\{ \frac{\log(q)}{n_0} \right\}^{1/2 - v/4} R_{\delta,v}^{1/2} \right\}^{1/2}.
$$

When further assuming $R_{\delta,v} = o({n_0 / \log(q)}^{1/2 - v/2})$, DetectErr = $o_p[n_0^{-3/8} {\log(q)}^{1/8}]$.

In Theorem [3,](#page-17-1) we show that the ℓ_2 -error of $\widehat{\boldsymbol{\beta}}_{MU}^{[0]}$ is ensured to achieve faster convergence rate between those of the minority-only $\hat{\beta}_{\text{Thr}}^{[0]}$ and the knowledge transfer $\hat{\beta}_{\text{KTr}}^{[0]}$, with the minority-only part additionally including DetectErr as the detection error incurred by model selection in Algorithm [3.](#page-12-0) We also show such a price satisfies $\text{DetectErr} = o_p[n_0^{-3/8} \{\log(q)\}^{1/8}]$ when further imposing some reasonable assumptions including that the higher-order nuisance error Δ_r is negligible as discussed in Remark [3](#page-15-3) and $R_{\delta,v} = o(\lbrace n_0/\log(q) \rbrace^{1/2-v/2})$, i.e., $\bar{\delta}$ is ultra-sparse when $v = 0$. This price is relatively light as $n_0^{-3/8}$ $_0^{-3/8}$ is not that far from the parametric rate $n_0^{-1/2}$ $\frac{-1/2}{0}$.

Remark 7. Note that our main results about knowledge transfer in Theorems [2](#page-16-2) and [3](#page-17-1) have a similar spirit as those established in recent transfer learning literature like [Li et al. \(2022\)](#page-26-3) and [Tian and Feng \(2023\)](#page-27-6). Nevertheless, our method and theory are more technically involved. This is mainly because our setup does not include actual labels on $\mathcal T$, and we leverage the dense $\widehat{\beta}_{\rm Deb}^{[0]}$ to approximate the covariate shift corrected objective for $\bar{\beta}^{[0]}$ with the needs in addressing nuisance model errors; see Lemma [3.](#page-17-0)

5 Simulation Studies

We investigate the finite-sample performance of the proposed method under various scenarios. Let $\mathbf{Z} = (Z_1, Z_2, \ldots, Z_{p+q})^{\dagger}$ be a vector of independent variables, where $Z_1 = 1$ and each Z_j for $j \geq 2$ follows a standard normal distribution truncated to the interval $(-1.5, 1.5)$. Let $g(a) = b(a) = \text{expit}(a) = e^a/(1+e^a), \, \mathbf{X} = (Z_1, \ldots, Z_q)^{\mathsf{T}}, \, \mathbf{W} = (Z_{q+1}, \ldots, Z_{q+p})^{\mathsf{T}}, \, \phi(\mathbf{Z}) = \mathbf{Z},$ and $\delta_t = (0, 0.2 \times \mathbf{1}_t^{\tau}, \mathbf{0}_{p+q-t-1}^{\tau})^{\tau}$, where t is a hyperparameter to be specified later. We consider the following three data-generating configurations for the outcome Y and the source indicator S, where $Y \mid (\mathbf{Z}, R)$ and $S \mid (\mathbf{Z}, R)$ follow Bernoulli distributions independently.

- (I) Both nuisance models are correctly specified: $\mathbb{P}(Y = 1 | \mathbf{Z}, R = r) = \text{expit}(\mathbf{Z}^{\mathsf{T}}(\boldsymbol{\gamma} + r\boldsymbol{\delta}_{t})),$ and $\mathbb{P}(S = 1 | \mathbf{Z}, R = r) = \text{expit}(\mathbf{Z}^T \boldsymbol{\alpha})$, where $\boldsymbol{\gamma} = (0, 1.2 \times \mathbf{1}_3^T, \mathbf{0}_{q-4}^T, 1, 0.8, 0.5, \mathbf{0}_{p-3}^T)^T$ and $\boldsymbol{\alpha} = (0, 0.2 \times \mathbf{1}_3^{\text{T}}, \mathbf{0}_{q-4}^{\text{T}}, 0.5 \times \mathbf{1}_3^{\text{T}}, \mathbf{0}_{p-3}^{\text{T}})^{\text{T}}$.
- (II) The density ratio model is correct while the imputation model is misspecified: $\mathbb{P}(Y =$ $1 | \mathbf{Z}, R = r$ = expit $(\psi_1(\mathbf{Z}) - r(0.2Z_4 + 0.5Z_{q+1} + \mathbf{Z}^T \delta_t))$ and $\mathbb{P}(S = 1 | \mathbf{Z}, R = r)$ = $\text{expit}(\mathbf{Z}^{\mathsf{T}}\boldsymbol{\alpha})$, where $\psi_1(\mathbf{Z}) = -1.5Z_1 + 1.2(|Z_2| + 0.9Z_2) + 1.2(|Z_3| + 0.9Z_3) + 1.2Z_4$ and $\boldsymbol{\alpha} = (0, -1, -1, \mathbf{0}_{q-3}^{\intercal}, 0.5, \mathbf{0}_{p-1}^{\intercal})^{\intercal}.$
- (III) The imputation model is correct while the density ratio model is misspecified: $\mathbb{P}(Y =$ $1 \mid \mathbf{Z}, R = r$ = expit $(\mathbf{Z}^{\mathsf{T}}(\boldsymbol{\gamma} + r\boldsymbol{\delta}_t))$ and $\mathbb{P}(S = 1 \mid \mathbf{Z}, R = r) = \text{expit}(\psi_2(\mathbf{Z})),$ where $\boldsymbol{\gamma} = (0, 0.7, 0.9, 0.8, \mathbf{0}_{q-4}^{\mathsf{T}}, 0.75, -0.75, \mathbf{0}_{p-2}^{\mathsf{T}})^{\mathsf{T}}$ and $\psi_2(\mathbf{Z}) = -0.2Z_1 - 0.8Z_2Z_{q+1} 0.5Z_3Z_{q+2}$.

In Settings (I) and (III), we introduce the data-generation parameter $t \in \mathbb{Z}$ to control the outcome model heterogeneity between the majority and minority groups. In specific, t is the number of non-zero entries of the model difference between the two sub-populations $\delta_t = \bar{\beta}^{[1]} - \bar{\beta}^{[0]}$. When $t = 0$, the majority and minority groups share the same datagenerating mechanism, resulting in $\beta^{[0]} = \beta^{[1]}$. When t gets larger, there will be stronger model heterogeneity between the two groups.

We consider our MAKEUP algorithm with protection against negative transfer (Algo-rithm [3\)](#page-12-0), where the temperature parameter a is set to 5, denoted as MU. We also consider the version without protection against negative transfer (Algorithm [2\)](#page-11-0), denoted as $MU_{\text{maj-g}}$.

We also include the minority-only estimator $\hat{\beta}_{\text{Thr}}^{[0]}$ denoted as MU_{min-o}. We compare our method against two simpler covariate shift correction strategies including: (i) the importance weighting method with ℓ_1 penalty (IW) and with weighted ℓ_1 penalty (IW_{aLasso}) detailed in Algorithm A1, and (ii) the imputation-only method with ℓ_1 penalty (IM) and with weighted ℓ_1 penalty (IM_{aLasso}) detailed in Algorithm A2 of Supplementary C.1. For comparison, we also include the TransGLM method [\(Tian and Feng, 2023\)](#page-27-6) with two variants according to our setup. The first version (TransGLM) directly uses labeled source data to learn parameters by transferring information from the majority to the minority group, without accounting for covariate shifts between the target and source populations. The second variant (Trans GLM_{iw}) adjusts for covariate shifts using importance weighting. The details of these two variants are provided in Algorithm A3 in Supplementary C.1.

We fix the sample size in the majority group as $n_{\text{s},1} = 3000$, $n_{\text{T},1} = 5000$, and the sample size of the underrepresented group in the unlabeled target population as $n_{\tau,0} = 1000$. We then vary the sample size of the labeled source population in the underrepresented group $n_{s,0} \in \{300, 400, 500, 600\}$. We set $q = 100, p \in \{100, 500\}$. Each setting is repeated 100 times to estimate the mean ℓ_2 error $\|\hat{\boldsymbol{\beta}} - \bar{\boldsymbol{\beta}}^{[0]}\|_2$ for any estimate $\hat{\boldsymbol{\beta}}$. We also vary the value of $t \in \{0, 1, 2, 3\}$ for Settings I and III to reflect the discrepancy between the majority and minority groups. The tuning parameters λ 's for these comparable methods are decided by five-fold cross-validation. We select the penalty parameters λ_{α} and λ_{γ} using cross-validation from the range $[0.01(\log p/n_{\tau,r})^{1/2}, 0.5(\log p/n_{\tau,r})^{1/2}]$, and select $\lambda_{\alpha_j}^{[r]}, \lambda_{\gamma}^j$, and τ using the tuning procedure introduced in Supplementary B.3.

The results for $q = 100$, $p = 500$, and $t = 1$ with different sample sizes of $n_{s,0}$ are summarized in Table [1.](#page-20-0) The results for $p = 100$ show a similar pattern to $p = 500$, and are provided in Table A.1 in Supplementary C.2. The ℓ_2 errors of most methods decrease as the sample size $n_{s,0}$ increases. Our proposed MU and MU_{maj-g} methods consistently show the best performance, with our final version, MU, moderately outperforming $MU_{\text{maj-g}}$. This is likely due to better handling of model heterogeneity since $t \neq 0$. MU_{min-o} improves with larger sample sizes but still lags behind MU and $MU_{\text{maj-g}}$.

The benchmark methods IW, IWaLasso, IM, and IMaLasso show better performance with increased sample sizes but remain less effective overall. In Setting II, where only the density ratio model is correct, IM performs poorly due to the incorrect imputation model specification. In Setting III, IWaLasso performs the worst due to the incorrect density ratio model. The performance of TransGLM remains relatively unchanged with increasing sample size, as it does not account for covariate shifts, leading to biased estimates. In contrast, Trans GLM_{iw} shows improved performance in Settings I and II when $n_{s,0}$ increases, where the density ratio model is correctly specified, due to its adjustment for covariate shift. However, in Setting III, where the density ratio model is misspecified, $TransGLM_{iw}$'s performance remains largely unchanged with increased sample size. Overall, our proposed MAKEUP method demonstrate superior performance across different settings, as it effectively handles both the covariate

Method	Setting I				Setting II				Setting III			
$n_{\mathcal{S},0}$	300	400	500	600	300	400	500	600	300	400	500	600
MU	0.10	0.07	0.06	0.05	0.48	0.42	0.30	0.31	0.14	0.09	0.08	0.06
MU_{maj-g}	0.16	0.12	0.08	0.07	0.48	0.40	0.32	0.29	0.20	0.14	0.10	0.08
$MU_{\text{min-o}}$	0.29	0.18	0.15	0.11	2.08	1.64	1.37	1.22	0.38	0.18	0.12	0.08
ΙW	0.77	0.58	0.53	0.44	1.31	1.09	0.86	0.87	0.68	0.57	0.50	0.47
IW_{aLasso}	1.20	0.99	0.61	0.64	1.73	1.09	0.93	0.74	1.06	0.83	0.68	0.55
ΙM	1.12	0.86	0.73	0.63	2.46	2.19	2.07	1.95	0.82	0.62	0.51	0.44
IM_{aLasso}	0.55	0.39	0.32	0.28	1.51	1.33	1.20	1.09	0.40	0.29	0.25	0.22
TransGLM	0.69	0.64	0.67	0.67	1.05	1.03	1.02	0.98	0.25	0.24	0.24	0.23
$TransGLM_{iw}$	0.36	0.30	0.29	0.28	0.69	0.67	0.60	0.58	0.28	0.26	0.26	0.25

Table 1: Empirical ℓ_2 error averaged over 100 repetitions for all methods with $q = 100$, $p = 500, t = 1.$

Method			Setting I		Setting III					
t	θ		$\overline{2}$	3	0		$\overline{2}$	3		
MU	0.07	0.07	0.08	0.08	0.07	0.09	0.10	0.11		
MU_{maj-g}	0.11	0.12	0.14	0.14	0.12	0.14	0.15	0.17		
MU_{min-o}	0.19	0.18	0.18	0.18	0.18	0.18	0.18	0.18		
IW	0.58	0.58	0.58	0.60	0.58	0.57	0.57	0.57		
IW_{aLasso}	1.03	0.99	0.98	0.98	0.83	0.83	0.80	0.83		
ĪМ	0.86	0.86	0.86	0.86	0.62	0.62	0.63	0.62		
IM_{aLasso}	0.39	0.39	0.39	0.39	0.29	0.29	0.29	0.29		
TransGLM	0.69	0.64	0.61	0.58	0.36	0.24	0.27	0.24		
$TransGLM_{iw}$	0.35	0.30	0.27	0.21	0.38	0.26	0.29	0.25		

Table 2: Empirical ℓ_2 error averaged over 100 repetitions for all methods with $q = 100$, $p = 500, n_{s,0} = 400$. The parameter t for δ_t varies from 0 to 3.

shift correction and model knowledge transfer.

In addition, we investigate the performance of the proposed method under different levels of model heterogeneity, focusing on settings where the imputation model is correctly specified. The results for $p = 500$ with $n_{s,0} = 400$ are summarized in Table [2,](#page-20-1) while the results for $p = 100$ are given in Table A2 in Supplementary C.2.

From Table [2,](#page-20-1) we can see that MU consistently performs the best across all settings, followed by $MU_{\text{maj-g}}$ and $MU_{\text{min-o}}$. As t increases, MU maintains a lower error, which implies the benefit of protecting negative transfer. When t is small, both MU and $MU_{\text{maj-g}}$ outperform MU_{min-o} , which only uses the minority data. For example, when $t = 0$, MU has empirical ℓ_2 errors less than 50% of MU_{min-o} in Settings I and III. Even with t as large as 3, MU and $MU_{\text{maj-g}}$ still perform better than $MU_{\text{min-o}}$ in Setting I. The performance of IW, $\text{IW}_{\text{aLasso}}$, IM, and $\text{IM}_{\text{aLasso}}$ remains relatively stable across different t values, as they only use the minority group data.

6 Real Data Analysis

Type II diabetes (T2D), a common chronic disease caused by insulin insufficiency, places a substantial economic and social burden on society. In the United States alone, the total estimated cost of diagnosed diabetes reached \$412.9 billion in 2022 [\(Parker et al., 2024\)](#page-27-10). Accurately predicting the risk of developing Type 2 Diabetes is crucial for early intervention, prevention, and reducing the long-term health and economic impacts associated with the disease. Extensive genetic studies have suggested that genetic factors play a significant role in the risk of T2D and some non-white racial groups including African Americans and hispanic and Latin Americans have a higher genetic predisposition to T2D [\(Mercader and](#page-27-11) [Florez, 2017;](#page-27-11) [Mahajan et al., 2018;](#page-26-8) [Armstrong et al., 2024,](#page-24-8) e.g.).

In our study, we aim to enable the use of EHR-linked biobank data from Mass General Brigham (MGB) biobank [\(Castro et al., 2022\)](#page-24-1) in deriving genetic risk prediction models optimized for underrepresented populations by transferring knowledge from the majority population. On one hand, both the demographic variables and single nucleotide polymorphisms (SNPs) X and the EHR features W are readily available for all MGB biobank patients. On the other hand, the gold standard label for the T2D status, $Y \in \{0, 1\}$, has only been collected for a subset of $n_{\text{s,i}} = 375$ for the majority (White race) and $n_{\text{s,o}} = 77$ for the minority (non-White race) patients, whose medical records were manually reviewed in a biomedical study in 2014. In this study, to obtain gold standard labels for multiple phenotypes more efficiently, patients with International Classification of Diseases (ICD) codes for several phenotypes, such as rheumatoid arthritis and coronary artery disease, were sampled and reviewed simultaneously, leading to sampling that was not completely at random. The target data consists of $n_{\text{s},1} = 5000$ and $n_{\text{s},0} = 1000$ patients drawn from the MGB biobank participants with their EHR features updated in 2021. Importantly, the EHR system at MGB, as well as its ICD version, changed around 2015. Thus, the shift in EHR and genomic features between the source and target samples can be attributed to both variation in the sampling method and the difference in the time window of data collection.

Our goal is to construct a genetic risk model for the T2D status Y using the highdimensional demographic and genetic variants $\mathbf{X} \in \mathbb{R}^{272}$. To adjust for the distributional shift between the source and target samples, We include 66 EHR features denoted as W , consisting of T2D-relevant diagnostic and procedure codes and a measure of total health utilization. These features are selected from the complete range of EHR variables by implementing the clinical knowledge extraction tool developed by [Hong et al. \(2021\)](#page-25-8), which is independent of the dataset in this study. In this way, the two nuisance models are fitted on $\mathbf{Z} = (\mathbf{X}^{\mathsf{T}}, \mathbf{W}^{\mathsf{T}})^{\mathsf{T}}$, which is clearly high dimensionality given the labeled sample sizes $n_{s,0}$ and $n_{s,1}$. Note that a similar transfer learning problem on the MGB data has been considered by [Zhou et al.](#page-28-4) [\(2024\)](#page-28-4). However, unlike their study, we aim to perform risk modeling using the original high-dimensional genetic variants, rather than a one-dimensional genetic score combining the SNPs with β -coefficients extracted from existing genome-wide association studies. This makes our study more flexible and data-driven but introduces new statistical challenges, as discussed in previous sections.

A validation dataset $\mathcal V$ is created by randomly selecting and labeling 47 individuals from the target minority population in order to evaluate the transfer learning methods. To measure predictive performance, we calculate the following metrics on \mathcal{V} : (1) Brier skill score (BSS), defined as $1 - \widehat{\mathbb{E}}_{\mathcal{V}} \{ Y - g(\mathbf{X}^{\mathsf{T}} \boldsymbol{\beta}) \}^2 / \widehat{\text{Var}}_{\mathcal{V}}(Y)$ for some estimator $\boldsymbol{\beta}$, where $\widehat{\mathbb{E}}_{\mathcal{V}}$ and $\text{Var}_{\mathcal{V}}$ respectively denote the empirical mean and variance operators on the validation samples; (2) Goodness-of-fit (GOF), measured by negative deviance on the validation data: $1-2\mathbb{E}_{\mathcal{V}}\{-Y\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta}+G(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})\};$ and (3) area under the receiver operating characteristic curve (AUC) of the predictor $g(\mathbf{X}^{\mathsf{T}}\boldsymbol{\beta})$ on \mathcal{V} .

Following our simulation studies, we applied our proposed MAKEUP method MU along with IW, IW_{aLasso}, IM, IM_{aLasso}, TransGLM, and TransGLM_{iw} to construct the genetic risk model for T2D. Additionally, we incorporated two unsupervised phenotyping algorithms, PheNorm [\(Yu et al., 2018\)](#page-28-8) and MAP [\(Liao et al., 2019\)](#page-26-9), for comparison. PheNorm first uses a regularized denoising regression to condense EHR features into low-dimensional risk scores and then performs clustering on them to derive the phenotype. MAP ensembles Gaussian and Poisson mixture models on the main surrogates of the target disease for phenotyping. We implemented these algorithms with the EHR features W on the unlabeled minority data and used their imputed phenotypes as the outcome to regress against X . This strategy is similar to the IM approach, and we denote the two benchmarks as $IM_{PheNorm}$ and IM_{MAP} .

We evaluated the performance of these methods on the validation set \mathcal{V} , with results presented in Table [3.](#page-22-0) Among all the methods tested, MU demonstrated the best performance across all three evaluation metrics. Specifically, it significantly outperformed methods that rely solely on a single nuisance model, achieving approximately 30% higher BSS compared to IMaLasso and much higher GOF than IWaLasso. Furthermore, MU showed a moderate improvement over TransGLM and TransGLM_{iw}, with an approximately 25% higher AUC. While the unsupervised algorithms PheNorm and MAP performed well in terms of AUC, they exhibited much lower BSS and GOF values compared to MU.

	ΜU	IW	$\rm{IW}_{\rm{aLasso}}$	IМ	$\rm IM_{aLasso}$	TransGLM	$TransGLM_{iw}$	$M_{PheNorm}$	$1M_{MAP}$
BSS	0.36	-0.11	-0.22	0.23	0.24	0.00	0.00	-0.19	-0.16
GOF	0.16	-0.85	-1.87	-0.02	-0.02	-0.82	-0.78	-0.45	-0.42
AUC	0.89	0.87	0.62	0.81	0.83	0.70	0.71	0.84	0.85

Table 3: Predictive performance of the T2D risk models evaluated on the validation data. The evaluation metrics BSS, GOF, and AUC are defined in Section [6.](#page-21-0) Descriptions of the benchmark methods can be found in Sections [5](#page-18-0) and [6.](#page-21-0)

7 Discussion

We develop MAKEUP, a novel transfer learning approach for high-dimensional regression on an underrepresented group without any labeled samples. MAKEUP simultaneously addresses covariate shift and model heterogeneity. To correct for covariate shift, our modelassisted method achieves both model-double-robustness and rate-double-robustness, making it insensitive to the misspecification or poor convergence rate of either nuisance model. Unlike recent literature on doubly robust estimation with high-dimensional nuisance models and low-dimensional target parameters [\(Tan, 2020;](#page-27-8) [Zhou et al., 2024,](#page-28-4) e.g.), our target estimator is sparse-regularized and lacks asymptotic linearity, which makes calibrating the nuisance models more challenging. Our key debiasing step address this issue and enables proper calibration, while the downstream sparsifying step removes error accumulated in the dense debiased estimator. This approach can be generalized to other important setting, such as estimating the conditional average treatment effect (CATE) with high-dimensional effect modifiers, as considered in [Kato \(2024\)](#page-25-6) and others.

Meanwhile, we incorporate the idea of knowledge-guided shrinkage to mitigate the disparity in data collection for the underrepresented subgroup. Compared to existing literature on model knowledge transfer [\(Li et al., 2022,](#page-26-3) e.g.), we tackle a more challenging setup where the target cohort is unlabeled, and the debiased coefficient vector obtained in the previous step is used as a substitute for actual labeled samples in sparse regression and negative transfer protection. This makes our method and theory in this part more technically involved.

In data-driven biomedical studies, as well as in fields like economics and social science, both covariate shift from unlabeled target and sampling disparity for underrepresented subpopulation are prominent challenges that impede generalizable and responsible statistical learning. For example, leveraging a clinical trial to infer treatment effects on an observational cohort will also encounter the covariate shift between the two cohorts [\(Colnet et al.,](#page-25-9) [2024,](#page-25-9) e.g.), and age and gender disparities have been frequently considered in recent studies [\(Ting et al., 2017;](#page-28-9) [Ludmir et al., 2019,](#page-26-10) e.g.). Though these two challenges often co-occur in practice, existing analytical tools typically address only one at a time. Our work fills this methodological gap by simultaneously addressing both challenges in a coherent and complete framework, demonstrating potential for wide application.

Lastly, we shall point out several limitations and future directions of our work. First, the computational costs of MAKEUP could be high, as it requires debiasing and calibration for each coefficient separately in the high-dimensional outcome model. This issue also significantly increases the number of tuning parameters in our method. It would be desirable to develop simpler and faster methods that can achieve similar robustness and efficiency properties to our current approach. Second, it is common to have multiple source cohorts in practice, e.g., more than one institution with labeled sample in their databases. Thus, there is interest and a need to extend our method to the federated transfer learning setup [\(Li et al., 2023;](#page-26-5) [Tian et al., 2023,](#page-27-12) e.g.). In this scenario, non-informative sources and privacy constraints need to be carefully addressed. Third, while we currently focus on transfer learning for risk prediction models, it could be useful to generalize our framework to other more complex learning tasks, such as reinforcement learning and individualized treatment effect estimation.

References

- Amarasingham, R., Patzer, R. E., Huesch, M., Nguyen, N. Q., and Xie, B. (2014). Implementing electronic health care predictive analytics: considerations and challenges. Health $a\frac{f}{a}$ affairs, 33(7):1148–1154.
- Armstrong, N. D., Patki, A., Srinivasasainagendra, V., Ge, T., Lange, L. A., Kottyan, L., Namjou, B., Shah, A. S., Rasmussen-Torvik, L. J., Jarvik, G. P., et al. (2024). Variant level heritability estimates of type 2 diabetes in african americans. Scientific Reports, 14(1):14009.
- Bastani, H. (2021). Predicting with proxies: Transfer learning in high dimension. Management Science, 67(5):2964–2984.
- Baybutt, A. and Navjeevan, M. (2023). Doubly-robust inference for conditional average treatment effects with high-dimensional controls. arXiv preprint arXiv:2301.06283.
- Braithwaite, J. (2018). Changing how we think about healthcare improvement. Bmj, 361.
- Cai, T., Li, M., and Liu, M. (2024). Semi-supervised triply robust inductive transfer learning. Journal of the American Statistical Association, (just-accepted).
- Castro, V. M., Gainer, V., Wattanasin, N., Benoit, B., Cagan, A., Ghosh, B., Goryachev, S., Metta, R., Park, H., Wang, D., et al. (2022). The mass general brigham biobank portal: an i2b2-based data repository linking disparate and high-dimensional patient data to support multimodal analytics. Journal of the American Medical Informatics Association, 29(4):643–651.
- Chakrabortty, A., Lu, J., Cai, T. T., and Li, H. (2019). High dimensional M-estimation with missing outcomes: A semi-parametric framework. arXiv preprint arXiv:1911.11345.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. The Econometrics Journal, 21(1):C1–C68.
- Colnet, B., Mayer, I., Chen, G., Dieng, A., Li, R., Varoquaux, G., Vert, J.-P., Josse, J., and Yang, S. (2024). Causal inference methods for combining randomized trials and observational studies: a review. Statistical Science, 39(1):165–191.
- Dalalyan, A. S. and Tsybakov, A. B. (2007). Aggregation by exponential weighting and sharp oracle inequalities. In *International Conference on Computational Learning Theory*, pages 97–111. Springer.
- Davis, S. E., Lasko, T. A., Chen, G., Siew, E. D., and Matheny, M. E. (2017). Calibration drift in regression and machine learning models for acute kidney injury. Journal of the American Medical Informatics Association, 24(6):1052–1061.
- De Lillo, A., D'Antona, S., Pathak, G. A., Wendt, F. R., De Angelis, F., Fuciarelli, M., and Polimanti, R. (2021). Cross-ancestry genome-wide association studies identified heterogeneous loci associated with differences of allele frequency and regulome tagging between participants of european descent and other ancestry groups from the uk biobank. Human Molecular Genetics, 30(15):1457–1467.
- Fan, Q., Hsu, Y.-C., Lieli, R. P., and Zhang, Y. (2022). Estimation of conditional average treatment effects with high-dimensional data. Journal of Business $\mathcal C$ Economic Statistics, 40(1):313–327.
- Guo, L. L., Pfohl, S. R., Fries, J., Posada, J., Fleming, S. L., Aftandilian, C., Shah, N., and Sung, L. (2021). Systematic review of approaches to preserve machine learning performance in the presence of temporal dataset shift in clinical medicine. Applied Clinical Informatics, 12(04):808–815.
- He, Z., Sun, Y., and Li, R. (2024). Transfusion: Covariate-shift robust transfer learning for high-dimensional regression. In International Conference on Artificial Intelligence and Statistics, pages 703–711. PMLR.
- Hong, C., Rush, E., Liu, M., Zhou, D., Sun, J., Sonabend, A., Castro, V. M., Schubert, P., Panickan, V. A., Cai, T., et al. (2021). Clinical knowledge extraction via sparse embedding regression (KESER) with multi-center large scale electronic health record data. NPJ digital medicine, $4(1):1-11$.
- Huang, J., Gretton, A., Borgwardt, K., Schölkopf, B., and Smola, A. J. (2007). Correcting sample selection bias by unlabeled data. In Advances in Neural Information Processing Systems, pages 601–608.
- Kato, M. (2024). Triple/debiased lasso for statistical inference of conditional average treatment effects. arXiv preprint arXiv:2403.03240.
- Kennedy, E. H. (2023). Towards optimal doubly robust estimation of heterogeneous causal effects. Electronic Journal of Statistics, 17(2):3008–3049.
- Lam, M., Chen, C.-Y., Li, Z., Martin, A. R., Bryois, J., Ma, X., Gaspar, H., Ikeda, M., Benyamin, B., Brown, B. C., et al. (2019). Comparative genetic architectures of schizophrenia in east asian and european populations. Nature genetics, 51(12):1670–1678.
- Li, R., Chen, Y., Ritchie, M. D., and Moore, J. H. (2020). Electronic health records and polygenic risk scores for predicting disease risk. Nature Reviews Genetics, 21(8):493–502.
- Li, S., Cai, T., and Duan, R. (2023). Targeting underrepresented populations in precision medicine: A federated transfer learning approach. The Annals of Applied Statistics, 17(4):2970–2992.
- Li, S., Cai, T. T., and Li, H. (2022). Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. Journal of the Royal Statistical Society Series B: Statistical Methodology, 84(1):149–173.
- Liao, K. P., Sun, J., Cai, T. A., Link, N., Hong, C., Huang, J., Huffman, J. E., Gronsbell, J., Zhang, Y., Ho, Y.-L., et al. (2019). High-throughput multimodal automated phenotyping (map) with application to phewas. Journal of the American Medical Informatics Association, 26(11):1255-1262.
- Lin, H. and Reimherr, M. (2024). Smoothness adaptive hypothesis transfer learning. *arXiv* preprint arXiv:2402.14966.
- Liu, M., Zhang, Y., Liao, K. P., and Cai, T. (2023). Augmented transfer regression learning with semi-non-parametric nuisance models. Journal of Machine Learning Research, $24(293):1-50.$
- Ludmir, E. B., Mainwaring, W., Lin, T. A., Miller, A. B., Jethanandani, A., Espinoza, A. F., Mandel, J. J., Lin, S. H., Smith, B. D., Smith, G. L., et al. (2019). Factors associated with age disparities among cancer clinical trial participants. JAMA oncology, 5(12):1769–1773.
- Mahajan, A., Taliun, D., Thurner, M., Robertson, N. R., Torres, J. M., Rayner, N. W., Payne, A. J., Steinthorsdottir, V., Scott, R. A., Grarup, N., et al. (2018). Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and isletspecific epigenome maps. Nature genetics, 50(11):1505–1513.
- Martin, A. R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B. M., and Daly, M. J. (2019). Clinical use of current polygenic risk scores may exacerbate health disparities. Nature genetics, 51(4):584–591.
- Matheny, M. E., Miller, R. A., Ikizler, T. A., Waitman, L. R., Denny, J. C., Schildcrout, J. S., Dittus, R. S., and Peterson, J. F. (2010). Development of inpatient risk stratification models of acute kidney injury for use in electronic health records. Medical Decision Making, 30(6):639–650.
- Mercader, J. M. and Florez, J. C. (2017). The genetic basis of type 2 diabetes in hispanics and latin americans: challenges and opportunities. Frontiers in Public Health, 5:329.
- Moons, K. G., Kengne, A. P., Grobbee, D. E., Royston, P., Vergouwe, Y., Altman, D. G., and Woodward, M. (2012). Risk prediction models: Ii. external validation, model updating, and impact assessment. Heart, 98(9):691–698.
- Parikh, R. B., Kakad, M., and Bates, D. W. (2016). Integrating predictive analytics into high-value care: the dawn of precision delivery. *JAMA*, 315(7):651–652.
- Parker, E. D., Lin, J., Mahoney, T., Ume, N., Yang, G., Gabbay, R. A., ElSayed, N. A., and Bannuru, R. R. (2024). Economic costs of diabetes in the us in 2022. Diabetes Care, $47(1):26-43.$
- Qiu, H., Tchetgen, E. T., and Dobriban, E. (2023). Efficient and multiply robust risk estimation under general forms of dataset shift. *arXiv preprint arXiv:2306.16406*.
- Reddi, S. J., Poczos, B., and Smola, A. (2015). Doubly robust covariate shift correction. In Twenty-Ninth AAAI Conference on Artificial Intelligence.
- Semenova, V. and Chernozhukov, V. (2021). Debiased machine learning of conditional average treatment effects and other causal functions. The Econometrics Journal, 24(2):264– 289.
- Smucler, E., Rotnitzky, A., and Robins, J. M. (2019). A unifying approach for doubly-robust ℓ_1 -regularized estimation of causal contrasts. arXiv preprint arXiv:1904.03737.
- Steyerberg, E. W., Moons, K. G., van der Windt, D. A., Hayden, J. A., Perel, P., Schroter, S., Riley, R. D., Hemingway, H., Altman, D. G., and Group, P. (2013). Prognosis research strategy (progress) 3: prognostic model research. *PLoS medicine*, 10(2):e1001381.
- Tan, Z. (2020). Model-assisted inference for treatment effects using regularized calibrated estimation with high-dimensional data. Annals of Statistics, 48(2):811–837.
- Tian, Y. and Feng, Y. (2023). Transfer learning under high-dimensional generalized linear models. Journal of the American Statistical Association, 118(544):2684–2697.
- Tian, Y., Gu, Y., and Feng, Y. (2023). Learning from similar linear representations: Adaptivity, minimaxity, and robustness. arXiv preprint arXiv:2303.17765.
- Tian, Y., Wu, P., and Tan, Z. (2024). Semi-supervised regression analysis with model misspecification and high-dimensional data. $arXiv$ preprint $arXiv:2406.13906$.
- Ting, P.-S., Chen, L., Yang, W.-C., Huang, T.-S., Wu, C.-C., and Chen, Y.-Y. (2017). Gender and age disparity in the initiation of life-supporting treatments: a population-based cohort study. BMC Medical Ethics, 18:1–9.
- Van de Geer, S., Bühlmann, P., Ritov, Y., Dezeure, R., et al. (2014). On asymptotically optimal confidence regions and tests for high-dimensional models. The Annals of Statistics, 42(3):1166–1202.
- Verma, A., Huffman, J. E., Rodriguez, A., Conery, M., Liu, M., Ho, Y.-L., Kim, Y., Heise, D. A., Guare, L., Panickan, V. A., et al. (2023). Diversity and scale: genetic architecture of 2,068 traits in the va million veteran program. $medRxiv$.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge university press.
- Wang, K. (2023). Pseudo-labeling for kernel ridge regression under covariate shift. $arXiv$ preprint arXiv:2302.10160.
- Wang, X., Liu, M., Nogues, I.-E., Chen, T., Xiong, X., Bonzel, C.-L., Zhang, H., Hong, C., Xia, Y., Dahal, K., et al. (2024). Heterogeneous associations between interleukin-6 receptor variants and phenotypes across ancestries and implications for therapy. Scientific Reports, 14(1):8021.
- West, K. M., Blacksher, E., and Burke, W. (2017). Genomics, health disparities, and missed opportunities for the nation's research agenda. JAMA, 317(18):1831–1832.
- Yu, S., Ma, Y., Gronsbell, J., Cai, T., Ananthakrishnan, A. N., Gainer, V. S., Churchill, S. E., Szolovits, P., Murphy, S. N., Kohane, I. S., et al. (2018). Enabling phenotypic big data with PheNorm. Journal of the American Medical Informatics Association, 25(1):54– 60.
- Zhou, D., Liu, M., Li, M., and Cai, T. (2024). Doubly robust augmented model accuracy transfer inference with high dimensional features. Journal of the American Statistical Association, $0(0):1-23$.