

A CHARGE MONOMIAL BASIS OF THE GARSIA-PROCESI RING

MITSUKI HANADA

ABSTRACT. We construct a basis of the Garsia-Procesi ring using the catabolizability type of standard Young tableaux and the charge statistic. This basis turns out to be equal to the descent basis defined in [3]. Our new construction connects the combinatorics of the basis with the well-known combinatorial formula for the modified Hall-Littlewood polynomials $\tilde{H}_\mu[X; q]$, due to Lascoux, which expresses the polynomials as a sum over standard tableaux that satisfy a catabolizability condition. In addition, we prove that identifying a basis for the antisymmetric part of R_μ with respect to a Young subgroup S_γ is equivalent to finding pairs of standard tableaux that satisfy conditions regarding catabolizability and descents. This gives an elementary proof of the fact that the graded Frobenius character of R_μ is given by the catabolizability formula for $\tilde{H}_\mu[X; q]$.

1. INTRODUCTION

The polynomial ring $\mathbb{C}[\mathbf{x}] = \mathbb{C}[x_1, \dots, x_n]$ has a natural S_n action permuting the variables. Taking the quotient of this polynomial ring by the ideal generated by S_n -invariant polynomials with no constant term defines the classical *coinvariant ring*:

$$R_{1^n} = \mathbb{C}[\mathbf{x}] / \langle e_1(\mathbf{x}), \dots, e_n(\mathbf{x}) \rangle,$$

where $e_d(\mathbf{x})$ is the d th elementary symmetric function. The coinvariant ring R_{1^n} is a graded version of the regular representation of S_n . As a graded algebra with an S_n action, it is isomorphic to the cohomology ring of the complex flag variety \mathcal{F}_n . There are two well-known monomial bases of R_{1^n} , the *Artin basis* and the *Garsia-Stanton descent basis*, both with elements indexed by permutations in S_n . In this paper, we will focus on subsets of the latter basis.

Certain quotients R_μ of R_{1^n} correspond to $\mathbb{C}S_n$ -modules that come from Springer fibers. Given a partition μ of n , the Springer fiber $\mathcal{F}_\mu \subset \mathcal{F}_n$ is the set of flags that are fixed by a unipotent matrix with Jordan type μ . The cohomology ring $H^*(\mathcal{F}_\mu)$ of the Springer fiber has an S_n action which is compatible with that on $H^*(\mathcal{F}_n)$ [12].

Though the definitions above can be extended to different Lie types, there are certain nice properties when we consider Type A. In particular, the map $H^*(\mathcal{F}_n) \rightarrow H^*(\mathcal{F}_\mu)$ is a surjection. From this, we have a concrete realization of $H^*(\mathcal{F}_\mu)$ as a quotient of R_{1^n} by an S_n invariant ideal. The following presentation is due to De Concini-Procesi [4] and Tanisaki [14]:

$$R_\mu = \mathbb{C}[\mathbf{x}] / \langle e_d(S) \mid S \subset \mathbf{x}, d > |S| - p_{n-|S|}(\mu) \rangle$$

where $p_k(\mu)$ is the number of boxes of the Young diagram of μ that are not in the first k columns. Garsia and Procesi [6] constructed a monomial basis of R_μ that is a subset

of the Artin monomials. In light of their work, we refer to R_μ as the *Garsia-Procesi ring* indexed by μ . The question of finding a subset of the Garsia-Stanton descent monomials that is a monomial basis of R_μ was unresolved until recently, when Carlsson and Chou gave a construction using shuffles of descent compositions [3].

The graded Frobenius character (see Section 2.3 for definition) of the Garsia-Procesi ring R_μ is given by the modified Hall-Littlewood polynomials $\tilde{H}_\mu[X; q]$ [7]. These polynomials have a combinatorial formula due to Lascoux [8] and Butler [2]:

$$(1.1) \quad \tilde{H}_\mu[X; q] = \sum_{\substack{T \in \text{SYT}_n \\ \text{ctype}(T) \succeq \mu}} q^{\text{cocharge}(T)} s_{\text{shape}(T)}(X).$$

The sum is over standard Young tableaux of size n that satisfy a condition related to an operation called *catabolism* (see Section 2.5 for definition). Catabolism, which was defined by Lascoux [8], is an operation on tableaux, or words, that lowers the cocharge value. Using catabolism, we can associate to each tableau T a partition called its *catabolizability type*, denoted $\text{ctype}(T)$. The sum in Equation (1.1) is over all standard tableaux T that have $\text{ctype}(T) \succeq \mu$, where \succeq denotes the dominance order on partitions.

In this paper, we give a monomial basis of R_μ , using standard tableaux of a certain catabolizability type. The set we construct is the natural subset of the descent basis to consider in light of the combinatorial formula (1.1) for the graded Frobenius series.

For any permutation w , we associate a word $c(w)$ of the same length called the *charge word* of w such that sum of the entries of $c(w)$ is $\text{charge}(w)$. We refer to the monomial $\mathbf{x}^{c(w)}$, where the exponents are given by the charge word of w , as the *charge monomial* corresponding to w . For technical reasons, as well as to match the notation of Carlsson-Chou, we use μ to index the monomial basis of R_{μ^t} , where μ^t is the *transpose* of μ (see Section 2.1 for definition).

Theorem A. *Let μ be a partition of n . The set of charge monomials*

$$\{\mathbf{x}^{c(w)} \mid w \in S_n, \text{ctype}(P(w)^t) \succeq \mu^t\}$$

is a monomial basis of R_{μ^t} . In fact, it coincides with the basis given by Carlsson-Chou [3].

Our basis consists of charge monomials corresponding to certain permutations in S_n . We later see that the set of charge monomials for all permutations in S_n is equal to the Garsia-Stanton descent basis of R_{1^n} . Thus our set of monomials is indeed a subset of the descent basis.

Although our basis is equal to that of Carlsson-Chou, our construction of it is different and provides the first direct connection between the structure of the Garsia-Procesi ring and the catabolizability formula for the modified Hall-Littlewood polynomial that gives its graded Frobenius character. It is immediate that our basis is of the correct size in each dimension, a fact that was nontrivial from the construction of Carlsson-Chou. The construction of our

basis is clearly compatible with the Hilbert series of R_{μ^t} that we compute from (1.1):

$$(1.2) \quad \text{Hilb}_q(R_{\mu^t}) = \sum_{\substack{T \in \text{SYT}_n \\ \text{ctype}(T) \supseteq \mu^t}} q^{\text{charge}(T)} |\text{SYT}(\text{shape}(T))|$$

where $\text{SYT}(\lambda)$ denotes the set of standard Young tableaux of shape λ .

Though it is easy to identify the ungraded Frobenius character of R_{μ} directly from its structure, as exhibited in Garsia–Procesi [6], it serves challenging to do the same for the graded character. This often requires heavy geometric machinery ([4]) or further combinatorial properties of the ring and character ([6]). However, our basis construction gives a way to easily identify the graded Frobenius character of R_{μ} as the modified Hall-Littlewood polynomial $\tilde{H}_{\mu}[X; q]$ that only depends on (1.1) and the ungraded Frobenius character. For any Young subgroup $S_{\gamma} \subset S_n$, we define $N_{\gamma} = \sum_{\sigma \in S_{\gamma}} \text{sgn}(\sigma)\sigma$ to be the antisymmetrizing element with respect to S_{γ} . The graded Frobenius character $\text{Frob}_q(R_{\mu^t})$ is determined by $\text{Hilb}_q(N_{\gamma}R_{\mu^t}) = \langle e_{\gamma}, \text{Frob}_q(R_{\mu^t}) \rangle$ for all Young subgroups S_{γ} , where $N_{\gamma}R_{\mu^t}$ is the antisymmetric part of R_{μ^t} with respect to $S_{\gamma} \subset S_n$. We show that $\text{Hilb}_q(N_{\gamma}R_{\mu^t})$ is easily determined using our basis construction and the ungraded Frobenius character of R_{μ^t} , by proving that the natural subset of our basis enumerated by $\langle e_{\gamma}, \text{Frob}_q(R_{\mu^t}) \rangle$ has the property that applying N_{γ} yields a basis for $N_{\gamma}R_{\mu^t}$.

Proposition B. *Let μ be a partition of n and $\gamma = (\gamma_1, \dots, \gamma_l)$ be a composition of n . The set*

$$\{N_{\gamma} \mathbf{x}^{c(w)} \mid w \in S_n, \text{ctype}(P(w)^t) \supseteq \mu^t, \text{Des}(Q(w)) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}\}$$

is a basis of $N_{\gamma}R_{\mu^t}$, where the pair $(P(w), Q(w))$ is the pair of standard tableaux in bijection with the permutation w via the RSK algorithm and $\text{Des}(Q(w))$ is the descent set of the tableau $Q(w)$.

The following corollary is immediate from Proposition B and the catabolizability formula (1.1).

Corollary C. *For any partition μ of n we have $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}[X; q]$.*

The paper is organized as follows. In Section 2, we review the definitions and combinatorial tools we will use throughout the paper, including RSK insertion, charge, and catabolism. In Section 3 we recall definitions regarding the coinvariant ring and the Garsia-Procesi rings and their monomial bases, as well as recall the construction of the Carlsson-Chou basis from [3]. In Section 4, we give the construction of our set and prove that it agrees with the Carlsson-Chou basis, which shows that our set is indeed a basis due to their results. In particular, we show that our set of monomials is contained in their basis and then conclude they are the same set since they have the same cardinality. In Section 5, we introduce an algorithm that can be used to prove the other direction of containment, as well as other results about catabolizability types. In Section 6, we use a result from the previous section to show that this basis gives us a simple description of the Hilbert series of $N_{\gamma}R_{\mu^t}$. From this, we show $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}[X; q]$.

2. BACKGROUND

2.1. Words and Partitions. For any word $w = w_1 w_2 \dots w_n$ we write $\text{rev}(w) := w_n w_{n-1} \dots w_1$ for the reverse word. A permutation $w \in S_n$ is a word of length n where each element in $\{1, \dots, n\}$ appears exactly once.

The *inversion set* and *descent set* of $w \in S_n$ are denoted

$$\text{Inv}(w) := \{(i, j) \mid i < j, w_i > w_j\},$$

$$\text{Des}(w) := \{i \mid w_i > w_{i+1}\}.$$

The statistics *inversion number* (inv) and *major index* (maj) are defined to be

$$\text{inv}(w) = |\text{Inv}(w)|,$$

$$\text{maj}(w) = \sum_{i \in \text{Des}(w)} i.$$

Both of these statistics are Mahonian, meaning that they are distributed in the following way:

$$\sum_{w \in S_n} q^{\text{inv}(w)} = \sum_{w \in S_n} q^{\text{maj}(w)} = [n]_q!,$$

where $[n]_q! = [1]_q [2]_q \dots [n]_q$ with $[k]_q = 1 + q + \dots + q^{k-1}$.

A *partition* λ of n , denoted $\lambda \vdash n$, is given by $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l > 0$ and $\lambda_1 + \dots + \lambda_l = n$. A *composition* γ of n , denoted $\gamma \models n$, is a sequence $(\gamma_1, \gamma_2, \dots, \gamma_l)$ where $\gamma_i > 0$ and $\gamma_1 + \gamma_2 + \dots + \gamma_l = n$.

The *Young diagram* of a partition $\lambda \vdash n$ is the partial grid of n boxes, where we have λ_i squares in the i th row. We use French notation, meaning that the first part corresponds to the bottom row. For example, the partition $(3, 1)$ corresponds to the Young diagram



The *transpose* of a partition λ , denoted λ^t , is the partition we get by switching the rows and columns of its Young diagram.

There is a partial ordering on the set of all the partitions of size n called the *dominance* ordering, denoted \trianglerighteq , defined by

$$\mu \trianglerighteq \lambda \Leftrightarrow \mu_1 + \dots + \mu_k \geq \lambda_1 + \dots + \lambda_k \text{ for all } k$$

It is well known that $\mu \trianglerighteq \lambda \Leftrightarrow \mu^t \trianglelefteq \lambda^t$. If we move a box of λ to a lower row so that the resulting shape μ is a partition, we have $\mu \trianglerighteq \lambda$. Dominance is the transitive closure of moving boxes down.

A *semistandard (Young) tableau* of shape λ is a filling of λ with positive integers such that the rows are weakly increasing from left to right and the columns are strictly increasing from bottom to top. The *weight* of a semistandard tableau T is the tuple (m_1, m_2, \dots) where m_i is the number of times i appears in T . We denote the shape of T by $\text{shape}(T)$.

A *standard (Young) tableau* is a semistandard tableau where each entry in $\{1, 2, \dots, n\}$ appears in the filling exactly once: that is, it is a semistandard tableau of weight (1^n) . For

any partition λ , we denote the set of standard Young tableaux of shape λ by $\text{SYT}(\lambda)$. We denote the set of standard Young tableaux of all shapes of size n by SYT_n .

The reading word $\text{rw}(T)$ of a tableau T is the word we get by concatenating the row words from top to bottom. The transpose of a tableau T with shape λ , denoted T^t , is the filling of λ^t we get by swapping the rows and columns of T .

2.2. RSK. The RSK (Robinson-Schensted-Knuth) correspondence gives a bijection from permutations $w \in S_n$ to pairs $(P(w), Q(w))$ of standard Young tableau of size n of the same shape. We say $P(w)$ is the *insertion tableau* and $Q(w)$ is the *recording tableau* of w . We note that for any given standard tableau T of shape λ , the number of permutations w with $P(w) = T$ is $|\text{SYT}(\text{shape}(T))|$, since we have one such permutation for each possible choice of $Q(w)$.

We recall some properties of RSK that we will use in following sections. For more details, see [13, Chapter 7]. For a standard Young tableau T , we define the descent set of T to be $\text{Des}(T) = \{i \mid i \text{ appears below } i+1 \text{ in } T\}$. Note that $\text{Des}(T) = \text{Des}(\text{rw}(T)^{-1})$.

Proposition 2.1 ([13, Lemma 7.23.11]). For any $w \in S_n$, we have $\text{Des}(w) = \text{Des}(Q(w))$.

Proposition 2.2 ([13, Corollary A1.2.11]). For any $w \in S_n$, we have $P(\text{rev}(w)) = (P(w))^t$.

2.3. Symmetric functions. For more detailed references on symmetric functions or the representation theory of the symmetric group, see [13, Chapter 7] or [10]. Let Λ denote the ring of symmetric functions in variables $X = (x_1, x_2, \dots)$. We follow Macdonald's notation [10] for the monomial symmetric functions $\{m_\lambda\}$, the elementary symmetric functions $\{e_\lambda\}$, the complete (homogeneous) symmetric functions $\{h_\lambda\}$, and the Schur functions $\{s_\lambda\}$.

The Schur functions s_λ can be represented with respect to the monomial symmetric functions:

$$s_\lambda = \sum_{\gamma \vdash |\lambda|} K_{\lambda, \gamma} m_\gamma$$

where the coefficient $K_{\lambda, \gamma}$, which we call the *Kostka number*, counts the number of semistandard Young tableaux of shape λ and weight γ . These coefficients can also be expressed in terms of standard Young tableaux:

$$(2.1) \quad K_{\lambda, \gamma} = |\{T \in \text{SYT}(\lambda) \mid \text{Des}(T) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}\}|$$

where $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_l)$. A symmetric function f is *Schur-positive* if all the coefficients in the Schur expansion of f are nonnegative integers.

The *Hall inner product* is the inner product defined on Λ by the relation $\langle m_\lambda, h_\gamma \rangle = \delta_{\lambda, \gamma}$. We have $\langle f, g \rangle = \langle g, f \rangle$ for any $f, g \in \Lambda$. The Schur functions are an orthonormal basis with respect to this inner product. The involution ω on Λ is defined by $\omega e_\lambda = h_\lambda$. This map ω is an isometry: that is, we have $\langle f, g \rangle = \langle \omega f, \omega g \rangle$ for any $f, g \in \Lambda$. We also have $\omega s_\lambda = s_{\lambda^t}$ for any λ .

Recall that the irreducible representations of S_n are indexed by partitions of size n . The *Frobenius characteristic map* is a map F from (virtual) character of S_n to symmetric functions of degree n that takes the irreducible character indexed by $\lambda \vdash n$ to the Schur

function s_λ . This map encodes the character of an S_n representation V into a Schur-positive symmetric function. The *Frobenius character* of V , denoted $\text{Frob}(V)$, is defined to be $F(\chi_V)$, where χ_V is the character of V . We have $\text{Frob}(V_\lambda) = F(\chi_{V_\lambda}) = s_\lambda$, where V_λ is the irreducible S_n representation indexed by λ .

For a graded vector space $V = \bigoplus_{d \geq 0} V_d$, the *Hilbert series* $\text{Hilb}_q(V)$ is

$$\text{Hilb}_q(V) = \sum_{d \geq 0} q^d \dim(V_d).$$

For a graded $\mathbb{C}S_n$ -module $V = \bigoplus_{d \geq 0} V_d$, the *graded Frobenius character* $\text{Frob}_q(V)$ is

$$\text{Frob}_q(V) = \sum_{d \geq 0} q^d \text{Frob}(V_d).$$

For the Young subgroup $S_\gamma = S_{\gamma_1} \times \cdots \times S_{\gamma_l} \subset S_n$ corresponding to $\gamma \models n$, we define $N_\gamma = \sum_{\sigma \in S_\gamma} \text{sgn}(\sigma)\sigma$ to be the antisymmetrizer with respect to γ . For any $\mathbb{C}S_n$ -module V , the vector space $N_\gamma V$ is the subspace of elements of V that are antisymmetric with respect to S_γ . Let $\mathcal{E} \uparrow_{S_\gamma}^{S_n}$ denote the induction of the sign representation \mathcal{E} of S_γ to S_n . It is well known that $\text{Frob}(\mathcal{E} \uparrow_{S_\gamma}^{S_n}) = e_\gamma$. Using Frobenius reciprocity (see [5, Chapter 3.3]), we get the following result:

Proposition 2.3. For any $\mathbb{C}S_n$ -module V and Young subgroup $S_\gamma \subset S_n$, we have

$$\dim(N_\gamma V) = \langle e_\gamma, \text{Frob}(V) \rangle.$$

The analogous statement for graded modules holds as well:

Proposition 2.4. For any graded $\mathbb{C}S_n$ -module $V = \bigoplus_{d \geq 0} V_d$ and Young subgroup $S_\gamma \subset S_n$, we have

$$\text{Hilb}_q(N_\gamma V) = \langle e_\gamma, \text{Frob}_q(V) \rangle.$$

2.4. Cocharge and Charge. The *transformed Hall-Littlewood polynomials* $H_\mu[X; q]$ are defined to be

$$H_\mu[X; q] = \sum K_{\lambda, \mu}(q) s_\lambda$$

where $K_{\lambda, \mu}(q)$ is the q -Kostka polynomial (see [10, Chapter III.6] for definition). There is a combinatorial formula for $K_{\lambda, \mu}(q)$ due to Lascoux-Schützenberger [9] using a statistic defined on semistandard Young tableaux called *charge*:

$$K_{\lambda, \mu}(q) = \sum_{T \in \text{SSYT}(\lambda, \mu)} q^{\text{charge}(T)}.$$

The sum is over all semistandard tableaux of shape λ and weight μ . In particular, $K_{\lambda, \mu}(1) = K_{\lambda, \mu}$. From this, we see that $H_\mu[X; 1]$ is equal to the complete homogeneous symmetric function h_μ .

The *modified Hall-Littlewood polynomials* $\tilde{H}_\mu[X; q]$ are defined by:

$$\tilde{H}_\mu[X; q] = q^{n(\mu)} H_\mu[X; q^{-1}]$$

where $n(\mu) = \sum(i - 1)\mu_i$. Then we can write

$$\tilde{H}_\mu[X; q] = \sum_{\lambda} \tilde{K}_{\lambda, \mu}(q) s_{\lambda}(X)$$

when $\tilde{K}_{\lambda, \mu}(q) := q^{n(\mu)} K_{\lambda, \mu}(q^{-1})$ is the modified q -Kostka polynomial. The polynomial $\tilde{K}_{\lambda, \mu}(q)$ can be expressed by the statistic *cocharge* :

$$\tilde{K}_{\lambda, \mu}(q) = \sum_{T \in \text{SSYT}(\lambda, \mu)} q^{\text{cocharge}(T)}$$

where for a semistandard tableau of weight μ we have $\text{cocharge}(T) = n(\mu) - \text{charge}(T)$.

We now define charge and cocharge on permutations. For $w \in S_n$, we define $\text{charge}(w) = \text{maj}(\text{rev}(w^{-1}))$ and $\text{cocharge}(w) = \binom{n}{2} - \text{charge}(w)$. Explicitly, we can compute cocharge of w by assigning labels to each letter in w . The *cocharge word* of w (denoted $\text{cc}(w)$) is a word of length n that gives us the cocharge labelling of w , which we define in the following way. We label 1 of w with 0. We proceed by reading the numbers in increasing order: if we label i with a k , then we label $(i + 1)$ with a k if it is to the right of i . We label $(i + 1)$ with a $(k + 1)$ if it is to the left of i . The statistic *cocharge*(w) is the sum of the letters in $\text{cc}(w)$.

Example 2.5. Let $w = 3 \ 5 \ 1 \ 6 \ 2 \ 4 \ 7$. The corresponding cocharge word is $\text{cc}(w) = 1 \ 2 \ 0 \ 2 \ 0 \ 1 \ 2$, hence $\text{cocharge}(w) = 1 + 2 + 0 + 2 + 0 + 1 + 2 = 8$.

We have the following classification of cocharge words of permutations.

Proposition 2.6. Let z be a word of nonnegative integers of length n containing a 0. We have $z = \text{cc}(w)$ for some $w \in S_n$ if and only if for all $i \in \{1, 2, \dots, n\}$ (at least) one of the following holds:

- (1) There exists $i' > i$ such that $z_i = z_{i'}$,
- (2) There exists $i' < i$ such that $z_i + 1 = z_{i'}$,
- (3) There is no entry in z larger than z_i .

Proof. If $z = \text{cc}(w)$, one of the three conditions will hold for any index $1 \leq i \leq n$. The first (resp. second) case corresponds to when $w_i + 1$ appears to the right (resp. left) of w_i . The last case corresponds to when w_i has the highest cocharge value in the word.

If we have such z we can easily recover $w \in S_n$ such that $\text{cc}(w) = z$. For any nonnegative integer k , let c_k denote the number of times k appears in z and $\{k_1 < k_2 < \dots < k_{c_k}\} := \{i \mid z_i = k\}$. From condition (2), if $c_m > 0$ but $c_{m+1} = 0$ for some nonnegative integer m , it follows that z consists of $\{0, 1, \dots, m\}$ and each of these letters appears at least once.

We construct w so that for each k such that $c_k \neq 0$, the subword $w_{k_1} \dots w_{k_{c_k}}$ is the increasing sequence

$$(c_0 + \dots + c_{k-1} + 1)(c_0 + \dots + c_{k-1} + 2) \dots (c_0 + \dots + c_{k-1} + c_k).$$

The resulting word w is a permutation by construction. It is a simple exercise to check that $\text{cc}(w) = z$. □

Lemma 2.7. *Let $w \in S_n$, x be a positive integer, and $1 \leq k \leq n$. We define $z^{(k)}$ to be a word of length $n + 1$ such that*

$$z_i^{(k)} = \begin{cases} \text{cc}(w)_i & 1 \leq i < k \\ x & i = k \\ \text{cc}(w)_{i-1} & k < i \leq n + 1. \end{cases}$$

- (i) *If $x = \text{cc}(w)_i$ for some i , then $z^{(k)}$ is a cocharge word for any k .*
- (ii) *Consider m such that $\text{cc}(w)_m$ is the rightmost, largest entry of $\text{cc}(w)$. If $x = \text{cc}(w)_m + 1$ and $k < m$, the word $z^{(k)}$ is a cocharge word.*

Proof. Both cases follow from Proposition 2.6. We know each entry in $\text{cc}(w)$ satisfies the condition in Proposition 2.6. We can also see that inserting a new letter into this word does not affect whether preexisting entries satisfy conditions (1) or (2), since it does not change the relative order of those entries within $z^{(k)}$. Hence it suffices to check that the new letter x and the letters in $\text{cc}(w)$ that only satisfied (3) still satisfy one of the three conditions when looking at $z^{(k)}$. Consider m such that the letter $\text{cc}(w)_m$ only satisfies (3). There is only one such letter: the rightmost, largest entry in $\text{cc}(w)$.

For (i), if $x = \text{cc}(w)_i$ for some i , it follows that $x \leq \text{cc}(w)_m$, thus $\text{cc}(w)_m$ is still the largest value in the word. Hence $\text{cc}(w)_m$ satisfies (1) (if $x = \text{cc}(w)_m$ and $m < k$) or (3) (otherwise). We can also see that x clearly satisfies either (1) or (2).

For (ii), we know $x = \text{cc}(w)_m + 1$ and $k < m$. Hence $\text{cc}(w)_m$ satisfies (2) and x satisfies (3). \square

We define the *charge* word $c(w)$ of permutation w to be the following:

$$(2.2) \quad c(w) = \text{rev}(\text{cc}(\text{rev}(w))).$$

We can see that $c(w)$ can be computed by the same algorithm we use to compute the cocharge word, except with left and right exchanged. The statistic $\text{charge}(w)$ is the sum of the letters in $c(w)$.

Lascoux-Schützenberger [9] proved that charge is the unique statistic on words that satisfies a set of properties, one of them being the following:

Theorem 2.8 (Lascoux-Schützenberger [9]). *If $w, w' \in S_n$ are such that $P(w) = P(w')$, then $\text{charge}(w) = \text{charge}(w')$.*

Note that the original statement is for words with partition weight that are Knuth equivalent; for our purposes, it suffices to consider the case above. From this, we can define charge (resp. cocharge) on standard Young tableau T by defining it on words and then extending the definition to tableaux by setting $\text{charge}(T) = \text{charge}(\text{rw}(T))$ (resp. $\text{cocharge}(T) = \text{cocharge}(\text{rw}(T))$).

One observation we can make from Proposition 2.2 and (2.2) is the following:

Proposition 2.9. *If T is a standard tableau, then $\text{charge}(T) = \text{cocharge}(T^t)$.*

To each $w \in S_n$, we associate a monomial $\mathbf{x}^{c(w)}$, where the exponents are given by the charge word of w . We refer to this monomial as the *charge monomial* of w .

Example 2.10. Consider $w' = 7 \ 4 \ 2 \ 6 \ 1 \ 5 \ 3$. Note that $w' = \text{rev}(w)$ from Example 2.5. From (2.2), we know $c(w') = \text{rev}(cc(w)) = 2 \ 1 \ 0 \ 2 \ 0 \ 2 \ 1$. Thus the corresponding charge monomial is $\mathbf{x}^{c(w')} = x_1^2 x_2 x_4^2 x_6^2 x_7$.

2.5. Catabolism and catabolizability. Let T be a standard tableau with $\text{rw}(T) = ww'$ where w' is the first row of T . *Catabolism* is defined to be the operation that takes $T = P(ww')$ to $K(T) := P(w'w)$. It is known that $\text{cocharge}(K(T)) = \text{cocharge}(T) - (|\lambda| - \lambda_1)$ where $\lambda = \text{shape}(T)$. Repeated applications of catabolism eventually produce a one row tableau with zero cocharge.

We can see that applying catabolism to T is equivalent to sliding the first row of T to the right until it detaches from the higher rows, and then swapping the two pieces and applying jeu-de-taquin (for a definition of jeu-de-taquin, see [13, Chapter 7.A1.2]) to the resulting skew shape until we get a partition shape tableau. This is illustrated in the example below.

Example 2.11. Consider the following tableau of shape $\lambda = (3, 2, 1)$:

$$T = \begin{array}{|c|c|c|} \hline 6 & & \\ \hline 2 & 5 & \\ \hline 1 & 3 & 4 \\ \hline \end{array}.$$

We can see that $\text{rw}(T) = 625134$ where $w = 625$ and $w' = 134$. Observe that

$$K(T) = P(w'w) = P(134625) = \begin{array}{|c|c|c|c|c|} \hline 3 & 6 & & & \\ \hline 1 & 2 & 4 & 5 & \\ \hline \end{array}.$$

Alternatively, we can use jeu-de-taquin to see

$$T = \begin{array}{|c|c|c|} \hline 6 & & \\ \hline 2 & 5 & \\ \hline 1 & 3 & 4 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline & 6 & \\ \hline & 2 & 5 \\ \hline \end{array} \xrightarrow{\text{jeu-de-taquin}} \begin{array}{|c|c|c|c|c|} \hline 3 & 6 & & & \\ \hline 1 & 2 & 4 & 5 & \\ \hline \end{array} = K(T).$$

Note that $cc(T) = 8$ and $cc(K(T)) = 5 = 8 - (6 - 3)$, where $|\lambda| = 6, \lambda_1 = 3$.

We use catabolism to define catabolizability type. For a standard tableau T , let $d(T)$ be the largest integer m such that the first row of T contains $1, 2, \dots, m$. The *catabolizability type* of T (denoted $\text{ctype}(T)$) is given by the sequence

$$(2.3) \quad \text{ctype}(T) = (d(T), d(K(T)) - d(T), \dots, d(K^i(T)) - d(K^{i-1}(T)), \dots).$$

Example 2.12. Consider $T = \begin{array}{|c|c|c|c|} \hline 6 & 8 & & \\ \hline 4 & 5 & & \\ \hline 1 & 2 & 3 & 7 \\ \hline \end{array}$. Since $d(T) = 3$, we have $\text{ctype}(T)_1 = 3$. The result of applying catabolism is

$$K(T) = P(12376845) = \begin{array}{|c|c|c|c|c|} \hline 7 & & & & \\ \hline 6 & 8 & & & \\ \hline 1 & 2 & 3 & 4 & 5 \\ \hline \end{array}.$$

Since $d(K(T)) = 5$, we have $\text{ctype}(T)_2 = d(K(T)) - d(T) = 2$. Repeating this process, we can compute $\text{ctype}(T) = (3, 2, 1, 1, 1)$.

We can think of $\text{ctype}(T)$ as encoding the sequence of catabolisms we use to go from T to the one row tableau. We have the following result regarding this sequence.

Proposition 2.13 (Lascoux [8, Lemma 9.6]). For standard tableau T , the sequence $\text{ctype}(T)$ is a partition.

An alternative way to view $\text{ctype}(T)$ is using m -catabolizability (see [11] for details). Let (m) denote the one row shape of length m . If $(m) \subset \text{shape}(T)$ and the first row of T contains the smallest m letters in T , we define the m -catabolism of T to be deleting the first m entries of the first row of T and then proceeding as we did for standard catabolism, where we denote the resulting tableau $\text{Cat}_m(T)$. We say that T of size n is λ -catabolizable for partition $\lambda = (\lambda_1, \hat{\lambda}) \vdash n$ if the first row of T contains the smallest λ_1 many entries of T and $\text{Cat}_{\lambda_1}(T)$ is $\hat{\lambda}$ -catabolizable. Catabolizability type of a tableau and λ -catabolizability are related in the following way.

Proposition 2.14 (Shimozono-Weyman [11, Proposition 46]). A standard tableau T is λ -catabolizable if and only if $\text{ctype}(T) \supseteq \lambda$.

From this, we can compute $\text{ctype}(T)$ by repeatedly applying maximal m -catabolisms, where at each step we have

$$\text{ctype}(T)_i = d(\text{Cat}_{\text{ctype}(T)_{i-1}} \cdots \text{Cat}_{\text{ctype}(T)_1}(T)).$$

This simplifies our process, since at each step we are dealing with smaller tableaux.

Example 2.15. Consider $T = \begin{array}{|c|c|} \hline 6 & 8 \\ \hline 4 & 5 \\ \hline 1 & 2 & 3 & 7 \\ \hline \end{array}$. We know $d(T) = 3$ and can compute $\text{Cat}_3(T)$:

$$\begin{array}{|c|c|} \hline 6 & 8 \\ \hline 4 & 5 \\ \hline 1 & 2 & 3 & 7 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|} \hline 7 \\ \hline 6 & 8 \\ \hline 4 & 5 \\ \hline \end{array} \xrightarrow{\text{jeu-de-taquin}} \begin{array}{|c|c|} \hline 7 \\ \hline 6 & 8 \\ \hline 4 & 5 \\ \hline \end{array} = \text{Cat}_3(T).$$

We see that $\text{ctype}(T)_2 = d(\text{Cat}_3(T)) = 2$. Repeating this process, we can compute $\text{ctype}(T) = (3, 2, 1, 1, 1)$, which is the same as what we get in Example 2.12.

In general, many things are unknown about catabolism and catabolizability, though the operation of catabolism itself is very easy to compute. However, there is a concrete algorithm, due to Blasiak [1], called the *catabolism insertion algorithm*, which computes the catabolizability type of a permutation w , where we define $\text{ctype}(w) := \text{ctype}(P(w))$.

Algorithm 2.16 (Blasiak [1, Algorithm 3.2]). We define a function f on pairs (x, ν) , where $x = ya$ is a word (a is the last letter of x) and ν is a partition, to be

$$f(x, \nu) = \begin{cases} (y, \nu + \epsilon_{a+1}) & \text{if } \nu + \epsilon_{a+1} \text{ is a partition,} \\ ((a+1)y, \nu) & \text{otherwise.} \end{cases}$$

where ϵ_{a+1} is the composition $(0, \dots, 0, 1, 0, \dots)$ with 1 in the $(a+1)$ th coordinate.

Let w be a permutation of length n with cocharge word $\text{cc}(w)$. We apply f to $(\text{cc}(w), \emptyset)$ repeatedly until we get (\emptyset, μ) , where $\mu \vdash n$.

We introduce terminology regarding this algorithm. When we refer to the i th letter of the input word, we also count the letters that we have deleted from the word. Hence the input word is always of length n , but potentially with empty spots. At each step of the algorithm, when the input is (ya, ν) , we say we *read* i if the letter a corresponds to the i th letter in the original input word $\text{cc}(w)_i$.

Proposition 2.17 (Blasiak [1]). When applying Algorithm 2.16 to $(cc(w), \emptyset)$, the resulting partition μ is $ctype(w)$.

As a slight modification to the original algorithm introduced in [1], we record the entries of $cc(w)$ that correspond to the boxes in μ as we build the partition. When reading i results in adding a box to the partition ν , we fill the new box with i . This gives us a standard filling (but not a SYT) of shape $ctype(w)$, meaning $\{1, \dots, n\}$ appear exactly once. We denote this filling as T_w . For the modified algorithm, the final tuple is (\emptyset, T_w) .

If the initial word z is not a cocharge word, this algorithm may not terminate. For example: if $z = 02$, the second letter will never be deleted, since we cannot add a box to the partition (1) in rows 2 or higher.

Example 2.18. Consider

$$w = 6 \ 3 \ 4 \ 1 \ 2 \ 5$$

$$cc(w) = 2 \ 1 \ 1 \ 0 \ 0 \ 1.$$

We will apply f to $(cc(w), \emptyset)$ repeatedly until we get an empty word in the first coordinate. In order to keep track of the indices, we do not rotate $cc(w)$: instead, we read the word from right to left. The position we are reading at each step of the algorithm is underlined.

$$\begin{array}{c}
 (2 \ 1 \ 1 \ 0 \ 0 \ \underline{1}, \ \emptyset) \\
 \downarrow \\
 (2 \ 1 \ 1 \ 0 \ \underline{0} \ 2, \ \emptyset) \\
 \downarrow \\
 (2 \ 1 \ 1 \ \underline{0} \ 2, \ \boxed{5}) \\
 \downarrow \\
 (2 \ 1 \ \underline{1} \ 2, \ \boxed{5 \ 4}) \\
 \downarrow \\
 (2 \ \underline{1} \ 2, \ \begin{array}{|c|c|} \hline 3 & \\ \hline 5 & 4 \\ \hline \end{array}) \\
 \downarrow \\
 (\underline{2} \ 2, \ \begin{array}{|c|c|} \hline 3 & 2 \\ \hline 5 & 4 \\ \hline \end{array}) \\
 \downarrow \\
 (\ \underline{2}, \ \begin{array}{|c|c|} \hline 1 & \\ \hline 3 & 2 \\ \hline 5 & 4 \\ \hline \end{array}) \\
 \downarrow \\
 (\ \ , \ \begin{array}{|c|c|} \hline 1 & 6 \\ \hline 3 & 2 \\ \hline 5 & 4 \\ \hline \end{array})
 \end{array}$$

From this, we conclude $T_{634125} = \begin{array}{|c|c|} \hline 1 & 6 \\ \hline 3 & 2 \\ \hline 5 & 4 \\ \hline \end{array}$ and $\text{ctype}(634125) = (2, 2, 2)$.

We now state some nice properties of the filling T_w .

Lemma 2.19. *If we add i to row r of T_w when reading i for the k th time, we have*

$$k + \text{cc}(w)_i = r.$$

Proof. Each time we read i without adding it to T_w , we increase the i th letter of the input word by 1. After reading through the word $(k-1)$ times without adding i to the filling, the i th letter in the input word is $\text{cc}(w)_i + (k-1)$. Since we add i the next time we read the i th letter, we know we add it to the $(\text{cc}(w)_i + (k-1) + 1)$ th row, where the last 1 is coming from the fact that $f(x, \nu) = (y, \nu + \epsilon_{a+1})$. \square

We now look at subwords of $\text{cc}(w)$ determined by the columns of T_w . Consider $w \in S_n$ such that $\text{ctype}(w) = \mu^t$. For $1 \leq j \leq (\mu^t)_1$ and $1 \leq r \leq \mu_j$, let

$$\{i \mid i \text{ is in row } r', \text{ column } j \text{ of } T_w, 1 \leq r' \leq r\} = \{j_1^{(r)} < j_2^{(r)} < \dots < j_r^{(r)}\}.$$

We are numbering the entries in column j of T_w in increasing order, though that order may not match the order in which they appear in the column.

We define the subword $\text{cc}(w)^{(j,r)}$ of $\text{cc}(w)$ to be

$$\text{cc}(w)^{(j,r)} = \text{cc}(w)_{j_1^{(r)}} \text{cc}(w)_{j_2^{(r)}} \dots \text{cc}(w)_{j_r^{(r)}}.$$

In this case, $\text{cc}(w)^{(j,r)}$ is the subword corresponding to the first r rows of column j in T_w . We write $\text{cc}(w)^{(j)} = \text{cc}(w)^{(j, \mu_j)}$ when r is equal to the height of column j .

Example 2.20. Using $w = 634125$ from Example 2.18, we can see that

$$\text{cc}(w)^{(1,3)} = \text{cc}(w)_1 \text{cc}(w)_3 \text{cc}(w)_5 = 210.$$

We can also see that $j_1^{(3)} = 1$ but 1 is not in the first row of T_w .

Proposition 2.21. For $w \in S_n$, consider j, r such that $1 \leq j \leq (\mu^t)_1$ and $1 \leq r \leq \mu_j$. Then $\text{cc}(w)^{(j,r)} = \text{cc}(\sigma)$ for some $\sigma \in S_r$.

Proof. We show this by induction on r . When $r = 1$, we know $\text{cc}(w)^{(j,1)} = 0$, since for any i that appears in the first row of T_w we must have $\text{cc}(w)_i = 0$. Hence the claim holds.

Assume the claim holds for $(r-1)$. We know that we obtain $\text{cc}(w)^{(j,r)}$ from $\text{cc}(w)^{(j,r-1)}$ by adding $\text{cc}(w)_a$ into $\text{cc}(w)^{(j,r-1)}$ in the correct position, where a is the entry in row r , column j of T_w . Let b denote the entry directly below a in T_w . Since b appears below a , we know that when we constructed T_w using Algorithm 2.16 to $\text{cc}(w)$ we first added b , and then a , to the filling. If we let k_a, k_b denote the number of times we read a, b before adding them to T_w , we must have $k_b \leq k_a$. If $k_a = k_b$, then since we add b before a , we must have $a < b$, since $\text{cc}(w)_b$ must be to the right of $\text{cc}(w)_a$ in $\text{cc}(w)$. We also have $\text{cc}(w)_a = \text{cc}(w)_b + 1$ from Lemma 2.19 since a is one row above b . If $k_b < k_a$, then $\text{cc}(w)_b \geq \text{cc}(w)_a$.

From this, we know that if $cc(w)_a$ is larger than every letter in $cc(w)^{(j,r-1)}$, we have $cc(w)_a = cc(w)_b + 1$ where $b > a$ is the entry directly below a in T_w and $cc(w)_b$ is the largest letter in $cc(w)^{(j,r-1)}$.

By Lemma 2.7, we conclude $cc(w)^{(j,r)}$ is a cocharge word by induction. \square

Remark 2.22. The fact that $cc(w)^{(j,r)}$ is a cocharge word is not obvious a priori, since it is not true that any subword of a cocharge word is a cocharge word. For example, consider $w = 21$, which has $cc(w) = 10$. Though 1 is a subword of $cc(w)$, it is not a cocharge word.

3. COINVARIANT RING AND GARSIA-PROCESI RING

The *coinvariant ring* R_{1^n} is defined to be $R_{1^n} = \mathbb{C}[\mathbf{x}]/I$, where I is the ideal generated by S_n invariant polynomials with zero constant term. More specifically, we can write $I = \langle e_1(\mathbf{x}), \dots, e_n(\mathbf{x}) \rangle$. It is clear that R_{1^n} is a $\mathbb{C}S_n$ -module under the action permuting the variables. It is well-known that this is a graded version of the regular representation of S_n : in particular, $\dim(R_{1^n}) = n!$. As a graded algebra with S_n action, it is isomorphic to $H^*(\mathcal{F}_n)$, where \mathcal{F}_n is the complete flag variety for GL_n . In particular, we have $\text{Hilb}_q(R_{1^n}) = [n]_q!$.

In Type A , we have that $H^*(\mathcal{F}_\mu)$, the cohomology ring of the Springer fiber indexed by $\mu \vdash n$, is a quotient of $H^*(\mathcal{F}_n)$. These cohomology rings also have an S_n action: we have the following concrete presentation of these $\mathbb{C}S_n$ -modules as quotients of R_{1^n} due to De Concini-Procesi [4] and Tanisaki [14]. Consider the transpose partition $\mu^t = (\mu_1^t \geq \mu_2^t \geq \dots \geq \mu_n^t \geq 0)$, where we pad the end of the partition with 0's until we have a tuple of length n . Let $p_k(\mu) := \mu_n^t + \dots + \mu_{n-k+1}^t$. Note that $p_k(\mu)$ is the number of boxes of the Young diagram of μ that are not in the first $n - k$ columns. Given a subset $S \subset \mathbf{x}$, we define $e_d(S)$ to be the d th elementary symmetric function in the set of variables S . The *Tanisaki ideal* I_μ is defined to be

$$I_\mu = \langle e_d(S) \mid S \subset \mathbf{x}, d > |S| - p_{n-|S|}(\mu) \rangle.$$

The *Garsia-Procesi ring* R_μ is the quotient $R_\mu = \mathbb{C}[\mathbf{x}]/I_\mu$. As a graded $\mathbb{C}S_n$ -module, it is isomorphic to $H^*(\mathcal{F}_\mu)$. From this, we know that the graded Frobenius character of R_μ is the modified Hall-Littlewood polynomial $\tilde{H}_\mu[\mathbf{x}; q]$. Using the combinatorial formula (1.1) for $\tilde{H}_\mu[\mathbf{x}; q]$, we compute $\text{Hilb}_q(R_\mu)$:

$$\begin{aligned} \text{Hilb}_q(R_\mu) &= \sum_{\substack{T \in \text{SYT} \\ \text{ctype}(T) \geq \mu}} q^{\text{cocharge}(T)} \dim(V^{\text{shape}(T)}) \\ &= \sum_{\substack{T \in \text{SYT} \\ \text{ctype}(T) \geq \mu}} q^{\text{cocharge}(T) + |\text{SYT}(\text{shape}(T))|} \\ (3.1) \quad &= \sum_{\substack{T \in \text{SYT} \\ \text{ctype}(P(w)) \geq \mu}} q^{\text{cocharge}(w)}. \end{aligned}$$

When $\mu = 1^n$, we have $I_{1^n} = \langle e_1(\mathbf{x}), \dots, e_n(\mathbf{x}) \rangle$, hence the Garsia-Procesi ring indexed by 1^n is the full coinvariant ring R_{1^n} .

3.1. Monomial bases of R_{1^n} and generalizations to R_μ . We recall two monomial bases of the coinvariant ring, both of which are indexed by permutations in S_n .

The *Artin basis* is given by the following set of monomials:

$$\{f_\sigma(\mathbf{x}) = \prod_{i < j, \sigma_i > \sigma_j} x_{\sigma_i} \mid \sigma \in S_n\} = \{x_1^{a_1} x_2^{a_2} \cdots x_n^{a_n} \mid a_i < i\}.$$

From the second description of the Artin monomials, we can easily see that

$$\sum_{w \in S_n} q^{\deg(f_w(\mathbf{x}))} = (1+q)(1+q+q^2) \cdots (1+q+q^2+\cdots+q^{n-1}) = [n]_q!$$

The *Garsia-Stanton descent basis* is given by the following set of monomials:

$$\{g_w(\mathbf{x}) = \prod_{i: w_i > w_{i+1}} x_{w_1} \cdots x_{w_i} \mid w \in S_n\}.$$

Let $D_n = \{\alpha : \mathbf{x}^\alpha = g_w(\mathbf{x}) \mid w \in S_n\}$ denote the set of exponents of monomials in the descent basis. We refer to these words as *descent words*.

Lemma 3.1. *We have $D_n = \{c(w) \mid w \in S_n\}$.*

Proof. For any $\sigma \in S_n$, the charge monomial is

$$\mathbf{x}^{c(\sigma)} = \prod_{j: (\sigma^{-1})_j < (\sigma^{-1})_{j+1}} x_{(\sigma^{-1})_{j+1}} \cdots x_{(\sigma^{-1})_n}.$$

Thus $\mathbf{x}^{c(\sigma)} = g_w(\mathbf{x})$ where $w = \text{rev}(\sigma^{-1})$. □

The *descent order* on monomials is defined by:

$$(3.2) \quad \mathbf{x}^\alpha \leq_{\text{des}} \mathbf{x}^\beta \Leftrightarrow \begin{cases} \text{sort}(\alpha) \preceq \text{sort}(\beta) \\ \text{sort}(\alpha) = \text{sort}(\beta) \text{ and } \alpha \preceq \beta. \end{cases}$$

where $\text{sort}(\alpha)$ reorders the vector α into a partition and \preceq is the lexicographic ordering on words.

The descent monomials correspond to the permutation statistic maj . That is, we have $\deg(g_w(\mathbf{x})) = \text{maj}(w)$. Since this statistic is Mahonian, we have:

$$\sum_{w \in S_n} q^{\deg(g_w(\mathbf{x}))} = \sum_{w \in S_n} q^{\text{maj}(w)} = [n]_q!.$$

Thus, either basis give a combinatorial explanation of $\text{Hilb}_q(R_{1^n}) = [n]_q!$.

Since R_μ is a quotient of R_{1^n} , the natural question to ask is whether there exists a subset of the Artin monomials or the descent monomials which is monomial basis of R_μ . Garsia-Procesi [6] constructed a monomial basis of R_μ which is a subset of the Artin basis of R_{1^n} . However, this basis does not give an analogous combinatorial explanation of the expression (3.1).

The problem of finding a subset of the descent basis that gives a monomial basis of R_μ was recently solved by Carlsson-Chou [3]. We now review the construction of the Carlsson-Chou descent basis. For two words $z^{(1)}, z^{(2)}$ of length l_1, l_2 we define $\text{Sh}(z^{(1)}, z^{(2)})$ to be the collection of all words u of length $(l_1 + l_2)$ such that $z^{(1)}$ and $z^{(2)}$ are two disjoint subwords of u . We construct such u by interleaving $z^{(1)}$ and $z^{(2)}$. An element u of $\text{Sh}(z^{(1)}, z^{(2)})$ is a *shuffle* of $z^{(1)}$ and $z^{(2)}$. We can extend this definition and let $\text{Sh}(z^{(1)}, \dots, z^{(l)})$ denote the set of all shuffles of $z^{(1)}, \dots, z^{(l)}$. We also define the set of *reverse shuffles* of $z^{(1)}, \dots, z^{(l)}$, denoted $\text{Sh}'(z^{(1)}, \dots, z^{(l)})$, to be equal to $\text{Sh}(\text{rev}(z^{(1)}), \dots, \text{rev}(z^{(l)}))$.

The Carlsson-Chou descent basis of the Garsia-Procesi ring is defined as follows. For $\mu = (\mu_1, \dots, \mu_l) \vdash n$, define \mathcal{D}_μ to be

$$(3.3) \quad \mathcal{D}_\mu = \bigcup_{(z^{(1)}, \dots, z^{(l)})} \text{Sh}(z^{(1)}, \dots, z^{(l)})$$

where $(z^{(1)}, \dots, z^{(l)})$ ranges over all l -tuples in $\mathcal{D}_{\mu_1} \times \dots \times \mathcal{D}_{\mu_l}$.

Theorem 3.2 (Carlsson-Chou [3]). *The set $\mathbf{x}^{\mathcal{D}_\mu} = \{\mathbf{x}^\alpha \mid \alpha \in \mathcal{D}_\mu\}$ is a monomial basis of R_{μ^t} .*

Example 3.3. Consider $\mu = (3, 1)$. Note that

$$\begin{aligned} D_3 &= \{012, 011, 101, 001, 010, 000\}, \\ D_1 &= \{0\}. \end{aligned}$$

Then we have

$$\mathcal{D}_{3,1} = \{0012, 0102, 0120, 0011, 0101, 1001, 1010, 0110, 0001, 0010, 0100, 0000\}.$$

The corresponding set of monomials is

$$\{x_3x_4^2, x_2x_4^2, x_2x_3^2, x_3x_4, x_2x_4, x_1x_4, x_1x_3, x_2x_3, x_4, x_3, x_2, 1\},$$

Carlsson-Chou show that their basis is a subset of the Garsia-Stanton descent basis. However, it is not obvious how to see directly for which $w \in S_n$ we have $g_w(\mathbf{x}) \in \mathbf{x}^{\mathcal{D}_\mu}$ without computing \mathcal{D}_μ . It is also not obvious that $|\mathcal{D}_\mu| = \dim(R_{\mu^t})$, since multiple shuffles can correspond to the same descent word. For example, $0101 \in \text{Sh}(101, \mathbf{0})$ and $0101 \in \text{Sh}(011, \mathbf{0})$ both correspond to the same element in \mathcal{D}_μ .

4. CHARGE MONOMIAL BASIS OF THE GARSIA-PROCESI RING

We now define sets of charge monomials that are monomial bases of Garsia-Procesi rings. Our construction involves charge words of permutations whose insertion tableaux satisfy a catabolizability condition.

Definition 4.1. The set \mathcal{C}_μ is defined to be

$$(4.1) \quad \mathcal{C}_\mu := \{c(w) \mid w \in S_n, \text{ctype}(P(w)^t) \supseteq \mu^t\}.$$

From Lemma 3.1, we know $\mathcal{C}_\mu \subset D_n$.

Theorem A. *The set*

$$(4.2) \quad \mathbf{x}^{\mathcal{C}_\mu} = \{\mathbf{x}^\alpha \mid \alpha \in \mathcal{C}_\mu\}$$

is a monomial basis of R_{μ^t} . In fact, it coincides with the basis $\mathbf{x}^{\mathcal{D}_\mu}$ given by Carlsson-Chou [3], ie., $\mathcal{C}_\mu = \mathcal{D}_\mu$.

Before we prove Theorem A, we point out the connections between this basis and the Hilbert series $\text{Hilb}_q(R_{\mu^t})$. Using combinatorial formula (1.1) for the modified Hall-Littlewood polynomials and Proposition 2.9 gives us the following expression for $\tilde{H}_{\mu^t}[X; q]$.

$$(4.3) \quad \tilde{H}_{\mu^t}[X; q] = \sum_{\substack{T \in \text{SYT}_n \\ \text{ctype}(T^t) \succeq \mu^t}} q^{\text{charge}(T)} s_{\text{shape}(T^t)}(X).$$

From this, we can see that the classification of insertion tableaux that appears in 4.1 is the natural one to consider when looking at subsets of descent monomials. In fact, it is evident from our construction that the degrees of the monomials in our set match what we expect from $\text{Hilb}_q(R_{\mu^t})$. Rephrasing (3.1) for $\tilde{H}_{\mu^t}[X; q]$ using Proposition 2.9, we have

$$(4.4) \quad \text{Hilb}_q(R_{\mu^t}) = \sum_{\substack{w \in S_n \\ \text{ctype}(P(w)^t) \succeq \mu^t}} q^{\text{charge}(w)}.$$

It is clear that our set has the correct number of monomials of each degree to be a basis of R_{μ^t} by construction. It is also apparent that $|\mathcal{C}_\mu| = \dim(R_{\mu^t})$, which was not obvious from the definition of \mathcal{D}_μ . This construction also gives us a direct way to determine for which $w \in S_n$ we have $g_w(\mathbf{x}) \in \mathbf{x}^{\mathcal{C}_\mu}$. In particular, we have $g_w(\mathbf{x}) \in \mathbf{x}^{\mathcal{C}_\mu}$ for $w = \text{rev}(\sigma^{-1})$ where $\text{ctype}(P(\sigma)^t) \succeq \mu^t$.

Example 4.2. Consider $\mu = (3, 1)$. There are five standard tableaux S such that $\text{ctype}(S^t) \succeq (2, 1, 1) = \mu^t$. We list them, along with all words w such that $P(w) = S$ and their charge monomials:

S	$\{w \mid P(w) = S\}$	$\{x^{c(w)} \mid P(w) = S\}$
$\begin{array}{ c c c } \hline 2 & & \\ \hline 1 & 3 & 4 \\ \hline \end{array}$	$\{2134, 2314, 2341\}$	$\{x_3x_4^2, x_2x_4^2, x_2x_3^2\}$
$\begin{array}{ c c } \hline 2 & 4 \\ \hline 1 & 3 \\ \hline \end{array}$	$\{2143, 2413\}$	$\{x_3x_4, x_2x_4\}$
$\begin{array}{ c } \hline 4 \\ \hline 2 \\ \hline 1 & 3 \\ \hline \end{array}$	$\{4213, 4231, 2431\}$	$\{x_1x_4, x_1x_3, x_2x_3\}$
$\begin{array}{ c } \hline 3 \\ \hline 2 \\ \hline 1 & 4 \\ \hline \end{array}$	$\{3214, 3241, 3421\}$	$\{x_4, x_3, x_2\}$
$\begin{array}{ c } \hline 4 \\ \hline 3 \\ \hline 2 \\ \hline 1 \\ \hline \end{array}$	$\{4321\}$	$\{1\}$

Note that all charge monomials for words with the same insertion tableau S have degree equal to $\text{charge}(S)$. The resulting set of charge monomials is

$$\{x_3x_4^2, x_2x_4^2, x_2x_3^2, x_3x_4, x_2x_4, x_1x_4, x_1x_3, x_2x_3, x_4, x_3, x_2, 1\},$$

which is the same as in Example 3.3.

Now we will prove Theorem A by showing $\mathcal{C}_\mu = \mathcal{D}_\mu$, which implies that our set is a basis of R_{μ^t} by the work of Carlsson-Chou. Since we know that \mathcal{D}_μ is a basis of R_{μ^t} and that \mathcal{C}_μ has the correct cardinality to be a basis, it suffices to show that $\mathcal{C}_\mu \subset \mathcal{D}_\mu$. We first make the following observation, which is a corollary of Lemma 3.1 using (2.2):

Corollary 4.3. $D_n = \{\text{rev}(\text{cc}(w)) \mid w \in S_n\}$.

To show that $\mathcal{C}_\mu \subset \mathcal{D}_\mu$, we translate the problem into one about cocharge words to make use of Algorithm 2.16. We first consider the case where $\text{ctype}(w) = \mu^t$.

Proposition 4.4. Let μ be a partition of n of length l and w be a permutation of n with $\text{ctype}(w) = \mu^t$. Then $\text{cc}(w) \in \text{Sh}'(z^{(1)}, \dots, z^{(l)})$ for some $(z^{(1)}, \dots, z^{(l)}) \in D_{\mu_1} \times \dots \times D_{\mu_l}$.

Proof. We obtain a filling T_w of shape $\text{ctype}(w) = \mu^t$ by applying Algorithm 2.16 to $\text{cc}(w)$. It is clear that $\text{cc}(w) \in \text{Sh}(\text{cc}(w)^{(1)}, \text{cc}(w)^{(2)}, \dots, \text{cc}(w)^{(l)})$, where $\text{cc}(w)^{(j)}$ is the subword of $\text{cc}(w)$ corresponding to column j in T_w . We know $\text{cc}(W)^{(j)}$ has length μ_j , which is the height of column j . Thus, from Proposition 2.21, we know $\text{cc}(w)^{(j)} = \text{cc}(\sigma)$ for $\sigma \in S_{\mu_j}$. By Corollary 4.3, this implies $\text{rev}(\text{cc}(w)^{(j)}) \in D_{\mu_j}$. \square

Example 4.5. Recall that in Example 2.18, we showed that for $w = 6\ 3\ 4\ 1\ 2\ 5$, with $\text{cc}(w) = 2\ 1\ 1\ 0\ 0\ 1$, we have $\text{ctype}(w) = (2, 2, 2) = (3, 3)^t$ and that the filling T_w of $(2, 2, 2)$ is

1	6
3	2
5	4

From this, we construct two subwords of lengths 3, 3 respectively:

$$\text{cc}(w)^{(1)} = \text{cc}(w)_1 \text{cc}(w)_3 \text{cc}(w)_5 = 210$$

$$\text{cc}(w)^{(2)} = \text{cc}(w)_2 \text{cc}(w)_4 \text{cc}(w)_6 = 101$$

We can see that $\text{cc}(w)^{(1)} = \text{cc}(321) = \text{rev}(c(123))$ and $\text{cc}(w)^{(2)} = \text{cc}(213) = \text{rev}(c(213))$. Thus $\text{cc}(w) \in \text{Sh}'(c(123), c(213))$, where $(c(123), c(213)) \in D_{(3,3)}$.

We rephrase Proposition 4.4 to match the language of Theorem A using (2.2):

Corollary 4.6. For any $w \in S_n$ such that $\text{ctype}(P(w)^t) = \mu^t$, we have $c(w) \in D_\mu$.

Proof. From (2.2) we know

$$\text{cc}(w) \in \text{Sh}'(z^{(1)}, \dots, z^{(l)}) \Leftrightarrow c(\text{rev}(w)) = \text{rev}(\text{cc}(w)) \in \text{Sh}(z^{(1)}, \dots, z^{(l)}).$$

We also have $\text{ctype}(\text{rev}(w)) = \text{ctype}(P(w)^t)$ using Theorem 2.2. \square

Note that Corollary 4.6 only considers permutations w with $\text{ctype}(P(w))^t = \mu^t$. To prove the inclusion $\mathcal{C}_\mu \subset \mathcal{D}_\mu$, we extend this argument to consider $\text{ctype}(P(w))^t \triangleright \mu^t$.

Proof of Theorem A. We know $|\mathcal{C}_\mu| = |\mathcal{D}_\mu|$, since $|\mathcal{C}_\mu| = \dim(R_{\mu^t})$ and \mathcal{D}_μ is a basis of R_{μ^t} . Hence to show equality of the two sets, we will show $\mathcal{C}_\mu \subset \mathcal{D}_\mu$. From Corollary 4.6, it suffices to consider $w \in S_n$ where $\text{ctype}(P(w)^t) \triangleright \mu^t$. First, consider $w \in S_n$ such that $\text{ctype}(P(w)^t) = \lambda^t \triangleright \mu^t$ where λ and μ differ by moving one box in the Young diagram. We write $\lambda = \mu - \varepsilon_{j_1} + \varepsilon_{j_2}$ (where $j_1 < j_2$). Note that j_1 and j_2 correspond to rows of λ, μ , which mean they are columns of λ^t, μ^t .

Since $\text{ctype}(P(w)^t) = \text{ctype}(\text{rev}(w))$, we consider the filling $T_{\text{rev}(w)}$ of λ^t that we get from Algorithm 2.16. Let a be the entry in the last box of column j_2 . Take the box containing a and append it into the end of column j_1 .

By assumption, we know the resulting shape is μ^t . Denote the resulting filling of shape μ^t by $T_{\text{rev}(w)}^{(\mu^t)}$. We claim that each subword of $\text{cc}(\text{rev}(w))$ that corresponds to a column of $T_{\text{rev}(w)}^{(\mu^t)}$ is still a valid cocharge word. The only columns we need to consider are columns j_1 and j_2 . Since the only modification to column j_2 is taking off the last entry we added, we see that the resulting word is still a valid cocharge word from Proposition 2.21.

Since $j_1 < j_2$ and $T_{\text{rev}(w)}$ is partition shape, there exists an entry b in column j_1 which is in the same row as a in $T_{\text{rev}(w)}$. Since $j_1 < j_2$, we know we first added b , and then a , to the filling $T_{\text{rev}(w)}$. If we let k_a, k_b denote the number of times we read a, b before adding them to T_w , we must have $k_b \leq k_a$. Thus, we know that $\text{cc}(\text{rev}(w))_a \leq \text{cc}(\text{rev}(w))_b$ from Lemma 2.19.

Thus adding a to $\text{cc}(\text{rev}(w))^{(j_1)}$ does not introduce a new cocharge value to this word since $\text{cc}(\text{rev}(w))_b$ is in $\text{cc}(\text{rev}(w))^{(j_1)}$. From case (i) of Lemma 2.7, we conclude the resulting word is a cocharge word of a permutation.

If $\text{ctype}(\text{rev}(w)) \triangleright \mu^t$ differ by moving multiple boxes, we repeat this process until we get a filling of μ^t where each of the columns correspond to valid cocharge words. It is clear that $\text{cc}(\text{rev}(w))$ is a shuffle of these cocharge words. Hence $c(w) \in D_\mu$. From this, we have $\mathcal{C}_\mu = \mathcal{D}_\mu$. Since $\mathbf{x}^{\mathcal{D}_\mu}$ is a basis of R_{μ^t} , we conclude that our set $\mathbf{x}^{\mathcal{C}_\mu}$ is a basis as well. \square

5. PROPERTIES OF CATABOLIZABILITY TYPE

Theorem A says that two conditions, one involving catabolizability type and the other involving shuffles, are equivalent. In particular, we get the following corollary,

Corollary 5.1. Let $u^{(1)}, u^{(2)}, \dots, u^{(l)}$ be permutations where $u^{(j)} \in S_{\mu_j}$. Then we have

$$\text{Sh}(\text{cc}(u^{(1)}), \dots, \text{cc}(u^{(l)})) \subset \{\text{cc}(w) \mid w \in S_n, \text{ctype}(w) \triangleright \mu^t\}.$$

Proof. This follows from rephrasing the relation $\mathcal{D}_\mu \subset \mathcal{C}_\mu$ by reversing the words and using Proposition 2.2 and (2.2). \square

However, we can show the following stronger result using properties of Algorithm 2.16.

Theorem 5.2. Let $u^{(1)}, u^{(2)}, \dots, u^{(l)}, w$ be permutations such that $\text{cc}(w) \in \text{Sh}(\text{cc}(u^{(1)}), \dots, \text{cc}(u^{(l)}))$. We have

$$\text{ctype}(w) \triangleright \text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)}) + \dots + \text{ctype}(u^{(l)}),$$




FIGURE 1

where $\text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)}) + \dots + \text{ctype}(u^{(l)})$ is the partition given by the partwise sum of $\text{ctype}(u^{(1)}), \text{ctype}(u^{(2)}), \dots, \text{ctype}(u^{(l)})$.

We can see that Corollary 5.1 follows from Theorem 5.2. For any $u \in S_n$, we know $\text{ctype}(u) \succeq (1)^n$, since $(1)^n$ is the unique smallest partition of size n with respect to dominance order. Hence, for permutations $u^{(1)}, u^{(2)}, \dots, u^{(l)}, w$ where $u^{(j)} \in S_{\mu_j}$ and $\text{cc}(w) \in \text{Sh}(\text{cc}(u^{(1)}), \dots, \text{cc}(u^{(l)}))$, we have

$$\text{ctype}(w) \succeq \text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)}) + \dots + \text{ctype}(u^{(l)}) \succeq (1)^{\mu_1} + (1)^{\mu_2} + \dots + (1)^{\mu_l} = \mu^t.$$

Hence Theorem 5.2 gives an improved lower bound for the catabolizability type of a shuffle.

We prove Theorem 5.2 by introducing a new modification of Algorithm 2.16 that keeps track of the difference between $\text{ctype}(w)$ and $\text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)})$. We first construct our inputs for the algorithm. We restrict to the case where $l = 2$, but the same arguments generalize to larger l . Let $u^{(1)}, u^{(2)}, w$ be permutations such that $\text{cc}(w) \in \text{Sh}(\text{cc}(u^{(1)}), \text{cc}(u^{(2)}))$. If we specify a shuffle of $\text{cc}(u^{(1)})$ and $\text{cc}(u^{(2)})$ that is equal to $\text{cc}(w)$, we have that each letter in $\text{cc}(w)$ corresponds uniquely to a letter in $\text{cc}(u^{(1)})$ or $\text{cc}(u^{(2)})$. Fix such shuffle.

We can create fillings $T_{u^{(1)}}^w, T_{u^{(2)}}^w$ of shapes $\text{ctype}(u^{(1)}), \text{ctype}(u^{(2)})$ using Algorithm 2.16. Similar to the modification we described in Section 2.5, we fill the boxes as we add them to the partition. However, we now fill the box with tuples (k, i) . Consider the box we add to $T_{u^{(1)}}^w$ when reading the letter $\text{ctype}(u^{(1)})_m$ for the k th time. There exists an i such $\text{ctype}(u^{(1)})_m$ corresponds to $\text{cc}(w)_{n-i+1}$ in the fixed shuffle. We fill the box with the tuple (k, i) .

Note that the convention for the index we put in the box is different from what we used in Section 4. The $(n - i + 1)$ will be useful when we define a total ordering on these entries. We are abusing notation here, since the fillings depend on how we shuffle the two cocharge words to get $\text{cc}(w)$, not just w .

Example 5.3. Let $w = 5 \ 9 \ 1 \ 2 \ 6 \ 7 \ 10 \ 3 \ 8 \ 4$. Then $\text{cc}(w) = \mathbf{1 \ 2 \ 0 \ 0 \ 1 \ 1 \ 2 \ 0 \ 1 \ 0} \in \text{Sh}(\mathbf{120012}, \mathbf{1010})$, where we have colored the entries to denote the shuffle we chose. The resulting fillings $T_{u^{(1)}}^w, T_{u^{(2)}}^w$ are depicted in Figure 1.

$$(T_{u^{(1)}} + T_{u^{(2)}})^w = \begin{array}{|c|c|c|c|} \hline (3,4) & & & \\ \hline (2,9) & & & \\ \hline (2,5) & & & \\ \hline (1,10) & (1,2) & (1,6) & \\ \hline (1,7) & (1,8) & (1,1) & (1,3) \\ \hline \end{array}$$

FIGURE 2

From Lemma 2.19, we know if (k, i) appears in row r of $T_{u^{(1)}}$, $T_{u^{(2)}}$ we must have $k + cc(w)_{n-i+1} = r$. Furthermore, since the second coordinate of an entry records which letter in $cc(w)$ created that box, for each $i \in \{1, \dots, n\}$, there exists a unique k such that (k, i) appears in $T_{u^{(1)}}^w$ or $T_{u^{(2)}}^w$. This (k, i) appears exactly once.

We combine $T_{u^{(1)}}^w, T_{u^{(2)}}^w$ to get a filling of shape $c\text{type}(u^{(1)}) + c\text{type}(u^{(2)})$, denoted $(T_{u^{(1)}} + T_{u^{(2)}})^w$, by taking the row wise sum of the two diagrams.

Example 5.4. We combine the two diagrams we get from Example 5.3 to get a filling of $(2 + 2, 1 + 2, 1, 1, 1)$, depicted in Figure 2.

The lexicographic ordering (\preceq) on tuples $(a, b) \in \mathbb{Z}_{>0}^2$ is defined by:

$$(a, b) \prec (c, d) \Leftrightarrow \begin{cases} a < c \\ a = c, b < d. \end{cases}$$

The lexicographic ordering of the tuples keeps track of the order in which we read the entries in Algorithm 2.16. That is: if $(k_1, i_1) \preceq (k_2, i_2)$, then we read $cc(w)_{n-i_1+1}$ for the k_1 th time before we read $cc(w)_{n-i_2+1}$ for the k_2 th time.

Let $T_{u^{(1)}+u^{(2)}}^w$ be the filling obtained by rearranging the entries within the rows of $(T_{u^{(1)}} + T_{u^{(2)}})^w$ so that the entries within the rows are increasing from left to right with respect to \preceq . We have the following result:

Lemma 5.5. *The filling $T_{u^{(1)}+u^{(2)}}^w$ satisfies the following conditions:*

- (i) Each $i \in \{1, 2, \dots, n\}$ appears in the second coordinate of an entry exactly once,
- (ii) The entries in each row and column are increasing with respect to the lexicographic ordering on tuples,
- (iii) For any entry (k, i) in row r of $T_{u^{(1)}+u^{(2)}}^w$, we have $k + cc(w)_{n-i+1} = r$.

Proof. The conditions (i) and (iii) hold by construction. It suffices to show the entries within each columns are increasing.

Let (k, i) be the entry in the row r , column j , where $r > 2$. We know (k, i) is the j th smallest thing in row r , since we reordered within the rows so that the entries were

increasing. For each entry in row r of $T_{u^{(1)}+u^{(2)}}^w$, there is an entry in row $r - 1$ that appeared directly below it in $T_{u^{(1)}}^w$ or $T_{u^{(2)}}^w$. Since $T_{u^{(1)}}^w, T_{u^{(2)}}^w$ were constructed through the Algorithm 2.16, we know that their columns are increasing with respect to the lexicographic ordering.

Thus for each entry (k', i) in row r such that $(k', i') \preceq (k, i)$, there is an entry in the row $(r - 1)$ that was directly below (k', i) in $T_{u^{(1)}}^w$ or $T_{u^{(2)}}^w$ which is smaller than (k', i') .

Therefore, if (k, i) is the j th smallest thing in row r , there are at least j many things in row $(r - 1)$ that are smaller than (k, i) . Hence (k, i) is larger than the entry directly below it. \square

Example 5.6. Continuing with our example, we have

$$T_{u^{(1)}+u^{(2)}} = \begin{array}{|c|} \hline (3,4) \\ \hline (2,9) \\ \hline (2,5) \\ \hline (1,2) & (1,6) & (1,10) \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} .$$

The entries are increasing within each row and column.

We now define the notion of row insertion when dealing with these tableaux. This is a modification of the classical row insertion used in RSK, except we also check the column increasing condition when inserting and we may modify more than one entry in the row, by sliding part of the row over to the right by one.

Algorithm 5.7. (Modified Row Insertion) Consider a filling T of a two row partition shape, where each entry is a tuple in $\mathbb{Z}_{>0}^2$ and the rows and columns are increasing with respect to the lexicographic ordering.

Let $(a, b) \in \mathbb{Z}_{>0}^2$. We *insert* (a, b) into the second row of this partition by doing the following. We first find the leftmost (c, d) in the second row that satisfies:

$$(5.1) \quad (a, b) \preceq (c, d),$$

$$(5.2) \quad (a, b) \succeq (e, f)$$

when (e, f) denotes the entry directly below (c, d) . If no such (c, d) exists, we say (a, b) *pops out* of this row and we are done.

If such (c, d) exists, replace (c, d) with (a, b) . The resulting filling is still increasing in the rows and columns. The only place where we could have a contradiction is directly to the left of (a, b) , but this cannot happen since (c, d) is the leftmost entry that satisfies (5.1) and (5.2).

We now repeat the steps above to insert (c, d) into the same row. We continue until an entry pops out or we insert into the last box of the row.

If we insert into the last box of the row, replacing (g, h) , we finish our insertion process by appending (g, h) to the end of the second row if the resulting filling has increasing columns

and is still a partition shape. This only happens if the second row was strictly shorter than the first. In this case, nothing pops out. If the two rows are the same length, (g, h) pops out and the insertion is complete.

Note that once we successfully insert (a, b) , the entry (c, d) gets inserted to the direct right of (a, b) or it pops out. From this, we can see that this process is equivalent to determining where we can put (a, b) and then moving the entries larger than it to the right by one until we find a contradiction with the bottom row or we reach the end of the row.

We illustrate this through the following examples. The first one results in an entry popping out.

Example 5.8. We insert $(a, b) = (2, 6)$ into the second row of the following partition:

(2,4)	(3,2)	(3,3)	(4,1)	
(1,1)	(2,5)	(2,7)	(3,8)	(5,2)

Since $(3, 2)$ is the leftmost entry satisfying (5.1) and (5.2), we replace $(3, 2)$ with $(2, 6)$:

(3,2)

(2,4)	(2,6)	(3,3)	(4,1)	
(1,1)	(2,5)	(2,7)	(3,8)	(5,2)

Now we repeat: we replace $(3, 3)$ with $(3, 2)$:

(3,3)

(2,4)	(2,6)	(3,2)	(4,1)	
(1,1)	(2,5)	(2,7)	(3,8)	(5,2)

However, we cannot replace $(4, 1)$ with $(3, 3)$ since $(3, 8) \not\geq (3, 3)$. Hence $(3, 3)$ pops out of this insertion and we are done.

We also give an example of when nothing pops out of the insertion process.

Example 5.9. We insert $(a, b) = (2, 6)$ into the second row of the following partition:

(2,4)	(3,2)	
(1,1)	(2,5)	(2,7)

Since $(3, 2)$ is the leftmost entry satisfying (5.1) and (5.2), we replace $(3, 2)$ with $(2, 6)$:

$$(3,2)$$

(2,4)	(2,6)	
(1,1)	(2,5)	(2,7)

Now, we can append $(3, 2)$ to the end of row 2 while maintaining the column increasing condition. Adding a box to the second row does not change the fact that the shape is a two row partition. Hence we insert $(3, 2)$ into the empty spot.

(2,4)	(3,2)	(3,2)
(1,1)	(2,5)	(2,7)

Note that nothing pops out at the end of this insertion.

We make the following observation.

Lemma 5.10. *If we insert (a, b) into row 2 of T and (x, y) pops out, we have $(a, b) \preceq (x, y)$.*

Proof. If we are unable to insert (a, b) , then $(x, y) = (a, b)$. Otherwise, the insertion replaces larger entries with smaller ones. \square

We can extend this definition of row insertion to any partition shape tableau T filled with tuples in $\mathbb{Z}_{>0}^2$ where the rows and columns are increasing. For any row $r > 1$ of T and $(a, b) \in \mathbb{Z}_{>0}^2$, we define $T \leftarrow_r (a, b)$ to be the result of inserting (a, b) into row r of T . Unlike RSK, we only modify row r ; we do not continue inserting into higher rows.

Lemma 5.11. *Consider a three row partition shape tableau T filled with tuples in $\mathbb{Z}_{>0}^2$ where the entries within the rows and columns are increasing. Consider two tuples $(a, b) \preceq (a', b')$. Let $T \leftarrow_2 (a, b)$ denote the tableau we get by inserting (a, b) into the second row of T .*

- (1) *Assume (x, y) pops out when we insert (a, b) into row 2 of T . If (x', y') pops out when we insert (a', b') into row 3 of $T \leftarrow_2 (a, b)$, then $(x, y) \preceq (x', y')$.*
- (2) *If nothing pops out when we insert (a, b) into row 2 of T , then nothing pops out when we insert (a', b') into row 3 of $T \leftarrow_2 (a, b)$.*

Proof. (1) Note that if $(x, y) = (a, b)$, the claim follows immediately. Otherwise, we placed (a, b) in the second row of T when inserting. Let (g, h) denote the entry in row 3 that is directly above (a, b) in $T \leftarrow_2 (a, b)$. Let (p, q) be the entry in row (a, b) directly above where (x, y) was in T , before it popped out. Since $(x, y) \preceq (p, q)$, it suffices to show $(p, q) \preceq (x', y')$.

When we inserted (a, b) into row 2 of T , we first put (a, b) below (g, h) . Then, we moved each entry in row 2 to the right by 1 until we put an entry under (p, q) . This resulted in (x, y) popping out.

Now, we insert (a', b') into row 3 of $T \leftarrow_2 (a, b)$. We proceed by cases. If $(a', b') \preceq (g, h)$, then (a', b') can be inserted in some spot to the left of (g, h) by just checking (5.1): the column condition follows automatically since $(a, b) \preceq (a', b')$, thus (a', b') is larger than any

entry to the left of (a, b) in row r . Any entry between (a', b') and (g, h) must also be larger than (a, b) , so we can keep inserting until we arrive at (g, h) .

We can also move all the entries from (g, h) up to (p, q) over to the right by one, because this is just equivalent to realigning rows 2 and 3 of $T \leftarrow_2 (a, b)$ to how they were in T . Thus the insertion process will continue until we reach (p, q) .

As a result, we either get through the entire row without anything popping out, or we pop out an entry (x', y') that is at least (p, q) . Thus $(p, q) \preceq (x', y')$.

If $(g, h) \preceq (a', b') \preceq (p, q)$, then we can always insert (a', b') between (g, h) and (p, q) . Consider $(g_1, h_1) \preceq (g_2, h_2)$ that appear consecutively in row 2 of $T \leftarrow_2 (a, b)$ such that $(g, h) \preceq (g_1, h_1) \preceq (a', b') \preceq (g_2, h_2) \preceq (p, q)$. Then, we know that the entry (s_1, t_1) that appears directly below (g_2, h_2) was originally below (g_1, h_1) in T . Thus $(s_1, t_1) \preceq (g_1, h_1) \preceq (a', b') \preceq (g_2, h_2)$, which means we can replace (g_2, h_2) with (a', b') . Once we insert (a', b') into row 2, we can use the same argument as above to conclude that $(x, y) \preceq (p, q) \preceq (x', y')$.

If $(p, q) \preceq (a', b')$, the claim automatically holds since $(a', b') \preceq (x', y')$ (with potential equality).

The second statement follows from the same argument: if such (x, y) does not exist, then (p, q) does not exist. Thus the two cases we consider are $(a', b') \preceq (g, h)$ and $(g, h) \preceq (a', b')$, which correspond to the first two cases in the previous argument. We can keep inserting into row 3 until we add a box at the end of row 3, which we know we can do since row 2 of $T \leftarrow_2 (a, b)$ is one longer than row 2 of T . \square

Now, we use this new notion of insertion to define a modification of Algorithm 2.16 that keeps track of the difference between $\text{ctype}(w)$ and $\text{ctype}(u^{(1)})$ and $\text{ctype}(u^{(2)})$. The input $(\text{cc}(w), \emptyset, T_{u^{(1)}+u^{(2)}}^w)$ depends on the shuffle of $\text{cc}(u^{(1)})$, $\text{cc}(u^{(2)})$ that is equal to $\text{cc}(w)$ that we fix.

Algorithm 5.12. The input is $(\text{cc}(w), \emptyset, S)$, where S is a filling of a partition shape that satisfies the conditions in Lemma 5.5. As we did for Algorithm 2.16, we apply the function f repeatedly to the first two coordinates until we get $(\emptyset, \text{ctype}(w))$ in the first two coordinates. We modify the last coordinate when we add a box to the partition.

At each step, we say we *read* (k, i) if we read the letter corresponding to $\text{cc}(w)_{n-i+1}$ for the k th time. Anything smaller than (k, i) in the lexicographic order has *already been read*. After each step of the algorithm, we get a triple (z, ν, T) where z is a word, ν is some partition, and T is a tableau of partition shape that satisfies the following conditions:

- (1) Each $i \in \{1, 2, \dots, n\}$ appears in the second coordinate of a tuple exactly once,
- (2) The entries in each row and column of T are increasing with respect to the lexicographic ordering on tuples,
- (3) As partitions, we have $\text{shape}(T) \supseteq \text{shape}(S)$,
- (4) For any entry (k, i) in the r th row of T , we have $k + \text{cc}(w)_{n-i+1} = r$,
- (5) A tuple (k, i) that appears in T has already been read if and only if it is contained in the subdiagram of T of shape ν .
- (6) A tuple (k, i) that appears in T has not been read if and only if the letter corresponding to $\text{cc}(w)_{n-b+1}$ has not been deleted from the input word.

We modify the filling T in the following way. Consider the step when we read (k, i) , where the input is (z, ν, T) , and it results in adding a box to row r of ν .

From (1), we know there exists m such that (m, i) appears in T . Still i corresponding to a letter in z that has not been deleted yet, we know (m, i) has not been read yet from (6). Thus $m \geq k$. From (4), this means (m, i) appears in row r' of T where $r \leq r'$. If (m, i) is in the r th row of T , then $k = m$. Note that from (2) and (5), we know that the box with (k, i) corresponds to the box that we just added to ν . In this case, we do not modify T and conditions (1) - (6) still hold.

If (m, i) is in row r' where $r < r'$, we will create a new tableau \tilde{T} of partition shape that satisfies conditions (1) - (6). Delete the box containing (m, i) as well as the ones directly above it in T . This results in deleting entries $\{(m, i), (m_1, i_1), \dots, (m_s, i_s)\}$, where (m_t, i_t) denotes the entry that was t boxes above (m, i) . Note that since the entries are increasing within the columns of T , we have

$$(5.3) \quad (m, i) \preceq (m_1, i_1) \preceq \dots \preceq (m_s, i_s)$$

The rows and columns are still increasing with respect to the lexicographic ordering if we ignore the empty spaces in the column that contained $\{(m, i), (m_1, i_1), \dots, (m_s, i_s)\}$.

We will create a new partition shape filling of n boxes by inserting the entries we removed back into this filling.

Let $d = r' - r$. We first insert $(m - d, i) = (k, i)$ into row r using Algorithm 5.7. We modify the first coordinate to maintain condition (4). Since (k, i) is the smallest unread tuple, we know that (k, i) goes in the first box in row r that contains an unread tuple. From (5) and (6), we know this is the box that we add to ν at this step.

If an entry pops out, we denote that entry by (x_0, y_0) . Now, insert $(m_1 - d, i_1)$ into row $r + 1$. If some entry (x_1, y_1) pops out, we know $(x_0, y_0) \preceq (x_1, y_1)$ from Lemma 5.11 (1) since $(m - d, i) \preceq (m_1 - d, i_1)$.

In this way, we insert $(m_1 - d, i_1), \dots, (m_s - d, i_s)$ into rows $r + 1, \dots, r + s$. If we insert into a row with a gap in the middle, we ensure that we maintain the empty spot by skipping over it in the insertion process. That is: we never insert in the empty spot in the middle of the row. The elements that pop out at each step have the following relation:

$$(5.4) \quad (k, i) \preceq (x_0, y_0) \preceq (x_1, y_1) \preceq \dots \preceq (x_j, y_j),$$

where either $j = s$ or nothing popped out when inserting $(m_{j+1} - d, i_{j+1})$ into row $r + j + 1$.

If no entry pops out when we insert (k, i) into row r we can insert $(m_1 - d, i_1), \dots, (m_s - d, i_s)$ into rows $r + 1, \dots, r + s$ without anything popping out by Lemma 5.11 (2). Similarly, if nothing popped out when inserting $(m_{j+1} - d, i_{j+1})$ into row $r + j + 1$, we can insert the remaining $(m_{j+2} - d, i_{j+2}), \dots, (m_s - d, i_s)$ into rows $r + j + 2, \dots, r + s$ without anything popping out.

Once we have inserted up to $(m_s - d, i_s)$, we still have an empty column where (m, i) used to be and above. Since the insertion algorithm replaces entries with smaller ones, we know that the entry below the empty space in row r' must still be smaller than (m, i) . If there exists an entry (x_0, y_0) that popped out of row r , we have the following inequalities

using (5.4):

$$(5.5) \quad (k+d, i) = (m, i) \preceq (x_0+d, y_0) \preceq (x_1+d, y_1) \preceq \dots (x_j+d, y_j).$$

Thus we can fill the empty column with the entries $(x_0+d, y_0), (x_1+d, y_1), (x_j+d, y_j)$ and preserve the column increasing condition for all the columns.

Now that we have put all the entries back into the filling, we rearrange the entries within rows r' and above so that the entries within the rows are increasing and we have a partition shape with no gaps. Let the resulting diagram be \tilde{T} .

We can see if $j < s$, this corresponds to lowering boxes in the shape, hence $\text{shape}(T) \trianglelefteq \text{shape}(\tilde{T})$. If $j = s$, we have that $\text{shape}(T) = \text{shape}(\tilde{T})$.

We check that the resulting diagram \tilde{T} satisfies conditions (1) - (6). Conditions (1),(4) hold by construction. Condition (3) holds from arguments above.

We see that (2) holds using the argument in the proof of Lemma 5.5.

At each point, we only move entries that have not been read yet: this follows from $(k, i) = (m-d, i)$ being the smallest unread entry and equations (5.3). We also see that throughout the insertion process we do not move any of the entries in the subdiagram of shape ν , since it consists of all the tuples that have been read. The box we add to ν contains (k, i) , which is the newly read tuple. Hence (5) and (6) holds.

At the end of the algorithm, we have $(\emptyset, \text{ctype}(w), T_w)$ where $\text{shape}(T_w) = \text{ctype}(w)$. We include an example of this algorithm in the Appendix. Theorem 5.2 immediately follows from this algorithm.

Proof of Theorem 5.2. We apply Algorithm 5.12 to the initial input $(\text{cc}(w), \emptyset, T_{u^{(1)}+u^{(2)}}^w)$. At any point in the algorithm, where the tuple is of the form (z, ν, T) , we have

$$\text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)}) = \text{shape}(T_{u^{(1)}+u^{(2)}}^w) \trianglelefteq \text{shape}(T)$$

. Since we end with tuple $(\emptyset, \text{ctype}(w), T_w)$ with $\text{shape}(T_w) = \text{ctype}(w)$, it follows that $\text{ctype}(u^{(1)}) + \text{ctype}(u^{(2)}) \trianglelefteq \text{ctype}(w)$. \square

Throughout this algorithm, we can think of T as keeping track of the “lower bounds” of insertion: that is, if we read a certain entry in T , we must add a box there. However, we may add boxes earlier, which is when we perform the chains of insertions.

We can also use this modified algorithm to prove the following statement about how certain modifications to permutations raise the catabolizability type.

Proposition 5.13. Consider $w \in S_n$ and index i such that $w_i + 1 < w_{i+1}$. If $\tilde{w} = w_1 \dots w_{i+1} w_i \dots w_n$, we have $\text{ctype}(\tilde{w}) \trianglerighteq \text{ctype}(w)$.

Proof. We know $\text{cc}(w)_i < \text{cc}(w)_{i+1}$, since they differ by more than one. This swap does not change the cocharge label of any of the letters. Thus $\text{cc}(\tilde{w})$ is the word we get by swapping $\text{cc}(w)_i$ and $\text{cc}(w)_{i+1}$ in $\text{cc}(w)$.

Let T_w be the filling of $\text{ctype}(w)$ we get from Algorithm 2.16, where we fill a box with tuple (k, i) if we add the box when reading the letter $\text{cc}(w)_{n-i+1}$ for the k th time. There exists unique k_1, k_2 such that $(k_1, n-i+1), (k_2, n-i)$ appear in T_w . Replace these two

tuples with $(k_1, n - i), (k_2, n - i + 1)$ to account for the swapping of $\text{cc}(w)_i$ and $\text{cc}(w)_{i+1}$. Denote the resulting filling by $T_w^{\tilde{w}}$.

The filling $T_w^{\tilde{w}}$ satisfies the conditions (1), (3) in Lemma 5.5 by construction. We can check that it satisfies (2) as well. Note that the only case where swapping the second coordinates changes the relative order on the elements in $T_w^{\tilde{w}}$ is if $k_1 = k_2$. In this case, the potential issue occurs if $(k_1, n - i), (k_2, n - i + 1)$ appear in the same row or column of T_w . However, this never happens. Since $\text{cc}(w)_i \neq \text{cc}(w)_{i+1}$, if the two appeared in the same row we would have $k_1 > k_2$ from Lemma 2.19. If $k_1 = k_2$, then $(k_1, n - i)$ appears in a higher row than $(k_2, n - i + 1)$. Since we add $(k_1, n - i)$ to T_w before we add $(k_2, n - i + 1)$, it is impossible that these two tuples appear in the same column of T_w . Hence $T_w^{\tilde{w}}$ satisfies (2).

Now, we apply Algorithm 5.12 to the initial input $(\text{cc}(\tilde{w}), \emptyset, T_w^{\tilde{w}})$. Since $\text{shape}(T_w^{\tilde{w}}) = \text{ctype}(w)$ and the resulting tuple is $(\emptyset, \text{ctype}(\tilde{w}), T_{\tilde{w}})$ with $\text{shape}(T_{\tilde{w}}) = \text{ctype}(\tilde{w})$, it follows that $\text{ctype}(\tilde{w}) \supseteq \text{ctype}(w)$. \square

6. ANTISYMMETRIC PART

In the work of Garsia–Procesi [6, Section 3], the ungraded Frobenius character of R_μ is identified to be the complete homogeneous symmetric function h_μ directly from the structure of R_μ . However, they require further properties of R_μ , as well as the modified Hall-Littlewood polynomials, to identify the graded Frobenius character.

In this section, we use properties of our charge monomial basis $\{\mathbf{x}^{c(w)} \mid \text{ctype}(P(w)^t) \supseteq \mu^t\}$ to give a direct, elementary proof of the fact that $\text{Frob}_q(R_\mu) = \tilde{H}_\mu[X; q]$ that only depends on the well-known combinatorial formula (1.1) and the ungraded character.

We do this by showing that for any composition $\gamma = (\gamma_1, \dots, \gamma_l) \models n$, we have $\text{Hilb}(N_\gamma R_{\mu^t}) = \langle e_\gamma, \tilde{H}_{\mu^t}[X; q] \rangle$, where N_γ is the antisymmetrizer with respect to the Young subgroup $S_\gamma = S_{\gamma_1} \times \dots \times S_{\gamma_l} \subset S_n$:

$$N_\gamma = \sum_{\sigma \in S_\gamma} \text{sgn}(\sigma) \sigma.$$

Recall that we have the following formula for $\tilde{H}_{\mu^t}[X; q]$ for $\mu \vdash n$, using Proposition 2.9:

$$(6.1) \quad \tilde{H}_{\mu^t}[X; q] = \sum_{\substack{S \in \text{SYT} \\ \text{ctype}(S^t) \supseteq \mu}} q^{\text{charge}(S)} s_{\text{shape}(S^t)}.$$

We apply ω to both terms. Since $\{m_\lambda\}, \{h_\gamma\}$ are dual with respect to the inner product, we get

$$\langle e_\gamma, \tilde{H}_{\mu^t}[X; q] \rangle = \langle h_\gamma, \omega \tilde{H}_{\mu^t}[X; q] \rangle = \langle m_\gamma, \omega \tilde{H}_{\mu^t}[X; q] \rangle,$$

where $\langle m_\gamma, \omega \tilde{H}_{\mu^t}[X; q] \rangle$ denotes the coefficient of m_γ in the monomial symmetric function expansion of $\omega \tilde{H}_{\mu^t}[X; q]$. We compute $\omega \tilde{H}_{\mu^t}[X; q]$ using $\omega s_\lambda = s_{\lambda^t}$ for any partition λ and

the combinatorial formula (6.1):

$$\omega \tilde{H}_{\mu^t}[X; q] = \sum_{\substack{S \in \text{SYT} \\ \text{ctype}(S^t) \supseteq \mu^t}} q^{\text{charge}(S)} s_{\text{shape}(S)}.$$

From the above, we can see

$$\begin{aligned} \langle e_\gamma, \tilde{H}_{\mu^t}[X; q] \rangle &= \sum_{\substack{S \in \text{SYT} \\ \text{ctype}(S^t) \supseteq \mu^t}} q^{\text{charge}(S)} (\langle m_\gamma \rangle s_{\text{shape}(S)}) \\ &= \sum_{\substack{S \in \text{SYT} \\ \text{ctype}(S^t) \supseteq \mu^t}} q^{\text{charge}(S)} K_{\text{shape}(S), \gamma}. \end{aligned}$$

Furthermore, we can rewrite expression (2.1) into a statement about permutations:

$$K_{\lambda, \gamma} = |\{w \in S_n \mid \text{shape}(P(w)) = \lambda, \text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}\}|.$$

Combining these observations, we have the following equality:

$$(6.2) \quad \langle e_\gamma, \tilde{H}_{\mu^t}[X; q] \rangle = \sum_{\substack{w \in S_n \\ \text{ctype}(P(w)^t) \supseteq \mu^t \\ \text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}}} q^{\text{charge}(w)}.$$

From this, proving $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}[X; q]$ is equivalent to showing that for any $\gamma \models n$, the Hilbert series $\text{Hilb}(N_\gamma R_{\mu^t})$ is equal to the right hand expression of (6.2). We show this by proving that the natural subset of charge monomials to expect from Equation (6.2) give a basis of $N_\gamma R_{\mu^t}$ by antisymmetrization.

Using the ungraded character of R_μ , given by h_μ , along with Proposition 2.3, we can make the following observation:

Lemma 6.1. *The dimension of $N_\gamma R_{\mu^t}$ is*

$$|\{w \in S_n, \text{ctype}(P(w)^t) \supseteq \mu^t, \text{Des}(Q(w)) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}\}|.$$

Proof. Since the ungraded Frobenius character of R_{μ^t} is h_{μ^t} , we know that $\dim(N_\gamma R_{\mu^t}) = \langle e_\gamma, h_{\mu^t} \rangle$. Using the fact that $\tilde{H}_{\mu^t}[X; 1] = h_{\mu^t}$, we have:

$$\dim(N_\gamma R_{\mu^t}) = \langle e_\gamma, h_{\mu^t} \rangle = \langle e_\gamma, \tilde{H}_{\mu^t}[X; 1] \rangle = \sum_{\substack{w \in S_n \\ \text{ctype}(P(w)^t) \supseteq \mu^t \\ \text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}}} 1. \quad \square$$

We now use this fact to determine a basis of $N_\gamma R_{\mu^t}$.

Proposition B. *Let $\mu \vdash n$ and $\gamma = (\gamma_1, \dots, \gamma_l) \models n$. The set*

$$(6.3) \quad \{N_\gamma \mathbf{x}^{c(w)} \mid w \in S_n, \text{ctype}(P(w)^t) \supseteq \mu^t, \text{Des}(Q(w)) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}\}.$$

is a basis of $N_\gamma R_{\mu^t}$.

Proof. We first show that the terms in (6.3) are nonzero. Consider $w \in S_n$ such that $\text{ctype}(\text{rev}(w)) \supseteq \mu^t$, and $\text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}$. We divide w into blocks of size $\gamma_1, \gamma_2, \dots, \gamma_l$ where the entries within each block are strictly increasing. This implies that no two entries in the same block have the same charge value. Note that S_γ permutes indices within the blocks, thus each monomial term of $N_\gamma x^{c(w)}$ is distinct. Hence $N_\gamma x^{c(w)} \neq 0$. From Lemma 6.1, this shows that our set has the correct cardinality to be a basis of $N_\gamma R_{\mu^t}$.

Now, we show that (6.3) spans $N_\gamma R_{\mu^t}$. The set $\{N_\gamma \mathbf{x}^\alpha \mid \alpha \in \mathcal{C}_\mu\}$ spans $N_\gamma R_{\mu^t}$ since $\{\mathbf{x}^\alpha \mid \alpha \in \mathcal{C}_\mu\}$ is a basis of R_{μ^t} . Consider $w \in S_n$ such that $\text{ctype}(\text{rev}(w)) \supseteq \mu^t$, $N_\gamma x^{c(w)} \neq 0$ but w does not have the correct descent set. We will show that $N_\gamma \mathbf{x}^{c(w)} = N_\gamma \mathbf{x}^{c(\text{sort}_\gamma(w))}$ for some $\text{sort}_\gamma(w) \in S_n$ with the correct descent set and catabolizability type.

If we divide $c(w)$ into blocks of size $\gamma_1, \gamma_2, \dots, \gamma_l$, the entries within a block all have distinct charge labels since $N_\gamma x^{c(w)} \neq 0$. From this, we can rearrange the entries in w so that they are strictly increasing within the blocks without changing the charge labels of the respective entries. That is: if $w_i > w_{i+1}$ with $c(w)_i \neq c(w)_{i+1}$, we know $w_i \neq (w_{i+1} + 1)$, hence we can swap w_i and w_{i+1} without changing the respective charge labels. We can repeat this until we sort w so that it is increasing within the blocks. Let $\text{sort}_\gamma(w)$ be the resulting permutation. We see that $\text{Des}(\text{sort}_\gamma(w)) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}$. By Proposition 5.13, we know $\text{ctype}(\text{rev}(\text{sort}_\gamma(w))) \supseteq \text{ctype}(\text{rev}(w)) \supseteq \mu^t$, hence $N_\gamma \mathbf{x}^{c(\text{sort}_\gamma(w))}$ is in (6.3). \square

Using the construction of our basis, we can translate questions about the structure of R_{μ^t} into questions about conditions on tableaux. In this case, choosing the elements that are a basis of $N_\gamma R_{\mu^t}$ is equivalent to looking at pairs of standard tableaux (P, Q) of the same shape with conditions on both P and Q . We illustrate this with an example:

Example 6.2. Consider $\mu = (3, 1)$ and $\gamma = (2, 2)$. All standard tableaux P that satisfy $\text{ctype}(P^t) \supseteq (2, 1, 1) = (3, 1)^t$ are listed below:

$$\begin{array}{|c|} \hline 2 \\ \hline \end{array} \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 2 & 4 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 3 \\ \hline \end{array}, \begin{array}{|c|} \hline 4 \\ \hline \end{array} \begin{array}{|c|} \hline 2 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 3 \\ \hline \end{array}, \begin{array}{|c|} \hline 3 \\ \hline \end{array} \begin{array}{|c|} \hline 2 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 4 \\ \hline \end{array}, \begin{array}{|c|} \hline 4 \\ \hline \end{array} \begin{array}{|c|} \hline 3 \\ \hline \end{array} \begin{array}{|c|} \hline 2 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array}$$

We also list all standard tableaux Q such that $\text{Des}(Q) \subset \{2\}$:

$$\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline \end{array}, \begin{array}{|c|} \hline 3 \\ \hline \end{array} \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & 4 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 2 \\ \hline \end{array}$$

Thus the pairs (P, Q) of standard tableaux of the same shape that satisfy $\text{ctype}(P^t) \supseteq (2, 1, 1)$ and $\text{Des}(Q) = (2, 2)$ are the pairs $(\begin{array}{|c|} \hline 2 \\ \hline \end{array} \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline \end{array}, \begin{array}{|c|} \hline 3 \\ \hline \end{array} \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline \end{array})$ and $(\begin{array}{|c|c|} \hline 2 & 4 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 3 \\ \hline \end{array}, \begin{array}{|c|c|} \hline 3 & 4 \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline 2 \\ \hline \end{array})$. These pairs corresponds to the words 2314, 2413, which give us the charge monomials $x_2 x_4^2, x_2 x_4$. From this, we know the basis of $N_\gamma R_{\mu^t}$ is given by the polynomials

$$\{x_2 x_4^2 - x_1 x_4^2 - x_2 x_3^2 + x_1 x_3^2, x_2 x_4 - x_1 x_4 - x_2 x_3 + x_1 x_3\}.$$

The following corollary is immediate from Proposition B.

Corollary 6.3. For any partition μ of n we have $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}[X; q]$.

Proof. The graded Frobenius character of a $\mathbb{C}S_n$ -module V is uniquely determined by the values $\langle e_\gamma, \text{Frob}_q(V) \rangle$ for all $\gamma \vdash n$. From Proposition B, we have

$$\text{Hilb}(N_\gamma R_{\mu^t}) = \sum_{\substack{w \in S_n \\ \text{ctype}(P(w)^t) \supseteq \mu^t \\ \text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\}}} q^{\text{charge}(w)}$$

which is equal to $\langle e_\gamma, \text{Frob}_q(R_{\mu^t}) \rangle$ by Equation (6.2). Thus $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}[X; q]$. \square

Carlsson-Chou [3] have a similar result on the antisymmetric component with respect to a certain Young subgroup. We define Sh_γ (resp. Sh'_γ) to be the set of all permutations such that $\{1, 2, \dots, \gamma_1\}, \{\gamma_1 + 1, \dots, \gamma_1 + \gamma_2\}, \dots, \{\gamma_1 + \dots + \gamma_{l-1} + 1, \dots, n\}$ appear in increasing (resp. decreasing) order.

Theorem 6.4 (Theorem 3.8.2, Carlsson-Chou [3]). *The set $\{N_\gamma g_\sigma(\mathbf{x}) \mid g_\sigma(\mathbf{x}) = \mathbf{x}^\alpha \text{ for } \alpha \in \mathcal{D}_\mu, \sigma \in \text{Sh}'_\gamma\}$ is a basis of $N_\gamma R_{\mu^t}$.*

Similar to what we do above, they use their result to show that $\text{Frob}_q(R_{\mu^t}) = \tilde{H}_{\mu^t}$ by deriving a combinatorial formula for $\langle m_\gamma, \omega \tilde{H}_{\mu^t} \rangle$ using parking functions and shuffle combinatorics which matches their description of the basis.

In contrast to their result, ours arises naturally from the catabolizability formula (1.1) for \tilde{H}_{μ^t} . Our proof of the result is also independent of theirs: we do not relate their classification to ours when we show Proposition B, though they use a similar argument to show that their basis is linearly independent. However, it is easy to see that the two sets coincide by using $x^{c(w)} = g_{\text{rev}(w^{-1})}(\mathbf{x})$ to show that the two conditions are equivalent:

$$\text{Des}(w) \subset \{\gamma_1, \gamma_1 + \gamma_2, \dots, \gamma_1 + \dots + \gamma_{l-1}\} \Leftrightarrow w^{-1} \in \text{Sh}_\gamma \Leftrightarrow \text{rev}(w^{-1}) \in \text{Sh}'_\gamma.$$

ACKNOWLEDGEMENTS

The author thanks Mark Haiman for helpful conversation and guidance.

APPENDIX

Example of 5.12. Let $w = 3 \ 9 \ 1 \ 2 \ 6 \ 7 \ 10 \ 3 \ 8 \ 4$.

Then $cc(w) = \mathbf{1} \ \mathbf{2} \ \mathbf{0} \ \mathbf{0} \ \mathbf{1} \ \mathbf{1} \ \mathbf{2} \ \mathbf{0} \ \mathbf{1} \ \mathbf{0} \in \text{Sh}(\mathbf{120012}, \mathbf{1010})$, where we have colored the entries to denote how the two words are shuffled.

The initial input when applying Algorithm 5.12 is given by:

$$\left(\begin{array}{c} \\ \\ 1 \ 2 \ 0 \ 0 \ 1 \ 1 \ 2 \ 0 \ 1 \ 0, \emptyset, \\ \\ \begin{array}{|c|} \hline (3,4) \\ \hline (2,9) \\ \hline (2,5) \\ \hline (1,2) \ (1,6) \ (1,10) \\ \hline (1,1) \ (1,3) \ (1,7) \ (1,8) \\ \hline \end{array} \end{array} \right)$$

We can see that the first three steps of algorithm correspond to adding boxes $(1, 1), (1, 2), (1, 3)$ to the partition in the second coordinate. We do not modify the filling in the last coordinate. The resulting triple we get is:

$$\left(\begin{array}{c} \\ \\ 1 \ 2 \ 0 \ 0 \ 1 \ 1 \ \mathbf{2} \ \dots, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \\ \\ \begin{array}{|c|} \hline (3,4) \\ \hline (2,9) \\ \hline (2,5) \\ \hline (1,2) \ (1,6) \ (1,10) \\ \hline (1,1) \ (1,3) \ (1,7) \ (1,8) \\ \hline \end{array} \end{array} \right).$$

The entries have been read up to this point are exactly those in the shape $\nu = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$. These boxes are shaded.

At the fourth step (which corresponds to reading $\mathbf{2}$), we can see that we will add a box to the partition in the second coordinate. However, we can see that $(3, 4)$ is in the fifth row. Thus, we delete $(3, 4)$ and insert $(1, 4)$ into the third row of our filling:

$$(1,4) \rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) \\ \hline (2,5) \\ \hline (1,2) \ (1,6) \ (1,10) \\ \hline (1,1) \ (1,3) \ (1,7) \ (1,8) \\ \hline \end{array} \Rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) \\ \hline (1,4) \ (2,5) \\ \hline (1,2) \ (1,6) \ (1,10) \\ \hline (1,1) \ (1,3) \ (1,7) \ (1,8) \\ \hline \end{array}$$

The resulting triple after the fourth step is:

$$\left(1 \ 2 \ 0 \ 0 \ 1 \ \underline{1} \ - \ - \ - \ - \ , \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} , \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & (2,5) & & \\ \hline (1,2) & (1,6) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right) .$$

At the fifth step (corresponding to reading $\underline{1}$), we add a box to the partition in the second coordinate, as well as modify the filling in the third coordinate by deleting (2,5) and inserting (1,5) into the second row:

$$(1,5) \rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & & & \\ \hline (1,2) & (1,6) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \Rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & & & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \rightarrow (\mathbf{1,6})$$

We see that (1,6) pops out of the second row. We then place (2,6) into the empty spot in the third row, where (2,5) was.

$$(2,6) \rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & & & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \Rightarrow \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & (2,6) & & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} .$$

The resulting triple is

$$\left(1 \ 2 \ 0 \ 0 \ \underline{1} \ - \ - \ - \ - \ - \ , \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} , \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & (2,6) & & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right) .$$

At the next step, we cannot add anything to the second row. So we add one to the element in the word without changing either of the shapes:

$$\left(1 \ 2 \ 0 \ \underline{0} \ 2 \ \dots \dots \dots, \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline (2,9) & & \\ \hline (1,4) & (2,6) & \\ \hline (1,2) & (1,5) & (1,10) \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right).$$

The next two steps just add boxes to the first row of the partition in the second coordinate without changing the filling. We get

$$\left(1 \ \underline{2} \ \dots \ 2 \ \dots \dots \dots, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline (2,9) & & & \\ \hline (1,4) & (2,6) & & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right).$$

At the next step, we delete (2,9) and insert (1,9) into the third row of the filling to get:

$$\left(1 \ \dots \ 2 \ \dots \dots \dots, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline (1,4) & (1,9) & (2,6) & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right).$$

Now, we can continue to cycle through the word and add boxes to the partition until we get

$$\left(\dots \dots \dots, \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline (1,4) & (1,9) & (2,6) & \\ \hline (1,2) & (1,5) & (1,10) & \\ \hline (1,1) & (1,3) & (1,7) & (1,8) \\ \hline \end{array} \right).$$

REFERENCES

1. Jonah Blasiak, *An insertion algorithm for catabolizability*, *Europ. J. Combinatorics* **33** (2012), no. 2, pp 267 – 276.
2. L.M. Butler and American Mathematical Society, *Subgroup lattices and symmetric functions*, American Mathematical Society: Memoirs of the American Mathematical Society, American Mathematical Society, 1994.
3. Erik Carlsson and Raymond Chou, *A descent basis for the Garsia-Procesi module*, *Advances in Mathematics* **457** (2024).
4. Corrado de Concini and Claudio Procesi, *Symmetric functions, conjugacy classes and the flag variety*, *Inventiones mathematicae* **64** (1981).
5. William Fulton and Joe Harris, *Representation Theory: A First Course*, Graduate texts in mathematics, Springer, 1991.
6. Adriano M. Garsia and Claudio Procesi, *On certain graded S_n -modules and the q -Kostka polynomials*, *Adv. Math* **94.1** (1992), pp 82–138.
7. Ryoshi Hotta and Tonny Albert Springer, *A specialization theorem for certain Weyl group representations and an application to the green polynomials of unitary groups*, *Inventiones mathematicae* (1977), 113–127.
8. Alain Lascoux, *Cyclic permutations on words, tableaux and harmonic polynomials*, *Proc. of the Hyderabad Conference on Algebraic Groups* (1989), pp. 323 – 347.
9. Alain Lascoux and Marcel-Paul Schützenberger, *Sur une conjecture de H. O. Foulkes*, *C. R. Acad. Sci. Paris Sér. I Math* **288** (1978), pp 95 – 9.
10. I.G. Macdonald, *Symmetric functions and hall polynomials*, Oxford classic texts in the physical sciences, Clarendon Press, 1998.
11. Mark Shimozono and Jerzy Weyman, *Graded characters of modules supported in the closure of a nilpotent conjugacy class*, *Europ. J. Combinatorics* **21** (2000), no. 2, pp 257 – 288.
12. Tonny Albert Springer, *A construction of representations of Weyl groups*, *Inventiones mathematicae* **44** (1978), 279–293.
13. Richard P. Stanley and Sergey Fomin, *Enumerative Combinatorics, volume 2*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1999.
14. Toshiyuki Tanisaki, *Defining ideals of the closures of the conjugacy classes and representations of the weyl groups*, *Tohoku Math. J. (2)* **34** (1982), no. 4, 575–585 (en).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY
 Email address: mhanada@berkeley.edu