

THE PROPERTIES OF BANACH ALGEBRAS AFFECTED BY DIFFERENTIAL IDENTITIES

MOHAMED MOUMEN AND LAHCEN TAOUFIQ

ABSTRACT. Let \mathcal{B} be a Banach algebra. The interest of this article lies in the study of the commutativity of \mathcal{B} if certain specific algebraic identities hold over a non-empty open subset of \mathcal{B} . The limitations imposed in the hypothesis of our results are justified by some examples.

1. Introduction

\mathcal{B} will denote a Banach algebra with center $Z(\mathcal{B})$. If for every $x, y \in \mathcal{B}$, $x\mathcal{B}y = 0$ means that either $x = 0$ or $y = 0$, then \mathcal{B} is said to be prime. The anti-commutator $xy + yx$ will be represented by the symbol $x \circ y$, whereas the commutator $[x, y]$ will stand for the Lie product $xy - yx$. An additive mapping d in \mathcal{B} is said to be a derivation if $d(xy) = d(x)y + xd(y)$ for all $x, y \in \mathcal{B}$. We recall that $f : \mathcal{B} \rightarrow \mathcal{B}$ is called a homomorphism if $f(xy) = f(x)f(y)$ and $f(x + y) = f(x) + f(y)$ holds for any $x, y \in \mathcal{B}$. An additive mapping $F : \mathcal{B} \rightarrow \mathcal{B}$ is said to be a generalized derivation associated with a derivation d if $F(xy) = F(x)y + xd(y)$ for all $x, y \in \mathcal{B}$. Basic examples are derivations and generalized inner derivations, i.e., maps of type $x \mapsto ax + xb$ for some $a, b \in \mathcal{B}$. A mapping $f : \mathcal{B} \rightarrow \mathcal{B}$ is centralizing on a part S of \mathcal{B} if $[f(x), x] \in Z(\mathcal{B})$ for all $x \in S$, in particular if $[f(x), x] = 0$ for all $x \in S$ then f is said to be commuting on S .

In 1955, Divinsky [1] established that a simple Artinian ring is commutative if it possesses a commuting non-trivial automorphism, which began the development of commuting and centralizing mappings. Posner [6] demonstrated two years later that a prime ring with a nonzero centralizing derivation must be commutative.

Over the last few decades, several authors have proved commutativity theorems for prime and semiprime rings admitting automorphisms or derivations that are commuting or centralizing on an appropriate subset of the ring (see [5] for a partial bibliography). In the direction of Banach algebras, Yood [7] showed that a semiprime complex Banach algebra \mathcal{B} must be commutative if there are non-empty open subsets \mathcal{H}_1 and \mathcal{H}_2 of \mathcal{B} such that for each $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ there are positive integers $n = n(x, y) > 1$ and $m = m(x, y) > 1$ depending on x and y such that either $[x^n, y^m] \in Z(\mathcal{B})$ or $x^n y^m \in Z(\mathcal{B})$. Recently, M. Moumen, L. Taoufiq and L. Oukhetite [8] proved that a prime Banach algebra \mathcal{B} must be commutative if there are non-empty open subsets $\mathcal{H}_1, \mathcal{H}_2$ and a non-injective continuous derivation d satisfying for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ there are strictly positive integers $n = n(x, y), m = m(x, y)$ such that $d(x^n y^m) + [x^n, y^m] \in Z(\mathcal{B})$. In [9] M. Moumen, L. Taoufiq and A. Boua proved similar results (see [9] and [10] for additional details).

The importance of commutativity in many disciplines such as spectral theory, information security, and especially decryption theory, is the reason why we chose to continue in this domain. Our method is based on some functional analysis theorems as Baire's category theorem. Among the results, we have demonstrated that a prime Banach algebra

2020 Mathematics Subject Classification: 46J10, 47B47, 16W25, 46J45, 16N60.

Key words and phrases: Banach algebras; continuity; closed subspace; derivation; open subset.

\mathcal{B} must be commutative if it has non-empty open subsets $\mathcal{H}_1, \mathcal{H}_2$ and a continuous automorphism f satisfying for any $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ there exist two strictly positive integers $p = p(x, y), q = q(x, y)$ such that $f(x^p y^q) + x^p \circ y^q \in Z(\mathcal{B})$. Further, we provide applications for our results.

Throughout this article, we shall make some use of the following well-known results (see [9] for further details).

FACT 1. *Let \mathcal{R} be a prime ring.*

1. *If $x \in Z(\mathcal{R})$ and $xy \in Z(\mathcal{R})$, then $x = 0$ or $y \in Z(\mathcal{R})$.*
2. *If \mathcal{R} has a nonzero centralizing derivation, then \mathcal{R} is commutative.*

2. Automorphism in Banach algebras

The development of the confirmation of our main ideas depends on the following lemma, attributed to Duncan and Bonsall [2].

LEMMA 1. *Let $s(t) = \sum_{i=0}^n t^i b_i$ be a polynomial with coefficients in \mathcal{B} (complex or real Banach algebra). If for an infinite set of real values of t , $s(t) \in C$, where C is a closed linear subspace of \mathcal{B} , then every b_i lies in C .*

THEOREM 1. *Let \mathcal{B} be a prime Banach algebra, \mathcal{H}_1 and \mathcal{H}_2 are non-void open subsets of \mathcal{B} . \mathcal{B} must be commutative if admits a continuous automorphism f satisfying: $f(x^p y^q) + x^p \circ y^q \in Z(\mathcal{B})$ for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ where p, q are not fixed but depends on the pair of elements x and y .*

Proof. We define the following set for any $(p, q) \in \mathbb{N}^* \times \mathbb{N}^*$:

$$O_{p,q} = \{(x, y) \in \mathcal{B}^2 \mid f(x^p y^q) + x^p \circ y^q \notin Z(\mathcal{B})\}.$$

We note that $(\cap O_{p,q}) \cap (\mathcal{H}_1 \times \mathcal{H}_2) = \emptyset$, indeed:

If they is $(a, b) \in \mathcal{H}_1 \times \mathcal{H}_2$ such that $(a, b) \in O_{p,q}$ for all $(p, q) \in \mathbb{N}^* \times \mathbb{N}^*$, then $f(x^p y^q) + x^p \circ y^q \notin Z(\mathcal{B})$ for all $(p, q) \in \mathbb{N}^* \times \mathbb{N}^*$, this contradicts the theorem's assumption. We now verify that every $O_{p,q}$ is open in $\mathcal{B} \times \mathcal{B}$. In other words, we demonstrate that the complement of $O_{p,q}$, $O_{p,q}^c$, is closed. In fact, to do this, we consider a sequence $((x_k, y_k))_{k \in \mathbb{N}} \subset O_{p,q}^c$ that converges to $(x, y) \in \mathcal{B} \times \mathcal{B}$. Since $((x_k, y_k))_{k \in \mathbb{N}} \subset O_{p,q}^c$, so

$$f((x_k)^p (y_k)^q) + (x_k)^p \circ (y_k)^q \in Z(\mathcal{B}) \text{ for all } k \in \mathbb{N}.$$

Given that f is a continuous maps, the sequence $(f((x_k)^p (y_k)^q) + (x_k)^p \circ (y_k)^q)_{k \in \mathbb{N}}$ converges to $f(x^p y^q) + x^p \circ y^q$, know that $Z(\mathcal{B})$ is closed, we obtain $f(x^p y^q) + x^p \circ y^q \in Z(\mathcal{B})$. As a result, $(x, y) \in O_{p,q}^c$ and $O_{p,q}^c$ is closed (i.e $O_{p,q}$ is open).

Whenever $O_{p,q}$ is dense, the Baire's category theorem informs us that their intersection is also dense, which conflicts with $(\cap O_{p,q}) \cap (\mathcal{H}_1 \times \mathcal{H}_2) = \emptyset$. Hence, there is $(n, m) \in \mathbb{N}^* \times \mathbb{N}^*$ such that $O_{n,m}$ is not a dense open and there exists a non-void open subset $\mathcal{O} \times \mathcal{O}'$ in $O_{p,q}^c$ such that:

$$f(x^n y^m) + x^n \circ y^m \in Z(\mathcal{B}) \text{ for all } x \in \mathcal{O}, y \in \mathcal{O}',$$

Fix $y \in \mathcal{O}'$. Let $x \in \mathcal{O}$ and $z \in \mathcal{B}$, then $x + tz \in \mathcal{O}$ for all sufficiently small real t . Therefore $P(t) = f((x + tz)^n y^m) + (x + tz)^n \circ y^m \in Z(\mathcal{B})$. We are capable of writing

$$P(t) = B_{n,0}(x, z, y) + tB_{n-1,1}(x, z, y) + t^2B_{n-2,2}(x, z, y) + \dots + t^n B_{0,n}(x, z, y).$$

By Lemma 1, we conclude that $B_{0,n}(x, z, y) = f(z^n y^m) + z^n \circ y^m \in Z(\mathcal{B})$. Accordingly, chosen to give $y \in \mathcal{O}'$ we have $f(x^n y^m) + x^n \circ y^m \in Z(\mathcal{B})$ for all $x \in \mathcal{B}$. We reverse the roles of \mathcal{O} and \mathcal{O}' in the above settings with x is fixed in \mathcal{B} , we find that $f(x^n y^m) + x^n \circ y^m \in Z(\mathcal{B})$ for all $(x, y) \in \mathcal{B}^2$.

Replacing x by x^m and y by y^n , we find that

$$f(x^{nm} y^{mn}) + x^{nm} \circ y^{mn} \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

By changing x and y , we obtain

$$f(y^{nm} x^{mn}) + y^{nm} \circ x^{mn} \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

Then

$$(f(x^{nm} y^{mn}) + x^{nm} \circ y^{mn}) - (f(y^{nm} x^{mn}) + y^{nm} \circ x^{mn}) \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

Since

$$y^{nm} \circ x^{mn} = x^{nm} \circ y^{mn} \text{ for all } x, y \in \mathcal{B},$$

we obtain

$$f([x^{nm}, y^{mn}]) \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

According to Theorem 2.2 of [11], we come to the conclusion that \mathcal{B} is commutative. \square

THEOREM 2. *Let \mathcal{H}_1 and \mathcal{H}_2 are non-void open subsets of a prime Banach algebra \mathcal{B} . If a continuous automorphism f of \mathcal{B} , satisfying the following condition:*

$$(\forall (x, y) \in \mathcal{H}_1 \times \mathcal{H}_2)(\exists (p, q) \in \mathbb{N}^* \times \mathbb{N}^* \text{ such that } f(x^p \circ y^q) + [x^p, y^q] \in Z(\mathcal{B})),$$

then \mathcal{B} is commutative.

Proof. In the same way as Theorem 1's proof, we arrive at the conclusion that there is $(n, m) \in \mathbb{N}^* \times \mathbb{N}^*$ such that

$$f(x^n \circ y^m) + [x^n, y^m] \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

If $y = x$, then $2f(x^{n+m}) \in Z(\mathcal{B})$ for all $x \in \mathcal{B}$, since \mathcal{B} is 2-torsion free therefore $f(x^{n+m}) \in Z(\mathcal{B})$ for all $x \in \mathcal{B}$. And if we replace x by $f^{-1}(x)$, we obtain $x^{n+m} \in Z(\mathcal{B})$ for all $x \in \mathcal{B}$ and \mathcal{B} is commutative (according to [8]). \square

APPLICATION 1. *Let \mathbb{R} be the field of real numbers. $\mathcal{B} = \mathcal{M}_2(\mathbb{R})$ endowed with usual's matrix addition and multiplication and the norm $\|\cdot\|_1$ defined by $\|A\|_1 = \sum_{1 \leq i, j \leq 2} |a_{i,j}|$ for*

all $A = (a_{i,j})_{1 \leq i, j \leq 2} \in \mathcal{B}$. Let f the automorphism of \mathcal{B} defined by $f(M) = AMA^{-1}$ where $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ (f is a continuous automorphism). Let \mathcal{H} be a nonempty open subset of \mathcal{B} included in $Z(\mathcal{B})$. For all $(A, B) \in \mathcal{H} \times \mathcal{H}$ and for all $(p, q) \in \mathbb{N}^ \times \mathbb{N}^*$ we have $A^p \in Z(\mathcal{B})$ and $B^q \in Z(\mathcal{B})$, then $f(A^p B^q) + A^p \circ B^q \in Z(\mathcal{B})$. By Theorem 1, we conclude that \mathcal{B} is commutative. Consequently $\mathcal{H} = \emptyset$.*

We conclude that the only open subset included in $Z(\mathcal{B})$ is the empty set.

APPLICATION 2. *Let E be a normed space over \mathbb{K} (\mathbb{R} or \mathbb{C}). The space $\mathcal{B} = \mathcal{L}_c(E)$ of continuous linear applications from E to E , endowed with addition and composition and the norm defined by $\|T\| = \sup_{\|x\| \leq 1} \|T(x)\|$ for all $T \in \mathcal{B}$, is a normed algebra over \mathbb{K} . Let*

G be the subspace of \mathcal{B} defined by $G = \{\lambda I_E \mid \lambda \in \mathbb{K}\}$, where I_E is the identity of E , we observe that $G \subset Z(\mathcal{B})$, according to Application 1 we conclude that the interior of G is empty because \mathcal{B} is not commutative and $\text{Int}(G) \subset \text{Int}(Z(\mathcal{B})) = \emptyset$.

The following theorem can be shown using the same techniques as the proof of Theorem 1.

THEOREM 3. Let \mathcal{B} be a prime Banach algebra, \mathcal{H}_1 and \mathcal{H}_2 two non-void open subsets of \mathcal{B} . If \mathcal{B} admits a continuous automorphism f satisfying one of the following conditions

- i. $(\forall (x, y) \in \mathcal{H}_1 \times \mathcal{H}_2)(\exists (p, q) \in \mathbb{N}^* \times \mathbb{N}^*)$ such that $f(x^p y^q) - x^p \circ y^q \in Z(\mathcal{B})$.
- ii. $(\forall (x, y) \in \mathcal{H}_1 \times \mathcal{H}_2)(\exists (p, q) \in \mathbb{N}^* \times \mathbb{N}^*)$ such that $f(x^p \circ y^q) - [x^p, y^q] \in Z(\mathcal{B})$.

Then \mathcal{B} is commutative.

APPLICATION 3. Let $n \geq 3$ be an integer and $\mathcal{B} = \mathcal{M}_n(\mathbb{K})$ where $\mathbb{K} (= \mathbb{R} \text{ or } = \mathbb{C})$. Let \mathcal{A} the set of all $n \times n$ strictly upper triangular real or complexes matrices. Endowed with the usual operations on matrices and the norm $\|\cdot\|_1$ defined by $\|A\|_1 = \sum_{1 \leq i, j \leq n} |a_{ij}|$ for all

$$A = (a_{i,j})_{1 \leq i, j \leq n} \in \mathcal{B}.$$

\mathcal{B} is a non-commutative real prime Banach algebra. Observe that

$$f(x^n y^n) - [x^n, y^n] = 0 \in Z(\mathcal{B}) \quad \forall (x, y) \in \mathcal{B}^2$$

where f is the continuous automorphism of \mathcal{B} defined by $f(M) = P^{-1}MP$ and $P \in \mathcal{B}$ is an invertible matrix. Hence, it follows from Theorem 1 that \mathcal{A} is not a open subset in \mathcal{B} .

COROLLARY 1. Let \mathcal{B} be a prime Banach algebra and D a dense part of \mathcal{B} . If \mathcal{B} admits a continuous automorphism f such that

$$\exists (p, q) \in \mathbb{N}^* \times \mathbb{N}^* : f(x^p \circ y^q) + [x^p, y^q] \in Z(\mathcal{B}) \text{ for all } x, y \in D,$$

then \mathcal{B} is commutative.

Proof. Let $x, y \in \mathcal{B}$, there exist two sequences $(x_k)_{k \in \mathbb{N}}$ and $(y_k)_{k \in \mathbb{N}}$ in D converging to x and y . Since $(x_k)_{k \in \mathbb{N}} \subset D$ and $(y_k)_{k \in \mathbb{N}} \subset D$, so

$$f((x_k)^p \circ (y_k)^q) + [(x_k)^p, (y_k)^q] \in Z(\mathcal{A}) \text{ for all } k \in \mathbb{N}.$$

Since d is continuous then the sequence $(f((x_k)^p \circ (y_k)^q) + [(x_k)^p, (y_k)^q])_{k \in \mathbb{N}}$ converges to $f(x^p \circ y^q) + [x^p, y^q]$, knowing that $Z(\mathcal{B})$ is closed, then $f(x^p \circ y^q) + [x^p, y^q] \in Z(\mathcal{B})$.

We conclude that :

$$\exists (p, q) \in \mathbb{N}^* \times \mathbb{N}^* : f(x^p \circ y^q) + [x^p, y^q] \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

By Theorem 2, we get the desired conclusion. \square

3. Generalized derivations in Banach algebras

In this section, we examine some analogous conclusions for generalized derivations in Banach algebras.

THEOREM 4. Let \mathcal{B} be a prime Banach algebra, \mathcal{H}_1 and \mathcal{H}_2 non-void open subsets of \mathcal{B} . If \mathcal{B} admits a continuous generalised derivation F associated with a non-injective derivation d satisfying for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ there are strictly positive integers p', q' such that $F(x^{p'} y^{q'}) + x^{p'} \circ y^{q'} \in Z(\mathcal{B})$, then \mathcal{B} is commutative or $d(Z(\mathcal{B})) = \{0\}$.

Proof. If $F = 0$, then our hypothesis reduces to $x^{p'} \circ y^{q'} \in Z(\mathcal{B})$ for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ so the required result follows from Theorem 1 [8]. Hence we may suppose that $F \neq 0$. For all $(p', q') \in \mathbb{N}^* \times \mathbb{N}^*$ let us define the following sets :

$$O_{p', q'} = \{(x, y) \in \mathcal{B}^2 \mid F(x^{p'} y^{q'}) + x^{p'} \circ y^{q'} \notin Z(\mathcal{B})\} \text{ and } F_{p', q'} = \{(x, y) \in \mathcal{B}^2 \mid F(x^{p'} y^{q'}) + x^{p'} \circ y^{q'} \in Z(\mathcal{B})\}.$$

Using a similar approach with Theorem 1, we arrive at (where $(n, m) \in \mathbb{N}^* \times \mathbb{N}^*$)

$$F(x^n y^m) + x^n \circ y^m \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}^2.$$

Changing x by x^m and y by y^n , We conclude that

$$F(x^{nm}y^{mn}) + x^{nm} \circ y^{mn} \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

By permuting x and y , we find that

$$F(y^{nm}x^{mn}) + y^{nm} \circ x^{mn} \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

Then

$$(F(x^{nm}y^{mn}) + x^{nm} \circ y^{mn}) - (F(y^{nm}x^{mn}) + y^{nm} \circ x^{mn}) \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

Since

$$y^{nm} \circ x^{mn} = x^{nm} \circ y^{mn} \text{ for all } x, y \in \mathcal{B},$$

we obtain

$$F([x^{nm}, y^{nm}]) \in Z(\mathcal{B}) \text{ for all } x, y \in \mathcal{B}.$$

Let $x \in \mathcal{B}$ and $a = x^{nm}$. We have $P(t) = F([a, (y + ta)^{nm}]) \in Z(\mathcal{B})$ for all $(y, t) \in \mathcal{B} \times \mathbb{R}$. This can be written as

$$P(t) = F([a, (y + ta)^{nm}]) = \sum_{k=0}^{mn} t^k F([a, A_{nm-k,k}]),$$

where $A_{nm-k,k}$ denotes the sum of all terms in which y appears exactly $nm - k$ times and a appears exactly k times. By Lemma 1, we conclude that $F([a, A_{nm-k,k}(x, y)]) \in Z(\mathcal{B})$ for all $0 \leq k \leq nm$. The coefficient of t in this polynomial is $F([a, A_{nm-1,1}])$ and $A_{nm-1,1} =$

$$\sum_{k=0}^{nm-1} a^{nm-1-k} y a^k. \text{ Since}$$

$$\begin{aligned} \sum_{k=0}^{nm-1} [a, a^{nm-1-k} y a^k] &= [a, a^{nm-1} y] + [a, a^{nm-2} y a] + [a, a^{nm-3} y a^2] + \dots + [a, y a^{nm-1}] \\ &= (a a^{nm-1} y - a^{nm-1} y a) + (a a^{nm-2} y a - a^{nm-2} y a a) + \dots + (a y a^{nm-1} - y a^{nm-1} a) \\ &= (a^{nm} y - a^{nm-1} y a) + (a^{nm-1} y a - a^{nm-2} y a^2) + \dots + (a y a^{nm-1} - y a^{nm}) \\ &= a^{nm} y - y a^{nm} \\ &= [a^{nm}, y]. \end{aligned}$$

Then

$$F([a, A_{nm-1,1}(x, y)]) = \sum_{k=0}^{nm-1} F([a, x^{nm-1-k} y x^k]) = F([a^{nm}, y]) \in Z(\mathcal{B}).$$

Thus

$$F([x^{n^2 m^2}, y]) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}^2.$$

Now, we prove that the restriction of d on $Z(\mathcal{B})$ is not injective.

For this, we consider $a \in \mathcal{B}$ such that $d(a) = 0$ and b be a non-zero element of $Z(\mathcal{A})$. For all $t \in \mathbb{R}$, we have

$$S(t) = F[(a + tb)^p, y^p] \in Z(\mathcal{B}) \text{ for all } y \in \mathcal{B},$$

where $p = n^2 m^2$. By reason of $b^k \in Z(\mathcal{B})$ ($\forall k \in \mathbb{N}^*$), we can write

$$S(t) = \sum_{k=0}^p \binom{k}{p} t^k F([a^{p-k} b^k, y^p]) = \sum_{k=0}^p t^k A_k(a, b, y) \in Z(\mathcal{B}) \text{ for all } y \in \mathcal{B}$$

where $A_k(a, b, y) = \binom{k}{p} F([a^{p-k} b^k, y^p])$, according to Lemma 1, we conclude that $A_k(a, b, y) \in Z(\mathcal{B})$ for all $0 \leq k \leq p$. In particular for $k = p - 1$, we get

$$p F([a b^{p-1}, y^p]) \in Z(\mathcal{B}),$$

we can reduce by p (because $Z(\mathcal{B})$ is a subspace of \mathcal{B}) and we obtain

$$F([ab^{p-1}, y^p]) \in Z(\mathcal{B}),$$

therefore $F(b^{p-1}[a, y^p]) \in Z(\mathcal{B})$ (because b^{p-1} is belongs to $Z(\mathcal{B})$). Since

$$F(b^{p-1}[a, y^p]) = b^{p-1}F([a, y^p]) + [a, y^p]d(b^{p-1}) \text{ for all } y \in \mathcal{B}$$

and $b^{p-1}F([a, y^p])$ is an element of $Z(\mathcal{B})$, then $d(b^{p-1})[a, y^p] \in Z(\mathcal{B})$ for all $y \in \mathcal{B}$. As $d(b^{p-1})$ is a non-zero element of $Z(\mathcal{B})$, by Fact 1, we conclude that

$$[a, y^p] \in Z(\mathcal{B}) \text{ for all } y \in \mathcal{B}.$$

Therefore, for all $t \in \mathbb{R}$ we have

$$Q(t) = [a, (a + ty)^p] \in Z(\mathcal{B}) \text{ for all } y \in \mathcal{B}.$$

That is

$$Q(t) = \sum_{k=0}^p t^k [a, B_{p-k,k}(a, y)] \text{ for all } y \in \mathcal{B}$$

where $B_{p-k,k}(a, y)$ denotes the sum of all terms in which a appears exactly $p-k$ times and y appears exactly k times. By Lemma 1, we deduce that $[a, B_{p-k,k}(a, y)] \in Z(\mathcal{B})$ for all $k \leq p$. The coefficient of t in this polynomial is $[a, B_{p-1,1}(a, y)]$ where $B_{p-1,1}(a, y) = \sum_{k=0}^{p-1} a^{p-1-k} y a^k$, therefore $[a, B_{p-1,1}(a, y)] = \sum_{k=0}^{p-1} [a, a^{p-1-k} y a^k] \in Z(\mathcal{B})$, we observe that

$$\begin{aligned} \sum_{k=0}^{p-1} [a, a^{p-1-k} y a^k] &= [a, a^{p-1} y] + [a, a^{p-2} y a] + [a, a^{p-3} y a^2] + \dots + [a, y a^{p-1}] \\ &= a a^{p-1} y - a^{p-1} y a + a a^{p-2} y a - a^{p-2} y a a + \dots + a y a^{p-1} - y a^{p-1} a \\ &= a^p y - a^{p-1} y a + a^{p-1} y a - a^{p-2} y a^2 + \dots + a y a^{p-1} - y a^p \\ &= a^p y - y a^p \\ &= [a^p, y]. \end{aligned}$$

Consequently,

$$[a^p, y] \in Z(\mathcal{B}) \text{ for all } y \in \mathcal{B}.$$

There are 2 cases:

Case 1: : If $a^p \in Z(\mathcal{B})$, since d is injective in $Z(\mathcal{B})$ and $d(a^p) = 0$ then $a^p = 0$, therefore $a = 0$ and d is injective.

Case 2: : If $a^p \notin Z(\mathcal{B})$ then $d(\mathcal{B}) \subset \mathcal{B}$ where d is the inner derivation associated to a^p , by Fact 1 we have $\mathcal{B} \subset Z(\mathcal{B})$, so d is injective.

Lastly, we conclude that the restriction of d on $Z(\mathcal{B})$ is non-injective, then there is a nonzero element a of $Z(\mathcal{B})$ such that $d(a) = 0$, we have $F([(x + ta)^p, y]) \in Z(\mathcal{B})$ for all $(x, y) \in \mathcal{B}$ and $t \in \mathbb{R}$. Since $a^k \in Z(\mathcal{B})$ for all $k \in \mathbb{N}^*$, we can write

$$F([(x + ta)^p, y]) = \sum_{k=0}^p \binom{p}{k} t^k F([a^{p-k} x^k, y]) \in Z(\mathcal{B}),$$

according to Lemma 1, we conclude that $\binom{p}{k} F([a^{p-k} x^k, y]) \in Z(\mathcal{B})$ for all $0 \leq k \leq p$. In particular for $k = 1$, we have $pF([a^{p-1} x, y]) \in Z(\mathcal{B})$ and $F([a^{p-1} x, y]) \in Z(\mathcal{B})$, since $a^{p-1} \in Z(\mathcal{B})$ we have $F(a^{p-1}[x, y]) \in Z(\mathcal{B})$, then $a^{p-1}F([x, y]) \in Z(\mathcal{B})$ (because $d(a^{p-1}) = 0$). So $a^{p-1} \in Z(\mathcal{B}) \setminus \{0\}$, according to Fact 1 we have

$$F([x, y]) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

If $d(Z(\mathcal{B})) \neq \{0\}$, then, there is $z \in Z(\mathcal{B})$ such that $d(z) \neq 0$ and we have

$$F([x, yz]) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

That is

$$F([x, y]z) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

And

$$F([x, y]z) + [x, y]d(z) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

Since

$$F([x, y]z) \in Z(\mathcal{B})$$

therefore

$$[x, y]d(z) \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}$$

by Fact 1 we conclude that

$$[x, y] \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

Replacing y be xy , we obtain

$$x[x, y] \in Z(\mathcal{B}) \text{ for all } (x, y) \in \mathcal{B}.$$

Then $x \in Z(\mathcal{B})$ or $[x, y] = 0$, both shows that \mathcal{B} is commutative. \square

We show the following result by traveling on the same path.

THEOREM 5. *Consider \mathcal{H}_1 and \mathcal{H}_2 as nonvoid open subsets of a prime Banach algebra \mathcal{B} . If \mathcal{B} admits a continuous generalised derivation F associated with a non-injective derivation d satisfying for all $(x, y) \in \mathcal{H}_1 \times \mathcal{H}_2$ there are strictly positive integers p and q such that $F(x^p \circ y^q) + [x^p, y^q] \in Z(\mathcal{B})$, then \mathcal{B} is commutative or $d(Z(\mathcal{B})) = \{0\}$.*

4. Some examples

The example that follows demonstrates that our theorems' assertion that " \mathcal{H}_1 and \mathcal{H}_2 are open" is not superfluous.

EXAMPLE 1. Let $\mathcal{B} = \mathcal{M}_2(\mathbb{R})$ the ring of square matrices with real inputs of order 2×2 endowed with usual's matrix addition and multiplication and the norm defined by

$$\|A\|_1 = \sum_{1 \leq i, j \leq 2} |a_{i, j}| \text{ for all } A = (a_{i, j})_{1 \leq i, j \leq 2} \in \mathcal{B}, \text{ is a real prime Banach algebra.}$$

$$\text{Let } \mathcal{F}_1 = \left\{ \begin{pmatrix} t & 0 \\ 0 & t \end{pmatrix} \mid t \in \mathbb{R} \right\} \text{ and } \mathcal{F}_2 = \left\{ \begin{pmatrix} t & 0 \\ 0 & t \end{pmatrix} \mid t \in \mathbb{R}^+ \right\}.$$

\mathcal{F}_1 is not open in \mathcal{B} , indeed, we have to show that the complement of \mathcal{F}_1 is not closed.

For this, we consider the sequence $\left(\begin{pmatrix} 1 + \frac{1}{n} & \frac{-1}{n} \\ \frac{1}{n} & 1 + \frac{1}{n} \end{pmatrix} \right)_{n \in \mathbb{N}^*}$ in \mathcal{F}_1^c complement of \mathcal{F}_1 who

converge to $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \notin \mathcal{F}_1^c$, then \mathcal{F}_1^c is not closed, that is \mathcal{F}_1 is not open in \mathcal{B} .

Let $A = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in \mathcal{F}_1$ and $B = \begin{pmatrix} b & 0 \\ 0 & b \end{pmatrix} \in \mathcal{F}_2$. For all $(p, q) \in \mathbb{N}^2$, we have

$$A^p = \begin{pmatrix} a^p & 0 \\ 0 & a^p \end{pmatrix} \text{ and } B^q = \begin{pmatrix} b^q & 0 \\ 0 & b^q \end{pmatrix}.$$

So

$$A^p B^q = \begin{pmatrix} a^p b^q & 0 \\ 0 & a^p b^q \end{pmatrix}, A^p \circ B^q = \begin{pmatrix} 2a^p b^q & 0 \\ 0 & 2a^p b^q \end{pmatrix} \text{ and } [A^p, B^q] = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

We conclude that

$$Id(A^p \circ B^q) = Id([A^p, B^q]) = Id(A^p B^q) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

(where Id is the identity application of \mathcal{B}). We obtain:

1. $Id(A^p B^q) + A^p \circ B^q \in Z(\mathcal{B})$ and $Id(A^p B^q) - A^p \circ B^q \in Z(\mathcal{B})$,
2. $Id(A^p \circ B^q) + [A^p, B^q] \in Z(\mathcal{B})$ and $Id(A^p \circ B^q) - [A^p, B^q] \in Z(\mathcal{B})$.

But \mathcal{B} is not commutative.

The example that follows demonstrates that we cannot change \mathbb{R} or \mathbb{C} by $\mathbb{F}_3 = \mathbb{Z}/3\mathbb{Z}$ in the statement of previous theorems.

EXAMPLE 2. Let $\mathcal{B} = (\mathcal{M}_2(\mathbb{Z}/3\mathbb{Z}), +, \times, \cdot)$ the 2-torsion free prime Banach algebra of square matrices of size 2 at coefficients in $\mathbb{Z}/3\mathbb{Z}$ with usual matrix addition and matrix multiplication.

The norm is defined by $\|A\|_1 = \sum_{1 \leq i, j \leq 2} |a_{i,j}|$ for all $A = (a_{i,j})_{1 \leq i, j \leq 2} \in \mathcal{B}$ with $|\cdot|$ is the norm

defined on $\mathbb{Z}/3\mathbb{Z}$ by:

$$|\bar{0}| = 0, |\bar{1}| = 1 \text{ and } |\bar{2}| = 2.$$

Observe that $\mathcal{H} = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \mid a \in \mathbb{Z}/3\mathbb{Z} \right\}$ is open in \mathcal{B} . Indeed, let $A \in \mathcal{H}$ the open ball $B(A, 1) = \{X \in \mathcal{B} \text{ such that } \|A - X\|_\infty < 1\} = \{A\} \subset \mathcal{H}$, then \mathcal{H} is a non-void open subset of \mathcal{B} . For all $(p, q) \in \mathbb{N}^* \times \mathbb{N}^*$ and for all $(A, B) \in \mathcal{H} \times \mathcal{H}$ we have :

1. $Id(A^p B^q) + A^p \circ B^q \in Z(\mathcal{B})$ and $Id(A^p B^q) - A^p \circ B^q \in Z(\mathcal{B})$.
2. $Id(A^p \circ B^q) + [A^p, B^q] \in Z(\mathcal{B})$ and $Id(A^p \circ B^q) - [A^p, B^q] \in Z(\mathcal{B})$.

So all the conditions are verified except $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$, but \mathcal{B} is not commutative.

References

- [1] N. Divinsky, On commuting automorphisms of rings, Trans. Roy. Soc. Canada. Sect. III. (3) 49 (1955), 19-22.
- [2] F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer-Verlag, New York, (1973).
- [3] I. N. Herstein, *A remark on rings and algebras*, Michigan Math. J. 10 (1963), 269-272.
- [4] J. Mayne, *Centralizing automorphism of prime rings*, Canad. Math. Bull. 19 (1976), 113-115.
- [5] J. H. Mayne, *Centralizing mappings of prime rings*, Canad. Math. Bull. 2 (1984), no. 1, 122-126
- [6] E.C.Posner, *Derivations in Prime rings*, Proc. Amer. Math. Soc., 8 (1957), 1093-1100.
- [7] B. Yood, *Some commutativity theorems for Banach algebras*, Publ. Math. Debrecen 45 (1994), no. 1-2, 29-33.
- [8] M. Moumen, L. Taoufiq and L. Oukhtite, *Some differential identities on prime Banach algebras*, Journal of Algebra and Its Applications(2022), doi:10.1142/S0219498823502584.
- [9] M. Moumen, L. Taoufiq and A. Boua, *On prime Banach algebras with continuous derivations*, Mathematica, 65(88), n 1, 2023, pp. 122-132..
- [10] M. Moumen and L. Taoufiq, *Commutativity Theorems and projection on the center of a Banach algebra*, Tatra Mt. Math. Publ. 83 (2023), 119–130, DOI: 10.2478/temp-2023-0009.
- [11] M. Moumen and L. Taoufiq, *On Commutativity of Banach algebra with endomorphisms*, Palestine Journal of Mathematics.Vol. 12(2023): 100-105.

National School of Applied Sciences

Department of Mathematics

Ibn Zohr University

Agadir, Morocco

E-mail: mohamed.moumen@edu.uiz.ac.ma

National School of Applied Sciences

Department of Mathematics

Ibn Zohr University

Agadir, Morocco

E-mail: l.taoufiq@uiz.ac.ma