# G-APPROXIMATE BEST PROXIMITY PAIRS IN METRIC SPACE WITH A DIRECTED GRAPH

S. A. M. MOHSENIALHOSSEINI\*, M. SAHELI

\* Faculty of Mathematics, Vali-e-Asr University, Rafsanjan, Iran. e-mail: amah@vru.ac.ir

Faculty of Mathematics, Vali-e-Asr University, Rafsanjan, Iran. e-mail: saheli@vru.ac.ir

Copyright © 2020 the authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. Let (X,d) be a metric space that has a directed graph G such that the sets V(G) and E(G) are respectively vertices and edges corresponding to X. We obtain sufficient conditions for the existence of an G-approximate best proximity pair of the mapping T in the metric space X endowed with a graph G such that the set V(G) of vertices of G coincides with X.

**Keywords:** *G*-approximate best proximity pairs, *G*-approximate best minimizing sequences, Connected graph, *GT*-minimizing.

2010 AMS Subject Classification: 47H10, 54H25, 46B20.

#### **1.** INTRODUCTION

Fixed point theory is a very popular tool in solving existence problems in many branches of Mathematical Analysis and its applications. In physics and engineering fixed point technique has been used in areas like image retrieval, signal processing and the study of existence and uniqueness of solutions for a class of nonlinear integral equations.

Graphs can be used to model many types of relations and processes in physical, biochemistry, electrical engineering, computer science and operations research, biological. The wide scope of these and other applications has been well-documented cf. [5, 14]

<sup>\*</sup>Corresponding author

E-mail address: amah@vru.ac.ir; mohsenialhosseini@gmail.com

Received May 21, 2011

Let X be a metric space and A and B nonempty subsets of X and d(A,B) is distance of A and B. If  $d(x_0, y_0) = d(A,B)$  then the pair  $(x_0, y_0)$  is called a best proximity pair for A and B and put

$$prox(A,B) := \{(x,y) \in A \times B : d(x,y) = d(A,B)\}$$

as the set of all best proximity pair (A,B). Best proximity pair evolves as a generalization of the concept of best approximation. That reader can find some important result of it in [6, 9, 10].

Now, as in [15] (see also[1, 2, 20, 21, 22, 23]), we can find the best proximity points of the sets A and B, by considering a map  $T : A \cup B \to A \cup B$  such that  $T(A) \subseteq B$  and  $T(B) \subseteq A$ . Best proximity pair also evolves as a generalization of the concept of fixed point of mappings. Because if  $A \cap B \neq \emptyset$  every best proximity point is a fixed point of *T*.

In 2011, Mohsenalhosseini et al [12], introduced to approximate best proximity pairs in metric space. In 2013, Mohsenalhosseini et al [11], introduced to approximate best proximity pairs in metric space for contraction maps.

Recently, two results have appeared, giving suficient conditions for f to be a Picard operator if (X,d) is endowed with a graph. The first result in this direction was given by J. Jakhymski [8] who also presented its applications to the Kelisky-Rivlin theorem on iterates of the Bernstein operators on the space C[0,1].

Reich [18], Ćirić [19] and Rus [17] proved that if (X, d) is complete, then every Ćirić– Reich–Rus operator has a unique fixed point.

In [3], the author considered the problem of existence of a fixed point for  $\varphi$ -contractions in metric spaces endowed with a graph. Also in [4], introduced to Fixed point theorems for Reich type contractions on metric spaces with a graph.

The aim of this paper is to study the existence of approximate best proximity pair for a map and two maps and their diameters for a cyclic map  $T: A \cup B \cup C \rightarrow A \cup B \cup C$  i.e.  $T(A) \subseteq B, T(B) \subseteq C$  and  $T(C) \subseteq A$  in metric spaces endowed with a graph G by defining the G-approximate best proximity pair. Moreover, we give some illustrative example of our main results.

#### **2. PRELIMINARIES**

This section recalls the following notations and the ones that will be used in what follows. We refer to Mohsenalhosseini et al [12, 11] for a detailed study of approximate best proximity pair for a map and two maps, also refer to Wayne Goddard et al [7] for a detailed study of Distance in Graphset.

Graphs are special examples of metric spaces with their intrinsic path metric. Now we express the relationship between distance, diameter and radius in the graph with the metric space. A path in a graph is a sequence of distinct vertices, such that adjacent vertices in the sequence are adjacent in the graph. For an unweighted graph, the length of a path is the number of edges on the path. For an (edge) weighted graph, the length of a path is the sum of the weights of the edges on the path. We assume that all weights are nonnegative and that all graphs are connected.

**Definition 2.1.** [13] Let  $T : X \to X$ ,  $\varepsilon > 0$ ,  $x_0 \in X$ . Then  $x_0 \in X$  is an  $\varepsilon$ -fixed point for T if  $d(x_0, Tx_0) < \varepsilon$ .

**Definition 2.2.** [13] Let  $T : X \to X$ . Then *T* has the approximate fixed point property (a.f.p.p) if

$$\forall \varepsilon > 0, F_{\varepsilon}(T) \neq \emptyset.$$

**Theorem 2.3.** [11] Let  $(X, \|.\|)$  be a complete norm space,  $T : X \to X$ ,  $x_0 \in X$  and  $\varepsilon > 0$ . If  $\|T^n(x_0) - T^{n+k}(x_0)\| \to 0$  as  $n \to \infty$  for some k > 0, then  $T^k$  has an  $\varepsilon$ -fixed point.

**Definition 2.4.** [8] We say that a mapping  $T : X \to X$  is a *G*-contraction or simply *G*-contraction if *T* preserves edges of *G*, i.e.,

(2.1) 
$$\forall x, y \in X((x, y) \in E(G) \Rightarrow (Tx, Ty) \in E(G)),$$

and T decreases weights of edges of G in the following way:

(2.2) 
$$\exists \alpha \in (0,1) \ \forall x, y \in X \ ((x,y) \in E(G) \Rightarrow d(Tx,Ty) \le \alpha d(x,y)).$$

### **3.** MAIN RESULTS

Throughout this section, we assume that *G* is a directed graph such that V(G) = X,  $E(G) \supseteq \triangle$ such that  $\triangle$  denote the diagonal of the Cartesian product  $X \times X$  and the graph *G* has no parallel edges. In this section, by using the idea of Jachymski [8], we will consider the existence of *G*-approximate best proximity points for the map  $T : A \cup B \rightarrow A \cup B$ , such that  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ , and its diameter.

**Definition 3.1.** Let *A* and *B* be nonempty subsets of a metric space endowed with a directed graph G = (V(G), E(G)) such that  $V(G) = A \cup B$ . The operator  $T : A \cup B \rightarrow A \cup B$  be a map such that  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ . The point  $x \in A \cup B$  is said to be a *G*-approximate best proximity point of the pair (A, B), if:

i)  $((x, y) \in E(G) \Rightarrow (Tx, Ty) \in E(G)), \forall x, y \in A \cup B;$ ii)  $\exists \varepsilon > 0 \ \forall x \in A \cup B \ ((x, Tx) \in E(G) \Rightarrow d(x, Tx) \le d(A, B) + \varepsilon).$ 

*Remark* 3.2. In this paper we will denote the set of all *G*-approximate best proximity point of the pair (A, B) of *T*, for a given  $\varepsilon$ , by :

$$P_T^{Ga}(A,B) = \{ x \in A \cup B : ((x,Tx) \in E(G) \Rightarrow d(x,Tx) \le d(A,B) + \varepsilon) \}.$$

**Example 3.3.** Suppose Let  $X = \mathbb{R}^2$  and  $A = \{(x, y) \in X : (x - y)^2 + y^2 \le 1\}$  and  $B = \{(x, y) \in X : (x + y)^2 + y^2 \le 1\}$  with T(x, y) = (-x, y) for  $(x, y) \in X$ . Define the graph *G* by V(G) = X. Then there exists  $((x, y), T(x, y)) \in E(G)$  such that  $d((x, y), T(x, y)) \le d(A, B) + \varepsilon$  for some  $\varepsilon > 0$ . Hence  $P_T^{Ga}(A, B) \neq \emptyset$ .

We say that the pair (A,B) is a *G*-approximate best proximity pair if  $P_T^{Ga}(A,B) \neq \emptyset$ .

**Proposition 3.4.** Let A and B be nonempty subsets of a metric space endowed with a directed graph G = (V(G), E(G)) such that  $V(G) = A \cup B$ . suppose that the mapping  $T : A \cup B \to A \cup B$ satisfying  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ . If  $\lim_{n\to\infty} d(T^nx, T^{n+1}x) = d(A, B)$ , for some  $x \in A \cup B$  satisfies the condition  $(x, Tx) \in E(G)$  then the pair (A, B) is a G-approximate best proximity pair.

**Proof:** Let  $\varepsilon > 0$  be given and  $x \in A \cup B$  with  $(x, Tx) \in E(G)$  such that  $\lim_{n \to \infty} d(T^n x, T^{n+1}x) = d(A, B)$ ; then there exists  $N_0 > 0$  such that for all  $n \ge N_0$ ,

$$d(T^n x, T^{n+1} x) < d(A, B) + \varepsilon.$$

If  $n = N_0$ , then  $d(T^{N_0}(x), T(T^{N_0}(x))) < d(A, B) + \varepsilon$ , then  $T^{N_0}(x) \in P_T^{Ga}(A, B)$  and  $P_T^{Ga}(A, B) \neq \emptyset$ .

**Definition 3.5.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G*. The operator  $T : A \cup B \to A \cup B$  satisfying  $T(A) \subseteq B$ ,  $T(B) \subseteq A$  is said to be a G-ciric-Rich-Rus-Moh operator if:

i)  $((x, y) \in E(G) \Rightarrow (Tx, Ty) \in E(G)), \forall x, y \in A \cup B;$ 

ii) there exists nonnegative numbers  $\alpha, \beta, \gamma$  with  $\alpha + 2\beta + \gamma < 1$ , such that, for each  $(x, y) \in E(G)$ , we have:

$$d(Tx,Ty) \le \alpha d(x,y) + \beta [d(x,Tx) + d(y,Ty)] + \gamma d(A,B)$$

**Theorem 3.6.** Let A and B be nonempty subsets of a metric space endowed with a graph G and  $T : A \cup B \rightarrow A \cup B$  be a G-ciric-Rich-Rus-Moh operator. If  $x \in A \cup B$  satisfies the condition  $(x, Tx) \in E(G)$  then the pair (A, B) is a G-approximate best proximity pair.

**Proof:** Let  $x \in A \cup B$  with  $(x, Tx) \in E(G)$ , then

$$d(Tx, T^2x) \le \alpha d(x, Tx) + \beta [d(x, Tx) + d(Tx, T^2x)] + \gamma d(A, B)$$

Therefore

$$d(Tx, T^2x) \leq \frac{\alpha + \beta}{1 - \beta} d(x, Tx) + \frac{\gamma}{1 - \beta} d(A, B).$$

Now if  $k = \frac{\alpha + \beta}{1 - \beta}$ , then

$$d(Tx, T^2x) \le kd(x, Tx) + (1-k)d(A, B)$$

also

$$d(T^{2}x, T^{3}x) \leq k^{2}d(x, Tx) + (1 - k^{2})d(A, B).$$

Therefore

$$d(T^{n}x, T^{n+1}x) \le k^{n}d(x, Tx) + (1-k^{n})d(A, B),$$

and so

$$d(T^nx,T^{n+1}x) \to d(A,B), as n \to \infty.$$

Therefore, by Theorem 3.4,  $P_T^{Ga}(A,B) \neq \emptyset$ , then pair(*A*,*B*) is a *G*-approximate best proximity pair.

**Definition 3.7.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G*. Suppose that the mapping  $T : A \cup B \to A \cup B$  satisfying  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ . We say that the sequence  $\{z_n\} \subseteq A \cup B$  with  $(z_n, Tz_n) \in E(G)$  is *GT*-minimizing if

$$\lim_{n\to\infty} d(z_n, Tz_n) = d(A, B).$$

**Theorem 3.8.** Let A and B be nonempty subsets of a metric space endowed with a graph G, suppose that the mapping  $T : A \cup B \to A \cup B$  satisfying  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ . If  $\{T^nx\}$  is a GT-minimizing for some  $x \in A \cup B$  satisfies the condition  $(x, Tx) \in E(G)$ , then (A, B) is a G-approximate best pair proximity.

**Proof.** Since

$$\lim_{n \to \infty} d(T^n x, T^{n+1} x) = d(A, B)$$

for some  $x \in A \cup B$  satisfies the condition  $(x, Tx) \in E(G)$ , then by Theorem 3.4  $P_T^{Ga}(A, B) \neq \emptyset$ . Therefore pair (A, B) is a *G*-approximate best proximity pair.

**Theorem 3.9.** Let A and B be nonempty subsets of a metric space endowed with a graph G such that E(G) is compact. Suppose that the mapping  $T : A \cup B \to A \cup B$  satisfying  $T(A) \subseteq B$ ,  $T(B) \subseteq A$ , T is continuous and  $||Tx - Ty|| \leq ||x - y||$ , where  $(x, y) \in E(G)$ . Then  $P_T^{Ga}(A, B)$  is nonempty and compact.

**Proof.** Since E(G) compact, there exists a  $z_0 \in E(G)$  such that

(3.1) 
$$||z_0 - Tz_0|| = \inf_{z \in E(G)} ||z - Tz||$$

If  $||z_0 - Tz_0|| > d(A, B)$ , then  $||Tz_0 - T^2z_0|| < ||z_0 - Tz_0||$  which contradict to the definition of  $z_0$ ,  $(Tz_0 \in E(G))$  and by 3.1. Therefore  $||z_0 - Tz_0|| = d(A, B) \le d(A, B) + \varepsilon$  for some  $\varepsilon > 0$  and  $z_0 \in P_T^{Ga}(A, B)$ . Therefore  $P_T^{Ga}(A, B)$  is nonempty.

Also, if  $\{z_n\} \subseteq P_T^{Ga}(A,B)$ , then  $||z_n - Tz_n|| < d(A,B) + \varepsilon$ , for some  $\varepsilon > 0$ , and by compactness of E(G), there exists a subsequence  $z_{n_k}$  and a  $z_0 \in E(G)$  such that  $z_{n_k} \to z_0$  and so

$$||z_0 - Tz_0|| = \lim_{k \to \infty} ||z_{n_k} - Tz_{n_k}|| < d(A, B) + \varepsilon$$

for some  $\varepsilon > 0$  hence  $P_T^{Ga}(A, B)$  is compact.

**Example 3.10.** If A = [-3, -1], B = [1, 3] and  $T : A \cup B \to A \cup B$  such that

$$T(x) = \begin{cases} \frac{1-x}{2} & x \in A \\ \\ \frac{-1-x}{2} & x \in B \end{cases}$$

Then  $P_T^{Ga}(A, B)$  is compact, we have

$$P_T^{Ga}(A,B) = \{x \in A \cup B : d(x,Tx) < d(A,B) + \varepsilon \text{ for some } \varepsilon > 0\}$$
$$= \{x \in A \cup B : d(x,Tx) < 2 + \varepsilon \text{ for some } \varepsilon > 0\}$$
$$= \{1,-1\}.$$

that is compact.

In the following, by  $diam(P_T^{Ga}(A,B))$  for a set  $P_T^{Ga}(A,B) \neq \emptyset$  we will understand the diameter of the set  $P_T^{Ga}(A,B)$ .

**Definition 3.11.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G* such that  $T(A) \subseteq B$ ,  $T(B) \subseteq A$  and  $\varepsilon > 0$ . We define diameter  $P_T^{Ga}(A, B)$  by

$$(3.2) \qquad \qquad diam(P_T^{Ga}(A,B)) = \sup\{d(x,y): x, y \in P_T^{Ga}(A,B)\}.$$

**Theorem 3.12.** Let A and B be nonempty subsets of a metric space endowed with a graph G. suppose that the mapping  $T : A \cup B \to A \cup B$ , such that  $T(A) \subseteq B$ ,  $T(B) \subseteq A$  and  $\varepsilon > 0$ . If T be a G-contraction then

$$diam(P_T^{Ga}(A,B)) \leq \frac{2\varepsilon}{1-\alpha} + \frac{2d(A,B)}{1-\alpha}.$$

**Proof.** If  $x, y \in P_T^{Ga}(A, B)$ , then

$$d(x,y) \leq d(x,Tx) + d(Tx,Ty) + d(Ty,y)$$
  
$$\leq \varepsilon_1 + \alpha d(x,y) + 2d(A,B) + \varepsilon_2.$$

put  $\varepsilon = Max\{\varepsilon_1, \varepsilon_2\}$ , Therefore  $d(x, y) \le \frac{2\varepsilon}{1-\alpha} + \frac{2d(A,B)}{1-\alpha}$ . Hence  $diam(P_T^{Ga}(A,B)) \le \frac{2\varepsilon}{1-\alpha} + \frac{2d(A,B)}{1-\alpha}$ .

In this section we will consider the existence of *G*-approximate best proximity points for two maps  $T : A \cup B \rightarrow A \cup B$ ,  $S : A \cup B \rightarrow A \cup B$ , and its diameter.

**Definition 3.13.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G* and  $T: A \cup B \to A \cup B$ ,  $S: A \cup B \to A \cup B$  be two maps such that  $T(A) \subseteq B$ ,  $S(B) \subseteq A$ . A point (x, y) in  $A \times B$  is said to be a *G*-approximate-pair fixed point for (T, S), if: i)  $\forall (x, y) \in A \times B$   $((x, y) \in E(G) \Rightarrow (Tx, Ty) \in E(G), (Sx, Sy) \in E(G));$ ii)  $\exists \alpha \in (0, 1) \forall (x, y) \in A \times B$   $((x, y) \in E(G) \Rightarrow d(Tx, Sy) \leq d(A, B) + \varepsilon)$ .

We say that the pair (T, S) has the *G*-approximate-pair fixed property in X if  $P_{(T,S)}^{Ga}(A,B) \neq \emptyset$ , where

$$P^{Ga}_{(T,S)}(A,B) = \{(x,y) \in E(G) : d(Tx,Sy) \le d(A,B) + \varepsilon \text{ for some } \varepsilon > 0\}.$$

**Theorem 3.14.** Let A and B be nonempty subsets of a metric space endowed with a graph G and  $T : A \cup B \to A \cup B$ ,  $S : A \cup B \to A \cup B$  be two maps such that  $T(A) \subseteq B$ ,  $S(B) \subseteq A$ . If, for every  $(x, y) \in A \times B$ ,

$$d(T^n(x), S^n(y)) \to d(A, B),$$

then (T,S) has the G-approximate-pair fixed property.

**Proof.** Let  $\varepsilon > 0$  be given and  $(x, y) \in A \times B$  with  $((x, y), (Tx, Ty)) \in E(G)$ . Since  $d(T^n(x), S^n(y)) \rightarrow d(A, B)$ , there existe  $n_0 > 0$  such that for all  $n \ge n_0$ ,

$$d(T^n(x), S^n(y)) < d(A, B) + \varepsilon.$$

Then  $d(T(T^{n-1}(x), S(S^{n-1}(y)) < d(A, B) + \varepsilon \text{ for every } n \ge n_0$ . Put  $x_0 = T^{n_0 - 1}(x)$  and  $y_0 = S^{n_0 - 1}(y)$ . Hence  $d(T(x_0), S(y_0)) \le d(A, B) + \varepsilon$  and  $P^{Ga}_{(T,S)}(A, B) \ne \emptyset$ .

**Definition 3.15.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G*. The operator  $T : A \cup B \to A \cup B$ ,  $S : A \cup B \to A \cup B$  satisfying  $T(A) \subseteq B$ ,  $S(B) \subseteq A$  is said to be a G-ciric-Rich-Rus-2map operator if:

i)  $\forall (x,y) \in A \times B \ ((x,y) \in E(G) \Rightarrow (Tx,Ty) \in E(G), (Sx,Sy) \in E(G));$ 

ii) there exists nonnegative numbers  $\alpha, \beta, \gamma$  with  $\alpha + 2\beta + \gamma < 1$ , such that, for each  $(x, y) \in E(G)$ , we have:

$$d(Tx,Sy) \le \alpha d(x,y) + \beta [d(x,Tx) + d(y,Sy)] + \gamma d(A,B).$$

**Theorem 3.16.** Let A and B be nonempty subsets of a metric space endowed with a graph G and  $T: A \cup B \to A \cup B$ ,  $S: A \cup B \to A \cup B$  are a G-ciric-Rich-Rus-2map operator. If  $x, y \in A \cup B$ satisfies the condition  $(x, Tx), (x, Sx) \in E(G), (y, Ty), (y, Sy) \in E(G)$  and if x is a G-approximate fixed point for T, or y is a G-approximate fixed point for S, then  $P_{(T,S)}^{Ga}(A, B) \neq \emptyset$ .

**Proof.** Let  $x, y \in A \cup B$  with  $(x, Tx), (x, Sx) \in E(G), (y, Ty), (y, Sy) \in E(G)$ , then

$$d(Tx,S(Tx)) \le \alpha d(x,Tx) + \beta [d(x,Tx) + d(Tx,S(Tx))] + \gamma d(A,B)$$

Therefore

$$d(Tx,S(Tx)) \leq \frac{\alpha+\beta}{1-\beta}d(x,Tx) + \frac{\gamma}{1-\beta}d(A,B).$$

Now if  $k = \frac{\alpha + \beta}{1 - \beta}$ , then

$$(3.3) d(Tx,S(Tx)) \le kd(x,Tx) + (1-k)d(A,B)$$

(3.4) 
$$d(Sy, T(Sy)) \le kd(y, Sy) + (1-k)d(A, B).$$

If x is a G-approximate fixed point for T, then there exists a  $\varepsilon > 0$  and by 3.3

$$d(Tx,S(Tx)) \leq kd(x,Tx) + (1-k)d(A,B)$$
  
$$\leq k(d(A,B) + \varepsilon) + (1-k)d(A,B)$$
  
$$= d(A,B) + k\varepsilon$$
  
$$< d(A,B) + \varepsilon.$$

and  $(x, Tx) \in P^{Ga}_{(T,S)}(A, B)$ , also if y is a *G*-approximate fixed point for *S*, then there exists a  $\varepsilon > 0$ and by 3.4

$$d(Sy, T(Sy)) \leq kd(y, Sy) + (1-k)d(A, B)$$
  
$$\leq k(d(A, B) + \varepsilon) + (1-k)d(A, B)$$
  
$$= d(A, B) + k\varepsilon$$
  
$$< d(A, B) + \varepsilon.$$

and  $(y, Sy) \in P^{Ga}_{(T,S)}(A, B)$ . Therefore  $P^{Ga}_{(T,S)}(A, B) \neq \emptyset$ .

**Theorem 3.17.** Let A and B be nonempty subsets of a metric space endowed with a graph G and  $T: A \cup B \rightarrow A \cup B$ ,  $S: A \cup B \rightarrow A \cup B$  be two continuous maps such that  $T(A) \subseteq B$ ,  $S(B) \subseteq A$ . We suppose that:

*i)* for every  $(x, y) \in E(G)$ ,  $d(Tx, Sy) \le \alpha d(x, y) + \gamma d(A, B)$  where  $\alpha, \gamma \ge 0$  and  $\alpha + \gamma = 1$ ; *ii)* for any  $\{x_n\}_{n \in N}$  and  $\{y_n\}_{n \in N}$  in  $A \cup B$  as flowing:

$$x_{n+1} = Sy_n$$
,  $y_{n+1} = Tx_n$  for some  $(x_1, y_1) \in E(G)$  for  $n \in N$ .

for any  $\{x_n\}_{n\in\mathbb{N}}$  in  $A\cup B$ , if  $x_n \to x$  and  $(x_n, x_{n+1}) \in E(G)$  for  $n \in \mathbb{N}$  then there is a subsequence convergent  $\{x_{n_k}\}_{k\geq 1}$  with  $(x_{n_k}, x) \in E(G)$  for  $n \in \mathbb{N}$ . Then there exists a  $x \in A$  such that d(x, Tx) = d(A, B).

## Proof. We have

$$d(x_{n+1}, y_{n+1}) = d(Tx_n, Sy_n)$$

$$\leq \alpha d(x_n, y_n) + \gamma(d(A, B))$$

$$\leq \dots$$

$$\leq \alpha^{n+1} d(x_0, y_0) + (1 + \alpha + \dots + \alpha^n) \gamma d(A, B).$$

If  $\{x_{n_k}\}_{k\geq 1}$  is converge to  $x_1 \in A$ , that is  $x_{n_k} \to x_1$  with  $(x_{n_k}, x_1) \in E(G)$  for all  $n \in N$ . Then

$$d(x_{n_{K+1}}, y_{n_{k+1}}) \leq \alpha^{n_{k+1}} d(x_0, y_0) + (1 + \alpha + \dots + \alpha^{n_k}) \gamma d(A, B)$$

Since T is continuous, then

$$d(x_{n_{k+1}},Tx_{n_k}) \rightarrow \frac{\gamma}{1-\alpha}d(A,B) = d(A,B).$$

Therefore  $d(x_1, Tx_1) = d(A, B)$ .

**Definition 3.18.** Let *A* and *B* be nonempty subsets of a metric space endowed with a graph *G* and Let  $T : A \cup B \to A \cup B$ ,  $S : A \cup B \to A \cup B$  be continues maps such that  $T(A) \subseteq B$ ,  $S(B) \subseteq A$ . We define diameter  $P_{(T,S)}^{Ga}(A,B)$  by,

$$diam(P^{Ga}_{(T,S)}(A,B)) = \sup\{d(x,y): d(Tx,Ty) \le \varepsilon + d(A,B) \text{ for some } \varepsilon > 0\}.$$

**Example 3.19.** Suppose  $A = \{(x,0): 0 \le x \le 1\}$ ,  $B = \{(x,1): 0 \le x \le 1\}$ ,  $T(x,0) = T(x,1) = (\frac{1}{2},1)$  and  $S(x,1) = S(x,0) = (\frac{1}{2},0)$ . Then d(T(x,0),S(y,1)) = 1 and  $diam(P_{(T,S)}^{Ga}(A,B) = diam(A \times B) = \sqrt{2}$ .

**Theorem 3.20.** Let A and B be nonempty subsets of a metric space endowed with a graph G and Let  $T : A \cup B \to A \cup B$ ,  $S : A \cup B \to A \cup B$  be continues maps such that  $T(A) \subseteq B$ ,  $S(B) \subseteq A$ . If, there exists a  $k \in [0, 1]$ ,

$$d(x,Tx) + d(Sy,y) \le kd(x,y).$$

Then

$$diam(P^{Ga}_{(T,S)}(A,B) \leq \frac{\varepsilon}{1-k} + \frac{d(A,B)}{1-k} \text{ for some } \varepsilon > 0.$$

**Proof.** If  $(x, y) \in P^{Ga}_{(T,S)}(A, B)$ , then

$$d(x,y) \leq d(x,Tx) + d(Tx,Sy) + d(Sy,y)$$
  
$$\leq \varepsilon + kd(x,y) + d(A,B).$$

Therefore  $d(x,y) \leq \frac{\varepsilon}{1-k} + \frac{d(A,B)}{1-k}$ . Then  $diam(P_{(T,S)}^{Ga}(A,B)) \leq \frac{\varepsilon}{1-k} + \frac{d(A,B)}{1-k}$ .

# **Conflict of Interests**

The authors declare that there is no conflict of interests.

# 4. Conclusions

Nowadays, fixed points, approximate fixed points and graphs in metric spaces play an important role in different areas of mathematics and its applications, particularly in mathematics, physics, biochemistry, electrical engineering and computer science. In this work, we introduced the new classes of operators and contraction maps and gave results about *G*-approximate best proximity points and diameter *G*-approximate best proximity points on the metric space *X* endowed with a graph *G* such that the set V(G) of vertices of *G* coincides with *X*. Also, by using two general Theorem regarding *G*-approximate best proximity points of cyclic maps on metric spaces we proved several *G*-approximate best proximity points theorems and diameter *G*-approximate best proximity points for a new class of operators and contraction mapping on the metric space *X* endowed with a graph *G*. We accompanied our theoretical results by some applied examples.

#### REFERENCES

- Eldred, A. Anthony; Veeramani, P. Existence and convergence of best proximity points. J. Math. Anal. Appl. 323 (2006), no. 2, 1001–1006.
- [2] Beer, Gerald; Pai, Devidas. Proximal maps, prox maps and coincidence points. Numer. Funct. Anal. Optim. 11 (1990), no. 5-6, 429–448.
- [3] F. Bojor, Fixed point of  $\phi$ -contraction in metric spaces endowed with a graph, Ann. Univ. Craiova Math. Comput. Sci. Ser. 37 (4) (2010) 85–92.
- [4] F. Bojor, Fixed point theorems for Reich type contractions on metric spaces with a graph, Nonlinear Analysis 75 (2012) 3895–3901.
- [5] L. Caccetta and K. Vijayan, Applications of graph theory, Fourteenth Australasian Conference on Combinatorial Mathematics and Computing (Dunedin, 1986), Ars. Combin., Vol. 23 (1987) 21-77.
- [6] Fan, Ky. Extensions of two fixed point theorems of F. E. Browder. Math. Z. 112 1969 234 240.
- [7] Wayne Goddard and Ortrud R. Oellermann, Distance in Graphs, Structural Analysis of Complex Networks, (2010) 49-72, DOI: 10.1007/978 - 0 - 8176 - 4789 - 6 - 3
- [8] J. Jachymski, The contraction principle for mappings on a metric space with a graph, Proc. Amer. Math. Soc. 1 (136) (2008) 1359–1373.
- [9] Kim, Won Kyu; Lee, Kyoung Hee. Corrigendum to: "Existence of best proximity pairs and equilibrium pairs" [J. Math. Anal. Appl. 316 (2006), no. 2, 433–446; MR2206681]. J. Math. Anal. Appl. 329 (2007), no. 2, 1482–1483.
- [10] Kirk, W. A.; Reich, Simeon; Veeramani, P. Proximinal retracts and best proximity pair theorems. Numer. Funct. Anal. Optim. 24 (2003), no. 7-8, 851–862.
- [11] S.A.M. Mohsenalhosseini, Approximate best proximity pairs in metric space for Contraction Maps, Advances in Fixed Point Theory, 4 (2014), pp. 310-324.
- [12] S.A.M. Mohsenalhosseini, H. Mazaheri and M.A. Dehghan, Approximate best proximity pairs in metric space, Abstract and Applied Analysis, (2011), Article ID 596971, 9 pages.
- [13] S. A. M. Mohsenalhosseini, H. Mazaheri, *Fixed Point for Completely Norm Space and Map T* $_{\alpha}$ , Mathematica Moravica Vol. **16** 2, (2012), 25-35.
- [14] F. S. Roberts, Graph theory and its applications to the problems of society, CBMS-NSF Monograph 29, SIAM Publications, Philadelphia (1978).
- [15] Singer, Ivan. Best approximation in normed linear spaces by elements of linear subspaces. Translated from the Romanian by Radu Georgescu. Die Grundlehren der mathematischen Wissenschaften, Band 171 Publishing

House of the Academy of the Socialist Republic of Romania, Bucharest; Springer-Verlag, New York-Berlin 1970 415 pp.

- [16] I. A. Rus, Generalized Contractions and Applications, Cluj University Press, Cluj-Napoca, 2001.
- [17] I.A. Rus, A. Petruşel, G. Petruşel, Fixed Point Theory, Cluj University Press, Cluj-Napoca, 2008.
- [18] S. Reich, Fixed points of contractive functions, Boll. Unione Mat. Ital. 5 (1972) 26-42.
- [19] L.B. Ćirić, A generalization of Banach's contraction principle, Proc. Amer. Math. Soc. 45 (1974) 267–273.
- [20] V. Vetrivel, P. Veeramani, P. Bhattacharyya, Some extensions of Fan's best approxomation theorem, Numer. Funct. Anal. Optim. 13 (1992) 397-402.
- [21] K. Włodarczyk, R. Plebaniak, A. Banach, Best proximity points for cyclic and noncyclic set-valued relatively quasi-asymptotic contractions in uniform spaces, Nonlinear Anal. 70 (2009) 3332-3341.
- [22] K. Włodarczyk, R. Plebaniak, A. Banach, Erratum to: "Best proximity points for cyclic and noncyclic setvalued relatively quasi-asymptotic contractions in uniform spaces" [Nonlinear Anal. 70 (2009) 3332-3341], Nonlinear Anal. 71 (2009) 3583-3586.
- [23] K. Włodarczyk, R. Plebaniak, C. Obczyński, Convergence theorems, best approximation and best proximity for set-valued dynamic systems of relatively quasi-asymptotic contractions in cone uniform spaces, Nonlinear Anal. 72 (2010) 794-805.

This figure "dis3.png" is available in "png" format from:

http://arxiv.org/ps/2411.03472v1

This figure "dis4.jpg" is available in "jpg" format from:

http://arxiv.org/ps/2411.03472v1

This figure "distance.png" is available in "png" format from:

http://arxiv.org/ps/2411.03472v1