Hyperrigidity II: R-dilations and ideals

Paweł Pietrzycki and Jan Stochel

ABSTRACT. In this paper, we study hyperrigidity for C^* -algebras. The absence of the unit in hyperrigid set creates the possibility of the existence of R-dilations with non-isometric R. This gives rise to the study of when a hyperrigid set is annihilated by a state, or more generally, by a UCP map, which in turn, is closely related to the concept of rigidity at 0 introduced by G. Salomon, who studied hyperrigid subsets of Cuntz-Krieger algebras. Moreover, we obtain a characterization of hyperrigid sets analogous to that obtained by Hansen and Pedersen for operator convex functions.

1. Introduction

The classical approximation theorem due to P. P. Korovkin [19] states that for any sequence of positive linear maps $\Phi_k : C[0,1] \to C[0,1], k \in \mathbb{N}$,

$$\lim_{k \to \infty} \|\Phi_k(x^j) - x^j\| = 0 \ \forall j \in \{0, 1, 2\} \implies \lim_{k \to \infty} \|\Phi_k(f) - f\| = 0 \ \forall f \in C[0, 1],$$

where \mathbb{N} stands for the set of all positive integers. In other words, the asymptotic behaviour of the sequence $\{\Phi_k\}_{k=1}^{\infty}$ on the C^* -algebra C[0,1] is uniquely determined by the vector space G spanned by $\{1,x,x^2\}$. This theorem unified many existing approximation processes such as the Bernstein polynomial approximation of continuous functions. It is worth mentioning that positive approximation process plays a fundamental role in the approximation theory and its applications. Another major achievement was the discovery of geometric theory of Korovkin sets by Y. A. Šaškin [26]. Namely, Šaškin observed that the key property of G is that its Choquet boundary (see Section 3 for the definition) coincides with [0,1]. Detailed surveys of most of these developments can be found in [6,1,2].

A natural non-commutative analogue of Korovkin-type rigidity would be a subset G of a unital C^* -algebra $\mathscr A$ with the property that for any sequence of unital completely positive (UCP) maps¹ $\Phi_k : \mathscr A \to \mathscr A$ $(k \in \mathbb N)$,

$$\lim_{k \to \infty} \|\Phi_k(g) - g\| = 0 \ \forall g \in G \implies \lim_{k \to \infty} \|\Phi_k(a) - a\| = 0 \ \forall a \in \mathscr{A}.$$

1

²⁰²⁰ Mathematics Subject Classification. Primary 46G10, 47B15; Secondary 47A63, 44A60. Key words and phrases. hyperrigidity, R-dilation, ideal, state.

The research of both authors was supported by the National Science Center (NCN) Grant OPUS No. DEC- $\frac{2021}{43}$ B/ST1/01651.

¹In this paper, it is assumed that completely positive maps are linear, representations of unital C^* -algebras preserve units and involutions, and C^* -subalgebras of a unital C^* -algebra $\mathscr A$ contain the unit of $\mathscr A$.

In fact, Arveson introduced even more non-commutativity in this picture. Namely, motivated both by the fundamental role of the classical Choquet boundary in classical approximation theory, and by the importance of approximation in the contemporary theory of operator algebras, he introduced hyperrigidity as a form of approximation that captures many important operator-algebraic phenomena. For the purposes of this paper, we use the concept of hyperrigidity in a more general context. Unlike Arveson's definition of hyperrigidity (see [5, Definition1.1]), we do not require that the set G, a candidate for a hyperrigid set, be separable or generate a given C^* -algebra \mathscr{A} . If G is finite or countably infinite and generates the C^* -algebra \mathscr{A} , then the current definition of hyperrigidity coincides with that given by Arveson. Through of the paper, $B(\mathcal{H}, \mathcal{K})$ stands for the Banach space of all bounded linear operators from a Hilbert space \mathcal{H} into a Hilbert space \mathcal{K} (all Hilbert spaces considered hereinafter are complex). We abbreviate $B(\mathcal{H}, \mathcal{H})$ to $B(\mathcal{H})$ and regard it as a C^* -algebra; I denotes the identity operator on \mathcal{H} .

DEFINITION 1.1. A nonempty subset G of a unital C^* -algebra \mathscr{A} is said to be hyperrigid (relative to \mathscr{A}) if for any faithful representation $\pi \colon \mathscr{A} \to B(\mathcal{H})$ on a Hilbert space \mathcal{H} and for any sequence $\Phi_k \colon B(\mathcal{H}) \to B(\mathcal{H})$ $(k \in \mathbb{N})$ of UCP maps,

$$\lim_{k \to \infty} \|\Phi_k(\pi(g)) - \pi(g)\| = 0 \ \forall g \in G \implies \lim_{k \to \infty} \|\Phi_k(\pi(a)) - \pi(a)\| = 0 \ \forall a \in \mathscr{A}.$$

Note that even in the case when \mathscr{A} is commutative, a priori this phenomenon is stronger than the one observed by Korovkin, as we allow the maps Φ_k to take values outside of \mathscr{A} . Arveson initiated a study of what might be called noncommutative approximation theory, focusing on the problem of when a finite or countably infinite set of generators of a unital C^* -algebra \mathscr{A} is hyperrigid. Arveson himself gave in [5, Theorem 2.1] a characterization of hyperrigidity that replaces the limit process by the so-called unique extension property. A more intrinsic characterization of the unique extension property can be found in [3, Proposition 2.4]). Arveson proved that if $T \in \mathcal{B}(\mathcal{H})$ is a selfadjoint operator and \mathscr{A} is the C^* -algebra generated by T, then $G = \{T, T^2\}$ is hyperrigid in $C^*(G)$, the unital C^* -subalgebra of $\mathcal{B}(\mathcal{H})$ generated by $G \cup \{I\}$. Moreover, if $V_1, \ldots, V_n \in \mathcal{B}(\mathcal{H})$ is a finite set of isometries that generates a C^* -algebra \mathscr{A} , then $G = \{V_1, \ldots, V_n, V_1 V_1^* + \cdots + V_n V_n^*\}$ is a hyperrigid set of generators for \mathscr{A} .

In accordance with Šaškin's insightful observation, Arveson [5] (see also [3, 4]) conjectured that hyperrigidity is equivalent to the non-commutative Choquet boundary of G being as large as possible, in the sense that every irreducible representation of $C^*(G)$ should be a boundary representation for G. This is now known as Arveson's hyperrigidity conjecture (see [5, Conjecture 4.3]). Some positive solutions of Arveson's hyperrigidity conjecture were found for certain classes of C^* -algebras (see [5, 18, 9, 11, 25, 16, 28, 23]). In full generality, Arveson's hyperrigidity conjecture turns out to have a negative solution. Recently, a counterexample has been found by B. Bilich and A. Dor-On in [8] (see also [7]). However, Arveson's hyperrigidity conjecture is still open for commutative C^* -algebras (even the singly generated case is not resolved).

In recent years, this issue has attracted considerable interest in various parts of operator algebras and operator theory [17, 18, 11, 12, 13, 33]. In particular, it is related to the Arveson-Douglas essential normality conjecture involving quotient modules of the Drury-Arveson space [17, Theorem 4.12].

2. Main results

In this section we will formulate the main results of this paper. The proofs of Theorems 2.3 and 2.11 appear in Section 4.

In a recent paper [23] we studied, in the context of a unital commutative C^* -algebra $\mathscr A$ generated by a single element $t\in\mathscr A$, which of the sets $G=\{t^{*m}t^n\colon (m,n)\in\Xi\}$ with $\Xi\subseteq\mathbb Z_+^2$ are hyperrigid in $\mathscr A$, where $\mathbb Z_+$ stands for the set of all nonnegative integers. We have shown that under some mild constraints imposed on Ξ , the assumption that the set G generates the C^* -algebra $\mathscr A$ implies the hyperrigidity of G (see [23, Theorem 2.4]). As a consequence, we obtained two criteria for hyperrigidity of G, one dependent on the geometry of the spectrum of the generating element t, the other independent (see [23, Theorem 2.5]). The independent case reads as follows.

THEOREM 2.1 ([23, Theorem 2.5(i)]). Let \mathscr{A} be a unital commutative C^* -algebra generated by $t \in \mathscr{A}$ and let Ξ be a set satisfying the following condition²:

$$\begin{array}{l} \{(p,q),(r,r)\}\subseteq \varXi\subseteq \mathbb{Z}_+^2 \ \textit{for some} \ p,q,r\in \mathbb{Z}_+ \ \textit{such that} \\ p\neq q \ \textit{and} \ p+q<2r, \ \textit{and} \ \gcd\{m-n\colon (m,n)\in \varXi\}=1. \end{array}$$

Then the set $G := \{t^{*m}t^n : (m,n) \in \Xi\}$ generates $\mathscr A$ and is hyperrigid in $\mathscr A$.

Regarding Theorem 2.1, it is worth pointing out that if a unital commutative C^* -algebra $\mathscr A$ is generated by an element t, then, by [24, Theorem 11.19], $\mathscr A$ can be identified (up to the C^* -algebra isomorphism) with C(X), where X is a nonempty compact subset of $\mathbb C$, and t can be identified with the function $\xi \colon X \to \mathbb C$, called the *coordinate function* on X, defined by

$$\xi(z) = z, \quad z \in X; \tag{2.1}$$

here C(X) stands for the C^* -algebra of all continuous complex functions on X equipped with supremum norm. Now, if $0 \in X$ and $(0,0) \notin \Xi$, then f(0) = 0 for every $f \in G$, or equivalently $G \subseteq \ker \chi$, where χ is the character of C(X) given by $\chi(f) = f(0)$ for $f \in C(X)$. In fact, as shown in Corollary 4.3, for any character χ of a singly generated commutative unital C^* -algebra \mathscr{A} , there exists a hyperrigid set G of generators of \mathscr{A} such that $G \subseteq \ker \chi$. This, in turn, is closely related to the concept of rigidity at 0 introduced by Salomon, who studied hyperrigid subsets of (unital and non-unital) Cuntz-Krieger algebras (see, e.g., [25, Example 5.5]). The original definition (see [25, Definition 3.1]) states that a set G generating a C^* -algebra \mathscr{A} is rigid at 0 in \mathscr{A} if for every sequence $\{\psi_n\}_{n=1}^{\infty}$ of contractive positive maps $\psi_n \colon \mathscr{A} \to \mathbb{C}$,

$$\lim_{n \to \infty} \psi_n(g) = 0 \ \forall g \in G \quad \Longrightarrow \quad \lim_{n \to \infty} \psi_n(a) = 0 \ \forall a \in \mathscr{A}.$$

If $\mathscr A$ is separable, then G is rigid at 0 if and only if there are no states on $\mathscr A$ that vanish on G (see [25, Theorem 3.3]). Therefore, if $\mathscr A$ is unital and separable and G is not rigid at 0, then G does not contain the unit of $\mathscr A$. For example, if G satisfies the assumptions of Theorem 2.1 with $\mathscr A=C(X)$ and $t=\xi$ and $(0,0)\notin \Xi$, then G contains the unit of C(X) if and only if X is a closed subset of the unit circle $\mathbb T:=\{z\in\mathbb C\colon |z|=1\}.$

²The abbreviation "gcd" stands for "the greatest common divisor" (always assumed to be positive). It follows from the well-ordering principle that if J is a nonempty set of integers, not all 0, then $\gcd(J)$ exists and there exists a finite nonempty subset J' of J such that $\gcd(J) = \gcd(J')$. For simplicity, writing $\gcd(J)$ means that J contains a nonzero integer.

Continuing the above discussion, let us recall the following dilation approach to characterizing hyperrigidity of subsets of C(X).

THEOREM 2.2 ([23, Theorem 2.2]). Let X be a nonempty compact subset of \mathbb{C} and G be a set of generators of C(X). Then the following conditions are equivalent:

- (i) G is hyperrigid,
- (ii) for all Hilbert spaces \mathcal{H} and \mathcal{K} such that $\mathcal{H} \subseteq \mathcal{K}$, and all normal operators $T \in \mathcal{B}(\mathcal{H})$ and $N \in \mathcal{B}(\mathcal{K})$ with spectra in X,

$$f(T) = Pf(N)|_{\mathcal{H}} \ \forall f \in G \implies PN = NP,$$

where P stands for the orthogonal projection of K onto \mathcal{H} ,

(iii) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all normal operators $T \in \mathbf{B}(\mathcal{H})$ and $N \in \mathbf{B}(\mathcal{K})$ with spectra in X, and every isometry $V : \mathcal{H} \to \mathcal{K}$,

$$f(T) = V^* f(N) V \ \forall f \in G \implies VT = NV.$$
 (2.2)

Moreover, conditions (i)-(iii) are still equivalent regardless of whether the Hilbert spaces considered in either of them are separable or not.

If we replace the isometry $V : \mathcal{H} \to \mathcal{K}$ in the if-clause of implication (2.2) by any operator $R \in \mathcal{B}(\mathcal{H}, \mathcal{K})$, i.e.,

$$f(T) = R^* f(N)R, \quad f \in G, \tag{2.3}$$

then R is an isometry if G contains the unit $\mathbf{1}$ of C(X). The absence of the unit $\mathbf{1}$ in G creates the possibility of the existence of non-isometric solutions R of (2.3). All of this together gives rise to the study of when a hyperrigid set G in $\mathscr A$ is annihilated by a state, or more generally, by a UCP map. This is explored in Section 4 (see, e.g., Theorem 2.8). The results given there and below are formulated for general unital C^* -algebras $\mathscr A$, so condition (2.3) should be replaced by the following:

$$\pi(g) = R^* \rho(g) R, \quad g \in G,$$

where $\pi: \mathscr{A} \to B(\mathcal{H})$ and $\rho: \mathscr{A} \to B(\mathcal{K})$ are representations of \mathscr{A} and $R \in B(\mathcal{H}, \mathcal{K})$. As discussed in [25], when considering hyperrigidity for non-unital C^* -algebras, UCP maps must be replaced by completely contractive completely positive (CCCP) maps. It turns out that even in the unital case, as long as the set G generating the C^* -algebra in question does not contain unit, CCCP maps naturally fit into the context of hyperrigidity. This is shown in Theorem 2.3 below. But first let us recall that according to [20, Theorem 2.1.7] and the Stinespring dilation theorem (see [29, Theorem 1]):

A linear map
$$\Psi \colon \mathscr{A} \to B(\mathcal{H})$$
 is CCCP if and only if Ψ is completely positive and $\|\Psi(e)\| \leqslant 1$.

Given a unital C^* -algebra \mathscr{A} , we denote by $\mathfrak{M}_{\mathscr{A}}$ the set of all characters of \mathscr{A} , i.e., the set of all one-dimensional representations $\chi \colon \mathscr{A} \to \mathbb{C}$.

Theorem 2.3. Let G be a nonempty subset of a unital C*-algebra $\mathscr A$ and ϕ be a state on $\mathscr A$. Consider the following conditions:

- (i) G is hyperrigid,
- (ii) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all representations $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ and every contraction $R \in B(\mathcal{H}, \mathcal{K})$,

$$\pi(g) = R^* \rho(g) R \ \forall g \in G \implies R\pi(a) = \rho(a) R \ \forall a \in \mathscr{A}, \tag{2.5}$$

(iii) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all representations $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ and every contraction $R \in B(\mathcal{H}, \mathcal{K})$,

$$\pi(g) = R^* \rho(g) R \ \forall g \in G \implies \pi(a) = R^* \rho(a) R \begin{cases} \forall a \in \mathscr{A} \ if \ R^* R = I \\ \forall a \in \ker \chi \ if \ R^* R \neq I \\ for \ some \ \chi \in \mathfrak{M}_{\mathscr{A}}, \end{cases}$$
 (2.6)

(iv) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all representations $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ and every contraction $R \in B(\mathcal{H}, \mathcal{K})$,

$$\pi(g) = R^* \rho(g) R \ \forall g \in G \implies \pi(a) = R^* \rho(a) R \ \forall a \in \ker \phi,$$

(v) for every Hilbert space \mathcal{H} , every representation $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and every CCCP map $\Psi \colon \mathscr{A} \to B(\mathcal{H})$,

$$\pi(g) = \Psi(g) \ \forall g \in G \implies \pi(a) = \Psi(a) \ \forall a \in \ker \phi.$$

Then

- (a) $(iv) \Rightarrow (iii) \Rightarrow (ii)$ and $(iv) \Leftrightarrow (v)$,
- (b) if G generates \mathscr{A} , then (ii) \Rightarrow (i) and (ii) \Leftrightarrow (iii),
- (c) if $\phi \in \mathfrak{M}_{\mathscr{A}}$, then (iv) \Rightarrow (ii),
- (d) if $G \subseteq \ker \phi$, then
 - (d.1) $(i) \Rightarrow (ii)$, $(i) \Rightarrow (iii)$, $(i) \Rightarrow (iv)$ and $(i) \Rightarrow (v)$,
 - (d.2) if G is hyperrigid, π and ρ are representations and R is a non-isometric contraction satisfying the if-clause of (2.5), then $\phi \in \mathfrak{M}_{\mathscr{A}}$,
 - (d.3) if $C^*(G) = \mathcal{A}$, then conditions (i)-(iii) are equivalent,
 - (d.4) if $C^*(G) = \mathscr{A}$ and ϕ is a character of \mathscr{A} , then (i)-(v) are equivalent.

COROLLARY 2.4. Let \mathscr{A} be a unital C^* -algebra which has no characters and let G be a nonempty subset of \mathscr{A} such that $G \subseteq \ker \psi$ for some state ψ on \mathscr{A} . Then the following conditions are equivalent:

- (i) G is hyperrigid,
- (ii) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all representations $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ and every contraction $R \in B(\mathcal{H}, \mathcal{K})$,

$$\pi(q) = R^* \rho(q) R \ \forall q \in G \implies \pi(a) = R^* \rho(a) R \ \forall a \in \mathscr{A},$$

(iii) for every Hilbert space \mathcal{H} , every representation $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and every CCCP map $\Psi \colon \mathscr{A} \to B(\mathcal{H})$.

$$\pi(g) = \Psi(g) \ \forall g \in G \implies \pi(a) = \Psi(a) \ \forall a \in \mathscr{A}.$$

Moreover: if (ii) holds, then R is an orthogonal projection; if (iii) holds, then Ψ is a UCP map.

PROOF. (i) \Rightarrow (ii) Let (π, ρ, R) be as in (ii) and $\pi(g) = R^*\rho(g)R$ for all $g \in G$. Since \mathscr{A} has no characters, implication (i) \Rightarrow (iii) of Theorem 2.3 shows that R is an isometry and the identity $\pi = R^*\rho R$ holds on the whole algebra \mathscr{A} .

- (ii) \Leftrightarrow (iii) This equivalence is a direct consequence of (2.4) and the Stinespring dilation theorem. In particular, by the previous paragraph, Ψ is a UCP map.
- (iii) \Rightarrow (i) In view of (2.4) and [23, Theorem B.2(iii)] ([5, Theorem 2.1]), this implication is obvious.

The existence of a non-isometric contraction R in the if-clause of implication (2.5), under the assumption of hyperrigidity of G, is closely related to the rigidity of G at 0 and the existence of a character of the C^* -algebra $\mathscr A$ vanishing on G. This is discussed in the two corollaries below.

COROLLARY 2.5. Let G be a hyperrigid set of generators of a unital C^* -algebra \mathscr{A} . Let $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ be representations on Hilbert spaces \mathcal{H} and \mathcal{K} respectively, and $R \in B(\mathcal{H}, \mathcal{K})$ be a non-isometric contraction such that $\pi(g) = R^* \rho(g) R$ for every $g \in G$. Then the following conditions are equivalent:

- (i) G is rigid at 0,
- (ii) there is no character of \mathscr{A} vanishing on G.

Moreover, there exists at most one state ψ on $\mathscr A$ such that $G \subseteq \ker \psi$, and if such a state exists, then it is a character of $\mathscr A$.

PROOF. In view of Theorem 2.3(b), only the "moreover" part requires the proof. Since R is not an isometry, we see that $e \notin G$, where e is the unit of \mathscr{A} . Suppose that ψ_1 and ψ_2 are states on \mathscr{A} vanishing on G. From Theorem 2.3(b), it follows that ψ_1 and ψ_2 are characters of \mathscr{A} that coincide on the set G. Since G generates the C^* -algebra \mathscr{A} , we conclude that $\psi_1 = \psi_2$.

Corollary 2.6. Let G be a hyperrigid subset of a unital C^* -algebra \mathscr{A} . Then the following conditions are equivalent:

- (i) there exists a character χ of \mathscr{A} such that $G \subseteq \ker \chi$,
- (ii) there exist representations $\pi: \mathscr{A} \to \mathbf{B}(\mathcal{H})$ and $\rho: \mathscr{A} \to \mathbf{B}(\mathcal{K})$ on Hilbert spaces \mathcal{H} and \mathcal{K} , a non-isometric contraction $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$ and a state ψ on \mathscr{A} such that $G \subseteq \ker \psi$ and $\pi(g) = R^* \rho(g) R$ for $g \in G$.

Moreover, if there exists a character of $\mathscr A$ vanishing on G, say χ , then any state on $\mathscr A$ vanishing on G coincides with χ .

PROOF. (i) \Rightarrow (ii) Take nonzero Hilbert spaces \mathcal{H} and \mathcal{K} . Define the representations $\pi \colon \mathscr{A} \to B(\mathcal{H})$ and $\rho \colon \mathscr{A} \to B(\mathcal{K})$ by

$$\pi(a) = \chi(a)I_{\mathcal{H}}$$
 and $\rho(a) = \chi(a)I_{\mathcal{K}}$ for every $a \in \mathscr{A}$.

Set $\psi = \chi$. Take an arbitrary $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$. Then $\pi(g) = R^* \rho(g) R$ for every $g \in G$ and $R\pi(a) = \rho(a)R$ for every $a \in \mathscr{A}$.

(ii)⇒(i) This implication is a direct consequence of Theorem 2.3(b).

The "moreover" part follows from the "moreover" part of Corollary 2.5 and the implication (i) \Rightarrow (ii).

REMARK 2.7. We have shown in the proof of implication (i) \Rightarrow (ii) of Corollary 2.6 that if there exists a character χ of $\mathscr A$ such that $G \subseteq \ker \chi$, then for all Hilbert spaces $\mathcal H$ and $\mathcal K$ there exist representations $\pi \colon \mathscr A \to B(\mathcal H)$ and $\rho \colon \mathscr A \to B(\mathcal K)$ such that for every $R \in B(\mathcal H, \mathcal K)$, $\pi(g) = R^* \rho(g) R$ for every $g \in G$.

The next two results model the objects appearing in assertion (b) of Theorem 2.3 and condition (ii) of Corollary 2.6. Below, given a contraction $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$, we write

$$\triangle = I - R^* R \quad \text{and} \quad \triangle_* = I - R R^*. \tag{2.7}$$

The operators $\triangle^{1/2}$ and $\triangle^{1/2}_*$ are called the *defect operators* of R (cf. [31]). Following W. Kaufman [15], we call an operator $R \in B(\mathcal{H})$ a pure contraction if ||Rh|| < ||h|| for every nonzero vector $h \in \mathcal{H}$.

THEOREM 2.8. Let G be a hyperrigid subset of a unital C*-algebra \mathscr{A} and ϕ be a state on \mathscr{A} such that $G \subseteq \ker \phi$. Let \mathcal{H} and \mathcal{K} be Hilbert spaces, $\pi \colon \mathscr{A} \to \mathbf{B}(\mathcal{H})$ and $\rho \colon \mathscr{A} \to \mathbf{B}(\mathcal{K})$ be representations and $R \in \mathbf{B}(\mathcal{H}, \mathcal{K})$ be a contraction such that $\pi(g) = R^* \rho(g) R$ for all $g \in G$. Then

 $R\pi(a) = \rho(a)R \ \forall a \in \mathscr{A} \ \& \ \ker \phi \subseteq J \ \& \ J \ is \ a \ closed *-ideal in \mathscr{A},$ (2.8) where $^3J := \{a \in \mathscr{A} : \pi(a) = R^*\rho(a)R\}$. Moreover, if R is non-isometric, then

- (i) $\ker \phi = J$ and ϕ is a character of \mathscr{A} ,
- (ii) \triangle commutes with π and \triangle_* commutes with ρ (see (2.7)),
- (iii) $\mathcal{H}_1 := \overline{\mathscr{R}(\Delta)} \neq \{0\} \text{ reduces } \pi \text{ and } \mathcal{K}_1 := \overline{\mathscr{R}(\Delta_* R)} \text{ reduces } \rho,$
- (iv) $R = R_0 \oplus R_1$, where $R_0 \in \boldsymbol{B}(\mathcal{H}_0, \mathcal{K}_0)$ is an isometry with $\mathcal{H}_0 := \mathcal{H}_1^{\perp}$ and $\mathcal{K}_0 := \mathcal{K}_1^{\perp}$, and $R_1 \in \boldsymbol{B}(\mathcal{H}_1, \mathcal{K}_1)$ is a pure contraction with dense range,
- (v) $\pi = \pi_0 \oplus \phi I_{\mathcal{H}_1}$ and $\pi_0 = R_0^* \rho_0 R_0$, where $\pi_0 := \pi|_{\mathcal{H}_0}$ and $\rho_0 := \rho|_{\mathcal{K}_0}$,
- (vi) $\rho = \rho_0 \oplus \phi I_{\mathcal{K}_1}$.

And *vice versa*, the following holds.

THEOREM 2.9. Let G be a nonempty subset of a unital C^* -algebra $\mathscr A$ and ϕ be a character of $\mathscr A$ such that $G \subseteq \ker \phi$. Let $R \in B(\mathcal H, \mathcal K)$ be a contraction of the form $R = R_0 \oplus R_1$ with $R_j \in B(\mathcal H_j, \mathcal K_j)$ for j = 0, 1, $\pi = \pi_0 \oplus \chi I_{\mathcal H_1}$, $\rho = \rho_0 \oplus \chi I_{\mathcal K_1}$ and $\pi_0 = R_0^* \rho_0 R_0$, where $\pi_0 \colon \mathscr A \to B(\mathcal H_0)$ and $\rho_0 \colon \mathscr A \to B(\mathcal K_0)$ are representations. Then $\pi(g) = R^* \rho(g) R$ for all $g \in G$ and R_0 is an isometry. Moreover, if R (equivalently R_1) is non-isometric, then

- (i) conditions (i) and (ii) of Theorem 2.8 hold,
- (ii) $\mathcal{H}_1 = \overline{\mathscr{R}(\Delta)}$ provided R_1 is a pure contraction,
- (iii) $\mathcal{K}_1 = \overline{\mathcal{R}(\Delta_* R)}$ provided R_1 is a pure contraction with dense range.

Our next result, Theorem 2.11, is inspired by the characterizations of operator convex functions given by Hansen and Pedersen.

THEOREM 2.10 ([14, Theorem 2.1]). If f is a continuous, real function on the half-open interval $[0, \alpha)$ (with $\alpha \leq \infty$), the following conditions are equivalent:

- (i) f is operator convex and $f(0) \leq 0$,
- (ii) $f(A^*XA) \leq A^*f(X)A$ for all A with $||A|| \leq 1$ and every selfadjoint X with spectrum in $[0, \alpha)$,
- (iii) $f(A^*XA+B^*YB) \leqslant A^*f(X)A+B^*f(Y)B$ for all A, B with $A^*A+B^*B \leqslant I$ and all X, Y with spectra in [0, a),
- (iv) $f(PXP) \leq Pf(X)P$ for every projection P and every selfadjoint X with spectrum in [0, a).

To be more precisely, conditions (ii) and (iii) of Theorem 2.11 and condition (ii) of Theorem 2.2 correspond to conditions (ii), (iii) and (iv) of Theorem 2.10, respectively. Condition (iii) of Theorem 2.11 is also related to [22, Theorem 1.5].

THEOREM 2.11. Suppose that X is a nonempty compact subset of \mathbb{C} , μ is a Borel probability measure on X and G is a finite or countably infinite set of generators of C(X) such that $\int_X f d\mu = 0$ for every $f \in G$. Fix an integer $n \ge 2$. Then the following conditions are equivalent:

(i) G is hyperrigid,

³By automatic continuity of representations of C^* -algebras, the ideal J is closed.

(ii) for all Hilbert spaces \mathcal{H} and \mathcal{K} , all normal operators $T \in B(\mathcal{H})$ and $N \in B(\mathcal{K})$ with spectra in X, and every contraction $R \in B(\mathcal{H}, \mathcal{K})$,

$$f(T) = R^* f(N)R \ \forall f \in G \implies RT = NR, \tag{2.9}$$

(iii) for all Hilbert spaces \mathcal{H} , $\mathcal{K}_1, \ldots, \mathcal{K}_n$, all normal operators $T \in \mathbf{B}(\mathcal{H})$, $N_1 \in \mathbf{B}(\mathcal{K}_1), \ldots, N_n \in \mathbf{B}(\mathcal{K}_n)$ with spectra in X, and all operators $R_1 \in \mathbf{B}(\mathcal{H}, \mathcal{K}_1), \ldots, R_n \in \mathbf{B}(\mathcal{H}, \mathcal{K}_n)$ such that $\sum_{i=1}^n R_i^* R_i \leq I$,

$$f(T) = \sum_{i=1}^{n} R_i^* f(N_i) R_i \ \forall f \in G \implies R_i T = N_i R_i \ \forall i \in \{1, \dots, n\},$$
 (2.10)

(iv) for all Hilbert spaces \mathcal{H} , $\mathcal{K}_1, \ldots, \mathcal{K}_n$, all normal operators $T \in \mathbf{B}(\mathcal{H})$, $N_1 \in \mathbf{B}(\mathcal{K}_1), \ldots, N_n \in \mathbf{B}(\mathcal{K}_n)$ with spectra in X, and all operators $R_1 \in \mathbf{B}(\mathcal{H}, \mathcal{K}_1), \ldots, R_n \in \mathbf{B}(\mathcal{H}, \mathcal{K}_n)$ such that $\sum_{i=1}^n R_i^* R_i \leq I$,

$$f(T) = \sum_{i=1}^{n} R_i^* f(N_i) R_i \ \forall f \in G \implies f(T) = \sum_{i=1}^{n} R_i^* f(N_i) R_i \ \forall f \in C_{\mu}(X),$$

where $C_{\mu}(X) := \{ f \in C(X) : \int_{X} f d\mu = 0 \}$. Moreover, the following hold:

- (a) conditions (i)-(iv) are still equivalent regardless of whether the Hilbert spaces considered in either of them are separable or not,
- (b) if G is hyperrigid and T, N, R (resp., T, N₁,..., N_n, R₁,..., R_n) satisfy the if-clause of (2.9) (resp., (2.10)) with $R^*R \leq I$ (resp., $\sum_{i=1}^n R_i^*R_i \leq I$), then μ is the Dirac measure at some (uniquely determined) point of X.

3. Prerequisites

In this paper, we use the following notation. The fields of real and complex numbers are denoted by \mathbb{R} and \mathbb{C} , respectively. The symbols \mathbb{Z}_+ , \mathbb{N} and \mathbb{R}_+ stand for the sets of nonnegative integers, positive integers and nonnegative real numbers, respectively. We write $\mathfrak{B}(X)$ for the σ -algebra of all Borel subsets of a topological Hausdorff space X.

Given a unital C^* -algebra $\mathscr A$ and a nonempty subset G of $\mathscr A$, we denote by $C^*(G)$ the unital C^* -subalgebra of $\mathscr A$ generated by $G \cup \{e\}$, where e is the unit of $\mathscr A$. If $t \in \mathscr A$, then we write $C^*(t) = C^*(\{t\})$. The spectrum of $a \in \mathscr A$ is denoted by $\sigma(a)$. By an ideal in an algebra we always mean a two-sided ideal. It is well-known that every ideal in a C^* -algebra is selfadjoint, i.e., it is closed under adjoints (see [20, Theorem 3.1.3]). However, for the sake of completeness, we will continue to use the term "*-ideal". If $\mathscr A$ is a unital C^* -algebra, the notation " $J \triangleleft \mathscr A$ " means "J is a closed *-ideal in $\mathscr A$ ". All representations of unital C^* -algebras are always assumed to preserve involution and units. A completely positive (linear) map between unital C^* -algebras that preserves units is called a unital completely positive (UCP) map. We will abbreviate "completely contractive completely positive map" between unital C^* -algebras to "CCCP map".

Let \mathcal{H} and \mathcal{K} be (complex) Hilbert spaces. Denote by $\mathbf{B}(\mathcal{H}, \mathcal{K})$ the Banach space of all bounded linear operators from \mathcal{H} to \mathcal{K} . If $A \in \mathbf{B}(\mathcal{H}, \mathcal{K})$, then A^* , $\mathcal{N}(A)$ and $\mathcal{R}(A)$ stand for the adjoint, the kernel and the range of A, respectively. We abbreviate $\mathbf{B}(\mathcal{H}, \mathcal{H})$ to $\mathbf{B}(\mathcal{H})$ and regard it as a C^* -algebra with unit I, where $I = I_{\mathcal{H}}$ denotes the identity operator on \mathcal{H} . We write $\sigma(A)$ and $\sigma_{\mathbf{p}}(A)$ for the spectrum and the point spectrum of $A \in \mathbf{B}(\mathcal{H})$, respectively. We say that $A \in \mathbf{B}(\mathcal{H})$ is normal if $A^*A = AA^*$, selfadjoint if $A = A^*$ and positive if $\langle Ah, h \rangle \geqslant 0$

for all $h \in \mathcal{H}$. Each positive operator $A \in \boldsymbol{B}(\mathcal{H})$ has a unique positive square root denoted by $A^{1/2}$. If $A \in \boldsymbol{B}(\mathcal{H})$, we write $|A| := (A^*A)^{1/2}$. Recall that $\mathcal{N}(A) = \mathcal{N}(|A|)$.

Let \mathscr{A} be a σ -algebra of subsets of a set X and let $F \colon \mathscr{A} \to \boldsymbol{B}(\mathcal{H})$ be a semispectral measure. Denote by $L^1(F)$ the vector space of all \mathscr{A} -measurable functions $f \colon X \to \mathbb{C}$ such that $\int_X |f(x)| \langle F(\mathrm{d}x)h,h \rangle < \infty$ for all $h \in \mathcal{H}$. Then for every $f \in L^1(F)$, there exists a unique operator $\int_X f \mathrm{d}F \in \boldsymbol{B}(\mathcal{H})$ such that (see e.g., $[\mathbf{30}, \mathrm{Appendix}]$)

$$\left\langle \int_X f dF h, h \right\rangle = \int_X f(x) \langle F(dx)h, h \rangle, \quad h \in \mathcal{H}.$$
 (3.1)

If F is a spectral measure, then $\int_X f dF$ coincides with the usual spectral integral. In particular, if F is the spectral measure of a normal operator $A \in \boldsymbol{B}(\mathcal{H})$, then we write $f(A) = \int_{\mathbb{C}} f dF$ for any F-essentially bounded Borel function $f \colon \mathbb{C} \to \mathbb{C}$; the map $f \mapsto f(A)$ is called the Stone-von Neumann calculus. For more information needed in this article on spectral integrals, including the spectral theorem for normal operators and the Stone-von Neumann calculus, we refer the reader to $[\mathbf{24}, \mathbf{32}, \mathbf{27}]$.

4. Proofs of Theorems 2.3, 2.8, 2.9 and 2.11

We begin by describing completely positive maps using UCP maps.

LEMMA 4.1. Let \mathscr{A} be a C^* -algebra with unit e, \mathcal{H} be a Hilbert space and $\Psi \colon \mathscr{A} \to \mathbf{B}(\mathcal{H})$ be a map. Then the following conditions are equivalent:

- (i) Ψ is completely positive,
- (ii) there exist a UCP map $\Phi: \mathscr{A} \to B(\mathcal{H})$ and a completely positive map $\widetilde{\Psi}: B(\mathcal{H}) \to B(\mathcal{H})$ such that

$$\Psi = \widetilde{\Psi} \circ \Phi, \tag{4.1}$$

(iii) there exist a UCP map $\Phi \colon \mathscr{A} \to \mathbf{B}(\mathcal{H})$ and $R \in \mathbf{B}(\mathcal{H})$ such that $R \geqslant 0$ and $\Psi(a) = R\Phi(a)R$, $a \in \mathscr{A}$.

Moreover, if (iii) holds, then $R = \Psi(e)^{1/2}$.

PROOF. (i) \Rightarrow (iii) By the Stinespring dilation theorem (see [29, Theorem 1]), there exist a Hilbert space \mathcal{K} , an operator $B \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ and a representation $\pi \colon \mathscr{A} \to \mathcal{B}(\mathcal{K})$ such that

$$\Psi(a) = B^* \pi(a) B, \quad a \in \mathscr{A}. \tag{4.2}$$

Since Ψ is positive, $\Psi(e) \ge 0$. We may assume without loss of generality that

$$\dim \overline{\mathscr{R}(\Psi(e))}^{\perp} \leqslant \dim \overline{\mathscr{R}(B)}^{\perp}. \tag{4.3}$$

Indeed, there exists a cardinal number $\mathfrak{n} \geqslant 1$ such that $\mathfrak{n} \cdot \dim \mathcal{K} \geqslant \dim \mathcal{H}$. Set $\mathcal{M} = \bigoplus_{\omega \in \Sigma} \mathcal{K}_{\omega}$ and $\mathcal{K}_{\mathcal{M}} = \mathcal{K} \oplus \mathcal{M}$, where Σ is a set of cardinality \mathfrak{n} and $\mathcal{K}_{\omega} = \mathcal{K}$ for every $\omega \in \Sigma$. Clearly, $\dim \mathcal{M} = \mathfrak{n} \cdot \dim \mathcal{K}$. Define the operator $B_{\mathcal{M}} \in \mathcal{B}(\mathcal{H}, \mathcal{K}_{\mathcal{M}})$ and the representation $\pi_{\mathcal{M}} : \mathscr{A} \to \mathcal{B}(\mathcal{K}_{\mathcal{M}})$ by $B_{\mathcal{M}}h = Bh \oplus 0$ for $h \in \mathcal{H}$ and $\pi_{\mathcal{M}} = \pi \oplus \bigoplus_{\omega \in \Sigma} \pi_{\omega}$ with $\pi_{\omega} = \pi$. Then

$$\dim \overline{\mathscr{R}(\Psi(e))}^{\perp} \leqslant \dim \left(\overline{\mathscr{R}(B)}^{\perp} \oplus \mathcal{M}\right) = \dim \left(\overline{\mathscr{R}(B)} \oplus \{0\}\right)^{\perp} = \dim \overline{\mathscr{R}(B_{\mathcal{M}})}^{\perp}.$$

Clearly (4.2) is valid with $\pi_{\mathcal{M}}$ and $B_{\mathcal{M}}$ in place of π and B, respectively.

Assume that (4.3) holds. Substituting a = e into (4.2), we get

$$||Bh||^2 = ||\Psi(e)^{1/2}h||^2, \quad h \in \mathcal{H}.$$
 (4.4)

Since $\overline{\mathscr{R}(\Psi(e)^{1/2})} = \overline{\mathscr{R}(\Psi(e))}$, we deduce from (4.4) that there exists a unique unitary operator $\widetilde{U} \in B(\overline{\mathscr{R}(\Psi(e))}, \overline{\mathscr{R}(B)})$ such that $Bh = \widetilde{U}\Psi(e)^{1/2}h$ for every $h \in \mathcal{H}$. Hence, by (4.3) there exist an isometry $U \in B(\mathcal{H}, \mathcal{K})$ such that

$$B = U\Psi(e)^{1/2}. (4.5)$$

Define the map $\Phi: \mathscr{A} \to \boldsymbol{B}(\mathcal{H})$ by $\Phi(a) = U^*\pi(a)U$ for $a \in \mathscr{A}$. The map Φ , being a composition of the representation π and the UCP map $\boldsymbol{B}(\mathcal{K}) \ni Y \mapsto U^*YU \in \boldsymbol{B}(\mathcal{H})$, is a UCP map. By (4.2) and (4.5) we have

$$\Psi(a) = B^*\pi(a)B = \Psi(e)^{1/2}\Phi(a)\Psi(e)^{1/2}, \quad a \in \mathscr{A}.$$

This means that (iii) holds with $R = \Psi(e)^{1/2}$.

(iii) \Rightarrow (ii) Let Φ and R be as in (iii). Then the map $\widetilde{\Psi}$ defined by $\widetilde{\Psi}(Y) = RYR$ for $Y \in \mathcal{B}(\mathcal{H})$ satisfies (4.1).

(ii)
$$\Rightarrow$$
(i) This implication is obvious.

PROOF OF THEOREM 2.8. If R is an isometry, then by the unique extension property [23, Theorem B.2(iii)] ([5, Theorem 2.1]) and Stinespring dilation theorem, $J = \mathcal{A}$, i.e., $\pi = R^* \rho R$. This implies that

$$(R^*\rho(a)R)^*(R^*\rho(a)R) = \pi(a)^*\pi(a) = \pi(a^*a) = R^*\rho(a)^*\rho(a)R, \quad a \in \mathscr{A}. \tag{4.6}$$

By [22, Lemma 3.2], we have

$$\rho(a)R = R(R^*\rho(a)R) = R\pi(a), \quad a \in \mathscr{A}, \tag{4.7}$$

so (2.8) is valid. Therefore, we can assume that R is non-isometric. Define the map $\Psi_0 \colon \mathscr{A} \to B(\mathcal{H})$ by $\Psi_0(a) = \phi(a) \triangle$ for $a \in \mathscr{A}$. Since states on C^* -algebras are UCP maps (see [21, Proposition 3.8]), we infer from [21, Lemma 3.10] that the map Ψ_0 is completely positive. Let us define the maps $\Psi_1, \Phi \colon \mathscr{A} \to B(\mathcal{H})$ by $\Psi_1(a) = R^*\rho(a)R$ and $\Phi(a) = \Psi_0(a) + \Psi_1(a)$ for $a \in \mathscr{A}$. By [29, Theorem 1], Ψ_1 is completely positive. Hence, since $\Psi_0(e) = \triangle$, Φ is a UCP map (e is the unit of \mathscr{A}). As $G \subseteq \ker \phi$, we see that $\pi(g) = \Phi(g)$ for all $g \in G$. Therefore, by [23, Theorem B.2(iii)] ([5, Theorem 2.1]), $\pi = \Phi$, i.e.,

$$\pi(a) = \Psi_0(a) + \Psi_1(a), \quad a \in \mathscr{A}. \tag{4.8}$$

Now we prove that R intertwines π and ρ . Take finite sequences $\{a_j\}_{j=1}^n \subseteq \mathcal{A}$ and $\{h_j\}_{j=1}^n \subseteq \mathcal{H}$. Since Ψ_0 is completely positive and the matrix $[a_k^*a_j]_{k,j=1}^n$ is positive, we deduce that $\sum_{j,k=1}^n \langle \Psi_0(a_k^*a_j)h_j,h_k\rangle \geqslant 0$. This together with (4.8) yields

$$\left\| \sum_{j=1}^{n} \pi(a_j) h_j \right\|^2 = \sum_{j,k=1}^{n} \langle \pi(a_k^* a_j) h_j, h_k \rangle$$

$$= \sum_{j,k=1}^{n} \langle \rho(a_k^* a_j) R h_j, R h_k \rangle + \sum_{j,k=1}^{n} \langle \Psi_0(a_k^* a_j) h_j, h_k \rangle$$

$$\geqslant \sum_{j,k=1}^{n} \langle \rho(a_k^* a_j) R h_j, R h_k \rangle = \left\| \sum_{j=1}^{n} \rho(a_j) R h_j \right\|^2.$$

Since $\pi(e) = I_{\mathcal{H}}$, the vectors $\{\pi(a)h : a \in \mathcal{A}, h \in \mathcal{H}\}$ span \mathcal{H} . As a consequence, there exists a unique contraction $\widehat{R} \in \mathcal{B}(\mathcal{H}, \mathcal{K})$ such that $\widehat{R}\pi(a) = \rho(a)R$ for every $a \in \mathcal{A}$. Substituting a = e, we see that $\widehat{R} = R$. Therefore, we have

$$R\pi(a) = \rho(a)R, \quad a \in \mathscr{A}.$$
 (4.9)

Our next goal is to show that J is a closed *-ideal in \mathscr{A} . Clearly, J is a closed selfadjoint vector subspace of \mathscr{A} . It follows from (4.9) that $R^*\rho(a)=\pi(a)R^*$ for every $a\in\mathscr{A}$, which implies that for all $a\in J$ and $b,c\in\mathscr{A}$,

$$\pi(bac) = \pi(b)\pi(a)\pi(c) = \pi(b)R^*\rho(a)R\pi(c) = R^*\rho(b)\rho(a)\rho(c)R = R^*\rho(bac)R.$$
 (4.10)

This shows that J is a closed *-ideal in \mathscr{A} . By (4.8), $\ker \phi \subseteq J$, so (2.8) is valid.

It remains to prove the "moreover" part.

- (i) Since the *-ideal J is proper (as R is non-isometric), $\ker \phi \subseteq J$ and $\ker \phi$ has codimension 1 in \mathscr{A} , we conclude that $\ker \phi = J$ and consequently that ϕ is a character of \mathscr{A} , which completes the proof of (i).
 - (ii) Since (4.9) implies that $R^*\rho(a)=\pi(a)R^*$ for all $a\in\mathscr{A}$, we get

$$R^*R\pi(a) = R^*\rho(a)R = \pi(a)R^*R, \quad a \in \mathscr{A}.$$

so \triangle commutes with π . Similarly, \triangle_* commutes with ρ because

$$\rho(a)RR^* = R\pi(a)R^* = RR^*\rho(a), \quad a \in \mathscr{A},$$

which completes the proof of (ii).

- (iii) This follows from (ii), (4.9) and the assumption that R is non-isometric.
- (iv) Since $R\triangle = \triangle_* R$, we get $R(\mathscr{R}(\triangle)) \subseteq \mathscr{R}(\triangle_* R)$, so $R(\mathcal{H}_1) \subseteq \mathcal{K}_1$. In turn, if $h \in \mathcal{H}_0 = \ker \triangle$, then

$$\langle Rh, \triangle_* Rg \rangle = \langle Rh, R\triangle g \rangle = \langle R^*Rh, \triangle g \rangle = \langle h, \triangle g \rangle = \langle \triangle h, g \rangle = 0, \quad g \in \mathcal{H},$$

which implies that $Rh \in \mathcal{K}_1^{\perp} = \mathcal{K}_0$. Hence $R(\mathcal{H}_0) \subseteq \mathcal{K}_0$. This yields $R = R_0 \oplus R_1$, where $R_j \in \boldsymbol{B}(\mathcal{H}_j, \mathcal{K}_j)$ for j = 0, 1 are given by $R_j h = Rh$ for $h \in \mathcal{H}_j$ and j = 0, 1. That R_0 is isometric follows from the equality $\mathcal{H}_0 = \ker(\triangle)$. If $h \in \mathcal{H}_1$ and $\|R_1h\| = \|h\|$, then $\langle \triangle h, h \rangle = 0$, which together with $\triangle \geqslant 0$ implies that $h \in \mathcal{H}_0$, so h = 0. Thus R_1 is a pure contraction. Since $R = R_0 \oplus R_1$ and $\triangle|_{\mathcal{H}_0} = 0$, we get

$$\triangle_* R = R \triangle = R_0 \triangle |_{\mathcal{H}_0} \oplus R_1 \triangle |_{\mathcal{H}_1} = 0 \oplus R_1 \triangle |_{\mathcal{H}_1}. \tag{4.11}$$

However, $\ker(\triangle|_{\mathcal{H}_1}) = \{0\}$, so $\overline{\mathscr{R}(\triangle|_{\mathcal{H}_1})} = \mathcal{H}_1$, which together with (4.11) yields

$$\mathcal{K}_1 = \{0\} \oplus \overline{R_1(\overline{\mathscr{R}(\triangle|_{\mathcal{H}_1})})} = \{0\} \oplus \overline{\mathscr{R}(R_1)}.$$

As a consequence, R_1 has dense range.

(v) & (vi) It follows from (4.8), (4.9) and (i) that

$$\pi(a) = R^* \rho(a) R + \chi(a) \triangle = \pi(a) R^* R + \chi(a) \triangle, \quad a \in \mathscr{A}. \tag{4.12}$$

This implies that $\pi(a)\triangle = \chi(a)\triangle$ for all $a \in \mathscr{A}$. Therefore, $\pi|_{\mathcal{H}_1} = \chi I_{\mathcal{H}_1}$ and consequently $\pi = \pi_0 \oplus \chi I_{\mathcal{H}_1}$. In turn, by (4.9), (4.12) and (ii), we have

$$R^*\rho(a) = \pi(a)R^* = R^*\rho(a)RR^* + \chi(a)\triangle R^* = R^*RR^*\rho(a) + \chi(a)\triangle R^*, \quad a \in \mathscr{A}.$$
 This yields

$$R^* \triangle_* \rho(a) = \chi(a) \triangle R^* = \chi(a) R^* \triangle_*, \quad a \in \mathscr{A}.$$

Taking adjoints, we see that

$$\rho(a)\triangle_*R = \chi(a)\triangle_*R, \quad a \in \mathscr{A}.$$

This implies that $\rho|_{\mathcal{K}_1} = \chi I_{\mathcal{K}_1}$ and thus $\rho = \rho_0 \oplus \chi I_{\mathcal{K}_1}$. Note that by (4.12),

$$\pi_{0}(a) \oplus \chi(a)I_{\mathcal{H}_{1}} = \pi(a) = R^{*}\rho(a)R + \chi(a)\triangle$$

$$= R_{0}^{*}\rho_{0}(a)R_{0} \oplus R_{1}^{*}\rho(a)|_{\mathcal{K}_{1}}R_{1} + \chi(a)0 \oplus \triangle|_{\mathcal{H}_{1}}$$

$$= R_{0}^{*}\rho_{0}(a)R_{0} \oplus (\chi(a)R_{1}^{*}R_{1} + \chi(a)\triangle|_{\mathcal{H}_{1}})$$

$$= R_{0}^{*}\rho_{0}(a)R_{0} \oplus \chi(a)(R^{*}R|_{\mathcal{H}_{1}} + \triangle|_{\mathcal{H}_{1}})$$

$$= R_{0}^{*}\rho_{0}(a)R_{0} \oplus \chi(a)I_{\mathcal{H}_{1}}, \quad a \in \mathscr{A}.$$

As a consequence, $\pi_0(a) = R_0^* \rho_0(a) R_0$ for all $a \in \mathscr{A}$.

Remark 4.2. It is easy to see that the conclusion of Theorem 2.8 remains valid if instead of $G \subseteq \ker \phi$ we assume that $G \subseteq \bigcap_{\iota \in A} \ker \phi_{\iota}$, where $\{\phi_{\iota}\}_{\iota \in A}$ is a family of states on \mathscr{A} . But then the expression " $\ker \phi \subseteq J$ " in (2.8) must be replaced by " $\ker \phi_{\iota} \subseteq J$ $\forall \iota \in A$ ", and condition (i) must now read " $\ker \phi_{\iota} = J$ and ϕ_{ι} is a character of \mathscr{A} for every $\iota \in A$ ", so $\phi_{\iota} = \phi$ for every $\iota \in A$. In particular, if $\Psi \colon \mathscr{A} \to B(\mathcal{M})$ is a nonzero completely positive map (e.g., a USP map) such that $G \subseteq \ker \Psi$, where \mathscr{M} is any nonzero Hilbert space, then $G \subseteq \bigcap_{f \in \Omega} \ker \phi_f$, where $\Omega := \{f \in \mathcal{H} \colon \|f\| = 1 \ \& \ \langle \Psi(e)f, f \rangle \neq 0\}$ and ϕ_f is the state on \mathscr{A} defined by

$$\psi_f(a) = \frac{1}{\langle \Psi(e)f, f \rangle} \langle \Psi(a)f, f \rangle, \quad a \in \mathcal{A}, f \in \Omega.$$

This is due to the fact that $\ker \Psi = \bigcap_{f \in \Omega} \ker \phi_f$. The inclusion " \subseteq " is obvious, while its converse " \supseteq " follows from the Cauchy-Schwarz inequality:

$$|\langle \Psi(a)f, f \rangle|^2 \leqslant \langle \Psi(a^*a)f, f \rangle \langle \Psi(e)f, f \rangle, \quad f \in \mathcal{H}, a \in \mathcal{A}.$$

PROOF OF THEOREM 2.9. Since $G \subseteq \ker \chi$, it is routine to verify that $\pi(g) = R^* \rho(g) R$ for all $g \in G$. Taking the values of both sides of the equality $\pi_0 = R_0^* \rho_0 R_0$ at the unit of \mathscr{A} , we see that R_0 is an isometry.

To prove the "moreover" part, assume that R_1 is non-isometric.

(i) Note that $a \in \mathcal{A}$ is in J if and only if

$$R_0^* \rho_0(a) R_0 \oplus \chi(a) I_{\mathcal{H}_1} = \pi_0(a) \oplus \chi(a) I_{\mathcal{H}_1}$$

= $\pi(a) = R^* \rho(a) R = R_0^* \rho_0(a) R_0 \oplus \chi(a) R_1^* R_1$,

or equivalently if and only if $\chi(a)(I_{\mathcal{H}_1} - R_1^*R_1) = 0$. Since R_1 is non-isometric, we conclude that $J = \ker \chi$.

Now we show that \triangle_* commutes with ρ . To this end, observe that

$$(R_0^* \rho_0(a) R_0)^* (R_0^* \rho_0(a) R_0) = \pi_0(a)^* \pi_0(a)$$

= $\pi_0(a^* a) = R_0^* \rho_0(a)^* \rho_0(a) R_0, \quad a \in \mathscr{A}.$

By [22, Lemma 3.2], $\rho_0(a)R_0 = R_0R_0^*\rho_0(a)R_0$ for all $a \in \mathscr{A}$. Multiplying by R_0^* on the right side, we see that $\rho_0(a)R_0R_0^* = R_0R_0^*\rho_0(a)R_0R_0^*$ for all $a \in \mathscr{A}$. Taking adjoints, we obtain $R_0R_0^*\rho_0(a) = \rho_0(a)R_0R_0^*$ for all $a \in \mathscr{A}$. Therefore, we have

$$RR^*\rho(a) = R_0 R_0^* \rho_0(a) \oplus \chi(a) R_1 R_1^*$$

= \rho_0(a) R_0 R_0^* \oplus \chi(a) R_1 R_1^* = \rho(a) R R^*, \quad a \in \mathscr{A},

so \triangle_* commutes with ρ . This implies that

$$\pi_0(a)R_0^*R_0 = R_0^*\rho_0(a)(R_0R_0^*)R_0 = R_0^*R_0(R_0^*\rho_0(a)R_0) = R_0^*R_0\pi_0(a), \quad a \in \mathscr{A}.$$

Arguing as above, we conclude that \triangle commutes with π .

(ii) Since R_0 is an isometry, we get

$$\Delta = (I_{\mathcal{H}_0} - R_0^* R_0) \oplus (I_{\mathcal{H}_1} - R_1^* R_1) = 0 \oplus (I_{\mathcal{H}_1} - R_1^* R_1). \tag{4.13}$$

However, R_1 is a pure contraction, so $\ker(I_{\mathcal{H}_1} - R_1^*R_1) = \{0\}$ and consequently $\overline{\mathscr{R}(I_{\mathcal{H}_1} - R_1^*R_1)} = \mathcal{H}_1$, which together with (4.13) yields $\overline{\mathscr{R}(\triangle)} = \mathcal{H}_1$.

(iii) Arguing as in (ii), we obtain

$$\triangle_* R = R \triangle = R_0 (I_{\mathcal{H}_0} - R_0^* R_0) \oplus R_1 (I_{\mathcal{H}_1} - R_1^* R_1) = 0 \oplus R_1 (I_{\mathcal{H}_1} - R_1^* R_1),$$

which implies that

$$\overline{\mathscr{R}(\triangle_*R)} = \{0\} \oplus \overline{R_1(\overline{\mathscr{R}(I_{\mathcal{H}_1} - R_1^*R_1)})} = \{0\} \oplus \overline{\mathscr{R}(R_1)} = \mathcal{K}_1.$$

This completes the proof.

PROOF OF THEOREM 2.3. (i) \Rightarrow (ii) & (i) \Rightarrow (iv) (Assuming $G \subseteq \ker \phi$) Apply Theorem 2.8(i).

(ii) \Rightarrow (i) if G generates \mathscr{A} (NOT ASSUMING $G \subseteq \ker \phi$). Let $\pi \colon \mathscr{A} \to B(\mathcal{H})$ be a representation of \mathscr{A} on a Hilbert space \mathcal{H} and $\widetilde{\Phi} \colon \mathscr{A} \to B(\mathcal{H})$ be a UCP map such that $\widetilde{\Phi}|_G = \pi|_G$. It follows from the Stinespring dilation theorem (see [29, Theorem 1]) that

$$\widetilde{\Phi}(a) = P\rho(a)|_{\mathcal{H}}, \quad a \in \mathscr{A},$$

$$\tag{4.14}$$

where $\rho \colon \mathscr{A} \to B(\mathcal{K})$ is a representation of \mathscr{A} on a Hilbert space \mathcal{K} such that $\mathcal{H} \subseteq \mathcal{K}$, and $P \in B(\mathcal{K})$ is the orthogonal projection of \mathcal{K} onto \mathcal{H} . Applying (ii) to the operator $R \in B(\mathcal{H}, \mathcal{K})$ defined by Rh = h for $h \in \mathcal{H}$, we deduce that $\pi(a) = \rho(a)|_{\mathcal{H}}$ for every $a \in \mathscr{A}$, and thus \mathcal{H} reduces ρ . Therefore, by (4.14), $\widetilde{\Phi}$ is a representation of \mathscr{A} . Since G generates the C^* -algebra \mathscr{A} , we infer from $\widetilde{\Phi}|_G = \pi|_G$ that $\widetilde{\Phi} = \pi$, which means that $\pi|_G$ has the unique extension property. Finally, using [23, Theorem B.2(iii)] ([5, Theorem 2.1]), we obtain (i).

(i) \Rightarrow (iii) (ASSUMING $G \subseteq \ker \phi$) Let (π, ρ, R) be as in (iii) and $\pi(g) = R^* \rho(g) R$ for all $g \in G$. If R is isometric, then using the Stinespring dilation theorem and the unique extension property [23, Theorem B.2(iii)] ([5, Theorem 2.1]) yields $\pi = R^* \rho R$. If R is non-isometric, then applying Theorem 2.8(i), we conclude that there exists $\chi \in \mathfrak{M}_{\mathscr{A}}$ such that $\ker \chi = J$, so $\pi(a) = R^* \rho(a) R$ for every $a \in \ker \chi$.

(ii) \Rightarrow (iii) if G generates \mathscr{A} (NOT ASSUMING $G \subseteq \ker \phi$) Indeed, let (π, ρ, R) be as in (iii) and $\pi(g) = R^*\rho(g)R$ for all $g \in G$. Set $J = \{a \in \mathscr{A} : \pi(a) = R^*\rho(a)R\}$. By (ii), $R\pi(a) = \rho(a)R$ for all $a \in \mathscr{A}$, so $R^*\rho(a) = \pi(a)R^*$ for all $a \in \mathscr{A}$. Hence, arguing as in (4.10), we see that J is a closed *-ideal in \mathscr{A} . If R is an isometry, then $e \in J$, so $J = \mathscr{A}$. Suppose now that R is non-isometric. Then the ideal J is proper. Denote by \mathscr{A}_G the non-unital *-algebra generated by G. By assumption $G \subseteq J$, so $\mathscr{A}_G \subseteq J$. We claim that $\mathscr{A} = J \dotplus \mathbb{C}e$. Take $a \in \mathscr{A}$. Since G generates the unital C^* -algebra \mathscr{A} , there exist sequences $\{b_n\}_{n=1}^\infty \subseteq \mathscr{A}_G \subseteq J$ and $\{\lambda_n\}_{n=1}^\infty \subseteq \mathbb{C}$ such that $a_n := b_n + \lambda_n e \to a$ as $n \to \infty$ (e stands for the unit of \mathscr{A}). First, consider the case when the sequence $\{\lambda_n\}_{n=1}^\infty$ is unbounded. Passing to a subsequence, if necessary, we can assume without loss of generality that $\lim_{n\to\infty} |\lambda_n| = \infty$. Then $\lim_{n\to\infty} \frac{1}{\lambda_n} a_n = 0$, which yields $e = -\lim_{n\to\infty} \frac{1}{\lambda_n} b_n \in J$, so the ideal J is not

proper, a contradiction. Therefore, the sequence $\{\lambda_n\}_{n=1}^{\infty}$ is bounded. In view of Bolzano-Weierstrass theorem, passing to a subsequence if necessary, we can assume without loss of generality that $\lim_{n\to\infty} \lambda_n = \lambda$ for some $\lambda \in \mathbb{C}$. Then the sequence $\{b_n\}_{n=1}^{\infty}$ is convergent and $a = b + \lambda e$, where $b = \lim_{n\to\infty} b_n \in J$. This shows that $\mathscr{A} = J \dotplus \mathbb{C}e$. Hence, there exists a unique character χ of \mathscr{A} such that $\ker \chi = J$.

(iii) \Rightarrow (ii) (NOT ASSUMING $G \subseteq \ker \phi$) Let (π, ρ, R) be as in (ii) and $\pi(g) = R^*\rho(g)R$ for all $g \in G$. By (iii) the then-clause of (2.6) is valid. Consider first the case when $\pi = R^*\rho R$. Then arguing as in (4.6) and (4.7) we see that $\rho(a)R = R\pi(a)$ for every $a \in \mathscr{A}$. The second and only possibility is that

$$\pi(a) = R^* \rho(a) R, \quad a \in \ker \chi, \tag{4.15}$$

where $\chi \in \mathfrak{M}_{\mathscr{A}}$. Since $\ker \chi$ is a *-ideal in \mathscr{A} , if $a \in \ker \chi$, then $a^*a \in \ker \chi$, which together with (4.15) implies that (cf. (4.6))

$$(R^*\rho(a)R)^*(R^*\rho(a)R) = \pi(a)^*\pi(a) = \pi(a^*a) = R^*\rho(a)^*\rho(a)R, \quad a \in \ker \phi.$$

By [22, Lemma 3.2], we get $\rho(a)R = R\pi(a)$ for all $a \in \ker \chi$. Since $\ker \chi$ is a subspace of \mathscr{A} of codimension 1, we deduce that $\rho(a)R = R\pi(a)$ for every $a \in \mathscr{A}$.

(iv) \Leftrightarrow (v) (NOT ASSUMING $G \subseteq \ker \phi$) This equivalence is a direct consequence of (2.4) and the Stinespring dilation theorem.

(iv) \Rightarrow (iii) (NOT ASSUMING $G \subseteq \ker \phi$) This can be verified using the fact that for any $\chi \in \mathfrak{M}_{\mathscr{A}}$, $\mathscr{A} = \ker \chi \dotplus \mathbb{C}e$.

(iv) \Rightarrow (ii) if $\phi \in \mathfrak{M}_{\mathscr{A}}$ (NOT ASSUMING $G \subseteq \ker \phi$) Let us combine implications (iv) \Rightarrow (iii) and (iii) \Rightarrow (ii).

Summarizing, we have proved that (a), (b), (c), (d.1), (d.3) and (d.4) hold. Condition (d.2) can be deduced from Theorem 2.8(i). \Box

COROLLARY 4.3. Let \mathscr{A} be the commutative unital C^* -algebra generated by a single element $t \in \mathscr{A}$. Then, for every character χ of \mathscr{A} , there exists at most a two-element hyperrigid set G of generators of \mathscr{A} such that $G \subseteq \ker \chi$. Moreover, if G is a hyperrigid subset of \mathscr{A} , then there exists at most one character χ of \mathscr{A} such that $G \subseteq \ker \chi$.

PROOF. By [24, Theorem 11.19], there is no loss of generality in assuming that $\mathscr{A}=C(X)$ and $t=\xi$, where X is a nonempty compact subset of $\mathbb C$ and ξ is as in (2.1). Given $\omega\in X$, we can define the map $\tau_\omega\colon X\to X-\omega$ by $\tau_\omega(z)=z-\omega$ for $z\in X$. Then $0\in\tau_\omega(X)$. Set $\widetilde G_\omega=\{\xi_\omega,\bar\xi_\omega\xi_\omega\}$, where ξ_ω is the coordinate function on $\tau_\omega(X)$. By Theorem 2.1, $\widetilde G_\omega$ is a hyperrigid set of generators of $C(\tau_\omega(X))$ and f(0)=0 for every $f\in\widetilde G_\omega$. Define the composition map $\pi_\omega\colon C(\tau_\omega(X))\to C(X)$ by $\pi_\omega(f)=f\circ\tau_\omega$ for $f\in C(\tau_\omega(X))$. Then π_ω is a unital C^* -algebra isomorphism. As a consequence, $G_\omega:=\pi_\omega(\widetilde G_\omega)$ is a hyperrigid set of generators of C(X) (see e.g., [23, Corollary B.3]). It is easy to see that $f(\omega)=0$ for every $f\in G_\omega$, or equivalently $G_\omega\subseteq\ker \chi_\omega$, where χ_ω is the character of C(X) given by $\chi_\omega(f)=f(\omega)$ for $f\in C(X)$. Since characters of C(X) are of the form χ_ω with $\omega\in X$ (see [24, p. 271, Example (a)]), this implies that for every character χ of C(X), there exists a hyperrigid set G of generators of C(X) such that $G\subseteq\ker\chi$.

The "moreover" part follows from Corollary 2.6.

PROOF OF THEOREM 2.11. (i) \Leftrightarrow (ii) First, observe that if $\pi: C(X) \to B(\mathcal{H})$ is a representation, then $\pi(\xi)$ is a normal operator and π takes the form

$$\pi(f) = f(\pi(\xi)), \quad f \in C(X). \tag{4.16}$$

Indeed, $\sigma(\pi(\xi)) \subseteq \sigma(\xi) = X$. Since both sides of the above equality are representations of C(X) which coincide at $f = \xi$ (see (2.1)) and, by the Stone-Weierstrass theorem, ξ generates C(X), we deduce that (4.16) is valid. Also, if $\rho \colon C(X) \to \boldsymbol{B}(\mathcal{K})$ is a representation and $R \in \boldsymbol{B}(\mathcal{H},\mathcal{K})$, then $R\pi(f) = \rho(f)R$ for all $f \in C(X)$ if and only if $R\pi(\xi) = \rho(\xi)R$. This means that with $\mathscr{A} := C(X)$, the conditions (ii) of Theorems 2.3 and 2.11 are equivalent. Defining the state ψ on C(X) by $\psi(f) = \int_X f \mathrm{d}\mu$ for $f \in C(X)$, we infer from Theorem 2.3 that conditions (i) and (ii) are equivalent.

- (ii) \Rightarrow (iii) Apply (ii) to the triplet (\mathcal{K}, R, N) defined by $\mathcal{K} := \mathcal{K}_1 \oplus \cdots \oplus \mathcal{K}_n$, $Rh := R_1 h \oplus \cdots \oplus R_n h$ for $h \in \mathcal{H}$, and $N := N_1 \oplus \cdots \oplus N_n$.
- (iii) \Rightarrow (ii) Set $\mathcal{K}_i = \mathcal{K}$ and $N_i = N$ for i = 1, ..., n, $R_1 = R$ and $R_i = 0$ for i = 2, ..., n. Substituting these objects into (iii) yields NR = RT.
- (i) \Leftrightarrow (iv) This follows from equivalence (i) \Leftrightarrow (iv) of Theorem 2.3 via the above arguments.
- (a) Let us denote by (i^*) - (iv^*) the versions of (i)-(iv) in which the considered Hilbert spaces are assumed to be separable. A close inspection of the above reasonings shows that $(ii^*)\Leftrightarrow (iii^*)$. It is easy to see that (ii^*) implies Theorem 2.2(iii^*), which, by the "moreover" part of this theorem, implies (i^*) , that is $(ii^*)\Rightarrow (i^*)$. Since (i)-(iii) are equivalent, $(iii)\Rightarrow (iii^*)$ and, by [23, Theorem B.2], $(i^*)\Rightarrow (i)$, we conclude that all the conditions (i)-(iii) and
- (b) This can be deduced from what was done above, and from Theorem 2.3(d.2) and the equality $\mathfrak{M}_{C(X)} = X$.

References

- [1] F. Altomare, M. Campiti, Korovkin type approximation theory and its applications, de Gruyter Studies in Mathmatics, Berlin, New York, 1994.
- [2] F. Altomare, Korovkin-type theorems and approximation by positive linear operators, Surveys in Approximation Theory, Vol. 5, 2010, pp. 92-164.
- [3] W. Arveson, The noncommutative Choquet boundary. J. Amer. Math. Soc. 21 (2008), 1065-1084.
- [4] W. Arveson, The noncommutative Choquet boundary III: operator systems in matrix algebras, Math. Scand. 106 (2010), 196-210
- [5] W. Arveson, The noncommutative Choquet boundary II: hyperrigidity, Israel J. Math. 184(2011), 349-385.
- [6] H. Berens, G. G. Lorentz, Geometric theory of Korovkin sets, J. Approx. Theory, 15 (1975), 161-189.
- [7] B. Bilich, Maximality of correspondence representations, arXiv: 2407.04278.
- [8] B. Bilich, A. Dor-On, Arveson's hyperrigidity conjecture is false, arXiv: 2404.05018.
- [9] L. G. Brown, Convergence of functions of self-adjoint operators and applications, Publ. Mat. 60 (2016), 551–564.
- [10] R. Clouâtre, Unperforated pairs of operator spaces and hyperrigidity of operator systems, Canad. J. Math 70 (2018), 1236-1260.
- [11] R. Clouâtre, M. Hartz, Multiplier algebras of complete Nevanlinna-Pick spaces: dilations, boundary representations and hyperrigidity, J. Funct. Anal. 274 (2018), 1690-1738.
- [12] K. Davidson, R. Kenneth, M. Kennedy, Noncommutative choquet theory, arXiv preprint arXiv:1905.08436 (2019).
- [13] K. R. Davidson, M. Kennedy, Choquet order and hyperrigidity for function systems, Adv. Math. 385 (2021), 107774.

- [14] F. Hansen, G. K. Pedersen, Jensen's inequality for operators and Löwner's theorem, Math. Ann. 258 (1982) 229–241.
- [15] W. E. Kaufman, Closed operators and pure contractions in Hilbert space, Proc. Amer. Math. Soc. 87 (1983), 83–87.
- [16] E. T. A. Kakariadis, O. M. Shalit, Operator algebras of monomial ideals in noncommuting variables, J. Math. Anal. Appl. 472 (2019), 738–813.
- [17] M. Kennedy, O. M. Shalit, Essential normality, essential norms and hyperrigidity, J. Funct. Anal. 268 (2015), 2990-3016.
- [18] C. Kleski, Korovkin-type properties for completely positive maps, Illinois J. Math. 58 (2014), 1107-1116.
- [19] P. P. Korovkin, On convergence of linear positive operators in the space of continuous functions. (Russian) Doklady Akad. Nauk SSSR (N.S.) 90 (1953), 961-964.
- [20] G. J. Murphy, C*-algebras and operator theory, Academic Press, Inc., Boston, MA, 1990.
- [21] V. I. Paulsen, Completely bounded maps and operator algebras, Cambridge Studies in Advanced Mathematics, 78, Cambridge University Press, Cambridge, 2002.
- [22] P. Pietrzycki, J. Stochel, On nth roots of bounded and unbounded quasinormal operators, Ann. Mat. Pura. Appl. 202 (2023), 1313-1333.
- [23] P. Pietrzycki, J. Stochel, Hyperrigidity I: singly generated commutative C^* -algebras, submitted (2024).
- [24] W. Rudin, Functional analysis, McGraw-Hill Series in Higher Math., McGraw-Hill Book Co., New York, 1973.
- [25] G. Salomon, Hyperrigid subsets of Cuntz-Krieger algebras and the property of rigidity at zero, J. Operator Theory 81 (2019), 61-79.
- [26] J. A. Šaškin, The Milman-Choquet boundary and the theory of approximations. Funkcional. Anal. i Priložen. 1 (1967), 95-96.
- [27] K. Schmüdgen, Unbounded self-adjoint operators on Hilbert space, Graduate Texts in Mathematics, 265, Springer, Dordrecht, 2012.
- [28] P. Shankar, Hyperrigid generators in C*-algebras, J. Anal. 28 (2020), 791–797.
- [29] W. F. Stinespring, Positive functions on C*-algebras, Proc. Amer. Math. Soc. 6 (1955), 211–216.
- [30] J. Stochel, Decomposition and disintegration of positive definite kernels on convex *-semigroups, Ann. Polon. Math. 56 (1992), 243–294.
- [31] B. Sz.-Nagy, C. Foias, H. Bercovici, L. Kérchy, Harmonic analysis of operators on Hilbert space, Springer, New York, 2010.
- [32] J. Weidmann, Linear operators in Hilbert spaces, Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [33] I. Thompson, An approximate unique extension property for completely positive maps, J. Funct. Anal. 286 (2024) 110193.

Wydział Matematyki i Informatyki, Uniwersytet Jagielloński, ul. Łojasiewicza 6, PL-30348 Kraków

Email address: pawel.pietrzycki@im.uj.edu.pl

Wydział Matematyki i Informatyki, Uniwersytet Jagielloński, ul. Łojasiewicza 6, PL-30348 Kraków

Email address: jan.stochel@im.uj.edu.pl