

A NEW CLASS OF GEOMETRICALLY DEFINED HYPERGRAPHS ARISING FROM THE HADWIGER-NELSON PROBLEM*

Sean Fiscus

Duke University
sean.fiscus@duke.edu

Eric Myzelev

University of Pennsylvania
myzelev@sas.upenn.edu

Hongyi Zhang

Haverford College
hzhang3@haverford.edu

Abstract

There is a famous problem in geometric graph theory to find the chromatic number of the unit distance graph on Euclidean space; it remains unsolved. A theorem of Erdős and De-Bruijn simplifies this problem to finding the maximum chromatic number of a finite unit distance graph. Via a construction built on sequential finite graphs obtained from a generalization of this theorem, we have found a class of geometrically defined hypergraphs of arbitrarily large edge cardinality, whose proper colorings exactly coincide with the proper colorings of the unit distance graph on \mathbb{R}^d . We also provide partial generalizations of this result to arbitrary real normed vector spaces.

1 Introduction

\mathbb{Z}^+ will denote the set of positive integers. A hypergraph is a pair $\mathcal{H} = (V, E)$ in which V is a non-empty set, the set of vertices of \mathcal{H} , and $E \subseteq 2^V$ (i.e. E is a set of subsets of V) satisfying $e \in E$ implies $|e| \geq 2$. (In other definitions, singletons may be allowed in E and/or E may be a multi-set.) E is the set of hyperedges, or just edges, of \mathcal{H} . A proper coloring of \mathcal{H} is a function $\varphi : V \rightarrow C =$ some set of colors, such that

*This work was supported by NSF DMS grant no. 1950563

no $e \in E$ is “monochromatic with respect to φ ”. This means that for each $e \in E$, $\varphi|_e$ is not constant. The chromatic number of \mathcal{H} is the smallest cardinality $|C|$ such that there is a proper coloring $\varphi : V \rightarrow C$.

If $\mathcal{H} = (V, E)$ is a hypergraph and $U \subseteq V$, $|U| \geq 2$, the subhypergraph of \mathcal{H} induced by U is $\mathcal{H}|_U = (U, E \cap 2^U)$. That is, the vertex set of $\mathcal{H}|_U$ is U and the edges of $\mathcal{H}|_U$ are the edges of \mathcal{H} that lie in U . A hypergraph $\mathcal{H} = (V, E)$ in which $|e| = 2$ for every $e \in E$ is a simple graph. The De Bruijn-Erdős theorem [1] concerning the chromatic numbers of infinite graphs is as follows:

If $m \in \mathbb{Z}^+$, $\mathcal{H} = (V, E)$ is a graph, and $\chi(\mathcal{H}|_F) \leq m$ for every finite set $F \subseteq V$, then $\chi(\mathcal{H}) \leq m$.

To prove our results we shall need a known generalization of this theorem to hypergraphs, which will be provided in the next section.

2 De Bruijn-Erdős for hypergraphs with finite edges

The following theorem generalizes the De Bruijn-Erdős Theorem to hypergraphs. A different proof may be found in [2], Chapter 26.

Theorem 2.1 (D-E). *Suppose that $m \in \mathbb{Z}^+$, $\mathcal{H} = (V, E)$ is a hypergraph with $2 \leq |e| < \infty$ for every $e \in E$, and $\chi(\mathcal{H}|_F) \leq m$ for every finite subset F of V . Then $\chi(\mathcal{H}) \leq m$.*

Remark 2.1.1. Since enlarging the edges of a hypergraph – i.e. putting new vertices in some of the edges to join those that were there to begin with – would seem to make it easier to avoid monochromatic edges when coloring the vertices, we are wondering if we really need the hypothesis, in Theorem D-E, that all edges of \mathcal{H} are finite subsets of V . In the proof to come the hypothesis is actually used, and we do not see a way to avoid that use, except by trading the hypothesis for other, clumsier hypotheses. Is there a mathematical logician in the house?

Proof. Suppose that $\mathcal{H} = (V, E)$, and $m \in \mathbb{Z}^+$ satisfy the hypothesis of Theorem D-E. Let $[m] = \{1, \dots, m\}$. A coloring of V with colors $1, \dots, m$ is an element of $X = [m]^V = \prod_{v \in V} [m]$, the Cartesian product of $[m]$ with itself $|V|$ times. Let $[m]$ have the discrete topology, in which every singleton is an open set. Since $[m]$ is finite, obviously $[m]$ is compact (in this or any other topology). By the Tychonoff theorem [3], X , with the product topology, is compact.

In the usual definition of the product topology on X , the basic open neighborhoods of a coloring $\varphi \in X$ are the sets $N(\varphi, F) = \{\psi \in X \mid \psi|_F = \varphi|_F\}$, $F \in \mathcal{F}(V) = \{\text{finite subsets of } V\}$. For $F \in \mathcal{F}(V)$, let $Y_F = \{\varphi \in X \mid \varphi|_F : F \rightarrow [m] \text{ is a proper coloring of } \mathcal{H}|_F\}$. Since $\chi(\mathcal{H}|_F) \leq m$, Y_F is nonempty. Also, Y_F is closed in X : if $\psi \in X \setminus Y_F$, then $\psi|_F$ is not a proper coloring of $\mathcal{H}|_F$, so for some $e \in E \cap 2^F$, $\psi|_e$ is constant (i.e., the vertices in e are all assigned the same $j \in [m]$ by ψ). Any coloring in $N(\psi, e)$ will assign that same j to each vertex of e ; therefore, $N(\psi, e) \subseteq X \setminus Y_F$. Thus $X \setminus Y_F$ is open in the product topology, so Y_F is closed.

If $F_1, \dots, F_t \in \mathcal{F}(V)$, then $F_1 \cup \dots \cup F_t \in \mathcal{F}(V)$, and we have

$$\emptyset \neq Y_{\cup_{i=1}^t F_i} \subseteq Y_{F_1} \cap \dots \cap Y_{F_t}$$

Thus $\{Y_F \mid F \text{ is in } \mathcal{F}(V)\}$ has the finite intersection property: any intersection of finitely many sets from the family is non-empty.

In a compact topological space, the intersection of all the sets in a collection of closed sets with the finite intersection property is non-empty. Therefore $\bigcap_{F \in \mathcal{F}(V)} Y_F \neq \emptyset$. Suppose $\varphi \in \bigcap_{F \in \mathcal{F}(V)} Y_F$. We claim that $\varphi : V \rightarrow [m]$ is a proper coloring of \mathcal{H} . Suppose that $e \in E$. Then e is a finite subset of V . Since e is an edge of $\mathcal{H} \mid_e$, $\varphi \in Y_e$ implies that φ assigns at least two colors from $[m]$ to the elements of e . \square

3 Hypergraphs Equivalent to the Unit Distance Graph

The motivation for this result is the following question: Does there exist a finite set of triangles S in \mathbb{R}^d such that the number of colors in a coloring of \mathbb{R}^d required to forbid monochromatic copies of triangles in S is the same as the chromatic number of the unit Euclidean distance graph on \mathbb{R}^d ? The answer is yes. In fact, we prove a stronger result: in addition to generalizing triangles to arbitrary m -point sets, we also show that there is such a set S so that a coloring φ of \mathbb{R}^d forbids congruent copies of m -gons in S if and only if φ forbids unit distance. To make this precise, we introduce a notion of equivalence of hypergraphs.

Definition 3.1. Let S be a set, and $\mathcal{H} = (S, E_{\mathcal{H}}), \mathcal{G} = (S, E_{\mathcal{G}})$ be hypergraphs where S is the vertex set. We say \mathcal{H} is equivalent to \mathcal{G} if $\chi(\mathcal{H}) = \chi(\mathcal{G})$ and $\varphi : S \rightarrow C$, such that $|C| = \chi(\mathcal{H}) = \chi(\mathcal{G})$, is a proper coloring of \mathcal{H} if and only if φ is a proper coloring of \mathcal{G} .

For the purposes of this paper, we define an m -gon as an arbitrary set of cardinality m . This means that some vertices of the m -gon can be collinear, which contradicts the standard geometric definition.

We also define a unit m -gon to be an m -gon in \mathbb{R}^d as defined above with the following additional property: there exists at least one pair of points x, y in m which are Euclidean distance 1 apart.

Theorem 3.1. Let $M \subset 2^{\mathbb{R}^d}$ be a non-empty finite set of m -gons, for some $m \geq 2$. Define $\mathcal{H}(M)$ as the hypergraph on \mathbb{R}^d with edge set $E = \{X \subset \mathbb{R}^d \mid X \text{ is congruent in } \mathbb{R}^d \text{ to some } T \in M\}$.

Then there must exist some finite set S of $(m+1)$ -gons, such that $\mathcal{H}(M)$ is equivalent to $\mathcal{H}(S)$, where $\mathcal{H}(S)$ is an $(m+1)$ -uniform hypergraph with edge set $E = \{Y \subset \mathbb{R}^d \mid Y \text{ is congruent in } \mathbb{R}^d \text{ to some element of } S\}$.

Proof. We shall proceed by recursively obtaining sets $S_1, S_2, \dots; F_1, F_2, \dots$ satisfying:

1. $S_1 \subseteq S_2 \subseteq \dots$
2. Each S_j is a finite set of $(m+1)$ -gons in \mathbb{R}^d , with each $(m+1)$ -gon containing some m -gon $X \in M$ as a subset of its points.
3. Defining $E_j := \{\text{congruent copies in } \mathbb{R}^d \text{ of the } (m+1)\text{-gons in } S_j\}$ and $\mathcal{H}_j := (\mathbb{R}^d, E_j)$, we obtain F_j , a finite subset of $\mathbb{R}^d \setminus M$ such that $\chi(\mathcal{H}_j \mid_{F_j}) = \chi(\mathcal{H}_j)$.
4. $S_{j+1} = S_j \cup \{X \cup \{z\} \mid z \in F_j, X \in M\}$.

Before giving the recursion, let us note that if the $S_j, F_j, \mathcal{H}_j = (\mathbb{R}^d, E_j)$ are as above then we have the following observations.

- (i) Since, for each $j = 1, 2, \dots$ and $e \in E_j$, e contains a copy of some $X \in M$, $\chi(\mathcal{H}_j) \leq \chi(\mathcal{H}(M))$.
- (ii) In view of the definition of E_j , $S_1 \subseteq S_2 \subseteq \dots$ implies that $E_1 \subseteq E_2 \subseteq \dots$, and thus $\chi(\mathcal{H}_1) \leq \chi(\mathcal{H}_2) \leq \dots$
- (iii) $\chi(\mathcal{H}(M))$ is finite. To see this, observe that in \mathbb{R}^d with the Euclidean norm $\| \cdot \|$, the sets $\{x, y\}$ congruent to a two-set $\{u, v\}$ are just the sets satisfying $\|x - y\| = \|u - v\|$. Therefore, a coloring of \mathbb{R}^d forbidding congruent copies of $\{u, v\}$ is, in other jargon, the same as a coloring which forbids the distance $\|u - v\|$. For every positive distance, the smallest number of colors needed to forbid that distance is $\chi(\mathbb{R}^d, 1)$, the chromatic number of the Euclidean unit distance graph on \mathbb{R}^d .
Suppose that $|M| = n$ (recall that M is finite). From each $T \in M$ select a 2-set D ; let the 2-sets selected be D_1, D_2, \dots, D_n and the distances determined by the 2 points in these sets be d_1, \dots, d_n . For each $i = 1, \dots, n$, let φ_i be a coloring of \mathbb{R}^d with $\chi(\mathbb{R}^d, 1)$ colors that forbids the distance d_i . Now color \mathbb{R}^d by an assignment ψ of n-tuples: $\psi(r) = (\varphi_1(r), \dots, \varphi_n(r))$. We now have colored \mathbb{R}^d with $\chi(\mathbb{R}^d, 1)^n < \infty$ colors, and it is easy to see that ψ is a proper coloring of $\mathcal{H}(M)$: For any $T' \subseteq \mathbb{R}^d$ congruent to some $T \in M$, T' will contain a doubleton congruent to one of the D_i ; to the two vectors in that doubleton, φ_i will assign different colors, which means that ψ will assign different n-tuples to the 2 vectors, which means that T' is not monochromatic.
- (iv) By (i), (iii), and Theorem D-E, it follows that for each j there is a finite set $F_j \in \mathbb{R}^d$ such that $\chi(\mathcal{H}_j|_{F_j}) = \chi(\mathcal{H}_j)$.
- (v) Suppose ρ is an isometry of \mathbb{R}^d and F is a finite non-empty subset of \mathbb{R}^d . Because ρ maps each $e \in E_j$ to an edge $\rho(e) \in E_j$, $\chi(\mathcal{H}_j|_F) = \chi(\mathcal{H}_j|_{\rho(F)})$.

The recursion:

Let $S_1 = \{X \cup \{a\} \mid X \in M\}$, for some $a \in \mathbb{R}^d \setminus \bigcup_{T \in M} T$. Let $F_1 \subseteq \mathbb{R}^d$ be a finite set such that $\chi(\mathcal{H}_1|_{F_1}) = \chi(\mathcal{H}_1)$; as explained in (iv) above, Theorem D-E guarantees the existence of such an F_1 , and by (v), we can assume $F_1 \cap (\bigcup_{T \in M} T) = \emptyset$; if $F_1 \cap (\bigcup_{T \in M} T) \neq \emptyset$, replace F_1 by a translate of itself.

From there, the recursion is dictated in 3 and 4: From S_j we get E_j ; the existence of F_j is guaranteed. Then we define S_{j+1} by 4, above, and roll on.

Since the integer sequence $(\chi(\mathcal{H}_j))_j$ is non-decreasing and bounded above by $\chi(\mathcal{H}(M))$, clearly it will be eventually constant. If that eventual constant value were $\chi(\mathcal{H}(M))$, we would be almost done, except for showing that every proper coloring of \mathcal{H}_k also properly colors $\mathcal{H}(M)$. All will be accomplished by the following.

Clearly $\chi(\mathcal{H}_1) \geq 2 > 1$ whereas $\chi(\mathcal{H}_k) \leq \chi(\mathcal{H}(M)) \leq k$ for $k \geq \chi(\mathcal{H}(M))$. Therefore there is a first value of $k \in \mathbb{Z}^+$ such that $\chi(\mathcal{H}_k) \leq k$.

We have $k > 1$ and

$$k - 1 < \chi(\mathcal{H}_{k-1}) \leq \chi(\mathcal{H}_k) \leq k$$

whence $\chi(\mathcal{H}_{k-1}) = \chi(\mathcal{H}_k) = k \leq \chi(\mathcal{H}(M))$.

Let $\varphi: \mathbb{R}^d \rightarrow \{1, \dots, k\}$ be a proper coloring of \mathcal{H}_k , and since $S_{k-1} \subset S_k$, φ is also a proper coloring of \mathcal{H}_{k-1} . If φ is a proper coloring of $\mathcal{H}(M)$, then $\chi(\mathcal{H}(M)) = k$, and, since φ is an arbitrarily chosen proper coloring of \mathcal{H}_k , the claim of this theorem will be affirmed, with $S = S_k$.

Suppose, on the contrary, that φ does not properly color $\mathcal{H}(M)$, implying that for some $X' = \{a_1, \dots, a_m\}$ a congruent copy of some $X \in M$, $\varphi(a_1) = \varphi(a_2) = \dots = \varphi(a_m)$. We can, without loss of generality, convene that $\varphi(a_1) = \dots = \varphi(a_m) = k$. Let $\rho: \mathbb{R}^d \rightarrow \mathbb{R}^d$ be an isometry such that $\rho(X) = X'$. Consider any point $\rho(z) \in \rho(F_{k-1})$, and note that if it was colored with color k then $X' \cup \{\rho(z)\}$ would be monochromatic and congruent to $X \cup \{z\}$, an edge in \mathcal{H}_{k-1} , which is impossible since φ properly colors \mathcal{H}_{k-1} . Thus φ is a proper coloring of $\mathcal{H}_{k-1} |_{F_{k-1}}$ with $k - 1$ colors. This means that $\chi(\mathcal{H}_{k-1}) = \chi(\mathcal{H}_{k-1} |_{F_{k-1}}) \leq k - 1$, and contradicts the fact that $\chi(\mathcal{H}_{k-1}) = k$. \square

Corollary 3.1.1. *For all integers $m \geq 2$, there exists a finite set S of unit m -gons, such that $\mathcal{H}(S)$ is equivalent to $(\mathbb{R}^d, 1)$, where $\mathcal{H}(S) = \mathcal{H}(\mathbb{R}^d, E)$ and $E = \{X \subset \mathbb{R}^d \mid X \text{ is congruent in } \mathbb{R}^d \text{ to some element of } S\}$ and $(\mathbb{R}^d, 1)$ is the Euclidean unit distance graph on \mathbb{R}^d .*

Proof. We prove the statement by induction. When $m = 2$, the corollary is trivially true.

Now consider $m > 2$ and suppose that the corollary holds true for $m - 1$. Then there exists a finite set S_{m-1} of $m - 1$ -gons as asserted in the Corollary. By the Theorem, there exists a finite set S_m of m -gons such that $\chi(\mathcal{H}(S_m)) = \chi(\mathcal{H}(S_{m-1}))$, and any proper coloring of $\mathcal{H}(S_m)$ with $\chi(\mathcal{H}(S_m))$ colors is also a proper coloring of $\mathcal{H}(S_{m-1})$.

According to the inductive hypothesis, $\chi(\mathcal{H}(S_{m-1})) = \chi(\mathbb{R}^d, 1)$, and any proper coloring of $\mathcal{H}(S_{m-1})$ with $\chi(\mathcal{H}(S_{m-1}))$ colors is also a proper coloring of the Euclidean unit distance graph on \mathbb{R}^d . Thus, $\chi(\mathcal{H}(S_m)) = \chi(\mathbb{R}^d, 1)$, and any proper coloring of $\mathcal{H}(S_m)$ with $\chi(\mathcal{H}(S_m))$ colors is also a proper coloring of the Euclidean unit distance graph on \mathbb{R}^d .

Also note that because there are two points distance 1 apart in all of the $m - 1$ -gons in S_{m-1} , for every m -gon in S_m , there must also be two points distance 1 apart. That is, S_m is a finite set of unit m -gons. \square

4 Generalizing to Non-Euclidean Norms on \mathbb{R}^d

In the preceding sections, distance in \mathbb{R}^d was provided by the Euclidean norm, hereinafter to be denoted as $\|\cdot\|_2$. Some, but not all, of Theorem 3.1 and its corollary survives generalization to the setting of a finite-dimensional normed vector space over

\mathbb{R}^d . Without loss of generality, the vector space will be \mathbb{R}^d and the norm will be denoted as $\|\cdot\|$.

Two sets $X, Y \subseteq \mathbb{R}^d$ are congruent copies of each other in $(\mathbb{R}^d, \|\cdot\|)$ if and only if one of them is the image of the other under a composition, in either order, of a surjective linear isometry of $(\mathbb{R}^d, \|\cdot\|)$ and a translation. With this definition of congruence, we lose one of the support beams to our geometric intuition that may seem essential to the proof of Theorem 3.1: there can exist $u, v, x, y \in \mathbb{R}^d$ such that $\|u - v\| = \|x - y\| > 0$ and yet $\{u, v\}$ and $\{x, y\}$ are not congruent.

However, note: if $\{u, v\}$ and $\{x, y\}$ are congruent, then $\|u - v\| = \|x - y\|$. Consequently, if φ is a coloring of $(\mathbb{R}^d, \|\cdot\|)$ which forbids a distance $a > 0$, then $|\varphi(e)| > 1$ if $e \subseteq \mathbb{R}^d$ contains two points a distance a apart.

For $a > 0$ let $\chi((\mathbb{R}^d, \|\cdot\|), a)$ denote the smallest $|C|$ such that some coloring $\varphi: \mathbb{R}^d \rightarrow C$ forbids the distance a . By the properties of norms, it is clear that $\chi((\mathbb{R}^d, \|\cdot\|), a) = \chi((\mathbb{R}^d, \|\cdot\|), 1)$ for all $a > 0$. Also very important for our generalizations:

Lemma 4.1. *For all $\|\cdot\|$, $\chi((\mathbb{R}^d, \|\cdot\|), 1) < \infty$.*

This is well known, but we will provide a proof outline in an Appendix.

From Lemma 4.1 and the definition of congruence we obtain, as in section 3, and by the same argument, the following.

Lemma 4.2. *Suppose that \mathcal{S} is a finite collection of subsets of \mathbb{R}^d , each subset with at least 2 elements, and $\mathcal{H}(\mathcal{S}) = (\mathbb{R}^d, E(\mathcal{S}))$ is defined by $E(\mathcal{S}) = \{T \subseteq \mathbb{R}^d \mid T \text{ is congruent to some } S \in \mathcal{S}\}$. Then $\chi(\mathcal{H}(\mathcal{S})) < \infty$.*

Theorem 4.3. *For any norm $\|\cdot\|$ on \mathbb{R}^d , Theorem 3.1 holds with the Euclidean norm replaced by $\|\cdot\|$, provided the phrase “congruent in \mathbb{R}^d ” is replaced by “congruent in $(\mathbb{R}^d, \|\cdot\|)$.”*

In view of Lemmas 4.1 and 4.2, the proof is straightforward; follow the path of argument in section 3.

However, generalizing Corollary 3.1.1 looks to us like a lost cause. In the case $\|\cdot\| = \|\cdot\|_2$, for any two unit vectors u, v , there is a linear isometry of $(\mathbb{R}^d, \|\cdot\|_2)$ that takes u into v , whence $\{0, u\}$ and $\{0, v\}$ are congruent. Thus the graph $((\mathbb{R}^d, \|\cdot\|_2), 1)$ is the hypergraph $\mathcal{H}(\{0, u\})$, which makes the base of the induction proof of the Corollary, when $m = 2$, “trivial”.

Given a non-Euclidean norm $\|\cdot\|$ on \mathbb{R}^d , we can, with reference to Theorem 4.3, find $\mathcal{S}_2 \subseteq \mathcal{S}_3 \subseteq \dots$ such that \mathcal{S}_m is a finite set of unit m -gons in $(\mathbb{R}^d, \|\cdot\|)$, $m = 2, 3, \dots$, and the hypergraphs $\mathcal{H}(\mathcal{S}_m)$ are all equivalent, but we can only be sure that $\chi(\mathcal{H}(\mathcal{S}_m)) \leq \chi((\mathbb{R}^d, \|\cdot\|), 1)$, and even if we happen to have equality, we see no way of assuring that every proper coloring of $\chi(\mathcal{S}_m)$ with number of colors equal to the chromatic number of graph will also properly color the unit distance graph on $(\mathbb{R}^d, \|\cdot\|)$.

On the other hand, we have no proof that there is no generalization of Corollary 3.1.1 for some or all non-Euclidean norms $\|\cdot\|$ on \mathbb{R}^d , $d > 1$. Perhaps the question is worth investigating. (Yes, we are aware that Corollary 3.1.1 will hold for every norm arising from an inner product on \mathbb{R}^d ; \mathbb{R}^d with such a norm is linearly and isometrically isomorphic to $(\mathbb{R}^d, \|\cdot\|_2)$.)

We can obtain m -uniform hypergraphs in $(\mathbb{R}^d, \|\cdot\|)$ with the same chromatic number as $((\mathbb{R}^d, \|\cdot\|), 1)$ if we abandon the practice of defining edge sets as the sets congruent to one of a finite set of m -gons. However, our method does not provably produce hypergraphs equivalent to the unit distance graph on $(\mathbb{R}^d, \|\cdot\|)$.

Let Q be a collection of finite subsets of \mathbb{R}^d , each with at least 2 elements, and let

$$E_0 = \{e \subseteq \mathbb{R}^d \mid \text{for some } f \in Q, e \text{ and } f \text{ are congruent}\}$$

For each $t \in \mathbb{Z}^+$, let

$$B_t = \{f \cup T \mid f \in Q, T \subseteq \mathbb{R}^d, f \cap T = \emptyset, \text{ and } |T| = t\}$$

$$E_t = \{e \subseteq \mathbb{R}^d \mid \text{for some } g \in B_t, e \text{ and } g \text{ are congruent}\}$$

It will be important to notice that because, for each $f \in Q$, we form infinitely many sets in B_t by taking the union of f with each and every t -subset of $\mathbb{R}^d \setminus f$, it follows that if $e \in E_0$, $T' \subseteq \mathbb{R}^d \setminus e$, and $|T'| = t$, then $e \cup T' \in E_t$.

Theorem 4.4. *With $t \in \mathbb{Z}^+$, Q, B_t, E_0 , and E_t as above, let $\mathcal{H}_0 = (\mathbb{R}^d, E_0)$ and $\mathcal{H}_t = (\mathbb{R}^d, E_t)$. If $\chi(\mathcal{H}_0) < \infty$ then $\chi(\mathcal{H}_t) = \chi(\mathcal{H}_0)$.*

Proof. Since each $e \in E_t$ contains an $e' \in E_0$, it follows that a proper coloring of \mathcal{H}_0 will also serve as a proper coloring of \mathcal{H}_t . Thus,

$$\chi(\mathcal{H}_t) \leq \chi(\mathcal{H}_0) < \infty$$

By the same argument,

$$\chi(\mathcal{H}_t) \leq \chi(\mathcal{H}_{t-1}) \leq \chi(\mathcal{H}_0)$$

It suffices to show that $\chi(\mathcal{H}_t) = \chi(\mathcal{H}_{t-1})$ for each $t \in \mathbb{Z}^+$.

Let $k = \chi(\mathcal{H}_t)$ and suppose that $k < \chi(\mathcal{H}_{t-1})$. Let $\varphi: \mathbb{R}^d \rightarrow \{1, \dots, k\}$ be such that no edge $g \in E_t$ is monochromatic, with reference to φ .

Since $k < \chi(\mathcal{H}_{t-1})$, there must exist some $B \in E_{t-1}$ such that φ assigns the same color to every element of B . Without loss of generality, we can assume that this color is k .

For every $u \in \mathbb{R}^d \setminus B$, $B \cup \{u\} \in E_t$. Therefore

$\varphi(u) \in \{1, \dots, k-1\}$; otherwise, $B \cup \{u\}$ would be monochromatic under the coloring φ . Thus, φ restricted to $\mathbb{R}^d \setminus B$ is a proper coloring of $\mathcal{H}_t|_{\mathbb{R}^d \setminus B}$ with $k-1$ colors.

We shall now use Theorem $D-E$ to prove the existence of a proper coloring of \mathcal{H}_t with colors $\{1, \dots, k-1\}$ which will contradict $k = \chi(\mathcal{H}_t)$. Since this contradiction descends from the assumption that

$k = \chi(\mathcal{H}_t) < \chi(\mathcal{H}_{t-1})$ it will follow that $\chi(\mathcal{H}_t) = \chi(\mathcal{H}_{t-1})$ and the theorem will be proven.

Suppose $F \subset \mathbb{R}^d$ is finite and $\chi(\mathcal{H}_t|_F) = \chi(\mathcal{H}_t)$. We aim to show that $\mathcal{H}_t|_F$ can be properly colored with no more than $k-1$ colors, which will imply that

$$\chi(\mathcal{H}_t) = \chi(\mathcal{H}_t|_F) \leq k-1 < k = \chi(\mathcal{H}_t).$$

Let $v \in \mathbb{R}^d$ be such that $(v+F) \cap B = \emptyset$. Then φ colors $v+F$ with no more than

$k - 1$ colors so that for each $\alpha \in 2^{v+F} \cap E_t$, φ assigns more than one color to the elements of α . Now color F as follows: color $f \in F$ with $\varphi(v + f)$. Since every translate of every $\alpha \in E_t$ is in E_t , and no $\alpha \in 2^{v+F} \cap E_t$ is monochromatic under coloring by φ , we have what we wanted, a proper coloring of $\mathcal{H}_t|_F$ with colors from $\{1, \dots, k - 1\}$. \square

Corollary 4.4.1. *Let $\|\cdot\|$ be a norm on \mathbb{R}^d . For each $m \in \mathbb{Z}^+$ such that $m > 2$, define $E_m = \{T \subseteq \mathbb{R}^d \mid |T| = m \text{ and } T \text{ contains 2 points } \|\cdot\| \text{-distance 1 apart}\}$. Let $\mathcal{G}_m = (\mathbb{R}^d, E_m)$. Then $\chi(\mathcal{G}_m) = \chi((\mathbb{R}^d, \|\cdot\|), 1)$ for all m .*

Proof. If $Q = \{\{\underline{0}, u\} \mid u \in \mathbb{R}^d \text{ and } \|u\| = 1\}$, then, in terms used in the Theorem's statement, $\mathcal{H}_0 = ((\mathbb{R}^d, \|\cdot\|), 1)$, the unit distance graph on $(\mathbb{R}^d, \|\cdot\|)$ and, for each $m > 2$, $\mathcal{H}_{1-2} = \mathcal{G}_m$. The conclusion follows from the Theorem. \square

5 Explicit Construction of Hypergraph

In this section, we are in \mathbb{R}^d with the usual Euclidean norm. We will give a “constructive” proof of a weaker version of Theorem 3.1.

By applying Theorem D-E for hypergraphs to Theorem 3.1, it can be shown that for every finite m -uniform hypergraph in \mathbb{R}^d , there must exist a finite $(m + 1)$ -uniform hypergraph of equal chromatic number. However, this deduction is non-constructive because all known proofs of Theorem D-E use the axiom of choice, and thus we have no control over what the finite hypergraph might look like. In this section, we show how to take a finite m -uniform hypergraph and construct a finite $(m + 1)$ -uniform hypergraph of equal chromatic number. This allows us to “construct”, for any $m \in \mathbb{Z}^+$, $m > 2$, a finite m -uniform hypergraph in \mathbb{R}^d with chromatic number equal to $\chi(\mathbb{R}^d, 1)$. We use “construct” in quotations, however, because this hypergraph construction will use a finite unit distance graph G with chromatic number $\chi(\mathbb{R}^d, 1)$.

Theorem 5.1. *Let \mathcal{H} be a finite, m -uniform hypergraph with vertices in \mathbb{R}^d . There exists a finite, $(m + 1)$ -uniform hypergraph \mathcal{H}' with vertices in \mathbb{R}^d such that $\chi(\mathcal{H}') = \chi(\mathcal{H})$.*

Proof. Let $k = \chi(\mathcal{H})$, and F be the vertex set $V(\mathcal{H})$. Note that for all translates $F + v \subset \mathbb{R}^d$ of F where $v \in \mathbb{R}^d$, the chromatic number of corresponding hypergraph $\mathcal{H} + v$, with vertex set $V(\mathcal{H} + v) = F + v$ and $E(\mathcal{H} + v) = \{e + v \mid e \in E(\mathcal{H})\}$, must also be k .

We now construct an $(m + 1)$ -uniform hypergraph \mathcal{H}' in the following way:

1. Let the vertex set of \mathcal{H}' be the union of k disjoint translates of $F = V(\mathcal{H})$, which we will call F_1, F_2, \dots, F_k . Denote by \mathcal{H}_i the translate of \mathcal{H} onto F_i .
2. Define the edge set of \mathcal{H}' by

$$E(\mathcal{H}') = \{\{e \cup v\} \mid e \in \mathcal{H}_i, v \in F_j, \text{ where } j > i\}$$

In other words, $E(\mathcal{H}')$ consists of all $(m + 1)$ -gons such that m points are from an edge in \mathcal{H}_i and the remaining point is from some F_j with $j > i$.

By construction, \mathcal{H}' is finite and $(m+1)$ -uniform. Also, any proper coloring φ of \mathcal{H} can be extended to a proper coloring of \mathcal{H}' by coloring each copy F_i of $V(\mathcal{H})$ in $V(\mathcal{H}')$ by the same colors assigned to $V(\mathcal{H})$. Doing so, all $e \in E(\mathcal{H}')$ will be properly colored, implying that all $e \in E(\mathcal{H}')$ must be properly colored by construction. Thus, we have shown that $k \geq \chi(H')$, and need only to confirm that $k \not\geq \chi(H')$. To show this, we demonstrate that any coloring of \mathcal{H}' with fewer than k colors must yield a monochromatic edge.

Consider a coloring $\varphi : \mathbb{R}^d \rightarrow \{1, \dots, k-1\}$. It is clear φ must not properly color \mathcal{H}_i for each $\mathcal{H}_i \in \mathcal{H}_1, \dots, \mathcal{H}_k$. Thus for each \mathcal{H}_i , there must be some edge $e = \{a_1, \dots, a_m\} \in E(\mathcal{H}_i)$ such that $\varphi(a_1) = \dots = \varphi(a_m)$.

We now show that there must be an edge $e \in \mathcal{H}'$ which φ monochromatically colors.

Let e_1 be a monochromatic edge in $E(\mathcal{H}_1)$. Without loss of generality suppose that $\varphi(e_1) = \{k-1\}$. If any vertex $v \in F_2 \cup \dots \cup F_k$ were colored by $k-1$, we would have found a monochromatic edge in \mathcal{H}' , by our construction of the edges in \mathcal{H}' . Thus, we can assume that $\varphi(v) \in \{1, \dots, k-2\}$ for $v \in F_2 \cup \dots \cup F_k$.

Likewise, there must be a monochromatic edge e_2 in $E(\mathcal{H}_2)$, because F_2 is colored now with even fewer colors than F_1 was. Because $\varphi(F_2) \in \{1, \dots, k-2\}$, we can without loss of generality suppose that $\varphi(e_2) = \{k-2\}$.

Continuing in this fashion until reaching F_k , we see that for each color $\alpha \in \{1, \dots, k-1\}$, there must appear a monochromatic edge in some \mathcal{H}_i for $i \in \{1, \dots, k-1\}$. Now consider $\varphi(v)$ for $v \in F_k$. No matter which color is chosen for v , there will appear a monochromatic edge in \mathcal{H}' of the form $e_i \cup v$ for some $e_i \in \mathcal{H}_i$. We see that it is impossible to properly color \mathcal{H}' with fewer than k colors, implying that $\chi(\mathcal{H}') = k$. \square

Corollary 5.1.1. *For all integers $m \geq 2$, there exists a finite, m -uniform hypergraph \mathcal{H}' with vertices in \mathbb{R}^d such that $\chi(\mathcal{H}') = \chi(\mathbb{R}^d, 1)$.*

Proof. We prove the statement by induction. When $m = 2$, the result follows directly from the D-E Theorem.

Now consider $m > 2$ and suppose the theorem holds true for $m-1$. Then there exists a finite, $(m-1)$ -uniform hypergraph \mathcal{H} with vertices in \mathbb{R}^d such that $\chi(\mathcal{H}) = \chi(\mathbb{R}^d, 1)$. By the previous theorem, there must exist an m -uniform hypergraph \mathcal{H}' with vertices in \mathbb{R}^d such that $\chi(\mathcal{H}') = \chi(\mathcal{H}) = \chi(\mathbb{R}^d, 1)$. \square

Remark 5.1.1. When $m = 2$, the hypergraph is not obtained constructively, but with this method, we still have control over the large-scale geometric structure of the m -uniform hypergraph based on how we position the finite hypergraphs when $m > 2$, and hence the term “construct”.

Appendix

Summary of the proof that for any norm $\|\cdot\|$ on \mathbb{R}^d , $\chi(\mathbb{R}^d, 1) < \infty$.

Let $\|\cdot\|$ be a norm on \mathbb{R}^d and let $\|\cdot\|_\infty$ be the norm on \mathbb{R}^d defined by $\|(a_1, \dots, a_d)\|_\infty = \max\{|a_i|; 1 \leq i \leq d\}$.

The proof of this well-known result will rely on the even better-known fact that any two norms on \mathbb{R}^d are equivalent. The equivalence of $\|\cdot\|$ and $\|\cdot\|_\infty$ means that there exist $c, C > 0$ such that for all $u \in \mathbb{R}^d$,

$$c\|u\|_\infty \leq \|u\| \leq C\|u\|_\infty.$$

The rightmost inequality can be used to prove the existence of $\varepsilon > 0$ such that the $\|\cdot\|$ -diameter of the d -dimensional cube $[0, \varepsilon]^d$ is < 1 . [Just take $0 < \varepsilon < \frac{1}{C}$.] Then the leftmost inequality above implies the existence of an integer $m > 0$ such that for each of the unit coordinate vectors $e_j = (S_{1j}, \dots, S_{dj})$ (in which S_{ij} is the Kronecker delta), $\|(m-1)e_j\| > 1$.

Next we partition the cube $Q = [0, m\varepsilon]^d$ into m^d little cubes, $\sum_{t=0}^{d-1} [j_t\varepsilon, (j_t+1)\varepsilon)$, $(j_0, \dots, j_{d-1}) \in \{0, 1, \dots, m-1\}^d$, and color Q with m^d colors; one to each little cube. Finally we color \mathbb{R}^d with those same colors by tiling \mathbb{R}^d with translates (with colors attached) of Q .

This will be a proper coloring of $(\mathbb{R}^d, \|\cdot\|, 1)$ because any two points bearing the same color are either within the same little cube—no two points of which are $\|\cdot\|$ -distance ≥ 1 apart,—or in two different little cubes, in which case the distance between them is > 1 .

Thus $\chi(\mathbb{R}^d, \|\cdot\|) \leq m^d < \infty$.

Acknowledgement

We would like to acknowledge that this paper was written with much advice from Dr. Peter Johnson, our mentor during the 2023 NSF-funded summer REU program at Auburn University. We extend our deepest gratitude for his generous guidance on this project and others throughout our time at Auburn.

References

- [1] N. G. De Bruijn and P. Erdős, A color problem for infinite graphs and a problem in the theory of relations, Indagationes Mathematicae 13 (1951), 369-373.
- [2] Soifer, A. (2008) *The Mathematical Coloring Book: Mathematics of Coloring and the Colorful Life of its Creators*. Springer New York. <https://doi.org/10.1007/978-0-387-74642-5>.
- [3] Munkres, J. (2000). Topology, Prentice Hall.