
SPECIAL FINSLER SPACES ADMITTING A SEMI-CONCURRENT VECTOR FIELD

A PREPRINT

M. R. Rajeshwari*, **S. K. Narasimhamurthy**** and **H. M. Manjunatha*****

Department of PG Studies and Research in Mathematics

Kuvempu University, Jnana Sahyadri

Shankaraghatta - 577 451, Shivamogga, Karnataka, India

*rajeshwarimr27@gmail.com

**nmurthysk@gmail.com

***manjunathahmmmt@gmail.com

ABSTRACT

The main objective of this paper is to study semi-concurrent vector fields on a Finsler manifold. We show that the quasi- C -reducible Finsler space, C^3 -like Finsler space, C^h -recurrent Finsler space, and P^2 -like Finsler space are equivalent to Riemannian if they admit a semi-concurrent vector field. Further, we prove the necessary and sufficient condition for a Finsler space satisfying C -conformal condition to become Riemannian.

Keywords Semi-concurrent vector fields · Special Finsler spaces · C -conformal condition.

Mathematics Subject Classification 2020 53B40 · 53Z05

1 Introduction

In the Finsler geometry, all geometrical quantities are direction-dependent. This feature makes Finsler geometry more suitable for studying physical theories like particle physics, relativistic optics, and general relativity. Finsler geometry has Riemannian geometry as a particular case. Recently, Finsler geometry was applied to study various astrophysical topics like wormhole models [5, 17, 20], compact star models [18], gravastar models [2, 21] and so on. The obtained results agree well with the accuracy of experimental observations. Nekouee et al. [14] have investigated the applications of the Finsler-Randers metric in cosmology. The Finslerian Schwarzschild-de sitter space-time is investigated in [6]. So, many researchers have been fascinated by this generalized geometry and studying the geometry of a Finsler manifold.

Brickell and Yano [3] studied the concept of concurrent vector fields on a Riemannian manifold. In 1950, Tachibana [22] defined and discussed the concurrent vector fields. Later, in 1974, Matsumoto and Eguchi [9] extended the study of concurrent vector fields. In 2004, Rastogi and Dwivedi [19] discussed the concurrent vector fields on a Finsler manifold.

Semi-concurrent vector fields are studied on a Finsler manifold in [24]. And we found from [24] that some special Finsler spaces admitting semi-concurrent vector fields reduce to Riemannian. Further, they [24] discussed some properties of concurrent vector fields. Youssef *et al.* [23] studied the concurrent π -vector fields on some special Finsler spaces. And found that the special Finsler spaces admitting a concurrent π -vector field reduce to a Riemannian.

In this paper, we study some results on the Finsler spaces satisfying the C -conformal condition. Also, we discuss some special Finsler spaces admitting a semi-concurrent vector field.

2 Notations and preliminaries

Let (M, F) be an n -dimensional smooth connected Finsler manifold; F being the Finsler function. Let (x^i) be the coordinates of any point of the base manifold M and (y^i) a supporting element at the same point. We use the following terminology and notations:

∂_i : partial differentiation with respect to x^i ,

$\dot{\partial}_i$: partial differentiation with respect to y^i (basis vector fields of the vertical bundle),

$g_{ij} := \frac{1}{2}\dot{\partial}_i\dot{\partial}_jF^2 = \dot{\partial}_i\dot{\partial}_jE$: the Finsler metric tensor, where $E = \frac{1}{2}F^2$ is the energy function,

$l_i := \dot{\partial}_iF = g_{ij}l^j = g_{ij}\frac{y^j}{F}$: the normalized supporting element; $l^i := \frac{y^i}{F}$,

$l_{ij} := \dot{\partial}_il_j$,

$h_{ij} := Fl_{ij} = g_{ij} - l_il_j$: the angular metric tensor,

$C_{ijk} := \frac{1}{2}\dot{\partial}_k g_{ij} = \frac{1}{4}\dot{\partial}_i\dot{\partial}_j\dot{\partial}_kF^2$: the Cartan tensor,

$C_{jk}^i := g^{ri}C_{rjk} = \frac{1}{2}g^{ri}\dot{\partial}_k g_{rj}$: the $h(hv)$ -torsion tensor,

G^i : the components of the geodesic spray associated with (M, F) ,

$N_j^i := \dot{\partial}_jG^i$: the Barthel (or Cartan nonlinear) connection associated with (M, F) ,

$\delta_i := \partial_i - N_i^r\dot{\partial}_r$: the basis vector fields of the horizontal bundle,

$G_{jh}^i := \dot{\partial}_hN_j^i = \dot{\partial}_h\dot{\partial}_jG^i$: the coefficients of Berwald connection,

$\Gamma_{jk}^i := \frac{1}{2}g^{ir}(\delta_jg_{kr} + \delta_kg_{jr} - \delta_rg_{jk})$: the Christoffel symbols with respect to δ_i ,

$(\Gamma_{jk}^i, N_j^i, C_{jk}^i)$: the Cartan connection.

For the Cartan connection $(\Gamma_{jk}^i, N_j^i, C_{jk}^i)$, we define

$X_{j|k}^i := \delta_kX_j^i + X_j^m\Gamma_{mk}^i - X_m^i\Gamma_{jk}^m$: the horizontal covariant derivative of X_j^i ,

$X_j^i|k := \dot{\partial}_kX_j^i + X_j^mC_{mk}^i - X_m^iC_{jk}^m$: the vertical covariant derivative of X_j^i .

Definition 1 [1] A Finsler structure on a manifold M is a function

$$F : TM \longrightarrow [0, \infty)$$

with the following properties:

1. F is C^∞ on the slit tangent bundle $TM := TM \setminus 0$.
2. $F(x, y)$ is positively homogenous of degree one in y : $F(x, \lambda y) = \lambda F(x, y)$ for all $y \in TM$ and $\lambda > 0$.
3. The Hessian matrix $g_{ij}(x, y) := \dot{\partial}_i\dot{\partial}_jE$ is positive-definite at each point of TM , where $E := \frac{1}{2}F^2$ is the energy function of the Lagrangian F .

The pair (M, F) is called a Finsler manifold and the symmetric bilinear form $g = g_{ij}(x, y)dx^i \otimes dx^j$ is called the Finsler metric tensor of the Finsler manifold (M, F) .

Sometimes, a function F satisfying the above conditions is said to be a regular Finsler metric.

Next, we will provide the definitions of some special Finsler spaces.

Definition 2 [7] A Finsler manifold (M, F) of dimension $n \geq 3$ is called a C -reducible if the Cartan tensor C_{ijk} has the following form:

$$C_{ijk} = \frac{1}{n+1}(h_{ij}C_k + h_{ki}C_j + h_{jk}C_i). \quad (1)$$

Definition 3 [10] A Finsler manifold (M, F) of dimension $n \geq 3$ is called a semi- C -reducible if the Cartan tensor C_{ijk} is written in the following form:

$$C_{ijk} = \frac{r}{n+1}(h_{ij}C_k + h_{ki}C_j + h_{jk}C_i) + \frac{t}{C^2}C_iC_jC_k, \quad (2)$$

where r and t are scalar functions such that $r + t = 1$.

Definition 4 [13] A Finsler manifold (M, F) of dimension $n \geq 3$ is called a quasi- C -reducible if the Cartan tensor C_{ijk} has the following form:

$$C_{ijk} = h_{ij}C_k + h_{ki}C_j + h_{jk}C_i. \quad (3)$$

In a three-dimensional Finsler space, F^3 , C_{ijk} is always written in the form

$$LC_{ijk} = Hm_i m_j m_k - JC_{ijk}(m_i m_j m_k) + IC_{ijk}(m_i m_j m_k) + Jn_i n_j n_k \quad (4)$$

where $L = L(x, y)$ is the fundamental function, $\mathcal{C}_{(ijk)}$ denote the cyclic permutation of indices i, j, k and addition. H, I and J are main scalars and (I_i, m_i, n_i) is Moor's frame [7, 13, 11]. Here $I_i = \dot{\partial}_i L$ is the unit vector along the element of support, m_i is the unit vector along C_i , that is, $m_i = C_i/C$, where $C^2 = g^{ij}C_i C_j$ and n_i is a unit vector orthogonal to the vectors I_i and m_i . Since the angular metric tensor h_{ij} in F^3 can be written as

$$h_{ij} = m_i m_j + n_i n_j, \quad (5)$$

we may write (4) as

$$C_{ijk} = \mathfrak{C}_{(ijk)} \{h_{ij} a_k + C_i C_j b_k\}. \quad (6)$$

where $a_k = \frac{1}{L}(Im_k + \frac{J}{3}n_k)$ and $b_k = \frac{1}{LC^2} \{(\frac{H}{3} - I)m_k + \frac{4J}{3}n_k\}$. Many authors [9, 7, 10, 15] have obtained various interesting special forms of C_{ijk} .

Definition 5 [16] A Finsler space F^n of dimension $n \geq 4$ is called a C3-like Finsler space if there exist covariant vector fields a_k and b_k in F^n such that its (h) hv-torsion tensor C_{ijk} can be written as

$$C_{ijk} = \mathfrak{C}_{(ijk)} \{h_{ij} a_k + C_i C_j b_k\}. \quad (7)$$

Since C_{ijk} is an indicatory tensor, it follows that a_k and b_k are indicatory tensors, that is, $a_0 = b_0 = 0$. If b_k is a null vector then contracting (7) with g^{ik} and putting $b_k = 0$, we get $a_k = \frac{1}{n+1}C_k$.

Definition 6 [1] A tangent vector field X of a Finsler space F^n is concurrent under the Cartan connection, if

$$X_{|j}^i = \partial_j X^i - N_j^h \dot{\partial}_h X^i + X^h F_{hj}^i = \partial_j X^i + X^h F_{hj}^i = -\delta_j^i \quad (8)$$

$$X_{|j}^i = \dot{\partial}_j X^i + X^h C_{hj}^i = X^h C_{hj}^i = 0. \quad (9)$$

where ∂_j and $\dot{\partial}_j$ denote partial differentiations by X^j and Y^j respectively.

The Ricci identities [8] in a Finsler space are as follows:

$$X^h R_{hijk} = 0, \quad (10)$$

$$X^h P_{hijk} + C_{ijk} = 0, \quad (11)$$

$$X^h S_{hijk} = 0, \quad (12)$$

where R_{hijk} , P_{hijk} and S_{hijk} are the components of the curvature tensors of CT. Since P_{hijk} are skew symmetric in h and i , we have from (9) that

$$X^h C_{ijk} = 0. \quad (13)$$

We know that the well-known identity,

$$P_{hijk} = C_{ijk|h} - C_{hjk|i} + C_{hjr} C_{ik|0}^r - C_{ijr} C_{hk|0}^r. \quad (14)$$

Substitute (14) in (9), we get

$$X^h (P_{hijk} - C_{ijk|h} + C_{hjk|i} - C_{hjr} C_{ik|0}^r + C_{ijr} C_{hk|0}^r) C_{hj}^i = 0. \quad (15)$$

We know that,

$$C_{ijk|h} = C_{jk|0}^r - C_{ih|0}^r + C_{kir} C_{jh|0}^r - C_{ijr} C_{hk|0}^r - P_{ijkh}. \quad (16)$$

Contracting the above equation, we get

$$X^h C_{ijk|h} = 0. \quad (17)$$

Definition 7 [4] A Finsler space F^n is called a C^h -recurrent if the torsion tensor satisfies the following equation:

$$C_{ijk|h} = C_{ijk} K_h, \quad (18)$$

where K_h is a covariant vector field.

Definition 8 [4] For a P2-like Finsler space F^n , we have

$$P_{hijk} = K_h C_{ijk} - K_i C_{kjh}, \quad (19)$$

where K_h is a covariant vector field.

Definition 9 [4] A Finsler space F^n is called a P-reducible if the torsion tensor P_{ijk} is defined as follows:

$$P_{ijk} = \frac{1}{n+1} (h_{ij} P_k + h_{jk} P_i + h_{ki} P_j), \quad (20)$$

where $P_i = P_{ir}^r = C_i$.

Definition 10 [4] A Finsler space F^n is said to satisfy T-condition if the T-tensor T_{hijk} vanishes identically, that is,

$$T_{hijk} = LC_{hij|k} + l_h C_{ijk} + l_i C_{hjk} + l_j C_{hik} + l_k C_{hij} = 0. \quad (21)$$

3 Semi-concurrent vector fields

Let (M, F) be an n -dimensional smooth Finsler manifold.

Definition 11 [22] A vector field $X^i(x)$ on M is said to be concurrent under the Cartan connection if it satisfies the following expression:

$$X^h(x)C_{hij} = 0, \quad X^i_{|j} = -\delta_j^i. \quad (22)$$

The condition (22) is called the C -condition.

Definition 12 [7] The manifold M fulfills the C -conformal condition if there exists a conformal transformation $\bar{F} = e^{\sigma(x)}F$ on M such that

$$\sigma_h(x)C_{ij}^h = 0, \quad (23)$$

where $\sigma_h := \frac{\partial \sigma}{\partial x^h}$. The condition (23) will be called the CC -condition.

Definition 13 [24] A vector field $B^i(x)$ on M is said to be semi-concurrent if it satisfies the following expression:

$$B^h(x)C_{hij} = 0. \quad (24)$$

The condition (24) will be called the SC -condition.

Lemma 1 [24] For the nonzero function B^i satisfying (24), if the scalars α and α' satisfy

$$\alpha B^i + \alpha' y^i = 0, \quad (25)$$

then $\alpha = \alpha' = 0$, which means that the two vector fields $B^i(x)$ and y^i are independent.

Theorem 1 A quasi- C -reducible Finsler space (M, F) admitting a semi-concurrent vector field is Riemannian.

Proof 1 For a quasi- C -reducible Finsler space, we have

$$C_{ijk} = h_{ij}C_k + h_{jk}C_i + h_{ki}C_j. \quad (26)$$

Contracting (26) by $B^i B^j$ and by using (24), we get

$$B^i B^j h_{ij} C_k = 0. \quad (27)$$

This implies that $C_k = 0$. Hence the space is Riemannian.

Theorem 2 A C_3 -like Finsler space satisfying the SC -condition is Riemannian provided $J = 0$.

Proof 2 For the C_3 -like Finsler space, we have

$$C_{ijk} = h_{ij}a_k + C_i C_j b_k + h_{jk}a_i + C_j C_k b_i + h_{ki}a_j + C_k C_i b_j. \quad (28)$$

Contracting (28) by $B^i B^j$ and from (24), we get

$$B^i B^j (h_{ij}a_k + h_{jk}a_i + h_{ki}a_j) = 0. \quad (29)$$

Equation (29) implies that

$$B^i B^j n_i n_j a_k + B^j B^k n_j n_k a_i + B^k B^i n_k n_i a_j = 0, \quad (30)$$

which yields

$$\frac{B^2 I}{LC} C_k + \frac{B^2 J}{L} n_k = 0. \quad (31)$$

In view of (31), we obtain that $C_k = 0$ if $J = 0$.

Theorem 3 A C^h -recurrent Finsler space admitting a semi-concurrent vector field is Riemannian if $1 + B^h K_h \neq 0$.

Proof 3 For the C^h -recurrent Finsler space, we have

$$C_{ijk|h} = C_{ijk} K_h, \quad (32)$$

where K_h is a covariant vector field. Consider

$$P_{hijk} = C_{ijk|h} - C_{hjk|i} + C_{hjr} C_{ik|0}^r - C_{ijr} C_{hk|0}^r. \quad (33)$$

Contracting (33) by B^h yields

$$B^h P_{hijk} = B^h C_{ijk|h}. \quad (34)$$

From Ricci identity, (34) implies that

$$B^h C_{ijk|h} = -C_{ijk}. \quad (35)$$

On contracting (32) by B^h and by using (35), we conclude that $C_{ijk} = 0$ and hence the space is Riemannian if $1 + B^h K_h \neq 0$.

Theorem 4 A P2-like Finsler space admitting a semi-concurrent vector field is Riemannian provided $1 + B^h K_h \neq 0$.

Proof 4 A P2-like Finsler space is characterized by

$$P_{hijk} = K_h C_{ijk} - K_i C_{kjh}, \quad (36)$$

where K^h is a covariant vector field. Consider

$$P_{hijk} = C_{ijk|h} - C_{hjk|i} + C_{hjr} C_{ik|0}^r - C_{ijr} C_{hk|0}^r. \quad (37)$$

Contracting (37) by B^h , we obtain

$$B^h P_{hijk} = B^h C_{ijk|h}. \quad (38)$$

From Ricci identity, (38) yields

$$B^h C_{ijk|h} = -C_{ijk}, \quad (39)$$

On contracting (36) by B^h and by using (39), we conclude that $C_{ijk} = 0$ if $1 + B^h K_h \neq 0$, which implies that the space is Riemannian.

Theorem 5 A P-reducible Finsler space F^n admitting a semi-concurrent vector field is a Landsberg space provided $B^2 F^2 - B_0^2 \neq 0$.

Proof 5 The torsion tensor P_{ijk} is given by

$$P_{ijk} = h_{ij} P_k + h_{jk} P_i + h_{ki} P_j. \quad (40)$$

Contracting (40) by $B^i B^j$ and since $P_{ijk} = C_{ijk}$, we obtain

$$B^i B^j h_{ij} C_k = 0, \quad (41)$$

which implies that

$$(B^2 F^2 - B_0^2) C_k = 0. \quad (42)$$

From (42), we deduce that $C_k = 0$, that is, $P_k = 0$ if $B^2 F^2 - B_0^2 \neq 0$, which implies that F^n is a Landsberg space.

The T -tensor is defined as follows [12]:

$$T_{hijk} = F C_{hij|k} + C_{hij} l_k + C_{hik} l_j + C_{hjk} l_i + C_{ijk} l_h \quad (43)$$

If (M, F) is Riemannian, then the T -tensor vanishes. If (M, F) satisfies the CC -condition, then the converse part of the result holds, which is shown in the following theorem.

Theorem 6 A Finsler manifold satisfying the CC -condition is Riemannian if and only if the T -tensor T_{hijk} vanishes.

Proof 6 First, we prove that the vertical covariant derivative of σ_h vanishes identically. Since $\sigma_h = \sigma_h(x)$, we have

$$\sigma_h|_k = \dot{\partial}_k \sigma_h + \sigma_i C_{hk}^i = 0.$$

Let the T -tensor vanish, then from (43), we have

$$F C_{hij|k} + C_{hij} l_k + C_{hik} l_j + C_{hjk} l_i + C_{ijk} l_h = 0.$$

Contracting by σ_m , and taking into account that $\sigma_m|_k = 0$, we get

$$\sigma_m C_{hij} l_k + \sigma_m C_{hik} l_j + \sigma_m C_{hjk} l_i + \sigma_m C_{ijk} l_h = 0.$$

Again contracting by g^{mh} , we obtain

$$\frac{\sigma_0}{F} C_{ijk} = 0.$$

Since $\sigma_0 \neq 0$, it follows that $C_{ijk} = 0$.

By defining the tensor T_{ij} ,

$$T_{ij} := T_{ijhk} g^{hk} = F C_i|_j + l_i C_j + l_j C_i,$$

we can state the following result.

Corollary 1 A Finsler manifold satisfying the CC -condition is Riemannian if and only if the tensor T_{ij} vanishes.

4 Conclusions

In Finsler geometry, special Finsler spaces play a significant role. In this context, we have studied semi-concurrent vector fields on some special Finsler spaces and obtained interesting results. We have shown that a quasi- C -reducible Finsler space, C^3 -like Finsler space, C^h -recurrent Finsler space, and P^2 -like Finsler space admitting a semi-concurrent vector field become Riemannian. And a P -reducible Finsler space satisfying the SC -condition becomes a Landsberg space. Further, we have obtained the necessary and sufficient condition for a Finsler space satisfying the C -conformal condition to become Riemannian.

References

- [1] P. L. Antonelli, R. S. Ingarden and M. Matsumoto, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology* (Springer, Dordrecht, 1993).
- [2] S. Banerjee, S. Ghosh, N. Paul and F. Rahaman, Study of gravastars in Finslerian geometry, *Eur. Phys. J. Plus* **135** (2020) 185.
- [3] F. Brickell and K. Yano, Concurrent vector fields and Minkowski structures, *Kōdai Math. Sem. Rep.* **26** (1974) 22–28.
- [4] M. Hashiguchi, On conformal transformations of Finsler metrics, *J. Math. Kyoto Univ.* **16** (1976) 25–50.
- [5] H. M. Manjunatha and S. K. Narasimhamurthy, The wormhole model with an exponential shape function in the Finslerian framework, *Chin. J. Phys.* **77** (2022) 1561–1578.
- [6] H. M. Manjunatha, S. K. Narasimhamurthy, S. K. Srivastava, Finslerian analogue of the Schwarzschild-de Sitter space-time, *Pramana-J. Phys.* **97** (2023) 90.
- [7] M. Matsumoto, On C -reducible Finsler spaces, *Tensor N.S.* **24** (1972) 29–37.
- [8] M. Matsumoto, A theory of three-dimensional Finsler spaces in terms of scalars, *Demonst. Math. Warszawa* **6** (1973) 223–251.
- [9] M. Matsumoto and K. Eguchi, Finsler spaces admitting a concurrent vector field, *Tensor N.S.* **28** (1974) 239–249.
- [10] M. Matsumoto and C. Shibata, On semi- C -reducibility, T-tensor= 0 and S_4 -likeness of Finsler spaces, *J. Math. Kyoto Univ.* **19** (1979) 301–314.
- [11] M. Matsumoto and S. Numata, On semi- C -reducible Finsler spaces with constant coefficients and C_2 -like Finsler spaces, *Tensor N.S.* **34** (1980) 218–222.
- [12] M. Matsumoto, *Foundations of Finsler Geometry and Special Finsler spaces* (Kaiseisha Press, Japan, 1986).
- [13] S. K. Narasimhamurthy, B. C. Chethana and D. M. Vasantha, Ricci flow equations on quasi- C -reducible Finsler space, *Int. J. Mathematics Trends and Technology* **29**(1) (2016) 28–32.
- [14] Z. Nekouee, S. K. Narasimhamurthy, H. M. Manjunatha and S. K. Srivastava, Finsler-Randers model for anisotropic constant-roll inflation, *Eur. Phys. J. Plus* **137** (2022) 1388.
- [15] T. Okada and S. Numata, On generalized C -reducible Finsler spaces, *Tensor N.S.* **35**(3) (1981) 313–318.
- [16] B. N. Prasad and J. N. Singh, On C^3 -like Finsler spaces, *Indian J. Pure Appl. Math.* **19** (1988) 423–428.
- [17] F. Rahaman, N. Paul, A. Banerjee, S. S. De, S. Ray and A. A. Usmani, The Finslerian wormhole models, *Eur. Phys. J. C* **76** (2016) 246.
- [18] F. Rahaman, N. Paul, S. S. De, S. Ray and Md. A. K. Jafry, The Finslerian compact star model, *Eur. Phys. J. C* **75** (2015) 564.
- [19] S. C. Rastogi and A. K. Dwivedi, On the existence of concurrent vector fields in a Finsler space, *Tensor N.S.* **65**(1) (2004) 48–54.
- [20] T. Sanjay, S. K. Narasimhamurthy, Z. Nekouee and H. M. Manjunatha, Traversable wormhole models in Finslerian geometry, *Pramana-J. Phys.* **98** (2024) 16.
- [21] T. Sanjay, S. K. Narasimhamurthy, Z. Nekouee and H. M. Manjunatha, Charged gravastars with conformal motion in the Finslerian space-time, *Eur. Phys. J. C* **84** (2024) 393.
- [22] S. Tachibana, On Finsler spaces which admit concurrent vector field, *Tensor N.S.* **1** (1950) 1–5.
- [23] N. L. Youssef, S. H. Abed and A. Soleiman, Concurrent π -vector fields and energy β -change, *Int. J. Geom. Methods Mod. Phys.* **6**(6) (2009) 1003–1031.
- [24] N. L. Youssef, S. G. Elgendi and E. H. Taha, Semi-concurrent vector fields in Finsler geometry, *Differ. Geom. Appl.* **65** (2019) 1–15.