

Nonlinear Hall effects with an exceptional ring

Fang Qin,^{1,*} Ruizhe Shen,^{2,†} and Ching Hua Lee^{2,‡}

¹*School of Science, Jiangsu University of Science and Technology, Zhenjiang 212100, China*

²*Department of Physics, National University of Singapore, Singapore 117551, Singapore*

(Dated: November 12, 2024)

In non-Hermitian band structures, exceptional points generically form gapless lines or loops that give rise to extensively many defective eigenstates. In this work, we investigate how they non-trivially contribute to higher-order nonlinear responses by introducing unique singularities in the Berry curvature dipole (BCD) or Berry connection polarizability (BCP). Using a tilted 2D dissipative Dirac model ansatz that harbors an exceptional ring, broken inversion symmetry is shown to give rise to extrinsic (BCD) and intrinsic (BCP) nonlinear Hall behaviors unique to systems with extensive exceptional singularities. In particular, when the non-Hermiticity is increased while keeping the ring radius fixed, the BCD response exhibits a power-law increase, while the BCP response correspondingly decreases. Our work sheds light on how non-Hermiticity can qualitatively control the extent and nature of higher harmonic generation in solids.

I. INTRODUCTION

Beyond the well-known Kubo formula paradigm that connects band topology to linear response properties [1–11], the topology and geometry of band crossings also intimately dictates the non-linear response, particularly in terms of technologically relevant higher harmonic generation properties [12–21].

A key nonlinear response phenomenon is the extrinsic nonlinear Hall effect, which arises due to broken inversion symmetry [22–27]. This unique response is proportional to the Berry curvature dipole (BCD), and has garnered significant experimental interest [22–27], being widely explored in a variety of materials including two-dimensional (2D) bilayer WTe₂ [28–30], 2D monolayer MoS₂ [31, 32], the three-dimensional (3D) Dirac semimetal Cd₃As₂ [33], Weyl semimetal materials [34–36], the topological insulator Bi₂Se₃ [37], the giant Rashba semiconductor BiTeI [38], bilayer graphene [39], twisted bilayer graphene [40, 41], twisted double bilayer graphene [42], strained twisted bilayer graphene [43, 44], and gapped bilayer graphene [45]. Recently, there has been also been a surge of interest in the intrinsic nonlinear Hall effect, which alternatively arise from the quantum geometric tensor [46–56] and has been experimentally probed in thick T_d-MoTe₂ samples [50] and the topological antiferromagnet material MnBi₂Te₄ [52–54].

At the same time, non-Hermitian physics has given rise to a new landscape for topological physics [57–90]. The infusion of non-Hermitian topology has greatly enriched the phenomenology of quantum response and transport. Previous studies on non-Hermitian linear response [91, 92] have unveiled anomalous coherent oscillations in wave packet dynamics [9, 93–100] induced by the non-Hermitian anomalous Berry connection. In particular, in non-Hermitian systems containing exceptional

point singularities, the Berry curvature exhibits singularities near the band crossings, leading to much heightened sensitivity around these singularities and qualitatively affecting both linear and high-order nonlinear responses. As such, in this work, we investigate a tilted two-dimensional massive dissipative Dirac model that represents an ansatz exceptional ring model. Our findings reveal that both the BCD and Berry connection polarizability (BCP) become markedly sensitive to non-Hermiticity near the singularities of an exceptional ring. Particularly, the distinct scaling laws of BCD and BCP in proximity to this exceptional ring lead to significantly different extrinsic and intrinsic nonlinear Hall responses.

The paper is organized as follows: In section II, we present the general formalism for both extrinsic and intrinsic nonlinear Hall effects. In section III, we introduce the Hamiltonian for an exceptional ring and provide a detailed analysis of its energy band structure. In section IV, we investigate the quantum geometry tensor, Berry curvature, quantum metric, and the extrinsic nonlinear Hall response described by the BCD. Moreover, we study the nonzero component intrinsic nonlinear Hall response described by the BCP.

II. GENERAL FORMALISM FOR NONLINEAR HALL EFFECTS

In this section, we introduce the foundational framework for understanding both extrinsic and intrinsic nonlinear Hall effects, with a focus on how such quantum geometric properties are modified in non-Hermitian systems.

A. Quantum geometry tensor

The quantum geometry tensor, which combines information about the Berry curvature and quantum metric, plays a crucial role in the topological characterization of

* qinfang@just.edu.cn

† ruizhe20@u.nus.edu

‡ phylch@nus.edu.sg

responses. Here, we introduce the concept of the quantum geometry tensor in non-Hermitian systems. Unlike the conventional Hermitian setup, the biorthogonal quantum geometry tensor in a non-Hermitian model is defined using the left and right eigenvectors $|\psi_R^{(n)}\rangle$ and $|\psi_L^{(n)}\rangle$, of the n -th energy band [60–90, 94, 101–104]. This tensor can be defined as a matrix in the r -dimensional subspace spanned by r selected occupied bands [48, 105–112]:

$$\begin{aligned}\mathcal{Q}_{ab}^{(nm)} &= \langle \partial_{k_a} \psi_L^{(n)} | (\mathbb{I} - \hat{P}) | \partial_{k_b} \psi_R^{(m)} \rangle \\ &= \mathcal{G}_{ab}^{(nm)} - \frac{i}{2} \Omega_{ab}^{(nm)},\end{aligned}\quad (1)$$

where $a, b \in \{x, y, z\}$ are for spatial coordinates, \mathbb{I} is the identity matrix, $\hat{P} = \sum_{n'=1}^r |\psi_R^{(n')}\rangle \langle \psi_L^{(n')}|$ is the biorthogonal projection operator [113] so that \mathcal{Q}_{ab} is a $r \times r$ matrix. Here, $\mathcal{G}_{ab}^{(nm)}$ and $\Omega_{ab}^{(nm)}$ are the biorthogonal quantum metric and Berry curvature respectively, which are given by [107, 108, 113–119]

$$\begin{aligned}\mathcal{G}_{ab}^{(nm)} &\equiv \frac{1}{2} \text{Re} \left[\mathcal{Q}_{ab}^{(nm)} + \mathcal{Q}_{ba}^{(mn)} \right] \\ &= \frac{1}{2} \text{Re} \left[\langle \partial_{k_a} \psi_L^{(n)} | \partial_{k_b} \psi_R^{(m)} \rangle + \langle \partial_{k_b} \psi_L^{(m)} | \partial_{k_a} \psi_R^{(n)} \rangle \right] \\ &\quad - \frac{1}{2} \sum_{n'} \left[\langle \partial_{k_a} \psi_L^{(n)} | \psi_R^{(n')} \rangle \langle \psi_L^{(n')} | \partial_{k_b} \psi_R^{(m)} \rangle \right. \\ &\quad \left. + \langle \partial_{k_b} \psi_L^{(m)} | \psi_R^{(n')} \rangle \langle \psi_L^{(n')} | \partial_{k_a} \psi_R^{(n)} \rangle \right],\end{aligned}\quad (2)$$

$$\begin{aligned}\Omega_{ab}^{(nm)} &\equiv \text{Re} \left\{ i \left[\mathcal{Q}_{ab}^{(nm)} - \mathcal{Q}_{ba}^{(mn)} \right] \right\} \\ &= \text{Re} \left\{ i \left[\langle \partial_{k_a} \psi_L^{(n)} | \partial_{k_b} \psi_R^{(m)} \rangle - \langle \partial_{k_b} \psi_L^{(m)} | \partial_{k_a} \psi_R^{(n)} \rangle \right] \right\},\end{aligned}\quad (3)$$

where we have the relations $\Omega_{ba}^{(nm)} = -\Omega_{ab}^{(nm)} = -\Omega_c^{(nm)}$.

B. Extrinsic nonlinear Hall effect

We now examine the extrinsic nonlinear Hall effect, where the direct response to an external force manifests in the velocity response of affected particles. Here, we consider a simple scenario involving a time-driven oscillating force

$$\mathbf{F}(t) = F_a^\omega(t) \mathbf{e}_a = \text{Re}(\mathcal{F}_a e^{i\omega t}) \mathbf{e}_a \quad (4)$$

along the a direction. $\mathbf{F}(t)$ has the unit of force, and it enters the non-equilibrium distribution function f through the Boltzmann equation within the relaxation time approximation, which is a semiclassical approximation. This force can induce an additional transverse velocity along the b direction in the particle clouds [10, 22–24, 120]:

$$\bar{v}_b(\mathcal{F}_a) \approx \text{Re}(\bar{v}_b^0 + \bar{v}_b^\omega e^{i\omega t} + \bar{v}_b^{2\omega} e^{i2\omega t}). \quad (5)$$

We first discuss the linear contribution $\bar{v}_b^\omega = \chi_{ba}^{(1)} \mathcal{F}_a$, where the coefficient $\chi_{ba}^{(1)} \approx \frac{\mathcal{S}}{2\pi\hbar} C_{ba} = \frac{\mathcal{S}}{2\pi\hbar} \varepsilon_{bac} C^c$

with [10]

$$C^c = \varepsilon^{bac} C_{ba} = -C_{ba} = -2\pi \sum_n \int \frac{d^2\mathbf{k}}{(2\pi)^2} \Omega_c^{(n)} f_0^{(n)}. \quad (6)$$

Here \mathcal{F}_a is the magnitude of force along the a direction, \mathcal{S} is the area of the system, \hbar is the reduced Planck's constant, ε^{bac} is the Levi-Civita anti-symmetric tensor, and a, b , and c represent the spatial coordinates x, y , and z . In Eq. (6), $\Omega_c^{(n)}$ is the c -th component of the Berry curvature $\boldsymbol{\Omega}^{(n)}$ computed for the n -th eigenstate, and $f_0^{(n)} = f_0^{(n)}(\epsilon_{\mathbf{k}}^{(n)} - E_F)$ is the equilibrium Fermi-Dirac distribution function of the n -th eigenenergy band $\epsilon_{\mathbf{k}}^{(n)}$ and Fermi energy E_F . Specifically, under the oscillating force $\mathbf{F} = F_a^\omega(t) \mathbf{e}_a$ with $x = a$ [Eq. (4)] [121–124], we obtain the linear velocity contribution in the y -direction

$$\bar{v}_y^\omega \approx \frac{\mathcal{S}}{2\pi\hbar} \mathcal{F}_x C^z. \quad (7)$$

Next, we focus on the nonlinear response, which arises from the following two contributions: $\bar{v}_b^0 = \chi_{baa}^{(0)} \mathcal{F}_a^2$ and $\bar{v}_b^{2\omega} = \chi_{baa}^{(2)} \mathcal{F}_a^2$. The coefficients are given by $\chi_{baa}^{(0)} \approx \chi_{baa}^{(2)} \approx \frac{\tau \mathcal{S} D_{baa}}{2\hbar^2(1+i\omega\tau)}$ [22–24], where τ is the relaxation time [22–24]. At the core of the extrinsic nonlinear response is the Berry curvature dipole (BCD) D_{baa} , as given by [22–24, 125]:

$$D_{baa} = \sum_n \int \frac{d^2\mathbf{k}}{(2\pi)^2} \Omega_{ba}^{(n)} (\partial_{k_a} \epsilon_{\mathbf{k}}^{(n)}) \frac{\partial f_0^{(n)}}{\partial \epsilon_{\mathbf{k}}^{(n)}}. \quad (8)$$

The derivative of the non-equilibrium distribution function $\partial f_0^{(n)} / \partial \epsilon_{\mathbf{k}}^{(n)}$ in the above integral for the BCD indicates that the extrinsic nonlinear Hall response is predominantly governed by the states near the Fermi surface [23–25, 125–127]. Setting $a = x$ in Eq. (4), the extrinsic nonlinear response manifesting as the y -direction nonlinear velocity can be obtained by solving the Boltzmann equation under the relaxation time approximation [22–24]

$$\bar{v}_{yxx}^{\text{BCD}} \approx \bar{v}_y^0 \approx \bar{v}_y^{2\omega} \approx \frac{\tau \mathcal{S} \mathcal{F}_x^2}{2\hbar^2} D_{yxx}, \quad (9)$$

where $\bar{v}_{yxx}^{\text{BCD}}$ is the extrinsic nonlinear velocity, and we have used $\omega\tau \rightarrow 0$. Furthermore, with the help of Eq. (8), we obtain

$$\begin{aligned}D_{yxx} &= \sum_n \int \frac{d^2\mathbf{k}}{(2\pi)^2} \left[\Omega_{yx}^{(n)} (\partial_{k_x} \epsilon_{\mathbf{k}}^{(n)}) \frac{\partial f_0^{(n)}}{\partial \epsilon_{\mathbf{k}}^{(n)}} \right] \\ &= -\sum_n \int \frac{d^2\mathbf{k}}{(2\pi)^2} \left[\Omega_z^{(n)} (\partial_{k_x} \epsilon_{\mathbf{k}}^{(n)}) \frac{\partial f_0^{(n)}}{\partial \epsilon_{\mathbf{k}}^{(n)}} \right] \\ &= -\sum_n \int \frac{dk_y}{2\pi} \left[f_0^{(n)} \Omega_z^{(n)} \right] \Big|_{k_x=-\pi}^{k_x=\pi} \\ &\quad + \sum_n \int \frac{d^2\mathbf{k}}{(2\pi)^2} \left[f_0^{(n)} \frac{\partial \Omega_z^{(n)}}{\partial k_x} \right],\end{aligned}\quad (10)$$

where we used $\Omega_{yx}^{(n)} = -\Omega_{xy}^{(n)} = -\Omega_z^{(n)}$. Note that the inversion-symmetry breaking is essential for the finite Berry curvature and non-vanishing BCD [23–25, 125–127].

C. Intrinsic nonlinear Hall effect

The second-order velocity tensor (nonlinear velocity tensor) is defined as the contribution to the velocity response that is quadratic to the applied force field \mathbf{F} as $\bar{v}_{abc} \propto \mathcal{F}_b \mathcal{F}_c$ [46], where \bar{v}_{abc} can be separated into an Ohmic-type nonlinear response and a Hall-type nonlinear response [128]. The Ohmic-type nonlinear response includes a second-order Drude conductivity, which is quadratically dependent on the relaxation time [129–132]. The Hall-type nonlinear response is independent of the relaxation time and is therefore called the intrinsic nonlinear Hall effect.

This intrinsic nonlinear response can be described by the Berry connection polarizability (BCP) [46, 47, 49–51], as given by the intrinsic nonlinear velocity $\bar{v}_{abc}^{\text{BCP}}$ [46, 47, 49, 51, 55, 56, 125]

$$\begin{aligned} \bar{v}_{abc}^{\text{BCP}} &= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \left[\frac{v_a^{(n)} \mathcal{G}_{bc}^{(n)} - v_b^{(n)} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \frac{\partial f_0^{(n)}}{\partial \epsilon_{\mathbf{k}}^{(n)}} \\ &= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_b}{2\pi} \left[\frac{f_0^{(n)} \mathcal{G}_{bc}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_a=-\pi}^{k_a=\pi} \\ &\quad + \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_a}{2\pi} \left[\frac{-f_0^{(n)} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_b=-\pi}^{k_b=\pi} \\ &\quad - \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} f_0^{(n)} \left[\frac{\partial_{k_a} \mathcal{G}_{bc}^{(n)} - \partial_{k_b} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right], \end{aligned} \quad (11)$$

where $\Gamma_{bc} = 2\mathcal{S}\mathcal{F}_b\mathcal{F}_c/\hbar$ with $\mathcal{F}_{b/c}$ the force magnitudes along the b/c direction, and $v_a^{(n)} = \partial \epsilon_{\mathbf{k}}^{(n)} / \partial k_a$ the group velocity of the n -th band [56]. Details on the derivation of Eq. (11) can be found in Appendix A.

III. EXCEPTIONAL RING MODEL

As previously introduced, non-Hermitian systems can exhibit heightened sensitivity near exceptional points [70, 93, 102, 104, 133–151]. Due to the greater number of non-Hermitian degrees of freedom, exceptional points occur far more generically than usual Hermitian nodal points [152], commonly resulting appearing as extended 1D exceptional lines or loops. Here, we investigate a specific scenario: a system with a ring of exceptional points in its Brillouin zone, such that its Hamiltonian matrix is not of full rank along a 1-dimensional locus.

Along this ring, the conduction and valence bands coalesce, and the absence of a complete set of left eigenvectors results in ill-defined Berry curvature with divergent behavior, as later given by Eq. (15). Prior research on exceptional structures has focused mainly on their band topology [74, 102, 136, 138, 140–144, 153–156], with limited insights into transport behaviors. This motivates our deeper investigation into the higher-order effects of Berry curvature, specifically the BCD and the BCP.

With this motivation, we introduce a tilted exceptional ring model to investigate the nonlinear exceptional response. Specifically, we consider a two-dimensional tilted massive Dirac model with gain and loss

$$\hat{H}(\mathbf{k}) = d_0 \sigma_0 + \sum_{i=x,y,z} d_i \sigma_i, \quad (12)$$

where $d_0 = t_0 k_x$, $d_x = v k_y$, $d_y = \eta v k_x$, and $d_z = i\gamma + M - \alpha k^2$. Here, $\sigma_{x,y,z}$ are the Pauli matrices and σ_0 denotes the 2×2 identity matrix. We set $\eta = -1$ throughout in our numerical calculations. The parameter γ represents the strength of gain or loss [157–161]. The tilted term $t_0 k_x \sigma_0$ breaks the inversion symmetry and breaks the rotational symmetry in the shape of the Fermi surface, playing a pivotal role in triggering the nonlinear Hall effect in our model.




FIG. 1. (a) Three-dimensional plot of the real energy band structure [Eq. (13)] of our exceptional ring model Eq. (12). (b) The $k_y = 0$ slice of the real part of the energy band structure, shown for parameters as $t_0 = 0.2$, $\gamma = v = 1$, $M = \alpha = 5$ that fixes the ring radius as $R = \gamma/v = \sqrt{M/\alpha} = 1$.

For our non-Hermitian Hamiltonian in Eq. (12), we can solve the eigenenergies for the upper (+) and lower (−) bands as follows:

$$\epsilon_{\mathbf{k}}^{(\pm)} = t_0 k_x \pm \sqrt{(i\gamma + M - \alpha k^2)^2 + v^2 k^2}, \quad (13)$$

where $k^2 = k_x^2 + k_y^2$. In this model, the bands touch along an exceptional ring, depicted in red as shown in Fig. 1(a). The corresponding $k_y = 0$ slice of the energy band structure is also plotted in Fig. 1(b) to show the gapless crossings more clearly. The exceptional ring in our model, where the degeneracy of eigenvalue vanishes

[74, 93, 133–136]: $(i\gamma + M - \alpha k^2)^2 + v^2 k^2 = 0$, is fixed at

$$k^2 = R^2 = M/\alpha = \gamma^2/v^2, \quad (14)$$

where R is the radius of the exceptional ring in momentum space, and can be written as $R = \sqrt{M/\alpha} = \gamma/v$. Physically, M represents the Dirac mass, with $2M$ describing the energy gap at $(k_x, k_y) = (0, 0)$. In particular, when $M = 0$, a half-quantized Hall conductance emerges at the center of the Dirac cone [8]. The parameter α accounts for the modified quadratic momentum-dependent Dirac mass, expressed as $M - \alpha k^2$, while the parameter $t_0 \pm v$, characterizes the slope of the exceptional point's linear energy spectrum, where v is the strength of the Rashba spin-orbit coupling.

Further properties of this exceptional ring are described in the appendices. Specifically, as detailed in Appendix B, we demonstrate that the Hamiltonian (12) retains inversion symmetry at $t_0 = 0$, which is however broken under $t_0 \neq 0$. Moreover, in Appendix C, we establish that the Hamiltonian (12) does not exhibit time-reversal symmetry, irrespective of the value of t_0 . To explore the properties of the model near the exceptional ring, we provide analytical expansions of both the Hamiltonian and the eigenenergies near these points in Appendix D. In the following section, we proceed to explore the nonlinear responses with such an exceptional ring.

A. Model Berry curvature and quantum metric

In our work, we investigate both the extrinsic and intrinsic nonlinear Hall effects. Here, we first focus on the extrinsic nonlinear response. This arises from the biorthogonal Berry curvature which, with the help of Eq. (3), can be expressed for our non-Hermitian model (12) as follows [1–5, 8–10]:

$$\begin{aligned} \Omega_z^{(\pm)} &= \varepsilon^{xyz} \Omega_{xy}^{(\pm)} \\ &= \text{Re} \left\{ i \left[\langle \partial_{k_x} \psi_L^{(\pm)} | \partial_{k_y} \psi_R^{(\pm)} \rangle - \langle \partial_{k_y} \psi_L^{(\pm)} | \partial_{k_x} \psi_R^{(\pm)} \rangle \right] \right\} \\ &= \pm \text{Re} \left\{ \frac{\eta v^2 (i\gamma + M + \alpha k^2)}{2 \left[(i\gamma + M - \alpha k^2)^2 + v^2 k^2 \right]^{3/2}} \right\}. \end{aligned} \quad (15)$$

Equation (15) indicates that the Berry curvature diverges at the exceptional ring where Eq. (14) is satisfied, greatly enhancing the BCD there. Details on the derivation of Eq. (15) can be found in Appendix E.

According to Eq. (11), the intrinsic nonlinear Hall effect arises from the biorthogonal quantum metric. This quantum metric for our two-band non-Hermitian Dirac model (12) can be expressed as [55, 56, 108, 113–115, 117]

$$\mathcal{G}_{ab}^{(\pm)} = \frac{1}{4} \text{Re} \left[(\partial_{k_a} \hat{\mathbf{d}}) \cdot (\partial_{k_b} \hat{\mathbf{d}}) \right], \quad (16)$$

where $\hat{\mathbf{d}} = (d_0/d, d_x/d, d_y/d, d_z/d)$ with $d = (d_x^2 + d_y^2 + d_z^2)^{1/2} = \sqrt{(i\gamma + M - \alpha k^2)^2 + v^2 k^2}$. derivation details of Eq. (16) can be found in Appendix F.

IV. NONLINEAR RESPONSES

A. Berry curvature dipole and extrinsic nonlinear response

The BCD contributes to the higher-order effects in the extrinsic nonlinear response. To calculate the BCD [Eq. (10)] in our exceptional ring system, a key term is the momentum gradient of the non-Hermitian Berry curvature [Eq. (15)] with respect to k_x , which is given by

$$\begin{aligned} P_{yxx}^{(\pm)} &\equiv \frac{\partial \Omega_z^{(\pm)}}{\partial k_x} \\ &= \pm \text{Re} \left\{ \frac{\eta k_x v^2 [2\alpha (g_-^2 + v^2 k^2) - 3g_+ (v^2 - 2\alpha g_-)]}{2 (g_-^2 + v^2 k^2)^{5/2}} \right\}, \end{aligned} \quad (17)$$

where $g_{\pm} = i\gamma + M \pm \alpha k^2$ and $\partial_{k_x} \Omega_z^{(\pm)}(-k_x) = -\partial_{k_x} \Omega_z^{(\pm)}(k_x)$.

In our work, we consider zero temperature, such that the corresponding equilibrium Fermi-Dirac distribution function can be expressed as $f_0^{(\pm)} = f_0^{(\pm)} [\text{Re}(\epsilon_{\mathbf{k}}^{(\pm)}) - E_F] = \Theta [E_F - \text{Re}(\epsilon_{\mathbf{k}}^{(\pm)})]$, with the Fermi energy E_F and the Heaviside function $\Theta(x)$. For simplicity, we choose the parameters that satisfy the condition $\sum_{\pm} [f_0^{(\pm)} \Omega_z^{(\pm)}]_{k_x = \pm\pi} = 0$. As such, the first term in the BCD [Eq. (10)] vanishes. To achieve this, we need a significant energy gap at $k_x = \pm\pi$, as illustrated in Fig. 1(a), i.e., when $k_x = \pm\pi$ is fixed, a significant gap naturally appears in the 2D slice energy spectrum as a function of k_y , as shown in Fig. 1(a). Given that the Berry curvature [Eq. (15)] is even about k_x , we then have $\Omega_z^{(\pm)}(k_x = \pi) = \Omega_z^{(\pm)}(k_x = -\pi)$. To achieve non-vanishing BCD, we need the conditions: $\Theta [E_F - \text{Re}(\epsilon_{k_x=\pi}^{(\pm)})] = \Theta [E_F - \text{Re}(\epsilon_{k_x=-\pi}^{(\pm)})]$ and $\text{Re}(\epsilon_{k_x=\pm\pi}^{(-)}) < E_F < \text{Re}(\epsilon_{k_x=\pm\pi}^{(+)})$ ($k_y \in [-\pi, \pi]$) as shown in Fig. 1(a). For our model, the BCD [Eq. (10)] hence does not vanish only at non-zero tilt ($t_0 \neq 0$).

According to Eq. (10), the predominant contribution to the nonlinear response stems from the states near the Fermi surface. Specifically, when the Fermi surface approaches the exceptional ring, these exceptional states can give rise to an intense response. In Figs. 2(a1)-2(d1), we plot the BCD near the exceptional ring ($R = 1$) with the Fermi level $E_F \in [-0.3, 0.3]$. Here, strong extrinsic nonlinear BCD responses are evident, highlighted by the two prominent peaks, particularly when the loss parameter γ increases from 0.1 to 1.5.

Next, we examine the BCD density profile $P_{yxx}^{(\pm)}(\mathbf{k})$ itself. As depicted in Figs. 2(a2)-2(d2), near the exceptional ring where bands coalesce at $k^2 = R^2 = M/\alpha$, the BCD density profile markedly diverges. The details for the scaling behavior of the BCD density profile near the exceptional ring can be found in Appendix G.




FIG. 2. **Extrinsic nonlinear response of our exceptional ring with fixed radius $R = \gamma/v = \sqrt{M/\alpha} = 1$, with $M = \alpha = 5$.** (a1-d1): The Berry curvature dipole (BCD) [Eq. (10)], showing strong peaks that increase with the non-Hermiticity γ . (a2)-(d2): The BCD density profile $P_{yx}^{(-)}(\mathbf{k})$ [Eq. (17)] in the k_x - k_y plane. The band structure tilt is $t_0 = 0.2$.




FIG. 3. **Maximum value of the extrinsic nonlinear response of an exceptional ring as a function of γ , for fixed radius $R = 1$.** We set $\gamma = v$ and $M = \alpha$ such that $R = 1$, and numerically plot the maximum value of the Berry curvature dipole (BCD) [Eq. (10)]. For different values of $M = \alpha = 3.5$ (black circle), 4.0 (blue cross), 4.5 (red triangle), and 5.0 (green square), the numerical data follows the same trend as the solid line, which is fitted from the ansatz $\text{Max}(D_{yx}) = \exp(a_0 \ln \gamma + b_0)$ ($a_0 = 0.5313$ and $b_0 = 9.4517$), demonstrating an approximate but universal power scaling law $\text{Max}(D_{yx}) \propto \sqrt{\gamma}$. The other parameters are the same as those in Fig. 2.

To understand how the BCD response scales with the non-Hermiticity for an exceptional ring at fixed radius (we set it to $R = 1$), we evaluate the maximum value of the BCD as a function of γ for different parameters $M = \alpha$, while keeping the radius $R = 1$ fixed. As shown in Fig. 3, the maximum BCD value increases like $\sqrt{\gamma}$ as non-Hermiticity γ increases, indicating that non-Hermiticity enhances the extrinsic nonlinear Hall effect. To understand this, we then consider the condition of $0 \ll |\gamma| = |v| \ll \alpha = M$ and expand Eq. (33) near the exceptional ring along $k = \sqrt{M/\alpha} + \delta k = 1 + \delta k$ and $k_x = (1 + \delta k) \cos \theta$ with $M = \alpha$ and $\delta k \rightarrow 0$ as follows:

$$\begin{aligned}
\frac{\partial \Omega_z^{(\pm)}}{\partial k_x} &= \pm \eta [(1+\delta k) \cos \theta] v^2 \operatorname{Re} \left\{ \frac{\alpha v [v(1+\delta k)^2 + (i\gamma + \alpha(1-(1+\delta k)^2))^2/v]}{v^{5/2} [v(1+\delta k)^2 + (i\gamma + \alpha(1-(1+\delta k)^2))^2/v]^{5/2}} \right. \\
&\quad \left. - \frac{3[i\gamma + \alpha(1+(1+\delta k)^2)] [v^2 - 2\alpha(i\gamma + \alpha(1-(1+\delta k)^2))]}{2v^{5/2} [v(1+\delta k)^2 + (i\gamma + \alpha(1-(1+\delta k)^2))^2/v]^{5/2}} \right\} \\
&\approx \pm \eta [(1+\delta k) \cos \theta] \operatorname{Re} \left\{ \frac{\sqrt{v}\alpha}{[v((1+\delta k)^2 - 1) + 2i\alpha(1-(1+\delta k)^2)]^{3/2}} \right. \\
&\quad \left. - \frac{3v^{3/2}[i\gamma + \alpha(1+(1+\delta k)^2)]}{2[v((1+\delta k)^2 - 1) + 2i\alpha(1-(1+\delta k)^2)]^{5/2}} - \frac{3\alpha v^{3/2}[1 - 2i\alpha/v - \alpha^2[1-(1+\delta k)^2]^2/v^2]}{[v((1+\delta k)^2 - 1) + 2i\alpha(1-(1+\delta k)^2)]^{5/2}} \right\} \quad (18) \\
&\approx \pm \eta [(1+\delta k) \cos \theta] \operatorname{Re} \left\{ \frac{(v/\alpha)^{1/2}}{[2i(1-(1+\delta k)^2)]^{3/2}} + \frac{6i(v/\alpha)^{1/2}}{[2i(1-(1+\delta k)^2)]^{5/2}} \right. \\
&\quad \left. - \frac{6(v/\alpha)^{3/2}}{[2i(1-(1+\delta k)^2)]^{5/2}} - \frac{3i(v/\alpha)^{5/2}}{2[2i(1-(1+\delta k)^2)]^{5/2}} \right\} \quad (19) \\
&\approx \pm \eta [(1+\delta k) \cos \theta] \operatorname{Re} \left\{ \frac{(v/\alpha)^{1/2}}{[2i(1-(1+\delta k)^2)]^{3/2}} + \frac{6i(v/\alpha)^{1/2}}{[2i(1-(1+\delta k)^2)]^{5/2}} \right\} \quad (20) \\
&\propto (v/\alpha)^{1/2} \propto \sqrt{\gamma}. \quad (21)
\end{aligned}$$

In the first line that approximates Eq. (18), higher-order small terms such as $\alpha^2[1-(1+\delta k)^2]^2/v$ in both the numerator and denominator are neglected. Subsequently, higher-order small terms such as $\alpha^2[1-(1+\delta k)^2]^2/v^2$ in the numerator and $v((1+\delta k)^2-1)$ in the denominator are also neglected in the second approximation (19). Furthermore, higher-order small terms such as $(v/\alpha)^{3/2}$ and $(v/\alpha)^{5/2}$ are neglected in the third approximation (20). Therefore, we obtain the derived analytical scaling behavior $\sqrt{\gamma}$ described by Eq. (21), which aligns with our numerical results shown in Fig. 3. However, this approximation is not entirely accurate if v is exceedingly small in the initial δk approximation. Numerically, the smallest δk is determined by the resolution of the k intervals, which in turn depends on the physical size of the system. For finite lattices, the maximum D_{yxx} scales as $\sqrt{\gamma}$ down to the spatial scale of lattices.

From Fig. 3, the maximum of the BCD decreases significantly as $\gamma = v \rightarrow 0$. In fact, it tends to zero, as can be seen by expanding the term $\partial \Omega_z^{(\pm)}/\partial k_x$ [Eq. (17)] in the BCD as follows:

$$\begin{aligned}
&\lim_{\gamma=v \rightarrow 0} \frac{\partial \Omega_z^{(\pm)}}{\partial k_x} \\
&\approx \pm \lim_{\gamma=v \rightarrow 0} \operatorname{Re} \left\{ \frac{\eta k_x v^2 [2\alpha(M - \alpha k^2)^2 + 6\alpha(M + \alpha k^2)(M - \alpha k^2)]}{2(M - \alpha k^2)^5} \right\} \\
&= \pm \lim_{\gamma=v \rightarrow 0} \operatorname{Re} \left\{ \frac{\eta k_x v^2 [\alpha(M - \alpha k^2) + 3\alpha(M + \alpha k^2)]}{(M - \alpha k^2)^4} \right\} \\
&= \pm \lim_{\gamma=v \rightarrow 0} \operatorname{Re} \left[\frac{2\alpha k_x v^2 (2M + \alpha k^2)}{(M - \alpha k^2)^4} \right] \\
&= \pm \lim_{\gamma \rightarrow 0} \operatorname{Re} \left[\frac{2\alpha k_x \gamma^2 (2 + k^2)}{M^3 (1 - k^2)^4} \right]. \quad (22)
\end{aligned}$$

The approximation Eq. (22) shows that $\lim_{\gamma=v \rightarrow 0} D_{yxx} =$

0, which is consistent with our numerical results in Fig. 3.

B. Berry connection polarizability and intrinsic nonlinear response

Here, we examine the intrinsic nonlinear response in our exceptional ring model (12). The yxx -component intrinsic nonlinear velocity response $\bar{v}_{yxx}^{\text{BCP}}$ [Eq. (11)] under the BCP is given by [46, 47, 49, 51]

$$\begin{aligned}
\bar{v}_{yxx}^{\text{BCP}} &= \sum_{n,n'=+,-}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{xx} \int \frac{dk_x}{2\pi} \left[\frac{f_0^{(n)} \mathcal{G}_{xx}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_y=\pi}^{k_y=-\pi} \\
&\quad + \sum_{n,n'=+,-}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{xx} \int \frac{dk_y}{2\pi} \left[\frac{-f_0^{(n)} \mathcal{G}_{yy}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_x=\pi}^{k_x=-\pi} \\
&\quad - \sum_{n,n'=+,-}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{xx} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} f_0^{(n)} \left[\frac{\partial_{k_y} \mathcal{G}_{xx}^{(n)} - \partial_{k_x} \mathcal{G}_{yy}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right], \quad (23)
\end{aligned}$$

where $\Gamma_{xx} = 2\mathcal{S}\mathcal{F}_x^2/\hbar$ with \mathcal{F}_x being the magnitude of the applied force, $v_a^{(n)} = \partial \epsilon_{\mathbf{k}}^{(n)}/\partial k_a$ the group velocity of the n -th band [56], and the quantum metric components given by ($\eta = -1$)

$$\mathcal{G}_{xx}^{(\pm)} = \operatorname{Re} \left\{ \frac{v^2}{4d^2} \left\{ 1 + \frac{k_x^2 [4\alpha(M + i\gamma) - v^2]}{d^2} \right\} \right\}, \quad (24)$$

$$\mathcal{G}_{yx}^{(\pm)} = \operatorname{Re} \left\{ \frac{k_x k_y v^2 [4\alpha(M + i\gamma) - v^2]}{4d^4} \right\}. \quad (25)$$




FIG. 4. **Fermi surface (red curves)** of our exceptional ring model [Eq. (12)] with model parameters $t_0 = 0.2$, $\gamma = v = 1$, and $M = \alpha = 5$, correspond to a ring radius $R = 1$. The Fermi region is slightly broadened due to the nonzero Fermi energy of $E_F = 0.5$, and is asymmetric about k_x but symmetric about k_y . The color denotes the BCP density profile $P_{xy}^{BCP(-)}(\mathbf{k})$ defined in Eq. (33).

Here we have written $d = (d_x^2 + d_y^2 + d_z^2)^{1/2} = \sqrt{(i\gamma + M - \alpha k^2)^2 + v^2 k^2}$. Furthermore, we can obtain

$$\frac{\partial \mathcal{G}_{xx}^{(\pm)}}{\partial k_y} = \text{Re} \left\{ \frac{k_y v^2 (2\alpha g_- - v^2)}{2(g_-^2 + v^2 k^2)^3} \times \left\{ (g_-^2 + v^2 k^2) + 2k_x^2 [4\alpha(M + i\gamma) - v^2] \right\} \right\}, \quad (26)$$

$$\frac{\partial \mathcal{G}_{yx}^{(\pm)}}{\partial k_x} = \text{Re} \left\{ \frac{k_y v^2 [4\alpha(M + i\gamma) - v^2]}{4(g_-^2 + v^2 k^2)^3} \times \left[(g_-^2 + v^2 k^2) + 4k_x^2 (2\alpha g_- - v^2) \right] \right\}, \quad (27)$$

where $g_- = i\gamma + M - \alpha k^2$.

First, the integrand $[f_0^{(\pm)} \mathcal{G}_{xx}^{(\pm)} / (\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)})]_{k_y = -\pi}^{k_y = \pi}$ in Eq. (23) is even about k_y and $\epsilon_{k_y = \pi}^{(\pm)} = \epsilon_{k_y = -\pi}^{(\pm)}$, so that it vanishes after the integration over the Fermi surface (as shown in Fig. 4). Figure 4 shows that the Fermi surface of our tilted massive 2D Dirac model [Eq. (12)] is asymmetric about k_x but symmetric about k_y . Second, the integrand $[-f_0^{(\pm)} \mathcal{G}_{yx}^{(\pm)} / (\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)})]_{k_x = -\pi}^{k_x = \pi}$ in Eq. (23) is odd about k_y so that it also vanishes after the integration over the Fermi surface. Indeed, these symmetry consideration dictate the vanish of some BCP components. From Eqs. (26) and (27), the integrand $[\partial_{k_y} \mathcal{G}_{xx}^{(\pm)} - \partial_{k_x} \mathcal{G}_{yx}^{(\pm)}] / [\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)}]$ is odd about k_y and even about k_x , leading to the vanishing of \bar{v}_{yx}^{BCP} after the integration over the Fermi surface. In the same way, we can find that the xyx -component $\bar{v}_{xyx}^{BCP} = -\bar{v}_{yxx}^{BCP}$ equals zero too. Besides, we also have $\bar{v}_{xyy}^{BCP} = 0$.

One nonvanishing component in the intrinsic nonlinear

response is the xyy -component intrinsic nonlinear velocity \bar{v}_{xyy}^{BCP} [Eq. (11)]:

$$\begin{aligned} \bar{v}_{xyy}^{BCP} &= \sum_{n, n' = +, -}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{yy} \int \frac{dk_y}{2\pi} \left[\frac{f_0^{(n)} \mathcal{G}_{yy}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right]_{k_x = -\pi}^{k_x = \pi} \\ &+ \sum_{n, n' = +, -}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{yy} \int \frac{dk_x}{2\pi} \left[\frac{-f_0^{(n)} \mathcal{G}_{xy}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right]_{k_y = -\pi}^{k_y = \pi} \\ &- \sum_{n, n' = +, -}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{yy} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} f_0^{(n)} \left[\frac{\partial_{k_x} \mathcal{G}_{yy}^{(n)} - \partial_{k_y} \mathcal{G}_{xy}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right], \end{aligned} \quad (28)$$

where $\Gamma_{yy} = 2\mathcal{S}\mathcal{F}_y^2/\hbar$ with the magnitude of force \mathcal{F}_y , and the quantum metric components are ($\eta = -1$)

$$\mathcal{G}_{yy}^{(\pm)} = \text{Re} \left\{ \frac{v^2}{4d^2} \left\{ 1 + \frac{k_y^2 [4\alpha(M + i\gamma) - v^2]}{d^2} \right\} \right\}, \quad (29)$$

$$\mathcal{G}_{xy}^{(\pm)} = \text{Re} \left\{ \frac{k_x k_y v^2 [4\alpha(M + i\gamma) - v^2]}{4d^4} \right\}, \quad (30)$$

giving rise to momentum-space derivatives

$$\frac{\partial \mathcal{G}_{yy}^{(\pm)}}{\partial k_x} = \text{Re} \left\{ \frac{k_x v^2 (2\alpha g_- - v^2)}{2(g_-^2 + v^2 k^2)^3} \times \left\{ (g_-^2 + v^2 k^2) + 2k_y^2 [4\alpha(M + i\gamma) - v^2] \right\} \right\}, \quad (31)$$

$$\frac{\partial \mathcal{G}_{xy}^{(\pm)}}{\partial k_y} = \text{Re} \left\{ \frac{k_x v^2 [4\alpha(M + i\gamma) - v^2]}{4(g_-^2 + v^2 k^2)^3} \times \left[(g_-^2 + v^2 k^2) + 4k_y^2 (2\alpha g_- - v^2) \right] \right\}. \quad (32)$$

Now, the integrand $[f_0^{(\pm)} \mathcal{G}_{yy}^{(\pm)} / (\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)})]_{k_y = -\pi}^{k_y = \pi}$ in Eq. (28) is even about k_x and $\Theta [E_F - \text{Re}(\epsilon_{k_x = \pi}^{(\pm)})] = \Theta [E_F - \text{Re}(\epsilon_{k_x = -\pi}^{(\pm)})]$ under present parameters as shown in Fig. 1(a). Under the current parameters illustrated in Fig. 1(a), a significant energy gap is present to satisfy this condition, so that it vanishes after the integration over the Fermi surface. Second, the integrand $\sum_{n, n' = +, -} [-f_0^{(n)} \mathcal{G}_{xy}^{(n)} / (\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')})]_{k_y = -\pi}^{k_y = \pi}$ in Eq. (28) vanishes after the summation of the band index $n, n' = +, -$ with the condition $\epsilon_{k_y = \pi}^{(\pm)} = \epsilon_{k_y = -\pi}^{(\pm)}$. From Eqs. (31) and (32), the integrand $[\partial_{k_x} \mathcal{G}_{yy}^{(\pm)} - \partial_{k_y} \mathcal{G}_{xy}^{(\pm)}] / [\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)}]$ is odd about k_x and even about k_y , hence a nonzero \bar{v}_{xyy}^{BCP} is permitted after the integration over the Fermi surface. Moreover, we have $\bar{v}_{yx}^{BCP} = -\bar{v}_{xyx}^{BCP} \neq 0$ and $\bar{v}_{yyx}^{BCP} = 0$. Therefore, we only need to focus on the nonzero component \bar{v}_{xyy}^{BCP} .




FIG. 5. **Intrinsic nonlinear response with different values of $\gamma = v$ for fixed radius $R = \gamma/v = \sqrt{M/\alpha} = 1$ with $M = \alpha = 5$.** (a1-d1): The intrinsic nonlinear velocity $\bar{v}_{xyy}^{\text{BCP}}$ [Eq. (28)] for the Berry connection polarizability, which is suppressed with increasing non-Hermiticity. (a2)-(d2): The BCP density profile $P_{xyy}^{\text{BCP}(-)}(\mathbf{k})$ [Eq. (33)] in the k_x - k_y plane. The other parameters are $t_0 = 0.2$.




FIG. 6. **Peak of the intrinsic nonlinear response $\bar{v}_{xyy}^{\text{BCP}}$ as a function of non-Hermiticity $\gamma = v$.** We plot the maximum value of $\bar{v}_{xyy}^{\text{BCP}}$ [Eq. (28)] with $\gamma = v$, such that $R = 1$. Results are similar across different values of $M = \alpha = 3.5$ (black circle), 4.0 (blue cross), 4.5 (red triangle), and 5.0 (green square). The solid line shows the power-law dependence of the maximum $\bar{v}_{xyy}^{\text{BCP}}$, which is fitted from the ansatz $\text{Max}(\bar{v}_{xyy}^{\text{BCP}}/\Gamma_{yy}) = \exp(a_1 \ln \gamma + b_1)$ in the small γ regime. For the case $M = \alpha = 3.5$, we have $a_1 = -0.5370$, $b_1 = 6.7711$ for the regime $\gamma = v \in [0.02, 0.21]$, approximately obeying the scaling law $\text{Max}(\bar{v}_{xyy}^{\text{BCP}}/\Gamma_{yy}) \propto \sqrt{1/\gamma}$. However, in the large γ regime, there is no such power-law scaling. The other parameters are the same as those in Fig. 5.

In Figs. 5(a1)-5(d1), we plot the intrinsic nonlinear velocity $\bar{v}_{xyy}^{\text{BCP}}$ [Eq. (28)] under different values of the tunable effective gain/loss γ at fixed radius $R = 1$ of the exceptional ring. Similar to the BCD, $\bar{v}_{xyy}^{\text{BCP}}$ exhibits a pair of peaks within the Fermi energy range $E_F \in [-0.3, 0.3]$.

However, in contrast to the BCD, Fig. 6 illustrates that the intrinsic nonlinear Hall effect is suppressed as non-Hermiticity increases. Here, we can also expand the integrand of $\bar{v}_{xyy}^{\text{BCP}}$ under large $\gamma = v$ as follows:

$$P_{xyy}^{\text{BCP}(\pm)}(\mathbf{k}) \equiv - \left[\frac{\partial_{k_x} \mathcal{G}_{yy}^{(\pm)} - \partial_{k_y} \mathcal{G}_{xy}^{(\pm)}}{\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)}} \right]$$

$$= \pm \text{Re} \left[\frac{k_x v^2 (v^2 + 4\alpha^2 k^2)}{8(g^2 + v^2 k^2)^{5/2}} \right] \quad (33)$$

$$\approx \pm \lim_{\gamma=v \rightarrow \infty} \text{Re} \left[\frac{k_x}{8v(k^2 - 1)^{5/2}} \right], \quad (34)$$

where $P_{xyy}^{\text{BCP}(\pm)}(\mathbf{k})$ is defined as the BCP density profile. This expansion (34) demonstrates that $\lim_{\gamma=v \rightarrow \infty} \bar{v}_{xyy}^{\text{BCP}} = 0$, indicating a complete suppression of the intrinsic response at large non-Hermiticity. Moreover, we plot the BCP density profile $P_{xyy}^{\text{BCP}(-)}(\mathbf{k})$ in Eq. (33) in the k_x - k_y plane as shown in Figs. 5(a2)-5(d2). Similar to the BCD density profile, near the exceptional ring at $k^2 = R^2 = M/\alpha = 1$ where the bands coalesce, the BCP density profile also markedly diverges. The details for the scaling behavior of the BCP density profile near the exceptional ring can be found as follows. Here, we illustrate that a detectable intrinsic nonlinear Hall response in our model requires a particular scaling law for the BCP density profile. We expand the BCP density profile near the exceptional ring along $k^2 = (\sqrt{M/\alpha} + \delta k)^2$ with $\gamma = v\sqrt{M/\alpha}$ and $\delta k \rightarrow 0$, where δk denotes a small deviation from the exceptional ring. Along the direction of $k_y = 0$, we further expand the BCP density profile under $k^2 = k_x^2 + k_y^2 = (\sqrt{M/\alpha} + \delta k_x)^2 + (0 + \delta k_y)^2$ to the lowest order of δk_x

as follows:

$$P_{xyy}^{\text{BCP}(\pm)}(\delta k_x) \approx \pm \text{Re} \left[\frac{\sqrt{M/\alpha}(v^2 + 4M\alpha)}{32\sqrt{2v} \left(v\sqrt{M/\alpha} - 2iM\right)^{5/2}} \right] \times (\delta k_x)^{-5/2}. \quad (35)$$

Besides, Fig. 6 also illustrates that the maximum $\bar{v}_{xyy}^{\text{BCP}}$ appears to diverge when $\gamma = v$ is relatively small but non-negligible. To understand this, we then consider the above condition of $0 \ll \gamma^2 = v^2 \ll 1$ and expand Eq. (33) near the exceptional ring along $k = \sqrt{M/\alpha} + \delta k = 1 + \delta k$ and $k_x = (1 + \delta k) \cos \theta$ with $M = \alpha$ and $\delta k \rightarrow 0$ as follows:

$$\begin{aligned} & - \left[\frac{\partial_{k_x} \mathcal{G}_{yy}^{(\pm)} - \partial_{k_y} \mathcal{G}_{xy}^{(\pm)}}{\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)}} \right] \\ &= \pm \text{Re} \left\{ \frac{[(1 + \delta k) \cos \theta] v^2 [v^2 + 4\alpha^2(1 + \delta k)^2]}{8v^{5/2} [v(1 + \delta k)^2 + [i\gamma + \alpha(1 - (1 + \delta k)^2)]^2/v]^{5/2}} \right\} \\ &\approx \pm \text{Re} \left\{ \frac{[(1 + \delta k) \cos \theta] \alpha^2 (1 + \delta k)^2}{2\sqrt{v} [v((1 + \delta k)^2 - 1) + 2i\alpha(1 - (1 + \delta k)^2)]^{5/2}} \right\} \quad (36) \\ &\approx \pm \text{Re} \left\{ \frac{[(1 + \delta k) \cos \theta] \alpha^2 (1 + \delta k)^2}{2\sqrt{v} [2i\alpha(1 - (1 + \delta k)^2)]^{5/2}} \right\} \propto \sqrt{\frac{1}{v}} = \sqrt{\frac{1}{\gamma}}. \quad (37) \end{aligned}$$

In the first line that approximates Eq. (36), higher-order small terms such as v^2 in the numerator and $\alpha^2[1 - (1 + \delta k)^2]^2/v$ in the denominator are neglected. Subsequently, the term $v((1 + \delta k)^2 - 1)$ in the denominator is also neglected in the second approximation (37). Therefore, we obtain the derived analytical scaling behavior $\sqrt{1/\gamma}$ described by Eq. (37), which aligns with our numerical results shown in Fig. 6. However, this approximation is not entirely accurate if v is exceedingly small in the initial δk approximation. Numerically, the smallest δk is determined by the resolution of the k intervals, which in turn depends on the physical size of the

system. For finite lattices, the maximum $\bar{v}_{xyy}^{\text{BCP}}$ scales as $\sqrt{1/\gamma}$ down to the spatial scale of lattices.

Interestingly beyond this regime, in the true limit of $\gamma = v \rightarrow 0$, the maximum $\bar{v}_{xyy}^{\text{BCP}}$ approaches zero, mirroring the behavior of the BCD. As such, we can also expand the integrand of Eq. (33) as follows:

$$- \lim_{\gamma=v \rightarrow 0} \left[\frac{\partial_{k_x} \mathcal{G}_{yy}^{(\pm)} - \partial_{k_y} \mathcal{G}_{xy}^{(\pm)}}{\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)}} \right] \approx \pm \lim_{\gamma=v \rightarrow 0} \text{Re} \left[\frac{k_x k^2 v^2 \alpha^2}{2(M - \alpha k^2)^5} \right], \quad (38)$$

which shows that $\lim_{\gamma=v \rightarrow 0} \bar{v}_{xyy}^{\text{BCP}} = 0$.

V. DISCUSSION

In this work, we investigate how an exceptional ring with macroscopic numbers of defective states can non-trivially influence the nonlinear response, focusing on both extrinsic and intrinsic nonlinear Hall effects. Based on an ansatz tilted dissipative Dirac model which possess an exceptional ring band structure, our findings reveal that the BCD in the extrinsic Hall effect intensifies with increasing non-Hermiticity. Conversely, the intrinsic nonlinear velocity, as quantified by the BCP, is suppressed significantly with growing non-Hermiticity, resulting in a subdued intrinsic nonlinear response. These behavior are unique to exceptional rings and represent new nonlinear response behaviors from the interplay of non-Hermitian defectiveness and extended exceptional band crossings.

ACKNOWLEDGMENTS

We acknowledge helpful discussions with Xiao-Bin Qiang, Hao-Jie Lin, and Rui Chen. C.H.L. and F.Q. acknowledge support from the QEP2.0 Grant from the Singapore National Research Foundation (Grant No. NRF2021-QEP2-02-P09) and the Singapore Ministry of Education Academic Research Fund Tier-II Grant (Award No. MOE-T2EP50222-0003).

Appendix A: Derivation of Eq. (11) for the BCP

In this appendix, we present the detailed derivation of Eq. (11) as follows:

$$\begin{aligned}
\bar{v}_{abc}^{\text{BCP}} &= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \left[\frac{v_a^{(n)} \mathcal{G}_{bc}^{(n)} - v_b^{(n)} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \frac{\partial f_0^{(n)}}{\partial \epsilon_{\mathbf{k}}^{(n)}} \\
&= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \left[\frac{(\partial_{k_a} f_0^{(n)}) \mathcal{G}_{bc}^{(n)} - (\partial_{k_b} f_0^{(n)}) \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \\
&= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_b}{2\pi} \left[\frac{f_0^{(n)} \mathcal{G}_{bc}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_a=-\pi}^{k_a=\pi} - \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \left[f_0^{(n)} \frac{\partial_{k_a} \mathcal{G}_{bc}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \\
&\quad + \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_a}{2\pi} \left[\frac{-f_0^{(n)} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_b=-\pi}^{k_b=\pi} - \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \left[-f_0^{(n)} \frac{\partial_{k_b} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \\
&= \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_b}{2\pi} \left[\frac{f_0^{(n)} \mathcal{G}_{bc}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_a=-\pi}^{k_a=\pi} + \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{dk_a}{2\pi} \left[\frac{-f_0^{(n)} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right] \Big|_{k_b=-\pi}^{k_b=\pi} \\
&\quad - \sum_{n,n'}^{\epsilon_{\mathbf{k}}^{(n)} \neq \epsilon_{\mathbf{k}}^{(n')}} \Gamma_{bc} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} f_0^{(n)} \left[\frac{\partial_{k_a} \mathcal{G}_{bc}^{(n)} - \partial_{k_b} \mathcal{G}_{ac}^{(n)}}{\epsilon_{\mathbf{k}}^{(n)} - \epsilon_{\mathbf{k}}^{(n')}} \right], \tag{A1}
\end{aligned}$$

where $\Gamma_{bc} = 2\mathcal{S}\mathcal{F}_b\mathcal{F}_c/\hbar$ with the system area being \mathcal{S} , $\mathcal{F}_{b/c}$ is the perturbing force field and $v_a^{(n)} = \partial \epsilon_{\mathbf{k}}^{(n)}/\partial k_a$ is the group velocity in the n -th band [56].

Appendix B: Inversion symmetry

The purpose of this section is to demonstrate that the Hamiltonian (12) in the main text retains inversion symmetry when $t_0 = 0$, and breaks inversion when $t_0 \neq 0$. Our Hamiltonian (12) under inversion transformation can be expressed as follows:

$$\mathcal{I} \hat{\mathcal{H}}(\mathbf{k}) \mathcal{I}^{-1} = t_0 k_x \sigma_0 - v k_y \sigma_x - \eta v k_x \sigma_y + (i\gamma + M - \alpha k^2) \sigma_z = \hat{\mathcal{H}}(-\mathbf{k}) + 2t_0 k_x \sigma_0, \tag{B1}$$

where $\mathcal{I} = \sigma_z$ [162] is the inversion operator, and we use

$$(\sigma_z)(\sigma_0)(\sigma_z)^{-1} = \sigma_0, \tag{B2}$$

$$(\sigma_z)(\sigma_x)(\sigma_z)^{-1} = -\sigma_x, \tag{B3}$$

$$(\sigma_z)(\sigma_y)(\sigma_z)^{-1} = -\sigma_y, \tag{B4}$$

$$(\sigma_z)(\sigma_z)(\sigma_z)^{-1} = \sigma_z. \tag{B5}$$

According to Eq. (B1), we find the condition $\mathcal{I} \hat{\mathcal{H}}(\mathbf{k}) \mathcal{I}^{-1} = \hat{\mathcal{H}}(-\mathbf{k})$ under $t_0 = 0$, which confirms the inversion symmetry. However, if $t_0 \neq 0$, Eq. (B1) indicates that this inversion symmetry is broken.

Appendix C: Time-reversal symmetry is broken

This section aims to establish that the Hamiltonian (12) in the main text does not exhibit time-reversal symmetry, regardless of whether t_0 is zero or not.

The Hamiltonian (12) under time-reversal transformation can be expressed as follows

$$\begin{aligned}
\mathcal{T} \hat{\mathcal{H}}(\mathbf{k}) \mathcal{T}^{-1} &= t_0 k_x \sigma_0 - v k_y \sigma_x - \eta v k_x \sigma_y - (i\gamma + M - \alpha k^2) \sigma_z \\
&= \hat{\mathcal{H}}(-\mathbf{k}) + 2t_0 k_x \sigma_0 - 2(i\gamma + M - \alpha k^2) \sigma_z, \tag{C1}
\end{aligned}$$

where $\mathcal{T} = i\sigma_y \mathcal{K}$ [162] is the time-reversal operator with the complex conjugate operator \mathcal{K} . Following this, we have $\mathcal{K}\hat{\mathcal{H}}(\mathbf{k})\mathcal{K}^{-1} = \hat{\mathcal{H}}^*(\mathbf{k})$, with following relations

$$(i\sigma_y)(\sigma_0)(i\sigma_y)^{-1} = \sigma_0, \quad (\text{C2})$$

$$(i\sigma_y)(\sigma_x)(i\sigma_y)^{-1} = -\sigma_x, \quad (\text{C3})$$

$$(i\sigma_y)(K\sigma_y K^{-1})(i\sigma_y)^{-1} = -\sigma_y, \quad (\text{C4})$$

$$(i\sigma_y)(\sigma_z)(i\sigma_y)^{-1} = -\sigma_z. \quad (\text{C5})$$

As a result, Eq. (C1) demonstrates that the time-reversal symmetry is broken, since $\mathcal{T}\hat{\mathcal{H}}(\mathbf{k})\mathcal{T}^{-1} \neq \hat{\mathcal{H}}(-\mathbf{k})$ whether t_0 is zero.

Appendix D: Perturbation near the exceptional ring

To investigate the behavior of our model near the exceptional ring, we expand the Hamiltonian (12) near the exceptional ring. Here, we focus on the expansion along $k^2 = (\sqrt{M/\alpha} + \delta k)^2$ with the radius of the ring given by $R = \sqrt{M/\alpha}$ and $\gamma = v\sqrt{M/\alpha}$. We consider $\delta \vec{k} \rightarrow 0$, representing a slight deviation from the ring.

Specifically, along the direction with $k_y = 0$, the Hamiltonian expansion at $k^2 = k_x^2 + k_y^2 = (\sqrt{M/\alpha} + \delta k_x)^2 + (0 + \delta k_y)^2$ (i.e., near the exceptional ring) to the lowest order of δk_x and δk_y is given by:

$$\hat{\mathcal{H}}(\delta k_x, \delta k_y) \approx t_0 \left(\sqrt{\frac{M}{\alpha}} + \delta k_x \right) \sigma_0 + v(\delta k_y) \sigma_x + \eta v \left(\sqrt{\frac{M}{\alpha}} + \delta k_x \right) \sigma_y + \left[i v \sqrt{\frac{M}{\alpha}} - 2\sqrt{M\alpha}(\delta k_x) \right] \sigma_z, \quad (\text{D1})$$

where $\delta k_x \rightarrow 0$ and $\delta k_y \rightarrow 0$.

Moreover, the corresponding eigenenergies for the model Hamiltonian (12) near the exceptional ring can be expanded with respect to the lowest order of δk_x as follows:

$$\epsilon_{\delta k_x}^{(\pm)} \approx t_0 \left(\sqrt{\frac{M}{\alpha}} + \delta k_x \right) \pm \sqrt{2v \left(v \sqrt{\frac{M}{\alpha}} - 2iM \right) \delta k_x}, \quad (\text{D2})$$

where we have ignored the higher-order $(\delta k_y)^2$ terms in the square root.

Appendix E: Derivations of Eq. (15) for the form of the Berry curvature

In this section, we show the detailed derivations of Eq. (15) as follows [163]:

$$\begin{aligned} \Omega_z^{(\pm)} &= \epsilon_{zxy} \Omega_{xy}^{(\pm)} = \epsilon_{zxy} \text{Re} \left(\partial_{k_x} \mathcal{A}_{\pm}^y - \partial_{k_y} \mathcal{A}_{\pm}^x \right) \\ &= \text{Re} \epsilon_{zxy} i \left[\left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_x} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_y} \right\rangle - \left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_y} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_x} \right\rangle \right] \\ &= \text{Re} \epsilon_{zxy} i \sum_{n'=+,-} \left[\left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_x} \middle| \psi_R^{(n')} \right\rangle \left\langle \psi_L^{(n')} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_y} \right\rangle - \left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_y} \middle| \psi_R^{(n')} \right\rangle \left\langle \psi_L^{(n')} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_x} \right\rangle \right] \\ &= \text{Re} \epsilon_{zxy} i \left[\left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_x} \middle| \psi_R^{(\mp)} \right\rangle \left\langle \psi_L^{(\mp)} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_y} \right\rangle - \left\langle \frac{\partial \psi_L^{(\pm)}}{\partial k_y} \middle| \psi_R^{(\mp)} \right\rangle \left\langle \psi_L^{(\mp)} \middle| \frac{\partial \psi_R^{(\pm)}}{\partial k_x} \right\rangle \right] \\ &= \text{Re} \epsilon_{zxy} i \left\{ \frac{\left\langle \psi_L^{(\pm)} \middle| \frac{\partial \hat{\mathcal{H}}}{\partial k_x} \middle| \psi_R^{(\mp)} \right\rangle \left\langle \psi_L^{(\mp)} \middle| \frac{\partial \hat{\mathcal{H}}}{\partial k_y} \middle| \psi_R^{(\pm)} \right\rangle}{\left[\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)} \right]^2} - \frac{\left\langle \psi_L^{(\pm)} \middle| \frac{\partial \hat{\mathcal{H}}}{\partial k_y} \middle| \psi_R^{(\mp)} \right\rangle \left\langle \psi_L^{(\mp)} \middle| \frac{\partial \hat{\mathcal{H}}}{\partial k_x} \middle| \psi_R^{(\pm)} \right\rangle}{\left[\epsilon_{\mathbf{k}}^{(\pm)} - \epsilon_{\mathbf{k}}^{(\mp)} \right]^2} \right\} \\ &= \pm \text{Re} \left\{ \frac{\eta v^2 (i\gamma + M + \alpha k^2)}{2 \left[(i\gamma + M - \alpha k^2)^2 + v^2 k^2 \right]^{3/2}} \right\}, \quad (\text{E1}) \end{aligned}$$

where $\mathcal{A}_+^a = i\langle\psi_L^{(+)}|\partial_{k_a}|\psi_R^{(+)}\rangle$ is the a -component of the biorthogonal Berry connection [163] $\mathcal{A}_+ = i\langle\psi_L^{(+)}|\nabla_{\mathbf{k}}|\psi_R^{(+)}\rangle$, and we have used $\langle\partial_{k_a}\psi_L^{(+)}|\psi_R^{(-)}\rangle = \langle\psi_L^{(+)}|\frac{\partial\hat{H}}{\partial k_a}|\psi_R^{(-)}\rangle / [\epsilon_{\mathbf{k}}^{(+)} - \epsilon_{\mathbf{k}}^{(-)}]$ and $\langle\psi_L^{(-)}|\partial_{k_a}\psi_R^{(+)}\rangle = \langle\psi_L^{(-)}|\frac{\partial\hat{H}}{\partial k_a}|\psi_R^{(+)}\rangle / [\epsilon_{\mathbf{k}}^{(+)} - \epsilon_{\mathbf{k}}^{(-)}]$.

Furthermore, we can expand the non-Hermitian Berry curvature (E1) under $k^2 = (\sqrt{M/\alpha} + \delta k)^2$ (i.e., near the exceptional ring) with respect to the lowest order of δk as follows:

$$\Omega_z^{(\pm)} \approx \pm \text{Re} \left[\frac{i\eta v(\delta k)^{-3/2}}{4\sqrt{2v\left(v\sqrt{\frac{M}{\alpha}} - 2iM\right)}} \right], \quad (\text{E2})$$

where we have used $\gamma = v\sqrt{M/\alpha}$ and $\delta k \rightarrow 0$.

Appendix F: Biorthogonal quantum metric

Here, we show the detailed derivations of the biorthogonal quantum metric for any two-band model Hamiltonian as follows:

$$\begin{aligned} \mathcal{G}_{ab}^{(+)} &\equiv \frac{1}{2} \text{Re} \left[\mathcal{Q}_{ab}^{(+)} + \mathcal{Q}_{ba}^{(+)} \right] \\ &= \frac{1}{2} \text{Re} \left[\langle\partial_{k_a}\psi_L^{(+)}|\partial_{k_b}\psi_R^{(+)}\rangle + \langle\partial_{k_b}\psi_L^{(+)}|\partial_{k_a}\psi_R^{(+)}\rangle \right] - \frac{1}{2} \text{Re} \left[\langle\partial_{k_a}\psi_L^{(+)}|\hat{P}^{(+)}|\partial_{k_b}\psi_R^{(+)}\rangle + \langle\partial_{k_b}\psi_L^{(+)}|\hat{P}^{(+)}|\partial_{k_a}\psi_R^{(+)}\rangle \right] \\ &= \frac{1}{2} \text{Re} \left[\langle\partial_{k_a}\psi_L^{(+)}|\hat{P}^{(-)}|\partial_{k_b}\psi_R^{(+)}\rangle + \langle\partial_{k_b}\psi_L^{(+)}|\hat{P}^{(-)}|\partial_{k_a}\psi_R^{(+)}\rangle \right] \\ &= \frac{1}{2} \text{Re} \left[(\langle\partial_{k_a}\psi_L^{(+)}|\hat{P}^{(-)}\rangle\langle\hat{P}^{(-)}|\partial_{k_b}\psi_R^{(+)}\rangle) + (\langle\partial_{k_b}\psi_L^{(+)}|\hat{P}^{(-)}\rangle\langle\hat{P}^{(-)}|\partial_{k_a}\psi_R^{(+)}\rangle) \right] \\ &= \frac{1}{2} \text{Re} \left[\langle\psi_L^{(+)}|\partial_{k_a}\hat{P}^{(+)}|\psi_R^{(-)}\rangle\langle\psi_L^{(-)}|\partial_{k_b}\hat{P}^{(+)}|\psi_R^{(+)}\rangle + \langle\psi_L^{(-)}|\partial_{k_a}\hat{P}^{(+)}|\psi_R^{(+)}\rangle\langle\psi_L^{(+)}|\partial_{k_b}\hat{P}^{(+)}|\psi_R^{(-)}\rangle \right] \\ &= \frac{1}{2} \text{Re} \text{Tr} \left[(\partial_{k_a}\hat{P}^{(+)})(\partial_{k_b}\hat{P}^{(+)}) \right] = \frac{1}{8} \text{Re} \text{Tr} \left[(\partial_{k_a}\hat{\mathbf{d}} \cdot \boldsymbol{\sigma})(\partial_{k_b}\hat{\mathbf{d}} \cdot \boldsymbol{\sigma}) \right] = \frac{1}{4} \text{Re} [(\partial_{k_a}\hat{\mathbf{d}}) \cdot (\partial_{k_b}\hat{\mathbf{d}})], \end{aligned} \quad (\text{F1})$$

where $\hat{P}^{(\pm)} = |\psi_R^{(\pm)}\rangle\langle\psi_L^{(\pm)}| = \frac{1}{2} (\mathbb{I} \pm \hat{\mathbf{d}} \cdot \boldsymbol{\sigma})$ [55] is the biorthogonal projection operator of the ‘ \pm ’ band, $\hat{\mathbf{d}} = (d_0/d, d_x/d, d_y/d, d_z/d)$ with $d = (d_x^2 + d_y^2 + d_z^2)^{1/2}$, $\boldsymbol{\sigma} = (\sigma_0, \sigma_x, \sigma_y, \sigma_z)$, $[\sigma_a, \sigma_b] = 2i\varepsilon_{abc}\sigma_c$, $\{\sigma_a, \sigma_b\} = 2\delta_{ab}\mathbb{I}$ with the identity matrix \mathbb{I} , $a, b \in \{x, y, z\}$, $\sum_{n'=\pm} |\psi_R^{(n')}\rangle\langle\psi_L^{(n')}| = \mathbb{I}$, we use $\hat{P}^{(-)}|\partial_{k_b}\psi_R^{(+)}\rangle = \partial_{k_b}\hat{P}^{(+)}|\psi_R^{(+)}\rangle$ [117], $\sigma_a\sigma_b = i\varepsilon_{abc}\sigma_c + \delta_{ab}\mathbb{I}$ and

$$\begin{aligned} \text{Tr} \left[(\partial_{k_a}\hat{\mathbf{d}} \cdot \boldsymbol{\sigma})(\partial_{k_b}\hat{\mathbf{d}} \cdot \boldsymbol{\sigma}) \right] &= \frac{1}{d} \text{Tr} [(\partial_{k_a}d_0\sigma_0 + \partial_{k_a}d_x\sigma_x + \partial_{k_a}d_y\sigma_y + \partial_{k_a}d_z\sigma_z)(\partial_{k_b}d_0\sigma_0 + \partial_{k_b}d_x\sigma_x + \partial_{k_b}d_y\sigma_y + \partial_{k_b}d_z\sigma_z)] \\ &= \frac{1}{d} \text{Tr} [(\partial_{k_a}d_0)(\partial_{k_b}d_0) + (\partial_{k_a}d_x)(\partial_{k_b}d_x) + (\partial_{k_a}d_y)(\partial_{k_b}d_y) + (\partial_{k_a}d_z)(\partial_{k_b}d_z)]\mathbb{I} \\ &= 2(\partial_{k_a}\hat{\mathbf{d}}) \cdot (\partial_{k_b}\hat{\mathbf{d}}). \end{aligned} \quad (\text{F2})$$

We also have $\mathcal{G}_{ab}^{(-)} = \mathcal{G}_{ab}^{(+)}$.

Appendix G: Scaling behavior of the BCD density profile

In this section, we will illustrate the scaling behavior of the BCD density profile near the exceptional ring. Here, we illustrate that a detectable extrinsic nonlinear Hall response in our model requires a particular scaling law for the BCD density profile. We expand the BCD density profile near the exceptional ring along $k^2 = (\sqrt{M/\alpha} + \delta k)^2$ with $\gamma = v\sqrt{M/\alpha}$ and $\delta k \rightarrow 0$, where δk denotes a small deviation from the exceptional ring. Along the direction of $k_y = 0$, we further expand the BCD density profile under $k^2 = k_x^2 + k_y^2 = (\sqrt{M/\alpha} + \delta k_x)^2 + (0 + \delta k_y)^2$ to the lowest order of δk_x as follows:

$$P_{yxx}^{(\pm)}(\delta k_x) \approx \mp \frac{3\eta}{8} \text{Re} \left[\frac{-v\left(v\sqrt{M/\alpha} + 2iM\right)}{2(M/\alpha)(v^2 + 4M\alpha)} \right]^{1/2} (\delta k_x)^{-5/2}. \quad (\text{G1})$$

- [1] Ryogo Kubo, “Statistical-mechanical theory of irreversible processes. I. General theory and simple applications to magnetic and conduction problems,” *Journal of the physical society of Japan* **12**, 570–586 (1957).
- [2] Ryogo Kubo, Mario Yokota, and Sadao Nakajima, “Statistical-mechanical theory of irreversible processes. II. Response to thermal disturbance,” *Journal of the Physical Society of Japan* **12**, 1203–1211 (1957).
- [3] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, “Quantized Hall Conductance in a Two-Dimensional Periodic Potential,” *Phys. Rev. Lett.* **49**, 405–408 (1982).
- [4] Shun-Qing Shen, *Topological insulators* (Springer Nature Singapore Pte Ltd., 2017).
- [5] Doru Sticlet, Balázs Dóra, and Cătălin Pașcu Moca, “Kubo Formula for Non-Hermitian Systems and Tachyon Optical Conductivity,” *Phys. Rev. Lett.* **128**, 016802 (2022).
- [6] Fang Qin, Shuai Li, Z. Z. Du, C. M. Wang, Wenqing Zhang, Dapeng Yu, Hai-Zhou Lu, and X. C. Xie, “Theory for the Charge-Density-Wave Mechanism of 3D Quantum Hall Effect,” *Phys. Rev. Lett.* **125**, 206601 (2020).
- [7] Fang Qin, Rui Chen, and Hai-Zhou Lu, “Phase transitions in intrinsic magnetic topological insulator with high-frequency pumping,” *Journal of Physics: Condensed Matter* **34**, 225001 (2022).
- [8] Fang Qin, Ching Hua Lee, and Rui Chen, “Light-induced half-quantized Hall effect and axion insulator,” *Phys. Rev. B* **108**, 075435 (2023).
- [9] Fang Qin, Ruizhe Shen, Linhu Li, and Ching Hua Lee, “Kinked linear response from non-Hermitian cold-atom pumping,” *Phys. Rev. A* **109**, 053311 (2024).
- [10] Fang Qin, Rui Chen, and Ching Hua Lee, “Light-enhanced nonlinear hall effect,” arXiv:2401.18038 (2024).
- [11] Mark O Goerbig, “Quantum Hall effects,” arXiv:0909.1998 (2009).
- [12] Takahiro Morimoto and Naoto Nagaosa, “Topological nature of nonlinear optical effects in solids,” *Science advances* **2**, e1501524 (2016).
- [13] Ching Hua Lee, Xiao Zhang, and Bochen Guan, “Negative differential resistance and characteristic nonlinear electromagnetic response of a topological insulator,” *Scientific reports* **5**, 18008 (2015).
- [14] Ching Hua Lee, Han Hoe Yap, Tommy Tai, Gang Xu, Xiao Zhang, and Jiangbin Gong, “Enhanced higher harmonic generation from nodal topology,” *Phys. Rev. B* **102**, 035138 (2020).
- [15] Tommy Tai and Ching Hua Lee, “Anisotropic nonlinear optical response of nodal-loop materials,” *Phys. Rev. B* **103**, 195125 (2021).
- [16] M. S. Mrudul and Gopal Dixit, “High-harmonic generation from monolayer and bilayer graphene,” *Phys. Rev. B* **103**, 094308 (2021).
- [17] Amar Bharti, M. S. Mrudul, and Gopal Dixit, “High-harmonic spectroscopy of light-driven nonlinear anisotropic anomalous Hall effect in a Weyl semimetal,” *Phys. Rev. B* **105**, 155140 (2022).
- [18] Amar Bharti, Misha Ivanov, and Gopal Dixit, “How massless are Weyl fermions in Weyl semimetals,” *Phys. Rev. B* **108**, L020305 (2023).
- [19] Yang-Yang Lv, Jinlong Xu, Shuang Han, Chi Zhang, Yadong Han, Jian Zhou, Shu-Hua Yao, Xiao-Ping Liu, Ming-Hui Lu, Hongming Weng, *et al.*, “High-harmonic generation in Weyl semimetal β -WP2 crystals,” *Nature Communications* **12**, 6437 (2021).
- [20] Shambhu Ghimire and David A Reis, “High-harmonic generation from solids,” *Nature physics* **15**, 10–16 (2019).
- [21] Zahra Nourbakhsh, Nicolas Tancogne-Dejean, Hamed Merdji, and Angel Rubio, “High Harmonics and Isolated Attosecond Pulses from MgO,” *Phys. Rev. Appl.* **15**, 014013 (2021).
- [22] Inti Sodemann and Liang Fu, “Quantum nonlinear Hall effect induced by Berry curvature dipole in time-reversal invariant materials,” *Phys. Rev. Lett.* **115**, 216806 (2015).
- [23] Z. Z. Du, C. M. Wang, Hai-Zhou Lu, and X. C. Xie, “Band Signatures for Strong Nonlinear Hall Effect in Bilayer WTe_2 ,” *Phys. Rev. Lett.* **121**, 266601 (2018).
- [24] Z. Z. Du, C. M. Wang, Shuai Li, Hai-Zhou Lu, and X. C. Xie, “Disorder-induced nonlinear Hall effect with time-reversal symmetry,” *Nature communications* **10**, 1–6 (2019).
- [25] Z. Z. Du, Hai-Zhou Lu, and X. C. Xie, “Nonlinear Hall effects,” *Nature Reviews Physics* **3**, 744–752 (2021).
- [26] Carmine Ortix, “Nonlinear Hall Effect with Time-Reversal Symmetry: Theory and Material Realizations,” *Advanced Quantum Technologies* **4**, 2100056 (2021).
- [27] Arka Bandyopadhyay, Nesta Benno Joseph, and Awadhesh Narayan, “Non-linear Hall Effects: Mechanisms and Materials,” arXiv:2401.02282 (2024).
- [28] Kaifei Kang, Tingxin Li, Egon Sohn, Jie Shan, and Kin Fai Mak, “Nonlinear anomalous Hall effect in few-layer WTe_2 ,” *Nature materials* **18**, 324–328 (2019).
- [29] Qiong Ma, Su-Yang Xu, Huitao Shen, David MacNeill, Valla Fatemi, Tay-Rong Chang, Andrés M Mier Valdivia, Sanfeng Wu, Zongzheng Du, Chuang-Han Hsu, *et al.*, “Observation of the nonlinear Hall effect under time-reversal-symmetric conditions,” *Nature* **565**, 337–342 (2019).
- [30] Xing-Guo Ye, Huiying Liu, Peng-Fei Zhu, Wen-Zheng Xu, Shengyuan A. Yang, Nianze Shang, Kaihui Liu, and Zhi-Min Liao, “Control over Berry Curvature Dipole with Electric Field in WTe_2 ,” *Phys. Rev. Lett.* **130**, 016301 (2023).
- [31] Jieun Lee, Zefang Wang, Hongchao Xie, Kin Fai Mak, and Jie Shan, “Valley magnetoelectricity in single-layer MoS_2 ,” *Nature materials* **16**, 887–891 (2017).
- [32] Joolee Son, Kyung-Han Kim, Y. H. Ahn, Hyun-Woo Lee, and Jieun Lee, “Strain Engineering of the Berry Curvature Dipole and Valley Magnetization in Monolayer MoS_2 ,” *Phys. Rev. Lett.* **123**, 036806 (2019).
- [33] Oleg Olegovich Shvetsov, Varnava Denisovich Esin, Anna Vladimirovna Timonina, Nikolay Nikolaevich Kolesnikov, and EV Deviatov, “Nonlinear Hall effect in three-dimensional weyl and dirac semimetals,” *JETP Letters* **109**, 715–721 (2019).
- [34] Yang Zhang, Yan Sun, and Binghai Yan, “Berry curvature dipole in Weyl semimetal materials: An ab initio

- study,” *Phys. Rev. B* **97**, 041101 (2018).
- [35] Dushyant Kumar, Chuang-Han Hsu, Raghav Sharma, Tay-Rong Chang, Peng Yu, Junyong Wang, Goki Eda, Gengchiao Liang, and Hyunsoo Yang, “Room-temperature nonlinear Hall effect and wireless radiofrequency rectification in Weyl semimetal TaIrTe₄,” *Nature Nanotechnology* **16**, 421–425 (2021).
- [36] Chuanchang Zeng, Snehasish Nandy, and Sumanta Tewari, “Nonlinear transport in weyl semimetals induced by berry curvature dipole,” *Phys. Rev. B* **103**, 245119 (2021).
- [37] Pan He, Hiroki Isobe, Dapeng Zhu, Chuang-Han Hsu, Liang Fu, and Hyunsoo Yang, “Quantum frequency doubling in the topological insulator Bi₂Se₃,” *Nature Communications* **12**, 698 (2021).
- [38] Jorge I. Facio, Dmitri Efremov, Klaus Koepf, Jih-Shih You, Inti Sodemann, and Jeroen van den Brink, “Strongly Enhanced Berry Dipole at Topological Phase Transitions in BiTeI,” *Phys. Rev. Lett.* **121**, 246403 (2018).
- [39] Sheng-Chin Ho, Ching-Hao Chang, Yu-Chiang Hsieh, Shun-Tsung Lo, Botsz Huang, Thi-Hai-Yen Vu, Carmine Ortix, and Tse-Ming Chen, “Hall effects in artificially corrugated bilayer graphene without breaking time-reversal symmetry,” *Nature Electronics* **4**, 116–125 (2021).
- [40] Junxi Duan, Yu Jian, Yang Gao, Huimin Peng, Jinrui Zhong, Qi Feng, Jinhai Mao, and Yugui Yao, “Giant Second-Order Nonlinear Hall Effect in Twisted Bilayer Graphene,” *Phys. Rev. Lett.* **129**, 186801 (2022).
- [41] Yue-Xin Huang, Xiaolong Feng, Hui Wang, Cong Xiao, and Shengyuan A. Yang, “Intrinsic Nonlinear Planar Hall Effect,” *Phys. Rev. Lett.* **130**, 126303 (2023).
- [42] Subhajit Sinha, Pratap Chandra Adak, Atasi Chakraborty, Kamal Das, Koyendril Debnath, LD Varma Sangani, Kenji Watanabe, Takashi Taniguchi, Umesh V Waghmare, Amit Agarwal, *et al.*, “Berry curvature dipole senses topological transition in a moiré superlattice,” *Nature Physics* **18**, 765–770 (2022).
- [43] Pierre A. Pantaleón, Tony Low, and Francisco Guinea, “Tunable large Berry dipole in strained twisted bilayer graphene,” *Phys. Rev. B* **103**, 205403 (2021).
- [44] Cheng-Ping Zhang, Jiewen Xiao, Benjamin T. Zhou, Jin-Xin Hu, Ying-Ming Xie, Binghai Yan, and K. T. Law, “Giant nonlinear Hall effect in strained twisted bilayer graphene,” *Phys. Rev. B* **106**, L041111 (2022).
- [45] Da Ma, Ying Xiong, and Justin CW Song, “Skew-scattering pockels effect and metallic electro-optics in gapped bilayer graphene,” arXiv:2407.12096 (2024).
- [46] Yang Gao, Shengyuan A. Yang, and Qian Niu, “Field induced positional shift of bloch electrons and its dynamical implications,” *Phys. Rev. Lett.* **112**, 166601 (2014).
- [47] Chong Wang, Yang Gao, and Di Xiao, “Intrinsic Nonlinear Hall Effect in Antiferromagnetic Tetragonal CuMnAs,” *Phys. Rev. Lett.* **127**, 277201 (2021).
- [48] Ching Hua Lee, Martin Claassen, and Ronny Thomale, “Band structure engineering of ideal fractional Chern insulators,” *Phys. Rev. B* **96**, 165150 (2017).
- [49] Huiying Liu, Jianzhou Zhao, Yue-Xin Huang, Weikang Wu, Xian-Lei Sheng, Cong Xiao, and Shengyuan A. Yang, “Intrinsic Second-Order Anomalous Hall Effect and Its Application in Compensated Antiferromagnets,” *Phys. Rev. Lett.* **127**, 277202 (2021).
- [50] Shen Lai, Huiying Liu, Zhaowei Zhang, Jianzhou Zhao, Xiaolong Feng, Naizhou Wang, Chaolong Tang, Yuanda Liu, KS Novoselov, Shengyuan A Yang, *et al.*, “Third-order nonlinear Hall effect induced by the Berry-connection polarizability tensor,” *Nature Nanotechnology* **16**, 869–873 (2021).
- [51] Huiying Liu, Jianzhou Zhao, Yue-Xin Huang, Xiaolong Feng, Cong Xiao, Weikang Wu, Shen Lai, Wei-bo Gao, and Shengyuan A. Yang, “Berry connection polarizability tensor and third-order Hall effect,” *Phys. Rev. B* **105**, 045118 (2022).
- [52] Anyuan Gao, Yu-Fei Liu, Jian-Xiang Qiu, Barun Ghosh, Thais V. Trevisan, Yugo Onishi, Chaowei Hu, Tiema Qian, Hung-Ju Tien, Shao-Wen Chen, *et al.*, “Quantum metric nonlinear Hall effect in a topological antiferromagnetic heterostructure,” *Science* **381**, 181 (2023).
- [53] Daniel Kaplan, Tobias Holder, and Binghai Yan, “Unification of Nonlinear Anomalous Hall Effect and Nonreciprocal Magnetoresistance in Metals by the Quantum Geometry,” *Phys. Rev. Lett.* **132**, 026301 (2024).
- [54] Naizhou Wang, Daniel Kaplan, Zhaowei Zhang, Tobias Holder, Ning Cao, Aifeng Wang, Xiaoyuan Zhou, Feifei Zhou, Zhengzhi Jiang, Chusheng Zhang, *et al.*, “Quantum-metric-induced nonlinear transport in a topological antiferromagnet,” *Nature* **621**, 487–492 (2023).
- [55] Zheng-Yang Zhuang and Zhongbo Yan, “Extrinsic and intrinsic nonlinear Hall effects across Berry-dipole transitions,” *Phys. Rev. B* **107**, L161102 (2023).
- [56] Zheng-Yang Zhuang and Zhongbo Yan, “Intrinsic nonlinear Hall effect in two-dimensional honeycomb topological antiferromagnets,” *Phys. Rev. B* **109**, 174443 (2024).
- [57] Naomichi Hatano and David R. Nelson, “Localization Transitions in Non-Hermitian Quantum Mechanics,” *Phys. Rev. Lett.* **77**, 570–573 (1996).
- [58] Naomichi Hatano and David R. Nelson, “Vortex pinning and non-Hermitian quantum mechanics,” *Phys. Rev. B* **56**, 8651–8673 (1997).
- [59] Naomichi Hatano and David R. Nelson, “Non-Hermitian delocalization and eigenfunctions,” *Phys. Rev. B* **58**, 8384–8390 (1998).
- [60] Shunyu Yao and Zhong Wang, “Edge States and Topological Invariants of Non-Hermitian Systems,” *Phys. Rev. Lett.* **121**, 086803 (2018).
- [61] Shunyu Yao, Fei Song, and Zhong Wang, “Non-Hermitian Chern Bands,” *Phys. Rev. Lett.* **121**, 136802 (2018).
- [62] Flore K. Kunst, Elisabet Edvardsson, Jan Carl Budich, and Emil J. Bergholtz, “Biorthogonal Bulk-Boundary Correspondence in Non-Hermitian Systems,” *Phys. Rev. Lett.* **121**, 026808 (2018).
- [63] Kazuki Yokomizo and Shuichi Murakami, “Non-Bloch Band Theory of Non-Hermitian Systems,” *Phys. Rev. Lett.* **123**, 066404 (2019).
- [64] Ching Hua Lee and Ronny Thomale, “Anatomy of skin modes and topology in non-Hermitian systems,” *Phys. Rev. B* **99**, 201103 (2019).
- [65] Tian-Shu Deng and Wei Yi, “Non-Bloch topological invariants in a non-Hermitian domain wall system,” *Phys. Rev. B* **100**, 035102 (2019).

- [66] Kohei Kawabata, Nobuyuki Okuma, and Masatoshi Sato, “Non-Bloch band theory of non-Hermitian Hamiltonians in the symplectic class,” *Phys. Rev. B* **101**, 195147 (2020).
- [67] Kazuki Yokomizo and Shuichi Murakami, “Non-Bloch band theory and bulk–edge correspondence in non-Hermitian systems,” *Progress of Theoretical and Experimental Physics* **2020**, 12A102 (2020).
- [68] Kohei Kawabata, Masatoshi Sato, and Ken Shiozaki, “Higher-order non-Hermitian skin effect,” *Phys. Rev. B* **102**, 205118 (2020).
- [69] Yifei Yi and Zhesen Yang, “Non-Hermitian Skin Modes Induced by On-Site Dissipations and Chiral Tunneling Effect,” *Phys. Rev. Lett.* **125**, 186802 (2020).
- [70] Kazuki Yokomizo and Shuichi Murakami, “Topological semimetal phase with exceptional points in one-dimensional non-Hermitian systems,” *Phys. Rev. Research* **2**, 043045 (2020).
- [71] Lei Xiao, Tianshu Deng, Kunkun Wang, Gaoyan Zhu, Zhong Wang, Wei Yi, and Peng Xue, “Non-Hermitian bulk–boundary correspondence in quantum dynamics,” *Nature Physics* **16**, 761 (2020).
- [72] Kai Zhang, Zhesen Yang, and Chen Fang, “Universal non-Hermitian skin effect in two and higher dimensions,” *Nature communications* **13**, 2496 (2022).
- [73] Zhesen Yang, Kai Zhang, Chen Fang, and Jiangping Hu, “Non-Hermitian Bulk-Boundary Correspondence and Auxiliary Generalized Brillouin Zone Theory,” *Phys. Rev. Lett.* **125**, 226402 (2020).
- [74] Xiao Zhang, Guangjie Li, Yuhan Liu, Tommy Tai, Ronny Thomale, and Ching Hua Lee, “Tidal surface states as fingerprints of non-Hermitian nodal knot metals,” *Communications Physics* **4**, 1–10 (2021).
- [75] Kazuki Yokomizo and Shuichi Murakami, “Non-Bloch band theory in bosonic Bogoliubov–de Gennes systems,” *Phys. Rev. B* **103**, 165123 (2021).
- [76] Balaganchi A. Bhargava, Ion Cosma Fulga, Jeroen van den Brink, and Ali G. Moghaddam, “Non-Hermitian skin effect of dislocations and its topological origin,” *Phys. Rev. B* **104**, L241402 (2021).
- [77] Kazuki Yokomizo, “Non-Bloch Band Theory of Non-Hermitian Systems and Bulk-Edge Correspondence,” in *Non-Bloch Band Theory of Non-Hermitian Systems* (Springer, 2022) pp. 35–56.
- [78] S. M. Rafi-Ul-Islam, Zhuo Bin Siu, Haydar Sahin, Ching Hua Lee, and Mansoor B. A. Jalil, “Critical hybridization of skin modes in coupled non-Hermitian chains,” *Phys. Rev. Res.* **4**, 013243 (2022).
- [79] Ruizhe Shen and Ching Hua Lee, “Non-Hermitian skin clusters from strong interactions,” *Communications Physics* **5**, 238 (2022).
- [80] Fang Qin, Ruizhe Shen, and Ching Hua Lee, “Non-Hermitian squeezed polarons,” *Phys. Rev. A* **107**, L010202 (2023).
- [81] Russell Yang, Jun Wei Tan, Tommy Tai, Jin Ming Koh, Linhu Li, Stefano Longhi, and Ching Hua Lee, “Designing non-Hermitian real spectra through electrostatics,” *Science Bulletin* **67**, 1865–1873 (2022).
- [82] Fang Qin, Ye Ma, Ruizhe Shen, and Ching Hua Lee, “Universal competitive spectral scaling from the critical non-Hermitian skin effect,” *Phys. Rev. B* **107**, 155430 (2023).
- [83] Linhu Li and Ching Hua Lee, “Non-Hermitian pseudogaps,” *Science Bulletin* **67**, 685–690 (2022).
- [84] Hui Jiang and Ching Hua Lee, “Dimensional Transmutation from Non-Hermiticity,” *Phys. Rev. Lett.* **131**, 076401 (2023).
- [85] Tommy Tai and Ching Hua Lee, “Zoology of non-Hermitian spectra and their graph topology,” *Phys. Rev. B* **107**, L220301 (2023).
- [86] Tuo Wan, Kai Zhang, Junkai Li, Zhesen Yang, and Zhaoju Yang, “Observation of the geometry-dependent skin effect and dynamical degeneracy splitting,” *Science Bulletin* **68**, 2330–2335 (2023).
- [87] Yu-Peng Wang, Chen Fang, and Jie Ren, “Absence of measurement-induced entanglement transition due to feedback-induced skin effect,” *Phys. Rev. B* **110**, 035113 (2024).
- [88] Ruizhe Shen, Fang Qin, Jean-Yves Desaulles, Zlatko Papić, and Ching Hua Lee, “Enhanced many-body quantum scars from the non-Hermitian Fock skin effect,” arXiv:2403.02395 (2024).
- [89] Alexander Poddubny, Janet Zhong, and Shanhui Fan, “Mesoscopic non-Hermitian skin effect,” *Phys. Rev. A* **109**, L061501 (2024).
- [90] Wen-Tan Xue and Ching Hua Lee, “Topologically protected negative entanglement,” arXiv:2403.03259 (2024).
- [91] Raphaël Bouganne, Manel Bosch Aguilera, Alexis Ghermaoui, Jérôme Beugnon, and Fabrice Gerbier, “Anomalous decay of coherence in a dissipative many-body system,” *Nature Physics* **16**, 21–25 (2020).
- [92] Lei Pan, Xin Chen, Yu Chen, and Hui Zhai, “Non-Hermitian linear response theory,” *Nature Physics* **16**, 767 (2020).
- [93] Yong Xu, Sheng-Tao Wang, and L-M Duan, “Weyl exceptional rings in a three-dimensional dissipative cold atomic gas,” *Physical Review Letters* **118**, 045701 (2017).
- [94] Ching Hua Lee, Linhu Li, Ronny Thomale, and Jiangbin Gong, “Unraveling non-Hermitian pumping: Emergent spectral singularities and anomalous responses,” *Phys. Rev. B* **102**, 085151 (2020).
- [95] Wang Jiong-Hao, Tao Yu-Liang, and Xu Yong, “Anomalous Transport Induced by Non-Hermitian Anomalous Berry Connection in Non-Hermitian Systems,” *Chinese Phys. Lett.* **39**, 010301 (2022).
- [96] Ching Hua Lee and Stefano Longhi, “Ultrafast and anharmonic Rabi oscillations between non-Bloch bands,” *Communications Physics* **3**, 147 (2020).
- [97] Yaashnaa Singhal, Enrico Martello, Shraddha Agrawal, Tomoki Ozawa, Hannah Price, and Bryce Gadway, “Measuring the adiabatic non-Hermitian Berry phase in feedback-coupled oscillators,” *Phys. Rev. Res.* **5**, L032026 (2023).
- [98] Hannah M Price and NR Cooper, “Mapping the Berry curvature from semiclassical dynamics in optical lattices,” *Physical Review A* **85**, 033620 (2012).
- [99] Sthitadhi Roy, Adolfo G. Grushin, Roderich Moessner, and Masudul Haque, “Wave-packet dynamics on Chern-band lattices in a trap,” *Phys. Rev. A* **92**, 063626 (2015).
- [100] Christopher Oliver, Aaron Smith, Thomas Easton, Grazia Salerno, Vera Guarrera, Nathan Goldman, Giovanni Barontini, and Hannah M. Price, “Bloch oscillations along a synthetic dimension of atomic trap states,” *Phys. Rev. Res.* **5**, 033001 (2023).
- [101] Dorje C Brody, “Biorthogonal quantum mechanics,” *Journal of Physics A: Mathematical and Theoretical* **47**,

- 035305 (2013).
- [102] Ching Hua Lee, “Exceptional Bound States and Negative Entanglement Entropy,” *Phys. Rev. Lett.* **128**, 010402 (2022).
- [103] Shi-Dong Liang and Guang-Yao Huang, “Topological invariance and global Berry phase in non-Hermitian systems,” *Phys. Rev. A* **87**, 012118 (2013).
- [104] Yue-Yu Zou, Yao Zhou, Li-Mei Chen, and Peng Ye, “Detecting bulk and edge exceptional points in non-Hermitian systems through generalized Petermann factors,” *Frontiers of Physics* **19**, 23201 (2024).
- [105] JP Provost and G Vallee, “Riemannian structure on manifolds of quantum states,” *Communications in Mathematical Physics* **76**, 289–301 (1980).
- [106] Raffaele Resta, “The insulating state of matter: a geometrical theory,” *The European Physical Journal B* **79**, 121–137 (2011).
- [107] Yu-Quan Ma, Shu Chen, Heng Fan, and Wu-Ming Liu, “Abelian and non-Abelian quantum geometric tensor,” *Phys. Rev. B* **81**, 245129 (2010).
- [108] Motohiko Ezawa, “Analytic approach to quantum metric and optical conductivity in Dirac models with parabolic mass in arbitrary dimensions,” arXiv:2408.02951 (2024).
- [109] Barun Ghosh, Yugo Onishi, Su-Yang Xu, Hsin Lin, Liang Fu, and Arun Bansil, “Probing quantum geometry through optical conductivity and magnetic circular dichroism,” arXiv:2401.09689 (2024).
- [110] Päivi Törmä, Sebastiano Peotta, and Bogdan A Bernevig, “Superconductivity, superfluidity and quantum geometry in twisted multilayer systems,” *Nature Reviews Physics* **4**, 528–542 (2022).
- [111] Sebastiano Peotta, Kukka-Emilia Huhtinen, and Päivi Törmä, “Quantum geometry in superfluidity and superconductivity,” arXiv:2308.08248 (2023).
- [112] Päivi Törmä, “Essay: Where Can Quantum Geometry Lead Us?” *Phys. Rev. Lett.* **131**, 240001 (2023).
- [113] Yugo Onishi and Liang Fu, “Fundamental bound on topological gap,” *Phys. Rev. X* **14**, 011052 (2024).
- [114] Shunji Matsuura and Shinsei Ryu, “Momentum space metric, nonlocal operator, and topological insulators,” *Phys. Rev. B* **82**, 245113 (2010).
- [115] Gero von Gersdorff and Wei Chen, “Measurement of topological order based on metric-curvature correspondence,” *Phys. Rev. B* **104**, 195133 (2021).
- [116] Mengjie Yang and Ching Hua Lee, “Percolation-Induced \mathcal{PT} Symmetry Breaking,” *Phys. Rev. Lett.* **133**, 136602 (2024).
- [117] Wei Chen, “Quantum geometrical properties of topological materials,” arXiv:2406.15145 (2024).
- [118] Motohiko Ezawa, “Intrinsic nonlinear conductivity induced by quantum geometry in altermagnets and measurement of the in-plane Néel vector,” arXiv:2409.09241 (2024).
- [119] Tianyu Liu, Xiao-Bin Qiang, Hai-Zhou Lu, and X. C. Xie, “Quantum geometry in condensed matter,” *National Science Review* **104**, nwa334 (2024).
- [120] Ching Hua Lee, Wen Wei Ho, Bo Yang, Jiangbin Gong, and Zlatko Papić, “Floquet Mechanism for Non-Abelian Fractional Quantum Hall States,” *Phys. Rev. Lett.* **121**, 237401 (2018).
- [121] Monika Aidelsburger, Michael Lohse, Christian Schweizer, Marcos Atala, Julio T. Barreiro, Sylvain Nascimbène, N. R. Cooper, Immanuel Bloch, and Nathan Goldman, “Measuring the Chern number of Hofstadter bands with ultracold bosonic atoms,” *Nature Physics* **11**, 162 (2015).
- [122] Marco Mancini, Guido Pagano, Giacomo Cappellini, Lorenzo Livi, Marie Rider, Jacopo Catani, Carlo Sias, Peter Zoller, Massimo Inguscio, Marcello Dalmonte, *et al.*, “Observation of chiral edge states with neutral fermions in synthetic Hall ribbons,” *Science* **349**, 1510–1513 (2015).
- [123] Nathan Goldman, Jan C Budich, and Peter Zoller, “Topological quantum matter with ultracold gases in optical lattices,” *Nature Physics* **12**, 639 (2016).
- [124] N. R. Cooper, J. Dalibard, and I. B. Spielman, “Topological bands for ultracold atoms,” *Reviews of Modern Physics* **91**, 015005 (2019).
- [125] Rui Chen, Z. Z. Du, Hai-Peng Sun, Hai-Zhou Lu, and X. C. Xie, “Nonlinear hall effect on a disordered lattice,” *Phys. Rev. B* **110**, L081301 (2024).
- [126] Z. Z. Du, C. M. Wang, Hai-Peng Sun, Hai-Zhou Lu, and X. C. Xie, “Quantum theory of the nonlinear Hall effect,” *Nature communications* **12**, 1–7 (2021).
- [127] Cong Xiao, Z. Z. Du, and Qian Niu, “Theory of nonlinear Hall effects: Modified semiclassics from quantum kinetics,” *Phys. Rev. B* **100**, 165422 (2019).
- [128] Stepan Tsirkin and Ivo Souza, “On the separation of Hall and Ohmic nonlinear responses,” *SciPost Physics Core* **5**, 039 (2022).
- [129] Tobias Holder, Daniel Kaplan, and Binghai Yan, “Consequences of time-reversal-symmetry breaking in the light-matter interaction: Berry curvature, quantum metric, and diabatic motion,” *Phys. Rev. Res.* **2**, 033100 (2020).
- [130] Hikaru Watanabe and Youichi Yanase, “Nonlinear electric transport in odd-parity magnetic multipole systems: Application to Mn-based compounds,” *Phys. Rev. Res.* **2**, 043081 (2020).
- [131] Hikaru Watanabe and Youichi Yanase, “Chiral Photocurrent in Parity-Violating Magnet and Enhanced Response in Topological Antiferromagnet,” *Phys. Rev. X* **11**, 011001 (2021).
- [132] J. Železný, Z. Fang, K. Olejník, J. Patches, F. Gerhard, C. Gould, L. W. Molenkamp, C. Gomez-Olivella, J. Zemen, T. Tichý, T. Jungwirth, and C. Ciccarelli, “Unidirectional magnetoresistance and spin-orbit torque in NiMnSb,” *Phys. Rev. B* **104**, 054429 (2021).
- [133] Alexander Cerjan, Sheng Huang, Mohan Wang, Kevin P Chen, Yidong Chong, and Mikael C Rechtsman, “Experimental realization of a Weyl exceptional ring,” *Nature Photonics* **13**, 623 (2019).
- [134] Tao Liu, James Jun He, Zhongmin Yang, and Franco Nori, “Higher-Order Weyl-Exceptional-Ring Semimetals,” *Phys. Rev. Lett.* **127**, 196801 (2021).
- [135] Bo Zhen, Chia Wei Hsu, Yuichi Igarashi, Ling Lu, Ido Kaminer, Adi Pick, Song-Liang Chua, John D Joannopoulos, and Marin Soljačić, “Spawning rings of exceptional points out of Dirac cones,” *Nature* **525**, 354–358 (2015).
- [136] Kohei Kawabata, Takumi Bessho, and Masatoshi Sato, “Classification of exceptional points and non-Hermitian topological semimetals,” *Phys. Rev. Lett.* **123**, 066405 (2019).
- [137] Kunkun Wang, Lei Xiao, Jan Carl Budich, Wei Yi, and Peng Xue, “Simulating Exceptional Non-Hermitian Metals with Single-Photon Interferometry,” *Phys. Rev.*

- Lett.* **127**, 026404 (2021).
- [138] Kun Ding, Guancong Ma, Meng Xiao, Z. Q. Zhang, and C. T. Chan, “Emergence, coalescence, and topological properties of multiple exceptional points and their experimental realization,” *Phys. Rev. X* **6**, 021007 (2016).
- [139] V. M. Martinez Alvarez, J. E. Barrios Vargas, and L. E. F. Foa Torres, “Non-Hermitian robust edge states in one dimension: Anomalous localization and eigenspace condensation at exceptional points,” *Phys. Rev. B* **97**, 121401 (2018).
- [140] Hengyun Zhou, Chao Peng, Yoseob Yoon, Chia Wei Hsu, Keith A Nelson, Liang Fu, John D Joannopoulos, Marin Soljačić, and Bo Zhen, “Observation of bulk Fermi arc and polarization half charge from paired exceptional points,” *Science* **359**, 1009–1012 (2018).
- [141] WD Heiss, “The physics of exceptional points,” *Journal of Physics A: Mathematical and Theoretical* **45**, 444016 (2012).
- [142] Lei Xiao, Tianshu Deng, Kunkun Wang, Zhong Wang, Wei Yi, and Peng Xue, “Observation of non-Bloch parity-time symmetry and exceptional points,” *Phys. Rev. Lett.* **126**, 230402 (2021).
- [143] Emil J. Bergholtz, Jan Carl Budich, and Flore K. Kunst, “Exceptional topology of non-Hermitian systems,” *Rev. Mod. Phys.* **93**, 015005 (2021).
- [144] Deyuan Zou, Tian Chen, Haiyu Meng, Yee Sin Ang, Xiangdong Zhang, and Ching Hua Lee, “Experimental observation of exceptional bound states in a classical circuit network,” *Science Bulletin* (2024), 10.1016/j.scib.2024.05.036.
- [145] Jan Wiersig, “Review of exceptional point-based sensors,” *Photonics Research* **8**, 1457–1467 (2020).
- [146] Midya Parto, Yuzhou GN Liu, Babak Bahari, Mercedeh Khajavikhan, and Demetrios N Christodoulides, “Non-Hermitian and topological photonics: optics at an exceptional point,” *Nanophotonics* **10**, 403–423 (2020).
- [147] Haiping Hu, Shikang Sun, and Shu Chen, “Knot topology of exceptional point and non-Hermitian no-go theorem,” *Phys. Rev. Res.* **4**, L022064 (2022).
- [148] Haiyu Meng, Lingling Wang, Ching Hua Lee, and Yee Sin Ang, “Terahertz polarization conversion from optical dichroism in a topological Dirac semimetal,” *Applied Physics Letters* **121** (2022), 10.1063/5.0122299.
- [149] Kun Ding, Chen Fang, and Guancong Ma, “Non-Hermitian topology and exceptional-point geometries,” *Nature Reviews Physics* **4**, 745–760 (2022).
- [150] Wenbo Mao, Zhoutian Fu, Yihang Li, Fu Li, and Lan Yang, “Exceptional-point-enhanced phase sensing,” *Science Advances* **10**, ead15037 (2024).
- [151] Haiyu Meng, Yee Sin Ang, and Ching Hua Lee, “Exceptional points in non-Hermitian systems: Applications and recent developments,” *Applied Physics Letters* **124** (2024), 10.1063/5.0183826.
- [152] Hua-Zhou Chen, Tuo Liu, Hong-Yi Luan, Rong-Juan Liu, Xing-Yuan Wang, Xue-Feng Zhu, Yuan-Bo Li, Zhong-Ming Gu, Shan-Jun Liang, He Gao, *et al.*, “Revealing the missing dimension at an exceptional point,” *Nature Physics* **16**, 571–578 (2020).
- [153] Vladyslav Kozii and Liang Fu, “Non-Hermitian topological theory of finite-lifetime quasiparticles: prediction of bulk Fermi arc due to exceptional point,” arXiv:1708.05841 (2017).
- [154] R. Arouca, C. H. Lee, and C. Morais Smith, “Unconventional scaling at non-Hermitian critical points,” *Phys. Rev. B* **102**, 245145 (2020).
- [155] Po-Yao Chang, Jhih-Shih You, Xueda Wen, and Shinsei Ryu, “Entanglement spectrum and entropy in topological non-Hermitian systems and nonunitary conformal field theory,” *Phys. Rev. Res.* **2**, 033069 (2020).
- [156] Sirui Liu, Hui Jiang, Wen-Tan Xue, Qingya Li, Jiangbin Gong, Xiaogang Liu, and Ching Hua Lee, “Non-Hermitian entanglement dip from scaling-induced exceptional criticality,” arXiv:2408.02736 (2024).
- [157] Jiaming Li, Andrew K Harter, Ji Liu, Leonardo de Melo, Yogesh N Joglekar, and Le Luo, “Observation of parity-time symmetry breaking transitions in a dissipative Floquet system of ultracold atoms,” *Nature Communications* **10**, 855 (2019).
- [158] Samantha Lapp, Fangzhao Alex An, Bryce Gadway, *et al.*, “Engineering tunable local loss in a synthetic lattice of momentum states,” *New Journal of Physics* **21**, 045006 (2019).
- [159] Zejian Ren, Dong Liu, Entong Zhao, Chengdong He, Ka Kwan Pak, Jensen Li, and Gyu-Boong Jo, “Chiral control of quantum states in non-Hermitian spin-orbit-coupled fermions,” *Nature Physics* **18**, 385 (2022).
- [160] Qian Liang, Dizhou Xie, Zhaoli Dong, Haowei Li, Hang Li, Bryce Gadway, Wei Yi, and Bo Yan, “Dynamic Signatures of Non-Hermitian Skin Effect and Topology in Ultracold Atoms,” *Physical Review Letters* **129**, 070401 (2022).
- [161] Wei Gou, Tao Chen, Dizhou Xie, Teng Xiao, Tian-Shu Deng, Bryce Gadway, Wei Yi, and Bo Yan, “Tunable nonreciprocal quantum transport through a dissipative Aharonov-Bohm ring in ultracold atoms,” *Physical Review Letters* **124**, 070402 (2020).
- [162] R. Y. Chen, Z. G. Chen, X.-Y. Song, J. A. Schneeloch, G. D. Gu, F. Wang, and N. L. Wang, “Magnetoinfrared Spectroscopy of Landau Levels and Zeeman Splitting of Three-Dimensional Massless Dirac Fermions in ZrTe₅,” *Phys. Rev. Lett.* **115**, 176404 (2015).
- [163] Wikipedia, “Berry connection and curvature,” *Wikipedia* (2023).