arXiv:2411.07027v1 [hep-ph] 11 Nov 2024 arXiv:2411.07027v1 [hep-ph] 11 Nov 2024

Functional renormalization group study of a four-fermion model with CP violation: implications to spontaneous CP violation models

Linlin Huang D[,](https://orcid.org/0000-0002-1013-8631)^{1,[∗](#page-0-0)} Mamiya Kawag[uch](https://orcid.org/0000-0001-9374-3716)i D,^{2,[†](#page-0-1)} Yadikaer Maitiniyazi D,^{1,[‡](#page-0-2)}

Shinya Matsuzaki \mathbf{D} ,^{1, [§](#page-0-3)} Akio Tomiya \mathbf{D} ,^{3, 4, [¶](#page-0-4)} and Masatoshi Yamada \mathbf{D} ^{1,**}

 1 ¹ Center for Theoretical Physics and College of Physics, Jilin University, Changchun 130012, China

² Anhui University of Science and Technology, Huainan, Anhui 232001, People's Republic of China

 3 Department of Mathematics, Tokyo Woman's Christian University, Tokyo 167-8585, Japan

⁴RIKEN Center for Computational Science, Kobe 650-0047, Japan

We work on the functional renormalization group analysis on a four-fermion model with the CP and P violation in light of nonperturbative exploration of the infrared dynamics of quantum chromodynamics (QCD) arising from the spontaneous CP violation models in a view of the Wilsonian renormalization group. The fixed point structure reveals that in the large- N_c limit, the CP $\bar{\theta}$ parameter is induced and approaches $\pi \cdot (N_f/2)$ (with the number of flavors N_f) toward the chiral broken phase due to the criticality and the large anomalous dimensions of the $U(1)$ axial violating four-fermion couplings. This trend seems to be intact even going beyond the large- N_c leading, as long as the infrared dynamics of QCD is governed by the scalar condensate of the quark bilinear as desired. This gives an impact on modeling of the spontaneous CP violation scenarios: the perturbatively irrelevant four-fermion interactions nonperturbatively get relevant in the chiral broken phase, implying that the neutron electric dipole moment becomes too big, unless cancellations due to extra CP and P violating contributions outside of QCD are present at a certain intermediate infrared scale.

I. INTRODUCTION

CP violation originated from the θ parameter in quantum chromodynamics (QCD) has currently been constrained by the neutron electric dipole moment (nEDM) measurements [\[1,](#page-15-0) [2\]](#page-15-1) to be vanishingly small, $\bar{\theta}$ < 10⁻¹¹. Here, $\theta = \theta - \arg \det(Y_u Y_d)$ is the physical CP violation angle with Y_u and Y_d the Yukawa matrices in the flavor basis. Although $\bar{\theta}$ (or the QCD θ parameter), in principle, can take an arbitrary value between 0 to 2π , its measured value is mysteriously quite small. This is the so-called strong CP problem which motivates us to introduce Beyond the Standard Model (BSM).

Models such as the Nelson-Barr (NB) model and the left-right symmetric model [\[3](#page-15-2)[–12\]](#page-15-3) based on spontaneous CP violation have been studied for solving the strong CP problem. In models of BSM physics with spontaneous CP violation, additional heavy quarks and heavy scalars are introduced. CP violation is initially forbidden by imposing parity (P) invariance, which sets the CP violating phases to zero at the level of the classical action. A heavy scalar field associated with a $U(1)$ axial symmetry (denoted as $U(1)_A$) is assumed to acquire a nonzero, complex vacuum expectation value, leading to spontaneous breaking of the $U(1)_A$ symmetry through quantum effects. This spontaneous breaking of $U(1)_A$ also induces spontaneous breaking of CP and P symmetries

in the model.

Integrating out the heavy scalar (at the perturbative loop level) radiatively generates CP- and P-violating effective operators including the QCD θ term and the weak CP-violation phase (i.e., the Kobayashi-Maskawa phase) $[3-12]$ $[3-12]$ $[3-12]$ ¹. The QCD θ parameter in the framework of the SM is renormalization group (RG) invariant due to its topological nature, while in the spontaneous CPviolation scenarios, a CP-violation phase could vary under RG transformations, associating with the RG evolution of the CP and P violating effective operators. The perturbative RG running of those CP and P violating operators has so far been clarified down til the QCD intrinsic scale (or the neutron mass scale) ~ 1 GeV based on the SM effective field theory approach [\[14–](#page-15-4)[17\]](#page-15-5).

The induced CP and P violating operators are typically higher-dimensional operators, e.g., four-fermion operators which are irrelevant in the perturbative arguments and therefore suppressed in the low energy dynamics. However, certain four-fermion operators play a crucial role in dynamical chiral symmetry breaking; see, e.g., Refs. [\[18–](#page-15-6)[20\]](#page-15-7). The effective couplings associated with these operators could be large, making these higher-dimensional terms relevant in low-energy regimes. In such cases of strong dynamics, the perturbative treatments fail, and nonperturbative techniques are necessary to accurately capture the RG evolution of CP-violation phases.

In this Letter, we make a first step based on the functional Renormalization Group (fRG) analysis on a four-

[∗] huangll22@mails.jlu.edu.cn

[†] mamiya@aust.edu.cn

[‡] ydqem22@mails.jlu.edu.cn

[§] synya@jlu.edu.cn

[¶] akio@yukawa.kyoto-u.ac.jp

^{∗∗} yamada@jlu.edu.cn

¹ The Vafa-Witten's theorem $[13]$ is inapplicable to the scenario of this type which leaves nonzero CP and P violation in the vacuum, because of the presence of extra CP- and P-violating interactions beyond the framework of QCD of the SM.

fermion model with CP (and P) violation and one quark flavor included. The model can capture the essential features of the low-energy description below the QCD intrinsic scale by integrating out QCD gluons from the spontaneous CP violation scenarios, in a view of the Wilsonian RG. The extension to two- or three-quark flavor cases is straightforwardly achievable, which is to deserve to another publication.

We discuss the fixed point structure and find that in the large- N_c limit, a finite value of $\bar{\theta}$ is induced and can approach $\pi/2$ toward the chiral broken phase due to the criticality and the large anomalous dimensions of the $U(1)_A$ -violating four-fermion couplings. The large- N_c subleading four-fermion term corrections seem to be suppressed enough when the model is referred to the underlying QCD theory, because those do not dominantly participate in the dynamical chiral symmetry breaking via the scalar condensate of the quark bilinear, which governs the infrared (IR) dynamics of QCD [\[21\]](#page-15-9). Thus the current conclusion would still be intact, and would phenomenologically make an impact on modeling of the spontaneous CP-violation scenarios: extra CP violating contributions at scales blow the QCD intrinsic scale from outside of the QCD sector would be necessary to be present to cancel $\bar{\theta} = \pi \cdot (N_f/2)$, otherwise the associated EDMs (the quark EDM, the chromo EDM, hence the nEDM) would all be dynamically amplified.

II. WILSONIAN RG VIEW OF SPONTANEOUS CP-VIOLATION SCENARIOS

We begin by addressing the low-energy description of the spontaneous CP-violation scenarios below the QCD intrinsic scale in a view of the Wilsonian RG. Just as a reference, we introduce an NB-like model which can share the universal low-energy features with other spontaneous CP-violation models. Instead of the original NB model [\[3–](#page-15-2) [7\]](#page-15-10) or the minimal models [\[22\]](#page-15-11), we consider a concise version with one flavor quark and one complex scalar, which still keeps the essential features of the NB-type model. The introduced relevant part of the Lagrangian is (see also Ref. $[17]$):

$$
-\mathcal{L}_{\text{NB-like}} = m_{\phi}^2 |\phi|^2 + y_q (\bar{q}_L q_R \phi + \bar{q}_R q_L \phi^*).
$$
 (1)

Here, ϕ is a complex scalar field charged under the P and $U(1)_A$ symmetry. This model is P and $U(1)_A$ invariant, hence at this moment the CP violating phases vanish. The scalar ϕ is assumed to develop nonzero complex vacuum expectation value, which spontaneously breaks the $U(1)_A$ symmetry as well as P.

In such a complex vacuum expectation value of ϕ , the P and/or $U(1)_A$ breaking effective operators are generated. Integrating ϕ out at the scalar mass scale $k = m_{\phi}$, the effective Lagrangian in the quark-gluon sector, up to mass dimension 6, takes the following form [\[14,](#page-15-4) [15,](#page-15-12) [17\]](#page-15-5):

$$
\mathcal{L}_{\text{eff}}(k \sim m_{\phi})
$$

$$
= \bar{q}i\mathcal{P}q - m_q(\bar{q}e^{i\gamma_5\theta_q/2}q + \text{h.c.}) - \frac{1}{2}G_{\mu\nu}G^{\mu\nu} + \theta_G \frac{g_s^2}{32\pi^2}G_{\mu\nu}\tilde{G}^{\mu\nu} + C_2^q g_s(\bar{q}\sigma_{\mu\nu}G^{\mu\nu}\gamma_5q) + \frac{1}{3}C_3 f^{abc}G^a_{\mu\nu}G^b_{\nu}\tilde{G}^{c\rho\mu} + C_4^q(\bar{q}q)(\bar{q}i\gamma_5q) + C_5^q(\bar{q}\sigma^{\mu\nu}q)(\bar{q}i\sigma_{\mu\nu}\gamma_5q) + \tilde{C}_2^{qq}\text{tr}(\bar{q}q\bar{q}i\gamma_5q) + \tilde{C}_4^{qq}\text{tr}(\bar{q}\sigma^{\mu\nu}q\bar{q}i\sigma_{\mu\nu}\gamma_5q) , \quad (2)
$$

where g_s stands for the QCD gauge coupling; $\sigma_{\mu\nu}$ = $\frac{i}{4}[\gamma_\mu,\gamma_\nu]$ is the Lorentz generators in spinor space with the Dirac's gamma matrices γ_{μ} ; $G_{\mu\nu} = G_{\mu\nu}^{a} T^{a}$ is the QCD gluon field strength with the gluon fields $A_{\mu} = A_{\mu}^{a}T^{a}$ along with the $SU(3)_c$ generators T^a ; f^{abc} denotes the structure constant of the $SU(3)_c$ group. The introduced coupling notation (in part) follows the literature [\[15\]](#page-15-12).

At the scales $k < m_{\phi}$, the induced CP and P violating operators (corresponding to terms except the minimal quark-gluon gauge coupling and the gluon kinetic term in the first line of Eq. (2)) logarithmically evolve in RG $[14, 15]$ $[14, 15]$ down til the QCD intrinsic scale, Λ_{QCD} . Those operators are mixed via the RG evolution [\[14,](#page-15-4) [15,](#page-15-12) [17\]](#page-15-5), hence all those CP and P violating contributions are potentially crucial.

By further integrating out the QCD gluon fields at $k \sim \Lambda_{\text{QCD}}$, in viewpoint of the Wilsonian RG, we would find the low-energy description of the NB-like model, which is given by the following local quark interaction terms up to dimension 6:

$$
\mathcal{L}_{\text{eff}}(k \lesssim \Lambda_{\text{QCD}}) = -m(\bar{q}e^{i\gamma_5 \theta/2}q + \text{h.c.}) \n+ \frac{G_S}{2}(\bar{q}q)^2 + \frac{G_P}{2}(\bar{q}i\gamma_5 q)^2 + \cdots \n+ C_4(\bar{q}q)(\bar{q}i\gamma_5 q) + C_5(\bar{q}\sigma^{\mu\nu}q)(\bar{q}i\sigma_{\mu\nu}\gamma_5 q) \n+ C'_2 \text{tr}(\bar{q}q\bar{q}i\gamma_5 q) + C'_4 \text{tr}(\bar{q}\sigma^{\mu\nu}q\bar{q}i\sigma_{\mu\nu}\gamma_5 q).
$$
\n(3)

Here and hereafter we omit the bar on θ . The G_S and G_P terms are P invariant, which are generated from the standard QCD gauge interaction among the quark and gluon (corresponding to the minimal coupling $\bar{q}Aq$ in the \mathcal{D} term of Eq.[\(2\)](#page-1-0)). They are also induced from the diagrams constructed from the even number of the P-violating nonminimal quark-gluon coupling (C_2^q) and the four-fermion couplings $(C_4^q, C_5^q, \widetilde{C}_2^{qq}, \widetilde{C}_4^{qq})$, involving the standard minimal quark-gluon coupling $(\bar{q}Aq)$. The discrepancy between G_S and G_P is generated from the $U(1)_A$ violation, including the underlying QCD instanton effect (i.e. the quantum anomaly), and the contributions from the $U(1)_A$ breaking operators with C_2^q , C_4^q , C_5^q , \widetilde{C}_2^{qq} , and \widetilde{C}_4^{qq} . The odd number of the P-violating nonminimal quark-gluon coupling (C_2^q) and the four-fermion couplings $(C_4^q, C_5^q, \tilde{C}_2^{qq}, \tilde{C}_4^{qq}),$ involving the standard minimal quarkgluon coupling $(\bar{q}Aq)$, induces the other P and $U(1)_A$ violating four-fermion interaction terms with the couplings $C_4, C_5, C'_2, \text{ and } C'_4.$

The ellipsis in the second line of Eq. [\(3\)](#page-1-1) abbreviates other P invariant four-fermion interaction terms such as

the forms of the vector $(\gamma_\mu \otimes \gamma^\mu)$, axialvector $(\gamma_\mu \gamma_5 \otimes$ $\gamma^{\mu}\gamma_{5})$, and tensor $(\sigma_{\mu\nu}\otimes\sigma^{\mu\nu})$ types, which can also in a similar way be induced from Eq. [\(2\)](#page-1-0). We will comment on potential contributions from those operators later, in the next section.

The full fRG analysis, which describes matching of the low-energy theory in Eq. [\(3\)](#page-1-1) with the ultraviolet (UV) theory including operators in Eq. (2) , can straightforwardly be done. In the present work, however, we shall focus particularly on the running of the CP and P violating couplings due to the nonperturbative four-fermion dynamics below the QCD intrinsic scale. The complete full fRG study will be pursued in another publication.

III. FOUR-FERMION MODEL WITH CP VIOLATION

Motivated by the NB-like models for the spontaneous CP-violation scenario, we consider an effective-theory approach. In this work, we study the following one-flavor $(N_f = 1)$ four-fermion theory as a toy model:

$$
S = \int d^4x \left[\bar{\psi} \left(i\gamma^{\mu} \partial_{\mu} - m e^{i\theta \gamma_5/2} \right) \psi + \frac{G_S}{2} \left(\bar{\psi} \psi \right)^2 + \frac{G_P}{2} \left(\bar{\psi} i \gamma_5 \psi \right)^2 + C_4 \left(\bar{\psi} \psi \right) \left(\bar{\psi} i \gamma_5 \psi \right) \right]. \tag{4}
$$

Here ψ and $\bar{\psi}$ are fermion fields with degrees of freedom of $SU(N_c)$ color symmetry. The first two four-fermion operators in the second line of Eq. [\(4\)](#page-2-0) are invariant under the P transformation, while the last one breaks the P invariance and consequently the CP invariance is broken.^{[2](#page-2-1)} Thus, when $C_4 = 0$, the theory is CP invariant. Besides, for $G_S = G_P$ and $C_4 = 0$, the $U(1)_A$ invariance is restored. In such a case, the phase θ becomes unphysical because it is erased by the $U(1)_A$ rotation of the fermion fields.

In comparison with Eq. (3) , the colored four-fermion interaction terms $(C'_2$ and C'_4 terms in the last line) have been discarded in Eq. [\(4\)](#page-2-0). This is simply because those color-nonsinglet channels will be irrelevant when the confinement phase of underlying QCD, or the bosonization of the color singlet fermion-bilinear fields is taken into account.

When the model is extended to multiple flavors cases (e.g., two, or three flavors), the global non-Abelian chiral symmetry can be reflected individually in the four-fermion interactions in Eq. [\(4\)](#page-2-0) in a way separated from the

$$
\begin{aligned}\n\bar{\psi}\psi \xrightarrow{\mathbf{P}} \bar{\psi}\psi, &\qquad \bar{\psi}i\gamma^5\psi \xrightarrow{\mathbf{P}} -\bar{\psi}i\gamma^5\psi, \\
\bar{\psi}\psi \xrightarrow{\mathbf{G}} \bar{\psi}\psi, &\qquad \bar{\psi}i\gamma^5\psi \xrightarrow{\mathbf{G}} \bar{\psi}i\gamma^5\psi.\n\end{aligned}
$$

Abelian $U(1)_A$ symmetry which is explicitly broken there (due to $G_S - G_P \neq 0$ and $C_4 \neq 0$): in the two-flavor case, say, the G_S and G_P terms can be extended like $G_{S}\left[(\bar{\psi}\psi)^{2}+(\bar{\psi}i\gamma_{5}\tau^{a}\psi)^{2}\right]^{*}+\left. G_{P}\left[(\bar{\psi}\tau^{a}\psi)^{2}+(\bar{\psi}i\gamma_{5}\psi)^{2}\right]\right.$ with the $SU(2)$ -doublet fermion field ψ and the Pauli matrices τ^a ($a = 1, 2, 3$), which is individually chiral $SU(2)$ invariant for each of the G_S and G_P terms, but breaks the $U(1)_A$ symmetry when $G_S \neq G_P$. In that sense, the discrepancy between G_S and G_P dictates the $U(1)_A$ anomaly of underlying QCD. The dynamical breaking of the chiral $SU(2)$ symmetry is triggered even solely through the strong G_S and/or G_P couplings like in the Nambu-Jona-Lasinio dynamics, not essentially due to the $U(1)_A$ anomaly $(G_S \neq G_P)$. Therefore, we shall generically define the chiral broken phase the case when G_S and/or G_P reach the criticality, irrespective to the presence of the $U(1)_A$ anomaly controlled by the condition $G_S \neq G_P$.

A. Functional Renormalization Group

To study the four-fermion model in the presence of the mass term with CP phase (4) , we employ the fRG which is the formulation of the Wilsonian RG in realm of quantum field theory. The Wilsonian coarse-graining process is described by the functional differential equation for an effective action. In this work, we utilize the Wetterich equation [\[23\]](#page-15-13) in which the central object is the cutoff-dependent one-particle irreducible effective action Γ_k where k is the IR cutoff scale. See Refs. [\[24–](#page-15-14)[37\]](#page-16-0) for a review on the fRG. The explicit form of the Wetterich equation (or flow equation) for a fermionic theory is given by

$$
\partial_t \Gamma_k = -\text{Tr}\left[\left(\Gamma_k^{(2)}(p) + \mathcal{R}_k(p) \right)^{-1} \partial_t \mathcal{R}_k(p) \right], \quad (5)
$$

where $\partial_t = k \partial_k$ is the derivative with respect to the dimensionless scale $t = \log(k/\Lambda)$ with a UV cutoff scale Λ. For this convention, the IR limit (k → 0) corresponds to $t \to -\infty$. In Eq. [\(5\)](#page-2-2),

$$
\Gamma_k^{(2)}(p) = \frac{\overrightarrow{\delta}}{\delta \overline{\psi}(p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \psi(p)} \tag{6}
$$

is the full two-point function (inverse full-propagator) for the fermion fields ψ and $\bar{\psi}$ in momentum space, "Tr" denotes the trace acting on all spaces, e.g., momentum, color, flavor etc., on which the fields ψ and ψ are defined. $\mathcal{R}_k(p)$ is the regulator function realizing the Wilsonian coarse-graining process in the path-integral formalism. In this work, we employ the Litim-type cutoff function [\[38\]](#page-16-1) for the fermion fields, i.e.,

$$
\mathcal{R}_k(p) = i\rlap/v \left(\sqrt{\frac{k^2}{p^2}} - 1 \right) \Theta(k^2 - p^2) \,, \tag{7}
$$

² Note here that the transformation laws under the P and C discrete symmetries are

where $\Theta(x)$ is the step function: $\Theta(x) = 1$ for $x > 0$, while $\Theta(x) = 0$ for $x < 0$. See Refs. [\[39](#page-16-2)[–42\]](#page-16-3) especially for the treatment of fermionic theories in the fRG.

By solving the Wetterich equation [\(5\)](#page-2-2) with the initial condition $\Gamma_{\Lambda} = S$ in Eq. [\(4\)](#page-2-0) at $k = \Lambda$, we obtain the full effective action $\Gamma = \Gamma_{k=0}$ at $k=0$. However, in the general Wilsonian viewpoint, Γ_k is expressed as an infinite series of effective operators, making it infeasible to handle the effective action without approximations. In this work, we make the following ansatz for the effective action for the model [\(4\)](#page-2-0) in Euclidean spacetimes:

$$
\Gamma_k \simeq \int d^4x \left[\bar{\psi} \left(\gamma^{\mu} \partial_{\mu} + m e^{i\theta \gamma_5/2} \right) \psi - \frac{G_S}{2} \left(\bar{\psi} \psi \right)^2 - \frac{G_P}{2} \left(\bar{\psi} i \gamma_5 \psi \right)^2 - C_4 \left(\bar{\psi} \psi \right) \left(\bar{\psi} i \gamma_5 \psi \right) \right].
$$
 (8)

Our convention for the Euclidean signature is summarized in Appendix [A.](#page-7-0) The couplings $(m, \theta, G_S, G_P \text{ and } C_4)$ are k-dependent parameters. Higher dimensional operators tend to be irrelevant in the IR regime due to their larger canonical dimensions, so that we can ignore them. However, four-fermion operators such as $(\bar{\psi}\gamma_{\mu}\psi)^{2}$ and $(\bar{\psi}\gamma_\mu\gamma^5\psi)^2$ as well as $(\bar{\psi}\sigma_{\mu\nu}\psi)^2$ may give potentially the same order contributions as of the four-fermion operators in the effective action [\(8\)](#page-3-0). Thus, the Fierz-complete basis should be considered; see Refs. [\[43–](#page-16-4)[46\]](#page-16-5). Nonetheless, these operators tend to be irrelevant even at a nontrivial fixed point [\[47\]](#page-16-6), so that we do not take into account these four-fermion operators in this work. In the current fermionic theory, the kinetic term does not receive quantum effects from the four-fermion operators thanks to the one-loop structure of the flow equation [\(5\)](#page-2-2).

B. Flow equations

Applying the Wetterich equation [\(5\)](#page-2-2) for the effective action (8) , we derive the flow equations for the five couplings. We list the flow equations for couplings in the effective action (8) within the large- N_c approximation. The detailed derivation is presented in Appendix [B.](#page-9-0) Here, in order to discuss the fixed point structure and the flow diagrams in the model in the following subsections, we define the dimensionless couplings as

$$
\tilde{m} = mk^{-1}, \qquad \tilde{G}_S = G_S k^2,
$$

\n
$$
\tilde{G}_P = G_P k^2, \qquad \tilde{C}_4 = C_4 k^2,
$$
\n(9)

while θ is already dimensionless.

In this work, we aim to discuss the basic properties of our model. Although the fRG formalism allows us to take into account the large- N_c subleading effects, it is expected that the main contributions come from the large- N_c leading parts. Thus, we study the flow equations within the large- N_c approximation in which the flow equations are obtained as

$$
\partial_t \tilde{m} = -\tilde{m} - 4N_c \left[(\tilde{G}_S + \tilde{G}_P) + 2\tilde{C}_4 \sin \theta + (\tilde{G}_S - \tilde{G}_P) \cos \theta \right] \tilde{m} \tilde{\mathcal{M}}, \tag{10}
$$

$$
\partial_t \theta = 8 \left[(\tilde{G}_S - \tilde{G}_P) \sin \theta - 2 \tilde{C}_4 \cos \theta \right] \tilde{\mathcal{M}}, \tag{11}
$$

$$
\partial_t \tilde{G}_S = 2\tilde{G}_S - 8N_c \Big[(\tilde{G}_S^2 + \tilde{C}_4^2)(1 - \tilde{m}^2) + 2\tilde{m}^2(\tilde{G}_S^2 - \tilde{C}_4^2) \cos\theta + 4\tilde{m}^2 \tilde{G}_S \tilde{C}_4 \sin\theta \Big] \tilde{\mathcal{I}},\tag{12}
$$

$$
\partial_t \tilde{G}_P = 2\tilde{G}_P - 8N_c \Big[(\tilde{G}_P^2 + \tilde{C}_4^2)(1 - \tilde{m}^2) - 2\tilde{m}^2(\tilde{G}_P^2 - \tilde{C}_4^2) \cos \theta + 4\tilde{m}^2 \tilde{G}_P \tilde{C}_4 \sin \theta \Big] \tilde{\mathcal{I}},\tag{13}
$$

$$
\partial_t \tilde{C}_4 = 2\tilde{C}_4 - 8N_c \Big[(\tilde{G}_S + \tilde{G}_P)\tilde{C}_4(1 - \tilde{m}^2) + 2\tilde{m}^2(\tilde{G}_S - \tilde{G}_P)\tilde{C}_4\cos\theta + 2\tilde{m}^2(\tilde{G}_S\tilde{G}_P + \tilde{C}_4^2)\sin\theta \Big] \tilde{\mathcal{I}}.
$$
 (14)

Here we have defined the (dimensionless) threshold functions as

$$
\widetilde{\mathcal{M}} = \frac{1}{k^2} \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^2} \n= \frac{1}{2(4\pi)^2} \frac{1}{(1 + \tilde{m}^2)^2},
$$
\n(15)

$$
\widetilde{\mathcal{I}} = \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^3}
$$

$$
=\frac{1}{2(4\pi)^2}\frac{1}{(1+\tilde{m}^2)^3},\tag{16}
$$

where in the second equality we have used the Litim-type cutoff function [\(7\)](#page-2-3).

The beta function for the mass parameter is proportional to the mass parameter itself, which reflects the concept of technical naturalness [\[48,](#page-16-7) [49\]](#page-16-8). In the case where both P and $U(1)_A$ symmetries are preserved, i.e., when $\tilde{G}_S = \tilde{G}_P$ and $\tilde{C}_4 = 0$, the beta function for θ vanishes, making θ invariant under renormalization group

(RG) transformations. Since the couplings \tilde{G}_S and \tilde{G}_P are P-invariant, their beta functions can depend on G_S^2 and G_P^2 . In contrast, \tilde{C}_4 breaks P symmetry, so its beta function must vary under P transformations and therefore cannot contain a \tilde{C}_4^2 term without another P breaking

C. Fixed point structure

effect, e.g., $\sin \theta \neq 0$.

In this subsection, we study the fixed-point structure. Thus, in the system, we solve $\partial_t \tilde{m} = \partial_t \theta = \partial_t \tilde{G}_S =$ $\partial_t \tilde{G}_P = \partial_t \tilde{C}_4 = 0$ and find their solutions. We start with the flow equation for θ . The fixed-point condition $\partial_t \theta = 0$ yields a relation as

$$
\tan \theta^* = \frac{2\tilde{C}_4^*}{\tilde{C}_S^* - \tilde{C}_P^*}.
$$
\n(17)

Inserting this into the other flow equations, the following nontrivial fixed points are found:

(FP1):
$$
\tilde{m}^* = 0
$$
, $\tilde{G}_S^* = \frac{8\pi^2}{N_c}$, $\tilde{G}_P^* = 0$, $\tilde{C}_4^* = 0$, (18)

(FP2):
$$
\tilde{m}^* = 0
$$
, $\tilde{G}_S^* = 0$, $\tilde{G}_P^* = \frac{8\pi^2}{N_c}$, $\tilde{C}_4^* = 0$, (19)

(FP3):
$$
\tilde{m}^* = 0
$$
, $\tilde{G}_S^* = G_P^* = \frac{8\pi^2}{N_c}$, $\tilde{C}_4^* = 0$. (20)

In addition to these nontrivial fixed points, there is the Gaussian (trivial) fixed point $\tilde{m}^* = \tilde{G}^*_{S} = \tilde{G}^*_{P} = \tilde{C}^*_{4} = 0.$ FP1 and FP2 are $U(1)_A$ -broken fixed points, while FP3 is $U(1)_A$ -symmetric one. For FP1 and FP2, from the fixed point condition [\(17\)](#page-4-0), $\tan \theta^* = 0$, i.e. $\theta^* = n\pi$ with n integers. In the case of FP1, the right-hand side of Eq. [\(17\)](#page-4-0) becomes an indeterminate form. This fact is consistent with the unphysical nature of θ when the system becomes $U(1)_{A}$ invariant at FP3.

Next, we analyze the critical exponents (scaling dimensions). To this end, we linearize the flow equations. More specifically, for a set of the flow equations $\partial_t \tilde{g}_i = \beta_i(\{\tilde{g}\})$ where $\{\tilde{g}\} = \{\tilde{m}, \theta, \tilde{G}_S, \tilde{G}_P, \tilde{C}_4\}$, we perform a Taylor expansion around a fixed point g_i^* and take up to the linear order of $\tilde{g} - \tilde{g}^*$ such that

$$
\partial_t \tilde{g}_i \simeq \frac{\partial \beta_i(\{\tilde{g}\})}{\partial \tilde{g}_j} \bigg|_{\tilde{g} = \tilde{g}^*} (\tilde{g}_j - \tilde{g}^*_j) \equiv -\mathcal{T}_{ij}(\tilde{g}_j - \tilde{g}^*_j), \quad (21)
$$

where $\beta({g^*}) = 0$ by definition of the fixed point. The solution to Eq. (21) is given by

$$
\tilde{g}_i = g_i^* + \sum_j c_j V_i^j \left(\frac{k}{\Lambda}\right)^{-\vartheta_j},\tag{22}
$$

where c_i are undetermined constants of integration, Λ is a reference scale and V_i^j are the right eigenvectors

of the stability matrix $\mathcal T$ with eigenvalues ϑ_i called the critical exponents or the scaling dimensions, namely, $\vartheta_i =$ $-eig(\mathcal{T})$. When we approximate eigenvectors as $V_i^j \approx \delta_i^j$, the critical exponents are approximately obtained from the diagonal parts of $\mathcal T$. Since at all the fixed points we find $\tilde{m}^* = \tilde{C}_4^* = 0$, such an approximation is reasonable and thus we have

$$
\vartheta_m = -\frac{\partial \beta_m}{\partial \tilde{m}}\Big|_{\tilde{g} = \tilde{g}^*}
$$

= $1 - \frac{N_c}{8\pi^2} \left((\tilde{G}_S^* + \tilde{G}_P^*) - (\tilde{G}_S^* - \tilde{G}_P^*) \cos \theta^* \right), (23)$

$$
\vartheta_{C_4} = -\frac{\partial \beta_{C4}}{\partial \tilde{C}_4} \bigg|_{\tilde{g} = \tilde{g}^*} = -2 + \frac{N_c}{4\pi^2} \left(\tilde{G}_S^* + \tilde{G}_P^* \right), \tag{24}
$$

$$
\vartheta_{\theta} = -\frac{\partial \beta_{\theta}}{\partial \theta} \bigg|_{\tilde{g} = \tilde{g}^*} = -\frac{N_c}{4\pi^2} \left(\tilde{G}_S^* - \tilde{G}_P^* \right) \cos \theta^*.
$$
 (25)

We exhibit the critical exponents at the Gaussian fixed point and FP3 for which the CP phase θ is unphysical. At the Gaussian fixed point, the critical exponents are identical with the canonical dimensions of couplings, i.e., we have $V_i^j = \delta_i^j$ and

$$
\vartheta_m = 1, \quad \vartheta_{G_S} = -2, \quad \vartheta_{G_P} = -2, \quad \vartheta_{C_4} = -2, \quad (26)
$$

while at FP3, we find

$$
\vartheta_m = 3, \quad \vartheta_{G_S} = 2, \quad \vartheta_{G_P} = 2, \quad \vartheta_{C_4} = 2.
$$
 (27)

All four-fermion couplings becomes relevant at FP3. This implies that a large anomalous dimension is induced by quantum effects which cannot be captured by the perturbative treatment. In particular, an important fact here is that \tilde{C}_4 becomes relevant at FP3 even if its fixed point value is zero. We will see in the next subsection that \tilde{C}_4 is driven by \tilde{G}_S and \tilde{G}_P and grows in the IR regime.

At the nontrivial fixed points (18) and (19) , we obtain eigenvalues such that we obtain, at FP1

$$
\vartheta_m = 3, \quad \vartheta_{\theta} = -2, \quad \vartheta_{G_S} = 2, \quad \vartheta_{G_P} = -2, \quad \vartheta_{C_4} = 0,
$$
\n(28)

and at FP2

$$
\vartheta_m = 1, \quad \vartheta_{\theta} = 2, \quad \vartheta_{G_S} = -2, \quad \vartheta_{G_P} = 2, \quad \vartheta_{C_4} = 0.
$$
\n(29)

Since the beta functions for \tilde{m} and θ depend on $\tilde{G}_S - \tilde{G}_P$, their critical exponents are also highly dependent on the differences of the fixed point values between \tilde{G}_{S}^{*} and \tilde{G}_{P}^{*} . As aforementioned, in QCD, the $U(1)_{A}$ symmetry is broken by the axial $U(1)$ anomaly and thus $\tilde{G}_S \neq \tilde{G}_P$. When a QCD hadron model with the lightest two quark flavors, which has a similar chiral and $U(1)_A$ breaking setup, is referred to (say, [\[50](#page-16-9)[–53\]](#page-16-10)), $G_S > G_P$ is ensured due to the positiveness of the η meson mass square. In this sense, FP1 (18) may be suitable as a low-energy effective field theory of QCD.

FIG. 1. Flow diagrams $(N_c = 3)$ on $\tilde{G}_S - \tilde{G}_P$ plane with $\tilde{C}_4 = 0$ (left) and $\tilde{G}_S - \tilde{C}_4$ plane with $\tilde{G}_P = 0$ (right) for $\tilde{m} = 0$. The red and purple points denotes the Gaussian fixed point and FP3 [\(20\)](#page-4-4), respectively, while blue points show FP1 [\(18\)](#page-4-2) and FP2 [\(18\)](#page-4-2). The dashed lines corresponds to the critical surface.

D. RG flow of parameters

We present the behavior of RG flows as solutions to Eqs. [\(10\)](#page-3-1)–[\(14\)](#page-3-2) with $N_c = 3$. To begin with, let us examine the case $\tilde{m} = 0$, in which the flow equations for the fourfermion couplings form a closed system. In Fig. [1,](#page-5-0) we display flow diagrams on the $\tilde{G}_S - \tilde{G}_P$ plane (with $\tilde{C}_4 = 0$) and on the $\tilde{G}_S - \tilde{C}_4$ plane (with $\tilde{G}_P = 0$). The arrows indicate the direction of RG flows from the UV to the IR. At the fixed point FP3, both \tilde{G}_S and \tilde{G}_P become relevant, meaning they are free parameters in this model. For initial values of \tilde{G}_S and \tilde{G}_P that exceed the fixedpoint values G_S^* and G_P^* , the RG flows diverge in the IR regime, suggesting dynamical breaking of the discrete axial symmetry; see, e.g., Refs. [\[18](#page-15-6)[–20\]](#page-15-7). Interestingly, the right-hand panel of Fig. [1](#page-5-0) shows that in this broken phase, \tilde{C}_4 tends to grow in the IR direction. This growth is essential for generating a finite value of θ . Thus, even if $\theta = 0$ at an initial scale $t = 0$ $(k = \Lambda)$, a finite value of \tilde{C}_4 can drive θ to become finite at IR scales. In Fig. [2,](#page-6-0) as an example, we present an RG flow solution to Eqs. (10) – (14) . with the initial conditions $\tilde{m} = 0.001, \ \theta = 0, \ \tilde{G}_S = 26.3,$ $\tilde{G}_P = 20$, and $\tilde{C}_4 = 1$ at $t = 0$. The RG flow of \tilde{m} does not change from the initial value, so we do not show it.

Here, we explore the possibilities of achieving maximum P/CP violation, specifically, $\theta = \pi/2$ in the IR regime in the current model. To see this, we set $\tilde{m} = 0$ and $\tilde{G}_P = 0$ for simplicity. Furthermore, we define $\tilde{m}_{\sigma}^2 = 1/\tilde{G}_S$ and $\rho = \tilde{C}_4/\tilde{G}_S$ whose flow equations read

$$
\partial_t \tilde{m}^2_{\sigma} = -2\tilde{m}^2_{\sigma} + 8N_c(1+\rho^2)\tilde{\mathcal{I}},\tag{30}
$$

$$
\partial_t \rho = 8N_c \frac{\rho^2}{\tilde{m}_\sigma^2} \tilde{\mathcal{I}}.\tag{31}
$$

Note here that \tilde{m}^2_{σ} corresponds to the mass parameter of an emergent bosonic field (corresponding to sigma meson) in the bosonization language [\[54,](#page-16-11) [55\]](#page-16-12). Thus, $m_{\sigma}^2 = 0$ corresponds to the critical value $\tilde{G}_S \to \infty$ at which the curvature of the effective potential for the bosonic field at the origin becomes zero. In the left-hand side panel of Fig. [3,](#page-6-1) we show the RG flow as a solution of Eqs. [\(30\)](#page-5-1) and [\(31\)](#page-5-2) with the initial condition $\tilde{m}_{\sigma}^2 = 3/8\pi^2$ and $\rho = 0.05$. In terms of the numerical treatment, the flow equation [\(30\)](#page-5-1) is convenient. We can solve the flow equation for \tilde{m}^2_{σ} beyond the critical scale, while the RG flow of ρ stops at $\tilde{m}^2_{\sigma} = 0$. Therefore, we cannot go beyond the critical scale in the current setup.

In Fig. [3,](#page-6-1) we depict an example of the RG flow of \tilde{m}_{σ}^2 , ρ and θ with the initial condition $\tilde{m}_{\sigma}^2 = 3/8\pi^2$, $\rho = 0.05$ and $\theta = 0$ at $t = 0$. The RG flow of \tilde{m}_{σ}^2 reaches zero around $t \approx 3$ which corresponds to the critical scale. Note here that ρ is a dimensionless parameter, so it tends to be a logarithmic running and does not change drastically under the RG evolution.

We see from the right-hand side panel of Fig. [3](#page-6-1) that a finite value of θ with the solution for \tilde{m}^2_{σ} and ρ is induced and reaches $\pi/2$. Once θ reaches $\pi/2$, the effect of \tilde{C}_4 in the flow equation [\(11\)](#page-3-3) for θ is switched off because of $\cos(\pi/2) = 0$. This fact is important to ensure that θ does not exceed $\pi/2$ and thus we can argue the P and CP violation within $0 \le \theta \le \pi/2$ in case of $N_f = 1$. In other words, the RG flow of θ does not behave as a limit cycle.

FIG. 2. RG flow as a solution to Eqs. [\(10\)](#page-3-1)–[\(14\)](#page-3-2) with $N_c = 3$ and the initial condition $\tilde{m} = 0.001$, $\theta = 0$, $\tilde{G}_S = 26.3$, $G_P = 20$ and $\tilde{C}_4 = 1$ at $t = 0$.

FIG. 3. RG flow as a solution to Eqs. [\(30\)](#page-5-1) and [\(31\)](#page-5-2) with $N_c = 3$ and the initial condition $\tilde{m}_{\sigma} = 3/8\pi^2$ and $\rho = 0.05$ at $t = 0$. The gray dashed line in the right-hand side panel indicates the value of $\pi/2$.

IV. SUMMARY

To summarize, the perturbatively irrelevant CP (and P) violating four-quark operators can be relevant due to the nonperturbatively yielded large anomalous dimensions triggered by the criticality of the dynamical chiral symmetry breaking. Consequently, the strong CP violation can be amplified in the chiral broken phase with the presence of the $U(1)_A$ anomaly and the CP phase approaches $\pi \cdot (N_f/2)$, even if it is tuned to vanish in the (seemingly) perturbative regime of QCD at the scales ≥ 1 GeV. This trend has been clarified in the large N_c limit by working on a four-fermion model, however, the $1/N_c$ subleading four-fermion term corrections seem to be suppressed enough, because those do not dominantly participate, in the dynamical chiral symmetry breaking via the scalar condensate of the quark bilinear, which governs the IR dynamics of QCD. Thus the current conclusion would still be intact.

Our findings would phenomenologically make an impact on modeling of the spontaneous CP violation scenarios: extra CP violating contributions at scales blow the QCD

intrinsic scale from outside of the QCD sector would be necessary to be present to cancel $\bar{\theta} = \pi \cdot (N_f/2)$, otherwise the associated EDMs (the quark EDM, the chromo EDM, hence the nEDM) would all be dynamically amplified.

In this work, we have investigated the RG flow of θ til the critical scale of dynamical chiral symmetry breaking. The further IR property of strong CP violation is, however, of importance. The measurement of the nEDM is indeed physics with the photon transfer energy (the applied electric field strength) $\sim 10^{-6} - 10^{-5}$ GeV [\[1,](#page-15-0) [2\]](#page-15-1), which is much less than the typical QCD scale $\sim 0.1-1$ GeV. The state-of-the-art theoretical estimates based on the lattice QCD at the physical point [\[56\]](#page-16-13) have been done for the transfer momentum in a range $\sim 200-400$ MeV, in which, however, the measurement is irrespective to $\bar{\theta}$, hence does not address its transfer momentum dependence, i.e., the RG scale. In order to follow the RG flow below the scales of the dynamical chiral symmetry breaking within the framework of the fRG, we may need to employ some techniques such as (dynamical) bosonization [\[57](#page-16-14)[–63\]](#page-17-0) and the weak formulation [\[21,](#page-15-9) [64\]](#page-17-1). These approaches to the CP-violating four-fermion model will be presented as our

future works.

ACKNOWLEDGMENTS

We thank Hiroyuki Ishida for useful comments on the NB scenario. The work of S. M. was supported in part by the National Science Foundation of China (NSFC) under Grant No.11747308, 11975108, 12047569, and the Seeds Funding of Jilin University. The work of A. T. was partially supported by JSPS KAKENHI Grant Numbers 20K14479, 22K03539, 22H05112, and 22H05111, and MEXT as "Program for Promoting Researches on the Supercomputer Fugaku" (Simulation for basic science: approaching the new quantum era; Grant Number JP-MXP1020230411, and Search for physics beyond the standard model using large-scale lattice QCD simulation and development of AI technology toward next-generation lattice QCD; Grant Number JPMXP1020230409). The work of M. Y. is supported by the National Science Foundation of China (NSFC) under Grant No. 12205116 and the Seeds Funding of Jilin University.

Appendix A: Eulidean spacetime

In this appendix, we provide our convention for the conversion from the Minkowski signature to the Euclidean one. The world line in Minkowski spacetime is given by

$$
ds^2 = dt^2 - d\mathbf{x}^2 = g_{\mu\nu}dx^{\mu}dx^{\nu},\tag{A1}
$$

where the metric is defined as $g_{\mu\nu} = g^{\mu\nu} = \text{diag}(1, -1, -1, -1)$. Now, we introduce the coordinate vectors $x_{E\mu}$ in Euclidean spacetime as

$$
t = x^0 = -ix_{E0} = -i\tau \,, \tag{A2}
$$

Then, the norm becomes

$$
x_{\mu}x^{\mu} = (x^{0})^{2} - (x^{i})^{2} = (-ix_{E0})^{2} - (x_{Ei})^{2} = -(x_{E0}^{2} + x_{Ei}^{2}) = -x_{E\mu}x_{E\mu}.
$$
 (A3)

In Euclidean spacetime, we do not have to take care about covariance and contravariance of vectors. Below, we summarize quantities in Euclidean spacetime

a) Vector $V_{\mu} = \{x_{\mu}, p_{\mu}, \dots\}$ $V^0 = -iV_{E0}$, $V^i = V_{Ei}$. (A4)

Then we have

$$
V_{\mu}V^{\mu} = -V_{E\mu}V_{E\mu}.
$$
\n(A5)

b) Derivative operator ∂_{μ}

$$
\partial_{\mu} = \left(\frac{\partial}{\partial x^{0}}, \frac{\partial}{\partial x^{i}}\right) = \left(\frac{\partial}{\partial (-ix_{E0})}, \frac{\partial}{\partial x_{Ei}}\right) = \left(i\frac{\partial}{\partial x_{E0}}, \frac{\partial}{\partial x_{Ei}}\right) = (i\partial_{E0}, \partial_{Ei}).
$$
\n(A6)

c) Gamma matrices γ_μ

$$
\gamma^0 = \gamma_{E0} , \qquad \gamma^i = i \gamma_{Ei} . \tag{A7}
$$

Then, the Clifford algebra $\{\gamma^{\mu}, \gamma^{\nu}\} = 2g^{\mu\nu}$ in Minkowski spacetime becomes

$$
\{\gamma^0, \gamma^0\} = \{\gamma_{E0}, \gamma_{E0}\} = 2, \qquad \{\gamma^i, \gamma^j\} = \{i\gamma_{Ei}, i\gamma_{Ej}\} = -\{\gamma_{Ei}, \gamma_{Ej}\} = -2\delta_{ij}.
$$
 (A8)

Thus, the Clifford algebra in Euclidean spacetime reads

$$
\{\gamma_{E\mu}, \gamma_{E\nu}\} = 2\delta_{\mu\nu} = 2\begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix} . \tag{A9}
$$

The hermitity of gamma matrices is

$$
(\gamma^0)^\dagger = \gamma^0 \Longrightarrow (\gamma_{E0})^\dagger = \gamma_{E0} , \qquad (\gamma^i)^\dagger = -\gamma^i \Longrightarrow (\gamma_{Ei})^\dagger = \gamma_{Ei} . \tag{A10}
$$

Thus, we have

$$
(\gamma_{E\mu})^{\dagger} = \gamma_{E\mu} \,. \tag{A11}
$$

The chirality matrix γ^5 is

$$
\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = i\gamma_{E0} i\gamma_{E1} i\gamma_{E2} i\gamma_{E3} = \gamma_{E0} \gamma_{E1} \gamma_{E2} \gamma_{E3} = \gamma_{E5}.
$$
 (A12)

d) Fields

– Scalar

$$
\phi(x) = \phi_E(x_E) \tag{A13}
$$

– Spinor

$$
\psi(x) = \psi_E(x_E), \qquad \bar{\psi}(x) = \psi^{\dagger} \gamma^0(x) = \psi_E^{\dagger} \gamma_{E0}(x_E) = \bar{\psi}_E(x_E). \tag{A14}
$$

– Vector

$$
A^{0}(x) = -iA_{E0}(x_{E}), \qquad A^{i}(x) = A_{Ei}(x_{E}). \qquad (A15)
$$

Note that $\bar{\psi}\gamma^{\mu}\psi$ is a vector; however, because of Eq. [\(A7\)](#page-7-1), we have

$$
\bar{\psi}\gamma^0\psi = \bar{\psi}_E\gamma_{E0}\psi_E, \qquad \bar{\psi}\gamma^i\psi = i\bar{\psi}_E\gamma_{Ei}\psi_E, \qquad (A16)
$$

which is different from Eqs. [\(A4\)](#page-7-2) and [\(A15\)](#page-8-0). The vector-type four-Fermion interaction becomes

$$
(\bar{\psi}\gamma^{\mu}\psi)^{2} = (\bar{\psi}\gamma^{0}\psi)(\bar{\psi}\gamma^{0}\psi) - (\bar{\psi}\gamma^{i}\psi)(\bar{\psi}\gamma^{i}\psi)
$$

\n
$$
= (\bar{\psi}_{E}\gamma_{E0}\psi_{E})(\bar{\psi}_{E}\gamma_{E0}\psi_{E}) - (\bar{\psi}_{E}i\gamma_{Ei}\psi_{E})(\bar{\psi}_{E}i\gamma_{Ei}\psi_{E})
$$

\n
$$
= (\bar{\psi}_{E}\gamma_{E0}\psi_{E})(\bar{\psi}_{E}\gamma_{E0}\psi_{E}) + (\bar{\psi}_{E}\gamma_{Ei}\psi_{E})(\bar{\psi}_{E}\gamma_{Ei}\psi_{E})
$$

\n
$$
= (\bar{\psi}_{E}\gamma_{E\mu}\psi_{E})^{2}.
$$
\n(A17)

e) Fourier transformations

$$
\phi_E(x_E) = \int \frac{\mathrm{d}^4 p_E}{(2\pi)^4} e^{ip_E \cdot x_E} \tilde{\phi}_E(p_E) , \qquad (A18)
$$

$$
\psi_E(x_E) = \int \frac{\mathrm{d}^4 p_E}{(2\pi)^4} e^{ip_E \cdot x_E} \tilde{\psi}_E(p_E) , \qquad (A19)
$$

$$
\bar{\psi}_E(x) = \int \frac{\mathrm{d}^4 p_E}{(2\pi)^4} e^{-ip_E \cdot x_E} \tilde{\bar{\psi}}_E(p_E) . \tag{A20}
$$

Hereafter, we omit tildes on the Fourier modes.

In this convention for the Euclidean signature, the action for the spinor field reads

$$
S = \int d^4x \left[\bar{\psi}(x) i \partial \psi(x) - m \bar{\psi}(x) \psi(x) - V(\bar{\psi}, \psi) \right]
$$

= $-i \int d^4x_E \left[-\bar{\psi}_E(x_E) (\gamma_{E\mu} \partial_{E\mu}) \psi_E(x_E) - m \bar{\psi}_E(x_E) \psi_E(x_E) - V_E(\bar{\psi}_E, \psi_E) \right]$
= iS_E , (A21)

where

$$
S_E = \int d^4x_E \left[\bar{\psi}_E(x_E) \left(\partial_{E\mu} \right) \psi_E(x_E) + m \bar{\psi}_E(x_E) \psi_E(x_E) + V_E(\bar{\psi}_E, \psi_E) \right]. \tag{A22}
$$

In the main text and hereafter, we omit the subscript "E" denoting the Euclidean signature.

Appendix B: Derivation of flow equations

The central method in this work is the Wetterich equation whose form is given by

$$
\partial_t \Gamma_k = \frac{1}{2} \text{STr} \left[\left(\Gamma_k^{(2)} + \mathcal{R}_k \right)^{-1} \partial_t \mathcal{R}_k \right] = \frac{1}{2} \tilde{\partial}_t \text{STr} \log \left[\Gamma_k^{(2)} + \mathcal{R}_k \right], \tag{B1}
$$

where $\tilde{\partial}_t = (\partial_t \mathcal{R}_k) \frac{\partial}{\partial \mathcal{R}_k}$ is the scale derivative acting only on the regulator function. In general, the Hessian $\Gamma_k^{(2)} + \mathcal{R}_k$ is given as the supermatrix in superfield space. Namely, we can write in the form of

$$
\Gamma_k^{(2)}(p) + \mathcal{R}_k(p) = \frac{\overrightarrow{\delta}}{\delta \Phi^T(-p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \Phi(p)} + \mathcal{R}_k(p) = \begin{pmatrix} M_{\rm BB} & M_{\rm BF} \\ M_{\rm FB} & M_{\rm FF} \end{pmatrix},
$$
\n(B2)

where $\Phi = (\phi, \psi, \bar{\psi}^T, \dots)$ is the superfield. In what follows, we consider the case of the fermionic system.

1. Supermatrix, superdeterminant and supertrace

We summarize the useful formulas for supermatrix to evaluate the Wetterich equation. We begin by considering a supermatrix in a form of

$$
M = \begin{pmatrix} M_{\rm BB} & M_{\rm BF} \\ M_{\rm FB} & M_{\rm FF} \end{pmatrix} . \tag{B3}
$$

Here, M_{BB} and M_{FF} are Grassmann-even elements, while M_{BF} and M_{FB} are Grassmann-odd. We define the supertrace for the supermatrix $(B3)$ as

$$
str M = tr M_{BB} - tr M_{FF}, \qquad (B4)
$$

which satisfies

$$
str\ (MN) = str\ (NM) \,. \tag{B5}
$$

With this definition, the superdeterminant is defined by

$$
s\det M = \exp(str \ln M),\tag{B6}
$$

such that

$$
sdet (MN) = sdet M \cdot sdet N. \tag{B7}
$$

Now, we deform M as

$$
M = \begin{pmatrix} M_{\rm BB} & 0 \\ M_{\rm FB} & 1 \end{pmatrix} \begin{pmatrix} 1 & M_{\rm BB}^{-1} M_{\rm BF} \\ 0 & N_{\rm FF} \end{pmatrix},
$$
 (B8)

where we have defined

$$
N_{\rm FF} = M_{\rm FF} - M_{\rm FB} M_{\rm BB}^{-1} M_{\rm BF} \,. \tag{B9}
$$

From Eq. [\(B7\)](#page-9-2), we have

$$
sdet M = sdet \begin{pmatrix} M_{BB} & 0 \\ M_{FB} & 1 \end{pmatrix} sdet \begin{pmatrix} 1 & M_{BB}^{-1} M_{BF} \\ 0 & N_{FF} \end{pmatrix}.
$$
 (B10)

Here, using Eq. [\(B6\)](#page-9-3) gives

$$
\mathrm{sdet}\begin{pmatrix} M_{\mathrm{BB}} & 0 \\ M_{\mathrm{FB}} & 1 \end{pmatrix} = \exp\left\{ \mathrm{str} \, \ln \left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \begin{pmatrix} M_{\mathrm{BB}} - 1 & 0 \\ M_{\mathrm{FB}} & 1 \end{pmatrix} \right] \right\}
$$

$$
= \exp\left\{\text{str}\sum_{n=1}^{\infty}\frac{(-)^{n-1}}{n}\begin{pmatrix}M_{\text{BB}}-1 & 0\\M_{\text{FB}} & 1\end{pmatrix}\right\}
$$

$$
= \exp\left\{\text{tr}\sum_{n=1}^{\infty}\frac{(-)^{n-1}}{n}(M_{\text{BB}}-1)^{n}\right\}
$$

$$
= \exp\{\text{tr}\ln M_{\text{BB}}\}
$$

$$
= \text{det} M_{\text{BB}}.
$$
(B11)

In the same manner, we obtain

$$
sdet\begin{pmatrix} 1 & M_{\rm BB}^{-1}M_{\rm BF} \\ 0 & N_{\rm FF} \end{pmatrix} = \exp\{-\text{tr}\ln N_{\rm FF}\} = (\det N_{\rm FF})^{-1}.
$$

Note here that the minus sign arises from Eq. [\(B4\)](#page-9-4). Finally, we arrive at

$$
s\det M = \frac{\det M_{\rm BB}}{\det N_{\rm FF}}.
$$
\n(B12)

This implies that

$$
\log \text{sdet } M = \log \text{det } M_{\text{BB}} - \log \text{det}(M_{\text{FF}} - M_{\text{FB}} M_{\text{BB}}^{-1} M_{\text{BF}})
$$

= tr $\log M_{\text{BB}} - \text{tr } \log(M_{\text{FF}} - M_{\text{FB}} M_{\text{BB}}^{-1} M_{\text{BF}}).$ (B13)

Instead of Eq. $(B8)$, the supermatrix $(B3)$ can also be deformed as

$$
M = \begin{pmatrix} N_{\text{BB}} & M_{\text{BF}} M_{\text{FF}}^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ M_{\text{FB}} & M_{\text{FF}} \end{pmatrix} . \tag{B14}
$$

Here, we have defined

$$
N_{\rm BB} = M_{\rm BB} - M_{\rm BF} M_{\rm FF}^{-1} M_{\rm FB}
$$
\n(B15)

Then, the superdeterminant for M can be written in the form as

$$
sdet M = \frac{det N_{BB}}{det M_{FF}},
$$
\n(B16)

for which we have

$$
\log \det M = \log \det \left(M_{\rm BB} - M_{\rm BF} M_{\rm FF}^{-1} M_{\rm FB} \right) - \log \det M_{\rm FF}
$$

= tr
$$
\log (M_{\rm BB} - M_{\rm BF} M_{\rm FF}^{-1} M_{\rm FB}) - \text{tr } \log M_{\rm FF}.
$$
 (B17)

2. General structure of flow equation

For the pure fermionic system, the supermatrix [\(B3\)](#page-9-1) is

$$
M \equiv \mathcal{G}_k^{-1}(p) = \frac{\overrightarrow{\delta}}{\delta \Phi^T(-p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \Phi(p)} + \mathcal{R}_k(p) = \begin{pmatrix} 0 & 0\\ 0 & M_{\text{FF}} \end{pmatrix} . \tag{B18}
$$

Here, we have the fermionic part in the field basis $\Phi(p) = \begin{pmatrix} \psi(p) \\ \tau(p) \end{pmatrix}$ $\bar{\psi}^T(-p)$ \setminus and $\Phi^T(-p) = (\psi^T(-p) \overline{\psi}(p))$ as

$$
M_{\rm FF} = \begin{pmatrix} \frac{\overrightarrow{\delta}}{\delta \psi^T(-p)} \mathcal{G}_k^{-1}(p) \frac{\overleftarrow{\delta}}{\delta \psi(p)} & \frac{\overrightarrow{\delta}}{\delta \psi^T(-p)} \mathcal{G}_k^{-1}(p) \frac{\overleftarrow{\delta}}{\delta \overline{\psi}^T(-p)} \\\frac{\overrightarrow{\delta}}{\delta \overline{\psi}(p)} \mathcal{G}_k^{-1}(p) \frac{\overleftarrow{\delta}}{\delta \psi(p)} & \frac{\overrightarrow{\delta}}{\delta \overline{\psi}(p)} \mathcal{G}_k^{-1}(p) \frac{\overleftarrow{\delta}}{\delta \overline{\psi}^T(-p)} \end{pmatrix} . \tag{B19}
$$

We briefly summarize several techniques to derive the beta functions. For a given effective action Γ_k , we compute the second-order functional derivative. Together with the regulator function, one schematically has

$$
\mathcal{G}_k^{-1}(p)\Big|_{\text{FF}} = (\mathcal{K}_k + \mathcal{R}_k) + \mathcal{V}_k = \mathcal{P}_k^{-1} + \mathcal{V}_k, \tag{B20}
$$

where \mathcal{K}_k is the field-independent part, while \mathcal{V}_k includes the vertex terms depending on field variables. More specifically, for an effective action, we obtain

$$
\mathcal{K}_{k}(p,-p) = \begin{pmatrix}\n\overrightarrow{\delta} & \overleftarrow{\delta} & \overrightarrow{\delta} \\
\overrightarrow{\delta\psi^{T}(-p)}^{\Gamma_{k}}\overrightarrow{\delta\psi(p)} & \overrightarrow{\delta\psi^{T}(-p)}^{\Gamma_{k}}\overrightarrow{\delta\overrightarrow{\psi^{T}(-p)}}\n\end{pmatrix}\n\Big|_{\psi=\overline{\psi}=0} = \begin{pmatrix}\n0 & K_{k}^{T}(p) \\
K_{k}(p) & 0\n\end{pmatrix}\n(2\pi)^{4}\delta^{4}(0).
$$
\n(B21)

The regulator matrix is given by

$$
\mathcal{R}_k(p) = \begin{pmatrix} 0 & R_k^f(p)^T \\ R_k^f(p) & 0 \end{pmatrix} (2\pi)^4 \delta^4(0) = \begin{pmatrix} 0 & i p^T r_k^f(p/k) \\ i p r_k^f(p/k) & 0 \end{pmatrix} (2\pi)^4 \delta^4(0).
$$
 (B22)

There are many choices for the regulator $r_k^f(p/k)$. In this work, we employ the Litim-type cutoff [\[38\]](#page-16-1) given as

$$
r_k^f(p/k) = \left(\sqrt{\frac{k^2}{p^2}} - 1\right) \Theta(k^2 - p^2).
$$
 (B23)

Then, \mathcal{P}_k is just the regulated inverse-propagator such that

$$
\mathcal{P}_k^{-1}(p,-p) = \begin{pmatrix} 0 & (P_k^{-1})^T \\ P_k^{-1} & 0 \end{pmatrix} = \begin{pmatrix} 0 & (K_k(p) + R_k^f(p))^T \\ K_k(p) + R_k^f(p) & 0 \end{pmatrix} (2\pi)^4 \delta^4(0).
$$
 (B24)

For the vertex matrix, we write

$$
\mathcal{V}_k = \begin{pmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{pmatrix} . \tag{B25}
$$

Note here that $V_{12} = -(V_{21})^T$.

Once we compute Eq. [\(B20\)](#page-11-0), the regulated full propagator is obtained by the expansion in terms of the vertex operator \mathcal{V}_k such that

$$
\mathcal{G}_k(p)\Big|_{\rm FF} = \left(\Gamma_k^{(2)} + \mathcal{R}_k\right)^{-1} = \mathcal{P}_k - \mathcal{P}_k \mathcal{V}_k \mathcal{P}_k + \mathcal{P}_k \mathcal{V}_k \mathcal{P}_k \mathcal{V}_k \mathcal{P}_k + \cdots
$$
\n(B26)

Inserting this into flow equation (5) and using Eqs. $(B24)$ and $(B25)$, we have

$$
\partial_t \Gamma_k = \frac{1}{2} \tilde{\partial}_t S \text{Tr} \log \left[\mathcal{G}_k^{-1} |_{\text{FF}} \right] = \frac{1}{2} \tilde{\partial}_t S \text{Tr} \log \left[\mathcal{P}_k^{-1} + \mathcal{V}_k \right]
$$

\n
$$
= \frac{1}{2} \text{STr} \left[\mathcal{P}_k \partial_t \mathcal{R}_k \right] - \frac{1}{2} \text{STr} \left[\mathcal{P}_k \mathcal{V}_k \mathcal{P}_k \partial_t \mathcal{R}_k \right] + \frac{1}{2} \text{STr} \left[\mathcal{P}_k \mathcal{V}_k \mathcal{P}_k \mathcal{V}_k \mathcal{P}_k \partial_t \mathcal{R}_k \right] + \cdots
$$

\n
$$
= - \text{Tr} \left[P_k \partial_t R_k^f \right] + \left(\frac{1}{2} \text{Tr} \left[P_k V_{21} P_k \partial_t R_k^f \right] + \frac{1}{2} \text{Tr} \left[P_k^T V_{12} P_k^T \partial_t (R_k^f)^T \right] \right)
$$

\n
$$
- \text{Tr} \left[(P_k V_{21} P_k V_{21} \mathcal{P}_k - P_k V_{22} P_k^T V_{11} \mathcal{P}_k) \partial_t R_k^f \right] + \cdots
$$

\n
$$
= - \text{Tr} \left[P_k \partial_t R_k^f \right] + \text{Tr} \left[P_k V_{21} P_k \partial_t R_k^f \right] - \text{Tr} \left[(P_k V_{21} P_k V_{21} P_k - P_k V_{22} P_k^T V_{11} P_k) \partial_t R_k^f \right] + \cdots
$$
 (B27)

Note that the minus sign in Eq. [\(B27\)](#page-11-3) reflects the definition of the supertrace [\(B4\)](#page-9-4). The first and second terms on the left-hand side of Eq. [\(B27\)](#page-11-3) correspond to the quantum corrections to the vacuum energy and the mass term. We are interested especially in the third term which gives the quantum corrections to the four-Fermi interactions.

3. Application to four-Fermion model

Here, based on the previous formulas, we show the explicit computation for deriving the flow equations in the four-fermion model in Eq. [\(8\)](#page-3-0). The truncated effective action in our work is given by

$$
\Gamma_k = \int d^4x \left[\bar{\psi} \left(\gamma^\mu \partial_\mu + m e^{i\theta \gamma_5/2} \right) \psi - \frac{G_S}{2} \left(\bar{\psi} \psi \right)^2 - \frac{G_P}{2} \left(\bar{\psi} i \gamma_5 \psi \right)^2 - C_4 \left(\bar{\psi} \psi \right) \left(\bar{\psi} i \gamma_5 \psi \right) \right]. \tag{B28}
$$

In momentum space, the kinetic term is written in the form as

$$
\int d^4x \,\overline{\psi} \left(\gamma^\mu \partial_\mu + m e^{i\theta \gamma_5/2}\right) \psi = \int \frac{d^4p}{(2\pi)^4} \frac{d^4q}{(2\pi)^4} \int d^4x \, e^{i(p-q)\cdot x} \overline{\psi}(q) \left(i\gamma^\mu p_\mu + m e^{i\theta \gamma_5/2}\right) \psi(p)
$$
\n
$$
= \int \frac{d^4p}{(2\pi)^4} \overline{\psi}(p) \left(i\gamma^\mu p_\mu + m e^{i\theta \gamma_5/2}\right) \psi(p), \tag{B29}
$$

from which the kinetic matrix \mathcal{K}_k in momentum space is found to be

$$
(\mathcal{K}_k(p,-p))_{IJ} = \begin{pmatrix} 0 & (i\rlap/v - me^{i\theta\gamma_5/2})_{IJ}^T \\ (i\rlap/v + me^{i\theta\gamma_5/2})_{IJ} & 0 \end{pmatrix} (2\pi)^4 \delta^4(0).
$$
 (B30)

Together with the regulator matrix [\(B22\)](#page-11-4), the regulated inverse propagator reads

$$
(\mathcal{P}_k^{-1}(p,-p))_{IJ} = \begin{pmatrix} 0 & \left(i\rlap{/}p(1+r_k^f) - me^{i\theta\gamma_5/2} \right)_{IJ}^T \\ \left(i\rlap{/}p(1+r_k^f) + me^{i\theta\gamma_5/2} \right)_{IJ} & 0 \end{pmatrix} (2\pi)^4 \delta^4(0). \tag{B31}
$$

Its inverse form is

$$
(\mathcal{P}_k(p,-p))_{IJ} = \begin{pmatrix} 0 & \frac{\left(-i\rlap{/}p(1+r_k^f) + me^{-i\theta\gamma_5/2}\right)_{IJ}}{p^2(1+r_k^f)^2 + m^2} \\ \frac{\left(-i\rlap{/}p(1+r_k^f) - me^{-i\theta\gamma_5/2}\right)_{IJ}}{p^2(1+r_k^f)^2 + m^2} & 0 \end{pmatrix} (2\pi)^4 \delta^4(0). \tag{B32}
$$

The vertex matrix [\(B25\)](#page-11-2) is calculated as

$$
(V_{11})_{IJ} = \frac{\overrightarrow{\delta}}{\delta \psi_I(-p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \psi_J(p)} = \frac{G_S}{2} \left[\bar{\psi}_{i,M} \delta_{MI} \bar{\psi}_N \delta_{NJ} \right] + \frac{G_P}{2} \left[\bar{\psi}_M(i\gamma_5)_{MI} \bar{\psi}_N(i\gamma_5)_{NJ} \right] + C_4 \left[\bar{\psi}_{i,M} \delta_{MI} \bar{\psi}_N(i\gamma_5)_{NJ} \right],
$$
(B33)

$$
(V_{22})_{IJ} = \frac{\overrightarrow{\delta}}{\delta \overline{\psi}_I(p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \overline{\psi}_J(-p)} = \frac{G_S}{2} \Big[\delta_{IM} \psi_M \delta_{JN} \psi_N \Big] + \frac{G_P}{2} \Big[(i\gamma_5)_{IM} \psi_M (i\gamma_5)_{JN} \psi_N \Big] + C_4 \Big[\delta_{IM} \psi_M (i\gamma_5)_{JN} \psi_N \Big] ,
$$
(B34)

$$
(V_{21})_{IJ} = \frac{\overrightarrow{\delta}}{\delta \overline{\psi}_I(p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \psi_J(p)} = -G_S \Big[\delta_{IJ} \overline{\psi} \psi \Big] - G_P \Big[(i\gamma_5)_{IJ} \overline{\psi} i\gamma_5 \psi \Big] - C_4 \Big[\delta_{IJ} \overline{\psi} i\gamma_5 \psi \Big] - C_4 \Big[\overline{\psi} \psi (i\gamma_5)_{IJ} \Big] - G_S \Big[\delta_{IM} \psi_M \overline{\psi}_N \delta_{NJ} \Big] - G_P \Big[(i\gamma_5)_{IM} \psi_M \overline{\psi}_N (i\gamma_5)_{NJ} \Big] - C_4 \Big[\delta_{IM} \psi_M \overline{\psi}_N (i\gamma_5)_{NJ} \Big] - C_4 \Big[(i\gamma_5)_{IM} \psi_M \overline{\psi}_N \delta_{NJ} \Big],
$$
(B35)

$$
(V_{12})_{IJ} = \frac{\overrightarrow{\delta}}{\delta \psi_I(-p)} \Gamma_k \frac{\overleftarrow{\delta}}{\delta \overline{\psi}_J(-p)} = -(V_{21})_{IJ} . \tag{B36}
$$

The large- N_c approximation corresponds to the following replacement: $(V_{11})_{IJ} \rightarrow 0$, $(V_{22})_{IJ} \rightarrow 0$,

$$
(V_{21})_{IJ} \rightarrow (V_{21}^{\text{LN}})_{IJ} = -G_P \left[\delta_{IJ} \bar{\psi} \psi \right] - G_S \left[(i\gamma_5)_{IJ} \bar{\psi} i\gamma_5 \psi \right] - C_4 \left[\delta_{IJ} \bar{\psi} i\gamma_5 \psi \right] - C_4 \left[\bar{\psi} \psi (i\gamma_5)_{IJ} \right],\tag{B37}
$$

$$
(V_{12})_{IJ} \rightarrow -\left[(V_{21}^{\text{LN}}) \right]_{IJ}^T.
$$

$$
{(2)}{IJ} \rightarrow -\left[(V_{21}^{\text{LN}}) \right]_{IJ}^T. \tag{B38}
$$

a. Vacuum energy

First, we evaluate the vacuum energy part which is independent of field operators. From the first term on the right-hand side of Eq. [\(B27\)](#page-11-3), we obtain

$$
\partial_t \rho_k = -\frac{1}{\Omega_4} \text{Tr} \left[P_k \partial_t R_k^f \right] = -\int \frac{\mathrm{d}^4 p}{(2\pi)^4} \text{tr} \left[\frac{\left(-i\rlap/v (1 + r_k^f) + m e^{-i\theta\gamma_5/2} \right)}{p^2 (1 + r_k^f)^2 + m^2} i(\rlap/v)_{JK} \partial_t r_k^f(p/k) \delta_{\alpha\beta}^{\text{color}} \right]
$$
\n
$$
= -4N_c \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \left[\frac{p^2 (1 + r_k^f)}{p^2 (1 + r_k^f)^2 + m^2} \partial_t r_k^f(p/k) \right]
$$
\n
$$
= -\frac{4N_c}{(4\pi)^2} \int_0^{k^2} \mathrm{d}p^2 \, p^2 \left[\frac{k^2}{k^2 + m^2} \right] \,, \tag{B39}
$$

where $\Omega_4 = \int d^4x = (2\pi)^4 \delta^4(0)$ is the four-dimensional spacetime volume and tr $\delta_{\alpha\beta}^{\text{color}} = \delta_{\alpha\alpha}^{\text{color}} = N_c$. This contribution however does not influence upon the spinor dynamics, so is irrelevant in the current work.

b. Mass term

From the second term on the right-hand side of Eq. [\(B27\)](#page-11-3), we read off quantum corrections for the mass term with the CP and P violating angle θ , such that

$$
\bar{\psi}\partial_t (me^{i\theta\gamma_5/2})\psi = \partial_t (m\cos(\theta/2))\bar{\psi}\psi + \partial_t (m\sin(\theta/2))\bar{\psi}\gamma^5\psi = \frac{1}{\Omega_4} \text{Tr} \left[P_k (V_{21}^{\text{LN}}) P_k \partial_t R_k^f \right]
$$
\n
$$
= \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \text{tr} \left[\frac{\left(-i\rlap{/}p(1+r_k^f) + me^{-i\theta\gamma_5/2} \right)_{IK}}{p^2(1+r_k^f)^2 + m^2} (V_{21}^{\text{LN}}) \kappa L \frac{\left(-i\rlap{/}p(1+r_k^f) + me^{-i\theta\gamma_5/2} \right)_{LM}}{p^2(1+r_k^f)^2 + m^2} i(\rlap{/}p)_{MJ} \partial_t r_k^f(p/k) \right]
$$
\n
$$
= \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{\partial_t r_k^f(p/k)}{(p^2(1+r_k^f)^2 + m^2)^2} \text{tr} \left[\left(-i\rlap{/}p(1+r_k^f) + me^{-i\theta\gamma_5/2} \right)_{IK} (V_{21}^{\text{LN}}) \kappa L \left(-i\rlap{/}p(1+r_k^f) + me^{-i\theta\gamma_5/2} \right)_{LM} i(\rlap{/}p)_{MJ} \right]
$$
\n
$$
= - \left[8N_c G_S \cos \left(\frac{\theta}{2} \right) + 8N_c C_4 \sin \left(\frac{\theta}{2} \right) \right] m \mathcal{M}(\bar{\psi}\psi) - \left[8N_c G_P \sin \left(\frac{\theta}{2} \right) + 8N_c C_4 \cos \left(\frac{\theta}{2} \right) \right] m \mathcal{M}(\bar{\psi}\gamma^5\psi), \qquad \text{(B40)}
$$

where we have defined the threshold function as

$$
\mathcal{M} = \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^2} = \frac{1}{2(4\pi)^2} \frac{k^6}{(k^2 + m^2)^2}.
$$
 (B41)

We can read off the flow equations as

$$
\partial_t (m \cos(\theta/2)) = -8N_c G_S m \cos\left(\frac{\theta}{2}\right) \mathcal{M} - 8N_c C_4 m \sin\left(\frac{\theta}{2}\right) \mathcal{M},\tag{B42}
$$

$$
\partial_t(m\sin(\theta/2)) = -8N_c G_P m \sin\left(\frac{\theta}{2}\right) \mathcal{M} - 8N_c C_4 m \cos\left(\frac{\theta}{2}\right) \mathcal{M},\tag{B43}
$$

from which we have the flow equations for m and θ as

$$
\partial_t m = \cos(\theta/2)\partial_t(m\cos(\theta/2)) + \sin(\theta/2)\partial_t(m\sin(\theta/2))
$$

= $-4N_c[(G_S + G_P) + 2C_4\sin\theta + (G_S - G_P)\cos\theta]m\mathcal{M},$ (B44)

$$
\partial_t \theta = \frac{2}{m} \left[-\sin(\theta/2) \partial_t (m \cos(\theta/2)) + \cos(\theta/2) \partial_t (m \sin(\theta/2)) \right]
$$

= $8N_c \left[(G_S - G_P) \sin \theta - 2C_4 \cos \theta \right] \mathcal{M}.$ (B45)

Here we define the dimensionless couplings

$$
\tilde{m} = mk^{-1}, \qquad \tilde{G}_S = G_S k^2, \qquad \tilde{G}_P = G_P k^2, \qquad \tilde{C}_4 = C_4 k^2. \qquad (B46)
$$

Then, the flow equations for \tilde{m} and θ read

$$
\partial_t \tilde{m} = -\tilde{m} - 4N_c \left[(\tilde{G}_S + \tilde{G}_P) + 2\tilde{C}_4 \sin \theta + (\tilde{G}_S - \tilde{G}_P) \cos \theta \right] \tilde{m} \tilde{\mathcal{M}}, \tag{B47}
$$

$$
\partial_t \theta = 8 \left[(\tilde{G}_S - \tilde{G}_P) \sin \theta - 2 \tilde{C}_4 \cos \theta \right] \tilde{\mathcal{M}}, \tag{B48}
$$

where the dimensionless threshold function is

$$
\widetilde{\mathcal{M}} = \frac{1}{k^2} \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^2} = \frac{1}{2(4\pi)^2} \frac{1}{(1 + \tilde{m}^2)^2} \,. \tag{B49}
$$

c. Four-Fermion interactions

f

The third term on the right-hand side of Eq. [\(B27\)](#page-11-3) corresponds to quantum corrections for the four-fermion interactions which are obtained from the flow equation as

$$
-\frac{\partial_t G_S}{2}(\bar{\psi}\psi)^2 - \frac{\partial_t G_P}{2}(\bar{\psi}i\gamma^5\psi)^2 - \partial_t C_4(\bar{\psi}\psi)(\bar{\psi}i\gamma^5\psi) = -\frac{1}{\Omega_4} \text{Tr} \left[P_k V_{21}^{\text{LN}} P_k V_{21}^{\text{LN}} P_k \partial_t R_k^f \right]. \tag{B50}
$$

We compute the right-hand side as

$$
-\frac{1}{\Omega_{4}} \text{Tr} \left[P_{k} V_{21}^{\text{LN}} P_{k} V_{21}^{\text{LN}} P_{k} \partial_{t} R_{k}^{f} \right]
$$
\n
$$
= -\frac{1}{\Omega_{4}} \text{Tr} \left[\frac{(-i\rlap/v (1 + r_{k}^{f}) + m e^{-i\theta \gamma_{5}/2})}{p^{2}(1 + r_{k}^{f})^{2} + m^{2}} V_{21} \frac{(-i\rlap/v (1 + r_{k}^{f}) + m e^{-i\theta \gamma_{5}/2})}{p^{2}(1 + r_{k}^{f})^{2} + m^{2}} V_{21} \frac{(-i\rlap/v (1 + r_{k}^{f}) + m e^{-i\theta \gamma_{5}/2})}{p^{2}(1 + r_{k}^{f})^{2} + m^{2}} \partial_{t} R_{k}^{f} \right]
$$
\n
$$
= -\int \frac{d^{4}p}{(2\pi)^{4}} \frac{p^{2}(1 + r_{k}^{f})\partial_{t} r_{k}^{f}(p/k)}{p^{2}(1 + r_{k}^{f})^{2} + m^{2})^{3}} \left[-N_{c} \left\{ (4G_{S}^{2} + 4C_{4}^{2})(k^{2} - m^{2}) - 2m^{2}(4G_{S}^{2} - 4C_{4}^{2})\cos\theta - 16m^{2}G_{S}C_{4}\sin\theta \right\} (\bar{\psi}\psi)^{2} - N_{c} \left\{ (4G_{P}^{2} + 4C_{4}^{2})(k^{2} - m^{2}) + 2m^{2}(4G_{P}^{2} - 4C_{4}^{2})\cos\theta - 16m^{2}G_{P}C_{4}\sin\theta \right\} (\bar{\psi}\psi)^{5}\psi \right\}
$$
\n
$$
- 4N_{c} \left\{ 2(G_{S} + G_{P})C_{4}(k^{2} - m^{2}) - 4m^{2}(G_{S} - G_{P})C_{4}\cos\theta - m^{2}(4G_{S}G_{P} + 4C_{4}^{2})\sin\theta \right\} (\bar{\psi}\psi)(\bar{\psi}\psi^{5}\psi) \right]
$$
\n
$$
= \left[4N_{c} \left\{ (G_{S}^{2} + C_{4}^{2})(k^{2} - m^{2}) + 2m^{2}(G_{S}^{2
$$

where we have defined the threshold function

$$
\mathcal{I} = \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^3} = \frac{1}{2(4\pi)^2} \frac{k^6}{(k^2 + m^2)^3}.
$$
 (B52)

Then, we obtain the flow equations

$$
-\frac{\partial_t G_S}{2} = 4N_c \left\{ (G_S^2 + C_4^2)(k^2 - m^2) + 2m^2(G_S^2 - C_4^2) \cos \theta + 4m^2 G_S C_4 \sin \theta \right\} \mathcal{I},\tag{B53}
$$

$$
-\frac{\partial_t G_P}{2} = 4N_c \left\{ (G_P^2 + C_4^2)(k^2 - m^2) - 2m^2(G_P^2 - C_4^2) \cos \theta + 4m^2 G_P C_4 \sin \theta \right\} \mathcal{I},\tag{B54}
$$

$$
-\partial_t C_4 = 8N_c \left\{ (G_S + G_P)C_4(k^2 - m^2) + 2m^2(G_S - G_P)C_4 \cos\theta + 2m^2(G_S G_P + C_4^2) \sin\theta \right\} \mathcal{I}.
$$
 (B55)

For the dimensionless couplings [\(B46\)](#page-14-0), we have

$$
\partial_t \tilde{G}_S = 2\tilde{G}_S - 8N_c \Big[(\tilde{G}_S^2 + \tilde{C}_4^2)(1 - \tilde{m}^2) + 2\tilde{m}^2(\tilde{G}_S^2 - \tilde{C}_4^2) \cos \theta + 4\tilde{m}^2 \tilde{G}_S \tilde{C}_4 \sin \theta \Big] \tilde{\mathcal{I}},\tag{B56}
$$

$$
\partial_t \tilde{G}_P = 2\tilde{G}_P - 8N_c \Big[(\tilde{G}_P^2 + \tilde{C}_4^2)(1 - \tilde{m}^2) - 2\tilde{m}^2(\tilde{G}_P^2 - \tilde{C}_4^2) \cos \theta + 4\tilde{m}^2 \tilde{G}_P \tilde{C}_4 \sin \theta \Big] \tilde{\mathcal{I}},\tag{B57}
$$

$$
\partial_t \tilde{C}_4 = 2\tilde{C}_4 - 8N_c \Big[(\tilde{G}_S + \tilde{G}_P)\tilde{C}_4 (1 - \tilde{m}^2) + 2\tilde{m}^2 (\tilde{G}_S - \tilde{G}_P)\tilde{C}_4 \cos\theta + 2\tilde{m}^2 (\tilde{G}_S \tilde{G}_P + \tilde{C}_4^2) \sin\theta \Big] \tilde{\mathcal{I}},
$$
(B58)

with the dimensionless threshold function

$$
\widetilde{\mathcal{I}} = \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{p^2 (1 + r_k^f) \partial_t r_k^f (p/k)}{(p^2 (1 + r_k^f)^2 + m^2)^3} = \frac{1}{2(4\pi)^2} \frac{1}{(1 + \tilde{m}^2)^3} \,. \tag{B59}
$$

- [1] C. Abel et al., Measurement of the Permanent Electric Dipole Moment of the Neutron, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.124.081803) 124 (2020) 081803 [[2001.11966](https://arxiv.org/abs/2001.11966)].
- [2] χ QCD collaboration, Nucleon electric dipole moment from the θ term with lattice chiral fermions, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.108.094512) 108 (2023) [094512](https://doi.org/10.1103/PhysRevD.108.094512) [[2301.04331](https://arxiv.org/abs/2301.04331)].
- [3] A. E. Nelson, *Naturally Weak CP Violation, [Phys. Lett. B](https://doi.org/10.1016/0370-2693(84)92025-2)* **136** (1984) 387.
- [4] S. M. Barr, Solving the Strong CP Problem Without the Peccei-Quinn Symmetry, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.53.329) 53 (1984) 329.
- [5] S. M. Barr, A Natural Class of Nonpeccei-quinn Models, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.30.1805) 30 (1984) 1805.
- [6] K. S. Babu and R. N. Mohapatra, A Solution to the Strong CP Problem Without an Axion, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.41.1286) 41 (1990) 1286.
- [7] S. M. Barr, D. Chang and G. Senjanovic, *Strong CP problem and parity, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.67.2765)* **67** (1991) 2765.
- [8] R. N. Mohapatra and G. Senjanovic, Natural Suppression of Strong p and t Noninvariance, [Phys. Lett. B](https://doi.org/10.1016/0370-2693(78)90243-5) 79 (1978) 283.
- [9] M. A. B. Beg and H. S. Tsao, Strong P, T Noninvariances in a Superweak Theory, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.41.278) 41 (1978) 278.
- [10] H. Georgi, A Model of Soft CP Violation, Hadronic J. 1 (1978) 155.
- [11] M. Dine, R. G. Leigh and A. Kagan, Supersymmetry and the Nelson-Barr mechanism, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.48.2214) 48 (1993) 2214 [[hep-ph/9303296](https://arxiv.org/abs/hep-ph/9303296)].
- [12] M. Dine and P. Draper, Challenges for the Nelson-Barr Mechanism, JHEP 08 [\(2015\) 132](https://doi.org/10.1007/JHEP08(2015)132) [[1506.05433](https://arxiv.org/abs/1506.05433)].
- [13] C. Vafa and E. Witten, Restrictions on Symmetry Breaking in Vector-Like Gauge Theories, [Nucl. Phys. B](https://doi.org/10.1016/0550-3213(84)90230-X) 234 (1984) 173.
- [14] E. E. Jenkins, A. V. Manohar and P. Stoffer, Low-Energy Effective Field Theory below the Electroweak Scale: Anomalous Dimensions, JHEP 01 [\(2018\) 084](https://doi.org/10.1007/JHEP01(2018)084) [[1711.05270](https://arxiv.org/abs/1711.05270)].
- [15] J. Hisano, K. Tsumura and M. J. S. Yang, QCD Corrections to Neutron Electric Dipole Moment from Dimension-six Four-Quark Operators, [Phys. Lett. B](https://doi.org/10.1016/j.physletb.2012.06.038) 713 (2012) 473 [[1205.2212](https://arxiv.org/abs/1205.2212)].
- [16] J. de Vries, P. Draper, K. Fuyuto, J. Kozaczuk and B. Lillard, Uncovering an axion mechanism with the EDM portfolio, Phys. Rev. D 104 [\(2021\) 055039](https://doi.org/10.1103/PhysRevD.104.055039) [[2107.04046](https://arxiv.org/abs/2107.04046)].
- [17] T. Banno, J. Hisano, T. Kitahara and N. Osamura, Closer look at the matching condition for radiative QCD θ parameter, JHEP 02 [\(2024\) 195](https://doi.org/10.1007/JHEP02(2024)195) [[2311.07817](https://arxiv.org/abs/2311.07817)].
- [18] K.-I. Aoki, K.-i. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, Nonperturbative renormalization group analysis of the chiral critical behaviors in QED, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.97.479) 97 (1997) 479 [[hep-ph/9612459](https://arxiv.org/abs/hep-ph/9612459)].
- [19] K.-I. Aoki, K. Takagi, H. Terao and M. Tomoyose, Nonladder extended renormalization group analysis of the dynamical chiral symmetry breaking, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.103.815) 103 (2000) 815 [[hep-th/0002038](https://arxiv.org/abs/hep-th/0002038)].
- [20] K.-I. Aoki and D. Sato, Solving the QCD non-perturbative flow equation as a partial differential equation and its application to the dynamical chiral symmetry breaking, PTEP 2013 [\(2013\) 043B04](https://doi.org/10.1093/ptep/ptt018) [[1212.0063](https://arxiv.org/abs/1212.0063)].
- [21] K.-I. Aoki, S.-I. Kumamoto and D. Sato, Weak solution of the non-perturbative renormalization group equation to describe dynamical chiral symmetry breaking, PTEP 2014 [\(2014\) 043B05](https://doi.org/10.1093/ptep/ptu039) [[1403.0174](https://arxiv.org/abs/1403.0174)].
- [22] L. Bento, G. C. Branco and P. A. Parada, A Minimal model with natural suppression of strong CP violation, [Phys. Lett. B](https://doi.org/10.1016/0370-2693(91)90530-4) 267 [\(1991\) 95.](https://doi.org/10.1016/0370-2693(91)90530-4)
- [23] C. Wetterich, *Exact evolution equation for the effective potential, [Phys. Lett.](https://doi.org/10.1016/0370-2693(93)90726-X)* **B301** (1993) 90 [[1710.05815](https://arxiv.org/abs/1710.05815)].
- [24] T. R. Morris, The Exact renormalization group and approximate solutions, [Int. J. Mod. Phys.](https://doi.org/10.1142/S0217751X94000972) A9 (1994) 2411 [[hep-ph/9308265](https://arxiv.org/abs/hep-ph/9308265)].
- [25] M. Reuter and C. Wetterich, *Effective average action for gauge theories and exact evolution equations*, *[Nucl. Phys.](https://doi.org/10.1016/0550-3213(94)90543-6)* **B417** [\(1994\) 181.](https://doi.org/10.1016/0550-3213(94)90543-6)
- [26] U. Ellwanger, Flow equations for N point functions and bound states, Proceedings, Workshop on Quantum field theoretical aspects of high energy physics: Bad Frankenhausen, Germany, September 20-24, 1993, Z. Phys. C62 [\(1994\) 503](https://doi.org/10.1007/BF01555911) [[hep-ph/9308260](https://arxiv.org/abs/hep-ph/9308260)].
- [27] T. R. Morris, Elements of the continuous renormalization group, Nonperturbative QCD: Structure of the QCD vacuum: Proceedings, Yukawa International Seminar, YKIS'97, Kyoto, Japan, December 2-12, 1997, [Prog. Theor. Phys. Suppl.](https://doi.org/10.1143/PTPS.131.395) 131 [\(1998\) 395](https://doi.org/10.1143/PTPS.131.395) [[hep-th/9802039](https://arxiv.org/abs/hep-th/9802039)].
- [28] J. Berges, N. Tetradis and C. Wetterich, Nonperturbative renormalization flow in quantum field theory and statistical physics, [Phys. Rept.](https://doi.org/10.1016/S0370-1573(01)00098-9) 363 (2002) 223 [[hep-ph/0005122](https://arxiv.org/abs/hep-ph/0005122)].
- [29] K. Aoki, Introduction to the nonperturbative renormalization group and its recent applications, [Int.J.Mod.Phys.](https://doi.org/10.1142/S0217979200000923) B14 (2000) [1249.](https://doi.org/10.1142/S0217979200000923)
- [30] C. Bagnuls and C. Bervillier, *Exact renormalization group equations. An Introductory review, [Phys. Rept.](https://doi.org/10.1016/S0370-1573(00)00137-X)* **348** (2001) 91 [[hep-th/0002034](https://arxiv.org/abs/hep-th/0002034)].
- [31] J. Polonyi, Lectures on the functional renormalization group method, [Central Eur. J. Phys.](https://doi.org/10.2478/BF02475552) 1 (2003) 1 [[hep-th/0110026](https://arxiv.org/abs/hep-th/0110026)].
- [32] J. M. Pawlowski, Aspects of the functional renormalisation group, [Annals Phys.](https://doi.org/10.1016/j.aop.2007.01.007) **322** (2007) 2831 [[hep-th/0512261](https://arxiv.org/abs/hep-th/0512261)].
- [33] H. Gies, Introduction to the functional RG and applications to gauge theories, [Lect.Notes Phys.](https://doi.org/10.1007/978-3-642-27320-9_6) 852 (2012) 287 [[hep-ph/0611146](https://arxiv.org/abs/hep-ph/0611146)].
- [34] B. Delamotte, An Introduction to the nonperturbative renormalization group, [Lect. Notes Phys.](https://doi.org/10.1007/978-3-642-27320-9_2) 852 (2012) 49 [[cond-mat/0702365](https://arxiv.org/abs/cond-mat/0702365)].
- [35] H. Sonoda, The Exact Renormalization Group: Renormalization theory revisited, 9, 2007, [0710.1662](https://arxiv.org/abs/0710.1662).
- [36] Y. Igarashi, K. Itoh and H. Sonoda, Realization of Symmetry in the ERG Approach to Quantum Field Theory, [Prog. Theor.](https://doi.org/10.1143/PTPS.181.1) [Phys. Suppl.](https://doi.org/10.1143/PTPS.181.1) **181** (2010) 1 [[0909.0327](https://arxiv.org/abs/0909.0327)].
- [37] O. J. Rosten, Fundamentals of the Exact Renormalization Group, [Phys. Rept.](https://doi.org/10.1016/j.physrep.2011.12.003) 511 (2012) 177 [[1003.1366](https://arxiv.org/abs/1003.1366)].
- [38] D. F. Litim, *Optimized renormalization group flows, Phys. Rev.* **D64** [\(2001\) 105007](https://doi.org/10.1103/PhysRevD.64.105007) [[hep-th/0103195](https://arxiv.org/abs/hep-th/0103195)].
- [39] M. Salmhofer and C. Honerkamp, Fermionic renormalization group flows: Technique and theory, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.105.1) 105 [\(2001\) 1.](https://doi.org/10.1143/PTP.105.1)
- [40] P. Kopietz, L. Bartosch and F. Schütz, *Introduction to the functional renormalization group*, vol. 798. Springer Berlin, Heidelberg, 2010, [10.1007/978-3-642-05094-7.](https://doi.org/10.1007/978-3-642-05094-7)
- [41] J. Braun, Fermion Interactions and Universal Behavior in Strongly Interacting Theories, J. Phys. **G39** [\(2012\) 033001](https://doi.org/10.1088/0954-3899/39/3/033001) [[1108.4449](https://arxiv.org/abs/1108.4449)].
- [42] W. Metzner, M. Salmhofer, C. Honerkamp, V. Meden and K. Schonhammer, Functional renormalization group approach to correlated fermion systems, [Rev. Mod. Phys.](https://doi.org/10.1103/RevModPhys.84.299) 84 (2012) 299 [[1105.5289](https://arxiv.org/abs/1105.5289)].
- [43] K.-I. Aoki and K. Miyashita, Evaluation of the spontaneous chiral symmetry breaking scale in general gauge theories with non-perturbative renormalization group, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.121.875) 121 (2009) 875.
- [44] J. Braun, M. Leonhardt and M. Pospiech, Fierz-complete NJL model study: Fixed points and phase structure at finite temperature and density, Phys. Rev. D 96 [\(2017\) 076003](https://doi.org/10.1103/PhysRevD.96.076003) [[1705.00074](https://arxiv.org/abs/1705.00074)].
- [45] J. Braun, M. Leonhardt and M. Pospiech, Fierz-complete NJL model study. II. Toward the fixed-point and phase structure of hot and dense two-flavor QCD, Phys. Rev. D 97 [\(2018\) 076010](https://doi.org/10.1103/PhysRevD.97.076010) [[1801.08338](https://arxiv.org/abs/1801.08338)].
- [46] J. Braun, M. Leonhardt and M. Pospiech, Fierz-complete NJL model study III: Emergence from quark-gluon dynamics, Phys. Rev. D 101 [\(2020\) 036004](https://doi.org/10.1103/PhysRevD.101.036004) [[1909.06298](https://arxiv.org/abs/1909.06298)].
- [47] K.-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, Wilson renormalization group equations for the critical dynamics of chiral symmetry, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.102.1151) 102 (1999) 1151 [[hep-th/9908042](https://arxiv.org/abs/hep-th/9908042)].
- [48] G. 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, [NATO Sci. Ser. B](https://doi.org/10.1007/978-1-4684-7571-5_9) 59 (1980) 135.
- [49] J. D. Wells, The Utility of Naturalness, and how its Application to Quantum Electrodynamics envisages the Standard Model and Higgs Boson, [Stud. Hist. Phil. Sci. B](https://doi.org/10.1016/j.shpsb.2015.01.002) 49 (2015) 102 [[1305.3434](https://arxiv.org/abs/1305.3434)].
- [50] M. Frank, M. Buballa and M. Oertel, Flavor mixing effects on the QCD phase diagram at nonvanishing isospin chemical potential: One or two phase transitions?, Phys. Lett. B 562 (2003) 221 [[hep-ph/0303109](https://arxiv.org/abs/hep-ph/0303109)].
- [51] D. Boer and J. K. Boomsma, Spontaneous CP-violation in the strong interaction at theta = pi, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.78.054027) 78 (2008) [054027](https://doi.org/10.1103/PhysRevD.78.054027) [[0806.1669](https://arxiv.org/abs/0806.1669)].
- [52] J. K. Boomsma and D. Boer, The High temperature CP-restoring phase transition at theta = pi, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.80.034019) 80 (2009) [034019](https://doi.org/10.1103/PhysRevD.80.034019) [[0905.4660](https://arxiv.org/abs/0905.4660)].
- [53] Y. Sakai, H. Kouno, T. Sasaki and M. Yahiro, Theta vacuum effects on QCD phase diagram, [Phys. Lett. B](https://doi.org/10.1016/j.physletb.2011.10.032) 705 (2011) 349 [[1105.0413](https://arxiv.org/abs/1105.0413)].
- [54] H. Kodama and J.-I. Sumi, Application of nonperturbative renormalization group to Nambu-Jona-Lasinio / Gross-Neveu model at finite temperature and chemical potential, [Prog. Theor. Phys.](https://doi.org/10.1143/PTP.103.393) 103 (2000) 393 [[hep-th/9912215](https://arxiv.org/abs/hep-th/9912215)].
- [55] K.-I. Aoki and M. Yamada, The RG flow of Nambu–Jona-Lasinio model at finite temperature and density, [Int. J. Mod.](https://doi.org/10.1142/S0217751X15501808) Phys. A 30 [\(2015\) 1550180](https://doi.org/10.1142/S0217751X15501808) [[1504.00749](https://arxiv.org/abs/1504.00749)].
- [56] C. Alexandrou, A. Athenodorou, K. Hadjiyiannakou and A. Todaro, Neutron electric dipole moment using lattice QCD simulations at the physical point, Phys. Rev. D 103 [\(2021\) 054501](https://doi.org/10.1103/PhysRevD.103.054501) [[2011.01084](https://arxiv.org/abs/2011.01084)].
- [57] K.-I. Aoki, K. Morikawa, J.-I. Sumi, H. Terao and M. Tomoyose, Analysis of the Wilsonian effective potentials in dynamical chiral symmetry breaking, Phys. Rev. D 61 [\(2000\) 045008](https://doi.org/10.1103/PhysRevD.61.045008) [[hep-th/9908043](https://arxiv.org/abs/hep-th/9908043)].
- [58] H. Gies and C. Wetterich, Renormalization flow of bound states, Phys. Rev. D 65 [\(2002\) 065001](https://doi.org/10.1103/PhysRevD.65.065001) [[hep-th/0107221](https://arxiv.org/abs/hep-th/0107221)].
- [59] H. Gies and C. Wetterich, Universality of spontaneous chiral symmetry breaking in gauge theories, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.69.025001) 69 (2004) [025001](https://doi.org/10.1103/PhysRevD.69.025001) [[hep-th/0209183](https://arxiv.org/abs/hep-th/0209183)].
- [60] S. Floerchinger and C. Wetterich, *Exact flow equation for composite operators*, *[Phys. Lett. B](https://doi.org/10.1016/j.physletb.2009.09.014)* 680 (2009) 371 [[0905.0915](https://arxiv.org/abs/0905.0915)].
- [61] M. Mitter, J. M. Pawlowski and N. Strodthoff, Chiral symmetry breaking in continuum QCD, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.91.054035) 91 (2015) [054035](https://doi.org/10.1103/PhysRevD.91.054035) [[1411.7978](https://arxiv.org/abs/1411.7978)].
- [62] J. Braun, L. Fister, J. M. Pawlowski and F. Rennecke, From Quarks and Gluons to Hadrons: Chiral Symmetry Breaking in Dynamical QCD, Phys. Rev. D 94 [\(2016\) 034016](https://doi.org/10.1103/PhysRevD.94.034016) [[1412.1045](https://arxiv.org/abs/1412.1045)].
- [63] T. Denz, M. Mitter, J. M. Pawlowski, C. Wetterich and M. Yamada, Partial bosonization for the two-dimensional Hubbard model, Phys. Rev. B 101 [\(2020\) 155115](https://doi.org/10.1103/PhysRevB.101.155115) [[1910.08300](https://arxiv.org/abs/1910.08300)].
- [64] K.-I. Aoki, S.-I. Kumamoto and M. Yamada, Phase structure of NJL model with weak renormalization group, [Nucl. Phys. B](https://doi.org/10.1016/j.nuclphysb.2018.04.005) 931 [\(2018\) 105](https://doi.org/10.1016/j.nuclphysb.2018.04.005) [[1705.03273](https://arxiv.org/abs/1705.03273)].