Cosmological Constraints on Dark Neutrino Towers

Luis A. Anchordoqui,^{1, 2, 3} Ignatios Antoniadis,^{4, 5} Dieter Lüst,^{6, 7} and Karem Peñaló Castillo¹

¹Department of Physics and Astronomy,

Lehman College, City University of New York, NY 10468, USA

²Department of Physics,

Graduate Center, City University of New York, NY 10016, USA

³Department of Astrophysics, American Museum of Natural History, NY 10024, USA

⁴High Energy Physics Research Unit, Faculty of Science,

Chulalongkorn University, Bangkok 1030, Thailand

⁵Laboratoire de Physique Théorique et Hautes Énergies - LPTHE

Sorbonne Université, CNRS, 4 Place Jussieu, 75005 Paris, France

⁶Max-Planck-Institut für Physik, Werner-Heisenberg-Institut, 80805 München, Germany

⁷Arnold Sommerfeld Center for Theoretical Physics,

Ludwig-Maximilians-Universität München, 80333 München, Germany

(Dated: November 2024)

We reexamine a dynamical dark matter model with Kaluza-Klein (KK) towers of gravitons and neutrinos fitting together in the dark dimension. We show that even though gravitational decays of neutrino KK towers have little impact in cosmology the weak decay channel could have significant cosmological effects. Taking conservative upper bounds on the dark matter decay rate into two photons before recombination and on the number of effective extra neutrino species ΔN_{eff} we derive constraints on the conversion rate from active to sterile species despite the dependence of the mixing angle on the KK mode mass. We also provide counterarguments to a recent claim suggesting that the bounds on ΔN_{eff} rule out micron-sized extra dimensions.

I. INTRODUCTION

The mechanism behind neutrino masses is now in limbo. One interesting proposal envisions the righthanded neutrinos as five-dimensional (5D) bulk states, with Yukawa couplings to the left-handed lepton and Higgs doublets that are localized states on the Standard Model (SM) brane [1–3]. The neutrino Yukawa couplings become tiny because of a volume suppression, yielding naturally light Dirac neutrinos. As a by-product of this type of construct, a neutrino tower of Kaluza-Klein (KK) modes arises with masses proportional to the inverse radius R_{\perp} of the fifth dimension.

Very recently, cosmological constraints on neutrino KK towers were reconsidered [4], with a deceptive conclusion that current bounds on the effective number of noninteracting relativistic species from measurements of the cosmic microwave background (CMB) [5] generically rule out micron-sized extra dimensions. In this paper we show that the limits inferred in [4] are not generic but rather model dependent. A major shortcoming of the analysis carried out in [4] is that the assumed cosmic evolution of the KK tower does not allow for intra-tower neutrino decays, viz. decays of a given KK mode in the tower into final states that include other, lighter KK excitations. Such a "dark-to-dark" decay process usually dominates the cosmological evolution of the tower and could be regarded as the telltale signature of the dynamical dark matter framework [6]. We also show that the limits on R_{\perp} inferred in [4] do not particularly apply to the dark dimension scenario [7].

The layout of the paper is as follows. We begin in Sec. II by summarizing phenomenological aspects of the dark dimension scenario. Along the way, we also provide a brief review of the cosmological evolution of dark gravitons and their interplay with dark neutrino towers [8]. In Sec. III we reexamine cosmological constraints on weak decays of bulk sterile neutrinos into active neutrinos and photons. The paper wraps up in Sec. IV with some conclusions.

II. DYNAMICAL DARK MATTER WITH TOWERS OF GRAVITONS AND NEUTRINOS

The dark dimension is a five-dimensional (5D) setup that has a compact space with characteristic lengthscale in the micron range. Most notably, this setup provides an economic explanation of the cosmological hierarchy problem, because the anti-de Sitter distance conjecture in de Sitter space [9] connects the size of the compact space R_{\perp} to the dark energy scale $\Lambda^{1/4}$ via $R_{\perp} \sim \lambda \Lambda^{1/4}$, where $\Lambda \sim 10^{-120} M_p^4$ is the cosmological constant, M_p the reduced Planck mass, and the proportionality factor is estimated to be within the range $10^{-1} < \lambda < 10^{-4}$ [7]. The KK tower of the dark dimension opens up at the mass scale $m_{\rm KK} \sim 1/R_{\perp}$. The species scale M_* where gravity becomes strong is linked to $m_{\rm KK}$ via $M_* \sim m_{\rm KK}^{1/3} M_p^{2/3}$ [10, 11]. In our calculations we adopt $m_{\rm KK} \sim 0.1$ eV which implies $M_* \sim 10^9$ GeV.

The dark dimension assembles a colosseum for dark matter (DM) contenders. Primordial black holes with Schwarzschild radius smaller than a micron provide one interesting DM candidate [12–17]. Massive spin-2 KK excitations of the graviton take the place of a second compelling possibility [8]. Herein we focus attention on

the second scenario whereby DM production proceeds through the coupling of SM fields with the 5D graviton. The cosmological chronicle begins with the matter fields in equilibrium at initial temperature $T_{\rm in} \sim {\rm GeV}$ [8]. These are localized recurrences in the compact space, which manifest as a tower of equally spaced dark gravitons, indexed by an integer l, and with mass $m_l = l m_{\rm KK}$. The production of graviton mode of mass m_l is dominantly at temperatures $T_l \sim m_l$. Thereby, the produced dark gravitons are initially in the GeV mass range.

A point worth noting at this juncture is that the cosmological overproduction of bulk graviton modes implies that in models with n large or mesoscopic extra dimensions the maximum temperatures must be less than

$$T_{\star} \sim 10^{3+(6n-15)/(n+2)} \left(\frac{M_{\star}}{10^9 \text{ GeV}}\right) \text{ GeV}$$
 (1)

which is an upper limit on the "normalcy" temperature at which the universe must be free of bulk modes [18]. Note that for n = 1, the assumed $T_{in} \sim 1$ GeV saturates the normalcy temperature.

The cosmic evolution of the dark sector is mostly driven by dark-to-dark decay processes, which regulate the decay of KK gravitons within the dark tower, realizing a particular version of the dynamical dark matter framework [6]. In the absence of isometries in the dark dimension, which is the common expectation, the KK momentum of the dark tower is not conserved [19]. This means that a dark graviton of KK quantum n can decay to two other ones, with quantum numbers n_1 and n_2 . If the KK quantum violation can go up to δn , the number of available channels is roughly $l \, \delta n$. In addition, because the decay is almost at threshold, the phase space factor is roughly the velocity of decay products, $v_{\rm r.m.s.} \sim \sqrt{m_{\rm KK} \, \delta n/m_l}$. All in all, the total decay width of graviton G_l is found to be,

$$\Gamma_{\text{tot}}^{G_l} \sim \sum_{l' < l} \sum_{0 < l'' < l-l'} \Gamma_{G_{l'}G_{l''}}^{G_l}$$
$$\sim \beta^2 \frac{m_l^3}{M_p^2} \times \frac{m_l}{m_{\text{KK}}} \, \delta n \times \sqrt{\frac{m_{\text{KK}}\delta n}{m_l}}$$
$$\sim \beta^2 \, \delta n^{3/2} \frac{m_l^{7/2}}{M_p^2 m_{\text{KK}}^{7/2}}, \qquad (2)$$

where β is parameter that controls the strength of the intra-tower decay amplitudes which correlates with the amplitudes on inhomogeneities in the dark dimension [8]. We further follow [8] to estimate the time evolution of the dark matter mass and assume that for times larger than $1/\Gamma_{\rm tot}^{G_l}$ dark matter which is heavier than the corresponding m_l has already decayed, yielding

$$m_l \sim \left(\frac{M_p^4 \ m_{\rm KK}}{\beta^4 \ \delta n^3}\right)^{1/7} t^{-2/7},$$
 (3)

where t indicates the time elapsed since the big bang.

The dark dimension scenario also provides visible signals on the brane, e.g., the decay of relic graviton KK states to photons. The partial decay width of such a process is estimated to be

$$\Gamma_{\gamma\gamma}^{G_l} = \frac{\tilde{\lambda}^2 \ m_{\rm KK}^3 \ l^3}{80\pi M_p^2} \,, \tag{4}$$

where the parameter λ measures the value of the dark graviton wave function at the SM brane and is expected to be $\mathcal{O}(1)$ [20, 21].

Hitherto, dark dimension cosmology can be characterized by a set of three parameters $\{\beta, \delta n, \lambda\}$, which must be constrained by experiment. In particular, CMB measurements are sensitive to processes that change the photon-baryon fluid between today and the last scattering surface. To be more specific, throughout the CMB epoch during the emission of the relic at redshift $z_{\rm CMB} \sim 1100$, matter dominates the energy density of the Universe. The number density of baryons is mostly composed of neutral hydrogen atoms (H_I) , together with a smaller Helium (He) component, $x_{\rm He} = n_{\rm He}/n_{\rm H_I} \simeq 1/13$, and a small percentage of free protons and electrons, $x_e = n_e/n_{\rm H_I} = n_p/n_{\rm H_I}$, which varies from about 20% at $z_{\rm CMB}$ to roughly 2×10^{-4} at $z \sim 20$ [22]. Most notably, the damping of the CMB power spectrum is driven by the integrated optical depth along the line of sight. Now, dark graviton towers decaying into photons within the redshift range $20 \lesssim z \lesssim 1100$ would bring in *exotic* energy that could ionize H_I and He. This in turn would increase the optical depth to recombination inducing a stronger damping of the CMB power spectrum on small scales. The non-observation of such a suppression in the spectrum places an upper limit on the energy delivered by DM decays into $\gamma\gamma$ [23], providing a direct constraint on λ [24].

In addition, we have seen that dark matter decay gives the daughter particles a velocity kick. Self-gravitating dark-matter halos that have a virial velocity smaller than this velocity kick may be disrupted by these particle decays. Combined cosmological zoom-in simulations of decaying dark matter with a model of the Milky Way satellite population rule out non-relativistic kick speeds $\gtrsim 10^{-4}$ for a dark matter lifetime $\tau_{\rm DM} \lesssim 29$ Gyr at 95%CL [25]. However, N-body simulations of isolated dark-matter halos seem to indicate that if $\tau_{\rm DM} \gtrsim 60$ Gyr and the kick speed $\lesssim 10^{-2}$ then the halos are essentially unchanged [26]. Herein, we adopt the 95%CL bound on today's dark matter kick velocity $\leq 2.2 \times 10^{-4}$ derived in [27], which directly constrain the β and δn parameters.

We adopt as benchmark the following parameter set $\{\beta \sim 500, \ \delta n \sim 0.2, \ \tilde{\lambda} \sim 0.5\}$ which is consistent with all available data; see Fig. 4 in [27]. Substituting these figures into (2) and (4) we obtain the following parametrization of the decay rates as a function of m_l ,

$$\Gamma_{\rm tot}^{G_l} \sim 5 \times 10^{-4} \left(\frac{m_l}{\rm GeV}\right)^{7/2} \, {\rm s}^{-1}$$
 (5)

$$\Gamma_{\gamma\gamma}^{G_l} \sim 2 \times 10^{-16} \left(\frac{m_l}{\text{GeV}}\right)^3 \text{ s}^{-1}, \qquad (6)$$

respectively. We can also conclude substituting the parameters β and δn into (3) that at present the DM has a mass $m_{l,\text{today}} \sim 78$ keV, where we have taken an age of the universe of 13.8 Gyr.

Thus far, we have assumed that DM is composed entirely of dark gravitons, but as it was already put forward in [8] the composition of the dynamical dark matter ensemble could call for other light modes in the bulk, which lead to new decay channels and additional DM components. To be specific, we do expect dark fermions to propagate in the bulk playing the role of right-handed neutrinos. Generally speaking, one would expect that the existence of other KK species would not affect the total abundance or cosmological evolution of the DM mass as the KK modes would just distribute among each other, as they decay from one to another. For example, the l'summed width of the decay $\nu_l \rightarrow \nu_{l'}G_{l-l'}$ derived in [28],

$$\sum_{l'} \Gamma^{\nu_l}_{\nu_{l'}G_{l-l'}} = \frac{m_l^4 \ R_\perp}{6\pi \ M_p^2} \sim 10^{-4} \ \left(\frac{m_l}{1 \ \text{GeV}}\right)^4 \ \text{s}^{-1} \,, \quad (7)$$

is comparable to the total decay width of the graviton (5), but somewhat smaller reflecting the tiny violation of the KK momentum conservation, i.e. $\delta n \sim 0.2$. However, for the particular case of dark neutrino towers we should exercise some caution, because these towers do not only couple gravitationally to the SM sector but also via Fermi's weak interaction. It is this weak decay channel that we now turn to study.

III. CONSTRAINTS ON BULK NEUTRINOS

Consider three 5D Dirac fermions Ψ_{α} , which are singlets under the SM gauge symmetries and interact in our brane with the three active left-handed neutrinos in a way that conserves lepton number. The S^1/\mathbb{Z}_2 symmetry in the dark dimension coordinate y contains y to -y, which acts as chirality (γ_5) on spinors. Then, in the Weyl basis each Dirac field can be decomposed into two two-component spinors $\Psi_{\alpha} \equiv (\psi_{\alpha L}, \psi_{\alpha R})^T$.

Neutrino masses arise in 5D bulk-brane interactions of the form

$$\mathscr{L} \supset h_{ij} \ \overline{L}_i \ H \ \psi_{jR}(y=0) \,, \tag{8}$$

where $H = -i\sigma_2 H^*$, L_i denotes the lepton doublets (localized on the SM brane), ψ_{jR} stands for the three bulk (right-handed) *R*-neutrinos evaluated at the position of the SM brane, y = 0 in the dark dimension, and h_{ij} are coupling constants. This gives a coupling with the *L*-neutrinos of the form $\langle H \rangle \ \overline{\nu}_{L_i} \ \psi_{jR}(y=0)$, where $\langle H \rangle = 175$ GeV is the Higgs vacuum expectation value. Expanding ψ_{jR} into modes canonically normalized leads for each of them to a Yukawa 3×3 matrix suppressed by the square root of the volume of the bulk $\sqrt{\pi R_{\perp} M_s}$, i.e.,

$$Y_{ij} = \frac{h_{ij}}{\sqrt{\pi R_\perp M_s}} \sim h_{ij} \frac{M_s}{M_p}, \qquad (9)$$

where $M_s \leq M_*$ is the string scale, and where in the second rendition we have dropped factors of π 's and of the string coupling.

Data analyses from short- and long-baseline neutrino oscillation experiments, together with observations of neutrinos produced by cosmic rays collisions in the atmosphere and by nuclear fusion reactions in the Sun, provide the most sensitive insights to determine the extremely small mass-squared differences. Neutrino oscillation data can be well-fitted in terms of two nonzero differences $\Delta m_{ij}^2 = m_i^2 - m_j^2$ between the squares of the masses of the three (i = 1, 2, 3) neutrino mass eigenvalues m_i , yielding $\Delta m_{\rm SOL}^2 = \Delta m_{21}^2 = (7.53 \pm 0.18) \times 10^{-5} \text{ eV}^2$ and $\Delta m_{\rm ATM}^2 = |\Delta m_{31}^2| \simeq \Delta m_{32}^2 = 2.453 \pm 0.033) \times 10^{-3} \text{ eV}^2$ or $\Delta m_{32}^2 = -2.536 \pm 0.034) \times 10^{-3} \text{ eV}^2$ [29]. A straightforward calculation shows that to obtain the correct order of magnitude of neutrino masses the coupling h_{ij} should be $\mathcal{O}(10^{-4})$ for our fiducial value $M_s \sim 10^9$ GeV [30].

Now, light modes in the bulk contribute to the effective number of relativistic neutrino-like species N_{eff} [37] and are bounded by experiment [5]. Using conservation of entropy, fully thermalized relics with g_* degrees of freedom contribute

$$\Delta N_{\rm eff} = g_* \left(\frac{43}{4g_s}\right)^{4/3} \begin{cases} 4/7 & \text{for bosons} \\ 1/2 & \text{for fermions} \end{cases}, \quad (10)$$

where q_s denotes the effective degrees of freedom for the entropy of the other thermalized relativistic species that are present when they decouple [38]. The 5D graviton has 5 helicities, but the spin-1 helicities do not have zero modes, because we assume the compactification has S^1/\mathbb{Z}_2 symmetry and so the ± 1 helicities are projected out. The spin-0 is the radion and the spin-2 helicities form the massless (zero mode) graviton. This means that for the 5D graviton, $g_* = 3$. The (bulk) left-handed neutrinos are odd, but the right-handed neutrinos are even and so each counts as a Weyl neutrino, for a total $q_* = 2 \times 3$. Assuming that the dark sector decouples from the SM sector before the electroweak phase transition we have $g_s = 106.75$. This gives $\Delta N_{\text{eff}} = 0.22$, in agreement with CMB observations [5]. Actually, $\Delta N_{\text{eff}} = 0.22$ roughly saturates the value from CMB observations.

The KK states of the neutrino towers are nonrelativistic during the CMB formation epoch. Indeed, in the presence of bulk masses [31, 32], the mixing of the first KK modes to active neutrinos can be suppressed [33, 34].¹ Thus, these KK states do not directly

¹ We note in passing that bulk neutrino masses relax the constraints derived in [35, 36] from neutrino oscillation experiments; for details, see [34].

contribute to $\Delta N_{\rm eff}$. However, these dark neutrinos can decay into SM fields (including neutrinos and photons) via weak interactions through their small active admixture. The active neutrinos that are produced through these decays after neutrino decoupling at $T_{\nu} \sim \text{MeV}$ but before the CMB epoch contribute to $\Delta N_{\rm eff}$.

Next, in line with our stated plan, we estimate the unsolicited contributions to $\Delta N_{\rm eff}$ from KK decays mediated by Fermi weak interaction dressed by an active-sterile mixing angle. The partial decay width of the process $\nu_l \rightarrow \nu \nu \nu$ is found to be [28]

$$\Gamma_{\nu\nu\nu}^{\nu_l} = \frac{G_F^2 \ m_l^5 \ \sin^2\theta_l}{192\pi^3} \sim \frac{\zeta^2 \ m_\nu^2 \ m_l^3}{\tau_\mu m_\mu^5} \,, \qquad (11)$$

where $\theta_l \sim \zeta m_{\nu}/m_l$ is the vacuum mixing angle between an active neutrino of mass $m_{\nu} \sim 0.1$ eV and a KK mode with mass m_l [39], ζ parametrizes our ignorance of KK decays and neutrino mixing in the dark dimension, and

$$G_F = \sqrt{\frac{192 \ \pi^3}{m_{\mu}^5 \ \tau_{\mu} \ (1 + \Delta q)}} \tag{12}$$

is the Fermi constant, with m_{μ} and τ_{μ} the muon mass and its lifetime, respectively, and where $\Delta q \ll 1$ parametrizes the effect of radiative corrections [40]. The partial width of the radiative decay $\nu_l \rightarrow \gamma \nu$ is smaller by a factor of $27\alpha/(8\pi)$, with α the fine structure constant [41].

Now, by demanding that $\Gamma_{\gamma\nu}^{\nu_l} \lesssim \Gamma_{\gamma\gamma}^{G_l}$ we find $\zeta \lesssim 0.01$. Indeed, for $\zeta \sim 0.01$ we obtain

$$\Gamma^{\nu_l}_{\gamma\nu} \sim 2.5 \times 10^{-16} \left(\frac{m_l}{\text{GeV}}\right)^3 \,\text{s}^{-1}\,,$$
 (13)

and

$$\Gamma^{\nu_l}_{\nu\nu\nu\nu} \sim 3 \times 10^{-14} \left(\frac{m_l}{\text{GeV}}\right)^3 \text{ s}^{-1}.$$
 (14)

Note that the partial decay widths (13) and (14) are much smaller than the inverse of the age of the universe at CMB $(t_{\rm CMB}^{-1} \sim 10^{-13} \text{ s}^{-1})$, and so there is no contribution to $\Delta N_{\rm eff}$ from KK weak decays. Moreover, by construction, the partial decay width of $\nu_l \rightarrow \gamma \nu$ is comparable to the size of the width derived in [24] for $G_l \rightarrow \gamma \gamma$, ensuring that for $z \gtrsim 20$, the KK weak decay would not inject enough energy to perturb the abundances of H_I and He. We conclude that nowadays micron-sized extra dimensions are consistent with observations.

IV. CONCLUSIONS

We have reexamined the dynamical dark matter model with KK towers of gravitons and neutrinos proposed in [8]. More specifically, we investigated the impact of weak decays from dark neutrino towers in cosmology. Our results are encapsulated in Fig. 1 and can be summarized as follows:



FIG. 1: Partial decay widths of various processes governing the cosmic evolution of the dynamical dark matter model with towers of gravitons and neutrinos proposed in [8]. For comparison, the horizontal lines indicate the inverse of the age of the universe at z = 20 and z = 1100, characterizing the reionization and CMB epochs, respectively.

- A thorough analysis of data from the *Planck* mission presented in [24] provides strict constraints on the partial decay width $\Gamma_{\gamma\gamma}^{G_l}$ not to perturb the reionization process of the Universe. Radiative decays $\nu_l \rightarrow \gamma \nu$, going via loop diagrams, can also discompose the reionization process. By demanding that $\Gamma_{\gamma\nu}^{\nu_l} \lesssim \Gamma_{\gamma\gamma}^{G_l}$ we have placed a bound on the vacuum sterile mixing, which is driven by the parameter ζ . We have shown that $\zeta \lesssim 0.01$ ensures that KK weak decays would not inject enough energy to perturb the abundances of H_I and He at $z \gtrsim 20$. As can be seen in Fig. 1 the decay widths of $\Gamma_{\gamma\nu}^{\nu_l}$ and $\Gamma_{\gamma\gamma}^{G_l}$ are always smaller than the inverse of the age of the universe during reionization $t_{\rm reio}^{-1} \sim 2 \times 10^{-16} \, {\rm s}^{-1}$ despite the dependence of the mixing angle on the KK mode mass.
- As can be deduced from Fig. 1 the upper bound on the parameter $\zeta \lesssim 0.01$ guarantees that decay widths of processes with active neutrinos in the final state are all smaller than $t_{\rm CMB}^{-1}$. This implies that the contribution to $\Delta N_{\rm eff}$ from weak decays of KK towers is negligible. Thus, the DM model proposed in [24] is consistent with upper bounds on $\Delta N_{\rm eff}$ from Planck data.
- In vacuum and with a seesaw model, $m_{\nu} \ll m_l$, the active-sterile mixing angle is generally approximated by $\theta_l \sim m_{\nu}/m_l$. The additional suppression factor which we have bounded to be $\zeta \lesssim 0.01$ is consistent with the estimates obtained in 4D sterile neutrino models; see e.g. [39, 42].
- The cosmological evolution of the KK towers is dominated by dark-to-dark decay processes. Actually, the intra-KK decays in the bulk require a spontaneous breakdown of the translational invariance in the compact space such that the 5D momenta are not conserved. The level of violation of KK momentum conservation is very small: $\delta n \sim 0.2$.

This is visible in Fig. 1 through a comparison of the total decay width $\Gamma_{\text{tot}}^{G_l}$ with $\sum_{l'} \Gamma_{\nu_{l'}G_{l-l'}}^{\nu_l}$.

- Alternatively, we could postulate that the violation of 5D momentum conservation is large such that KK right-handed-neutrino modes would be able decay fast to the zero mode, before neutrino decoupling. If this were the case, there would be no cosmological effects. Of course, in such a scenario DM cannot be realized by KK gravitons. DM can instead be due to 5D primordial black holes [12–17] or/and a fuzzy radion [43].
- One major assumption of the dynamical dark matter model proposed in [8] is that the SM brane is in a thermal state at T_{in} and the KK modes remain essentially unexcited. As the SM brane begins cooling off, its universal coupling to bulk fields gives inevitable KK production. Now, different pieces of experimental data point to an initial temperature just above the QCD phase transition. As noted in [8], such a happenstance may actually be en-
- K. R. Dienes, E. Dudas and T. Gherghetta, Neutrino oscillations without neutrino masses or heavy mass scales: A Higher dimensional seesaw mechanism, Nucl. Phys. B 557, 25 (1999) doi:10.1016/S0550-3213(99)00377-6 [arXiv:hep-ph/9811428 [hep-ph]].
- [2] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and J. March-Russell, Neutrino masses from large extra dimensions, Phys. Rev. D 65, 024032 (2001) doi:10.1103/PhysRevD.65.024032 [arXiv:hepph/9811448 [hep-ph]].
- G. R. Dvali and A. Y. Smirnov, Probing large extra dimensions with neutrinos, Nucl. Phys. B 563, 63-81 (1999) doi:10.1016/S0550-3213(99)00574-X [arXiv:hepph/9904211 [hep-ph]].
- [4] D. McKeen, J. N. Ng and M. Shamma, Signatures of bulk neutrinos in the early Universe, Phys. Rev. D 110, no.8, 8 (2024) doi:10.1103/PhysRevD.110.083507 [arXiv:2406.05266 [hep-ph]].
- [5] N. Aghanim et al. [Planck], Planck 2018 results VI: Cosmological parameters, Astron. Astrophys. 641, A6 (2020) [erratum: Astron. Astrophys. 652, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [6] K. R. Dienes and B. Thomas, Dynamical Dark Matter: I. Theoretical Overview, Phys. Rev. D 85, 083523 (2012) doi:10.1103/PhysRevD.85.083523 [arXiv:1106.4546 [hepph]].
- [7] M. Montero, C. Vafa and I. Valenzuela, The dark dimension and the Swampland, JHEP 02, 022 (2023) doi:10.1007/JHEP02(2023)022 [arXiv:2205.12293 [hepth]].
- [8] E. Gonzalo, M. Montero, G. Obied and C. Vafa, Dark Dimension Gravitons as Dark Matter, JHEP 11, 109 (2023) doi:10.1007/JHEP11(2023)109 [arXiv:2209.09249 [hep-ph]].
- [9] D. Lüst, E. Palti and C. Vafa, AdS and the Swampland, Phys. Lett. B **797**, 134867 (2019)

crypted in

$$T_{\rm in} \sim \left(\Lambda/M_p^4\right)^{1/6} M_p \sim 1 \,\,\mathrm{GeV}\,,\tag{15}$$

suggesting a possible UV/IR mixing at the interplay between dark energy and $T_{\rm in}$. An investigation along these lines is obviously important to be done.

Acknowledgements

The work of L.A.A. and K.P.C. is supported by the U.S. National Science Foundation (NSF Grant PHY-2412679). I.A. is supported by the Second Century Fund (C2F), Chulalongkorn University. The work of D.L. is supported by the Origins Excellence Cluster and by the German-Israel-Project (DIP) on Holography and the Swampland.

doi:10.1016/j.physletb.2019.134867 [arXiv:1906.05225 [hep-th]].

- [10] G. Dvali, Black holes and large N species solution to the hierarchy problem, Fortsch. Phys. 58, 528-536 (2010) doi:10.1002/prop.201000009 [arXiv:0706.2050 [hep-th]].
- G. Dvali and M. Redi, Black hole bound on the number of species and quantum gravity at LHC, Phys. Rev. D 77, 045027 (2008) doi:10.1103/PhysRevD.77.045027 [arXiv:0710.4344 [hep-th]].
- [12] L. A. Anchordoqui, I. Antoniadis and D. Lüst, Dark dimension, the swampland, and the dark matter fraction composed of primordial black holes, Phys. Rev. D 106, no.8, 086001 (2022) doi:10.1103/PhysRevD.106.086001 [arXiv:2206.07071 [hep-th]].
- [13] L. A. Anchordoqui, I. Antoniadis and D. Lüst, The dark universe: Primordial black hole \Rightarrow dark graviton gas connection, Phys. Lett. B **840**, 137844 (2023) doi:10.1016/j.physletb.2023.137844 [arXiv:2210.02475 [hep-th]].
- [14] L. A. Anchordoqui, I. Antoniadis and D. Lüst, Dark dimension, the swampland, and the dark matter fraction composed of primordial near-extremal black holes, Phys. Rev. D 109, no.9, 095008 (2024) doi:10.1103/PhysRevD.109.095008 [arXiv:2401.09087 [hep-th]].
- [15] L. A. Anchordoqui, I. Antoniadis and D. Lüst, More on black holes perceiving the dark dimension, Phys. Rev. D 110, no.1, 015004 (2024) doi:10.1103/PhysRevD.110.015004 [arXiv:2403.19604 [hep-th]].
- [16] L. A. Anchordoqui, I. Antoniadis, D. Lüst and K. Peñaló Castillo, Bulk black hole dark matter, Phys. Dark Univ. 46, 101714 (2024) doi:10.1016/j.dark.2024.101714 [arXiv:2407.21031 [hep-th]].
- [17] L. A. Anchordoqui, I. Antoniadis, D. Lüst and K. Peñaló Castillo, Through the looking glass into the dark dimension: Searching for bulk black hole dark matter with mi-

crolensing of X-ray pulsars, Phys. Dark Univ. **46**, 101681 (2024) doi:10.1016/j.dark.2024.101681 [arXiv:2409.12904 [hep-ph]].

- [18] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phenomenology, astrophysics and cosmology of theories with submillimeter dimensions and TeV scale quantum gravity, Phys. Rev. D 59, 086004 (1999) doi:10.1103/PhysRevD.59.086004 [arXiv:hepph/9807344 [hep-ph]].
- [19] R. N. Mohapatra, S. Nussinov and A. Perez-Lorenzana, Large extra dimensions and decaying *KK* recurrences, Phys. Rev. D 68, 116001 (2003) doi:10.1103/PhysRevD.68.116001 [arXiv:hepph/0308051 [hep-ph]].
- [20] T. Han, J. D. Lykken and R. J. Zhang, On Kaluza-Klein states from large extra dimensions, Phys. Rev. D 59, 105006 (1999) doi:10.1103/PhysRevD.59.105006 [arXiv:hep-ph/9811350 [hep-ph]].
- [21] L. J. Hall and D. Tucker-Smith, Cosmological constraints on theories with large extra dimensions, Phys. Rev. D 60, 085008 (1999) doi:10.1103/PhysRevD.60.085008
 [arXiv:hep-ph/9904267 [hep-ph]].
- [22] Y. Ali-Haimoud and C. M. Hirata, Ultrafast effective multi-level atom method for primordial hydrogen recombination, Phys. Rev. D 82, 063521 (2010) doi:10.1103/PhysRevD.82.063521 [arXiv:1006.1355 [astro-ph.CO]].
- [23] T. R. Slatyer and C. L. Wu, General Constraints on Dark Matter Decay from the Cosmic Microwave Background, Phys. Rev. D 95, no.2, 023010 (2017) doi:10.1103/PhysRevD.95.023010 [arXiv:1610.06933 [astro-ph.CO]].
- [24] J. A. P. Law-Smith, G. Obied, A. Prabhu and C. Vafa, Astrophysical constraints on decaying dark gravitons, JHEP 06, 047 (2024) doi:10.1007/JHEP06(2024)047 [arXiv:2307.11048 [hep-ph]].
- [25] S. Mau et al. [DES], Milky Way Satellite Census IV: Constraints on decaying dark matter from observations of Milky Way satellite galaxies, Astrophys. J. 932, no.2, 128 (2022) doi:10.3847/1538-4357/ac6e65 [arXiv:2201.11740 [astro-ph.CO]].
- [26] A. H. G. Peter, C. E. Moody and M. Kamionkowski, Dark-Matter Decays and Self-Gravitating Halos, Phys. Rev. D 81, 103501 (2010) doi:10.1103/PhysRevD.81.103501 [arXiv:1003.0419 [astro-ph.CO]].
- [27] G. Obied, C. Dvorkin, E. Gonzalo and C. Vafa, Dark dimension and decaying dark matter gravitons, Phys. Rev. D 109, no.6, 063540 (2024) doi:10.1103/PhysRevD.109.063540 [arXiv:2311.05318 [astro-ph.CO]].
- [28] K. Abazajian, G. M. Fuller and M. Patel, The cosmological bulk neutrino catastrophe, Phys. Rev. Lett. 90, 061301 (2003) doi:10.1103/PhysRevLett.90.061301 [arXiv:hep-ph/0011048 [hep-ph]].
- [29] S. Navas *et al.* [Particle Data Group], Review of particle physics, Phys. Rev. D **110**, no.3, 030001 (2024) doi:10.1103/PhysRevD.110.030001
- [30] L. A. Anchordoqui, I. Antoniadis and D. Lüst, Aspects of the dark dimension in cosmology, Phys. Rev. D 107, no.8, 083530 (2023) doi:10.1103/PhysRevD.107.083530

[arXiv:2212.08527 [hep-ph]].

- [31] A. Lukas, P. Ramond, A. Romanino and G. G. Ross, Solar neutrino oscillation from large extra dimensions, Phys. Lett. B 495, 136-146 (2000) doi:10.1016/S0370-2693(00)01206-5 [arXiv:hep-ph/0008049 [hep-ph]].
- [32] A. Lukas, P. Ramond, A. Romanino and G. G. Ross, Neutrino masses and mixing in brane world theories, JHEP 04, 010 (2001) doi:10.1088/1126-6708/2001/04/010 [arXiv:hep-ph/0011295 [hep-ph]].
- [33] M. Carena, Y. Y. Li, C. S. Machado, P. A. N. Machado and C. E. M. Wagner, Neutrinos in Large Extra Dimensions and Short-Baseline ν_e Appearance, Phys. Rev. D **96** (2017) no.9, 095014 doi:10.1103/PhysRevD.96.095014 [arXiv:1708.09548 [hep-ph]].
- [34] L. A. Anchordoqui, I. Antoniadis and J. Cunat, The dark dimension and the standard model landscape, Phys. Rev. D 109, no.1, 016028 (2024) doi:10.1103/PhysRevD.109.016028 [arXiv:2306.16491 [hep-ph]].
- [35] P. A. N. Machado, H. Nunokawa and R. Zukanovich Funchal, Testing for large extra dimensions with neutrino oscillations, Phys. Rev. D 84, 013003 (2011) doi:10.1103/PhysRevD.84.013003 [arXiv:1101.0003 [hepph]].
- [36] D. V. Forero, C. Giunti, C. A. Ternes and O. Tyagi, Large extra dimensions and neutrino experiments, Phys. Rev. D 106, no.3, 035027 (2022) doi:10.1103/PhysRevD.106.035027 [arXiv:2207.02790 [hep-ph]].
- [37] G. Steigman, D. N. Schramm and J. E. Gunn, Cosmological limits to the number of massive leptons, Phys. Lett. B 66 (1977), 202-204 doi:10.1016/0370-2693(77)90176-9
- [38] L. A. Anchordoqui and H. Goldberg, Neutrino cosmology after WMAP 7-Year data and LHC first Z' bounds, Phys. Rev. Lett. 108, 081805 (2012) doi:10.1103/PhysRevLett.108.081805 [arXiv:1111.7264 [hep-ph]].
- [39] S. Dodelson and L. M. Widrow, Sterile-neutrinos as dark matter, Phys. Rev. Lett. **72**, 17-20 (1994) doi:10.1103/PhysRevLett.72.17 [arXiv:hep-ph/9303287 [hep-ph]].
- [40] D. M. Webber et al. [MuLan], Measurement of the positive muon lifetime and determination of the Fermi constant to part-per-million precision, Phys. Rev. Lett. 106, 041803 (2011) doi:10.1103/PhysRevLett.106.079901 [arXiv:1010.0991 [hep-ex]].
- [41] V. D. Barger, R. J. N. Phillips and S. Sarkar, Remarks on the KARMEN anomaly, Phys. Lett. B 352, 365-371 (1995) [erratum: Phys. Lett. B 356, 617-617 (1995)] doi:10.1016/0370-2693(95)00486-5 [arXiv:hep-ph/9503295 [hep-ph]].
- [42] A. D. Dolgov and S. H. Hansen, Massive sterile neutrinos as warm dark matter, Astropart. Phys. 16, 339-344 (2002) doi:10.1016/S0927-6505(01)00115-3 [arXiv:hep-ph/0009083 [hep-ph]].
- [43] L. A. Anchordoqui, I. Antoniadis and D. Lüst, Fuzzy dark matter and the dark dimension, Eur. Phys. J. C 84, no.3, 273 (2024) doi:10.1140/epjc/s10052-024-12622y [arXiv:2307.01100 [hep-ph]].