arXiv:2411.07164v1 [gr-qc] 11 Nov 2024

Gravitational wave propagation beyond General Relativity: geometric optic expansion and lens-induced dispersion

Nicola Menadeo
1, 2, 3, $\ast\,$ and Miguel Zumalacár
regui
3, †

¹Scuola Superiore Meridionale, Largo San Marcellino 10, I-80138, Napoli, Italy

²INFN Sezione di Napoli, Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, I-80126, Napoli, Italy

³Max Planck Institute for Gravitational Physics (Albert Einstein Institute)

Am Mühlenberg 1, D-14476 Potsdam-Golm, Germany

The nature of gravity can be tested by how gravitational waves (GWs) are emitted, detected, and propagate through the universe. Propagation tests are powerful, as small deviations compound over cosmological distances. However, GW propagation tests of theories beyond Einstein's general relativity (GR) are limited by the high degree of symmetry of the average cosmological spacetime. Deviations from homogeneity, *i.e.* gravitational lenses, allow for new interactions, *e.g.*, between standard GW polarization and new scalar or vector fields, with different spin. Therefore, GW lensing beyond GR offers novel tests of cosmological gravity. Here we present the theory of GW propagation beyond GR in the short-wave expansion, including corrections to the leading-order amplitude and phase for the first time. As an example, we compute the dispersive (frequency-dependent) corrections to all metric and scalar field perturbations in Brans-Dicke, the simplest modified theory exhibiting GW dispersion. GW lensing effects are too small to observe in Brans-Dicke theories compatible with solar system and binary pulsar limits. Nevertheless, our formalism opens the possibility of novel tests of gravity, including dark-energy theories and screening mechanisms.

CONTENTS

I.	Introduction	1
II.	General theory for GW propagation	3
	A. Short wave Expansion	3
	1. Geometric optics	4
	2. Beyond geometric optics	4
	3. Fully-luminal theories	5
III.	General Relativity	5
	A. Short-wave expansion	6
	1. Geometric optics	6
	2. Beyond geometric optics	7
	B. Tetrad decomposition	7
	1. Geometric optics	7
	2. Beyond geometric optics	8
IV.	Brans-Dicke	9
	A. Short-wave expansion	10
	1. Geometric optics	10
	2. Beyond geometric optics	11
V.	Point-like lens	12

^{*} nicola.menadeo@aei.mpg.de

A. General relativity	13
B. Massless Brans-Dicke	13
C. Jordan frame	15
VI. Conclusion	16
Acknowledgments	17
A. Weak field limit	17
B. Brans-Dicke background functions	17
C. Functions for bGO corrections	18
1. Results from analytic integration	19
References	20

I. INTRODUCTION

Einstein's General Relativity (GR) has been remarkably successful in describing gravitational phenomena on a wide range of systems, across vastly different scales. However, open questions remain on both extremely small and large scales. Small-scale/high-energy problems of GR include the nature of spacetime singularities and the quantum completion of the theory. On cosmological scales, GR has led to the need to include dark matter and dark energy, accounting for 95% of the universe's

 $^{^{\}dagger}\ miguel.zumalacarregui@aei.mpg.de$

content today [1–4]. In particular dark energy, responsible for cosmic acceleration [5–11], could be interpreted as a breakdown of the attractive nature of gravity on the largest scales. This hypothesis has resulted in a widespread investigation of alternative theories [12–14], as well as an ambitious observational program to test them using cosmological observations [15–23].

The detection of GWs [24] opened a new frontier in the study of gravitational physics, both illuminating the foundations of GR and casting away alternative theories to the shadows. Observed GW signals are emitted by relativistic compact objects, thus probing the regime of strong-field and dynamical gravity [25–27]. In addition to emission, GW propagation across the universe are highly sensitive to other properties of gravity, such as the GW speed [28] and the graviton mass [29]. GW propagation tests probe gravitational interactions directly and can achieve high sensitivity, as anomalous effects accumulate over cosmological distances. Most crucially, they apply directly to theories that modify cosmological dynamics, constraining many dark-energy theories [30–41]. Many GW propagation tests, including measuring the anomalous speed and amplitude, are limited by require an electromagnetic counterpart or otherwise determining the redshift of the source. Moreover, only a handful of effects exist on the homogeneous and isotropic Friedman-Robertson-Walker (FRW) cosmological background [42, 43] that describes the average universe.

GW propagation over inhomogeneous spacetimes, *i.e.* GW lensing, drastically extend the range of phenomena that can be used to test gravity. The basic principle is that lenses/inhomogeneities break the FRW symmetries, allowing interactions between fields of different spin. In alternative theories, this means that the GR standard degrees of freedom (d.o.f.), the $+, \times$ metric polarizations, mix with new fields such as scalars or vectors. In addition to the many gravitational lensing effects in GR [44– 57], lens-induced effects provide a powerful discriminant between theories through novel wave-optic pheonomena. Interactions with new fields involving two derivatives cause lens-induced birefringence (LIB), a difference in speed between the + and \times components of GWs [58]. The absence of birefringence in GW data provided constraints on cosmological theories of gravity comparable to those based on the GW speed [59]. Related birefringent effects are predicted in strong gravitational fields in GR [52, 53, 60], as well as in parity-violating theories [61]. Although progress has been made to characterize GW

lensing beyond GR [58, 62–66], computational difficulties and the wide landscape of theories have prevented a systematic characterization of these phenomena.

One of the ways in which the theory of GW lensing beyond GR needs to be developed is by including frequency-dependent propagation effects. Most analyses rely on the *geometric optics* (GO) approximation [60, 62, 63, 67-71], which holds when the signal's wavelength is significantly smaller than the local curvature scale, the *i.e.* short-wave approximation. In this regime, GWs travel along null geodesics, and their polarization is parallel-transported along them [51, 60, 69]. Corrections beyond geometric optics (bGO) [55, 72–76] include lensinduced dispersion (LID) on GW signals. Such modifications, by the nature of the short-wave approximation, are frequency-dependent and can be probed by GW interferometers. The characterization of the bGO regime beyond GR remains an open problem. A significant step forward was the first explicit computation of bGO corrections in GR [72, 73], showing how bGO effects modify the amplitude and phase of GWs and discussing the emergence of apparent additional scalar polarizations modes, absent in the geometric optics limit. GW lensing beyond GR offers potential for even richer phenomena due to the presence of additional degrees of freecom [77].

In this work, we extend the study of GW lensing beyond GR, developing a formalism that incorporates bGO corrections that describe dispersive phenomena. For simplicity, we focus on the Brans-Dicke theory as an example of scalar-tensor models [78, 79]. Using the short-wave expansion approximation, we compute the bGO corrections to the leading-order scalar observables for tensor and scalar waves passing through a point-like gravitational lens.

The work is organized as follows:

In Sec. II we will introduce a general framework describing the full-propagation of the gravitational and scalar radiation. By employing the short-wave approximation we will show the general equations, order by order in the expansion parameter, the equations governing the geometric optics regime and the first corrections to it. Sec. III will be devoted to review calculations for GR, thus introducing the null-tetrad formalism and thereby presenting the geometric and beyond geometric optics equations and the respective formal solutions for the gravitational radiation. In Sec. IV we will present the full propagation in the BD theory and show the solution in the geometric optics regime of the leading-order scalar and tensor amplitude. Then we will present, for the first time up to our knowledge, the general expression describing beyond geometric optics (bGO) corrections. In Sec. V, we will explicitly evaluate the dispersive bGO corrections, analytically, in the special case of a point-like lens, reviewing the GR case shown in Ref. [73] and subsequently extending to BD.

Notation. Table. I sumarizes the main definitions that we will use throughout the work. Note that we will employ two equivalent descriptions of the theory, the Jordan (JF) and Einstein frame (EF). We will work with $c = \hbar = 1$. Symmetrized and anti-symmetrized indices will be denoted as $T_{(\mu\nu)} \equiv (T_{\mu\nu} + T_{\nu\mu})/2$ and $T_{[\mu\nu]} \equiv (T_{\mu\nu} - T_{\nu\mu})/2$.

	Fields	Amplitude decomposition
GR	$\tilde{h}_{\mu\nu}$, Eqs. (21)	$\tilde{h}^{(n)}_{\mu\nu} \equiv \tilde{\alpha}^{(n)}_{AB} \Theta^{AB}_{\mu\nu}$, Eq. (36)
EF	$(\tilde{h}_{\mu\nu}, \delta\tilde{\phi})$, Eqs. (60)	$\tilde{h}^{(n)}_{\mu\nu} \equiv \tilde{\alpha}^{(n)}_{AB} \Theta^{AB}_{\mu\nu}$, Eq. (36)
JF	$(h_{\mu u},\delta\phi)$	$h^{(n)}_{\mu\nu}\equiv\alpha^{(n)}_{AB}\Theta^{AB}_{\mu\nu}$

TABLE I: Notation used in the work.

II. GENERAL THEORY FOR GW PROPAGATION

In any theory of gravity, the propagation of GWs is determined by the equations of motion (EoM) for linearized perturbations, obtained by expanding around the background metric

$$g_{\mu\nu}^{\text{tot}} = g_{\mu\nu} + h_{\mu\nu}.$$
 (1)

Scalar-tensor theories of gravity include an additional scalar field which can be similarly expanded around the background solution as

$$\phi^{\text{tot}} = \bar{\phi} + \delta\phi. \tag{2}$$

The evolution of both gravitational and scalar waves can be then described by a set of coupled differential equations, which can be compactly written as

$$\mathbf{D}_{IJ}V^J = 0, \tag{3}$$

with

$$\mathbf{D}_{IJ} \equiv \mathcal{K}_{IJ}^{\alpha\beta} \nabla_{\alpha} \nabla_{\beta} + \mathcal{A}_{IJ}^{\alpha} \nabla_{\alpha} + \mathcal{M}_{IJ}, \qquad (4)$$

where ∇_{μ} identifies the covariant derivative compatible with the background metric tensor. $V^{J} \equiv (h_{\mu\nu}, \delta\phi)$ is a vector whose components are the dynamical field perturbations of the metric and the scalar field, after imposing the constraints on the non-dynamical components of $h_{\mu\nu}$, e.g., [58, Sec. V-B]. Here, Latin capital indices denote components of the field perturbation, while Greek indices denote spacetime components.

The propagation equation (4) is naturally split into three parts: the *kinetic matrix*, $\mathcal{K}_{IJ}^{\alpha\beta}$ encodes the secondorder differential operator that acts on field perturbations. Amplitude matrix, $\mathcal{A}_{IJ}^{\alpha}$, contains terms with firstorder derivatives. Finally, the mass matrix, \mathcal{M}_{IJ} , involves the contributions of zeroth-order derivative terms. More precisely, the vector V^J formally has 11 components (the usual 10 components of the metric plus one of the scalar field), and the respective matrices will be 11×11 .

The propagating d.o.f. are determined by diagonalizing the kinetic matrix, and are known as *propagation eigenstates*. The diagonalization is position-dependent and can be quite involved in general. For the scalartensor theories of interest in this work, the *i.e.* BD theory, diagonalization can be performed covariantly [58, 63]. Moreover, the transformation that achieves the diagonalization is equivalent to the well-known mapping between the Jordan and Einstein frames [80–86]. Once the kinetic matrix has been diagonalized, the eigenvectors of the evolving system correspond to the propagating eigenstates. The eigenvalues provide the dispersion relations which determine the propagation speed for each propagating d.o.f. . Hence, one can formally write

$$\mathcal{K}_{IJ}^{\alpha\beta}e^{I} = \mathcal{G}_{I}^{\alpha\beta}e^{I}, \qquad (5)$$

where e^{I} as an eigenstate (or polarization vector) of the kinetic matrix satisfying $e^{I}e^{J} = \delta^{IJ}$ and $\mathcal{G}_{I}^{\alpha\beta}$ is the *effective metric* of the *I-th* propagating eigenstate.

A. Short wave Expansion

As a working hypothesis, we focus on the case in which the GW wavelength λ is much smaller than the scale over which the spacetime varies significantly, $\mathcal{R}_{\text{back}}$. In such conditions, we can perform the short-wave expansion (also known as WKB approximation) by expanding the perturbation fields as

$$V^{I} = e^{i\theta^{I}/\epsilon} \sum_{n} \epsilon^{n} A^{(n)I}, \qquad (6)$$

where $\epsilon \equiv 2\lambda/\mathcal{R}_{\text{back}}$ is a dimensioneless book-keeping parameter and θ^I is the wave's phase related to the prop-

Modified		Geometric optics	eometric optics		
GW lensing	$\mathcal{O}(f^2)$	$\mathcal{O}(f^0)$	$\mathcal{O}(f^{-2})$	Arbitrary f	
Effect	speed	amplitude	phase	all	
Observable	birefringence	oscillations	dispersion	diffraction	
Interactions Tested	2 derivatives	1 derivatives	0 derivatives	all	
Developed in	Refs. [58, 59]	this work (for	r Brans-Dicke)	future work	

TABLE II: Novel effects allowed by lens-induced interactions between h_+, h_{\times} and additional fields. "Interactions tested" refers to the number of derivatives in the couplings between GW polarizations (h_+, h_{\times}) and additional fields: lower (higher) numbers are more generic (restricted).

agating d.o.f. . The quantity $A^{(n)I}$ contains information on both the amplitude and the polarization, and can be splitted as

$$A^{(n)I} \equiv a^{(n)}e^I,\tag{7}$$

where $a^{(n)}$ is a complex amplitude and e^{I} the polarization vector. Moreover, the short-wave expansion naturally defines the scalar and tensor wavevectors, respectively

$$k^I_\mu \equiv \nabla_\mu \theta^I. \tag{8}$$

By inserting the ansatz (6) into Eq. (4) and retaining terms of equal order in ϵ , we obtain the evolution equations for the phase and amplitude in the geometric optics regime, as well as the correction for these quantities in the bGO limit.

1. Geometric optics

The leading and next-to-leading order (NLO) equations in the short-wave expansion describe the geometric optic limit. The leading order equation is

$$\mathcal{G}_{I}^{\alpha\beta}k_{\alpha}^{I}k_{\beta}^{I} = 0.$$
⁽⁹⁾

It determines the evolution of the phase, and the speed of the propagation eigenstates, which in turn determine the theory's causal structure.

The diagonalization procedure (5) might lead to the case in which the resulting eigenvalues exhibit the same dispersion relation for each propagating eigenstate, thereby defining the theory as fully luminal.¹ In this work we choose to focus on fully luminal theories.

The NLO contribution provides a differential equation for the evolution of the leading order amplitude

$$\left[\mathcal{K}_{IJ}^{\alpha\beta}\left(2k_{\alpha}^{J}\nabla_{\beta}+\nabla_{\beta}k_{\alpha}^{J}\right)+\mathcal{A}_{IJ}^{\alpha}k_{\alpha}^{J}\right]A^{(0)J}=0.$$
 (10)

2. Beyond geometric optics

Let us now discuss the NLO corrections to the amplitude. This contribution is given by the equation at order ϵ^0 ,

$$\left[\mathcal{K}_{IJ}^{\alpha\beta}\left(2k_{\alpha}^{J}\nabla_{\beta}+\nabla_{\beta}k_{\alpha}^{J}\right)+\mathcal{A}_{IJ}^{\alpha}k_{\alpha}^{J}\right]A^{(1)J}=i\mathbf{D}_{IJ}A^{(0)J}.$$
(11)

The LHS of the above equation is identical to that of Eq. (10). The RHS, instead, is given by the full propagation equation acting on the leading order quantities. In other words, the failure of the leading order to satisfy the full propagation equation sources the NLO amplitude. Higher-order corrections follow the same equation, by simply substituting $A^{(1)J}$, $A^{(0)J}$ for $A^{(n)J}$, $A^{(n-1)J}$.

The leading bGO term describes a correction to the signal's phase. This interpretation follows from having an imaginary correction, $iA^{(1)J}$, to a real amplitude $A^{(0)J}$, *i.e.* the imaginary unit in Eq. (11). This motivates rewrit-

¹ The term luminal commonly denotes the speed at which a tensor wave propagates, on FRW, is equal to the speed of light.

$$V^{J} = e^{i\theta^{I}/\epsilon} A^{(0)J} \left(1 + i\epsilon \frac{A^{(1)J}}{A^{(0)J}}\right) \approx$$
(12)

$$\approx e^{i\theta^{1}/\epsilon} A^{(0)J} e^{i\epsilon(A^{(1)J}/A^{(0)J})} \tag{13}$$

$$\equiv e^{i\theta^I/\epsilon} A^{(0)J} \exp\left(i\frac{\beta^J}{GMf}\right),\qquad(14)$$

where the second line follows from $\mathcal{A}^{(1)J} \ll \mathcal{A}^{(0)J}$. The last line defines the dimensionless *LID* phasing parameter β^{J} , where the signal frequency f and lens' scale M have been factored out [52, 53].

LID produces a frequency- and polarization-dependent modulation of the waveform. Fig. 1 shows the effect on a typical source for positive/negative phasing, and assuming equal values for both polarizations, $\beta^+ = \beta^{\times}$. Deviations are stronger in the inspiral phase, which corresponds to lower frequencies. The frequency-dependence of LID allows the effect to be tested on any GW signal, without the need of a electromagnetic counterpart. On a given event, the correction is similar to a specific modified dispersion relation, *i.e.* an apparent violation of Lorentz invariance. This deviation from GR is routinely tested on GW catalogs [25–27], and for $\beta^J > 0$ also equivalent to an effective graviton mass [29, 87] (see Ref. [52, Sec. IV-A] for details).

Two key differences allow one to distinguish LID from other deviations from GR. First, a violation of Lorentz invariance or graviton mass is universal: it will appear in all GW events, rather than on the fraction that is affected by a gravitational lens. Second, LID is polarizationdependent in general, $\beta^I \neq \beta^J$. LID effects in GR appear when GWs propagating in strong curvature regions, *i.e.* a massive or intermediate-mass black hole acting as a lens [52–54]. LID due to strong-field GR effects can be distinguished by the detection of multiple images [88, 89] and the fact that the polarization dependence in GR (between the left-right polarized GW components) is suppressed by two additional powers of the frequency [52].

In this work we will focus on gravity theories with a single additional scalar degree of freedom. However the framework can be generalized to any type of theory, regardless of the number and type of degrees of freedom. In the following sections, we will specialize the calculations to GR and BD theory.

3. Fully-luminal theories

Note that Eqs. (10) and (11) can be re-written by employing Eq. (5). Specifically, in certain scalar-tensor theories, the structure of $\mathcal{G}_{I}^{\alpha\beta}$ depends only on the background metric tensor; in this case, using Eq. (9) along with Eq. (5), the resulting eigenvalues yield the same dispersion relation for each propagating eigenstate, thus defining the theory as *fully luminal*. In other words, such a definition states that scalar and tensor waves will propagate at the same speed. To this purpose, Eq. (9) simplifies to

$$g^{\mu\nu}k^{I}_{\mu}k^{I}_{\nu} = 0, \quad \forall I.$$
 (15)

Further, Eqs. (10) and (11), respectively, take the following form

$$\left[2k^J_{\alpha}\nabla^{\alpha} + \nabla^{\alpha}k^J_{\alpha} + \mathcal{A}^{\alpha}_{IJ}k^J_{\alpha}\right]A^{(0)J} = 0, \qquad (16)$$

$$\left[2k^{J}_{\alpha}\nabla^{\alpha} + \nabla^{\alpha}k^{J}_{\alpha} + \mathcal{A}^{\alpha}_{IJ}k^{J}_{\alpha}\right]A^{(1)J} = i\mathbf{D}_{IJ}A^{(0)J}.$$
 (17)

In this work, we choose to focus on fully luminal theories.

III. GENERAL RELATIVITY

Let us start by describing the standard case of GR. We consider the perturbation on the Einstein tensor $G_{\mu\nu} \equiv R_{\mu\nu} - g_{\mu\nu}R/2$ providing the following linearized equation

$$\delta G_{\mu\nu} = \mathsf{K}^{\alpha\beta}{}_{\mu\nu}h_{\alpha\beta} + \mathsf{K}_{\mu\nu}h + \frac{1}{2}g_{\mu\nu}R^{\alpha\beta}h_{\alpha\beta} - \frac{1}{2}h_{\mu\nu}R = 0,$$
(18)

where $h \equiv g^{\alpha\beta}h_{\alpha\beta}$, $R_{\mu\nu}$ and R are the trace, the Ricci tensor and scalar, respectively. Moreover, $\mathsf{K}^{\alpha\beta}_{\mu\nu}$, $\mathsf{K}_{\mu\nu}$ are the component of the kinetic matrix² containing second order covariant derivative contribution

$$\mathsf{K}^{\alpha\beta}{}_{\mu\nu} = -\frac{1}{2}\Box\delta^{\alpha}{}_{\mu}\delta^{\beta}{}_{\nu} + \nabla^{\alpha}\nabla_{(\mu}\delta^{\beta}{}_{\nu)} - \frac{1}{2}g_{\mu\nu}\nabla^{\alpha}\nabla^{\beta}, \quad (19)$$

$$\mathsf{K}_{\mu\nu} = \frac{1}{2}\Box - \frac{1}{2}\nabla_{\nu}\nabla_{\mu}.\tag{20}$$

Following the standard lore, the GWs propagation equation in vacuum is usually simplified by introducing the

 $^{^2}$ We will denote all terms contributing to the kinetic matrix as K: different terms are uniquely specified by the number of spacetime indices, cf. Eqs. (19), (20). A similar notation will be applied to A, M.



FIG. 1: Dispersive effects on GW signals. The plots show the LID corrections for positive (left) and negative (right) values of the phasing parameter β , and assuming that the effect is equal for both polarizations, $\beta^+ = \beta^{\times}$. Deviations are most apparent in the inspiral phase, corresponding to lower frequencies. The signals correspond to a $30 + 30M_{\odot}$, non-spinning quasi-circular binary system.

well-known trace-reversed metric perturbation and imposing the transverse and traceless (TT) gauge conditions on the latter [90, 91], namely

$$\tilde{h}_{\mu\nu} \equiv h_{\mu\nu} - \frac{1}{2}g_{\mu\nu}h, \qquad (21)$$

$$\nabla^{\mu}\tilde{h}_{\mu\nu} = 0, \qquad (22)$$

$$\tilde{h} = 0. \tag{23}$$

By plugging the above relations into Eq. (18) and using the background Einstein field equation, one recovers the usual form of the wave equation

$$\frac{1}{2}\Box\tilde{h}_{\mu\nu} - R_{\alpha\mu\nu\beta}\tilde{h}^{\alpha\beta} = 0, \qquad (24)$$

with $R_{\alpha\mu\nu\beta}$ as the Riemann tensor.

Note that Eq. (24) does not contain the trace h, which is removed by redefining the field as given by Eq. (21). As we will see later, similar redefinitions will help simplify the analysis of GW propagation in theories beyond GR. The introduction of the trace-reversed perturbation metric, indeed, is essentially the field transformation leading to the kinetic matrix diagonalization discussed in Sec. II A, thereby formally decoupling the evolution of the perturbation tensor field and the additional degree of freedom, specifically the trace in this case. Regardless of this, when working in vacuum (without the presence of matter), the trace can always be set to zero [91].

A. Short-wave expansion

Let us now expand the field $h_{\mu\nu}$ using the WKB approximation as

$$\tilde{h}_{\mu\nu} = \left(\tilde{h}_{\mu\nu}^{(0)} + \epsilon \tilde{h}_{\mu\nu}^{(1)} + ...\right) e^{i\theta^T/\epsilon} \,.$$
(25)

Here $\{\tilde{h}_{\mu\nu}^{(0)}, \tilde{h}_{\mu\nu}^{(1)}, ...\}$ are the tensor amplitudes that can be further decomposed into a scalar coefficient and a polarization tensor, and θ^T is the phase related to the tensor wavevector k_{μ}^T , according to Eq. (8). In the following we will denote $k_{\mu}^T \equiv k_{\mu}$ for simplicity.

1. Geometric optics

By plugging the ansatz (25) into Eq. (24) one gets, at the leading order in ϵ

$$k^{\mu}k_{\mu} = 0. (26)$$

This determines that k^{μ} is a null-vector and GWs propagate at the speed of light.

The NLO equation provides

$$\mathcal{D}\tilde{h}^{(0)}_{\mu\nu} = 0, \qquad (27)$$

where

$$\mathcal{D} \equiv \left(2k^{\alpha}\nabla_{\alpha} + \nabla_{\alpha}k^{\alpha}\right),\tag{28}$$

is the differential transport operator along the null ray, tangent to k^{α} . Further, plugging Eq. (25) into the transverse gauge condition (22) and keeping the leading-order terms, one obtains

$$k^{\mu}\tilde{h}^{(0)}_{\mu\nu} = 0.$$
 (29)

This indicates that the polarizations contained in $\tilde{h}^{(0)}_{\mu\nu}$ are orthogonal to the wavevector, and thus to the GWs propagation direction.

2. Beyond geometric optics

The equation describing the first order correction to the geometric optics is

$$\mathcal{D}\tilde{h}^{(1)}_{\mu\nu} = i \left(\Box \tilde{h}^{(0)}_{\mu\nu} - 2R_{\alpha\mu\nu\beta}\tilde{h}^{(0)\alpha\beta} \right).$$
(30)

Moreover, the transverse condition, at such an order, translates to

$$k^{\mu}\tilde{h}^{(1)}_{\mu\nu} = i\nabla^{\mu}\tilde{h}^{(0)}_{\mu\nu}.$$
 (31)

From the aforementioned equation it can be observed that bGO corrections are sourced from the leading-order amplitudes, which implies a deviation from orthogonality between polarizations, encoded in the bGO correction $h_{\mu\nu}^{(1)}$, and the propagation direction.

B. Tetrad decomposition

In general, it is possible to decompose the amplitude in Eq. (25) by introducing a tetrad of null vectors adapted to k^{μ} , where adapted means that every vector is parallely transported along the geodesic associated to k^{μ} . The tetrad basis is given by

$$e^{\mu}_{A} \equiv \{k^{\mu}, m^{\mu}, l^{\mu}, n^{\mu}\}, \qquad (32)$$

where n^{μ} is real, m^{μ} and l^{μ} are complex such that

$$l^{\mu} = \bar{m}^{\mu}, \quad g_{\mu\nu}m^{\mu}l^{\nu} = -g_{\mu\nu}k^{\mu}n^{\nu} = 1, \qquad (33)$$

with all the other contraction vanishing. The dual tetrad basis \hat{e}^{μ}_{A} can be defined as

$$\hat{k}^{\mu} \equiv -n^{\mu}, \ \hat{n}^{\mu} \equiv -k^{\mu}, \ \hat{m}^{\mu} \equiv l^{\mu}, \ \hat{l}^{\mu} \equiv m^{\mu},$$
 (34)

such that $e^{\mu}_{A}\hat{e}^{B}_{\nu} = \delta^{B}_{A}\delta^{\mu}_{\nu}$. Further, the background metric is decomposed as

$$\bar{g}_{\mu\nu} = 2m_{(\mu}l_{\nu)} - 2n_{(\mu}k_{\nu)}.$$
(35)

Generally, a rank-2 symmetric tensor can be decomposed along the null tetrad basis: in such a context, indeed, it is possible to split each tensor amplitude component (at any order n of the short-wave expansion) as

$$\tilde{h}^{(n)}_{\mu\nu} \equiv \tilde{\alpha}^{(n)}_{AB} \Theta^{AB}_{\mu\nu}, \qquad (36)$$

where $\tilde{\alpha}_{AB}^{(n)}$ are the complex coefficients and $\Theta_{\mu\nu}^{AB}$ are the polarization tensors, constructed from the null tetrads and constituting the basis of the decomposition, defined as

$$\Theta^{AB}_{\mu\nu} \equiv \frac{1}{2} \left(e^{A}_{\mu} e^{B}_{\nu} + e^{A}_{\nu} e^{B}_{\mu} \right).$$
(37)

The expression in Eq. (36) can be simplified by constraining some of the expansion coefficients using the TT gauge conditions.

1. Geometric optics

Specifically, by decomposing the leading order tensor amplitude $\tilde{h}^{(0)}_{\mu\nu}$ according to Eq. (36) and plugging it into the leading order transversality condition (29), one can readily verify that

$$\tilde{\alpha}_{nk}^{(0)} = \tilde{\alpha}_{nn}^{(0)} = \tilde{\alpha}_{nm}^{(0)} = \tilde{\alpha}_{nl}^{(0)} = 0.$$
(38)

The traceless condition (23), further provides

$$\tilde{\alpha}_{ml}^{(0)} = 0. \tag{39}$$

Still, we retain the freedom to fully fix the gauge. It is reasonable to expect that the spacetime is asymptotically flat, as the effect from the inhomogeneity is only localized around the lens and does not extend to infinity. Consequently, one can transform the amplitude tensor to ensure the equation remains fully gauge invariant [72], as follows

$$\tilde{h}_{\mu\nu} \to \tilde{h}_{\mu\nu} + 2C_{(\mu}k_{\nu)}, \qquad (40)$$

where C_{μ} is an arbitrary complex vector orthogonal to k_{μ} .

Such a requirement is satisfied by imposing $n^{\mu}\tilde{h}^{(0)}_{\mu\nu} = 0$, which, along with Eq. (36), provides

$$\tilde{\alpha}_{kk}^{(0)} = \tilde{\alpha}_{km}^{(0)} = \tilde{\alpha}_{kl}^{(0)} = 0.$$
(41)

After this process, the only unconstrained contributions are $\tilde{\alpha}_{mm}^{(0)}$ and $\tilde{\alpha}_{ll}^{(0)}$, thus yielding

$$\tilde{h}^{(0)}_{\mu\nu} = \tilde{\alpha}^{(0)}_{mm} m_{\mu} m_{\nu} + \tilde{\alpha}^{(0)}_{ll} l_{\mu} l_{\nu}.$$
(42)

These coefficients are the amplitudes associated with the two independent left-handed and right-handed helicity polarizations, typical for a massless spin-2 field, expressed in the null tetrad basis. The latter can be directly related to the usual GW amplitude polarizations h_+ and h_{\times} .

It is possible to express the above equation in terms of the usual tetrad basis [92, 93], upon which the standard GWs polarizations h_+ and h_{\times} are defined, by performing a transformation on the tetrads

$$\varepsilon_{\mu\nu}^{+} \equiv \frac{m_{\mu}m_{\nu} + l_{\mu}l_{\nu}}{\sqrt{2}},\tag{43}$$

$$\varepsilon_{\mu\nu}^{\times} \equiv i \frac{m_{\mu}m_{\nu} - l_{\mu}l_{\nu}}{\sqrt{2}},\tag{44}$$

where $\varepsilon_{\mu\nu}^{\pm}$ are the usual polarization tensors. Simultaneously, the coefficients change as follows

$$\tilde{\alpha}_{mm}^{(0)} \equiv \frac{\tilde{h}_{+}^{(0)} - i\tilde{h}_{\times}^{(0)}}{\sqrt{2}},\tag{45}$$

$$\tilde{\alpha}_{ll}^{(0)} \equiv \frac{\tilde{h}_{+}^{(0)} + i\tilde{h}_{\times}^{(0)}}{\sqrt{2}}.$$
(46)

The evolution of $\tilde{\alpha}_{mm}^{(0)}$ and $\tilde{\alpha}_{ll}^{(0)}$ is obtained by inserting Eq. (42) into Eq. (27) and subsequently projecting along the dual tetrad basis $\hat{e}_{A}^{\mu} \hat{e}_{B}^{\nu}$, yielding a scalar equation

$$\mathcal{D}\left(\tilde{\alpha}_{mm}^{(0)}\right) = \mathcal{D}\left(\tilde{\alpha}_{ll}^{(0)}\right) = 0.$$
(47)

By recalling $k^{\mu}\nabla_{\mu} \equiv d/d\xi$ as the directional derivative along k^{μ} , with ξ being an affine parameter, and using the relation $\nabla_{\mu}k^{\mu} = 2d \ln D/d\xi$ where D is the comoving distance along the geodesic (see Refs. [46, 94, 95] for a complete discussion on optical scalars), one can express Eq. (47) as a first order differential equation in the affine parameter

$$\frac{d}{d\xi} \left[\tilde{\alpha}_{\rm o}^{(0)} D \right] = 0. \tag{48}$$

The aforementioned equation can be easily integrated from the source point, labeled as ξ_s , to a generic point ξ (*i.e.* the observation point ξ_o) on the geodesic, thus providing

$$\tilde{\alpha}_{\circ}^{(0)}(\xi) = \frac{D(\xi_s)}{D(\xi)} \tilde{\alpha}_{\circ}^{(0)}(\xi_s), \qquad (49)$$

where we can observe the standard evolution of the amplitudes decaying with increasing distance. By plugging the above solution into Eq. (42), one can finally obtain the evolution, in the affine parameter, of the leading order tensor amplitude

$$\tilde{h}^{(0)}_{\mu\nu}(\xi) = \frac{D(\xi_s)}{D(\xi)} \left(\tilde{\alpha}^{(0)}_{mm}(\xi_s) m_{\mu} m_{\nu} + \tilde{\alpha}^{(0)}_{ll}(\xi_s) l_{\mu} l_{\nu} \right).$$
(50)

2. Beyond geometric optics

The evolution for the bGO coefficients $\tilde{\alpha}_{AB}^{(1)}$ of the tensor amplitude $h_{\mu\nu}^{(1)}$ (Eq. (36)) can be computed with a similar approach. An important difference is that some steps in the computation of $\tilde{h}_{\mu\nu}^{(0)}$ do not apply to $h_{\mu\nu}^{(1)}$. Specifically, the transversality gauge condition in the bGO regime is determined by Eq. (31), which accounts for an orthogonality deviation between the polarizations of $h_{\mu\nu}^{(1)}$ and the propagation direction. By inserting the decomposition given by Eq. (36) for $h_{\mu\nu}^{(1)}$ into Eq. (31), one gets

$$2\tilde{\alpha}_{nn}^{(1)}n_{\mu} + \tilde{\alpha}_{nl}^{(1)}l_{\mu} + \tilde{\alpha}_{mn}^{(1)}m_{\mu} + \tilde{\alpha}_{nk}^{(1)}k_{\mu} = -2i\nabla^{\nu}\tilde{h}_{\mu\nu}^{(0)},$$
(51)

which means that the above coefficients cannot be straightforwardly constrained to zero as it happens in Eq. (38) for $\tilde{h}^{(0)}_{\mu\nu}$. Moreover, the residual gauge condition $n^{\mu}\tilde{h}^{(1)}_{\mu\nu} = 0$ and traceless one, $\tilde{h} = 0$, provide

$$\tilde{\alpha}_{kk}^{(1)} = \tilde{\alpha}_{km}^{(1)} = \tilde{\alpha}_{kl}^{(1)} = \tilde{\alpha}_{kn}^{(1)} = \tilde{\alpha}_{ml}^{(1)} = 0, \qquad (52)$$

ending up with

$$\tilde{h}^{(1)}_{\mu\nu} = \tilde{\alpha}^{(1)}_{mm} m_{\mu} m_{\nu} + \tilde{\alpha}^{(1)}_{ll} l_{\mu} l_{\nu} + \tilde{\alpha}^{(1)}_{nn} n_{\mu} n_{\nu} + \\ + \tilde{\alpha}^{(1)}_{nm} n_{(\mu} m_{\nu)} + \tilde{\alpha}^{(1)}_{nl} n_{(\mu} l_{\nu)}.$$
(53)

The coefficients $\tilde{\alpha}_{nn}^{(1)}$, $\tilde{\alpha}_{nm}^{(1)}$, and $\tilde{\alpha}_{nl}^{(1)}$ are not constrained by the gauge conditions, unlike the leading-order amplitude in Eq. (38). This suggests these non-vanishing elements due to bGO/dispersive effects appear as novel polarizations spreading the gravitational radiation.

The evolution of the remaining $\tilde{\alpha}_{AB}^{(1)}$ coefficients is derived by inserting Eq. (36) into Eq. (30) and projecting along the dual tetrad basis $\hat{e}_{A}^{\mu}\hat{e}_{B}^{\nu}$, it is possible to derive and subsequently integrate the equation for the $\tilde{\alpha}_{AB}^{(1)}$ modes contributing to the leading order correction in the bGO regime, thus obtaining

$$\tilde{\alpha}_{AB}^{(1)}(\xi) = \frac{D(\xi_s)}{D(\xi)} \tilde{\alpha}_{AB}^{(1)}(\xi_s) + \frac{i}{D(\xi)} \int_{\xi_s}^{\xi} d\xi \hat{e}_A^{\mu} \hat{e}_B^{\nu} D(\xi) \left[\Box \tilde{h}_{\mu\nu}^{(0)} - 2R_{\alpha\mu\nu\beta} \tilde{h}^{(0)\alpha\beta} \right]$$
(54)

The first term describes the usual behavior of the mode propagating as 1/D, while the second contribution encodes the dispersive, frequency-dependent corrections (as it will be explicitly evaluated later in Sec. V). It is sourced by the background curvature, which can induce novel effects in the polarization tensor along the geodesic path. Further, the imaginary unit preceding the above integral indicates that the bGO corrections represent a modification of the GWs phase, rather than the (real-valued) leading-order amplitude (Eq. (14)). This causes GW dispersive phenomena, *i.e. spectrum frequency decomposition* analog to the phenomenon observed when a light ray passes through an optical prism.

IV. BRANS-DICKE

The BD theory of gravity is arguably the simplest among scalar-tensor theories, featuring a scalar field nonminimally coupled to the Ricci scalar. Assuming no interactions with any other matter fields, the theory is described by the action

$$S = \frac{1}{16\pi G} \int d^4x \sqrt{-g} \left(\phi R - \frac{\omega}{\phi} \phi_\mu \phi^\mu \right) + \int d^4x \sqrt{-g} \mathcal{L}_m,$$
(55)

where g is the determinant of the metric tensor $g_{\mu\nu}$, \mathcal{L}_m the matter Lagrangian, ϕ the scalar field coupled to gravity and ω the Brans-Dicke constant. Further, $\phi_{\mu} \equiv \nabla_{\mu}\phi$. The field equations are easily obtained by varying the action w.r.t. the metric tensor and the scalar field, thus yielding

$$\phi\left(R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R\right) - \frac{\omega}{\phi}\phi_{\mu}\phi_{\nu} - \phi_{\mu\nu} + g_{\mu\nu}\left(\Box\phi + \frac{\omega}{2}\phi_{\alpha}\phi^{\alpha}\right) = 8\pi G T_{\mu\nu},$$
(56)

$$R + \frac{2\omega}{\phi} \Box \phi - \frac{\omega}{\phi^2} \phi_\alpha \phi^\alpha = 0, \qquad (57)$$

where

$$T_{\mu\nu} \equiv \frac{-2}{\sqrt{-g}} \frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g^{\mu\nu}},\tag{58}$$

is the energy-momentum tensor and $T \equiv g^{\mu\nu}T_{\mu\nu}$ its trace [84]. By plugging Eq. (56) and its contraction into Eq. (57), the scalar field equation simply reduces to

$$\Box \phi = \frac{8\pi G}{3+2\omega} T. \tag{59}$$

In this work we will specialize to vacuum propagation $T_{\mu\nu} = 0.$

The study of GWs propagation can be performed by perturbing the background metric and scalar field, as shown in Eqs. (1) and (2) and linearizing the above field equations, thus finding a coupled system of second-order differential equations governing the dynamics of the fields $(h_{\mu\nu}, \delta\phi)$. In such a context, the diagonalization process of the kinetic matrix is feasible and can be performed covariantly by re-defining the metric perturbation as

$$\tilde{h}_{\alpha\beta} \equiv h_{\mu\nu} - \frac{1}{2}g_{\mu\nu}h - g_{\mu\nu}\frac{\delta\phi}{\bar{\phi}},\tag{60}$$

$$\delta \tilde{\phi} = \delta \phi. \tag{61}$$

This redefinition is the linearized version of Einsteinframe metric, in which the scalar couples to matter, rather than curvature.

Moreover, having thus mapped the theory to the Einstein frame with this transformation, the TT gauge conditions, on $\tilde{h}_{\mu\nu}$, hold (See Ref. [96]). The diagonalized propagation equation of perturbation fields in the Brans-Dicke theory, has the following structure

$$\begin{bmatrix} \left(\tilde{\mathsf{K}}_{\mu\nu}{}^{\alpha\beta\gamma\rho} & 0 \\ 0 & \tilde{\mathsf{K}}^{\gamma\rho} \right) \nabla_{\gamma}\nabla_{\rho} + \left(\tilde{\mathsf{A}}_{\mu\nu}{}^{\alpha\beta\gamma} & \tilde{\mathsf{A}}_{\mu\nu}{}^{\gamma} \\ 0 & \tilde{\mathsf{A}}^{\gamma} \end{bmatrix} \nabla_{\gamma} + \left(\tilde{\mathsf{M}}_{\mu\nu}{}^{\alpha\beta} & \tilde{\mathsf{M}}_{\mu\nu}{}^{h} \\ \tilde{\mathsf{M}}_{\phi}{}^{\alpha\beta} & \tilde{\mathsf{M}} \end{bmatrix} \end{bmatrix} \begin{pmatrix} \tilde{h}_{\alpha\beta} \\ \delta\phi \end{bmatrix} = 0.$$
(62)

We present the full expressions for the different K, A, M

tensors in Appendix B.

A. Short-wave expansion

Similarly to the short-wave metric expansion, Eq. (25), the perturbation of the scalar field can be also expanded as

$$\delta\phi \equiv \left(\delta\phi^{(0)} + \epsilon\delta\phi^{(1)} + \dots\right)e^{i\theta^S/\epsilon},\tag{63}$$

where $\{\delta\phi^{(0)}, \delta\phi^{(1)}, ...\}$ are a set of scalar amplitudes and θ^S is the scalar phase. One can define k_{μ}^S as the wavevector related to θ^S following Eq. (8). Recalling the discussion in Sec. II A, BD theory is classified as a fully luminal theory, which means that the scalar and tensor sectors share the same causal structure; once the gauge is fixed, the components of the diagonalized kinetic matrix, *i.e.* $\tilde{K}_{\mu\nu}{}^{\alpha\beta\gamma\rho}$ and $\tilde{K}^{\gamma\rho}$, are built only using the metric tensor. This leads to the same dispersion relation for scalar and tensor waves, resulting in the same propagation speed for all propagating eigenstates. As a consequence, the scalar and tenefore we can set $\theta^T = \theta^S = \theta$, implying $k_{\mu}^T = k_{\mu}^S = k_{\mu}$.

In the following paragraphs, by inserting Eqs. (25) and (63) into Eq. (62), we will analyze in detail the equations for both the scalar and tensor sectors, order by order in ϵ .

1. Geometric optics

Let us start by studying the scalar wave propagation equation at the leading and next-to-leading order in ϵ . The relation at leading order is

$$\tilde{\mathsf{K}}^{\alpha\beta}k_{\alpha}k_{\beta} = k_{\alpha}k^{\alpha} = 0, \qquad (64)$$

where in the second equality we used (B2). As discussed in Sec III, Eq. (64) ensures that k^{μ} is a null vector propagating at the speed of light.

The NLO contribution describes the evolution of the leading order scalar amplitude and provides a differential equation for $\delta \phi^{(0)}$

$$\left[\tilde{\mathsf{K}}^{\alpha\beta}\left(2k_{\alpha}\nabla_{\beta}+\nabla_{\alpha}k_{\beta}\right)-k_{\alpha}\tilde{\mathsf{A}}^{\alpha}\right]\delta\phi^{(0)}=0.$$
 (65)

Using Eqs. (B2) and (B5), the above equation can be re-written in a more compact form

$$\mathcal{D}\left(\frac{\delta\phi^{(0)}}{\sqrt{\phi}}\right) = 0. \tag{66}$$

Following the same procedure shown in Sec. III B, the latter can be expressed as

$$\frac{d}{d\xi} \left(\frac{\delta \phi^{(0)} D}{\sqrt{\phi}} \right) = 0, \tag{67}$$

which, in turn, can be easily integrated from the source of the wave ξ_s to a generic point ξ , thus providing

$$\delta\phi^{(0)}(\xi) = \delta\phi^{(0)}(\xi_s) \frac{D(\xi_s)}{\sqrt{\phi}(\xi_s)} \left(\frac{\sqrt{\phi}(\xi)}{D(\xi)}\right).$$
(68)

Let us now describe the geometric optics regime for the gravitational sector. The leading-order equation provides the usual dispersion relation $k_{\mu}k^{\mu} = 0$.

The equation describing the evolution of the leading order tensor amplitude has a richer structure

$$\begin{bmatrix} \mathsf{K}_{\mu\nu}{}^{\rho\sigma\alpha\beta}(2k_{\alpha}\nabla_{\beta}+\nabla_{\alpha}k_{\alpha})-2k_{\alpha}\tilde{\mathsf{A}}_{\mu\nu}{}^{\alpha\beta\gamma}\delta^{\rho}_{\beta}\delta^{\sigma}_{\gamma}\end{bmatrix}\tilde{h}^{(0)}_{\rho\sigma} = 2\tilde{\mathsf{A}}_{\mu\nu}{}^{\alpha}k_{\alpha}\delta\phi^{(0)}.$$
(69)

By plugging the tensor decomposition (36) into the above relation along with Eqs. (B1), (B3) and (B4), and subesequently contracting the result with $\hat{e}^{\mu}_{A}\hat{e}^{\nu}_{B}$, one easily obtains

$$\mathcal{D}\left(\sqrt{\bar{\phi}}\tilde{\alpha}^{(0)}_{AB}\right) = \frac{d}{d\xi} \left[\sqrt{\bar{\phi}}D\tilde{\alpha}^{(0)}_{AB}\right] = 0.$$
(70)

It is worth stressing that each amplitude evolves independently and the evolution can be obtained by integrating from the source of the waves to a generic point, ξ , thus yielding

$$\tilde{\alpha}_{AB}^{(0)}(\xi) = \tilde{\alpha}_{AB}^{(0)}(\xi_s) \frac{\sqrt{\bar{\phi}(\xi_s)}D(\xi_s)}{\sqrt{\bar{\phi}(\xi)}D(\xi)}.$$
(71)

Being the TT gauge conditions valid in such a theory, the constraints on the coefficients $\tilde{\alpha}_{AB}^{(0)}$ and $\tilde{\alpha}_{AB}^{(1)}$ are identically the same as those discussed for GR in Sec. III B: in the GO limit only the mm and ll components are non-zero, cf. Eqs. (42) and (53) hold. Specifically, by inserting the solution (71) in Eq. (42), one gets the evolution of the leading order tensor amplitude for the Brans-Dicke theory

$$\tilde{h}_{\mu\nu}^{(0)}(\xi) = \frac{\sqrt{\bar{\phi}(\xi_s)}D(\xi_s)}{\sqrt{\bar{\phi}(\xi)}D(\xi)} \left[\tilde{\alpha}_{mm}^{(0)}(\xi_s)m_{\mu}m_{\nu} + \tilde{\alpha}_{ll}^{(0)}(\xi_s)l_{\mu}l_{\nu} \right].$$
(72)

The result is similar to that of GR, where the leadingorder modes evolve inversely with the distance. However, one can observe a novel modulation factor, $\sqrt{\phi(\xi)}$, due to the presence of the scalar field. This is related to the background scalar value modulating the effective Planck's mass, and hence the amplitude of GWs. These results are in complete agreement with Refs. [62, 63, 67].

2. Beyond geometric optics

The bGO equations are significantly more intricate than that of GR due to additional terms and interactions between the scalar and tensor sectors. This differences lead to novel effects and require a detailed treatment.

For the scalar wave, the NNLO equation captures the

$$\begin{bmatrix} \tilde{\mathsf{K}}^{\alpha\beta} \left(2k_{\alpha} \nabla_{\beta} + \nabla_{\beta} k_{\alpha} \right) - k_{\gamma} \tilde{\mathsf{A}}^{\gamma} \end{bmatrix} \delta \phi^{(1)} = \\ = i \left[\left(\tilde{\mathsf{K}}^{\alpha\beta} \nabla_{\alpha} \nabla_{\beta} + \tilde{\mathsf{A}}^{\gamma} \nabla_{\gamma} + \tilde{\mathsf{M}} \right) \delta \phi^{(0)} + \tilde{\mathsf{M}}^{\alpha\beta}_{\phi} \tilde{h}^{(0)}_{\alpha\beta} \right].$$
(73)

The LHS of the above equation has the same structure of the NLO propagation, Eq. (65), and can be re-written as Eq. (66). The main difference is that the RHS introduces a source term, given by the full propagation equation acting on the leading-order scalar amplitude. In other words, the bGO correction is sourced by the failure of the GO prediction to satisfy the full equation, as expected from the general framework discussed in Sec II A. The general solution for $\delta \phi^{(1)}$ can therefore be expressed as

$$\delta\phi^{(1)}(\xi) = \delta\phi^{(1)}(\xi_s) \frac{D(\xi_s)}{\sqrt{\phi}(\xi_s)} \left(\frac{\sqrt{\phi}(\xi)}{D(\xi)}\right) + i\frac{\sqrt{\phi}(\xi)}{D(\xi)} \int_{\xi_s}^{\xi} d\xi \frac{D(\xi)}{\bar{\phi}^{3/2}} \left[\left(\tilde{\mathsf{K}}^{\alpha\beta}\nabla_{\alpha}\nabla_{\beta} + \tilde{\mathsf{A}}^{\gamma}\nabla_{\gamma} + \tilde{\mathsf{M}}\right) \delta\phi^{(0)} + \tilde{\mathsf{M}}^{\alpha\beta}_{\phi}\tilde{h}^{(0)}_{\alpha\beta} \right]. \tag{74}$$

The result clearly shows how tensor polarizations interact with the scalar sector due to mass-like terms with zero derivatives, $\tilde{M}^{\alpha\beta}\tilde{h}^{(0)}_{\alpha\beta}$. These interactions will source the scalar polarization, even if only tensor modes were emitted.

The tensor equation, at order ϵ^0 reads

$$\begin{bmatrix} \mathsf{K}_{\mu\nu}{}^{\rho\sigma\alpha\beta}(2k_{\alpha}\nabla_{\beta}+\nabla_{\alpha}k_{\alpha})-2k_{\alpha}\tilde{\mathsf{A}}_{\mu\nu}{}^{\alpha\beta\gamma}\delta^{\rho}_{\beta}\delta^{\sigma}_{\gamma}\end{bmatrix}\tilde{h}^{(1)}_{\rho\sigma}=i\mathsf{F}_{\mu\nu}$$
(75)

where

$$\mathsf{F}_{\mu\nu} \equiv \tilde{\mathsf{M}}^{h}_{\mu\nu} \delta \phi^{(0)} - k_{\gamma} \tilde{\mathsf{M}}_{\mu\nu}{}^{\gamma} \delta \phi^{(1)} + \\ + \left[-\frac{1}{2} \tilde{\mathsf{K}}_{\mu\nu}{}^{\rho\sigma\alpha\beta} \nabla_{\alpha} \nabla_{\beta} + \delta^{\rho}_{\alpha} \delta^{\sigma}_{\beta} \tilde{\mathsf{A}}_{\mu\nu}{}^{\alpha\beta\gamma} \nabla_{\gamma} + \tilde{\mathsf{M}}_{\mu\nu}{}^{\rho\sigma} \right] \tilde{h}^{(0)}_{\rho\sigma}$$

$$(76)$$

The structure of the LHS of Eq. (75) is identical to that of the GO, Eq. (69), and can be re-written as Eq. (70). Further, the evolution of the coefficients $\alpha_{AB}^{(1)}$, is obtained by plugging Eq. (36) into Eq. (75) and projecting the result along the dual tetrads basis $\hat{e}_{A}^{\mu}\hat{e}_{B}^{\nu}$, leading to

$$\frac{d}{d\xi} \left(\tilde{\alpha}_{AB}^{(1)} D \sqrt{\bar{\phi}} \right) = i D(\xi) \hat{e}_A^{\mu} \hat{e}_B^{\nu} \mathsf{F}_{\mu\nu}. \tag{77}$$

By integrating the above equation, one obtains

$$\tilde{\alpha}_{AB}^{(1)}(\xi) = \tilde{\alpha}_{AB}^{(1)}(\xi_s) \frac{\sqrt{\bar{\phi}(\xi_s)}D(\xi_s)}{\sqrt{\bar{\phi}(\xi)}D(\xi)} + \frac{i}{\sqrt{\bar{\phi}(\xi)}D(\xi)} \int_{\xi_s}^{\xi} d\xi' \frac{D(\xi')}{\sqrt{\bar{\phi}}} \left(\hat{e}_A^{\mu}\hat{e}_B^{\nu}\mathsf{F}_{\mu\nu}\right),$$
(78)

Note that the F tensor includes both leading order metric amplitudes and scalar field perturbations, cf. Eq. (76).

At this point, we can set without loss of generality $\tilde{\alpha}_{AB}^{(1)}(\xi_s) = \tilde{\alpha}_{AB}^{(1)}(\xi_o) = \delta \phi^{(1)}(\xi_s) = \delta \phi^{(1)}(\xi_o) = 0$. It is also reasonable to assume that bGO corrections triggering novel (either apparent or additional) polarizations forbidden in the GO limit do not contribute either at the source or at the observation point, as the lens-induced effects are entirely negligible asymptotically.

In any theory of gravity, mass-like terms, which involve zero-order derivatives, inevitably contribute to wave propagation, making them universal features regardless of the specific framework. In the bGO regime, in particular, the propagating perturbation fields gain a dispersive nature, thus leading to frequency-dependent modifications in the GW signal. Such dispersive effects introduce modifications to both the phase and amplitude of the signal, thus producing corrections that vary with frequency and providing a potential signature that may be detectable across cosmological distances. Importantly, the interactions between the scalar and tensor sectors rely on the presence of an inhomogeneous background; if the propagation medium were homogeneous, such as in a cosmological setting, the interactions would effectively vanish. Consequently, the presence of a gravitational lens or a similar inhomogeneous structure is necessary for these effects to manifest.

V. POINT-LIKE LENS

In this section, we apply the formalism to the point-like lens case, discussing the bGO dispersive corrections to the scalar and tensor amplitudes in Brans-Dicke theory. In presence of a point-like lens, the line element reads

$$ds^{2} = -(1+2\Psi)dt^{2} + (1-2\Psi)(dx^{2} + dy^{2} + dz^{2}), \quad (79)$$

where $\Psi \equiv -R_s/2R$ is the gravitational potential such that $\Psi \ll 1$, $R_s \equiv 2GM_L$ is the Schwarzschild radius with M_L as the lens' mass and $R \equiv \sqrt{x^2 + y^2 + z^2}$ the radial distance from the origin of the coordinate system, in which the lens is located (See Fig. 2). We will follow the perturbative approach, up to linear order in Ψ , presented in Ref. [73]. The first step is to expand the tensor amplitude (36) to the first order in the gravitational potential, yielding

$$\tilde{h}^{(n)}_{\mu\nu} = \bar{\tilde{\alpha}}^{(n)}_{AB} \bar{e}^{A}_{(\mu} \bar{e}^{B}_{\nu)} + 2\bar{\tilde{\alpha}}^{(n)}_{AB} \delta e^{A}_{(\mu} e^{B}_{\nu)} + \delta \tilde{\alpha}^{(n)}_{AB} \bar{e}^{A}_{(\mu} \bar{e}^{B}_{\nu)}, \quad (80)$$

where the bar identifies the background quantities constructed on the Minkowski metric, and the variations encode the effects of the gravitational potential characterizing the lens.

We will consider propagation along the \hat{z} -axis in the (x-z)-plane, *i.e.* y = 0. Specifically, given that $\Psi \ll 1$, the expected deflection angle between the lens-perturbed trajectory. Therefore, we can compute the deflection as leading-order corrections to the unperturbed trajectory (Born approximation) and choose the affine parameter to be $d\xi = dz/\Omega$, where $\Omega \equiv 2\pi/\lambda$ is a constant amplitude for the 4-vector momentum and λ is the signal's wavelenght (see Refs. [72, 73]). The background tetrads can be constructed to be constants

$$\bar{k}^{\mu} \equiv \Omega(1,0,0,1), \tag{81}$$

$$\bar{n}^{\mu} \equiv \frac{1}{2\Omega} (1, 0, 0, -1),$$
 (82)

$$\bar{m}^{\mu} \equiv \frac{1}{\sqrt{2}}(0, 1, i, 0),$$
(83)

$$\bar{l}^{\mu} \equiv \frac{1}{\sqrt{2}}(0, 1, -i, 0).$$
(84)

Further, it is also straightforward to verify that this construction fulfills all the properties highlighted in Sec. III B. The perturbation to the tedrads at the first order in Ψ can be explicitly evaluated by integrating the linearized geodesic equation given by [97]

$$\delta e^{\mu}_{A} = -\int_{z_{s}}^{z_{o}} \frac{dz}{\Omega} \delta \Gamma^{\mu}_{\alpha\beta} \bar{e}^{\alpha}_{A} \bar{k}^{\beta} = -\int_{-\infty}^{+\infty} \frac{dz}{\Omega} \delta \Gamma^{\mu}_{\alpha\beta} \bar{e}^{\alpha}_{A} \bar{k}^{\beta},$$
(85)



FIG. 2: Schematic visualization of the (x, z) plane illustrating the propagation process along the z-axis of scalar and tensor waves. In the presence of a lens located at the origin (black star) and a background scalar field around it (ideally represented by a halo around the lens), the waves trajectory is weakly deflected (solid purple line). The points z_s and z_o identify the source and observer, respectively. In our case of interest, $z_s \to -\infty$ and $z_o \to +\infty$.

where, in the second equality, we restricted to the special case in which the lens is far away from both the source and the observer $z_s \to -\infty$, $z_o \to +\infty$. Here $\delta \Gamma^{\mu}_{\ \alpha\beta}$ are the linerized Christoffel symbols, evaluated using Eq. (79) (whose expressions are shown in Appendix A), up to the first order in Ψ . The subscripts *s* and *o* denote the points in which the source and the observer are located, respectively. By plugging Eqs. (81–84) into the above integral, one gets

$$k^{\mu} = \bar{k}^{\mu} + \delta k^{\mu} = \Omega\left(1, -\frac{2R_s}{b}, -\frac{2R_s}{b}, 1\right), \quad (86)$$

$$n^{\mu} = \bar{n}^{\mu} + \delta n^{\mu} = \frac{1}{2\Omega} \left(1, 0, 0, -1 \right), \tag{87}$$

$$m^{\mu} = \bar{m}^{\mu} + \delta m^{\mu} = \frac{1}{\sqrt{2}} \left(-\frac{R_s}{b}, 1, i, \frac{R_s}{b} \right), \qquad (88)$$

$$l^{\mu} = \bar{l}^{\mu} + \delta l^{\mu} = \frac{1}{\sqrt{2}} \left(-\frac{R_s}{b}, 1, -i, \frac{R_s}{b} \right),$$
(89)

where $b \equiv \sqrt{x^2 + z^2}$ is the impact parameter.

A. General relativity

Let us first analyze how the GO amplitude (50) is modified at first order in Ψ . In particular, one has

$$\tilde{h}_{\mu\nu}^{(0)}(z_{o}) = \frac{\bar{D}(z_{s})}{\bar{D}(z_{o})} \left[\tilde{\alpha}_{mm}^{(0)}(z_{s})\bar{m}_{\mu}\bar{m}_{\nu} + \tilde{\alpha}_{ll}^{(0)}(z_{s})\bar{l}_{\mu}\bar{l}_{\nu} \right] \Delta(z_{o}) + \frac{\bar{D}(z_{s})}{\bar{D}(z_{o})} \left[2\tilde{\alpha}_{mm}^{(0)}(z_{s})\bar{m}_{(\mu}\delta m_{\nu)} + 2\tilde{\alpha}_{ll}^{(0)}(z_{s})\bar{l}_{(\mu}\delta l_{\nu)} \right],$$
(90)

with

$$\Delta(z_o) \equiv \left(1 - \frac{\delta D(z_o)}{\bar{D}(z_o)}\right),\tag{91}$$

where δD represents the first-order perturbation in the comoving distance along the geodesic [98]; the latter can be neglected at the first order in Ψ (as shown in Ref. [73], Appendix A). Moreover, for simplicity, we assumed a Euclidean distance parameterized as $\bar{D}(z) = z - z_s$. Recalling Eq. (80), the correction to the tensor amplitude $\tilde{h}^{(1)}_{\mu\nu}$, at the linear order in Ψ , is

$$\tilde{h}^{(1)}_{\mu\nu}(z_o) = \delta \tilde{\alpha}^{(1)}_{nm}(z_o) \bar{n}_{(\mu} \bar{m}_{\nu)} + \delta \tilde{\alpha}^{(1)}_{nl}(z_o) \bar{n}_{(\mu} \bar{l}_{\nu)} + \\
+ \delta \tilde{\alpha}^{(1)}_{mm}(z_o) \bar{m}_{\mu} \bar{m}_{\nu} + \delta \tilde{\alpha}^{(1)}_{ll}(z_o) \bar{l}_{\mu} \bar{l}_{\nu} + \\
+ \delta \tilde{\alpha}^{(1)}_{nn}(z_o) \bar{n}_{\mu} \bar{n}_{\nu}.$$
(92)

In GR, the only coefficient surviving is $\delta \tilde{\alpha}_{nn}^{(1)}$, discussed in Ref. [73], thus providing

$$\tilde{h}^{(1)}_{\mu\nu}(z_o) = -i \frac{D(z_s)}{\bar{D}(z_o)} \frac{4\Omega R_s}{b^2} \tilde{\zeta}^{(0)}_+(z_s) \bar{n}_{\mu} \bar{n}_{\nu}, \qquad (93)$$

where we defined the quantities

$$\tilde{\zeta}_{+}^{(0)}(z_s) \equiv \tilde{\alpha}_{mm}^{(0)}(z_s) + \tilde{\alpha}_{ll}^{(0)}(z_s), \tag{94}$$

$$\tilde{\zeta}_{-}^{(0)}(z_s) \equiv \tilde{\alpha}_{mm}^{(0)}(z_s) - \tilde{\alpha}_{ll}^{(0)}(z_s).$$
(95)

The latter are nothing but symmetric and antisymmetric combinations of the left- and right-handed modes contributing to the GO tensor amplitude. The above definitions can be further related to standard GW polarization via Eqs. (45) and (46), thus yielding

$$\tilde{\zeta}_{+}^{(0)}(z_s) = \sqrt{2}\tilde{h}_{+}^{(0)}(z_s), \qquad (96)$$

$$\tilde{\zeta}_{-}^{(0)}(z_s) = -i\sqrt{2}\tilde{h}_{\times}^{(0)}(z_s).$$
(97)

Note that Eq. (93) does not have components along $m_{\mu}m_{\nu}$ and $l_{\mu}l_{\nu}$, implying that there are no dispersive corrections affecting the left- and right-handed polarizations. However, the coupling between GWs and the background curvature induced by the lens triggers the existence of a new apparent polarization. In particular, this

effect actually arises from the failure of the left- and righthanded polarizations to remain orthogonal to the wave's propagation direction.

Recalling also Eq. (82), Eq. (93) shows an overall factor of $1/\Omega$, thus suggesting an explicit frequency dependence, which scales as 1/f as expected for bGO corrections leading to LID. In the high-frequency limit, $\Omega \to \infty$, these corrections vanish identically.

B. Massless Brans-Dicke

In this section, we discuss the bGO corrections for the massless Brans-Dicke theory. For transparency, we will restrict our analysis to a simple spherically symmetric background scalar field configuration: [99]

$$\bar{\phi}(R) \equiv \bar{\phi}_{\infty} + q \frac{GM_L}{R},\tag{98}$$

where ϕ_{∞} is an asymptotic costant value as $R \to \infty$ and q a dimensionless parameter. Furthermore, given its constant nature, we can set $\phi_{\infty} = 1$, thus recovering GR at the asymptotic limit when R is large.

Let us now start by analyzing the contribution of bGO arising from the scalar sector. By evaluating Eq. (74) using the BD matrices coefficients shown in Appendix B along with Eqs. (68), (90), and (98), the scalar amplitude correction can be re-written in a more compact form as

$$\delta\phi^{(1)}(z_{o}) = i \frac{\sqrt{\phi(z_{o})}}{\bar{D}(z_{o})} \int_{-\infty}^{+\infty} \frac{dz}{\Omega} \left[f_{\infty}^{\phi}(b,\tau;z) \delta\phi^{(0)}(z_{s}) + f_{\infty}^{+}(b,\tau;z) \tilde{\zeta}_{+}^{(0)}(z_{s}) + f_{\infty}^{-}(b,\tau;z) \tilde{\zeta}_{-}^{(0)}(z_{s}) \right],$$
(99)

where $\tau \equiv qM_L$ and the subscript ∞ refers to the form of the functions when the limit $z_s \to -\infty$ is performed.³

Moreover, the superscript denotes the GO variable to which the function is associated (e.g., f^{ϕ} is the function associated with $\delta\phi^{(0)}$, f^{\pm} to $\tilde{\zeta}^{(0)}_{\pm}(\xi_s)$, respectively). The functions $f^{\phi,\pm}_{\infty}(b,\tau;z)$ are presented in the Appendix C.

The result shows that the dispersive scalar correction is sourced by GO amplitudes, $\delta \phi^{(0)}(z_s)$ and $\tilde{\zeta}^{(0)}_{\pm}(z_s)$, thus encoding $\tilde{\alpha}^{(0)}_{mm}(z_s)$ and $\tilde{\alpha}^{(0)}_{ll}(z_s)$. The integration of the

³ In general, $f^{\phi,\pm}$ show an explicit dependence on z_s as well (refer to Appendix C) Note because the limit $z_s \to -\infty$ and $z_o \to +\infty$, affects not only the integration range, but also the integrand $f^{\phi,\pm}(b, z_s, \tau; z) \to f_{\infty}^{\phi,\pm}(b, \tau; z)$. It can be verified that the limit is well behaved.

aforementioned equation can be performed analytically, thus obtaining

$$\delta\phi^{(1)}(z_o) = \frac{i}{\Omega} \frac{\sqrt{\phi(z_o)}}{\bar{D}(z_o)} \left[F^{\phi}(b,\tau) \delta\phi^{(0)}(\xi_s) + F^{+}(b,\tau) \tilde{\zeta}^{(0)}_+(z_s) + F^{-}(b,\tau) \tilde{\zeta}^{(0)}_-(z_s) \right],$$
(100)

where $F^{\phi,\pm}(b,\tau)$ are the resulting integrals of the corresponding functions $f^{\phi,\pm}_{\infty}(b,\tau;z)$ whose explicit form is shown in the Appendix C1. It can be easily observed that such a correction is a frequency-dependent quantity, due to the presence of Ω , which behaves as 1/f and contributes as a modification to the phase of the scalar wave, due to the presence of the imaginary unit. The high-frequency limit causes this correction to vanish.

Let us derive the bGO corrections to the leading-order tensor amplitude by computing the coefficients appearing in Eq. (92). By perturbing Eq. (78) at the first order in Ψ along with the BD matrix coefficients appearing in Appendix B to make $F_{\mu\nu}$ explicit in terms of the metric and the scalar field, one gets

$$\delta\tilde{\alpha}_{AB}^{(1)}(z_{o}) = \frac{i}{\sqrt{\phi(z_{o})}\bar{D}(z_{o})} \int_{-\infty}^{+\infty} \frac{dz}{\Omega} \hat{e}_{A}^{\alpha} \hat{e}_{B}^{\beta} \frac{\bar{D}(z)}{\sqrt{\phi}} \left\{ -2(\delta R_{\alpha\beta})\delta\phi^{(0)} - 2\delta\tilde{h}_{\alpha\beta}^{(0)}\Box\bar{\phi} + \frac{2\omega X\delta\tilde{h}_{\alpha\beta}^{(0)}}{\bar{\phi}} - \frac{2(\delta\Gamma^{\gamma}{}_{\alpha\beta})\bar{\phi}_{\gamma}\delta\phi^{(0)}}{\bar{\phi}} + 2\delta\Gamma^{\rho}{}_{\alpha\beta}\tilde{h}_{\alpha\beta}^{(0)}\bar{\phi}^{\gamma} + 2\delta\Gamma^{\rho}{}_{\gamma[\beta}\tilde{h}_{\alpha]\rho}^{(0)}\bar{\phi}^{\gamma} - 2\tilde{h}_{\alpha\beta}^{(0)}\delta\Gamma^{\rho}{}_{\gamma\rho}\bar{\phi}^{\gamma} - 2\partial_{\beta}(\delta\tilde{h}_{\alpha\gamma}^{(0)})\bar{\phi}^{\gamma} + \partial_{\gamma}(\delta\tilde{h}_{\alpha\beta}^{(0)})\bar{\phi}^{\gamma} + \bar{\phi}\left[(\delta R)\tilde{h}_{\alpha\beta}^{(0)} - 2(\delta R^{\gamma}{}_{\beta})\tilde{h}_{\alpha\gamma}^{(0)} - 2(\delta R_{\gamma\alpha\beta\rho})\tilde{h}^{(0)\gamma\rho} + \Omega^{\gamma}{}_{\gamma\rho}\partial^{\rho}\tilde{h}_{\alpha\beta}^{(0)}\right] \right\}.$$

$$(101)$$

Subsequently plugging Eqs. (68), (90), and (98) into the above relation, each perturbed mode in the null tetrad basis can be expressed as

$$\delta \tilde{\alpha}_{AB}^{(1)}(z_o) = \frac{i}{\sqrt{\phi(z_o)}\bar{D}(z_o)} \int_{-\infty}^{+\infty} \frac{dz}{\Omega} \times \left[f_{AB_{\infty}}^{\phi}(b,\tau;z) \delta \phi^{(0)}(z_s) + f_{AB_{\infty}}^{\phi}(b,\tau;z) \tilde{\zeta}_{+}^{(0)}(z_s) + f_{AB_{\infty}}^{+}(b,\tau;z) \tilde{\zeta}_{-}^{(0)}(z_s) + f_{AB_{\infty}}^{-}(b,\tau;z) \tilde{\zeta}_{-}^{(0)}(z_s) \right],$$
(102)

where we refer to Appendix C for $f_{AB_{\infty}}^{\phi,\pm}(b,\tau;z)$. The subscript AB indicates that these functions contain, in their definition, the contraction with the appropriate dual tetrad $\hat{e}_{A}^{\alpha}\hat{e}_{B}^{\beta}$. Similarly to the scalar case, the result can be written in terms of leading-order scalar amplitude and tensor modes

$$\begin{pmatrix} \delta \tilde{\alpha}_{mm}^{(1)}(z_o) \\ \delta \tilde{\alpha}_{ll}^{(1)}(z_o) \\ \delta \tilde{\alpha}_{nn}^{(1)}(z_o) \end{pmatrix} = \frac{i}{\sqrt{\phi(z_o)} \bar{D}(z_o)} \begin{bmatrix} F_{mm}^{\phi}(b,\tau) & F_{mm}^{+}(b,\tau) & F_{mm}^{-}(b,\tau) \\ F_{ll}^{\phi}(b,\tau) & F_{ll}^{+}(b,\tau) & F_{ll}^{-}(b,\tau) \\ F_{nn}^{\phi}(b,\tau) & F_{nn}^{+}(b,\tau) & 0 \end{bmatrix} \begin{pmatrix} \delta \phi^{(0)}(z_s) \\ \tilde{\zeta}_{+}^{(0)}(z_s) \\ \tilde{\zeta}_{-}^{(0)}(z_s) \end{pmatrix},$$
(103)

$$\delta \tilde{\alpha}_{nm}^{(1)}(z_o) = \delta \tilde{\alpha}_{nl}^{(1)}(z_o) = 0.$$
(104)

where the functions $F_{AB}^{\phi,\pm}(b,\tau)$ are shown in Appendix C1. The above equation, together with (100),

describes the bGO corrections in the massless BD theory. The existence of scalar radiation and the non-minimal coupling of the scalar field with curvature trigger, from Eq. (101), $\delta \tilde{\alpha}_{mm}^{(1)}$ and $\delta \tilde{\alpha}_{ll}^{(1)}$ to become new non-zero coefficients. Note that these terms arise exclusively in the BD scenario and are completely absent in the GR case (refer to table III for a summary). Recalling that the mm and ll modes are related to the + and × polarizations, it is important to emphasize that these corrections have a direct impact on the GW polarizations.

Secondly, the $\delta \tilde{\alpha}_{nn}^{(1)}$ -mode emerge as an apparent scalar polarization, and while it is also predicted by GR, it exhibits distinct differences in comparison. In the GR case, one can verify that most of the integrals in Eq. (101) vanish and the only term yielding a non-zero outcome is $(\delta R_{\gamma\alpha\beta\rho})\tilde{h}^{(0)\gamma\rho}$, thus providing the result shown in Eq. (93).⁴ In BD instead, it can be observed that there are two additional contributions from Eq. (101), namely $(\delta\Gamma^{\gamma}{}_{\alpha\beta})\bar{\phi}_{\gamma}\delta\phi^{(0)}/\bar{\phi}$ and $\delta\Gamma^{\rho}{}_{\alpha\beta}\tilde{h}^{(0)}_{\gamma\rho}\bar{\phi}^{\gamma}$.

The first term is related to the existence of scalar waves that influence the evolution of tensor amplitude, whereas the second one pertains to the coupling between scalar fields and the gravitational dynamics. Essentially, this means that even if we disregard scalar waves and focus exclusively on how the background scalar field impacts the propagation of tensor waves, there would nevertheless appear a discrepancy from what GR results predict. Note that $\delta \tilde{\alpha}_{nm}^{(1)}(z_o)$ and $\delta \tilde{\alpha}_{nl}^{(1)}(z_o)$ are vanishing even for BD. In general, the limit to GR is straightforward to perform by setting $\delta \phi^{(0)}, \tau, \beta \to 0$.

In BD, the bGO regime not only induces a new apparent polarization (with novel contributions due to the existence of the scalar wave and scalar-gravitational coupling) but also leads to corrections along the left- and right-handed polarizations, thus producing additional phase modifications to the GWs phase. LID corrections to standard GW polarizations are a smoking gun of BD: Detecting a non-zero $\delta \tilde{\alpha}_{mm}^{(1)}$ and $\delta \tilde{\alpha}_{ll}^{(1)}$ would be a clear signature of deviations from GR.

Finally, once all coefficients are plugged into Eq. (92), along with Eqs. (81–84), the result will account for an overall bGO dispersive correction / LID scaling as $1/\Omega$, thus as 1/f.

Summarizing, we derived analytical expressions for the LID corrections for a point-like lens. We considered a spherically symmetric background scalar field, performing the calculation perturbatively up to the first order in the gravitational potential, assuming a Euclidean comoving distance on the geodesic. We restricted the analysis to the special case in which source and observation points are pushed to infinity, *i.e.* $z_s \to -\infty, z_o \to +\infty$. This situation in BD triggers the nn component, similarly to GR but with a different amplitude and dependence on the parameters. This can be interpreted as an apparent polarization caused by the fact that, in the bGO regime, the polarizations are no longer orthogonal to the propagation direction. Additionally, there are corrections to the mm and ll components, which directly impact the standard GW polarizations: these corrections are absent in GR and provide a smoking gun for BD or other alternative theories.

C. Jordan frame

The above results were obtained in the Einstein frame, in which the dynamical variables decouple. To connect with observables, we need to rewrite them in the Jordan frame, in which matter couples minimally to the metric perturbation. In order to do that, we need to reverse Eq. (60) by obtaining

$$h_{\mu\nu} = \tilde{h}_{\mu\nu} - \frac{1}{2}g_{\mu\nu}\tilde{h} - g_{\mu\nu}\frac{\delta\phi}{\bar{\phi}}.$$
 (105)

By recalling $\tilde{h} = 0$, the above equation reduces to

$$h_{\mu\nu} = \tilde{h}_{\mu\nu} - g_{\mu\nu} \frac{\delta\phi}{\bar{\phi}}.$$
 (106)

We note that the short-wave approximation (25) holds for $h_{\mu\nu}$ as well. By plugging the latter ansatz into the above equation, the relation on the amplitude, at any order n, yields

$$h_{\mu\nu}^{(n)} = \tilde{h}_{\mu\nu}^{(n)} - g_{\mu\nu} \frac{\delta\phi^{(n)}}{\bar{\phi}}.$$
 (107)

Applying the general rank-2 tensor decomposition (36) on both sides of Eq. (107), one gets the general transformation of the expansion coefficients between the Einstein and Jordan frames

$$\alpha_{CD}^{(n)}\Theta_{\mu\nu}^{AB} = \tilde{\alpha}_{AB}^{(n)}\Theta_{\mu\nu}^{AB} - g_{\mu\nu}\frac{\delta\phi^{(n)}}{\bar{\phi}}.$$
 (108)

⁴ To study the GR case from Eq. (101), one simply needs to discard the contributions from the scalar field and the scalar wave by setting $\bar{\phi}$ = constant. Consequently, the derivatives of the field will vanish, and the amplitudes at every order of the WKB expansion will be zero.

At the leading order in the short-wave expansion, by using Eq. (35), employing the appropriate gauge conditions, in both Jordan and Einstein frames, and subsequently contracting the above result with dual tetrads, one gets

$$\alpha_{mm}^{(0)} = \tilde{\alpha}_{mm}^{(0)}, \tag{109}$$

$$\alpha_{ll}^{(0)} = \tilde{\alpha}_{ll}^{(0)},\tag{110}$$

$$\alpha_{ml}^{(0)} = -\frac{\delta\phi^{(0)}}{\bar{\phi}},\tag{111}$$

with all the other coefficients vanishing.

Coefficients describing bGO corrections are evaluated in the same way. The nonzero LID terms read

$$\alpha_{mm}^{(1)} = \tilde{\alpha}_{mm}^{(1)}, \qquad (112)$$

$$\alpha_{u}^{(1)} = \tilde{\alpha}_{u}^{(1)}.$$
 (113)

$$\alpha_{nn}^{(1)} = \tilde{\alpha}_{nn}^{(1)}, \tag{114}$$

$$\alpha_{ml}^{(1)} = -\frac{\delta\phi^{(1)}}{\bar{\phi}}\,,\tag{115}$$

while $\alpha_{nm}^{(1)} = \tilde{\alpha}_{nm}^{(1)}$ and $\alpha_{nl}^{(1)} = \tilde{\alpha}_{nl}^{(1)}$ vanish identically. Moreover, as the bGO corrections are derived at first order in the gravitational potential Ψ , the aforementioned relations apply to the terms $\delta \tilde{\alpha}_{AB}^{(1)}$ by replacing $\tilde{\alpha}_{AB}^{(1)}$ with $\delta \tilde{\alpha}_{AB}^{(1)}$.

From the above equations, it can be observed that the scalar field perturbation excites the ml-mode, while mm, ll, and nn coefficients remain invariant under the mapping between the Einstein and Jordan frames. Ultimately, we can express the GO and bGO tensor amplitudes in the Jordan frame as

$$h_{\mu\nu}^{(0)} = \alpha_{mm}^{(0)} \bar{m}_{\mu} \bar{m}_{\nu} + \alpha_{ll}^{(0)} \bar{l}_{\mu} \bar{l}_{\nu} - \frac{\delta\phi^{(0)}}{\bar{\phi}} \bar{m}_{(\mu} \bar{l}_{\nu)}, \qquad (116)$$

$$h_{\mu\nu}^{(1)} = \delta\alpha_{mm}^{(1)}\bar{m}_{\mu}\bar{m}_{\nu} + \delta\alpha_{ll}^{(1)}\bar{l}_{\mu}\bar{l}_{\nu} + \delta\alpha_{nn}^{(1)}\bar{n}_{\mu}\bar{n}_{\nu} - \frac{\delta\phi^{(1)}}{\bar{\phi}}\bar{m}_{(\mu}\bar{l}_{\nu)}$$
(117)

VI. CONCLUSION

This work presents a novel framework to describe frequency-dependent corrections to the propagation of gravitational waves (GWs) in theories beyond Einstein's General Relativity (GR). These represent dispersive phenomena, analog to how a prism splits light into its component frequencies. The role of the prism is played by a gravitational lens, which allows the standard GW polarizations $(+, \times)$ to interact with fields with lower spin

	$\delta \alpha_{nn}^{(1)}$	$\delta \alpha_{mm}^{(1)}$	$\delta \alpha_{ll}^{(1)}$	$\delta \alpha_{ml}^{(1)}$	$\delta \alpha_{nl}^{(1)}$	$\delta \alpha_{nm}^{(1)}$
GR	\checkmark	×	×	×	×	×
BD	\checkmark	\checkmark	\checkmark	\checkmark	×	×

TABLE III: Schematic summary of the zero (\checkmark) vs non-zero (\checkmark) bGO corrections in Einstein's GR and Brans-Dicke (BD), in the Jordan frame. In addition to introducing dispersive effects in the standard GW polarizations ($\delta \alpha_{ll}^{(1)}, \delta \alpha_{mm}^{(1)} \neq 0$), BD modifies the $\delta \alpha_{nn}^{(n)}$. Moreover, the scalar field perturbation excites $\delta \alpha_{ml}^{(1)}$.

(e.g., a scalar). Lens-induced dispersion (LID) phenomena appear beyond geometric optics (bGO) and can be tested on any GW signal, not requiring an electromagnetic counterpart.

We begin presenting a general, model-independent description of linearized GW propagation in inhomogeneous spacetime, relying only on a short-wave expansion. For simplicity, we focus on scalar-tensor theories, which include an additional scalar field. The propagation of the radiative d.o.f. is described by a system of tensorial differential equations, coupled due to the interaction between gravity and the scalar field. We express this system compactly in matrix form and introduce the kinetic, amplitude and mixing matrices, which respectively encode interactions with two, one and zero derivatives of the perturbations, and whose coefficients are functions of the background quantities. We then describe how to define *propagation eigenstates*, for which the kinetic matrix is diagonal [58, 63], and present the general structure of the equations in the short-wave expansion. The analogy here is similar to the way neutrinos change states as they travel. In the realm of beyond GR theories, the shift from interaction to propagation eigenstates involves interactions involving two derivatives, rather than zero.

We present the computation of bGO corrections in GR before addressing scalar-tensor theories (Sec. III). This also serves as an introduction of the tetrad decomposition, affine parameter and luminosity distance. We review how bGO corrections introduce an apparent new polarization, the nn mode, via dispersive corrections. We then apply the same concepts to Brans-Dicke (BD) theory (Sec. IV), presenting for the first time the structure of the equations that describe dispersive phenomena in GW propagation beyond GR. Although complex, two important properties of BD make these computations feasible: First, the diagonalization of the kinetic matrix is simple (60), and equivalent to the well-known definition of the Einstein frame metric. Second, BD is a fully-luminal theory, with all excitations sharing the same propagation speed, wavevector and geodesics (Sec IV A).

We then present an explicit computation of lensinduced diffraction (LID) in BD in the presence of a point lens (Sec V). Additional, well-motivated approximations allow us to present analytical results: we assume the lens to be isolated and that deviations w.r.t. flat spacetime are small for the trajectories (Born approximation) and negligible at large separations from the lens. As expected, all corrections reduce to GR results when the scalar field is constant, and LID vanish as 1/f at high frequencies, recovering the GO limit. To facilitate the interpretation of data in the context of BD theory, we rewrite our results in terms of the minimally-coupled Jordan frame metric, Eqs. (116) and (117).

Our results exemplify how LID provides a smoking gun for deviations from GR:

- Additional physical polarizations are present and receive dispersive corrections, which depend on the theory parameters. For BD, the scalar field perturbation sources the *ml* component, additional breathing mode which is not present in GR.
- Like in GR, *apparent* new polarizations (*nn*) are sourced by bGO corrections. The amplitude of this correction depends on the BD parameter.
- The standard metric polarizations (+, ×) receive novel frequency-dependent corrections. These are absent in GR and would be a clear signature of new gravitational dynamics.

This rich structure emerges even for the relatively simple case of BD, despite important simplifications like fullluminality and the absence of kinetic interactions. More complex theories of gravity will lead to more and/or stronger observational signatures.

Three important aspects of LID make them promising for testing gravity. First, the frequency-dependent corrections can be tested on all GW sources (irrespective of their intrinsic properties, electromagnetic counterpart, etc.). Second, they stem from inhomogeneous backgrounds, which distinguish between the standard GW polarizations $(+, \times)$. Finally, they arise from interactions with zero derivatives, which are expected in any modified theory. LID phenomena are therefore a universal prediction in theories beyond GR.

This work bridges important gaps in the theoretical understanding of GWs propagation beyond GR. Future research should address more complex gravity theories, including different speeds for scalar and tensor waves, screening mechanisms, or non-trivial cosmological dynamics, likely leading to enhanced dispersive effects on the gravitational wave signal. Because dispersive effects are stronger at low frequencies, this program needs to not only consider ground detectors, but also space-borne observatories and ultra-low frequency GWs observable through pulsar-timing arrays. This program will enable novel tests of gravity and dark energy theories, leveraging the full potential of the GW spectrum.

Acknowledgments

We thank Guillerme Brando, Han Gil Choi, Jose Maria Ezquiaga, Srashti Goyal, Serena Giardino, Stefano Savastano, and Hiroki Takeda for the discussions and comments on this work. N.M. thanks D. Usseglio for his valuable insights into the project and is especially grateful to Prof. S. Capozziello for his support and encouragement throughout all stages of this work. N.M. also acknowledges the Scuola Superiore Meridionale for sponsoring his visiting period to the Albert Einstein Institute (AEI) in Potsdam, which provided an invaluable opportunity for collaboration and research.

Appendix A: Weak field limit

Given the line element of Eq. (79), one gets the Christoffel symbols up to first order in Ψ

$$\delta \Gamma^{0}_{\ 00} = \delta \Gamma^{0}_{\ ij} = \delta \Gamma^{i}_{\ j0} = 0,$$
 (A1)

$$\partial \Gamma^0_{\ i0} = \partial_i \Psi,$$
 (A2)

$$\delta\Gamma^{i}{}_{jk} = \delta_{jk}\partial^{i}\Psi - \delta^{i}_{k}\partial_{j}\Psi - \delta^{i}_{j}\partial_{k}\Psi, \qquad (A3)$$

where δ_{ij} is the Kronecker delta. Further, the linearized Riemann and Ricci tensors and scalar, respectively, are then provided

$$\delta R^{\rho}_{\ \mu\lambda\nu} = \partial_{\lambda} \delta \Gamma^{\rho}_{\ \mu\nu} - \partial_{\nu} \delta \Gamma^{\rho}_{\ \mu\lambda}, \tag{A4}$$

$$\delta R_{\mu\nu} = \delta^{\lambda}_{\rho} \delta R^{\rho}_{\ \mu\lambda\nu}, \tag{A5}$$

$$\delta R = g^{\mu\nu} \delta R_{\mu\nu}.\tag{A6}$$

Appendix B: Brans-Dicke background functions

Here we present the non-zero components of the kinetic, amplitude, and mass matrices, appearing in Eq. (62), which constitute the system of differential equations describing the propagation of GWs and scalar waves in the TT gauge. In particular, we have

$$\tilde{\mathsf{K}}_{\mu\nu}{}^{\alpha\beta\gamma\rho} \equiv -\frac{1}{2}\bar{\phi}\delta^{\alpha}_{\mu}\delta^{\beta}_{\nu}g^{\gamma\rho}, \tag{B1}$$

$$\tilde{\mathsf{K}}^{\gamma\rho} \equiv -\bar{\phi}g^{\gamma\rho},\tag{B2}$$

$$\tilde{\mathsf{A}}_{\mu\nu}^{\ \alpha\beta\gamma} \equiv \frac{1}{2} \left(2\bar{\phi}^{\alpha}\delta^{\beta}_{(\mu}\delta^{\gamma}_{\nu)} - \bar{\phi}^{\gamma}\delta^{\alpha}_{\mu}\delta^{\beta}_{\nu} \right),\tag{B3}$$

$$\tilde{\mathsf{A}}_{\mu\nu}{}^{\gamma} \equiv \frac{3+2\omega}{2\bar{\phi}} \left(g_{\mu\nu}\bar{\phi}^{\gamma} - 2\bar{\phi}_{(\mu}\delta_{\nu)}^{\gamma} \right),\tag{B4}$$

$$\tilde{\mathsf{A}}^{\gamma} \equiv \bar{\phi}^{\gamma},\tag{B5}$$

$$\begin{split} \tilde{\mathsf{M}}_{\mu\nu}{}^{\alpha\beta} &\equiv \frac{\phi}{2} \left(g_{\mu\nu} R^{\alpha\beta} + 2R_{(\mu}{}^{\alpha}\delta^{\beta}_{\nu)} - \delta^{\alpha}_{\mu}\delta^{\beta}_{\nu}R + 2R^{\alpha}{}_{\mu\nu}{}^{\beta} \right) + \\ &+ \delta^{\alpha}_{\mu}\delta^{\beta}_{\nu} \left(\Box \bar{\phi} - \frac{\omega X}{\bar{\phi}} \right) - g_{\mu\nu} \left(\bar{\phi}^{\beta\alpha} + \frac{1}{2} \bar{\phi}^{\alpha} \bar{\phi}^{\beta} \right), \end{split} \tag{B6}$$

$$\tilde{\mathsf{M}}^{h}_{\mu\nu} \equiv R_{\mu\nu} - g_{\mu\nu} \left[\frac{1}{2} R - \frac{\Box \bar{\phi}}{\bar{\phi}} - \frac{(3+\omega)X}{\bar{\phi}^2} \right] + \\
+ \frac{(3+\omega)}{\bar{\phi}^2} \bar{\phi}_{\mu} \bar{\phi}_{\nu} - \frac{\bar{\phi}_{\nu\mu}}{\bar{\phi}},$$
(B7)

$$\tilde{\mathsf{M}}^{\alpha\beta}_{\phi} \equiv \bar{\phi}(\bar{\phi}_{\alpha\beta}),\tag{B8}$$

$$\tilde{\mathsf{M}} \equiv \frac{2X}{\bar{\phi}},\tag{B9}$$

where $X \equiv -\nabla_{\mu} \bar{\phi} \nabla^{\mu} \bar{\phi}/2$ is the canonical kinetic term of the scalar field.

Appendix C: Functions for bGO corrections

Here, we present the functions to be integrated, along the z-axis, to obtain the bGO corrections. For clarity, we have set $\bar{\phi}_{\infty} = 1$, as discussed in Sec. V B. Let us start by showing the functions of Eq. (99)

$$f^{\phi}(b, z_s, \tau; z) = \frac{\bar{D}(z_s)}{\sqrt{\bar{\phi}(z_s)}} \frac{p(b, z_s, \tau; z)}{q(b, z_s, \tau; z)}, \qquad (C1)$$

with

$$p(b, z_s, \tau; z) \equiv \tau^2 \left(8b^2 + 9z^2 - 2zz_s + z_s^2\right) + 16\tau (b^2 + z^2)^{3/2} + 8 \left(b^2 + z^2\right)^2,$$
(C2)

$$q(b, z_s, \tau; z)] \equiv 4 (b^2 + z^2) (z - z_s)^2 \times \\ \times \left[(b^2 + z^2) + 2\tau (b^2 + z^2)^{1/2} + \tau^2 \right].$$
(C3)

The functions multiplying $\tilde{\zeta}^{(0)}_+$ and $\tilde{\zeta}^{(0)}_-,$ respectively, are

$$f^{+}(b, z_{s}, \tau; z) = -\bar{D}(z_{s})\sqrt{\bar{\phi}(z_{s})}\frac{3\tau(b^{2} + 2zR_{s})}{2g(b, \tau; z)}, \quad (C4)$$

$$f^-(b, z_s, \tau; z) = 0, \tag{C5}$$

with

$$g(b,\tau;z) \equiv (b^2 + z^2)^2 \left[\tau + (b^2 + z^2)^{1/2}\right].$$
 (C6)

We proceed by presenting the elements that compose Eq. (102), only considering the coefficients appearing in Eq. (92). The quantity $\delta \alpha_{nn}^{(1)}$ is build up by

$$f_{nn}^{\phi}(b,\tau;z) = 2\Omega^2 R_s \frac{D(z_s)}{\sqrt{\phi(z_s)}} \times \left[\frac{b^2 \tau}{(b^2 + z^2)^{1/2} g(b,\tau;z)} - \frac{(b^2 - 2z^2)}{(b^2 + z^2)^{5/2}} \right],$$
(C7)

$$f_{nn}^{+}(b,\tau;z) = -b^{2}\Omega^{2}R_{s}\bar{D}(z_{s})\sqrt{\phi(z_{s})} \times \left[\frac{\tau}{(b^{2}+z^{2})^{1/2}g(b,\tau;z)} + \frac{3}{(b^{2}+z^{2})^{5/2}}\right],$$
(C8)

$$f_{nn}^{-}(b,\tau;z) = 0.$$
 (C9)

Proceeding with $\delta \alpha_{mm}^{(1)}$, we have

$$f_{mm}^{\phi}(b, z_s, \tau; z) = b^2 R_s \frac{\bar{D}(z_s)}{\sqrt{\phi(z_s)}} \times \left[-\frac{\tau}{(b^2 + z^2)^{1/2} g(b, \tau; z)} + \frac{3}{(b^2 + z^2)^{5/2}} \right],$$
(C10)

$$\begin{aligned} f_{mm}^{+}(b, z_{s}, \tau; z) &= f_{mm}^{-}(b, z_{s}, \tau; z) = \\ &= R_{s} \bar{D}(z_{s}) \sqrt{\bar{\phi}(z_{s})} \left\{ \frac{5 \left(b^{2} - 2z^{2}\right)}{2 \left(b^{2} + z^{2}\right)^{5/2}} - \frac{\tau (b^{2} + z^{2})^{1/2}}{g(b, \tau; z)} + \right. \\ &\left. + \frac{z\tau^{2} \left(b^{2} + z^{2}\right)^{2}}{4g(b, \tau; z)^{2}} + \frac{b^{2}\tau}{2(b^{2} + z^{2})^{1/2}g(b, \tau; z)} \right\}, \end{aligned}$$

$$(C11)$$

Given that the tetrads m^{μ} and l^{μ} differ only by a complex conjugation, the functions characterizing $f_{ll}^{\phi,\pm}(b, z_s, \tau; z)$ exhibit the same functional dependence as those of $\delta \alpha_{mm}^{(1)}$, following

$$f^{\phi}_{mm}(b, z_s, \tau; z) = f^{\phi}_{ll}(b, z_s, \tau; z),$$

$$f^{+}_{mm}(b, z_s, \tau; z) = f^{+}_{ll}(b, z_s, \tau; z),$$

$$f^{-}_{mm}(b, z_s, \tau; z) = -f^{-}_{ll}(b, z_s, \tau; z).$$
(C12)

The functions for $\delta \alpha_{nm}^{(1)}$ are instead

$$f_{nm}^{\phi}(b, z_s, \tau; z) = \sqrt{2}\Omega b z R_s \frac{D(z_s)}{\sqrt{\phi(z_s)}} \times \left[\frac{\tau}{(b^2 + z^2)^{1/2} g(b, \tau; z)} - \frac{3}{(b^2 + z^2)^{5/2}}\right].$$
 (C13)

$$\begin{aligned} f_{nm}^{+}(b,z_{s},\tau;z) &= f_{nm}^{-}(b,z_{s},\tau;z) = \\ &- \frac{\Omega R_{s}b}{2\sqrt{2}} \bar{D}(z_{s}) \sqrt{\bar{\phi}(z_{s})} \left[\frac{3z}{(b^{2}+z^{2})^{5/2}} + \right. \\ &+ \frac{1}{(b^{2}+z^{2})^{3/2}(z-z_{s})} + \frac{z\tau}{(b^{2}+z^{2})^{1/2}g(b,\tau;z)} \right], \end{aligned}$$
(C14)

Analogously, the functions $f_{nm}^{\phi,\pm}(b, z_s, \tau; z)$ follows

$$\begin{aligned} f_{nl}^{\phi}(b, z_{s}, \tau; z) &= f_{nl}^{\phi}(b, z_{s}, \tau; z), \\ f_{nl}^{+}(b, z_{s}, \tau; z) &= f_{ll}^{+}(b, z_{s}, \tau; z), \\ f_{nl}^{-}(b, z_{s}, \tau; z) &= -f_{nl}^{-}(b, z_{s}, \tau; z). \end{aligned} \tag{C15}$$

We now consider the case where $z_s \to -\infty$ and $z_o \to +\infty$: apart from the integration boundaries, this operation also applies to the functions $f^{\phi,\pm}(b, z_s, \tau; b)$ and $f_{AB}^{\phi,\pm}(b, z_s, \tau; b)$. In particular, we observe that the functional dependence on z_s appears only in the functions $f^{\phi}(b, z_s, \tau; b)$ and $f_{nm}^{\pm}(b, z_s, \tau; b)$. Performing the aforementioned limit on these two, we obtain

$$f_{\infty}^{\phi}(b,\tau;z) = \frac{\bar{D}(z_s)}{\sqrt{\bar{\phi}(z_s)}} \frac{\tau^2(b^2+z^2)}{4g(b,\tau;z)^2},$$
(C16)

$$\begin{split} f_{nm_{\infty}}^{\pm}(b,\tau;z) &= -\frac{\Omega R_{s}b}{2\sqrt{2}}\bar{D}(z_{s})\sqrt{\bar{\phi}(z_{s})} \times \\ & \times \left[\frac{3z}{(b^{2}+z^{2})^{5/2}} + \frac{z\tau}{(b^{2}+z^{2})^{1/2}g(b,\tau;z)}\right]. \end{split}$$
(C17)

1. Results from analytic integration

We define $F^{\phi,\pm}(b,\tau)$ and $F^{\phi,\pm}_{AB}(b,\tau)$ as

$$F^{\phi,\pm}(b,\tau) \equiv \int_{-\infty}^{+\infty} \frac{dz}{\Omega} f_{\infty}^{\phi,\pm}(b,\tau;z), \qquad (C18)$$

$$F_{AB}^{\phi,\pm}(b,\tau) \equiv \int_{-\infty}^{+\infty} \frac{dz}{\Omega} f_{AB_{\infty}}^{\phi,\pm}(b,\tau;z).$$
(C19)

The scalar bGO correction $\delta \phi^{(1)}$ is obtained by employing the definition (C18) along with Eqs. (C4), (C5) and (C16), thus obtaining

$$F^{\phi}(b,\tau) \equiv \frac{\bar{D}(z_s)}{\sqrt{\phi(z_s)}} \left[\frac{\pi b^2 - 2b\tau - \pi\tau^2}{4b(b^2 - \tau^2)} + \frac{b^2 - 2\tau^2}{(b^2 - \tau^2)}u(b,\tau) \right]$$
(C20)

$$F^{+}(b,\tau) \equiv -\bar{D}(z_{s})\sqrt{\bar{\phi}(z_{s})} \times \\ \times \frac{3b^{2}}{4\tau^{2}} \left[\frac{2\pi b^{2} - 4b\tau + \pi\tau^{2}}{b^{3}} + 8u(b,\tau)\right],$$
(C21)

$$F^{-}(b,\tau) \equiv 0, \tag{C22}$$

with

$$u(b,\tau) \equiv \frac{1}{(b^2 - \tau^2)^{1/2}} \times \\ \times \left[\tan^{-1} \left(\frac{\tau}{\sqrt{b^2 - \tau^2}} \right) - \cot^{-1} \left(\sqrt{\frac{b - \tau}{b + \tau}} \right) \right].$$
(C23)

The limit to GR is achieved by setting $\tau \to 0$, which leads $F^{\phi}(b,\tau)$ and $F^{+}(b,\tau)$ to identically vanish. This result is consistent since, in GR, the scalar wave does not exist.

The results regarding the coefficients $\delta \tilde{\alpha}_{AB}^{(1)}$ composing the tensor bGO correction $h_{\mu\nu}^{(1)}$ are derived by using Eq. (C19). By starting from the $\delta \tilde{\alpha}_{nn}^{(1)}$, the result is

$$F_{nn}^{\phi}(b,\tau) = \frac{D(z_s)}{\sqrt{\phi}(z_s)} \frac{\Omega R_s}{\tau^3} \times \left\{ \frac{8\tau^3}{3b^2} - \frac{\pi\tau^2}{b} + 4\tau - 2b\left[\pi + 4bu(b,\tau)\right] \right\},$$
(C24)

$$F_{nn}^{+}(b,\tau) = -\bar{D}(z_s)\sqrt{\bar{\phi}(z_s)}\frac{4\Omega R_s}{b^2} \times \left\{\frac{4}{3} - \frac{\pi b}{8\tau} + \frac{b^2}{2\tau^2} - \frac{\pi b^3}{4\tau^3} - \frac{b^4}{\tau^3}u(b,\tau)\right\},$$
(C25)

$$F_{nn}^{-}(b,\tau) = 0.$$
 (C26)

The $\tau \to 0$ limit makes $F_{nn}^{\phi}(b,\tau)$ and $F_{nn}^{-}(b,\tau)$ vanish but not $F_{nn}^{+}(b,\tau)$ whose terms in the parenthesis resemble to provide the result in Eq. (93), with $\bar{\phi}(z_s) \to 1$.

The result for the mm and ll components acts as

$$F_{mm}^{\phi}(b,\tau) = F_{ll}^{\phi}(b,\tau) = \frac{D(z_s)}{\sqrt{\phi(z_s)}} \frac{R_s}{\Omega} \times \left[\frac{8}{3b^2} + \frac{\pi}{2b\tau} - \frac{2}{\tau^2} + \frac{\pi b}{\tau^3} - \frac{4b^2 u(b,\tau)}{\tau^3} \right],$$
(C27)

$$F_{mm}^{+}(b,\tau) = F_{mm}^{-}(b,\tau) = F_{ll}^{+}(b,\tau) = -F_{ll}^{-}(b,\tau) =$$

$$= \bar{D}(z_s)\sqrt{\bar{\phi}(z_s)}\frac{R_s}{\Omega} \times$$

$$\times \left[-\frac{4}{3b^2} + \frac{3\pi}{4b\tau} + \frac{1}{\tau^2} - \frac{\pi b}{2\tau^3} - \frac{2(b^2 - 2\tau^2)u(b,\tau)}{\tau^3}\right].$$
(C28)

To conclude, the integration of Eqs. (C13) and (C14) vanishes identically, thus providing $\delta \tilde{\alpha}_{nm}^{(1)} = \delta \tilde{\alpha}_{nl}^{(1)} = 0$.

- N. Aghanim <u>et al.</u> (Planck), Astron. Astrophys. **641**, A6 (2020), [Erratum: Astron.Astrophys. 652, C4 (2021)], arXiv:1807.06209 [astro-ph.CO].
- [2] E. Oks, New Astron. Rev. 93, 101632 (2021), arXiv:2111.00363 [astro-ph.CO].
- [3] A. K. Drukier, K. Freese, and D. N. Spergel, Phys. Rev. D 33, 3495 (1986).
- [4] K. Arun, S. B. Gudennavar, and C. Sivaram, Advances in Space Research 60, 166 (2017), arXiv:1704.06155 [physics.gen-ph].
- [5] A. G. Riess <u>et al.</u> (Supernova Search Team), Astron. J. 116, 1009 (1998), arXiv:astro-ph/9805201.
- S. Perlmutter et al. (Supernova Cosmology Project), Astrophys. J. 517, 565 (1999), arXiv:astro-ph/9812133.
- [7] M. Li, X.-D. Li, S. Wang, and Y. Wang, Front. Phys. (Beijing) 8, 828 (2013), arXiv:1209.0922 [astro-ph.CO].
- [8] M. Li, X.-D. Li, S. Wang, and Y. Wang, Commun. Theor. Phys. 56, 525 (2011), arXiv:1103.5870 [astro-ph.CO].
- [9] J. Frieman, M. Turner, and D. Huterer, Ann. Rev. Astron. Astrophys. 46, 385 (2008), arXiv:0803.0982 [astroph].
- [10] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, Phys. Rep. **530**, 87 (2013), arXiv:1201.2434 [astro-ph.CO].
- [11] E. J. Copeland, M. Sami, and S. Tsujikawa, Int. J. Mod. Phys. D 15, 1753 (2006), arXiv:hep-th/0603057.
- [12] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, Phys. Rept. **513**, 1 (2012), arXiv:1106.2476 [astroph.CO].
- [13] S. Capozziello and M. De Laurentis, Phys. Rept. 509, 167 (2011), arXiv:1108.6266 [gr-qc].
- [14] A. Joyce, B. Jain, J. Khoury, and M. Trodden, Phys. Rept. 568, 1 (2015), arXiv:1407.0059 [astro-ph.CO].
- [15] D. H. Weinberg, M. J. Mortonson, D. J. Eisenstein, C. Hirata, A. G. Riess, and E. Rozo, Phys. Rept. 530, 87 (2013), arXiv:1201.2434 [astro-ph.CO].
- [16] P. A. R. Ade <u>et al.</u> (Planck), Astron. Astrophys. **594**, A14 (2016), arXiv:1502.01590 [astro-ph.CO].
- [17] Y. Mellier <u>et al.</u> (Euclid), (2024), arXiv:2405.13491 [astro-ph.CO].
- [18] G. Cusin, M. Lewandowski, and F. Vernizzi, JCAP 04, 061, arXiv:1712.02782 [astro-ph.CO].
- [19] G. Cusin, M. Lewandowski, and F. Vernizzi, JCAP 04, 005, arXiv:1712.02783 [astro-ph.CO].
- [20] E. Bellini and I. Sawicki, JCAP 07, 050, arXiv:1404.3713 [astro-ph.CO].
- [21] M. Zumalacárregui, E. Bellini, I. Sawicki, J. Lesgourgues, and P. G. Ferreira, JCAP 08, 019, arXiv:1605.06102 [astro-ph.CO].
- [22] M. Ishak, Living Rev. Rel. 22, 1 (2019), arXiv:1806.10122 [astro-ph.CO].

- [23] E. Abdalla et al., JHEAp 34, 49 (2022), arXiv:2203.06142
 [astro-ph.CO].
- [24] B. P. Abbott <u>et al.</u> (LIGO Scientific, Virgo), Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [25] B. P. Abbott <u>et al.</u> (LIGO Scientific, Virgo), Phys. Rev. D 100, 104036 (2019), arXiv:1903.04467 [gr-qc].
- [26] R. Abbott <u>et al.</u> (LIGO Scientific, Virgo), Physical Review D 103, 122002 (2021).
- [27] R. Abbott <u>et al.</u> (LIGO Scientific, VIRGO, KAGRA), (2021), arXiv:2112.06861 [gr-qc].
- [28] B. P. Abbott <u>et al.</u> (LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL), Astrophys. J. Lett. 848, L13 (2017), arXiv:1710.05834 [astro-ph.HE].
- [29] C. de Rham, J. T. Deskins, A. J. Tolley, and S.-Y. Zhou, Rev. Mod. Phys. 89, 025004 (2017), arXiv:1606.08462 [astro-ph.CO].
- [30] J. M. Ezquiaga and M. Zumalacárregui, Phys. Rev. Lett. 119, 251304 (2017), arXiv:1710.05901 [astro-ph.CO].
- [31] P. Creminelli and F. Vernizzi, Phys. Rev. Lett. 119, 251302 (2017), arXiv:1710.05877 [astro-ph.CO].
- [32] T. Baker, E. Bellini, P. G. Ferreira, M. Lagos, J. Noller, and I. Sawicki, Phys. Rev. Lett. **119**, 251301 (2017), arXiv:1710.06394 [astro-ph.CO].
- [33] L. Lombriser and A. Taylor, JCAP 03, 031, arXiv:1509.08458 [astro-ph.CO].
- [34] D. Bettoni, J. M. Ezquiaga, K. Hinterbichler, and M. Zumalacárregui, Phys. Rev. D 95, 084029 (2017), arXiv:1608.01982 [gr-qc].
- [35] K. Bamba, S. Capozziello, S. Nojiri, and S. D. Odintsov, Astrophys. Space Sci. **342**, 155 (2012), arXiv:1205.3421 [gr-qc].
- [36] J. Sakstein and B. Jain, Phys. Rev. Lett. 119, 251303 (2017), arXiv:1710.05893 [astro-ph.CO].
- [37] P. Creminelli, M. Lewandowski, G. Tambalo, and F. Vernizzi, JCAP 12, 025, arXiv:1809.03484 [astroph.CO].
- [38] P. Creminelli, G. Tambalo, F. Vernizzi, and V. Yingcharoenrat, JCAP 05, 002, arXiv:1910.14035 [gr-qc].
- [39] P. Creminelli, G. Tambalo, F. Vernizzi, and V. Yingcharoenrat, JCAP 10, 072, arXiv:1906.07015 [gr-qc].
- [40] M. Lagos, L. Jenks, M. Isi, K. Hotokezaka, B. D. Metzger, E. Burns, W. M. Farr, S. Perkins, K. W. K. Wong, and N. Yunes, Phys. Rev. D 109, 124003 (2024), arXiv:2402.05316 [gr-qc].
- [41] I. D. Saltas, I. Sawicki, L. Amendola, and M. Kunz, Phys. Rev. Lett. 113, 191101 (2014), arXiv:1406.7139 [astroph.CO].
- [42] J. M. Ezquiaga and M. Zumalacárregui, Front. Astron. Space Sci. 5, 44 (2018), arXiv:1807.09241 [astro-ph.CO].

- [43] J. M. Ezquiaga, W. Hu, M. Lagos, and M.-X. Lin, JCAP 11 (11), 048, arXiv:2108.10872 [astro-ph.CO].
- [44] M. Bartelmann, Class. Quant. Grav. 27, 233001 (2010), arXiv:1010.3829 [astro-ph.CO].
- [45] R. Takahashi, T. Suyama, and S. Michikoshi, Astron. Astrophys. 438, L5 (2005), arXiv:astro-ph/0503343.
- [46] P. Schneider, J. Ehlers, and E. E. Falco, <u>Gravitational Lenses</u>, Astronomy and Astrophysics Library (Springer, 1992).
- [47] V. Bozza, S. Capozziello, G. Iovane, and G. Scarpetta, Gen. Rel. Grav. 33, 1535 (2001), arXiv:gr-qc/0102068.
- [48] B. Bonga, J. Feldbrugge, and A. Ribes Metidieri, (2024), arXiv:2410.03828 [gr-qc].
- [49] R. Takahashi and T. Nakamura, Astrophys. J. 595, 1039 (2003), arXiv:astro-ph/0305055.
- [50] G. Tambalo, M. Zumalacárregui, L. Dai, and M. H.-Y. Cheung, Phys. Rev. D 108, 043527 (2023), arXiv:2210.05658 [gr-qc].
- [51] M. A. Oancea, J. Joudioux, I. Y. Dodin, D. E. Ruiz, C. F. Paganini, and L. Andersson, Phys. Rev. D 102, 024075 (2020), arXiv:2003.04553 [gr-qc].
- [52] M. A. Oancea, R. Stiskalek, and M. Zumalacárregui, Phys. Rev. D 109, 124045 (2024), arXiv:2209.06459 [gr-qc].
- [53] M. A. Oancea, R. Stiskalek, and M. Zumalacárregui, Mon. Not. Roy. Astron. Soc. 535, L1 (2024), arXiv:2307.01903 [gr-qc].
- [54] K.-i. Kubota, S. Arai, H. Motohashi, and S. Mukohyama, (2024), arXiv:2408.03289 [gr-qc].
- [55] G. Braga, A. Garoffolo, A. Ricciardone, N. Bartolo, and S. Matarrese, (2024), arXiv:2405.20208 [astro-ph.CO].
- [56] K.-i. Kubota, S. Arai, and S. Mukohyama, Phys. Rev. D 109, 044027 (2024), arXiv:2309.11024 [gr-qc].
- [57] H. Villarrubia-Rojo, S. Savastano, M. Zumalacárregui, L. Choi, S. Goyal, L. Dai, and G. Tambalo, (2024), arXiv:2409.04606 [gr-qc].
- [58] J. M. Ezquiaga and M. Zumalacárregui, Phys. Rev. D 102, 124048 (2020), arXiv:2009.12187 [gr-qc].
- [59] S. Goyal, A. Vijaykumar, J. M. Ezquiaga, and M. Zumalacarregui, Phys. Rev. D 108, 024052 (2023), arXiv:2301.04826 [gr-qc].
- [60] L. Andersson, J. Joudioux, M. A. Oancea, and A. Raj, Phys. Rev. D 103, 044053 (2021), arXiv:2012.08363 [grqc].
- [61] Y.-F. Wang, S. M. Brown, L. Shao, and W. Zhao, Phys. Rev. D 106, 084005 (2022), arXiv:2109.09718 [astroph.HE].
- [62] C. Dalang, P. Fleury, and L. Lombriser, Phys. Rev. D 102, 044036 (2020), arXiv:1912.06117 [gr-qc].
- [63] C. Dalang, P. Fleury, and L. Lombriser, Phys. Rev. D 103, 064075 (2021), arXiv:2009.11827 [gr-qc].
- [64] H. Takeda and T. Tanaka, (2024), arXiv:2404.10809 [gr-qc].

- [65] J. Streibert, H. O. Silva, and M. Zumalacárregui, (2024), arXiv:2404.07782 [gr-qc].
- [66] V. Faraoni and E. Gunzig, Astron. Astrophys. 332, 1154 (1998), arXiv:astro-ph/9801172.
- [67] A. Garoffolo, G. Tasinato, C. Carbone, D. Bertacca, and S. Matarrese, JCAP 11, 040, arXiv:1912.08093 [gr-qc].
- [68] K.-i. Kubota, S. Arai, and S. Mukohyama, Phys. Rev. D 107, 064002 (2023), arXiv:2209.00795 [gr-qc].
- [69] S. M. Koksbang and S. Räsänen, JCAP 04 (04), 030, arXiv:2108.06163 [astro-ph.CO].
- [70] G. Tasinato, A. Garoffolo, D. Bertacca, and S. Matarrese, JCAP 06, 050, arXiv:2103.00155 [gr-qc].
- [71] V. Faraoni, Astrophys. Lett. Commun. 35, 305 (1996), arXiv:astro-ph/9602154.
- [72] G. Cusin and M. Lagos, Phys. Rev. D 101, 044041 (2020), arXiv:1910.13326 [gr-qc].
- [73] C. Dalang, G. Cusin, and M. Lagos, Phys. Rev. D 105, 024005 (2022), arXiv:2104.10119 [gr-qc].
- [74] A. I. Harte, Gen. Rel. Grav. 51, 14 (2019), arXiv:1808.06203 [gr-qc].
- [75] A. I. Harte, Gen. Rel. Grav. 51, 160 (2019), arXiv:1906.10708 [gr-qc].
- [76] L. T. Santana, J. a. C. Lobato, M. O. Calvão, and R. R. R. Reis, Phys. Rev. D 110, 044031 (2024), arXiv:2407.04627 [gr-qc].
- [77] S. Goyal, K. Haris, A. K. Mehta, and P. Ajith, Phys. Rev. D 103, 024038 (2021), arXiv:2008.07060 [gr-qc].
- [78] C. Brans and R. H. Dicke, Phys. Rev. 124, 925 (1961).
- [79] V. Faraoni, Phys. Rev. D 59, 084021 (1999), arXiv:grqc/9902083.
- [80] E. E. Flanagan, Class. Quant. Grav. 21, 3817 (2004), arXiv:gr-qc/0403063.
- [81] M. Zumalacarregui, T. S. Koivisto, and D. F. Mota, Phys. Rev. D 87, 083010 (2013), arXiv:1210.8016 [astroph.CO].
- [82] D. Bettoni and S. Liberati, Phys. Rev. D 88, 084020 (2013), arXiv:1306.6724 [gr-qc].
- [83] M. Zumalacárregui and J. García-Bellido, Phys. Rev. D 89, 064046 (2014), arXiv:1308.4685 [gr-qc].
- [84] D. Bettoni and M. Zumalacárregui, Phys. Rev. D 91, 104009 (2015), arXiv:1502.02666 [gr-qc].
- [85] S. Capozziello, R. de Ritis, and A. A. Marino, Class. Quant. Grav. 14, 3243 (1997), arXiv:gr-qc/9612053.
- [86] V. Faraoni and E. Gunzig, Int. J. Theor. Phys. 38, 217 (1999), arXiv:astro-ph/9910176.
- [87] C. M. Will, Physical Review D 57, 2061 (1998).
- [88] L. Gondán and B. Kocsis, Mon. Not. Roy. Astron. Soc. 515, 3299 (2022), arXiv:2110.09540 [astro-ph.HE].
- [89] S. H. W. Leong, J. Janquart, A. K. Sharma, P. Martens, P. Ajith, and O. A. Hannuksela, (2024), arXiv:2408.13144 [astro-ph.HE].
- [90] M. Maggiore, (Oxford University Press, 2007) 10.1093/acprof:oso/9780198570745.001.0001.

- [91] C. W. Misner, K. S. Thorne, and J. A. Wheeler, <u>Gravitation</u> (W. H. Freeman, San Francisco, 1973).
- [92] M. Isi, Class. Quant. Grav. 40, 203001 (2023), arXiv:2208.03372 [gr-qc].
- [93] C. M. Will, (Cambridge University Press, 2018).
- [94] E. Poisson, (Cambridge University Press, 2009) 10.1017/CBO9780511606601.
- [95] P. Fleury, (2015), arXiv:1511.03702 [gr-qc].
- [96] J. a. C. Lobato and M. O. Calvão, Phys. Rev. D 109,

044004 (2024), arXiv:2402.02497 [gr-qc].

- [97] G. Cusin, C. Pitrou, and J.-P. Uzan, Phys. Rev. D 96, 103019 (2017), arXiv:1704.06184 [astro-ph.CO].
- [98] C. Bonvin, R. Durrer, and M. A. Gasparini, Phys. Rev. D 73, 023523 (2006), [Erratum: Phys.Rev.D 85, 029901 (2012)], arXiv:astro-ph/0511183.
- [99] V. Faraoni, Class. Quant. Grav. 26, 145014 (2009), arXiv:0906.1901 [gr-qc].