# Rational solutions of Painlevé V from Hankel determinants and the asymptotics of their pole locations

M. Balogoun<sup>†1</sup> M. Bertola<sup>†2</sup>,

<sup>†</sup> Department of Mathematics and Statistics, Concordia University 1455 de Maisonneuve W., Montréal, Québec, Canada H3G 1M8

#### Abstract

In this paper we analyze the asymptotic behaviour of the poles of certain rational solutions of the fifth Painlevé equation. These solutions are constructed by relating the corresponding tau function with a Hankel determinant of a certain sequence of moments. This approach was also used by one of the authors and collaborators in the study of the rational solutions of the second Painlevé equation. More specifically we study the roots of the corresponding polynomial tau function, whose location corresponds to the poles of the associated rational solution. We show that, upon suitable rescaling, the roots fill a well-defined region bounded by analytic arcs when the degree of the polynomial tau function tends to infinity. Moreover we provide an approximate location of these roots within the region in terms of suitable quantization conditions.

## Contents

1	Introduction	<b>2</b>
2	Semiclassical OPs and tau functions of Painlevé type2.1Special case of semiclassical functional of PV type2.2From Orthogonal Polynomials to Lax Pairs	<b>5</b> 6 7
3	Asymptotic analysis	12
4	Construction of the g-function         4.1 Outside the EoT: genus 0 g-function and effective potential         4.2 Inside the EoT: genus one g-function and effective potential	<b>13</b> 16 19
5	Deift-Zhou steepest descent analysis         5.1       Asymptotic analysis for s outside of the EoT         5.1.1       Conclusion of the steepest descent analysis         5.2       Asymptotic analysis inside the EoT         5.2.1       Second transformation, model problem and its solution: the case Re (s) > 0         5.2.2       Second transformation, model problem and its solution: the case Re (s) < 0	<ul> <li>20</li> <li>21</li> <li>24</li> <li>25</li> <li>27</li> <li>34</li> <li>37</li> <li>38</li> </ul>
6	Approximation of the Hamiltonian6.1Hamiltonian outside EoT	<b>38</b> 39
7	Conclusion	40
<b>A</b>	Airy parametrices         A.1 The Airy parametrix for a three main-arc intersection <sup>1</sup> Malik Balogoun@concordia.ca	<b>40</b> 41

<sup>2</sup>Marco.Bertola@concordia.ca

## 1 Introduction

The six Painlevé equations were classified by Painlevé and his student Gambier [36, 18] more than a century ago. This was a result of the search for second order ODEs in the complex plane whose solutions, roughly speaking, have the property that all movable singularities are isolated poles. This property has now become known, and referred to, as the *Painlevé property*.

While this might have remained a purely mathematical investigation, it was much later recognized that these equations have significant applications in mathematical physics, with the resurgence in the late '70s with the works connecting with Ising model and conformal field theory [32, 33]. Another momentous resurgence happened in the '90s when Tracy and Widom [39] used a special second Painlevé transcendent (the Hastings-McLeod solution [20]) to describe the fluctuations of the larges eigenvalue of a large random Hermitean matrix.

Amongst special solutions of the Painlevé equation the rational ones attract a natural interest; the literature is extensive and seems to start with [42] who discussed rational solution of the second Painlevé equation and defined a special sequence of polynomials that are now called *Vorob'ev-Yablonskii* after their discoverers (it appears that Yablonskii defined them slightly earlier but the reference is difficult to find [43]). Rational solutions also appear in semiclassical limits of integrable PDEs; in the one-dimensional sine-Gordon equation near a separatrix, for example, one finds that a suitable scaling of the solution is expressible in terms of a rational solution of the second Painlevé equation [9].

Rational solutions exist for all but the first Painlevé equation: although there does not seem to be a full and complete classification in all six cases, for particular cases either the full classification of rational solutions exist or there are constructions of special families of rational solutions (for the Painlevé II [42, 43, 24], for the Painlevé III,V, VI [31, 30, 40] Painlevé IV [35]

The literature that investigates the asymptotic behaviour of the rational solutions, and the pole distribution thereof, is more recent, probably due to the interest spurred by numerical investigations and the appearance of well defined patterns; for the zeros of Okamoto polynomials (which are poles of rational solutions of PIV) see [34], for the zeros of Vorobev–Yablonskii polynomials and Painlevé II see [10, 11, 2], for the second Painlevé hierarchy see [1].

The approach to asymptotic analysis relies on the formulation of an associated Riemann–Hilbert problem, namely, a boundary value problem for a piecewise analytic matrix valued function matrix. Within this framework there are two logical distinct approaches that can be used. We can categorize them under the following banners:

- 1. the isomonodromic approach;
- 2. the orthogonal polynomial (OP) approach.

The isomonodromic approach relies on the general fact that any Painlevé equation appears as the compatibility between a  $2 \times 2$  system of ODEs with rational coefficient in the complex plane and an additional PDE in an auxiliary parameter [21]. The different solutions are parametrized by (generalized) monodromy data of the ODE, which is the starting point for the Riemann–Hilbert analysis. Typically the degree of the rational solution appears explicitly as one of the parameters in the monodromy data, and can be used as large parameter in the asymptotics. This is the philosophy behind the works [10, 11].

The second approach was used, possibly for the first time, in [2] and then also applied to the generalized Vorob'ev–Yablonski polynomials in [1]. It is also the approach we follow in this paper. The main connection between OPs and equations of Painlevé type was established in [3], where it was shown that Hankel determinants built out of the moments of "semiclassical" moment functionals are always isomonodromic tau functions in the sense of Jimbo-Miwa-Ueno [21]. It was a remark (Rem 5.3 in [3]) that special choices of semiclassical moment functionals lead automatically to tau functions of Painlevé equations (all, except possibly for Painlevé I). In genereal, however, these solutions correspond to transcendental solutions like, for example, the solutions of PII constructed out of determinants of derivatives of Airy functions, see [23].

It is possible to further restrict the setup of orthogonal polynomials in such a way that the moments of the moment functional become *polynomials* in a parameter, which then guarantees that the Hankel determinant (automatically an isomonodromic tau function) is a polynomial tau function of an equation of Painlevé type. This is what works "behind the scenes" in [2].

The advantage of this reformulation in terms of associated Orthogonal Polynomials is that there is a solid and well developed framework for studying their large degree asymptotics, with an extensive literature that starts with the seminal work of Deift et al. [15].

Before going into any further detail let us discuss the known literature and results about the rational solutions of the fifth Painlevé equation.

**Rational solutions of PV.** The fifth Painlevé equation is the following nonlinear, second order ODE in the complex domain for the unknown function y(t)

$$y'' = \left(\frac{1}{2y} + \frac{1}{y-1}\right)(y')^2 - \frac{y'}{t} + \frac{(y-1)^2}{t^2}\left(\alpha y + \frac{\beta}{y}\right) + \frac{\gamma y}{t} + \frac{\delta y(y+1)}{y-1},$$
(1.1)

where  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$  are parameters: we shall refer to (1.1) as  $P_5(\alpha, \beta, \gamma, \delta)$ . The equation admits certain symmetries that change the value of the parameters; if y(t) is a solution of  $P_5(\alpha, \beta, \gamma, \delta)$  then

- 1. y(-t) is a solution of  $P_5(\alpha, \beta, -\gamma, \delta)$ ;
- 2.  $\frac{1}{u(t)}$  is a solution of  $P_5(-\beta, -\alpha, -\gamma, \delta)$ ;
- 3.  $y(\lambda t)$  is a solution of  $P_5(\alpha, \beta, \lambda\gamma, \lambda^2 \delta)$ , for any  $\lambda \in \mathbb{C}^{\times}$ .

Using the above symmetries the analysis is reduced to only two families; the family where  $\delta = 0$  (which is called "degenerate" and can be reduced to Painlevé III) and the case  $\delta \neq 0$  which, by virtue of the last of the above symmetries, is customarily set to  $\delta = -\frac{1}{2}$ .

The classification of rational solution is contained in [25] where the authors show that rational solutions exist only if the parameters satisfy certain relations. More precisely (paraphrasing and condensing their results)

**Theorem 1.1 ([25], Thm. 1.1, Thm. 1.2)** The equation  $P_5(\alpha, \beta, \gamma, -\frac{1}{2})$  (1.1) admits rational solutions if and only if there are integers  $k, m \in \mathbb{Z}$  such that

- (I)  $\alpha = \frac{1}{2}(\gamma + k)^2$ ;  $\beta = -\frac{m^2}{2}$ , k + m odd, and  $\alpha \neq 0$  when |k| < m;
- (II)  $\alpha = \frac{1}{2}m^2$ ;  $\beta = -\frac{(\gamma+k)^2}{2}$ , k+m odd, and  $\beta \neq 0$  when |k| < m;
- (III)  $\beta = -\frac{1}{2}(\alpha_1 + m)^2, \gamma = k$  with  $\alpha_1^2 = 2\alpha$  so that  $m \ge 0$  and k + m even;
- (IV)  $\alpha = \frac{k^2}{4}, \ \beta = -\frac{m^2}{8}, \ \gamma \notin \mathbb{Z}$  where k, m > 0 and k, m both odd.

In cases (I) and (II) the solution is unique if  $\gamma \notin \mathbb{Z}$  and there are at most two rational solutions otherwise.

Observe that the cases (I) and (II) of the above result are really the same family up to the application of the symmetry  $y(t) \leftrightarrow \frac{1}{y(-t)}$  which transforms solution of  $P_5(\alpha, \beta, \gamma, -\frac{1}{2})$  into solutions of  $P_5(-\alpha, -\beta, \gamma, -\frac{1}{2})$ . In the recent [13] the authors construct the rational solutions corresponding to the case (II) of [25] above

In the recent [13] the authors construct the rational solutions corresponding to the case (II) of [25] above (which is case (i) in Thm. 4.1 of [13]). More precisely they construct the tau functions

$$\tau_{m,n}^{(\mu)} := \det\left[\left(t\frac{\mathrm{d}}{\mathrm{d}t}\right)^{j+k} L_{m+n}^{(n+\mu)}(t)\right]_{j,k=0}^{n-1}$$
(1.2)

where  $L_n^{(\alpha)}(t)$  are the associated Laguerre polynomials (see DLMF: 18.5.12)

$$L_n^{(\alpha)}(t) = \sum_{\ell=0}^n \frac{(\alpha+\ell+1)_{n-\ell}}{(n-\ell)!\,\ell!} (-x)^\ell = \frac{(\alpha+1)_n}{n!} {}_1F_1\left( {-n \atop \alpha+1}; t \right).$$
(1.3)

Then their result implies that (1.2) is the tau function of  $P_5\left(\frac{m^2}{2}, -\frac{(m+2n+1+\mu)^2}{2}, \mu, -\frac{1}{2}\right)$ . Many interesting properties are discussed but no description of the asymptotic behaviour for large degree is undertaken.

**Results.** In the present paper we will provide an alternative description for the same tau function; more precisely we construct the tau function for case (I) in Thm. 1.1 corresponding to the parameters

 $\alpha = 3N + \mathbf{K} - 1, \quad m = N + \mathbf{K}, \quad \gamma = \boldsymbol{\rho}, \quad N \in \mathbb{N}, \quad \mathbf{K} \in \mathbb{Z}, \quad \boldsymbol{\rho} \in \mathbb{C}, \tag{1.4}$ 

which corresponds to the parameters  $\mu, n, m$  of [13] as follows

$$m = N + K, \quad n = N - 1, \quad \mu = 1 - 2N - K + \rho.$$
 (1.5)

To construct this  $\tau$ -function we consider the following sequence of *moments* depending on the parameters  $K \in \mathbb{Z}, \rho \in \mathbb{C}$ 

$$\mu_j(t) := \oint_{|z|=1} z^j z^K \left( 1 - \frac{1}{z} \right)^{\mu} e^{\frac{t}{z}} dz, \qquad j = 0, 1, 2, \dots$$
(1.6)

We interpret the  $\mu_j$ 's as the moments of the complex measure  $d\mu(z) = z^K \left(1 - \frac{1}{z}\right)^{\rho} e^{\frac{t}{z}} dz$  on |z| = 1. Observe that  $\mu_j(t)$  are polynomials in t, and can also be obtained from a generating function

$$\sum_{j=0}^{\infty} \mu_j(t) \frac{s^j}{j!} = \oint_{|z|=1} z^K \left( 1 - \frac{1}{z} \right)^{\rho} e^{\frac{t}{z} + sz} dz$$
(1.7)

or also (by replacing  $z = \frac{1}{w}$  in the definition of moments (1.6) and then interpreting them as residues)

$$F(s;t) := -2i\pi s^{-K-1} (1-s)^{\rho} e^{st} = \sum_{j=0}^{\infty} \mu_j(t) \frac{s^j}{j!}$$
(1.8)

The map that associates to any polynomial p(z) the value  $\mathcal{M}[p] := \oint_{|z|=1} p(z) d\mu(z)$  is an example of *semiclassical moment functional* [27, 28, 29, 37] and thus fits naturally in the general theory of [3] which guarantees that the Hankel determinant

$$\tau_n(t; \mathbf{K}, \boldsymbol{\rho}) := \det \left[ \mu_{a+b-2}(t) \right]_{a,b=1}^n, \tag{1.9}$$

is a tau function of an equation of Painlevé type. The matching with PV is explained in Section 2.2. More explicitly, from the identification it will follow that it satisfies the  $\sigma$ -form of the PV equation ([22], App. C) in the following way: let us define the quantities

$$H_V := \frac{\mathrm{d}}{\mathrm{d}t} \ln \tau_n(t; \boldsymbol{K}, \boldsymbol{\rho}) + \frac{\boldsymbol{\rho}}{2}$$
(1.10)

$$\sigma(t) := tH_V + \frac{t}{2}(\theta_0 + \theta_\infty) + \frac{(\theta_0 + \theta_\infty)^2 - \theta_1^2}{4}$$
(1.11)

$$\theta_0 := 2n + \mathbf{K}; \qquad \theta_1 := -\boldsymbol{\rho}; \qquad \theta_\infty := \boldsymbol{\rho} - \mathbf{K}.$$
 (1.12)

Then the  $\sigma$ -form of PV is the following ODE for  $\sigma$  defined in (1.11)

$$\left(t\frac{\mathrm{d}^{2}\sigma}{\mathrm{d}t^{2}}\right)^{2} = \left(\sigma - t\frac{\mathrm{d}\sigma}{\mathrm{d}t} + 2\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)^{2} - (\theta_{\infty} + 2\theta_{0})\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)^{2} + -4\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \frac{\theta_{0} - \theta_{1} + \theta_{\infty}}{2}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \theta_{0}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \frac{\theta_{0} + \theta_{1} + \theta_{\infty}}{2}\right)$$
(1.13)

In the second part of the paper we exploit the connection between  $\tau_n(t)$  and Hankel determinants to describe the asymptotic location of its zeros as  $n \to \infty$ .

More precisely we describe the asymptotic behaviour of the zeros of the polynomials

$$P_n(s) := \tau_n(ns; \mathbf{K}, \boldsymbol{\rho}), \quad \mathbf{K} \in \mathbb{Z}, \quad \boldsymbol{\rho} \notin \mathbb{Z}, \quad n \to \infty,$$
(1.14)

in the s-plane (these zeroes are the same as the zeroes of  $\tau_n$  but homothetically rescaled by a factor 1/n). We observe that we do not consider a double-scaling limit where the either one of the parameters K,  $\rho$  or



Figure 1: The zeros of several instances of polynomial tau functions (all for K = 0).

both are proportional to  $n \text{ as } n \to \infty$ . While the general setup is conducive to such an analysis, the details of the construction of the *g*-function would have to be changed significantly and this is deferred to a future publication. The logic of the analysis is parallel to the one employed in [2, 1] and the main goal is to explain the emerging shape which clearly appears, see Fig. 1

In the figure an almond-shaped region, which we call "Eye of the Tiger" (EoT), will be shown to be the asymptotic region where all zeros of  $P_n(s)$  (1.14) lie. The boundary of the region (marked in black) is determined by the implicit equation in the *s*-plane

$$2\ln\left|\frac{2i}{s} + i\sqrt{\frac{4}{s^2} + 1}\right| - 2\operatorname{Re}\left(\sqrt{1 + \frac{s^2}{4}}\right) = 0 \tag{1.15}$$

which follows from (4.20), (4.26) below. Within the EoT the zeros of  $P_n(s)$  (1.14) arrange themselves in a semi-regular lattice. We give a description of this lattice in terms of a "quantization condition" described in Section 5.2.4, in particular (5.98) (for Re (s) > 0) or (5.122) (for Re (s) < 0). The two quantization conditions (5.98), (5.122) determine an asymptotic grid (shown in Fig. 2 in green and blue thin lines) at whose vertices the zeroes are approximately located. We do not estimate rigorously the rate of convergence, but the numerical evidence is quite striking even for relatively small values of n.

## 2 Semiclassical OPs and tau functions of Painlevé type

Let us recall the notion of semiclassical orthogonal polynomials (SOPs) [27, 28, 29, 37].

**Definition 2.1** Given a pair of polynomials (A, B), with B monic, a semiclassical moment functional of type (A, B) is a linear map  $\mathcal{M} : \mathbb{C}[z] \to \mathbb{C}$  on the space of polynomials in the indeterminate variable z such that for all  $p \in \mathbb{C}[z]$ 

$$\mathcal{M}[A(z)p(z)] = \mathcal{M}[B(z)p'(z)].$$
(2.1)

A polynomial  $p_n(z)$  of degree n is called orthogonal for the moment functional  $\mathcal{M}$  if

$$\mathcal{M}[p_n(z)z^j] = 0, \quad j = 0, 1, \dots, n-1.$$
 (2.2)

Given a moment functional  $\mathcal{M}$  its *moments* are the (in general complex) numbers

$$\mu_j := \mathcal{M}[z^j]. \tag{2.3}$$

It is possible to show that these moment functionals admit an integral representation as follows: define the  $symbol^3$  as the function  $\theta$  such that

$$\theta'(z) := -\frac{B'(z) + A(z)}{B(z)}.$$
(2.4)

Then we can express a semiclassical moment functional of type (A, B) in terms of an integral of the form

$$\mathcal{M}[p(z)] = \int_{\gamma} p(z) \mathrm{e}^{\theta(z)} \mathrm{d}z.$$
(2.5)

Here the contour of integration can be chosen in several homology classes, each providing a linearly independent moment functional of type (A, B). The allowable contours have the defining property that the integration of the right side of (2.1) can be performed by parts and with vanishing boundary terms. It is known that there are  $d = \max\{\deg A, \deg B - 1\}$  independent such homology classes (generically) and a description of them can be found, for example in [3]. It was also shown in loc.cit. that semiclassical moment functionals are inextricably related with the theory of isomonodromic deformations, and hence in particular with the theory of Painlevé equations, as we briefly recall.

Indeed it was shown in [3] that any deformation of the coefficients of A, B that preserves all the residues of  $\theta'$  is an *isomonodromic deformation* for a suitably defined differential equation in the complex plane of rank 2. It was determined by the Japanese school several decades ago [21, 22] that to any such isomonodromic deformation we can associate a tau function, namely a function of the isomonodromic deformation parameters (i.e. of the coefficients of A, B in this case). While this is not an appropriate venue to review all applications of tau functions, suffices to say here (see Rem. 5.3 of [3]) that by appropriate choices of the symbol and corresponding contours of integration  $\gamma$  we can construct tau function for all the Painlevé equations II–VI. We will make below one such choice which has the additional property of producing *polynomial* moments, and hence polynomial tau functions.

### 2.1 Special case of semiclassical functional of PV type

We specialize the previous description to the case of the following symbol depending on the parameters  $t, \mathbf{K}, \boldsymbol{\rho}$ 

$$\theta(z) := \frac{t}{z} + \rho \ln \left( 1 - \frac{1}{z} \right) + \mathbf{K} \ln z \quad \mathrm{e}^{\theta(z)} = \left( 1 - \frac{1}{z} \right)^{\rho} z^{\mathbf{K}} \mathrm{e}^{\frac{t}{z}}.$$
(2.6)

In (2.6) all the logarithms are principal and  $\theta$  is analytic in  $\mathbb{C} \setminus (-\infty, 1]$ . Since K,  $\rho$  are the residues of  $\theta' dz$ , the only isomonodromic parameter is t in the above expression. Note that if  $K \in \mathbb{Z}$  (which is going to be our main focus shortly) the weight function  $e^{\theta(z)}$  is analytic in  $\mathbb{C} \setminus [0, 1]$ . Furthermore we will avail ourselves of the freedom of deforming the segment [0, 1] to a smooth arc with the same endpoints when needed. The corresponding moment functionals, in the terminology of Def. 2.1 above, are of type

$$(A,B) = \left(\boldsymbol{K}\boldsymbol{z}^{2} + (\boldsymbol{\rho} + \boldsymbol{K} + t)\boldsymbol{z} + t, \ \boldsymbol{z}^{2}(\boldsymbol{z} - 1)\right)$$
(2.7)

We can choose the contours of integration for defining the specific moment functional as follows:

<sup>&</sup>lt;sup>3</sup>We adopt the terminology common in the literature on Töplitz operators.

- 1.  $\gamma$  a closed contour originating at z = 0 along the direction  $-\arg(-t)$ , looping around z = 0 and terminating at z = 0 along the same direction and leaving z = 1 in the inside.
- 2.  $\tilde{\gamma}$  the same as above but leaving z = 1 on the outside.

There are some special instances, depending on the values of the parameters  $K, \rho$ ; if  $\rho = 0, 1, 2, ...$ , then we can replace  $\tilde{\gamma}$  with a contour terminating at z = 1.

Most important for our consideration is the case  $K \in \mathbb{Z}, \rho \notin \mathbb{Z}$ ; in these cases, we can replace  $\gamma$  with the homotopy class of a circle of radius R > 1 in  $\mathbb{C} \setminus [0, 1]$ .

The latter case is relevant for the construction of rational solutions of the fifth Painlevé equation: indeed, if  $K \in \mathbb{Z}$  the moments are *polynomials* in the variable t

$$\mu_j(t) = \oint_{|z|=R} z^j e^{\theta(z)} dz, \quad \deg \mu_j(t) = j - K - 1.$$
(2.8)

If  $j \leq K-2$ , the moment  $\mu_j$  vanishes identically because the integration correspond to  $2i\pi$  times the residue at infinity, which vanishes by inspection.

We consider the Hankel determinants:

$$D_n(t) := \det \left[ \mu_{a+b-2}(t) \right]_{a,b=1}^n.$$
(2.9)

They are also polynomials of degree at most

$$\deg D_n(t) \le n(n - \mathbf{K} - 2). \tag{2.10}$$

The results of [3] imply that these Hankel determinants are tau functions of an isomonodromic system, and we are going to identify this with the Lax pair of the fifth Painlevé equation.

### 2.2 From Orthogonal Polynomials to Lax Pairs

The bridge is provided by the Riemann–Hilbert formulation of OP [17] which, in our case reads as follows.

**Problem 2.1 (RHP**<sub>Y</sub>) Given  $n \in \mathbb{N}$ , find a 2 × 2 matrix-valued function  $Y(z) = Y_n(z)$  such that

- 1. Y(z) and  $Y^{-1}(z)$  are holomorphic and bounded in  $\mathbb{C} \setminus \gamma$ ;
- 2. The boundary values along  $z \in \gamma$  satisfy

$$Y(z_{+}) = Y(z_{-}) \begin{bmatrix} 1 & e^{\theta(z)} \\ 0 & 1 \end{bmatrix}, \qquad \forall z \in \gamma$$
(2.11)

with  $\theta$  as in (2.6).

3. As  $z \to \infty$  the matrix  $Y_n(z)$  admits an asymptotic expansion of the form

$$Y_n(z) = \left(\mathbf{1} + \sum_{\ell=1}^{\infty} \frac{Y_n^{(\ell)}}{z^{\ell}}\right) z^{n\sigma_3}, \quad \sigma_3 := \begin{bmatrix} 1 & 0\\ 0 & -1 \end{bmatrix}$$
(2.12)

where the coefficient matrices  $Y_n^{(\ell)}$  are independent of z.

In the following theorem we condense the essential results that are at the core of much of the theory of orthogonal polynomial. The theorem is mostly the reformulation of results of [17] together with notational adaptations to the current context.

**Theorem 2.1** ([17]) Consider the RHP 2.1;

1. if the solution exists it is unique and det  $Y_n(z) \equiv 1$ ;

2. the solution exists if and only if the Hankel determinant  $D_n$  in (2.9) is different from zero;

\_

3. the solution has the form

$$Y_n(z) = \begin{bmatrix} p_n(z) & \oint_{\gamma} \frac{p_n(w)e^{\theta(w)}dw}{(w-z)2i\pi} \\ \tilde{p}_{n-1}(z) & \oint_{\gamma} \frac{\tilde{p}_{n-1}(w)e^{\theta(w)}dw}{(w-z)2i\pi} \end{bmatrix}$$
(2.13)

where  $p_n(z)$  is the monic orthogonal polynomial of degree n and  $\tilde{p}_{n-1}$  a polynomial of degree deg  $\tilde{p}_{n-1} \leq n-1$ ; they are expressible explicitly in terms of the moments as follows

$$p_{n}(z) = \frac{1}{D_{n}} \det \begin{bmatrix} \mu_{0} & \mu_{1} & \mu_{2} & \cdots & \mu_{n} \\ \mu_{1} & \mu_{2} & \cdots & & \mu_{n+1} \\ \mu_{2} & \cdots & & & \mu_{n+2} \\ \vdots & & & & \\ \mu_{n-1} & \mu_{n} & \cdots & & \mu_{2n-1} \\ 1 & z & z^{2} & \cdots & z^{n} \end{bmatrix}$$
(2.14)  
$$\widetilde{p}_{n-1}(z) = \frac{-2i\pi}{D_{n}} \det \begin{bmatrix} \mu_{0} & \mu_{1} & \cdots & \mu_{n-1} \\ \mu_{1} & \mu_{2} & \cdots & \mu_{n} \\ \mu_{2} & \cdots & & \mu_{n+1} \\ \vdots & & & \\ \mu_{n-2} & \cdots & & \mu_{2n-3} \\ 1 & z & \cdots & z^{n-1} \end{bmatrix}$$
(2.15)

with  $D_n$  the Hankel determinant of moments (2.9).

The matrix  $Y_n(z;t)$  (we highlight its dependence on t as well) satisfies a first order ODE; indeed let us define

$$\Psi(z;t) := Y_n(z;t) e^{\frac{1}{2}\theta(z)\sigma_3} = Y_n(z;t) z^{\frac{\kappa}{2}\sigma_3} \left(1 - \frac{1}{z}\right)^{\frac{\mu}{2}\sigma_3} e^{\frac{t}{2z}\sigma_3}.$$
(2.16)

We note that near  $z = 0, 1, \infty$ , the matrix  $\Psi$  has the following expansion, which follows from the properties of  $Y_n$  and the definition of  $\theta$  (2.6)

$$\Psi(z;t) = \begin{cases} \mathcal{O}^{\times}(1)z^{\frac{K-\rho}{2}\sigma_3}\mathrm{e}^{\frac{t}{2z}\sigma_3}, & z \to 0\\ \mathcal{O}^{\times}(1)(z-1)^{\frac{\rho}{2}\sigma_3}, & z \to 1\\ \left(\mathbf{1} + \mathcal{O}(z^{-1})\right)z^{\left(n + \frac{K}{2}\right)\sigma_3}, & z \to \infty \end{cases}$$
(2.17)

where  $\mathcal{O}^{\times}(1)$  means a locally analytic and invertible matrix.

**Proposition 2.1** The matrix  $\Psi(z;t)$  in (2.16) satisfies the pair of first order PDEs

$$\begin{cases} \partial_z \Psi(z;t) = A(z;t)\Psi(z;t) \\ \partial_t \Psi(z;t) = B(z;t)\Psi(z;t). \end{cases}$$
(2.18)

where the matrices A, B have the form

$$A(z;t) := -\frac{t}{z^2} G_0 \sigma_3 G_0^{-1} + G_0 \left( \frac{K - \rho}{z} \sigma_3 - \frac{t}{z} \Big[ G_0^{-1} G_1, \sigma_3 \Big] \right) G_0^{-1} + \frac{K}{z - 1} H_0 \sigma_3 H_0^{-1} + \frac{n + K}{z} \sigma_3 \quad (2.19)$$

$$B(z;t) := \frac{1}{z} G_0 \sigma_3 G_0^{-1} \tag{2.20}$$

**Proof.** It follows from the RHP 2.1 that  $\Psi$  satisfies a boundary value problem

$$\Psi(z_{+};t) = \Psi(z_{-};t) \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$
(2.21)

Differentiating the above relation with respect to z on both sides we conclude that both  $\Psi$  and  $\partial_z \Psi$  satisfy the same jump relation and hence the matrix

$$A(z;t) := \partial_z \Psi(z;t) \Psi^{-1}(z;t)$$

$$(2.22)$$

extends analytically to an analytic function with at most isolated singularities at z = 0, 1. We thus have

$$A(z;t) = Y'_n Y_n^{-1} + Y_n \left(\frac{-t}{z^2} + \frac{\rho}{z(z-1)} + \frac{K}{z}\right) \sigma_3 Y_n^{-1}.$$
(2.23)

This shows that the only singularities of A(z) are a double pole at z = 0 and a simple pole at z = 1; near  $z = \infty$ , using the asymptotic behaviour of  $Y_n$  as in (2.12), we conclude that

$$A(z;t) = -\frac{t}{z^2}G_0\sigma_3G_0^{-1} + G_0\left(\frac{K-\rho}{z}\sigma_3 - \frac{t}{z}\left[G_0^{-1}G_1,\sigma_3\right]\right)G_0^{-1} + \frac{K}{z-1}H_0\sigma_3H_0^{-1} + \frac{n+K}{z}\sigma_3 \quad (2.24)$$
$$G_0 := Y_n(0), \quad G_1 := Y'_n(0); \quad H_0 := Y_n(1).$$

A similar argument produces the expression for B(z; t).

In the pair of PDE's (2.18) the z-equation is an ODE (considering t as parameter) with an irregular singularity of Poincaré rank 2 at z = 0 and two Fuchsian singularities at  $z = 1, \infty$ , respectively.

**Comparison with the Japanese Lax Pair.** In the work of the Japanese school [22] the z-component of the Lax pair for Painlevé V has also two Fuchsian and one second rank singularities, but with the positions reversed. More specifically, the Lax pair proposed in [22] is as follows<sup>4</sup> (see formulas (C.38–C.45) in loc. cit.)

$$\begin{aligned} \partial_{\zeta} \Phi(\zeta;t) &= A_{JMU}(\zeta;t) \Phi(\zeta;t) , \qquad \partial_{t} \Phi(\zeta;t) = B_{JMU}(\zeta;t) \Phi(\zeta;t) \\ A_{JMU}(\zeta;t) &= \frac{t}{2} \sigma_{3} + \frac{1}{\zeta} \begin{bmatrix} Z + \frac{\theta_{0}}{2} & -U(Z + \theta_{0}) \\ \frac{Z}{U} & -Z - \frac{\theta_{0}}{2} \end{bmatrix} + \frac{1}{\zeta - 1} \begin{bmatrix} -Z - \frac{\theta_{0} + \theta_{\infty}}{2} & UY\left(Z + \frac{\theta_{0} - \theta_{1} + \theta_{\infty}}{2}\right) \\ -\frac{Z + \frac{\theta_{0} + \theta_{1} + \theta_{\infty}}{2}}{UY} & Z + \frac{\theta_{0} + \theta_{\infty}}{2} \end{bmatrix} \\ B_{JMU}(\zeta;t) &= \frac{\zeta}{2} \sigma_{3} + \begin{bmatrix} 0 & -\frac{U}{t}\left(Z + \theta_{0} - Y\left(Z + \frac{\theta_{0} - \theta_{1} + \theta_{\infty}}{2}\right)\right) \\ \frac{1}{tUY}\left((Y - 1)Z + \frac{\theta_{0} + \theta_{1} + \theta_{\infty}}{2}\right) & 0 \end{bmatrix}$$
(2.25)

where Z = Z(t), Y = Y(t), U = U(t) satisfy a nonlinear first order system of ODEs in t (C.40 in loc. cit.), which implies the fifth Painlevé equation for Y:

$$\frac{\mathrm{d}^2 Y}{\mathrm{d}t^2} = \left(\frac{1}{2Y} + \frac{1}{Y-1}\right) \left(\frac{\mathrm{d}Y}{\mathrm{d}t}\right)^2 - \frac{1}{t} \frac{\mathrm{d}Y}{\mathrm{d}t} + \frac{(Y-1)^2(\alpha Y + \frac{Y}{\beta})}{t^2} + \frac{\gamma Y}{t} + \frac{\delta Y(Y+1)}{Y-1}$$
$$\alpha = \frac{1}{2} \left(\frac{\theta_0 - \theta_1 + \theta_\infty}{2}\right)^2; \ \beta = -\frac{1}{2} \left(\frac{\theta_0 - \theta_1 - \theta_\infty}{2}\right)^2; \ \gamma = 1 - \theta_0 - \theta_1; \ \delta = -\frac{1}{2}.$$
(2.26)

The solution  $\Phi(\zeta; t)$  has the following formal expansions near  $\zeta = 0, 1, \infty$ :

$$\Phi(\zeta;t) = \mathcal{O}(1)\zeta^{\frac{\theta_0\sigma_3}{2}} , \qquad \zeta \to 0$$

$$\Phi(\zeta;t) = \mathcal{O}(1)(\zeta - 1)^{\frac{1}{2}}, \qquad \zeta \to 1$$

$$\Phi(\zeta;t) = \left(\mathbf{1} + \frac{\Phi_1}{\zeta} + \mathcal{O}(\zeta^{-2})\right) \zeta^{-\frac{\theta_\infty}{2}\sigma_3} e^{\frac{t}{2}\zeta\sigma_3} ,\qquad \qquad \zeta \to \infty.$$
(2.27)

The Hamiltonian function for the Painlevé equation is given by

$$H_V = -\frac{1}{2} \operatorname{tr}(\Phi_1 \sigma_3) \tag{2.28}$$

<sup>&</sup>lt;sup>4</sup>We transcribe the results of [22] but we adapt their notation to our conventions. Note that the paper contains a couple of small typos: in (C.39) there should be an x in front of the first term of B and the sign of the (1, 2) entry of the next term should be the opposite.

and the equation admits the so-called sigma-form: indeed, introducing the new function

$$\sigma(t) = tH_V - \frac{t}{2}(\theta_0 + \theta_\infty) + \frac{(\theta_0 + \theta_\infty)^2 - \theta_1^2}{4},$$
(2.29)

it can be verified that it satisfies ([22] formula (C.45))

$$\left(t\frac{\mathrm{d}^{2}\sigma}{\mathrm{d}t^{2}}\right)^{2} = \left(\sigma - t\frac{\mathrm{d}\sigma}{\mathrm{d}t} + 2\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)^{2} - \left(\theta_{\infty} + 2\theta_{0}\right)\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)^{2} + -4\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \frac{\theta_{0} - \theta_{1} + \theta_{\infty}}{2}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \theta_{0}\right)\left(\frac{\mathrm{d}\sigma}{\mathrm{d}t} - \frac{\theta_{0} + \theta_{1} + \theta_{\infty}}{2}\right).$$
(2.30)

In order to identify our Lax pair (2.18) with the Japanese one (2.25) it suffices to map  $\zeta = \frac{1}{z}$  and suitably normalize our matrix  $\Psi(z;t)$ .

**Lemma 2.1** The map  $\zeta = \frac{1}{z}$  and

$$\Phi(\zeta;t) = Y_n(0;t)^{-1} \Psi\left(\frac{1}{\zeta};t\right) e^{-i\frac{\pi}{2}\rho\sigma_3}$$
(2.31)

transforms the Lax pair (2.18) into (2.25) with parameters

 $\theta_0 = -2n - \mathbf{K}; \qquad \theta_1 = \boldsymbol{\rho}; \qquad \theta_\infty = \mathbf{K} - \boldsymbol{\rho}.$  (2.32)

This corresponds to the parameters  $\alpha, \beta, \gamma$  in (2.26) as follows

$$\alpha = \frac{(N+\rho)^2}{2}; \quad \beta = -\frac{(N+K)^2}{2}; \quad \gamma = 1+2N+K-\rho.$$
(2.33)

**Proof.** The map  $\zeta = \frac{1}{z}$  maps z = 0 to  $\zeta = \infty$ ,  $z = \infty$  to  $\zeta = 0$  and z = 1 to  $\zeta = 1$ . Thus the exponents of (formal) monodromy  $\theta_{\{0,1,\infty\}}$  are the read off by matching the exponents in (2.17) and (2.27).

The above lemma allows us to identify the Hamiltonian with the logarithmic derivative of the Hankel determinant and, thus, the  $\tau$  function with the Hankel determinant itself. This is, a priori, a result of [3], but we can here derive it directly from the formulas already reported.

**Proposition 2.2** The Hankel determinant  $D_n(t)$  in (2.9) is a polynomial tau function of the fifth Painlevé equation. In particular the Hamiltonian is

$$H_V = -\frac{1}{2} \operatorname{tr} \left( Y_n^{-1}(0) Y_n'(0) \sigma_3 \right) + \frac{\rho}{2} = -\left( Y_n^{-1}(0) Y_n'(0) \right)_{11} + \frac{\rho}{2} = \frac{\mathrm{d}}{\mathrm{d}t} \ln D_n(t) + \frac{\rho}{2}.$$
 (2.34)

**Proof.** We need to identify  $\Phi_1$  in the expansion at  $\zeta = \infty$  in (2.27) with the suitable expansion of  $\Psi$  at z = 0. Recalling the definition of  $\Psi$  (2.16) we see that

$$Y_{n}(0)^{-1}\Psi\left(\frac{1}{\zeta};t\right)e^{-\frac{i\pi}{2}\rho\sigma_{3}} = Y_{n}(0)^{-1}\left(Y_{n}(0) + \frac{1}{\zeta}Y_{n}'(0) + \mathcal{O}(\zeta^{-2})\right)\left(\zeta - 1\right)^{\frac{\rho}{2}\sigma_{3}}\zeta^{-\frac{\kappa}{2}\sigma_{3}}e^{\frac{i}{2}\zeta\sigma_{3}} = \\ = \left(\mathbf{1} + \frac{Y_{n}(0)^{-1}Y_{n}'(0)}{\zeta} + \mathcal{O}(\zeta^{-2})\right)\left(\mathbf{1} - \frac{\rho}{2\zeta}\sigma_{3} + \mathcal{O}(\zeta^{-2})\right)\zeta^{\frac{\rho-\kappa}{2}\sigma_{3}}e^{\frac{i}{2}\zeta\sigma_{3}} = \\ = \left(\mathbf{1} + \frac{Y_{n}(0)^{-1}Y_{n}'(0) - \frac{\rho}{2}\sigma_{3}}{\zeta} + \mathcal{O}(\zeta^{-2})\right)\zeta^{\frac{\rho-\kappa}{2}\sigma_{3}}e^{\frac{i}{2}\zeta\sigma_{3}}.$$
(2.35)

Thus we conclude that

$$\Phi_1 = Y_n(0)^{-1} Y'_n(0) - \frac{\rho}{2} \sigma_3.$$
(2.36)

Then, according to (2.28), we must compute

$$H_{V} = -\frac{1}{2} \operatorname{tr} \left( \Phi_{1} \sigma_{3} \right) = -\frac{1}{2} \operatorname{tr} \left( Y_{n}(0)^{-1} Y_{n}'(0) \sigma_{3} \right) + \frac{\rho}{2} = -\left( Y_{n}(0)^{-1} Y_{n}'(0) \right)_{11} + \frac{\rho}{2},$$
(2.37)

where we have used that  $Y_n(0)Y'_n(0)$  is a traceless matrix (due to the fact that det  $Y_n \equiv 1$ ).

Computation of  $(Y_n(0)^{-1}Y'_n(0))_{11}$ . Using that det  $Y_n \equiv 1$  and the formula (2.13) we deduce

$$(Y_n(0)^{-1}Y'_n(0))_{11} = p'_n(0) \oint_{\gamma} \frac{\tilde{p}_{n-1}(w)e^{\theta(w)}dw}{w2i\pi} - \tilde{p}'_{n-1}(0) \oint_{\gamma} \frac{p_n(w)e^{\theta(w)}dw}{w2i\pi}.$$
(2.38)

The derivatives at z = 0 of  $p_n, \tilde{p}_{n-1}$  are obtained from their determinantal representations (2.14), (2.15):

$$p'_{n}(0) = (-1)^{n+1} \frac{\begin{vmatrix} \mu_{0} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n+1} & \dots & \mu_{2n-1} \end{vmatrix}}{D_{n}}$$

$$\tilde{p}'_{n-1}(0) = \frac{(-1)^{n+1} 2\pi i}{D_{n}} \begin{vmatrix} \mu_{0} & \mu_{2} & \dots & \mu_{n-1} \\ \mu_{1} & \mu_{3} & \dots & \mu_{n} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-2} & \mu_{n} & \dots & \mu_{2n-3} \end{vmatrix}$$

$$(2.39)$$

The contour integrals in (2.38) can be expressed as well in terms of determinants of the moments  $\mu_j$  provided we extend their definition to all  $j \in \mathbb{Z}$ :

$$\oint_{\gamma} \frac{\tilde{p}_{n-1}(w)e^{\theta(w)}dw}{w2i\pi} = -\frac{\begin{vmatrix} \mu_{0} & \mu_{1} & \mu_{2} & \dots & \mu_{n-1} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-2} & \mu_{n-1} & \mu_{n} & \dots & \mu_{2n-3} \\ \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-2} \end{vmatrix}}{D_{n}} = (-1)^{n} \frac{\begin{vmatrix} \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-2} & \mu_{n-1} & \mu_{n} & \dots & \mu_{2n-3} \end{vmatrix}}{D_{n}}$$

$$(2.41)$$

$$\oint_{\gamma} \frac{p_{n}(w)e^{\theta(w)}dw}{w2i\pi} = \frac{1}{2\pi i} \frac{\begin{vmatrix} \mu_{0} & \mu_{1} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n} & \mu_{n+1} & \dots & \mu_{2n-1} \\ \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \end{vmatrix}}{D_{n}} = \frac{(-1)^{n}}{2\pi i} \frac{\begin{vmatrix} \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \\ \mu_{0} & \mu_{1} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n} & \mu_{n+1} & \dots & \mu_{2n-1} \\ D_{n} & & & & & \\ \end{pmatrix}$$

$$(2.42)$$

Inserting (2.39), (2.40), (2.41), (2.42) into (2.38), simplifying and rearranging the rows of the resulting determinants we obtain

$$(Y_{n}(0)^{-1}Y_{n}'(0))_{11} = -\frac{\begin{vmatrix} \mu_{0} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n+1} & \dots & \mu_{2n-1} \end{vmatrix} \begin{vmatrix} \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-2} & \mu_{n-1} & \mu_{n} & \dots & \mu_{2n-3} \end{vmatrix}} + \\ + \frac{\begin{vmatrix} \mu_{0} & \mu_{2} & \dots & \mu_{n-1} \\ \mu_{1} & \mu_{3} & \dots & \mu_{n} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-2} & \mu_{n} & \dots & \mu_{2n-3} \end{vmatrix}}{D_{n}} \begin{vmatrix} \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \\ \mu_{0} & \mu_{1} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n} & \mu_{n+1} & \dots & \mu_{2n-1} \end{vmatrix}}{D_{n}}$$
(2.43)

We now use the Desnanot-Jacobi identity, which can be stated as follows; given a square matrix M denote by  $M^{[a_1,\ldots,a_r][b_1,\ldots,b_r]}$  the matrix obtained by deleting the rows  $a_1,\ldots,a_r$  and columns  $b_1,\ldots,b_r$ ; then

$$\det M^{[a,b][c,d]} \det M = \det M^{[a][c]} \det M^{[b][d]} - \det M^{[a][d]} \det M^{[b][c]}.$$
(2.44)

We apply the identity to the matrix

$$M := \begin{vmatrix} \mu_{-1} & \mu_{0} & \mu_{1} & \dots & \mu_{n-1} \\ \mu_{0} & \mu_{1} & \mu_{2} & \dots & \mu_{n} \\ \mu_{1} & \mu_{2} & \mu_{3} & \dots & \mu_{n+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_{n} & \mu_{n+1} & \dots & \mu_{2n-1} \end{vmatrix} \in \operatorname{gl}(n+1).$$

$$(2.45)$$

Then using (2.44) we have

$$(Y_n(0)^{-1}Y'_n(0))_{11} = \frac{-\det M^{[1][2]} \det M^{[n+1][n+1]} + \det M^{[1,n+1][2,n+1]} \det M}{D_n^2} \stackrel{(2.44)}{=} \\ = -\frac{\det M^{[1][n+1]} \det M^{[n+1][2]}}{D_n^2} = -\frac{\det M^{[n+1][2]}}{D_n} \tag{2.46}$$

where the last equality is on account of the fact that  $D_n = \det M^{[1][n+1]}$ . The final part of the verification is based on the observation that

$$\partial_t \mu_j(t) = \partial_t \oint_{\gamma} z^j \left( 1 - \frac{1}{z} \right)^K z^{K-1} \mathrm{e}^{\frac{t}{z}} \mathrm{d}z = \mu_{j-1}(t), \quad j \in \mathbb{Z}.$$
(2.47)

This allows us to show that

$$\partial_t \det D_n = \det M^{[n+1][2]} \tag{2.48}$$

and hence

$$(Y_n(0)^{-1}Y'_n(0))_{11} = -\partial_t \ln D_n(t).$$
(2.49)

This concludes the proof.

## 3 Asymptotic analysis

The goal of this second part of the paper is to study the behaviour of the poles of the rational solution or, which is the same, the zeroes of the Hankel determinant  $D_n(t)$  as  $n \to \infty$ . More precisely, and in line with similar investigations done for rational solution of PII [10, 2, 11] we are going to re-scale the zeros concurrently with n and study the behaviour of the zeros in the  $s = \frac{t}{n}$ -plane. Namely we study the sequence of functions (polynomials)

$$\tau_n(s) := D_n(ns; \boldsymbol{K}, \boldsymbol{\rho}). \tag{3.1}$$

where

$$D_n(t;K,\boldsymbol{\rho}) := \det\left[\oint_{\gamma} z^{a+b+\boldsymbol{K}} \left(1 - \frac{1}{z}\right)^{\boldsymbol{\rho}} e^{\frac{t}{z}} \frac{\mathrm{d}z}{2i\pi}\right]_{a,b=0}^{n-1}$$
(3.2)

We point out that we are not scaling the parameters  $\rho \in \mathbb{C} \setminus \mathbb{Z}$  nor  $K \in \mathbb{Z}$  as  $n \to \infty$ ; a different asymptotic analysis would be needed if we were to set  $\rho \propto n$  and/or  $K \propto n$ . We will briefly point out the main differences when appropriate.

The Theorem 2.1 can be re-formulated for convenience in the following form:

**Theorem 3.1** The function  $\tau_n(s) := D_n(ns; K, \rho)$  is zero if and only if the RHP 2.1 has no solution (with t = ns).

The asymptotic analysis of the RHP 2.1 with t = ns falls within the purview of the Deift–Zhou asymptotic method that was brought to fruition in [15] for orthogonal polynomials, see also [7].

It consists of several, by now more or less standardized, steps that we can summarize as follows before delving into the details:

- 1. construct a "g-function"; this is a function g(z; s) whose domain of analyticity  $\mathscr{D}_s$  depends (continuously) on the parameter s. It can be expressed as an Abelian integral on a hyperelliptic Riemann surface  $\mathcal{R}$  whose genus depends on s. The g-function is uniquely characterized by a free-boundary value problem and certain inequalities which will be described in Sec. 4.
- 2. Use the constructed g-function to normalize the RHP 2.1 and express a new matrix valued function W(z) which satisfies a new RHP.
- 3. The RHP for W then undergoes a sequence of *transformations* (or reformulations) into equivalent RHPs.
- 4. The final reformulation is then amenable to a nonlinear steepest descent analysis using a standard *small norm theorem* for Riemann–Hilbert problems.

The key concept is that the solvability of the initial RHP 2.1 is equivalent to the solvability of its final reformulation hinted at in the list above. Thus, if we can guarantee that the final reformulation is solvable for s in a suitable domain, we can conclude that there are no zeros of  $\tau_n(s)$  in that domain (at least for n sufficiently large).

Let us give a visual overview of the result; in Figure 1 we can see several instances of plots of the zeros of  $\tau_n(s)$  for various values of  $\rho$ ; the common feature, which will be proved, is that all the zeros lie, asymptotically for large n, within the "Eye of the Tiger" region (EoT), marked by the black arcs; the equation of these black arcs is given implicitly in (4.26).

Also indicated is a grid of lines within the EoT; their intersection is the *approximate* location of the zeros as it follows from the asymptotic analysis, and they represent the vanishing of a theta function for an elliptic curve, see Section 5.2.

Consequently we are going to split the analysis in two cases;

- the Outside of the EoT, Sections 4.1 and 5.1;
- the Inside of the EoT, Sections 4.2 and 5.2.

## 4 Construction of the *g*-function

The method of the steepest descent analysis requires the construction of a scalar function with certain properties that we list below. This function is universally known in the literature as the g-function. Here we recall that the symbol of the moment functional is

$$\theta(z;s) = \frac{ns}{z} + \boldsymbol{\rho} \ln\left(1 - \frac{1}{z}\right) + \boldsymbol{K} \ln z.$$
(4.1)

Since  $K, \rho$  will not be scaling in our setup, we introduce  $\theta_0(z; s) = \frac{s}{z}$  (the "scaling" part) so that

$$\theta(z;s) = n\theta_0(z;s) + \boldsymbol{\rho} \ln\left(1 - \frac{1}{z}\right) + \boldsymbol{K} \ln z.$$
(4.2)

If  $\mathbf{K} = n\mathbf{K}_0$  and/or  $\boldsymbol{\rho} = n\boldsymbol{\rho}_0$ , namely, if we were to scale  $\mathbf{K}, \boldsymbol{\rho}$ , we would accordingly re-define  $\theta_0$  to contain all terms proportional to n.

**Definition 4.1 (The** *g*-function and its properties) The *g*-function is a locally bounded analytic function on  $\mathbb{C} \setminus \Gamma$  where  $\Gamma$  is a union of oriented contours (to be determined) extending to infinity satisfying the properties listed hereafter.



Figure 2: From top to bottom, left to right:  $(n, \rho) = (16, \frac{3}{100} + \frac{13i}{100}), (16, \frac{101}{100} + \frac{13i}{100}), (17, 3 + 1i), (17, \frac{1}{2} + \frac{i}{2}), (26, \frac{101}{100}), (40, \frac{3}{100})$ . Observe that the accuracy of the localization of the zeros by the grid depends on the ratio  $|\rho|/n$ , the smaller the better the approximation. For example in the case in the top right  $(n, \rho) = (17, 3 + 1i)$  the ratio is significant (approximately  $|\rho|/n \simeq 0.19$ ) and an analysis where  $\rho$  is considered as scaling would undoubtedly be more appropriate. In all cases we have  $\mathbf{K} = 0$ .

- 1. The contour  $\Gamma$  can be written as  $\Gamma = \Gamma_m \cup \Gamma_c \cup \Gamma_\infty$  (with  $\Gamma_m$  denoting the "main  $\operatorname{arc}(s)$ ", and  $\Gamma_c$ the "complementary  $\operatorname{arc}(s)$ ") where each of the components have pairwise disjoint relative interiors and both  $\Gamma_m, \Gamma_c$  consist of a finite union of compact arcs:  $\Gamma_{\{m,c\}} = \bigsqcup \Gamma_{\{m,c\}}^{(j)}$ . Finally  $\Gamma_\infty$  is a simple contour extending to infinity from a finite point, traversing eventually the negative real axis and oriented from infinity.
- 2. the contour  $\gamma = \{|z| = R, R > 1\}$  can be homotopically retracted to  $\Gamma_m \cup \Gamma_c$  in  $\mathbb{C} \setminus [0, 1]$ , where [0, 1] here denotes a smooth simple arc connecting z = 0, 1 (not necessarily the straight segment).
- 3. for each  $z \in \Gamma_m \cup \Gamma_c$  we have

$$g(z_{+}) + g(z_{-}) = -\theta_0(z) - \ell + i\varpi_j, \qquad \varpi_j \in \mathbb{R}, \qquad z \in \Gamma_m^{(j)}$$

$$(4.3)$$

$$g(z_{+}) - g(z_{-}) = i\widehat{\varpi}_{j}, \qquad \widehat{\varpi}_{j} \in \mathbb{R}, \qquad z \in \Gamma_{c}^{(j)}$$

$$(4.4)$$

for some constants  $\ell, \varpi_j, \widehat{\varpi}_j$  (different on each of the connected components<sup>5</sup> of  $\Gamma_m, \Gamma_c$ ), while

$$g(z_{+}) - g(z_{-}) = 2i\pi, \quad z \in \Gamma_{\infty}.$$
(4.5)

4. as  $z \to \infty$  in  $\mathbb{C} \setminus \Gamma$  we have

$$g(z) = \ln(z) + \mathcal{O}(z^{-1}).$$
 (4.6)

- 5. the real part of the g-function is continuous on  $\mathbb{C}$  (including  $\Gamma$ ) and harmonic on  $\mathbb{C} \setminus \Gamma_m$  and satisfies the following inequalities:
  - (i) for all  $z \in \Gamma_c$  we have

Re 
$$(g(z_+) + g(z_-) + \theta_0(z) + \ell)$$
 = Re  $(2g(z) + \theta_0(z) + \ell) \le 0$  (4.7)

with the equality holding only at the endpoints of each component of  $\Gamma_c$  and possibly at isolated points within the relative interior of  $\Gamma_c$ ;

(ii) for  $z \in \Gamma_m$  we have

Re 
$$(g(z_+) + g(z_-) + \theta_0(z) + \ell) =$$
 Re  $(2g(z) + \theta_0(z) + \ell) \equiv 0.$  (4.8)

(iii) the inequality

$$\operatorname{Re}\left(2g(z) + \theta_0(z) + \ell\right) \ge 0 \tag{4.9}$$

holds in an open neighbourhood U of  $\dot{\Gamma}_m$  (the indicating the interior set in the relative topology of the collection of arcs  $\Gamma_m$ , i.e., the arcs minus the end-points) with the equality holding only on  $\Gamma_m$  itself.

It is convenient to reformulate Def. 4.1 in terms of the so-called effective potential

$$\varphi(z;s) = 2g(z;s) + \theta_0(z;s) + \ell = 2g(z;s) + \frac{s}{z} + \ell.$$
(4.10)

**Corollary 4.1** There exist constants  $\varpi_j$ ,  $\varsigma_j$  such that the effective potential  $\varphi$  satisfies:

- 1. Re  $(\varphi(z))$  is continuous on  $\mathbb{C} \setminus \{0\}$ , harmonic on  $\mathbb{C} \setminus \Gamma_m \cup \{0\}$ ;
- 2. Re  $(\varphi \theta_0)$  is harmonic near z = 0 and Re  $(\varphi 2\ln(z) \theta_0(z))$  is harmonic at infinity;
- 3. the following equalities and inequalities hold

$$\varphi(z_{+}) + \varphi(z_{-}) = 2i\varpi_{j}, \qquad z \in \Gamma_{m}^{(j)}$$
(4.11)

$$\varphi(z_{+}) - \varphi(z_{-}) = 2i\varsigma_{j}, \qquad z \in \Gamma_{c}^{(j)}$$
(4.12)

$$\varphi(z_+) - \varphi(z_-) = 4i\pi, \qquad z \in \Gamma_\infty$$
(4.13)

 $\operatorname{Re}\varphi(z) \equiv 0, \quad z \in \Gamma_m \tag{4.14}$ 

$$\operatorname{Re}\varphi(z) \le 0, \quad z \in \Gamma_c \tag{4.15}$$

$$\operatorname{Re}\varphi(z) \ge 0, \quad z \in U,\tag{4.16}$$

<sup>&</sup>lt;sup>5</sup>We will use different notation for these constants in the specific cases we discuss below.

with U as in Def. 4.1, 5(iii).

In the next two sections we are going to construct  $\varphi$  directly and verify those properties.

### 4.1 Outside the EoT: genus 0 *g*-function and effective potential

For |s| sufficiently large we are going to postulate first, and then verify, the form of the effective potential. We will see that the conditions in Corollary 4.1 are fulfilled for s ranging in an unbounded region and up to the boundary of a compact almond-shaped region that we are going to informally call the "eye of the tiger" (EoT). This is the region bounded by the black arcs in Fig 2.

We start by postulating the following expression;

$$\varphi(z;s) := 2 \int_{-\frac{is}{2}}^{z} \frac{\sqrt{w^2 + \frac{s^2}{4}}}{w^2} \mathrm{d}w \; \Rightarrow \; \varphi'(z;s) = 2 \frac{\sqrt{z^2 + \frac{s^2}{4}}}{z^2} \tag{4.17}$$

The function  $\varphi(z;s)$  can be written explicitly

$$\varphi(z;s) = 2\ln\left(\frac{2iz}{s} + i\sqrt{\frac{4z^2}{s^2} + 1}\right) - 2\sqrt{1 + \frac{s^2}{4z^2}}$$
(4.18)

We need to describe the domain of analyticity; we start from the domain of analyticity of  $\varphi'$ . A great simplification is achieved by observing that  $\varphi$  is really a function only of  $\frac{z}{s}$ , namely

$$\varphi(z;s) = \varphi\left(\frac{z}{s};1\right),\tag{4.19}$$

and hence it suffices to describe the domain and properties of  $\varphi_0(z) := \varphi\left(\frac{z}{s}; 1\right)$  which is given by

$$\varphi_0(z) = 2\ln\left(2iz + i\sqrt{4z^2 + 1}\right) - 2\sqrt{1 + \frac{1}{4z^2}}, \quad \varphi_0'(z) = \frac{\sqrt{4z^2 + 1}}{z^2}.$$
(4.20)

The determination of the root is chosen such that  $\varphi'_0(z) \simeq \frac{1}{z}$  at  $z = \infty$ , with a branch-cut connecting the branchpoints  $\pm \frac{i}{2}$  to be determined below. The language of vertical trajectories of quadratic differentials of [38] is useful in this discussion: by definition these are the arcs of curves where Re  $\varphi_0$  is constant, which, in the plane of the variable  $\xi(z) := \varphi_0(z) = \int^z \varphi'_0(w) dw$  are (by definition) vertical segments, whence the terminology. We start by observing that  $\operatorname{res}_{z=\infty} \varphi'_0(z) dz = -2i\pi$  and  $\operatorname{res}_{z=0} \varphi'_0(z) dz = \pm 2i\pi$  (with the sign depending on whether the branch-cut leaves z = 0 to the left or to the right) and hence, no matter how we choose the branch-cut  $\Gamma_m$  (connecting the branchpoints) we have that

the function  $\operatorname{Re} \varphi_0(z)$  is single-valued, harmonic in  $\mathbb{C} \setminus \Gamma \cup \{0\}$  and continuous in  $\mathbb{C} \setminus \{0\}$ ; (4.21)

for |z| sufficiently large  $\operatorname{Re} \varphi_0(z) = \ln |z| + \text{harmonic and bounded.}$  (4.22)

The observation (4.22) implies that the level-curves of  $\operatorname{Re} \varphi_0$  are deformed circles for |z| sufficiently large. One can verify that changing the determination of both radicals in (4.20) has the effect of flipping the sign of  $\operatorname{Re} \varphi_0$  and hence that  $\operatorname{Re} \varphi_0$  is a well-defined harmonic function on the Riemann surface of the radical

$$w^2 = 1 + 4z^2. \tag{4.23}$$

Furthermore  $\operatorname{Re} \varphi_0$  is an odd function under the holomorphic involution that maps (w, z) to (-w, z). This means that the level sets  $\operatorname{Re} \varphi_0 = 0$  are well defined on the z-plane; they consist of "vertical trajectories" (in the terminology of [38]) issuing from the points  $\pm \frac{i}{2}$  and forming the pattern illustrated in Fig. 3.

We choose the branch-cut of the radical as the arc of Fig. 3 joining  $\pm \frac{i}{2}$  in the right half-plane. With this choice we have that

$$\varphi_0'(z) \simeq -\frac{1}{z^2} + \mathcal{O}(1), \quad z \to 0.$$
 (4.24)

and in general  $\varphi'(z;s) = \frac{1}{s}\varphi'_0\left(\frac{z}{s}\right)$  satisfies

$$\varphi'(z;s) = -\frac{s}{z^2} + \mathcal{O}(1).$$
 (4.25)

Verification of the properties of  $\varphi$  and range of validity. It suffices to verify the properties for s = 1since changing  $s \in \mathbb{C}$  just amounts to a complex homotethy  $z \mapsto sz$ . We choose  $\Gamma_m$  as the arc joining  $\pm \frac{i}{2}$  in the right plane,04  $\Gamma_c$  as an arc joining the two points  $\pm \frac{i}{2}$  in the left plane, and inside the region bounded by the imaginary axis and the contour  $-\Gamma_m$  (see Fig. 3). Finally we choose  $\Gamma_\infty$  as the ray  $(-i\infty, -\frac{i}{2}]$ . We then proceed with the verification of the properties in Corollary 4.1:

- 1. on the sole connected component  $\Gamma_m$ , we have  $\varphi_0(z_+) + \varphi_0(z_-) = 0$  since the two boundary values differ by a vanishing period of  $\varphi'_0$ ;
- 2. on  $\Gamma_c$  we similarly have  $\varphi(z_+) = \varphi(z_-)$ ;
- 3. on  $\Gamma_{\infty}$  we have  $\varphi_0(z_+) = \varphi_0(z_-) \operatorname{res}_{w=\infty} \varphi'_0(w) dw = \varphi_0(z_-) + 4i\pi$
- 4. Since  $\Gamma_m$  is defined as the zero level set of  $\operatorname{Re} \varphi_0$ , we have  $\operatorname{Re} \varphi_0 \equiv 0$  on  $\Gamma_m$  by definition;
- 5. in the unbounded doubly-connected region outside of the "apricot" in Fig. 3 we have  $\operatorname{Re} \varphi_0 = \ln |z| + \mathcal{O}(1)$  near  $z \to \infty$ ; thus inevitably  $\operatorname{Re} \varphi_0 > 0$  in the whole region (which, we remind, is bounded by the zero levelsets of  $\operatorname{Re} \varphi_0$ );
- 6. In the right hemi-apricot, the sign must be also positive because  $\operatorname{Re} \varphi_0(z) = \operatorname{Re} \left(\frac{1}{z} + \mathcal{O}(1)\right);$
- 7. by the same token, the sign is negative in the left hemi-apricot.

Thus all conditions except possibly the condition no. 2 in Def. 4.1 are verified, namely, we still need to verify that the union  $\Gamma_m \cup \Gamma_c$  is homotopic to a circle |z| = R, R > 1 in the cut plane  $\mathbb{C} \setminus [0, 1]$ .

Since the levelsets in Fig. 3 are scaled by s, this latter condition is fulfilled as long as the point z = 1 lies inside the re-scaled apricot. This holds clearly for |s| sufficiently large, and it fails precisely when the point z = 1 lies on either  $\Gamma_m$  or  $\Gamma_c$ , namely when

$$\operatorname{Re}\varphi(1;s) = 0 = \operatorname{Re}\varphi_0\left(\frac{1}{s}\right). \tag{4.26}$$

Observing (4.26) we conclude that the shape of the locus (4.26) is simply the image of the apricot in Fig. 3 (left pane) under the inversion  $z \mapsto \frac{1}{z}$  (see right pane in Fig. 3). In Fig. 4 we illustrate the contour for a generic value of s outside of the EoT.

The discussion of this section can be summarized by the following theorem.

**Theorem 4.1** For s outside the region EoT the effective potential is given by (4.18) (and (4.17)). The contour  $\gamma$  can be chosen as the contour  $\Gamma_m \cup \Gamma_c$  consisting of the arc  $\Gamma_m$  from  $-\frac{is}{2}$  to  $\frac{is}{2}$  lying on the left of the straight segment, passing through the origin, from  $\frac{is}{2}$  to  $-\frac{is}{2}$  (in this orientation). The arc  $\Gamma_c$  is an arc in the right "lobe" keeping z = 0 to its left. See Fig. 4.

The Robin constant  $\ell$  appearing in (4.10) is given by

$$\ell = 2\ln\left(-\frac{4}{s}\right).\tag{4.27}$$

The only statement that has not been proven yet is the expression (4.27). The expression for g(z;s) is derived from that of the effective potential (4.18) and the relation (4.10) expressing the effective potential in terms of the *g*-function. Since  $g(z) \sim \ln(z) + \mathcal{O}(z^{-1})$  (note the absence of a constant term in the asymptotic expansion) we can deduce  $\ell$  by the expansion at  $z = \infty$  of  $g(z;s) - \ln(z)$ .

From (4.18), (4.10) we have

$$g(z;s) - \ln z = \ln\left(\frac{2}{s}\right) + \ln\left(-1 - \sqrt{1 + \frac{s^2}{4z^2}}\right) - \frac{\ell}{2} + \mathcal{O}(z^{-1}) = \ln\left(-\frac{4}{s}\right) - \frac{\ell}{2} + \mathcal{O}(z^{-1}).$$
(4.28)

Setting to zero the constant term in the expansion yields the statement (4.27).



Figure 3: Left: The levelsets of  $\operatorname{Re} \varphi_0(z) = 0$ . Right: the boundary of the region of validity of the genus–zero assumption (ignore the vertical rays issuing from  $\pm 2i$ ). It is the locus of  $\operatorname{Re} \varphi_0(1/s) = 0$ . The inside of this region we refer to as the "Eye of the Tiger" (EoT).



Figure 4: The contours  $\Gamma_m$ ,  $\Gamma_c$  and the regions where  $\operatorname{Re} \varphi(z; s) < 0$  (shaded). Indicated also the boundaries,  $\mathscr{L}_{\pm}$ , of the lens regions  $\Lambda_{\pm}$ 

### 4.2 Inside the EoT: genus one *g*-function and effective potential

The boundary of EoT is precisely the condition that the point z = 1 belongs to the two sub-arcs of the zero-level set of  $\operatorname{Re} \varphi(z; s)$  forming the "rind" of the apricot. As we move s inside the EoT we cannot use the same effective potential described in the previous section because the second condition in Def. 4.1 ceases to be verified, namely the contour of integration  $\gamma$  cannot be homotopically retracted to  $\Gamma_m \cup \Gamma_c$  within  $\mathbb{C} \setminus [0, 1]$  since either  $\Gamma_c$  or  $\Gamma_m$  intersect the segment [0, 1].

The idea to resolve the impasse is to treat z = 1 as a "hard–edge", using the terminology that has come to pass in the random matrix theory literature [5, 8]. We thus postulate the following form for  $\varphi'(z;s)$ 

$$\varphi'(z;s) = \frac{2}{z^2}\sqrt{z^2 + \frac{s^2}{4} + \frac{Az^2}{z-1}} = \frac{2\sqrt{z^2(z-1+A) + \frac{s^2}{4}(z-1)}}{z^2\sqrt{z-1}}$$
(4.29)

The parameter A = A(s) is chosen by the condition that all periods of  $\varphi'(z; s)dz$  on the Riemann surface of the radical are purely imaginary (this is called *Boutroux condition*), which is the necessary condition so that Re  $\varphi$  is continuous across the cuts; the Riemann surface of  $\varphi'(z; s)$  is an elliptic curve branched at z = 1 and the other three roots of the radical in the numerator:

$$\mu^{2} = (z-1)\left(z^{2}(z-1+A) + \frac{s^{2}}{4}(z-1)\right).$$
(4.30)

We denote these roots as  $b, a_+, a_-$  with b the closest root to z = 1; an expression in terms of Cardano's formulæ is possible but not necessary.

Now the complex parameter A(s) is determined implicitly by the two real equations

$$\operatorname{Re} \int_{b}^{1} \varphi'(z;s) dz = 0 \qquad \operatorname{Re} \int_{b}^{a_{+}} \varphi'(z;s) dz = 0.$$
(4.31)

Under these conditions it then follows that the real part of

$$\varphi(z;s) = \int_{a_{-}}^{z} \varphi'(w;s) \mathrm{d}w \tag{4.32}$$

is a well defined (single valued) harmonic function on the Riemann surface minus the preimages of the points z = 0 on the two sheets.

**Determination of**  $\Gamma_m$  and  $\Gamma_c$ . By the same argument already used in the genus zero case, the zero level sets Re  $\varphi(z; s) = 0$  are well defined; they consists of the critical vertical trajectories of the quadratic differential  $Q = \varphi'(z; s)^2 dz^2$  [38]

$$Q = 4 \frac{z^2(z-1+A) + \frac{s^2}{4}(z-1)}{z^4(z-1)}.$$
(4.33)

The following discussion is best followed by referring to Fig. 5, in particular the smaller inset vignettes. The main arcs  $\Gamma_m$  are sub-arcs of the zero levelset of Re  $\varphi$  and we need to discuss their qualitative topology before proceeding.

The critical points are the three (generically) simple zeros and the simple pole z = 1; there are three vertical trajectories that issue from each simple zero, while from the simple pole there is only one. The union of the trajectories is a connected planar graph and the unbounded region is put in conformal equivalence with the puncture unit disk by the map

$$\zeta = e^{-\frac{\varphi(z;s)}{4i\pi}} \tag{4.34}$$

which maps the exterior region into the disk  $|\zeta| < 1$ , with  $z = \infty$  mapped to  $\zeta = 0$ . Some observations are in order:

1. the level sets of  $\operatorname{Re} \varphi$  depend only on  $s^2$  and they are conjugated if we conjugate s;

2. one of the zeros of  $\varphi'$  is connected by a vertical trajectory to z = 1; we denote this zero by z = b; the other two zeros are one in the upper and one in the lower half plane. We denote them by  $a_{\pm}$ , respectively.

While the level sets of  $\operatorname{Re} \varphi$  depend on  $s^2$  alone, we must choose the branch-cuts  $\Gamma_m$  differently according to the cases  $\operatorname{Re} s > 0$  or  $\operatorname{Re} s < 0$ ; the reason is that the sign distribution of  $\operatorname{Re} \varphi$  differs in the two cases. This is seen by the following reasoning:

- In the outside region  $\operatorname{Re} \varphi \simeq 2 \ln |z| + \mathcal{O}(1)$  and hence  $\operatorname{Re} \varphi > 0$ ;
- near the origin we must have  $\varphi(z;s) = -\frac{s}{z} + \mathcal{O}(1)$  and hence for  $\operatorname{Re} s > 0$  the right "lobe" is where  $\operatorname{Re} \varphi < 0$ ; viceversa it is the left one if  $\operatorname{Re} s < 0$ .

The branch-cut  $\Gamma_m$  is then singled out by the fact that across it the function  $\operatorname{Re} \varphi$  is continuous but not differentiable, namely,  $\operatorname{Re} \varphi$  has the same sign (positive) on both sides.

Collecting these observations, we thus have determined that

- 1. For Re s > 0 the branch-cut  $\Gamma_m$  consists of the three arcs of the vertical trajectories connecting z = b, 1and  $z = b, a_+$  and  $z = b, a_-$ .
- 2. For Re s < 0 the branch-cut  $\Gamma_m$  consists of the two arcs of the vertical trajectories connecting z = b, 1and  $z = a_+, a_-$  (passing to the left of the origin).

See Fig. 5.

The specification of the constants that appear for the boundary values of  $\varphi$  in Corollary 4.1 will be given in Sections 5.2.1 and 5.2.2 for Re (s) > 0 and Re (s) < 0, respectively.

### 5 Deift–Zhou steepest descent analysis

We split the asymptotic analysis in three subsections, according to the following cases:

- 1.  $s \in \mathcal{K}_{out}$ , where  $\mathcal{K}_{out}$  is a closed subset contained outside of the EoT;
- 2.  $s \in \mathcal{K}_{in,+}$ , where  $\mathcal{K}_{in,+}$  is a compact subset of  $\{\operatorname{Re}(s) > 0\} \cap EoT$ ;
- 3.  $s \in \mathcal{K}_{in,-}$ , where  $\mathcal{K}_{in,-}$  is a compact subset of  $\{\operatorname{Re}(s) < 0\} \cap EoT$ .

Note that for  $s \in i\mathbb{R} \cap EoT$  the elliptic curve is degenerate and a different analysis would be needed, analogous to what would be needed on the boundary of the EoT. Near the corners  $s = \pm \frac{i}{2}$  a yet different analysis would be needed, which involves the construction of special local parametrices based on the first Painlevé transcendent. We do not discuss these transitional regions.

The main take-away of the analysis is the following theorem.

**Theorem 5.1** For any closed subset  $\mathcal{K}$  on the outside of the EoT, the RHP 2.1 (with t = ns) is solvable for n sufficiently large, and hence the poles of the rational solutions must be inside a neighbourhood of the EoT.

The Deift-Zhou steepest descent analysis, formalized in [15] and in countless literature thereafter requires a number of transformations of the problem into equivalent ones. We describe briefly below these problems. We recall that our starting point is the following

**Problem 5.1** Find a  $2 \times 2$  matrix-valued function  $Y(z) = Y_n(z; s)$  analytic in  $\mathbb{C} \setminus \gamma$ , with analytic bounded inverse and such that

$$Y(z_{+}) = Y(z_{-}) \begin{bmatrix} 1 & z^{\mathbf{K}} \left(1 - \frac{1}{z}\right)^{\boldsymbol{\rho}} e^{\frac{ns}{z}} \\ 0 & 1 \end{bmatrix} \qquad z \in \gamma$$

$$(5.1)$$

$$Y(z) = \left(\mathbf{1} + \mathcal{O}(z^{-1})\right) z^{n\sigma_3}, \quad z \to \infty$$
(5.2)



Figure 5: Illustration of the zero level-sets of  $\operatorname{Re} \varphi(z; s)$  for various values of s; the inset vignettes have their centre at the value of s to which they correspond.

We remind the reader that  $K \in \mathbb{Z}$  while we assume  $\rho \notin \mathbb{Z}$ . We will use the notation

$$Q(z) = z^{K} \left(1 - \frac{1}{z}\right)^{\rho}, \qquad (5.3)$$

where the domain is  $\mathbb{C} \setminus \Gamma_0^1$ , where  $\Gamma_0^1$  denotes an arc homotopic to [0, 1] at fixed endpoints. For s outside the EoT we shall choose  $\Gamma_0^1 = [0, 1]$  (the segment).

It will be necessary for the analysis inside the EoT to partially homotopically retract the integration contour  $\gamma$  along a subarc of  $\Gamma_0^1$  (from the left and right of it): in that case the RHP (5.1) will take a slightly different form due to the fact that the function Q (5.3) has a jump-discontinuity such that  $Q(z_+) = Q(z_-)e^{2i\pi\rho}$ . In this case the jump matrix of the RHP (5.1) along such a subarc of  $\Gamma_0^1$  needs to be replaced by

$$\begin{bmatrix} 1 & Q(z_{+}) \left( e^{-2i\pi\rho} - 1 \right) e^{\frac{ns}{z}} \\ 0 & 1 \end{bmatrix}.$$
 (5.4)

### 5.1 Asymptotic analysis for *s* outside of the EoT

We recall that  $\Gamma_m$ ,  $\Gamma_c$  have been defined in Section 4.1

Normalization:  $Y \to W$ . We define

$$W(z) = e^{n\frac{\ell}{2}\sigma_3}Y(z)e^{-n\left(g(z)+\frac{\ell}{2}\right)\sigma_3}$$
(5.5)

A direct verification shows that W solves the following

**Problem 5.2** (*RHP<sub>W</sub>*) The matrix W(z) is analytic in  $\mathbb{C} \setminus \gamma = \mathbb{C} \setminus (\Gamma_m \cup \Gamma_c)$  and satisfies

$$W(z_{+}) = W(z_{-})J_{W}(z), \quad z \in \gamma$$

$$(5.6)$$

$$W(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{z}\right), \quad z \to \infty$$
 (5.7)

where

$$J_W(z) = \begin{bmatrix} e^{-n(g(z_+) - g(z_-))} & Q(z)e^{n\left(\frac{s}{z} + g(z_+) + g(z_-) + \ell\right)} \\ 0 & e^{n(g(z_+) - g(z_-))} \end{bmatrix} = \begin{bmatrix} e^{-\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} & Q(z)e^{\frac{n}{2}(\varphi(z_+) + \varphi(z_-))} \\ 0 & e^{\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} \end{bmatrix}$$
(5.8)

**Lens Opening:**  $W \to T$ . Refer to Fig. 4. The lens regions  $\Lambda_{\pm}$  are the two regions bounded by  $\Gamma_m$  and the arcs  $\mathscr{L}_{\pm}$  chosen arbitrarily in the regions where  $\operatorname{Re} \varphi > 0$ . The process of "opening the lenses" consist in re-defining the matrix W within those regions. We thus define

$$T_{0}(z) := \begin{cases} W(z) & z \notin \Lambda_{\pm} \\ W(z) \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{-} \\ W(z) \begin{bmatrix} 1 & 0 \\ -\frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{+} \end{cases}$$
(5.9)

A direct computation shows that the matrix  $T_0(z)$  satisfies the following RHP

**Problem 5.3** (*RHP*<sub> $T_0$ </sub>) The matrix  $T_0(z)$  satisfies the conditions

$$T_0(z_+) = T_0(z_-)J_{T_0}(z), \qquad z \in \gamma$$
(5.10)

$$T_0(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{z}\right), \quad z \to \infty$$
 (5.11)

where

$$J_{T_0}(z) = \begin{cases} J_W(z) & z \in \Gamma_c \\ \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \mathscr{L}_{\pm} \\ \begin{bmatrix} 0 & Q(z) \\ -\frac{1}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_m \end{cases}$$
(5.12)

At this point we have obtained a RHP where the jump matrices on  $\Gamma_c \cup \mathscr{L}_{\pm}$  converge pointwise in the relative interior to the identity matrix, but not uniformly.

To further normalize the problem we need to construct the *outer parametrix*, namely the (explicit) solution of an auxiliary RHP where we simply drop the jump conditions on  $\Gamma_c \cup \mathscr{L}_{\pm}$ .

Outer parametrix. We seek the solution of the following "model problem"

**Problem 5.4** (*RHP*<sub> $M_{O}$ </sub>) The matrix  $M_{Q}(z)$  is analytic and analytically invertible in  $\mathbb{C} \setminus \Gamma_{m}$  and satisfies

$$M_{Q}(z_{+}) = M_{Q}(z_{-}) \begin{bmatrix} 0 & Q(z) \\ -\frac{1}{Q(z)} & 0 \end{bmatrix} \qquad z \in \Gamma_{m}$$
(5.13)

$$M_Q(z) = \mathbf{1} + \mathcal{O}(z^{-1}) \qquad |z| \to \infty \tag{5.14}$$

$$M_Q(z) = \mathcal{O}\left(\frac{1}{\left(z \mp \frac{is}{2}\right)^{\frac{1}{4}}}\right) \qquad z \to \pm \frac{is}{2}.$$
(5.15)

We now solve the RHP 5.4. To this end we need to construct a special scalar function S(z), often called the *Szegö* function, satisfying the following scalar boundary value problem

$$S(z_{+}) + S(z_{-}) = \ln Q(z), \quad z \in \Gamma_m$$
 (5.16)

$$\sup_{z \in \mathbb{C} \setminus \Gamma_m} |S(z)| < +\infty.$$
(5.17)

The solution is given by the following expression

$$S(z) = R(z) \int_{\Gamma_m} \frac{\ln Q(w) dw}{R(w_+)(w-z)2i\pi}, \qquad R(z) := \sqrt{z^2 + \frac{s^2}{4}}$$
(5.18)

where the branch-cut of the radical R(z) is chosen to run along  $\Gamma_m$ . We leave to the reader the verification that the proposed expression (5.18) fulfills all the required conditions.

We can actually simplify the Szegö function by a contour deformation recalling that  $\ln Q(w_+) = \ln Q(w_-) + 2i\pi\rho$  for z on the segment [0,1]; indeed by a contour deformation we can rewrite S(z) as follows

$$S(z) = R(z) \int_{\Gamma_m} \frac{\ln Q(w) dw}{R(w_+)(w-z)2i\pi} =$$
(5.19)

$$= -\frac{R(z)}{2} \oint_{\odot\Gamma_m} \frac{\ln Q(w) \mathrm{d}w}{R(w)(w-z)2i\pi},\tag{5.20}$$

where the symbol  $\odot \Gamma_m$  stands for a counterclockwise loop leaving  $\Gamma_m$  in its interior region and the segment [0, 1] on the exterior. Using then the Cauchy's residue theorem we get

$$S(z) = \frac{1}{2} \ln Q(z) - \frac{R(z)}{2} \int_{0}^{1} \frac{(\ln Q(w_{+}) - \ln Q(w_{-})) dw}{(w - z)R(w)2i\pi} =$$

$$= \frac{\ln Q(z)}{2} - \frac{R(z)}{2} \int_{0}^{1} \frac{\rho dw}{R(w)(w - z)} - \frac{R(z)}{2} \int_{-\infty}^{0} \frac{\mathbf{K} dw}{R(w)(w - z)} =$$

$$= \frac{\mathbf{K}}{2} \ln \left( \frac{(R(z) - z) (R(0) + R(z))}{s/2} \right) + \frac{\rho}{2} \ln \left( \frac{z + \frac{s^{2}}{4} + R(z)R(1)}{\frac{s}{2} (R(z) + R(0))} \right)$$
(5.21)
(5.22)

The simplest way to verify this latest formula is to verify  $S(z_+) + S(z_-) = \ln Q(z)$  using that  $R(z_+) = -R(z_-)$ ; also one needs to verify that the expression has no singularities except the discontinuity across the branch-cut of R(z), and that it is bounded at infinity. We leave the instructive but tedious verification to the reader.

**Remark 5.1** The functon S(z) has a finite value at  $z = \infty$  given by

$$S(\infty) = \frac{\mathbf{K}}{2} \ln\left(\frac{s}{4}\right) + \frac{\mathbf{\rho}}{2} \ln\left(\frac{2}{s} + \sqrt{1 + \frac{4}{s^2}}\right)$$
(5.23)

With these preparations we state

Proposition 5.1 (Solution of the RHP 5.4) The solution of the RHP 5.4 is given by

$$M_Q(z) := e^{-S(\infty)\sigma_3} M(z) e^{S(z)\sigma_3}$$
(5.24)

$$M(s) := \left(\frac{z - \frac{is}{2}}{z + \frac{is}{2}}\right)^{\frac{-2}{4}} = \frac{1}{2} \begin{bmatrix} F + \frac{1}{F} & -i\left(F - \frac{1}{F}\right)\\ i\left(F - \frac{1}{F}\right) & F + \frac{1}{F} \end{bmatrix}$$
(5.25)

$$F := \left(\frac{z - \frac{is}{2}}{z + \frac{is}{2}}\right)^{\frac{1}{4}} \tag{5.26}$$

where the branchcut of F runs along  $\Gamma_m$  and the determination is chosen so that  $F(z) \to 1$  as  $|z| \to \infty$ . The proof is left to the reader.

#### 5.1.1 Conclusion of the steepest descent analysis

The final steps of the analysis requires the construction of *local parametrices* near the endpoints  $\pm \frac{is}{2}$ . Namely, one fixes two disks  $\mathbb{D}_{\pm}$  centered at each of the two points: this part is a quite standard construction and therefore we only sketch the main points. Let  $\mathbb{D}_{\pm}$  be two disjoint disks centered at  $\pm \frac{is}{2}$  (respectively) and of radii, say,  $r = \frac{|s|}{4}$  (the radius is not important for the discussion as long as they are small enough so as not to contain the origin):

$$\mathbb{D}_{\pm} := \left\{ \left| z \mp \frac{is}{2} \right| < \frac{|s|}{4} \right\}.$$
(5.27)

In a neighbourhood of  $a_{\pm} = \frac{\pm is}{2}$  the effective potential  $\varphi$  (4.18) has the following behaviour

$$\varphi(z;s) = \frac{2^{\frac{5}{2}}}{3} \left(\frac{z}{a_{\pm}} - 1\right)^{\frac{3}{2}} (1 + \mathcal{O}(z - a_{\pm})) \mod 2i\pi, \qquad z \sim a_{\pm}$$
(5.28)

In fact the additive constant is 0 in the case of  $a_{-}$  and  $-2i\pi$  for  $a_{+}$ , but since  $\varphi$  appears always in the exponent, this is irrelevant.

We define the **local coordinates**  $\zeta_{\pm}$  by the formulas

$$-\frac{4}{3}\zeta_{-}^{\frac{3}{2}} = n\varphi(z) = n\frac{2^{\frac{5}{2}}}{3}\left(\frac{z}{a_{-}} - 1\right)^{\frac{3}{2}}\left(1 + \mathcal{O}(z - a_{-})\right)$$
(5.29)

$$-\frac{4}{3}\zeta_{+}^{\frac{3}{2}} = n\left(\varphi(z) + 2i\pi\right) = n\frac{2^{\frac{5}{2}}}{3}\left(\frac{z}{a_{+}} - 1\right)^{\frac{3}{2}}\left(1 + \mathcal{O}(z - a_{+})\right).$$
(5.30)

From the above formulas it appears that both  $\zeta_{\pm}$  define a conformal map from the two disks  $\mathbb{D}_{\pm}$  (respectively) to a neighbourhood  $\mathscr{D}_n$  of the origin which is homothetically expanding with  $n^{\frac{3}{2}}$ . The determination of the fractional root can be chosen so that  $\zeta_{\pm}$  map the main arc  $\Gamma_m$  (where  $\operatorname{Re} \varphi = 0$  and  $\operatorname{Im} \varphi(z_+)$ ) is decreasing from 0 at  $a_-$  to  $-2i\pi$  at  $a_+$ ) to the negative real axis in the respective  $\zeta_{\pm}$  planes, with the points  $a_{\pm}$  being mapped to  $\zeta_{\pm} = 0$ . The arc  $\Gamma_c$  can be chosen so that it is mapped to the positive  $\zeta_{\pm}$ -axis while the arcs of  $\mathscr{L}_{\pm} \cap \mathbb{D}_{\pm}$  are mapped to two straight segments with slopes  $\pm 3\pi/2$ .

Then, the jump matrices  $J_{T_0}$  in the RHP 5.3 restricted to  $\mathbb{D}_+$  can be rewritten as (we focus on the case of  $\mathbb{D}_+$  for definiteness)

$$J_{T_0}(z) = \begin{cases} \begin{bmatrix} 1 & Q(z)e^{-\frac{4}{3}\zeta_+^{\frac{3}{2}}} \\ 0 & 1 \end{bmatrix} & z \in \Gamma_c \cap \mathbb{D}_+ \\ \begin{bmatrix} 1 & 0 \\ \frac{e^{\frac{4}{3}\zeta_+^{\frac{3}{2}}}}{Q(z)} & 1 \end{bmatrix} & z \in \mathscr{L}_{\pm} \cap \mathbb{D}_+ \\ \begin{bmatrix} 0 & Q(z) \\ -\frac{1}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_m \cap \mathbb{D}_+ \end{cases}$$
(5.31)

Observe that  $Q(z), \frac{1}{Q}(z)$  are both locally analytic at  $z = a_{\pm}$ , see (5.3). Then the following matrix

$$\mathcal{P}_{+}(z) := \mathbf{A}(\zeta_{+}(z)) Q(z)^{-\frac{\sigma_{3}}{2}} e^{\frac{2}{3}\zeta_{+}(z)^{\frac{3}{2}}\sigma_{3}}$$
(5.32)

exhibits discontinuities across  $(\Gamma_c \cup \mathscr{L}_+ \cup \mathscr{L}_{-m}) \cap \mathbb{D}_+$  with jump matrices given exactly by (5.31), as a consequence of the RHP satisfied by the matrix  $\mathbf{A}(\zeta)$  shown on Fig. 10. Completely analogous expressions hold for  $\mathcal{P}_-(z)$ .

The final approximation to the matrix  $T_0(z)$  (5.9) is then given by (see the definition of  $M_Q$  in (5.24))

$$\mathbf{G}(z) := \begin{cases} M_Q(z) & z \notin \mathbb{D}_+ \cup \mathbb{D}_-\\ M_Q(z)Q(z)^{\frac{\sigma_3}{2}} \begin{bmatrix} 1 & -1\\ i & i \end{bmatrix} \frac{\zeta_{\pm}(z)^{-\sigma_3/4}}{4\sqrt{\pi}} \mathcal{P}_{\pm}(z) & z \in \mathbb{D}_{\pm} \end{cases}$$
(5.33)



Figure 6: The support of the RHP for the error matrix  $\mathscr{E}(z)$ .

The matrix  $\mathbf{G}(z)$  has the same jumps as  $T_0(z)$  within  $\mathbb{D}_{\pm}$  and on  $\Gamma_m$ ; it has additional jump discontinuities across the boundaries  $\partial \mathbb{D}_{\pm}$  of the form

$$\mathbf{G}(z_{\pm}) = \mathbf{G}(z_{\pm}) \left( \mathbf{1} + \mathcal{O}(n^{-1}) \right), \quad z \in \partial \mathbb{D}_{\pm}.$$
(5.34)

Consequently the matrix

$$\mathscr{E}(z) := T_0(z)\mathbf{G}(z)^{-1} \tag{5.35}$$

has only jump discontinuities on  $\partial \mathbb{D}_+ \cup \partial \mathbb{D}_- \cup \mathscr{L}_+ \cup \mathscr{L}_-$  with all jump matrices that can be verified to be of the form  $\mathbf{1} + \mathcal{O}(n^{-1})$ , see Fig. 6.

It is then a standard result referred to in the literature as "small norm theorem" [14, 15] that the solution of this last RHP  $\mathscr{E}(z)$  exists and is uniformly close to the identity matrix. Reversing the order of the various transformations, this proves that the solution  $Y_n(z;s)$  of the RHP 5.1, exists (at least for n large enough).

### 5.2 Asymptotic analysis inside the EoT

With reference to Fig. 7 we have already observed that the levelsets of  $\operatorname{Re} \varphi(z; s)$  depend only on  $s^2$  and hence are invariant under the map  $s \mapsto -s$ ; the same is not true for the branch-cuts  $\Gamma_m$  and the regions of positivity of  $\operatorname{Re} \varphi$ . The decision of where to place the branch-cuts is forced by the "sign distribution" (i.e. in which regions we have  $\operatorname{Re} \varphi > 0$ ): this is mandated by the behaviour near infinity  $\operatorname{Re} \varphi = 2 \ln |z| + \mathcal{O}(1)$ for the outer region, and by the behaviour near the origin  $\operatorname{Re} \varphi = \operatorname{Re} \frac{s}{z} + \mathcal{O}(1)$  for the two bounded regions. Before we fork the analysis according to the sign of  $\operatorname{Re}(s)$  we make some common preparation.

As shown in Fig. 7 one of the arcs of  $\Gamma_m$  consists of an arc joining z = 1, b, denoted  $\Gamma_b^1$  (the orientation of the arc is suggested by the index placements). Thus, we need to preemptively modify the initial RHP 2.1 so that the contour  $\gamma$  is "retracted" in part along  $\Gamma_b^1$ . We leave the details to the reader and we report the resulting RHP, which is now the starting point of the following analysis.

**Problem 5.5** Let  $\gamma = \Gamma_b^1 \cup \gamma_b$  consist of the arc  $\Gamma_b^1$  and a loop,  $\gamma_b$ , starting and ending at z = b and containing the origin. Let Q(z) in (5.3) be defined with the branchcut extending from z = 0 to z = b and then along  $\Gamma_b^1$ , as shown in Fig. 7 by the dashed line. Given  $n \in \mathbb{N}$ , find a  $2 \times 2$  matrix-valued function  $Y(z) = Y_n(z)$  such that

1. Y(z) and  $Y^{-1}(z)$  are holomorphic and bounded in  $\mathbb{C} \setminus \gamma$ ;



Figure 7: The contours  $\Gamma_m$  (red) and  $\Gamma_c$  (green) for two antipodal values  $s_1 = -s_2$  of s with Re  $(s_1) > 0$ (left) and Re  $(s_2) < 0$  (right). Specifically here  $s_1 = 0.48 + 0.48i = -s_2$ . The dashed line indicates the branch-cut of the function Q(z), which runs partially along  $\Gamma_b^1$ . The components of  $\Gamma_c$  (green arcs) can be chosen freely within the shaded region, which indicates the region where Re  $\varphi < 0$ .

2. The boundary values along  $z \in \gamma$  satisfy

$$Y(z_{+}) = Y(z_{-}) \begin{bmatrix} 1 & Q(z) e^{\frac{\pi s}{z}} \\ 0 & 1 \end{bmatrix}, \qquad \forall z \in \gamma_{b}$$
(5.36)

$$Y(z_{+}) = Y(z_{-}) \begin{bmatrix} 1 & \varkappa Q(z_{+}) e^{\frac{ns}{z}} \\ 0 & 1 \end{bmatrix} \qquad \qquad \forall z \in \Gamma_{b}^{1}, \tag{5.37}$$

where

$$\varkappa := \left( e^{-2i\pi\rho} - 1 \right). \tag{5.38}$$

3. As  $z \to \infty$  the matrix  $Y_n(z)$  admits an asymptotic expansion of the form (2.12)

The first transformation is the same as in (5.5) and leads to the RHP for W as follows

**Problem 5.6** (*RHP<sub>W</sub>*) The matrix W(z) is analytic in  $\mathbb{C} \setminus \gamma = \mathbb{C} \setminus (\Gamma_m \cup \Gamma_c)$  and satisfies

$$W(z_{+}) = W(z_{-})J_{W}(z), \qquad z \in \gamma$$
(5.39)

$$W(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{z}\right), \quad z \to \infty$$
 (5.40)

where

$$J_W(z) = \begin{bmatrix} e^{-\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} & Q(z)e^{\frac{n}{2}(\varphi(z_+) + \varphi(z_-))} \\ 0 & e^{\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} \end{bmatrix}, \qquad z \in \Gamma \setminus \Gamma_b^1$$
(5.41)

$$J_W(z) = \begin{bmatrix} e^{-\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} & \varkappa Q(z_+) e^{\frac{n}{2}(\varphi(z_+) + \varphi(z_-))} \\ 0 & e^{\frac{n}{2}(\varphi(z_+) - \varphi(z_-))} \end{bmatrix}, \qquad z \in \Gamma_b^1$$
(5.42)

The second transformation is also similar to (5.9). However, from this point onwards, the details of the transformation depend on the case  $\operatorname{Re}(s) > 0$  or  $\operatorname{Re}(s) < 0$  and thus are given in Sections 5.2.1 and 5.2.2, respectively.

We will not carry out completely the error analysis, which would require the construction of appropriate local parametrices near the points  $z = b, a_{\pm}, 1$ . These are more or less known in the literature; we mention, for the sake of the reader with experience in the Deift-Zhou method, that

- 1. Near the points  $z = a_{\pm}, b$  the local parametrices are constructed from Airy functions [15, 2];
- 2. Near the point z = 1 the local parametrix is contructed in terms of Bessel functions, [41].

We instead focus on the construction of the global (outer) parametrix which solves a model problem; the pragmatic reason is that this computation will produce a formula for the approximate location of the zeros of  $\tau_n(s)$  which can be actually tested numerically, and whose result is evident in Fig. 2. Some details on the construction of the local parametrices are contained in Section 5.2.3 and App. A.1.

### 5.2.1 Second transformation, model problem and its solution: the case $\operatorname{Re}(s) > 0$

With reference to Fig. 7, we denote by  $\Lambda_{\pm}^{(x,y)}$  the region bounded by  $\Gamma_x^y$  and  $\mathscr{L}_{\pm}^{(x,y)}$ , for  $x, y \in \{a_+, a_-, b, 1\}$  and we call them *lens regions* (as opposed to their boundaries, which we refer to as the lens arcs).

**Definition of**  $T_0$  for  $\operatorname{Re}(s) > 0$ . We refer to the left pane of Fig. 7 and then set

$$T_{0}(z) := \begin{cases} W(z) \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{-}^{(a_{\pm},b)} \\ W(z) \begin{bmatrix} 1 & 0 \\ -\frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{+}^{(a_{\pm},b)} \end{cases}$$
(5.43)

however for the lens  $\Lambda^{(b,1)}_{\pm}$  adjacent to  $\Gamma^1_b$  we have instead

$$T_{0}(z) := \begin{cases} W(z) \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi(z)-2i\pi\rho}}{\varkappa Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{-}^{(b,1)} \\ W(z) \begin{bmatrix} 1 & 0 \\ -\frac{e^{-n\varphi(z)}}{\varkappa Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{+}^{(b,1)} \end{cases}$$
(5.44)

The main arcs are as in Fig. 7, left pane, thus

$$\Gamma_m = \Gamma_b^1 \cup \Gamma_{a_-}^b \cup \Gamma_b^{a_+}, \qquad \Gamma_c = \Gamma_{a_+}^{a_-}.$$
(5.45)

Recall from Sec. 4.2 that we have defined

$$\varphi(z;s) := \int_{a_{-}}^{z} \frac{2}{w^{2}} \sqrt{w^{2} + \frac{s^{2}}{4} + \frac{Aw^{2}}{w - 1}} \mathrm{d}w, \qquad (5.46)$$

with the constant A determined by the condition of all periods being purely imaginary and the integration path chosen in  $\mathbb{C} \setminus \Gamma_m$ . The effective potential satisfies the following boundary value relations

$$\varphi(z_{+}) + \varphi(z_{-}) = 0 \qquad \qquad z \in \Gamma_{a_{-}}^{b} \tag{5.47}$$

$$\varphi(z_{+}) + \varphi(z_{-}) = 2\Omega_1 \qquad \qquad z \in \Gamma_b^1 \tag{5.48}$$

$$\varphi(z_{+}) + \varphi(z_{-}) = 2\Omega_2 \qquad \qquad z \in \Gamma_{\iota}^{a_{+}} \tag{5.49}$$

- $\varphi(z_{+}) + \varphi(z_{-}) = 2\Omega_{2}$  $\varphi(z_{+}) \varphi(z_{-}) = 4i\pi$  $z \in \Gamma_c$ (5.50)

where the constants  $\Omega_1, \Omega_2$  are given by

$$\Omega_1 := \int_b^{a_+} \varphi'(z_+) \mathrm{d}z \in i\mathbb{R}_- \qquad \Omega_2 := \int_b^1 \varphi'(z_-) \mathrm{d}z \in i\mathbb{R}_+.$$
(5.51)

From the Cauchy residue theorem one also has

$$2\int_{a_{-}}^{b}\varphi'(z_{-})\mathrm{d}z + 2\int_{b}^{1}\varphi'(z_{-})\mathrm{d}z + 2\int_{b}^{a_{+}}\varphi'(z_{-})\mathrm{d}z = 4i\pi,$$
(5.52)

with all the integrals in  $i\mathbb{R}_+$ .

A direct computation, using also the properties (5.47)–(5.50) shows that the matrix  $T_0(z)$  satisfies the following RHP

**Problem 5.7** (*RHP*<sub> $T_0$ </sub>) The matrix  $T_0(z)$  satisfies the conditions

$$T_0(z_+) = T_0(z_-)J_{T_0}(z), \quad z \in \gamma$$
(5.53)

$$T_0(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{z}\right), \quad z \to \infty$$
 (5.54)

where

$$J_{T_{0}}(z) = \begin{cases} J_{W}(z) & z \in \Gamma_{c} = \Gamma_{a_{+}}^{a_{-}} \\ \begin{bmatrix} \frac{1}{e^{-n\varphi(z)}} & 0 \\ \frac{1}{Q(z)} & 1 \end{bmatrix} & z \in \mathscr{L}_{+}^{(b,a_{+})} \cup \mathscr{L}_{-}^{(a_{-},b)} \cup \mathscr{L}_{+}^{(a_{-},b)} \\ \begin{bmatrix} \frac{1}{e^{-n\varphi(z)}} & 0 \\ \frac{1}{Q(z)} & 1 \end{bmatrix} & z \in \mathscr{L}_{+}^{(b,1)} \\ \begin{bmatrix} \frac{1}{e^{-n\varphi(z)}} & 0 \\ \frac{1}{Q(z)} & (1+\frac{1}{\varkappa}) & 1 \end{bmatrix} & z \in \mathscr{L}_{b}^{c_{+}} \\ \begin{bmatrix} \frac{1}{e^{-n\varphi(z)}} & 0 \\ \frac{1}{Q(z)} & (\frac{e^{-2i\pi\varphi}}{\varkappa} - 1) & 1 \end{bmatrix} & z \in \mathscr{L}_{b}^{c_{-}} \\ \begin{bmatrix} \frac{1}{e^{-n\varphi(z)}} & 0 \\ \frac{1}{\varkappa Q(z)} & 1 \end{bmatrix} & z \in \mathscr{L}_{b}^{c_{-}} \\ \begin{bmatrix} 0 \\ \frac{e^{-n\varphi(z)}}{\varkappa Q(z)} & 0 \end{bmatrix} & z \in \mathscr{L}_{-}^{(b,1)} \\ \begin{bmatrix} 0 \\ \frac{e^{-n\Omega_{1}}}{\varkappa Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{b}^{1} \\ \begin{bmatrix} 0 \\ -\frac{e^{-n\Omega_{2}}}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{b}^{1} \\ \begin{bmatrix} 0 \\ -\frac{e^{-n\Omega_{2}}}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{a_{-}}^{b_{+}} \end{cases}$$

Model problem. The model problem that we have to solve is the following one Problem 5.8 Find a matrix valued function  $M_Q$ , analytic in  $\mathbb{C} \setminus \Gamma_m$  such that

$$\begin{split} M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} 0 & Q(z) \\ \frac{-1}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{a_{-}}^{b} \\ M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} 0 & Q(z) e^{n\Omega_{2}} \\ \frac{-e^{-n\Omega_{2}}}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{b}^{a_{+}} \\ M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} 0 & \varkappa Q(z_{+}) e^{n\Omega_{1}} \\ \frac{-e^{-n\Omega_{1}}}{\varkappa Q(z_{+})} & 0 \end{bmatrix} & z \in \Gamma_{b}^{1} \end{split}$$
(5.56)

with  $\varkappa$  as in (5.38) and Q as in (5.3). Furthermore the following local behaviours hold

$$M_Q(z) = \mathbf{1} + \mathcal{O}(z^{-1}) \qquad |z| \to \infty \tag{5.57}$$

$$M_Q(z) = \mathcal{O}\left(\frac{1}{(z-q)^{\frac{1}{4}}}\right), \quad q = 1, b, a_{\pm} \qquad z \to q.$$
 (5.58)

To further normalize the problem we need to construct a Szegö function, along the line of what we already have done in the case of the outside of the EoT.

The Szegö function. Let R(z) denote the radical function

$$R(z) = \sqrt{(z-1)(z-b)(z-a_{+})(z-a_{-})} = \sqrt{z^4 + (A-2)z^3 + \left(\frac{s^2}{4} - A + 1\right)z^2 - \frac{s^2}{2}z + \frac{s^2}{4}}$$
(5.59)

with the branch-cuts chosen along the three arcs of  $\Gamma_m = \Gamma_b^1 \cup \Gamma_b^{a_+} \cup \Gamma_{a_-}^b$ , and with the overall determination such that  $R(z) \simeq z^2$  as  $z \to \infty$ . Consider the following expression:

$$S(z) = R(z) \left( \int_{b}^{1} \frac{\ln Q(w_{+}) + \ln \varkappa - \nu}{R(w_{+})(w-z)} \frac{\mathrm{d}w}{2i\pi} + \int_{a_{-}}^{b} \frac{\ln Q(w)}{R(w_{+})(w-z)} \frac{\mathrm{d}w}{2i\pi} + \int_{b}^{a_{+}} \frac{\ln Q(w)}{R(w_{+})(w-z)} \frac{\mathrm{d}w}{2i\pi} \right)$$
(5.60)

The constant  $\nu$  is chosen so that the function S(z) is bounded as  $|z| \to \infty$ . We can actually simplify (5.60) quite a bit; consider, for example the integral

$$I_1 := \int_b^1 \frac{\ln Q(w_+) + \ln \varkappa - \nu}{R(w_+)(w-z)} \frac{\mathrm{d}w}{2i\pi}$$
(5.61)

Since, along  $\Gamma_b^1$  we have  $Q(w_+) = Q(w_-)e^{2i\pi\rho}$  and  $R(w_+) = -R(w_-)$  we can convert  $I_1$  into

$$I_1 = \frac{1}{2} \int_b^1 \frac{\ln Q(w_+) \mathrm{d}w}{R(w_+)(w-z)2i\pi} + \frac{1}{2} \int_1^b \frac{\ln Q(w_-) \mathrm{d}w}{R(w_-)(w-z)2i\pi} + \int_b^1 \frac{i\pi \rho + \ln \varkappa - \nu \mathrm{d}w}{R(w_+)(w-z)2i\pi}$$
(5.62)

Along the same lines, denoting the other two integrals in (5.60) by  $I_2, I_3$  we have

$$I_2 = \frac{1}{2} \int_{a_-}^{b} \frac{\ln Q(w) \mathrm{d}w}{R(w_+)(w-z)2i\pi} + \frac{1}{2} \int_{b}^{a_-} \frac{\ln Q(w) \mathrm{d}w}{R(w_-)(w-z)2i\pi}$$
(5.63)

$$I_{3} = \frac{1}{2} \int_{b}^{a_{+}} \frac{\ln Q(w) \mathrm{d}w}{R(w_{+})(w-z)2i\pi} + \frac{1}{2} \int_{a_{+}}^{b} \frac{\ln Q(w) \mathrm{d}w}{R(w_{-})(w-z)2i\pi}.$$
(5.64)

Adding  $I_1 + I_2 + I_3$  we observe that the integrals involving  $\ln Q(w)$  form a closed loop from b that goes around the whole  $\Gamma_m$  in the clockwise direction without intersecting the cut of Q(z). Using (5.3) we observe that we have

$$\ln Q(z_{+}) = \ln Q(z_{-}) + \begin{cases} 2i\pi \mathbf{K} & z \in (-\infty, 0) \\ 2i\pi \boldsymbol{\rho} & z \in (0, 1). \end{cases}$$
(5.65)

By using Cauchy's theorem we conclude that

$$S(z) = \frac{1}{2}\ln Q(z) + R(z) \left( \int_{-\infty}^{0} \frac{-\mathbf{K} dw}{2R(w)(w-z)} - \frac{1}{2} \int_{b}^{0} \frac{\mathbf{\rho} dw}{R(w)(w-z)} + \int_{b}^{1} \frac{(i\pi\mathbf{\rho} + \ln\varkappa - \nu) dw}{R(w_{+})(w-z)2i\pi} \right).$$
(5.66)

Observe that the branch-cut of  $\ln Q(z)$  on  $(-\infty, 0)$  is compensated by the branch-cut of the first integral in the above expression, so that S(z) is actually analytic across the ray  $(-\infty, 0)$ . From (5.66) it is then simple to ascertain that to make S(z) bounded at infinity we need to choose the value of  $\nu$  as follows (see (5.38) for  $\varkappa$ )

$$\nu = i\pi\rho \left(1 - \frac{\int_b^0 \frac{\mathrm{d}w}{R(w)}}{\int_b^1 \frac{\mathrm{d}w}{R(w_+)}}\right) + \ln\varkappa.$$
(5.67)

The expression (5.60) makes it clear, through the use of the Sokhostki–Plemelji formula, that the following properties of the Szegö function hold:

**Proposition 5.2 (Szegö function for** Re(s) > 0) The function S(z) defined in (5.60) or equivalently (5.66) it analytic and locally bounded on  $\mathbb{C} \cup \{\infty\} \setminus \Gamma_m$ , and with boundary conditions

$$S(z_{+}) + S(z_{-}) = \begin{cases} \ln (\varkappa Q(z_{+})) - \nu & z \in \Gamma_{b}^{1} \\ \ln Q(z) & z \in \Gamma_{a_{-}}^{b} \cup \Gamma_{b}^{a_{+}}. \end{cases}$$
(5.68)

Furthermore S(z) is bounded near  $z = b, a_{\pm}$  and near z = 1 it has the behaviour ([19] Ch. 5)

$$S(z) = -\frac{\rho}{2}\ln(z-1) + \mathcal{O}(1).$$
(5.69)

**Remark 5.2 (Alternative description/derivation of the Szegö function)** Proceeding from the desired properties in Proposition (5.2) and differentiating we obtain

$$S'(z_{+}) + S'(z_{-}) = \frac{Q'(z)}{Q(z)}, \quad z \in \Gamma_{m}.$$
(5.70)

Then one can seek a solution of the form

$$S'(z) = \frac{1}{2} \frac{Q'(z)}{Q(z)} + \eta(z)$$
(5.71)

where  $\eta(z)$  must be a function analytic off  $\Gamma_m$  with  $\eta(z_+) + \eta(z_-) = 0$  on  $\Gamma_m$  and such as to cancel out the singularities of  $\frac{Q'}{Q}$  in (5.71) outside of  $\Gamma_m$ . Since  $\frac{Q'}{Q} = \frac{\rho}{z-1} + \frac{K-\rho}{z}$  we see that

$$\eta(z) = \frac{\rho - K}{2} \frac{R(0)}{zR(z)} - \frac{C}{R(z)}.$$
(5.72)

which cancels the pole at z = 0 in the expression (5.71). Here R is given by (5.59). The constant C is determined by the condition that  $S(z_+) + S(z_-) = \ln Q(z)$  on the arcs  $\Gamma_b^{a_{\pm}}$ , which implies that integral of  $\eta$  along the contour  $\tilde{\gamma}$  on the Riemann surface of  $W^2 = R(z)^2$  (see Fig. 8, left pane) vanishes.

$$C = \frac{(\boldsymbol{\rho} - \boldsymbol{K})R(0)}{2} \frac{\oint_{\widetilde{\gamma}_{+}} \frac{\mathrm{d}z}{zW}}{\oint_{\widetilde{\gamma}} \frac{\mathrm{d}z}{W}}.$$
(5.73)

Using the explicit expression of R in (5.59) we note that  $R(0) = -\frac{s}{2}$  and can simplify to

$$C = C_{+} = \frac{(\mathbf{K} - \mathbf{\rho})}{4} \frac{s \oint_{\widetilde{\gamma}_{+}} \frac{dz}{zW}}{\int_{b}^{1} \frac{dz}{R(z_{+})}},$$
(5.74)

where we have used the fact that  $\oint_{\widetilde{\gamma}} \frac{\mathrm{d}z}{R(z)} = -\int_b^1 \frac{\mathrm{d}z}{R(z_+)}$  (see Fig. 8, left pane).

When  $\operatorname{Re}(s) < 0$  instead we have to ask that the antiderivative of (5.71) is continuous on  $\mathbb{C} \setminus \Gamma_m$  and then the contour  $\tilde{\gamma}$  is different. In this case then the constant C is given by

$$C = C_{-} = \frac{(\boldsymbol{\rho} - \boldsymbol{K})R(0)}{2} \frac{\oint_{\widetilde{\gamma}_{-}} \frac{\mathrm{d}z}{zW}}{\oint_{\widetilde{\gamma}_{-}} \frac{\mathrm{d}z}{W}} = \frac{(\boldsymbol{K} - \boldsymbol{\rho})s}{4} \frac{\oint_{a_{+}}^{a_{-}} \frac{\mathrm{d}z}{R(z_{+})}}{\oint_{a_{+}}^{a_{-}} \frac{\mathrm{d}z}{R(z_{+})}}.$$
(5.75)

We observe that, in fact, the contours  $\tilde{\gamma}_{\pm}$  are homologically equivalend on the Riemann surface punctured at the two points z = 0 on both sheets. However the definition of the domain of R(z) has been chosen with different branch-cuts. Keeping this in mind, we can use for C the expression (5.75) in both cases with the understanding that in the case  $\operatorname{Re}(s) > 0$  the indication of the boundary value  $R(z_{+})$  is irrelevant.

In either case, for future reference, we observe that (recalling the definition of R (5.59))

$$S'(0) = -\frac{\rho}{2} \frac{R'(0)}{R(0)} - \frac{C_{\pm}}{R(0)} = \frac{\rho}{2} + \frac{(K-\rho)}{2} \frac{\oint_{a_{+}}^{u_{-}} \frac{dz}{zR(z_{+})}}{\oint_{a_{+}}^{a_{-}} \frac{dz}{R(z_{+})}}.$$
(5.76)

~ a



Figure 8: The alternative computation of the Szego function and the contour  $\tilde{\gamma}$ . For Re (s) > 0 in the left pane, for Re (s) < 0 in the right pane. The dashed part of the contour  $\tilde{\gamma}$  correspond to the second sheet of the radical function R(z) (the one where  $R(z) \simeq -z^2$  at infinity). The two contours are homologically equivalent.

With the Szegö function at hand we can normalize the RHP 5.8. We define a new model problem

$$M(z) := e^{S(\infty)\sigma_3} M_{_{O}}(z) e^{-S(z)\sigma_3}.$$
(5.77)

The effect of this normalization is to turn the jump matrices (5.56) into constant ones as detailed in the following problem

**Problem 5.9** (*RHP<sub>M</sub>*) Find a matrix-valued function M(z) analytic in  $\mathbb{C} \setminus \Gamma_m$  such that

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \qquad z \in \Gamma_{a_{-}}^{b}$$

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} 0 & e^{n\Omega_{2}} \\ -e^{-n\Omega_{2}} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{a_{+}}$$

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} 0 & e^{n\Omega_{1}+\nu} \\ -e^{-n\Omega_{1}-\nu} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{1} \qquad (5.78)$$

and satisfying

$$M(z) = \mathbf{1} + \mathcal{O}(z^{-1}) \qquad |z| \to \infty \tag{5.79}$$

$$M(z) = \mathcal{O}\left(\frac{1}{(z-q)^{\frac{1}{4}}}\right), \quad q = 1, b, a_{\pm} \qquad z \to q.$$
 (5.80)

The solution of the RHP 5.9 is relatively standard and it requires the use of the Jacobi theta function.

Solution of the RHP 5.9. With the same definition of R in (5.59) we also define

$$h(z) := \frac{1}{\left((z-1)(z-b)(z-a_{+})(z-a_{-})\right)^{\frac{1}{4}}},$$
(5.81)

with the same branch-cuts as R along  $\Gamma_m$  and the determination chosen so that  $h(z) \simeq \frac{1}{z}$  as  $z \to \infty$ . A careful analysis of the phases shows that

$$h(z_{+}) = ih(z_{-}) \qquad \qquad z \in \Gamma_{a_{-}}^{b}$$

$$(5.82)$$

$$h(z_{+}) = -ih(z_{-}) \qquad \qquad z \in \Gamma_b^1 \cup \Gamma_b^{a_{+}}.$$
(5.83)

We also need the **Abel map**: let

$$\omega_1 := \int_b^1 \frac{\mathrm{d}z}{R(z_+)}, \qquad \omega_2 := \int_b^{a_+} \frac{\mathrm{d}z}{R(z_-)}, \tag{5.84}$$

$$\mathfrak{u}(z) := \int_{a_{-}}^{z} \frac{\mathrm{d}w}{2\omega_{1}R(w)}, \quad \tau := \frac{\omega_{2}}{\omega_{1}}, \tag{5.85}$$

where the contour of integration runs in the simply connected domain  $\mathbb{C} \setminus \Gamma$ . A direct inspection reveals Lemma 5.1 (Properties of the Abel map) *The following relations hold:* 

$$\mathfrak{u}(z_{+}) = -\mathfrak{u}(z_{-}) + \begin{cases} 0 & z \in \Gamma_{a_{-}}^{b} \\ -1 & z \in \Gamma_{b}^{a_{+}} \\ -\tau & z \in \Gamma_{b}^{1}. \end{cases}$$
(5.86)

Let  $\vartheta(u;\tau)$  be the Riemann (Jacobi) theta function (also denoted  $\theta_4$ ):

$$\vartheta(u;\tau) := \sum_{n \in \mathbb{Z}} e^{in^2 \pi \tau + 2i\pi nu}.$$
(5.87)

The function  $\vartheta$  vanishes at  $u = \frac{\tau+1}{2} + k + \ell \tau$  for all  $k, \ell \in \mathbb{Z}$  and satisfies the quasi-periodicity properties

$$\vartheta(u+k+\ell\tau;\tau) = e^{-2i\pi\ell u - i\pi\ell^2\tau} \vartheta(u;\tau).$$
(5.88)

Consider the following two row-vectors;

$$\phi(z;\infty) = [\phi_1(z;\infty), \phi_2(z;\infty)],$$

$$\phi_1(z;\infty) = \frac{\vartheta \left(\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{i\pi K\mathfrak{u}(z)}$$

$$\phi_2(z;\infty) = \frac{-i\vartheta \left(-\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(-\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{-i\pi K\mathfrak{u}(z)}$$
(5.89)

and

$$\psi(z;\infty) = \left[\psi_1(z;\infty), \psi_2(z;\infty)\right]$$
  

$$\psi_1(z;\infty) = \frac{-i\vartheta\left(\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right)h(z)}{\vartheta\left(\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{i\pi K\mathfrak{u}(z)}$$
  

$$\psi_2(z;\infty) = \frac{-\vartheta\left(-\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right)h(z)}{\vartheta\left(-\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{-i\pi K\mathfrak{u}(z)}.$$
(5.90)

Furthermore all entries are bounded by (5.80) by the very definition of h(z) (5.81) and the fact that  $\vartheta$  is an entire function with zeros only at the half-period.

Let us investigate the behaviour of  $\phi, \psi$  near  $z = \infty$ : a direct computation shows that

$$\lim_{z \to \infty} \begin{bmatrix} \phi \\ \psi \end{bmatrix} = \begin{bmatrix} \frac{\vartheta \left(G - \frac{\tau+1}{2}\right)}{\omega_1 \vartheta' \left(-\frac{\tau+1}{2}\right)} e^{i\pi K \mathfrak{u}(\infty)} & 0\\ 0 & \frac{\vartheta \left(G - \frac{\tau+1}{2}\right)}{\omega_1 \vartheta' \left(-\frac{\tau+1}{2}\right)} e^{-i\pi K \mathfrak{u}(\infty)} \end{bmatrix}$$
(5.91)

Thus, as long as the common factor  $\vartheta \left(G - \frac{\tau+1}{2}\right) \neq 0$  we can define (the dependence on the parameters G, K is understood in the right side)

$$\widehat{M}(z;G,K,\infty) := \frac{\omega_1 \vartheta'\left(-\frac{\tau+1}{2}\right) e^{-i\pi K \mathfrak{u}(\infty)\sigma_3}}{\vartheta\left(G - \frac{\tau+1}{2}\right)} \begin{bmatrix} \boldsymbol{\phi}(z;\infty) \\ \boldsymbol{\psi}(z;\infty) \end{bmatrix}$$
(5.92)

**Remark 5.3** We have emphasized in the notation that  $\infty$  as a point plays a role in the expression. We will use later the same formula, but replacing  $\infty$  with the point z = 0. It is important to point out that, irrespectively of what point we replace instead of  $\infty$ , the matrix satisfies the same boundary relations (5.94) below, and also the same behaviour near the branch-points  $z = 1, b, a_{\pm}$ . The only difference is that if we replace  $\infty$  by a point  $z_0$ , the matrix  $\widehat{M}(z; G, K, z_0)$  will then vanish at infinity and have a simple pole at  $z = z_0$  with singular part proportional to the identity matrix:

$$\lim_{z \to \infty} \widehat{M}(z; G, K, z_0) = \mathbf{0}, \qquad \widehat{M}(z; G, K, z_0) = \frac{\omega_1 h(0)}{\mathfrak{u}'(z_0)} \frac{\mathbf{1}}{z - z_0} + \mathcal{O}(1), \quad z \to z_0.$$
(5.93)

We are going to use this observation later.

It is a direct verification using the definition of the Abel map (5.85), the properties of the theta function (5.88), the jump relation of h (5.82) that each row  $\phi, \psi$  satisfy the three boundary value relations similar to those in (5.78):

$$\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} 0 & 1\\ -1 & 0 \end{bmatrix} \qquad z \in \Gamma_{a_{-}}^{b} 
\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} 0 & -e^{i\pi K} \\ e^{-i\pi K} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{a_{+}} 
\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} 0 & -e^{-2i\pi \left(G - \frac{K\tau}{2}\right)} \\ e^{2i\pi \left(G - \frac{K\tau}{2}\right)} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{1} \qquad (5.94)$$

By matching these to the boundary relations (5.78) we deduce that

**Proposition 5.3** The solution of the RHP (5.9) is given by  $M(z) = \widehat{M}(z; G, K, \infty)$  in (5.92) with the values of the constants G, K given by

$$G = \frac{1}{2i\pi} \left( -n\Omega_1 - \nu + n\Omega_2 \tau \right) + \frac{\tau + 1}{2}, \qquad K = \frac{n\Omega_2}{i\pi} + 1$$
(5.95)

**Proof.** Comparing the boundary relations (5.94) and (5.78) we obtain the system

$$\begin{cases} G - \frac{K\tau}{2} = -\frac{n\Omega_1 + \nu}{2i\pi} + \frac{1}{2} \\ K = \frac{n\Omega_2}{i\pi} + 1. \end{cases}$$
(5.96)

from which the relations follow.

The solvability of the RHP 5.9 and hence of the RHP 5.8 depend entirely on the non-vanishing of the expression

$$\vartheta\left(\frac{n\Omega_1 + \nu - n\Omega_2\tau}{2i\pi}\right) \tag{5.97}$$

with  $\Omega_1, \Omega_2$  defined by (5.51) and  $\nu$  by (5.67) Considering that the zeros of  $\vartheta$  are for  $u = \frac{\tau+1}{2} + \ell + k\tau$ , with  $k, \ell \in \mathbb{Z}$ , we obtain the quantization conditions for the periods of  $\varphi$ 

$$(H_1, H_2) \in \mathbb{Z}^2$$

$$H_1 := \frac{n\Omega_1}{2i\pi} + \operatorname{Re}\left(\frac{\nu}{2i\pi}\right) - \frac{\operatorname{Re}\tau}{\operatorname{Im}\tau}\operatorname{Im}\left(\frac{\nu}{2i\pi}\right) + \frac{1}{2}$$

$$H_2 := \frac{n\Omega_2}{2i\pi} - \frac{\operatorname{Im}\left(\frac{\nu}{2i\pi}\right)}{\operatorname{Im}\tau} + \frac{1}{2}.$$
(5.98)

Observe that both  $H_1, H_2$  are real functions of s since  $\Omega_1, \Omega_2 \in i\mathbb{R}$ . The modular parameter  $\tau$  is also, in a very implicit way, a function of s since the branch-points of the radical R (5.59) are determined by the Boutroux conditions (4.31). Ditto for  $\nu$ , which depends on s implicitly via the formula (5.67). **Explanation of Fig. 2.** The two quantization conditions (5.98) can be interpreted as describing a mesh of level-sets of the two functions  $H_1, H_2$  (both functions of the complex parameter s); in the various panels in Fig. 2 these are precisely forming the mesh of curves that populate the interior of EoT. The intersection points of this mesh are the points where the quantization conditions (5.98) hold, and hence where the solution of the model problem RHP 5.8 ceases to exist. They also approximate very precisely the zeros of the rational solutions (in fact better than expected), with only obvious deviations near the boundaries of the two halves of the EoT, since there the elliptic curve of the radical R (5.59) degenerates (i.e., two branch-points come together).

#### 5.2.2 Second transformation, model problem and its solution: the case $\operatorname{Re}(s) < 0$

With reference to Fig. 7, we denote by  $\Lambda_{\pm}^{(x,y)}$  the region bounded by  $\Gamma_x^y$  and  $\mathscr{L}_{\pm}^{(x,y)}$ , for  $x, y \in \{a_+, a_-, b, 1\}$  and we call them *lens regions* (as opposed to their boundaries, which we refer to as the lens arcs).

**Definition of**  $T_0$  for  $\operatorname{Re}(s) < 0$ . We refer to the right pane of Fig. 7 and then set

$$T_{0}(z) := \begin{cases} W(z) \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{-}^{(a_{+},a_{-})} \\ W(z) \begin{bmatrix} 1 & 0 \\ -\frac{e^{-n\varphi(z)}}{Q(z)} & 1 \end{bmatrix} & z \in \Lambda_{+}^{(a_{+},a_{-})} \end{cases}$$
(5.99)

while, for the lens  $\Lambda_{\pm}^{(b,1)}$  adjacent to  $\Gamma_b^1$  we have the same as in (5.44). The construction is similar to the previous case, but with the notable difference that now the main arcs  $\Gamma_m$  consists of two disjoint arcs (see Fig. 7, right pane)

$$\Gamma_m = \Gamma_b^1 \cup \Gamma_{a_+}^{a_-}, \qquad \Gamma_c = \Gamma_b^{a_+} \cup \Gamma_{a_-}^{b}.$$
(5.100)

We define the effective potential as before

$$\varphi(z;s) := \int_{a_{-}}^{z} \frac{2}{w^2} \sqrt{w^2 + \frac{s^2}{4} + \frac{Aw^2}{w-1}} \mathrm{d}w, \qquad (5.101)$$

where now, however, the branch-cuts of the radical in the integrand consist of the two arcs  $\Gamma_b^1$ ,  $\Gamma_{a_+}^{a_-}$  of Fig. 7, right pane, and  $\varphi$  has additionally a branchcut  $[1,\infty)$  and another one running along  $\Gamma_b^{a_+}$ . Now the effective potential satisfies the following boundary value relations:

$$\begin{aligned}
\varphi(z_{+}) + \varphi(z_{-}) &= 0 & z \in \Gamma_{a_{+}}^{a_{-}} \\
\varphi(z_{+}) + \varphi(z_{-}) &= 2\Omega_{1} & z \in \Gamma_{b}^{1} \\
\varphi(z_{+}) - \varphi(z_{-}) &= 0 & z \in \Gamma_{a_{-}}^{b} \\
\varphi(z_{+}) - \varphi(z_{-}) &= 2\Omega_{2} & z \in \Gamma_{b}^{a_{+}} \\
\varphi(z_{+}) - \varphi(z_{-}) &= 4i\pi & z \in [1,\infty)
\end{aligned}$$
(5.102)

where now

$$\Omega_1 := \int_{a_-}^b \varphi'(w) \mathrm{d}w, \quad \Omega_2 = \int_{a_-}^{a_+} \varphi'(w_+) \mathrm{d}w$$
 (5.103)

A direct computation shows that the matrix  $T_0(z)$  satisfies the following RHP

**Problem 5.10** (*RHP*<sub> $T_0$ </sub>) The matrix  $T_0(z)$  satisfies the conditions

$$T_0(z_+) = T_0(z_-)J_{T_0}(z), \quad z \in \gamma$$
(5.104)

$$T_0(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{z}\right), \quad z \to \infty$$
 (5.105)

where (we set for brevity  $Q_{\pm} := Q(z_{\pm}), \varphi_{\pm} := \varphi(z_{\pm})$  below)

$$J_{T_{0}}(z) = \begin{cases} J_{W}(z) = \begin{bmatrix} e^{-n\Omega_{2}} & Qe^{\frac{n}{2}(\varphi_{+}+\varphi_{-})} \\ 0 & e^{n\Omega_{2}} \end{bmatrix} & z \in \Gamma_{b}^{a_{+}} \\ J_{W}(z) = \begin{bmatrix} 1 & Qe^{n\varphi} \\ 0 & 1 \end{bmatrix} & z \in \Gamma_{a_{-}}^{b} \\ \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi}}{Q} & 1 \end{bmatrix} & z \in \mathscr{L}_{+}^{(a_{+},a_{-})} \cup \mathscr{L}_{-}^{(a_{+},a_{-})} \\ \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi}}{\varkappa Q} & 1 \end{bmatrix} & z \in \mathscr{L}_{+}^{(b,1)} \\ \begin{bmatrix} 1 & 0 \\ \frac{e^{-n\varphi-2i\pi\rho}}{\varkappa Q} & 1 \end{bmatrix} & z \in \mathscr{L}_{-}^{(b,1)} \\ \begin{bmatrix} 0 & \varkappa Q_{+}e^{n\Omega_{1}} \\ -\frac{e^{-n\Omega_{1}}}{\varkappa Q_{+}} & 0 \end{bmatrix} & z \in \Gamma_{b}^{b} \\ \begin{bmatrix} 0 & Q \\ -\frac{1}{Q} & 0 \end{bmatrix} & z \in \Gamma_{a_{+}}^{a_{-}} \end{cases}$$

**Problem 5.11** Find a matrix valued function  $M_Q$ , analytic in  $\mathbb{C} \setminus \Gamma_m$  such that

$$\begin{split} M_{Q}(z_{+}) &= M_{Q}(z_{-}) & z \in \Gamma_{a_{-}}^{b} \cup [1, \infty) \\ M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} 0 & Q(z) \\ \frac{-1}{Q(z)} & 0 \end{bmatrix} & z \in \Gamma_{a_{+}}^{a_{-}} \\ M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} e^{-n\Omega_{2}} & 0 \\ 0 & e^{n\Omega_{2}} \end{bmatrix} & z \in \Gamma_{b}^{a_{+}} \\ M_{Q}(z_{+}) &= M_{Q}(z_{-}) \begin{bmatrix} 0 & \varkappa Q(z_{+})e^{n\Omega_{1}} \\ \frac{-e^{-n\Omega_{1}}}{\varkappa Q(z_{+})} & 0 \end{bmatrix} & z \in \Gamma_{b}^{1} \end{split}$$
(5.107)

with  $\varkappa$  as in (5.38) and Q as in (5.3). Furthermore the following local behaviours hold

$$M_Q(z) = \mathbf{1} + \mathcal{O}(z^{-1}) \qquad |z| \to \infty \tag{5.108}$$

$$M_{Q}(z) = \mathcal{O}\left(\frac{1}{(z-q)^{\frac{1}{4}}}\right), \quad q = 1, b, a_{\pm} \qquad z \to q.$$
 (5.109)

Like before, to further normalize the problem we need to construct an appropriate Szegö function.

**The Szegö function.** Let R(z) denote the radical function (5.59), but now with the branch-cuts of the radical chosen along  $\Gamma_m = \Gamma_b^1 \cup \Gamma_{a_+}^{a_-}$  (right pane of Fig. 7). Consider the following expression:

$$S(z) = R(z) \left( \int_{b}^{1} \frac{\ln Q(w_{+}) + \ln \varkappa - \nu}{R(w_{+})(w-z)} \frac{\mathrm{d}w}{2i\pi} + \int_{a_{+}}^{a_{-}} \frac{\ln Q(w)}{R(w_{+})(w-z)} \frac{\mathrm{d}w}{2i\pi} \right)$$
(5.110)

Following analogous considerations as in the case  $\operatorname{Re}(s) > 0$  (Sec. 5.2.1) we finally obtain the same expression (5.66), where the condition that S(z) is bounded at infinity imposes the same constraint on  $\nu$  as in (5.67). Similarly to Prop. 5.2 we have now:

**Proposition 5.4 (Szegö function for** Re(s) < 0) The function S(z) defined in (5.110) or equivalently (5.66) it analytic and bounded on  $\mathbb{C} \cup \{\infty\} \setminus \Gamma_m$ , and with boundary conditions

$$S(z_{+}) + S(z_{-}) = \begin{cases} \ln (\varkappa Q(z_{+})) - \nu & z \in \Gamma_{b}^{1} \\ \ln Q(z) & z \in \Gamma_{a_{+}}^{a_{-}}. \end{cases}$$
(5.111)

Furthermore S(z) is bounded near  $z = b, a_{\pm}$  and near z = 1 it has the behaviour

$$S(z) = -\frac{\rho}{2}\ln(z-1) + \mathcal{O}(1).$$
(5.112)

With this new Szegö function we normalize the RHP 5.11 and define the new model problem

$$M(z) := e^{S(\infty)\sigma_3} M_Q(z) e^{-S(z)\sigma_3}.$$
(5.113)

The new matrix M solves now the following problem.

**Problem 5.12** (*RHP<sub>M</sub>*) Find a matrix-valued function M(z) analytic in  $\mathbb{C} \setminus \Gamma_m$  such that

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \qquad z \in \Gamma_{a_{+}}^{a_{-}}$$

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} e^{-n\Omega_{2}} \\ 0 & e^{n\Omega_{2}} \end{bmatrix} \qquad z \in \Gamma_{b}^{a_{+}}$$

$$M(z_{+}) = M(z_{-}) \begin{bmatrix} 0 & e^{n\Omega_{1}+\nu} \\ -e^{-n\Omega_{1}-\nu} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{1} \qquad (5.114)$$

and satisfying the same (5.79), (5.80).

**Solution of the RHP 5.12.** With the same definition of R in (5.59), we also set the same definition of h as in (5.81) where however now the domain consists of  $\mathbb{C}$  minus the branchcuts  $\Gamma_b^1 \cup \Gamma_b^{a_+} \cup \Gamma_{a_+}^{a_-}$ , and the determination chosen so that  $h(z) \simeq \frac{1}{z}$  as  $z \to \infty$ . A careful analysis of the phases of h shows that

$$\begin{aligned} h(z_{+}) &= -ih(z_{-}) & z \in \Gamma_{b}^{1} \\ h(z_{+}) &= -ih(z_{-}) & z \in \Gamma_{a_{+}}^{a_{-}} \\ h(z_{+}) &= -h(z_{-}) & z \in \Gamma_{b}^{a_{+}}. \end{aligned}$$
 (5.115)

The Abel map is defined as in (5.85) but without the boundary value in the definition of  $\omega_2$  in (5.84) since now the contour  $\Gamma_b^{a_+}$  is not a branch-cut of the radical R. This time the Abel map satisfies somewhat different relations, a consequence of the different choice of branchcuts.

Lemma 5.2 (Properties of the Abel map) The following relations hold:

$$\mathfrak{u}(z_{+}) = -\mathfrak{u}(z_{-}) + \begin{cases} 0 & z \in \Gamma_{a_{+}}^{a_{-}} \\ -\tau & z \in \Gamma_{b}^{1}. \end{cases}$$

$$\mathfrak{u}(z_{+}) = \mathfrak{u}(z_{-}) - 1 & z \in \Gamma_{b}^{a_{+}}. \qquad (5.116)$$

Consider now the following two row-vectors;

$$\phi(z) = [\phi_1(z), \phi_2(z)],$$

$$\phi_1(z) = \frac{\vartheta \left(\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{i\pi K\mathfrak{u}(z)}$$

$$\phi_2(z) = \frac{i\vartheta \left(-\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(-\mathfrak{u}(z) - \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{-i\pi K\mathfrak{u}(z)}$$
(5.117)

and

$$\boldsymbol{\psi}(z) = [\psi_1(z), \psi_2(z)]$$
  

$$\psi_1(z) = \frac{\vartheta \left(\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{i\pi K \mathfrak{u}(z)}$$
  

$$\psi_2(z) = \frac{i\vartheta \left(-\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2} + G\right) h(z)}{\vartheta \left(-\mathfrak{u}(z) + \mathfrak{u}(\infty) - \frac{\tau+1}{2}\right)} e^{-i\pi K \mathfrak{u}(z)}.$$
(5.118)

These are essentially the same formulæ as (5.89) (5.90) except for a minor modification of the normalization. Accordingly we define  $\widehat{M}(z; G, K, \infty)$  exactly as in (5.92). The boundary relations for this matrix  $\widehat{M}$  are now:

$$\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \qquad \qquad z \in \Gamma_{a_{+}}^{a_{-}}$$
$$\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} -e^{-i\pi K} & 0 \\ 0 & -e^{i\pi K} \end{bmatrix} \qquad \qquad z \in \Gamma_{b}^{a_{+}}$$

$$\widehat{M}(z_{+}) = \widehat{M}(z_{-}) \begin{bmatrix} 0 & e^{-2i\pi \left(G - \frac{K\tau}{2}\right)} \\ -e^{2i\pi \left(G - \frac{K\tau}{2}\right)} & 0 \end{bmatrix} \qquad z \in \Gamma_{b}^{1}$$
(5.119)

By matching these to the boundary relations (5.114) we deduce that

**Proposition 5.5** The solution of the RHP (5.12) is given by  $\widehat{M}(z;G,K)$  in (5.92) with the values of the constants G, K given by

$$G = \frac{1}{2i\pi} \left( n\Omega_1 + \nu + n\Omega_2 \tau \right) + \frac{\tau}{2}, \qquad K = \frac{n\Omega_2}{i\pi} + 1$$
(5.120)

The proof is entirely similar to Prop. 5.3. The same observation holds and that the solvability of the RHP 5.12 and hence of the RHP 5.12 depend entirely on the non-vanishing of the expression

$$\vartheta\left(\frac{n\Omega_1 + \nu - n\Omega_2\tau}{2i\pi} + \frac{1}{2}\right) \tag{5.121}$$

with  $\Omega_1, \Omega_2$  defined by (5.103) and  $\nu$  by (5.67). Note the slight difference from (5.97). The quantization conditions for the periods of  $\varphi$  now read

$$\widetilde{H}_{1} := \frac{n\Omega_{1}}{2i\pi} + \operatorname{Re}\left(\frac{\nu}{2i\pi}\right) - \frac{\operatorname{Re}\tau}{\operatorname{Im}\tau}\operatorname{Im}\left(\frac{\nu}{2i\pi}\right) \in \mathbb{Z}$$
$$\widetilde{H}_{2} := \frac{n\Omega_{2}}{2i\pi} + \frac{\operatorname{Im}\left(\frac{\nu}{2i\pi}\right)}{\operatorname{Im}\tau} + \frac{1}{2} \in \mathbb{Z}.$$
(5.122)

The mesh of levelsets  $\widetilde{H}_1 \in \mathbb{Z} \ni \widetilde{H}_2$  are the gridlines shown in Fig. 2 in the portion  $\operatorname{Re}(s) < 0$  of each pane.

#### 5.2.3 Local parametrices

The construction of the local parametrices is similar to that of Section 5.1.1. The only parametrix that is less commonly encountered (but see [6]) is the one that needs to be defined at the point b in the case  $\operatorname{Re}(s) > 0$ , when three main arcs meet at the same point (see Fig. 7, left pane). Let  $\mathbb{D}_x$  denote a small disk centered at  $x \in \{a_+, a_-, b, 1\}$  and not containing any of the other branchpoints  $a_+, a_-, b, 1$ . Inside each of these disk one has to define an appropriate local solution of the RHP 5.7 or 5.10 (according to the sign of  $\operatorname{Re}(s) > 0$  or  $\operatorname{Re}(s) < 0$ , respectively). Such a local solution (a.k.a. "local parametrix") is constructed in terms of Airy functions or Bessel functions. More specifically:

- 1. inside  $\mathbb{D}_{a_+}, \mathbb{D}_{a_-}$  the local parametrix is constructed in a manner analogous to what we illustrated in Section 5.1.1;
- 2. inside  $\mathbb{D}_b$  the local parametrix is constructed also in terms of Airy functions, arranged however in a different formula along the lines explained in App. A.1;
- 3. inside  $\mathbb{D}_1$  the local parametrix is constructed in terms of Bessel functions; this can be found in ([41], Section 3, pag. 155–156), or ([4], Section 4.2), and we will not enter into details since it is mainly irrelevant to the formulas that are of interest in this paper.

#### 5.2.4 Summary; the approximating mesh of location of zeros

The jump matrices of the error matrix  $\mathscr E$  are of the form

$$J_{\mathscr{E}}(z) = M_{Q}(z) \left( \mathbf{1} + \mathcal{O}(n^{-1}) \right) M_{Q}^{-1}(z).$$
(5.123)

While in the region outside of EoT the model solution  $M_Q$  does not depend on n (see (5.1)), when  $s \in EoT$  the solution is instead given by Prop. 5.3 and then M (and consequently  $M_Q$  (5.77)) and hence depends on n. This has the effect that the conjugation in (5.123) may have the effect, if any of the entries of  $M_Q$  is large, of negating the validity of the estimate  $J_{\mathscr{E}} = \mathbf{1} + \mathcal{O}(n^{-1})$ . Given that  $\det(M) \equiv 1$ , the source of the potential divergence are the terms in the denominators in (5.92). Of these the only term that may potentially vanish is the term  $\vartheta \left(G - \frac{\tau+1}{2}\right)$ , which becomes (5.97) or (5.121) (depending on the sign of Re(s)), so that the correct error estimate is

$$J_{\mathscr{E}}(z) = \mathbf{1} + \mathcal{O}\left(\frac{1}{n\vartheta\left(G - \frac{\tau+1}{2}\right)^2}\right).$$
(5.124)

with G given by either (5.95) or (5.120) (depending on the sign of  $\operatorname{Re}(s)$ ).

Thus the sufficient condition of asymptotic solvability of the RHP 5.1 is the condition that

$$\vartheta\left(\frac{n\Omega_1(s) - n\Omega_2(s)\tau(s) + \nu(s)}{2i\pi} + \frac{1}{2}\mathcal{X}_{\{\operatorname{Re} s > 0\}}(s)\right) = o\left(\frac{1}{\sqrt{n}}\right),\tag{5.125}$$

with  $\mathcal{X}_{\{\operatorname{Re} s>0\}}(s)$  denoting the indicator function of the right half-plane. This condition fails inside the union of small disks of radii  $\mathcal{O}(n^{-\frac{3}{2}})$  in the *s*-plane centered at the points where  $\vartheta = 0$ , namely, the intersection of the mesh of lines indicated by (5.98) (for  $\operatorname{Re}(s) > 0$ ) or (5.122) (for  $\operatorname{Re}(s) < 0$ ).

The actual numerics shows a seemingly much faster convergence of the mesh to the actual position of the zeroes (see Fig. 2), which appears to be one of those serendipitous situations where the approximation works "better than expected".



Figure 9: The RHPs for the error terms in the two cases  $\operatorname{Re}(s) > 0$  (left) and  $\operatorname{Re}(s) < 0$ . Compare with Fig. 7.

## 6 Approximation of the Hamiltonian

The Hamiltonian is obtained from the Taylor expansion near z = 0 of the solution of the RHP 5.1 outside of the EoT, and RHP 5.5 inside the EoT, according to Proposition 2.2.

In either cases the chain of transformations  $(Y \to W \to T_0 \sim M_Q)$  and the subsequent approximation imply that in a neighbourhood of z = 0 we have

$$Y_n(z) = e^{-n\frac{\ell}{2}\sigma_3} W(z) e^{n\left(g(z) + \frac{\ell}{2}\right)\sigma_3} = e^{-n\frac{\ell}{2}\sigma_3} \mathscr{E}(z) M_Q(z) e^{n\left(g(z) + \frac{\ell}{2}\right)\sigma_3}$$
(6.1)

where  $\mathscr{E}(z)$ , as discussed in Sec. 5.2.4, is a matrix that is  $\mathbf{1} + \mathcal{O}\left(\frac{1}{n\vartheta\left(G-\frac{\tau+1}{2}\right)^2}\right)$  near the origin (as long as a bas finite distance from the boundary of the FoT and the imaginary arithm it)

 $\boldsymbol{s}$  has finite distance from the boundary of the EoT and the imaginary axis within it).

We thus have

$$Y_{n}^{-1}Y_{n}'\Big|_{z=0} = \left(e^{-ng\sigma_{3}}\left(M_{Q}^{-1}M_{Q}' + M_{Q}^{-1}\mathscr{E}^{-1}\mathscr{E}'M_{Q}\right)e^{ng\sigma_{3}} + ng'(z)\sigma_{3}\right)\Big|_{z=0}$$
(6.2)

Given the expression of M in either Prop. 5.1 (for s outside the EoT), Prop. 5.3 for  $s \in EoT \cap \{\operatorname{Re}(s) > 0\}$ , or Prop. 5.5  $s \in EoT \cap \{\operatorname{Re}(s) < 0\}$  we have

$$Y_n^{-1}Y_n'\Big|_{z=0} = \left(e^{(-ng-S)\sigma_3} \left(M^{-1}M' + M^{-1}\mathscr{E}^{-1}\mathscr{E}'M\right)e^{(S+ng)\sigma_3} + (S'+ng'(z))\sigma_3\right)\Big|_{z=0},$$
(6.3)

where clearly the expressions for g, S also depend on which region we are considering.

We are interested in the (1, 1) entry and hence the conjugation in the first term by the diagonal matrix  $e^{(S-ng)\sigma_3}$  is immaterial. Furthermore the term containing  $\mathscr{E}$  yields a sub-leading contribution which we ignore for the purpose of this computation. In principle we should fork the computation according to the three regions (i) outside EoT; (ii) inside EoT with  $\operatorname{Re}(s) > 0$ ; (iii) inside EoT with  $\operatorname{Re}(s) < 0$ . However the computation inside EoT would be rather formal because we have not specified the correct orders of approximations when s is in a neighbourhood of one of the zeros. Additionally the explicit approximation is of no particular interest to us and thus for simplicity we decided to forego it entirely in this paper.

#### 6.1 Hamiltonian outside EoT

Using the explicit expression of M(z) in Prop. 5.1 we have

$$M^{-1}M' = \frac{i\sigma_2}{s} \tag{6.4}$$

and hence this term does not contribute being off-diagonal. Using (5.22) and the fact that  $R(0) = -\frac{s}{2}$ ,  $R(1) = -\sqrt{1 + \frac{s^2}{4}}$  with the root's determination such that  $R(1;s) \simeq -\frac{s}{2}$  as  $|s| \to \infty$ , we find

$$S'(0) = \frac{K}{s} + \frac{2\rho}{s^2 + s\sqrt{s^2 + 4}}.$$
(6.5)

Using then the explicit expression of g(z) derived from (4.17)

$$g'(z) = \frac{1}{2}\varphi'(z;s) + \frac{s}{2z^2} = \frac{\sqrt{z^2 + \frac{s^2}{4}}}{z^2} + \frac{s}{2z^2}.$$
(6.6)

Since g'(z) is regular at z = 0 we need to take the determination of the root near z = 0 that tends to  $-\frac{s}{2}$  and hence

$$g'(z) = -s\frac{\sqrt{1 + \frac{4z^2}{s^2}}}{2z^2} + \frac{s}{2z^2} = -\frac{1}{s} + \mathcal{O}(z) \quad \Rightarrow \quad g'(0;s) = -\frac{1}{s}.$$
(6.7)

We thus conclude that

**Proposition 6.1** For s in closed subsets outside of EoT we have the uniform approximation

$$H_V(s) = \frac{n - \mathbf{K}}{s} - \frac{2\rho}{s^2 + s\sqrt{s^2 + 4}} + \frac{\rho}{2} + \mathcal{O}(n^{-1}).$$
(6.8)

# 7 Conclusion

In this paper we have considered the particular scaling where we send  $n \to \infty$  and rescale only the independent variable t = ns (4.2). However, as evidenced also in Figure 2, if  $\rho$  or K or both are large relative to n, the approximation needs to be modified.

If we let  $\rho = n\rho$ ,  $\mathbf{K} = n\varkappa$  then we would need to construct a different *g*-function where  $\theta$  in (4.2) is replaced by

$$\theta(z;s) = n\left(\frac{s}{z} + \rho \ln\left(1 - \frac{1}{z}\right) + \varkappa \ln z\right).$$
(7.1)

Consequently the construction of the g-function, even under the one-cut assumption would significantly change. In practice this means that the EoT would have a different shape that depends on  $\rho, \varkappa$ . While conceptually there is no major difference, we found that there are practical and significant obstacles in obtaining an effective description of the g-function under these assumptions. A separate analysis is needed but it is deferred to a future publication.

A separate, long term question is whether the other families of rational solution described in Theorem 1.1 can be similarly framed in terms of semiclassical orthogonal polynomials. Irrespectively, the isomonodromic approach discussed in the Introduction should be available and hence the corresponding asymptotic analysis should be accessible. These are also issue that we defer to future investigations.

## A Airy parametrices

The complete construction of the approximation to the RHP of the main body of the paper requires the definition of a local solution to the final RHP which is known in the literature as "Airy Parametrix". While this is quite standard, it may be useful for the reader to find here its complete and self contained definition. The origin of these definitions can be traced back to [15] but here we refer to appendix A in [2]. We define

$$A_{0}(\zeta) = \begin{bmatrix} \frac{\mathrm{d}}{\mathrm{d}\zeta} \operatorname{Ai}(\zeta) & \mathrm{e}^{i\frac{\pi}{3}} \frac{\mathrm{d}}{\mathrm{d}\zeta} \operatorname{Ai}\left(\mathrm{e}^{-i\frac{2\pi}{3}}\zeta\right) \\ \operatorname{Ai}(\zeta) & \mathrm{e}^{i\frac{\pi}{3}} \operatorname{Ai}\left(\mathrm{e}^{-i\frac{2\pi}{3}}\zeta\right) \end{bmatrix}, \quad \zeta \in \mathbb{C}$$
(A.1)

where  $Ai(\zeta)$  is the Airy function, namely, a particular solution to Airy's equation

$$f(\zeta)'' = \zeta f(\zeta)$$

satisfying the following asymptotic behaviour as  $\zeta \to \infty$  in the sector  $-\pi < \arg \zeta < \pi$ 

$$\operatorname{Ai}(\zeta) = \frac{\zeta^{-1/4}}{2\sqrt{\pi}} e^{-\frac{2}{3}\zeta^{3/2}} \left( 1 - \frac{5}{48}\zeta^{-3/2} + \frac{385}{4608}\zeta^{-6/2} + O\left(\zeta^{-9/2}\right) \right).$$

We finally define the following piecewise-analytic matrix-valued function;

$$\mathbf{A}(\zeta) = \begin{cases} A_0(\zeta), & \arg \zeta \in (0, \frac{2\pi}{3}), \\ A_0(\zeta) \begin{bmatrix} 1 & 0\\ -1 & 1 \end{bmatrix}, & \arg \zeta \in (\frac{2\pi}{3}, \pi), \\ A_0(\zeta) \begin{bmatrix} 1 & -1\\ 0 & 1 \end{bmatrix}, & \arg \zeta \in (-\frac{2\pi}{3}, 0), \\ A_0(\zeta) \begin{bmatrix} 0 & -1\\ 1 & 1 \end{bmatrix}, & \arg \zeta \in (-\pi, -\frac{2\pi}{3}), \end{cases}$$
(A.2)

which solves the RHP with jumps for  $\arg \zeta = -\pi, -\frac{2\pi}{3}, 0, \frac{2\pi}{3}$  as depicted in Figure 10.

The definition (A.2) is crafted in such a way that the following asymptotic expansion holds in any direction

$$\mathbf{A}(\zeta) = \frac{\zeta^{\sigma_3/4}}{2\sqrt{\pi}} \begin{bmatrix} -1 & i \\ 1 & i \end{bmatrix} \left\{ I + \frac{1}{48\zeta^{3/2}} \begin{bmatrix} 1 & 6i \\ 6i & -1 \end{bmatrix} + \mathcal{O}\left(\zeta^{-6/2}\right) \right\} e^{-\frac{2}{3}\zeta^{3/2}\sigma_3}.$$
 (A.3)



Figure 10: The jump behaviour of the Airy Parametrix.

### A.1 The Airy parametrix for a three main-arc intersection

The basic building block discussed in the above Sec. A can be used to construct a more complicated local parametrix that needs to be used in a neighbourhood of a 3-arc intersection, like the case of the point z = b for the asymptotic in the EoT for Re (s) > 0, Sec 5.2.1.

We thus first state and solve a model RHP and then show how it proves of use to the case at hand.

Consider the jump matrices of the RHP 5.7 on the arcs that originate or terminate at z = b (see (5.55)): they are summarized in the left pane of Fig. 11, where we have denoted by  $\varphi_0, \varphi_1, \varphi_2$  the restrictions of  $\varphi$ to the sectors in  $\mathbb{D}_b$  defined as follows:  $\mathscr{S}_0$  bounded by the arcs  $\Gamma_b^1, \Gamma_b^{a_+}, \mathscr{S}_1$  bounded by the arcs  $\Gamma_b^{a_+}, \Gamma_{a_-}^{b_-}$ and  $\mathscr{S}_2$  bounded by the arcs  $\Gamma_{a_-}^b, \Gamma_b^1$ , respectively (see Fig. 7, left pane). They are related to each other as follows

$$\varphi_0(z) + \varphi_1(z) = 2\Omega_2, \quad z \in \Gamma_h^{a_+}, \tag{A.4}$$

$$\varphi_0(z) + \varphi_2(z) = 2\Omega_1, \quad z \in \Gamma_b^1, \tag{A.5}$$

$$\varphi_1(z) + \varphi_2(z) = 0, \quad z \in \Gamma_{a_-}^b. \tag{A.6}$$

Observing that  $\varphi_0(b) = -\Omega_1 - \Omega_2$  we can define the local conformal map  $\zeta(z)$  by the expression.

$$\frac{4}{3}\zeta^{\frac{3}{2}} = n\varphi_0(z) + n(\Omega_1 + \Omega_2), \quad z \in \mathscr{S}_0, \tag{A.7}$$

where the radical of  $\zeta$  is intended in the sense of principal determination (with the branch-cut extending along  $\zeta \in \mathbb{R}_{-}$ ). If we perform the analitic extension of this definition to the sectors  $\mathscr{S}_1, \mathscr{S}_2$  we obtain the following relations;

$$n\varphi_1(z) = -n\varphi_0(z) + 2\Omega_2 = -\frac{4}{3}\zeta^{\frac{3}{2}} - n\Omega_1 + n\Omega_2, \quad z \in \mathscr{S}_1$$
(A.8)

$$n\varphi_2(z) = -n\varphi_0(z) + 2\Omega_1 = -\frac{4}{3}\zeta^{\frac{3}{2}} - n\Omega_2 + n\Omega_1, \quad z \in \mathscr{S}_2.$$
(A.9)

With this definition of  $\zeta$ , the arc  $\Gamma_{a_{-}}^{b} \cap \mathbb{D}_{b}$  is mapped to the negative  $\zeta$  axis.

Moreover we have denoted by  $\widetilde{Q}(z)$  the analytic extension of Q(z) in the full disk  $\mathbb{D}_b$  from the upper part, so that

$$\widetilde{Q}(z) = \begin{cases} Q(z) & \left\{ z : \operatorname{Im} \zeta(z) > 0 \right\} \cap \mathbb{D}_b \\ e^{2i\pi\rho}Q(z) & \left\{ z : \operatorname{Im} \zeta(z) < 0 \right\} \cap \mathbb{D}_b. \end{cases}$$
(A.10)

With these notation in place and recalling the relation ship between  $\varkappa$  and  $e^{2i\pi\rho}$  (5.38), the jump matrices are as indicated in Fig. 11, left pane; note that we have also re-oriented some of the arcs; note that the jump matrix on the left ray is the result of the multiplication of the jumps on the two arcs  $\mathscr{L}^{(a_-,b)}_+, \mathscr{L}^{(b,a_+)}_+$ .



Figure 11: The local RHP 5.7 in  $\mathbb{D}_b$  (left pane) and its transformation to the  $\zeta$  plane.

With these preparations, the solution of the RHP with the jump matrices indicated in the right pane of Fig. 11 is written as follows in terms of the matrix  $\mathbf{A}(z)$  defined in (A.2):

$$\widetilde{\mathbf{P}}(\zeta) = \begin{cases} \mathbf{A}(\zeta) \mathrm{e}^{\left(\frac{2}{3}\zeta^{\frac{3}{2}} + \frac{\Omega_{1} - \Omega_{2}}{2}\right)\sigma_{3}} \left(\frac{\varkappa}{(\varkappa + 1)\widetilde{Q}(z)}\right)^{\frac{\sigma_{3}}{2}} \mathrm{e}^{\frac{i\pi\sigma_{3}}{2}} & \arg(\zeta) \in \left(\frac{\pi}{3}, \pi\right) \\ \mathbf{A}(\zeta) \mathrm{e}^{\left(\frac{2}{3}\zeta^{\frac{3}{2}} + \frac{\Omega_{2} - \Omega_{1}}{2}\right)\sigma_{3}} \left(\frac{1}{\varkappa(\varkappa + 1)\widetilde{Q}(z)}\right)^{\frac{\sigma_{3}}{2}} \mathrm{e}^{-\frac{i\pi\sigma_{3}}{2}} & \arg(\zeta) \in \left(-\pi, -\frac{\pi}{3}\right) \\ \mathbf{A}(\zeta) \mathrm{e}^{\left(\frac{2}{3}\zeta^{\frac{3}{2}} + \frac{\Omega_{2} + \Omega_{1}}{2}\right)\sigma_{3}} \left[\frac{0}{\frac{1}{\widetilde{Q}(z)}}\right] \left(\frac{\varkappa + 1}{\varkappa\widetilde{Q}(z)}\right)^{\frac{\sigma_{3}}{2}} \mathrm{e}^{\frac{-i\pi\sigma_{3}}{2}} & \arg(\zeta) \in \left(-\frac{\pi}{3}, \frac{\pi}{3}\right) \end{cases}$$
(A.11)

## References

- F. Balogh, M. Bertola, T. Bothner, "Hankel determinant approach to generalized Vorob'ev-Yablonski polynomials and their roots", *Constr. Approx.* 44 (2016), no. 3, 417–453.
- [2] M. Bertola, T. Bothner. "Zeros of large degree Vorob'ev-Yablonski polynomials via a Hankel determinant identity." Int. Math. Res. Not. IMRN, (19):9330–9399, 2015.
- [3] M. Bertola, B. Eynard, and J. Harnad. "Semiclassical orthogonal polynomials, matrix models and isomonodromic tau functions." Comm. Math. Phys., 263(2):401–437, 2006.
- [4] M. Bertola, A. Katsevich, A. Tovbis. "Singular value decomposition of a finite Hilbert transform defined on several intervals and the interior problem of tomography: the Riemann-Hilbert problem approach." *Comm. Pure Appl. Math.*, 69(3):407–477, March 2016.
- [5] M. Bertola and S. Y. Lee. "First colonization of a hard-edge in random matrix theory." Constr. Approx., 31(2):231–257, 2010.
- [6] M. Bertola and M. Y. Mo. "Commuting difference operators, spinor bundles and the asymptotics of orthogonal polynomials with respect to varying complex weights." Adv. Math., 220(1):154–218, 2009.
- [7] Pavel Bleher and Alexander Its. Double scaling limit in the random matrix model: the Riemann-Hilbert approach. *Comm. Pure Appl. Math.*, 56(4):433–516, 2003.
- [8] A. Borodin, P. J. Forrester, "Increasing subsequences and the hard-to-soft edge transition in matrix ensembles", J. Phys. A, 36 (2003), no. 12, 2963–2981.

- [9] R. J. Buckingham, P. D. Miller, "The sine-Gordon equation in the semiclassical limit: critical behavior near a separatrix", J. Anal. Math. 118 (2012), no. 2, 397–492.
- [10] R. Buckingham, P. Miller, "Large-degree asymptotics of rational Painlevé-II functions: noncritical behaviour", *Nonlinearity* 27 (2014), 2489-2577.
- [11] R. Buckingham, P. Miller, "Large-degree asymptotics of rational Painlevé-II functions: critical behaviour", Nonlinearity 28 (2015), 1539–1596.
- [12] R. Buckingham, P. Miller, "Large-degree asymptotics of rational Painlevé-IV solutions by the isomonodromy method", Constr. Approx. 56 (2022), no. 2, 233–443
- [13] P. A. Clarkson, C. Dunning, "Rational Solutions of the Fifth Painlevé Equation. Generalised Laguerre Polynomials", arXiv:2304:01579
- [14] P. Deift, Orthogonal polynomials and random matrices: A Riemann-Hilbert approach, Courant lecture notes, 1999.
- [15] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides and X. Zhou, Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory, *Commun. Pure Appl. Math.* **52** (1999), 1335-1425.
- [16] P. Deift, X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Ann. of Math., 137 (1993), 295-368.
- [17] A. Fokas, A. Its and A. Kitaev, Discrete Painlevé equations and their appearance in quantum gravity, Comm. Math. Phys. 142 (2) (1991), 313-344.
- [18] " Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est a points critiques fixes", Acta Math. 33, (1910), no. 1, 1–55.
- [19] F. D. Gakhov. "Boundary value problems", Dover Publications, Inc., New York, 1990. Translated from the Russian, Reprint of the 1966 translation.
- [20] S. P. Hastings, J. B. McLeod, "A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation", Arch. Rational Mech. Anal. 73 (1980), no. 1, 31?51.
- [21] Michio Jimbo, Tetsuji Miwa, and Kimio Ueno. "Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. I. General theory and  $\tau$ -function." <u>Phys. D</u>, 2(2):306–352, 1981.
- [22] Michio Jimbo and Tetsuji Miwa. "Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II." Phys. D, 2(3):407–448, 1981.
- [23] N. Joshi, K. Kajiwara, M. Mazzocco, "Generating function associated with the determinant formula for the solutions of the Painlevé II equation", Astérisque (2004), no. 297, 67–78.
- [24] K. Kajiwara, Y. Ohta, Determinant structure of the rational solutions for the Painlevé II equation, Journal of Mathematical Physics 37 (1996), 4693-4704.
- [25] A. V. Kitaev, C. K. Law, and J. B. McLeod. Rational solutions of the fifth Painlevé equation. Differential Integral Equations, 7(3-4):967–1000, 1994.
- [26] H. L. Krall and Orrin Frink. A new class of orthogonal polynomials: The Bessel polynomials. Trans. Amer. Math. Soc., 65:100–115, 1949.
- [27] F. Marcellán, I.A. Rocha, "On semiclassical linear functionals: integral representations", Proceedings of the Fourth International Symposium on Orthogonal Polynomials and their Applications (Evian-Les-Bains, 1992), 57, no. 1-2, 239–249 (1995).

- [28] F. Marcellán, I.A. Rocha, "Complex path integral representation for semiclassical linear functionals", J. Appr. Theory 94 (1998) 107–127.
- [29] P. Maroni, "Prolégomènes á l'étude des polynômes semiclassiques", Ann. Mat. Pura Appl. 149 (1987) 165–184.
- [30] T. Masuda, Y. Ohta, K. Kajiwara, "A determinant formula for a class of rational solutions of Painlevé V equation", Nagoya Math. J., 168 (2002), 1–25.
- [31] Masatoshi Noumi and Yasuhiko Yamada. Umemura polynomials for the Painlevé V equation. Phys. Lett. A, 247(1-2):65–69, 1998.
- [32] B. M. McCoy, C. Tracy, T. T. Wu, "Connection between the KdV equation and the two-dimensional Ising model", *Phys. Lett. A* 61 (1977), no. 5, 283?284.
- [33] B. M. McCoy, C. Tracy, T. T. Wu, "Painlevé functions of the third kind", J. Mathematical Phys. 18 (1977), no. 5, 1058?1092.
- [34] V. Yu. Novokshenov, A.A. Shchelkonogov, "Double scaling limit in the Painlevé IV equation and asymptotics of the Okamoto polynomials", *Amer. Math Soc. Transl. Ser. 2, 233*, Adv. Math. Sci, 66, (2014), 199–210.
- [35] K. Okamoto, "Studies on the Painlevé equations III, second and fourth Painlevé equations, PII and PIV", Math. Ann. 275 (1986) 222-254.
- [36] P. Painlevé, "Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme", Acta Math. 25 (1902), no. 1, 1–85.
- [37] J. Shohat, "A differential equation for orthogonal polynomials", Duke Math. J., 5 (1939), 401–417
- [38] Kurt Strebel. Quadratic differentials, volume 5 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1984.
- [39] C. Tracy, H. Widom, "Level-spacing distributions and the Airy kernel", Comm. Math. Phys., 159 (1994), no. 1, 151–174.
- [40] H. Umemura, "Special polynomials associated with the Painlevé equations I", Ann. Fac. Sci. Toulouse Math. (6) 29 (2020), no. 5, 1063–1089
- [41] M. Vanlessen. Strong asymptotics of Laguerre-type orthogonal polynomials and applications in random matrix theory. Constr. Approx., 25(2):125–175, 2007.
- [42] A. Vorob'ev, "On rational solutions of the second Painlevé equation", Diff. Eqns 1 (1965), 58-9 (in Russian).
- [43] A. Yablonskii, On rational solutions of the second Painlevé equation, Vesti Akad. Navuk. BSSR Ser. Fiz. Tkh. Nauk. 3 (1959), 30-5 (in Russian).