Epidemic outbreaks in structured host populations

Horst R. Thieme*

School of Mathematical and Statistical Sciences Arizona State University, Tempe, AZ 85287-1804, USA

November 14, 2024

Abstract

For a heterogeneous host population, the basic reproduction number of an infectious disease, \mathcal{R}_0 , is defined as the spectral radius of the next generation operator (NGO). The threshold properties of the basic reproduction number are typically established by imposing conditions that make \mathcal{R}_0 an eigenvalue of the NGO associated with a positive eigenvector and a positive eigenfunctional (eigenvector of the dual of the NGO). More general results can be obtained by imposing conditions that associate \mathcal{R}_0 just with a positive eigenfunctional. The next generation operator is conveniently expressed by a measure kernel or a Feller kernel which enables the use of analytic rather than functional analytic methods.

Keywords: force of infection, tight measure kernels, topologically irreducible kernels, dominated kernels

MSC Classification: 92D30, 92D25, 28A25, 45H05, 45M99, 47J05, 47N20

1 Introduction

The emergence of Covid-19 has reignited the interest in epidemic models of Kermack–McKendrick type. See [2, 3, 5, 6, 8, 10, 11, 12, 15, 16, 19, 21, 30, 33, 34, 35, 36, 40, 42, 45, 46] and the references therein. The host populations typically are heterogenous, and it is of interest how the spread of the infection is affected by the structure of the host population. It is one of the mathematical consequences of host heterogeneity that the basic reproduction number, \mathcal{R}_0 , is formulated in a functional analytic way, as the

^{*(}hthieme@asu.edu)

spectral radius of an appropriate positive bounded linear operator (called the "next generation operator", NGO) on the state space of the force of infection. In order to explore the threshold character of \mathcal{R}_0 , the existing literature typically uses that, under appropriate assumptions, the spectral radius is an eigenvalue of the next generation operator associated with a positive eigenvector and a positive eigenfunctional (eigenvector of the dual opeator). Persistence theory [44, 54] teaches the lesson that it can be sufficient if the spectral radius is just associated with a positive eigenfunctional [25, Thm.7] [43, App.2.6].

It is one of the aims of this article to show how the existence of an eigenfunctional plays out for epidemic models in structured host populations, in particular in getting away with less compactness assumptions and, with the exception of one fundamental theorem (Section 3.1), the use of real rather than functional analysis.

1.1 Scent of the model

The structure of the host population is described by a nonempty set Ω of structural characteristics of the hosts, x, also called traits, which do not change over time. One of the first population structures that have been considered appears to be associated with the spatial spread of the epidemic, and the trait of the host is spatial location. Some of the respective models have been formulated so generally that they apply to general structures [7, 47].

Let us assume that there is a function $s : \mathbb{R}_+ \times \Omega \to (0, 1]$ such that s(t, x) is the probability that a typical host with trait x that is susceptible at time 0 (the begin of the epidemic) will still be susceptible at time t > 0 [2]. In an epidemic scenario, where infected hosts do not become susceptible, $s(\cdot, x)$ is a decreasing function of t and s(0, x) = 1. Define

$$J(t,x) = -\ln s(t,x).$$
 (1.1)

Then $J(\cdot, x)$ is increasing, J(0, x) = 0, and

$$s(t,x) = e^{-J(t,x)}.$$
 (1.2)

J is called the *cumulative force of infection*, cFoI, up to time t. Under appropriate assumptions,

$$J(t,x) = \int_0^t I(r,x) dr,$$
 (1.3)

resulting in the differential equation

$$\partial_t s(t,x) = -s(t,x)I(t,x), \qquad (1.4)$$

with a non-negative function $I : \mathbb{R}_+ \times \Omega \to \mathbb{R}$. I(t, x) is called the *force* of infection, FoI, exerted on susceptible hosts of trait x at time t by all presently infected hosts. Cf. [4, 9, 22, 23, 38].

Some more equations will connect s and I (Section 5), which we do not present here because they are quite complex. In the spirit of Kendrick and McKermack [26, 27, 28], we will assume that the infectivity of an infected host does not only depend on its trait but also on its infection age (a special form of class age [22, 47], the time that has passed since the moment of being infected [2, 9, 23, 24, 32, 37, 38]). Differently from [2], we do not assume that there is a prescribed prehistory of the epidemic, but similarly to [7, 24, 41, 47] we assume the scenario that the disease is introduced into the population at some time t_0 which is normalized to $t_0 = 0$. This seems to be more appropriate for emerging diseases like Covid-19.

By its interpretation, it is suggestive that the cFoI J(t, x) is an increasing function of t. If is bounded, we can define the final size of the cFoI,

$$w(x) = \lim_{t \to \infty} J(t, x), \qquad x \in \Omega.$$
 (1.5)

By (1.2),

$$s_{\infty}(x) := \lim_{t \to \infty} s(t, x) = e^{-w(x)}, \qquad s \in \Omega.$$
(1.6)

1.2 Overview

The final size of the cFoI, w in (1.5) satisfies a Hammerstein equation with a measure kernel. See Section 2, where we also introduce the concept of a next generation operator (NG0) and of the basis reproduction number \mathcal{R}_0 as its spectral radius In Section 3, we give a preview of threshold results for \mathcal{R}_0 , which are discussed in Section 4. We also give the elementary proof of a fundamental threshold theorem. In Section 5, we derive integral equations of Volterra Hammerstein form for the FoI and for the cFoI. In Section 6, following [47], we show the existence of minimal solutions to these equations. Minimal solutions are unique by their nature, and we argue that they are the epidemiologically relevant solutions. In Section 7, again following [47], we derive Hammerstein equations with measure kernels for the final size of the cFoI and prove the existence of minimal solutions. In Section 8 and 9, we show threshold theorems for dominated and semi-separable measure kernels, and in Section 10 for Feller kernels in case that Ω is a metric space.

2 Some basic concepts

In this section, we get to see the equation for the final size of the cFoI, w in (1.5), and we introduce the concept of a next generation operator (NG0) and of the basis reproduction number \mathcal{R}_0 as its spectral radius.

2.1 The final size of the epidemic and measure kernels

Whereas the equation for the cumulative force of infection J is a quite complicated Volterra Hammerstein equation, the equation for its final size w boils down to a Hammerstein equation

$$w(x) - w^{\circ}(x) = \int_{\Omega} f(w(\xi))\kappa(d\xi, x) =: F(w)(x), \qquad x \in \Omega.$$
 (2.1)

Cf. [24, (31)]. While w is the final size of the cFoI exerted by all infected hosts, w° is the final size of the cFoI exerted by the hosts that were infected at the beginning of the epidemic. Here,

$$f(y) = 1 - e^{-y}, \qquad y \in \mathbb{R}_+.$$
 (2.2)

 Ω is a measurable space with a σ -algebra \mathcal{B} , e.g., an open or closed subset of \mathbb{R}^n . Further, $\kappa : \mathcal{B} \times \Omega \to \mathbb{R}_+$ is a measure kernel. Let us explain.

Let $\mathcal{M}(\Omega)$ be the vector space of real-valued measures on \mathcal{B} and $\mathcal{M}_+(\Omega)$ the order cone of nonnegative measures. See [52][54, Chap.33] for a population oriented introduction.

Further, let $M^b(\Omega)$ be the Banach space of bounded measurable functions with the supremum norm, $\|\cdot\|_{\infty}$, and $M^b_+(\Omega)$ be the closed order cone of nonnegative functions. The final sizes w and w° are elements of $M^b_+(\Omega)$.

Now, κ is a measure kernel [52][54, Sec.13.2] if

$$\kappa(\cdot, x) \in \mathcal{M}_+(\Omega), \quad x \in \Omega, \qquad \kappa(\omega, \cdot) \in M^b_+(\Omega), \quad \omega \in \mathcal{B}.$$
 (2.3)

Measure kernels are considered for a uniform treatment of the initial-value problem version of the model in [2] and of some special cases of [47, Sec.3]. Notice that both frameworks allow the consideration of continuous and discrete population structure. We will show (Section 7) that the final size wdefined by (1.5) is the minimal solution of (2.1), given as the pointwise limit $w = \lim_{n\to\infty} w_n$ of the recursion

$$w_n = F(w_{n-1}) + w^{\circ}, \quad n \in \mathbb{N}, \qquad w_0 = w^{\circ}.$$
 (2.4)

By induction, (w_n) is a increasing sequence of functions in $M^b_+(\Omega)$. Cf. [47, Sec.3].

2.2 Next generation operator and basic reproduction number

The event of no infection is represented by the zero function. The linearization of F in (2.1) at the zero function is the positive bounded linear operator $K: M^b(\Omega) \to M^b(\Omega)$ given by

$$(Kg)(x) = \int_{\Omega} g(\xi)\kappa(d\xi, x), \qquad x \in \Omega, \quad g \in M^{b}(\Omega).$$
 (2.5)

More precisely, K is the order-derivative of F [52][54, Sec.14.2]: By the mean value theorem applied to f in (2.2), since f' is decreasing,

$$f(y) \ge y f'(y) = y e^{-y}, \qquad y \in \mathbb{R}_+, \tag{2.6}$$

$$f'(\|g\|_{\infty}) Kg \le F(g) \le Kg, \quad g \in M^b_+(\Omega), \qquad f'(0) = 1.$$
 (2.7)

Remark 2.1. Even more precisely, by (2.6),

$$F(g)(x) \ge \int_{\Omega} f'(g(\xi)) g(\xi) \kappa(d\xi, x), \qquad x \in \Omega, \quad g \in M^b_+(\Omega).$$

By (1.6), if w is the final size of the cFoI,

$$F(w)(x) \ge \int_{\Omega} w(\xi) s_{\infty}(\xi) \kappa(d\xi, x), \qquad x \in \Omega.$$

In our epidemiological context, K is called the *next generation operator* (with generation to be understood in an epidemiological sense).

The basic reproduction number \mathcal{R}_0 is defined as the spectral radius of K, which is given by the Gelfand formula

$$\mathcal{R}_0 = \mathbf{r}(K) = \inf_{n \in \mathbb{N}} \|K^n\|^{1/n} = \lim_{n \to \infty} \|K^n\|^{1/n}.$$
 (2.8)

Here K^n is the *n*-fold composition (iterate) of K with itself and $||K^n||$ its operator norm. See [54, A.3.2] for a proof of the last equality that also holds if K is not linear but only homogeneous. We also call \mathcal{R}_0 the spectral radius of the measure kernel κ ,

$$\mathcal{R}_0 = \mathbf{r}(\kappa). \tag{2.9}$$

 \mathcal{R}_0 will turn out to be an epidemic threshold parameter for the type of models we will consider. Cum grano salis, epidemic outbreaks occur if $\mathcal{R}_0 > 1$, and do not occur if $\mathcal{R}_0 < 1$.

3 Preview of threshold results

In the following, we will give a glimpse of the epidemic threshold character of the basic reproduction number, \mathcal{R}_0 .

3.1 A fundamental result from positive operator theory

It follows from the definition of a measure kernel, (2.3), that $u: \Omega \to \mathbb{R}_+$,

$$u(x) = \kappa(\Omega, x), \quad x \in \Omega, \tag{3.1}$$

is an element of $M^b_+(\Omega)$. The linear operator K on $M^b(\Omega)$ given by (2.5) is uniformly *u*-bounded, i.e.,

$$Kg \le ||g||_{\infty} u, \qquad g \in M^b_+(\Omega).$$
 (3.2)

Further, the constant function $u_1(x) = 1$ is an interior point of $M^b_+(\Omega)$,

$$|g(x)| \le ||g||_{\infty} u_1(x), \qquad x \in \Omega, \quad g \in M^b(\Omega).$$
(3.3)

After these preparations, the following theorem is a special case of more abstract ones for positive bounded linear operators on ordered Banach spaces with a solid normal cone [25, Thm.7] [31] [43, App.2.6][54, Cor.11.17].

Theorem 3.1. If $\mathcal{R}_0 > 0$, there exists a bounded linear positive eigenfunctional θ : $M^b(\Omega) \to \mathbb{R}$ of K (and of κ) such that $\theta(Kg) = \mathcal{R}_0 g$ for all $g \in M^b(\Omega), \theta(g) > 0$ if $g \in M^b_+(\Omega)$ and $g \ge \delta u$ for some $\delta > 0$.

The inequality $\theta(u) > 0$ follows from (3.2). Otherwise, since θ is increasing and linear,

$$\mathcal{R}_0\theta(g) = \theta(Kg) \le \|g\|_{\infty} \,\theta(u) = 0, \qquad g \in M^b_+(\Omega),$$

and θ would be the zero functional.

The functional θ can rightfully be called also an eigenfunctional of the measure kernel κ because it is characterized by

$$\theta(\kappa(\omega, \cdot)) = \mathcal{R}_0 \,\theta(\chi_\omega), \qquad \omega \in \mathcal{B}, \tag{3.4}$$

where χ_{ω} is the indicator or characteristic function of the set ω , $\chi_{\omega}(\xi) = 1$ if $\xi \in \omega$ and $\chi_{\omega}(\xi) = 0$ if $\xi \in \Omega \setminus \omega$.

Generalizations of Theorem 3.1 to increasing bounded homogeneous operators can be found in [50] [54, Chap.11].

3.2 A general but weak threshold result

Persistence theory teaches that eigenfunctionals of the NGO are as important tools as eigenvectors [44, 54].

Theorem 3.2. Let $w \in M^b_+(\Omega)$ be the minimal solution of $w = F(w) + w^\circ$ which is uniquely determined by $w^\circ \in M^b_+(\Omega)$.

- (a) Let $\mathcal{R}_0 < 1$. Then there exists a constant c > 0 (independent of w and w°) such that $||w||_{\infty} \leq c ||w^\circ||_{\infty}$.
- (b) Let $\mathcal{R}_0 > 1$ and θ be the eigenfunctional of K from Theorem 3.1. If $\theta(w) > 0$ (in particular if $\theta(w^\circ) > 0$), then $||w w^\circ||_{\infty} \ge \ln \mathcal{R}_0$ and $\inf_{\Omega} s_{\infty} \le 1/\mathcal{R}_0$ for s_{∞} in (1.6).

Notice that there are no irreducibility or compactness assumptions for K. As a trade-off, there is not so much information as in other threshold results (see [2, 24] and the Sections (3.3) to (3.5). But one definitely sees that it makes a difference whether $\mathcal{R}_0 < 1$ or $\mathcal{R}_0 > 1$.

This result is fundamental for our approach and the proof is easy (once we take Theorem 3.1 for granted) so that we give it right here.

Proof. (a) By (2.7) and (2.4),

$$w_{n+1} \le Kw_n + w^\circ, \quad n \in \mathbb{N}.$$

By induction,

$$w_n \le \sum_{j=0}^n K^j w^\circ, \quad n \in \mathbb{N}.$$

Since $\mathcal{R}_0 < 1$, $K_{\infty} = \sum_{j=0}^{\infty} K^j$ exists with the series converging even in operator norm and is a bounded linear operator on $M^b(\Omega)$,

$$w_n(x) \le (K_\infty w^\circ)(x), \quad n \in \mathbb{N}, \quad x \in \Omega.$$

Taking pointwise limits, $w = \lim_{n \to \infty} w_n$,

$$w \le K_{\infty} w^{\circ}$$
 and $||w||_{\infty} \le ||K_{\infty}|| ||w^{\circ}||_{\infty}$.

(b) Assume $\theta(w) > 0$. Let $\tilde{w} = w - w^{\circ}$. By (2.7),

$$\tilde{w} = F(w) \ge f'(\|w\|_{\infty})K(w).$$

Since θ is an eigenfunctional of K associated with \mathcal{R}_0 and θ is increasing,

$$\theta(\tilde{w}) \ge f'(\|w\|_{\infty})\theta K(w) = e^{-\|w\|_{\infty}} \mathcal{R}_0 \theta(w) > 0.$$

Since F is increasing, again by (2.7),

$$\tilde{w} \ge F(\tilde{w}) \ge f'(\|\tilde{w}\|_{\infty})K(\tilde{w})$$

and, by applying θ ,

$$\theta(\tilde{w}) \ge f'(\|\tilde{w}\|_{\infty})\mathcal{R}_0\,\theta(\tilde{w}).$$

Since $\theta(\tilde{w}) > 0$, $f'(\|\tilde{w}\|_{\infty}) \le 1/\mathcal{R}_0$ and $\|\tilde{w}\|_{\infty} \ge \ln \mathcal{R}_0$. Since θ is increasing (linear and nonnegative), $\theta(w) \ge \theta(w^\circ)$.

Corollary 3.3. If $\mathcal{R}_0 > 1$, $\tilde{w} \in M^b_+(\Omega)$, $\tilde{w} = F(\tilde{w})$ and $\theta(\tilde{w}) > 0$, then $\|\tilde{w}\|_{\infty} \ge \ln \mathcal{R}_0$.

The next observation seems to be mainly the same as the one in [2, Thm.5.6] that the basic reproduction number associated with the situation after the epidemic outbreak should not exceed one.

Remark 3.4. Let the final sizes of the cumulative forces of infection be described by the minimal solution of $w = F(w) + w^{\circ}$.

Then $s_{\infty}(x) = e^{-w(x)}$ is the probability that a host with trait x that is susceptible at the beginning is still susceptible at the end of the epidemic, (1.6). Define the measure kernel κ_{∞} by

$$\kappa_{\infty}(\omega, x) = \int_{\omega} s_{\infty}(\xi) \kappa(d\xi, x), \qquad \omega \in \mathcal{B}, \quad x \in \Omega.$$

Let \mathcal{R}_0 be the spectral radius of κ , (2.9), and \mathcal{R}_∞ be the spectral radius of κ_∞ . Then, if $\mathcal{R}_0 > 0$ and $\theta(w^\circ) > 0$, we have $\mathcal{R}_\infty \leq 1$.

See Remark 7.3 for more details. In [2, Thm.5.6], the strict inequality $\mathcal{R}_{\infty} < 1$ is obtained because irreducibility of the kernel κ is assumed.

Proof. We can assume that $\mathcal{R}_{\infty} > 0$, which is the spectral radius of the operator induced by κ_{∞} analogously to (2.5), and let θ_{∞} be the respective eigenfunctional of that operator associated with \mathcal{R}_{∞} via Theorem 3.1 and characterized by (3.4) with κ_{∞} replacing κ . Since s_{∞} is bounded away from 0 by (1.6), $\theta_{\infty}(w) \ge \theta_{\infty}(w^{\circ}) > 0$. By Remark 2.1,

$$w(x) \ge \int_{\Omega} w(\xi) \kappa_{\infty}(d\xi, x) + w^{\circ}.$$

We apply the linear positive functional θ_{∞} to this equation and obtain that $\mathcal{R}_{\infty} \leq 1$.

In the following, we explore stronger assumptions and stronger results in the cases that the kernel κ is dominated or even semi-separable or that Ω is a metric space and κ is a tight Feller kernel that is topologically irreducible [52][54, Chap.13].

3.3 Dominated measure kernels

A measure kernel κ is called *dominated* by a measure $0 \neq \nu \in \mathcal{M}_+(\Omega)$ if

$$\kappa(\omega, x) \le \nu(\omega), \qquad \omega \in \mathcal{B}, \quad x \in \Omega.$$
 (3.5)

The model we will consider in Sections 5 to 7 typically involves a dominated measure kernel.

Theorem 3.5. Let the measure kernel κ be dominated by a measure ν and $\mathcal{R}_0 > 1$. Let $w^{\circ} \in M^b_+(\Omega)$ and w be the minimal solution to $w = F(w) + w^{\circ}$.

Then there exists some solution $\tilde{w} = F(\tilde{w})$ in $M^b_+(\Omega)$ with the following properties:

- (a) $\int_{\Omega} (f \circ \tilde{w}) d\nu \ge \ln \mathcal{R}_0 \text{ and } \|\tilde{w}\|_{\infty} \ge \ln \mathcal{R}_0.$
- (b) If $\theta(w^{\circ}) > 0$, then $w w^{\circ} \ge \tilde{w}$ and $\ln \mathcal{R}_0 \le \int_{\Omega} (1 s_{\infty}) d\nu$.

The last inequality follows from (a) and the first part of (b) and (1.6).

Choosing $w^{\circ} \equiv 1$ on Ω in the previous theorem such that $\theta(w^{\circ}) > 0$, we obtain the following fixed point result (notice the scarcity of assumptions).

Corollary 3.6. Let the measure kernel κ be dominated by a measure ν and $\mathcal{R}_0 > 1$. Then there exists some solution $\tilde{w} = F(\tilde{w})$ in $M^b_+(\Omega)$ with the following properties:

 $\int_{\Omega} (f \circ \tilde{w}) d\nu \ge \ln \mathcal{R}_0 \text{ and } \|\tilde{w}\|_{\infty} \ge \ln \mathcal{R}_0.$

For the proofs see Section 8.

3.4 Semi-separable measure kernels

In Theorem 3.5, the nonzero fixed point \tilde{w} may depend on w° . To remove this dependence, we consider the following concept [24].

A measure kernel κ is called *semi-separable* if there are some nonzero function $k_0 \in M^b_+(\Omega)$ and some nonzero measure $\nu \in \mathcal{M}_+(\Omega)$ and some number $\delta \in (0, 1]$ such that

$$\delta\nu(\omega)k_0(x) \le \kappa(\omega, x) \le \nu(\omega)k_0(x), \qquad x \in \Omega, \quad \omega \in \mathcal{B}.$$
(3.6)

Notice that the function k_0 is not assumed to be strictly positive on Ω and no positivity assumption is made for ν except that it is not the zero measure. Assuming strict positivity of k_0 would be an easy way, though, to guarantee that the infection reaches hosts of all traits, in other words, the occurrence of a pandemic in case that $\mathcal{R}_0 > 1$. Cf. [24]. Notice that, if κ is semi-separable, it is dominated by the measure $||k_0||_{\infty} \nu$.

Theorem 3.7. Let κ be a semi-separable measure kernel. Then $\mathcal{R}_0 > 0$ if and only if

$$\int_{\Omega} k_0 \, d\nu > 0.$$

Let w (and w°) be the final sizes of the cumulative (initial) forces of infection, i.e., w is the minimal solution to $w = F(w) + w^{\circ}$.

- (a) If $w^{\circ} \in M^b_+(\Omega)$ and $\int_{\Omega} w^{\circ} d\nu = 0$, then $w = w_0$.
- (b) Assume that $\mathcal{R}_0 > 1$. Then there exists a unique non-zero solution $\tilde{w} \in M^b_+(\Omega)$ to $\tilde{w} = F(\tilde{w})$.

Further, $\int_{\Omega} (f \circ \tilde{w}) d\nu \geq \ln \mathcal{R}_0 / \|k_0\|_{\infty}$ and $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$. Finally, if $\int_{\Omega} w^\circ d\nu > 0$, we have $w - w^\circ \geq \tilde{w}$.

Notice that, in this theorem, for $\mathcal{R}_0 > 1$, an epidemic outbreak occurs if and only if $\int_{\Omega} w^{\circ} d\nu > 0$ for the final size of the cumulative initial force of infection, w° . The next theorem explores how the threshold property of \mathcal{R}_0 plays out for the final size of the epidemic in terms of the cFoI if the number of initial infectives is very small.

Theorem 3.8. Let κ be a semi-separable measure kernel. Let $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ be a sequence in $M^b_+(\Omega)$ and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω .

For each $\ell \in \mathbb{N}$, let $w_{\ell} \in M^b_+(\Omega)$ be the minimal solution of

$$w_\ell = F(w_\ell) + w_\ell^\circ.$$

- (a) If $\mathcal{R}_0 \leq 1$, then $w_\ell w_\ell^\circ \to 0$ as $\ell \to \infty$, uniformly on Ω .
- (b) Let $\mathcal{R}_0 > 1$ and $\int_{\Omega} w_\ell^\circ d\nu > 0$ for all $\ell \in \mathbb{N}$ and \tilde{w} be the unique nonzero solution to $\tilde{w} = F(\tilde{w})$ from Theorem 3.7 (b).

Then, as $\ell \to \infty$, $w_{\ell} \to \tilde{w}$ pointwise on Ω and $w_{\ell} - w_{\ell}^{\circ} \to \tilde{w}$ uniformly on Ω .

For the proofs see Section 9. We mention that, on the way from the general but less informative result in Theorem 3.2 and the detailed results in this section, there are intermediate results that are of their own interest, but will be presented later (Section 8).

3.5 Metric spaces of traits and Feller kernels

We now assume that Ω is a metric space and the σ -algebra \mathcal{B} is the one generated by the open subsets of Ω . Then, the vector space of bounded continuous real-valued functions on Ω , $C^b(\Omega)$, is a closed subspace of $M^b(\Omega)$ with the supremum norm.

Definition 3.9 ([1, Chap.19.3][53][54, Chap.13]). A measure kernel κ is called a *Feller kernel* if the operator K on $M^b(\Omega)$ induced by (2.5) maps $C^b(\Omega)$ into itself.

A Feller kernel κ is called *topologically irreducible* if for any nonempty open strict subset U of Ω there exist some $x \in \Omega \setminus U$ such that $\kappa(U, x) > 0$.

A function $g \in M^b_+(\Omega)$ is called *topologically positive* if $\inf_U g > 0$ for some nonempty open subset U of Ω .

Theorem 3.10. Let κ be a topologically irreducible Feller kernel and $w^{\circ} \in M^b_+(\Omega)$ be topologically positive and $w \in M^b_+(\Omega)$ be the minimal solution to $w = F(w) + w^{\circ}$. Then $w(x) - w^{\circ}(x) > 0$ for all $x \in \Omega$.

So there is potential for a pandemic. We mention that topological irreducibility of κ is necessary for this result to hold (Remark 10.7).

3.5.1 Tight Feller kernels

Definition 3.11 ([52][54, Chap.13]). A measure $\mu \in \mathcal{M}_+(\Omega)$ is called *tight* if for any $\epsilon > 0$ there exists a compact subset W of Ω such that $\mu(\Omega \setminus W) < \epsilon$.

A Feller kernel κ is called *tight* if for any $\epsilon > 0$ there is a compact subset W of Ω such that

$$\kappa(\Omega \setminus W, x) < \epsilon, \qquad x \in \Omega. \tag{3.7}$$

A Feller kernel κ is called *quasi-tight* if $\kappa = \kappa_1 + \kappa_2$ where the κ_j are Feller kernels, κ_1 is tight, $\mathbf{r}(\kappa) > \mathbf{r}(\kappa_2)$, and for any $x \in \Omega$, there exists a separable subset ω of Ω such that $\kappa_2(x, \Omega \setminus \omega) = 0$.

If κ is a tight or quasi-tight Feller kernel, the eigenfunctional θ in Theorem 3.1 is given by a measure. See [52][54, Thm.13.39] and [54, Thm.13.42].

Theorem 3.12. Let κ be a quasi-tight Feller kernel and $\mathcal{R}_0 > 0$. Then there exists a eigenmeasure μ of κ associated with \mathcal{R}_0 ,

$$\mathcal{R}_0\mu(\omega) = \int_\Omega \kappa(\omega, x)\mu(dx), \qquad \omega \in \mathcal{B}.$$

If, in addition, κ is a topologically irreducible Feller kernel, $\int_{\Omega} g \, d\mu > 0$ for every topologically positive $g \in M^b_+(\Omega)$. If κ is a tight Feller kernel, μ is tight. Now we are in the situation that the eigenfunctional θ used in Theorem 3.2 (b) is given by a measure,

$$\theta(g) = \int_{\Omega} g(x)\mu(dx), \qquad g \in M^b(\Omega)$$

In particular, θ is continuous with respect to pointwise convergence of increasing sequences of functions. Theorem 3.2 in conjunction with Theorem 3.12 and 3.10 imply the following result.

Theorem 3.13. Let κ be a quasi-tight Feller kernel that is topologically irreducible and $\mathcal{R}_0 > 1$. Then we have a pandemic situation:

Let $w^{\circ} \in M^{b}_{+}(\Omega)$ be topologically positive and $w \in M^{b}_{+}(\Omega)$ be the minimal solution of the final size equation $w = F(w) + w^{\circ}$.

Then, $w - w^{\circ}$ is strictly positive on Ω and $||w - w^{\circ}||_{\infty} \ge \ln \mathcal{R}_0$ and $\inf_{\Omega} s_{\infty} \le 1/\mathcal{R}_0$.

3.5.2 Strong Feller kernels

Again, the last result shows the threshold property of \mathcal{R}_0 , though not in a very strong sense. To compare the final size solutions to nonzero fixed points of $\tilde{w} = F(\tilde{w})$, we strengthen the Feller property.

Definition 3.14. A measure kernel κ is called a *strong Feller kernel* if $\kappa(\omega, \cdot)$ is continuous on Ω for all $\omega \in \mathcal{B}$.

A dominated Feller kernel (Section 3.3) is a strong Feller kernel (Proposition 10.11). See Example 4.2 for a measure kernel that under weak assumptions is a Feller kernel though not a strong Feller kernel, but is a strong Feller kernel under stronger assumptions. As we will show (Lemma 10.10), the maps K and F associated with a strong Feller kernel map $M^b_+(\Omega)$ into $C^b_+(\Omega)$.

Theorem 3.15. Let κ be a strong Feller kernel that is tight and topologically irreducible and let $\mathcal{R}_0 > 1$. Let $w^{\circ} \in M^b_+(\Omega)$ be topologically positive.

Let $w \in M^b_+(\Omega)$ be the minimal solution to $w = F(w) + w^\circ$.

Then there exists some $\tilde{w} \in C^b_+(\Omega)$ such that \tilde{w} strictly positive on Ω , $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$ and $w - w^{\circ} \geq \tilde{w} = F(\tilde{w})$.

For the proof of this and related results see Theorem 10.12 and its corollaries.

3.5.3 Comparability kernels

In the last result, the fixed point \tilde{w} depends on w° . To enforce independence, we introduce the following concept (Section 10.5). Cf. [48, (K-2)].

Definition 3.16. A Feller kernel κ is called a *comparability kernel* if for any continuous function $g: \Omega \to (0, \infty)$ there exists some $\delta > 0$ such that

$$\int_{\Omega} g(\xi)\kappa(d\xi, x) \ge \delta \kappa(\Omega, x), \qquad x \in \Omega.$$
(3.8)

Sufficient conditions and examples will be presented in Section 10.5.1.

The next two theorems explores how the threshold property of \mathcal{R}_0 plays out for the final size of the epidemic in terms of the cFoI if the number of initial infectives is very small. The results parallel those of Theorem 3.8, but even if Ω is a metric space the assumptions are different. See the end of the discussion in Section 4.

Theorem 3.17. Let κ be a strong Feller kernel that is topologically irreducible and tight. Assume that κ is a comparability kernel and $\mathcal{R}_0 > 1$.

- (a) Then there exists a unique nonzero solution $\tilde{w} \in C^b_+(\Omega)$ to the equation $\tilde{w} = F(\tilde{w})$; further $\|\tilde{w}\| \ge \ln \mathcal{R}_0$.
- (b) If (w^o_ℓ)_{ℓ∈ℕ} is a decreasing sequence of topologically positive functions in M^b₊(Ω) and w^o_ℓ → 0 as ℓ → ∞ uniformly on all compact subsets of Ω, then w_ℓ - w^o_ℓ → w̃ uniformly on Ω for the minimal solutions w_ℓ of w_ℓ = F(w_ℓ) + w^o_ℓ.

Theorem 3.18. Let κ be a Feller kernel that is topologically irreducible. Assume that κ is a comparability kernel and $\mathcal{R}_0 \leq 1$.

- (a) Then there exists no solution $\tilde{w} \in \dot{C}^b_+(\Omega)$ to the equation $\tilde{w} = F(\tilde{w})$.
- (b) If κ is a strong Feller kernel and $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ is a sequence in $M^b_+(\Omega)$ and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω , then $w_{\ell} \to 0$ as $\ell \to \infty$ pointwise for the minimal solutions w_{ℓ} of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

For the proofs of these results see Section 10.5.

4 Discussion

The results in this paper are based on finding a positive eigenfunctional of a bounded linear positive operator associated with its spectral radius (Theorem 3.1). The existing literature mostly attempts to find a positive eigenvector in addition, [2, 24], e.g. In the context of measure kernels, one can take that route, too, if one imposes a uniform Feller property.

A measure kernel κ is called a *uniform Feller kernel* [54, Sec.13.6] if

$$\sup_{\omega \in \mathcal{B}} \left| \kappa(\omega, x) - \kappa(\omega, x_0) \right| \to 0, \qquad x \to x_0 \in \Omega.$$
(4.1)

If κ is given in the form

$$\kappa(\omega, x) = \int_{\omega} k(x, \xi) S_0(d\xi), \qquad \omega \in \mathcal{B}, \quad x \in \Omega,$$
(4.2)

as it is from Section 5 to Section 7 following [2], (4.1) is equivalent to

$$\int_{\Omega} \left| k(x,\xi) - k(x_0,\xi) \right| S_0(d\xi) \longrightarrow 0, \qquad x \to x_0.$$

Cf. hypothesis H_{A_1} in [2, Sec.5]. See also assumption (41) in [24].

General measure kernels appear in the final size equation for the epidemic model in [47, Sec.3]. If κ is a uniform Feller kernel, existence of a (strictly positive) eigenvector of K follows from [53], [54, Thm.13.52] and [54, Thm.13.58].

Theorem 4.1. Let κ be a tight uniform Feller kernel and $\mathcal{R}_0 = \mathbf{r}(\kappa) > 0$. Then there exists an eigenvector $v \in C^b_+(\Omega)$ such that $\mathcal{R}_0 v = Kv$. If κ is topologically irreducible in addition, v is strictly positive on Ω .

A somewhat more general result is proved in [53][54, Thm.13.58]. If κ is given by (4.2), we could neither find a strong Feller kernel that is not a uniform Feller kernel nor could we prove that any strong Feller kernel is a uniform Feller kernel. If we do not insist on an epidemiologic interpretation, we can consider the following measure kernel.

Example 4.2. Let $\Omega = \mathbb{R}$ with the standard topology induced by the absolute value. Consider

$$\kappa(\omega, x) = \int_{\mathbb{R}} \chi_{\omega}(\xi \phi(x)) k_0(\xi) d\xi, \qquad \omega \in \mathcal{B}, \quad x \in \mathbb{R},$$

where $k_0 : \mathbb{R} \to \mathbb{R}_+$ is finitely integrable and $\phi : \mathbb{R} \to (0, \infty)$ is continuous.

Here, χ_{ω} is the characteristic or indicator function of the set ω , $\chi_{\omega}(x) = 1$ if $x \in \omega$ and $\chi_{\omega}(x) = 0$ if $x \in \Omega \setminus \omega$. By Lebesgue's theorem of dominated convergence, κ is a Feller kernel. After a substitution,

$$\kappa(\omega, x) = \int_{\omega} k_0(\eta/\phi(x)) (1/\phi(x)) d\eta, \qquad \omega \in \mathcal{B}, \quad x \in \mathbb{R}.$$

Assume that k_0 is lower semicontinuous. Let $x_n \to x$ as $n \to \infty$. By Fatou's lemma,

$$\liminf_{n \to \infty} \kappa(\omega, x_n) \ge \int_{\omega} \liminf_{n \to \infty} k_0 \left(\eta / \phi(x_n) \right) \left(1 / \phi(x_n) \right) d\eta.$$

Since k_0 is lower semicontinuous and ϕ is continuous,

$$\liminf_{n \to \infty} k_0(\eta/\phi(x_n)) (1/\phi(x_n)) \ge k_0(\eta/\phi(x))(1/\phi(x)), \qquad \eta \in \mathbb{R}.$$

See [54, L.A.49] which also implies that $\kappa(\omega, \cdot)$ is lower semicontinuous. By Proposition 10.9, κ is a strong Feller kernel.

 κ is a uniform Feller kernel if and only if, for any sequence $x_n \to x$ $(n \to \infty)$,

$$\int_{\Omega} \left| k_0 \big(\eta / \phi(x_n) \big) \left(1 / \phi(x_n) \right) - k_0 \big(\eta / \phi(x) \big) \left(1 / \phi(x) \right) \right| d\eta \to 0, \qquad n \to \infty.$$

We cannot present a lower semicontinuous $k_0 \in L^1(\mathbb{R})$ that does not satisfy this condition, but if one would like to come up with sufficient conditions they would presumably entail Lebesgue's theorem of dominated convergence involving a.e. continuity of k_0 and some domination condition stronger than just k_0 being finitely integrable.

Our epidemic model considers density-dependent alias mass action incidence as do the models in [2, 24]. See (1.4). More generally, we could consider

$$\partial_t s(t, x) = -\phi(s(t, x))I(t, x), \qquad s(0, x) = 1.$$
 (4.3)

Here $\phi : \mathbb{R}_+ \to \mathbb{R}_+$ is continuous and strictly increasing, positive on $(0, \infty)$, $\phi(0) = 0$, $\phi(1) = 1$ and $\int_0^1 (1/\phi(r)) dr = \infty$. Then

$$1 - s(t, x) = f(J(t, x)),$$

where f is an increasing concave function with strictly decreasing derivative, f(0) = 0, f'(0) = 1. Cf. [47, (5)(6)].

Then the result $||w - w^{\circ}||_{\infty} \ge \ln \mathcal{R}_0 > 0$ in Theorem 3.2 and other theorems has the more general form $f'(||w - w^{\circ}||_{\infty}) \le 1/\mathcal{R}_0 < 1$. See the proof of Theorem 3.2 (b).

The generalization

$$\partial_t s(t, x) = -s(t, x) \psi(I(t, x)), \qquad s(0, x) = 1,$$
(4.4)

with a continuous increasing $\psi : \mathbb{R}_+ \to \mathbb{R}_+$ seems to be much more difficult to handle. For piecewise linear ψ see [35], and for general incidences (and no host population structure) see [20].

There are two different though not disjoint sets of assumptions under which the threshold properties of \mathcal{R}_0 for epidemics with small numbers of initial infectives can be rather completely described: if κ is a semi-separable measure kernel (Theorem 3.8) or if Ω is a metric space and κ is a strong Feller kernel that is tight and a topologically irreducible comparability kernel (Theorem 3.17 and 3.18).

On the one hand, every semi-separable measure kernel is a comparability kernel (Proposition 10.22).

On the other hand, a semi-separable Feller kernel is tight if and only if the measure ν in (3.6) is tight. Moreover, a semi-separable Feller kernel is topologically irreducible (Definition 3.9) only if $\nu(U) > 0$ for the measure ν in (3.6) and for every non-empty open subset U of Ω .

5 The epidemic model

We consider a host population which is structured by characteristic traits $x \in \Omega$, where Ω is a measurable space with σ -algebra \mathcal{B} . The model starts at time t = 0. Our derivation follows [2], but we consider an epidemic which is triggered by initial infectives.

5.1 Susceptible hosts, incidence and force of infection

Let I(t, x) be the force of infection that affects susceptible hosts with trait $x \in \Omega$ at time $t \ge 0$. Let us consider a typical host with trait x and s(t, x) be the probability that this host is still susceptible at time t provided it was susceptible at time 0 [2]. Then

$$s(t,x) = e^{-J(t,x)}, \qquad J(t,x) = \int_0^t I(r,x)dr, \qquad t \ge 0, \quad x \in \Omega,$$
 (5.1)

provided that the integral makes sense. J(t, x) is the cumulative force of infection affecting susceptible hosts with trait x, accumulated from time 0

to time t. J is certainly well-defined if $I(\cdot, x)$ is continuous and, by the fundamental theorem of calculus and the chain rule,

$$\partial_t s(t,x) = -s(t,x)I(t,x). \tag{5.2}$$

J is also well-defined if $I(\cdot, x)$ is finitely integrable on bounded subintervals of \mathbb{R}_+ . Then $s(\cdot, x)$ is absolutely continuous and the last differential equation holds for a.a. $t \ge 0$ [39, L.8.1]. In other words,

$$1 - s(t, x) = \int_0^t s(r, x) I(r, x) dr, \qquad t \ge 0, \quad x \in \Omega.$$
 (5.3)

Following [2, 3], we assume that the structural distribution of susceptible hosts at time 0 is described by a non-negative measure $S_0 : \mathcal{B} \to \mathbb{R}_+$. Since the size of the susceptible population is finite, $S_0(\Omega) < \infty$. The structural distribution of susceptible hosts at time $t \ge 0$ is given by

$$S(t,\omega) = \int_{\omega} s(t,x) S_0(dx), \qquad (5.4)$$

where $S(t, \omega)$ is the number of susceptible hosts with trait in the set $\omega \in \mathcal{B}$ at time $t \geq 0$.

5.2 Incidence

We redo the modeling steps because differently from [2] we do not consider an epidemic with prehistory but an emerging epidemic, for instance if a pathogen overcomes a species barrier.

 $S_0(\omega) - S(t,\omega)$ is the number of infected host with trait in $\omega \in \mathcal{B}$ at time t. By (5.4), (5.1), (5.3) and Tonelli's theorem, changing the order of integration,

$$S_0(\omega) - S(t,\omega) = \int_{\Omega} (1 - s(t,x)) S_0(dx) = \int_0^t B(r,\omega) dr, \quad (5.5)$$

$$B(t,\omega) = \int_{\omega} s(t,x)I(t,x)S_0(dx), \qquad t \ge 0, \quad \omega \in \mathcal{B}.$$
 (5.6)

 $B(t, \omega)$ is the incidence of the disease, the rate of new infections with trait in ω at time t. So to speak, B is the birth rate of the disease.

5.3 Force of infection and infection age

The infection age a of an infected host is the time since its infection. Let $u(t, \cdot, \omega)$ be the infection-age density of infected hosts at time t and trait in $\omega \in \mathcal{B}$. Recall the incidence B, the rate of infections,

$$u(t, a, \omega) = \int_{\omega} P(a, x) B(t - a, dx), \qquad t > a, \quad \omega \in \mathcal{B}.$$
(5.7)

Here, $P(a, x) \in [0, 1]$ is the probability that a host with trait x that has been infected in the past is still infected (in particular alive) at infection age a, P(0, x) = 1 and $P(\cdot, x)$ decreasing on \mathbb{R}_+ for each $x \in \Omega$. By (5.6),

$$u(t,a,\omega) = \int_{\omega} P(a,x)s(t-a,x)I(t-a,x)S_0(dx), \qquad t > a, \omega \in \mathcal{B}.$$
(5.8)

Let $u_0(a, \omega)$ be the hosts with trait in the set $\omega \in \mathcal{B}$ and infection age a at time 0. Then

$$u(t,a,\omega) = \int_{\omega} Q(a,a-t,x)u_0(a-t,dx), \qquad a > t, \quad \omega \in \mathcal{B}.$$
(5.9)

Here $Q(a, s, x) \in [0, 1]$ is the probability that an infected host with trait x and with infection age s at the beginning is still alive and infected at age a > s. Often, one chooses

$$Q(a, s, x) = \frac{P(a, x)}{P(s, x)}, \qquad a > s > 0.$$
(5.10)

However, this assumes that the infection of the initially infected hosts occurred in the same way as after the start of the epidemic.

The force of infection is given by

$$I(t,x) = \int_0^\infty \left(\int_\Omega \eta(x,a,\xi) u(t,a,d\xi) \right) da, \qquad t \ge 0, \quad x \in \Omega, \tag{5.11}$$

where $\eta(x, a, \xi)$ indicates how a susceptible host with trait $x \in \Omega$ is affected by an infected host with trait $\xi \in \Omega$ and infection age a. By (5.8) and (5.9),

$$I(t,x) = \int_0^t \left(\int_\Omega A(x,a,\xi) s(t-a,\xi) I(t-a,\xi) S_0(d\xi) \right) da + I_0(t,x), \quad (5.12)$$
$$A(x,a,\xi) = \eta(x,a,\xi) P(a,\xi), \qquad x,\xi \in \Omega, \quad a \ge 0, \quad (5.13)$$

and I_0 is the force of infection due to the initially infected hosts,

$$I_0(t,x) = \int_0^\infty \left(\int_\Omega \eta(x,t+a,\xi) Q(t+a,a,\xi) u_0(a,d\xi) \right) da.$$
(5.14)

5.4 The cumulative force of infection

Let

$$J_0(t,x) = \int_0^t I_0(r,x)dr, \qquad t \ge 0, \quad x \in \Omega,$$
 (5.15)

be the cumulative force of infection due to the initially infected hosts. By Tonelli's theorem and (5.1) and (5.12),

$$J(t,x) - J_0(t,x) = \int_0^t \left(I(r,x)dr - I_0(r,x) \right) dr$$
$$= \int_\Omega S_0(d\xi) \int_0^t A(x,a,\xi) \left(\int_a^t s(r-a,\xi)I(r-a,\xi)dr \right) da$$

After a change of variables and by (5.3),

$$J(t,x) - J_0(t,x) = \int_{\Omega} S_0(d\xi) \int_0^t A(x,a,\xi) \Big(\int_0^{t-a} s(r,\xi) I(r,\xi) dr \Big) da.$$

= $\int_{\Omega} S_0(d\xi) \int_0^t A(x,a,\xi) \Big(1 - s(t-a,\xi) \Big) da.$

By (5.1),

$$J(t,x) - J_0(t,x) = \int_{\Omega} S_0(d\xi) \int_0^t A(x,a,\xi) f(J(t-a,\xi)) da, \qquad (5.16)$$

with $f(J) = 1 - e^{-J}$, $J \in \mathbb{R}$, (2.2). Cf. [24, (10)]. By (5.14) and (5.15),

$$J_0(t,x) = \int_0^\infty \int_\Omega \Big(\int_0^t \eta(x,r+a,\xi)Q(r+a,a,\xi)dr \Big) u_0(a,d\xi)da.$$
(5.17)

Assumption 5.1. The functions η, P, Q are nonnegative and are measurable on their respective domains equipped with the appropriate product σ -algebras. $u_0 : \mathbb{R}_+ \times \mathcal{B} \to \mathbb{R}_+$ has the analogous properties of a measure kernel,

$$\int_0^\infty u_0(a,\Omega)da < \infty.$$
(5.18)

The integrals

$$\int_0^\infty \eta(x, r+a, \xi) Q(r+a, a, \xi) dr$$
(5.19)

provide a bounded function of $(x, \xi, a) \in \Omega^2 \times \mathbb{R}_+$. The integrals

$$\int_{\Omega} S_0(d\xi) \int_0^{\infty} A(x, a, \xi) da$$
(5.20)

with A from (5.13) provide a bounded function of $x \in \Omega$.

Remark 5.2. Assumption (5.19) holds, e.g., if $\eta : \Omega \times \mathbb{R}_+ \times \Omega \to \mathbb{R}_+$ is measurable and bounded and $D : \mathbb{R}_+ \times \Omega \to \mathbb{R}_+$ given by

$$D(a,\xi) = \int_0^\infty Q(r+a,a,\xi)dr, \qquad a \in \mathbb{R}_+, x \in \Omega, \tag{5.21}$$

is a bounded function. If Q is given by (5.10), $D(a,\xi)$ is the expected duration of remaining infected life at infection-age a with trait ξ . See [49, Sec.12.4].

6 Existence of minimal solutions

As one can expect from (5.15, $J_0(\cdot, x)$ is increasing on \mathbb{R}_+ for all $x \in \Omega$. See (5.17). Guided by (5.16), for $n \in \mathbb{Z}_+$, we define inductively

$$J_{n+1}(t,x) = J_0(t,x) + \int_{\Omega} S_0(d\xi) \int_0^t A(x,a,\xi) f(J_n(t-a,\xi)) da.$$
(6.1)

By induction, since f in (2.2), is increasing, $J_{n+1}(t,x) \ge J_n(t,x), t \in \mathbb{R}_+, x \in \Omega$, and $J_n(\cdot, x)$ is increasing on \mathbb{R}_+ for all $n \in \mathbb{Z}_+, x \in \Omega$. Further, for $n \in \mathbb{N}$,

$$J_n(t,x) \le J_0(t,x) + \int_{\Omega} S_0(d\xi) \int_0^t A(x,a,\xi) da, \qquad t \in \mathbb{R}_+, x \in \Omega.$$
 (6.2)

For all (t, x), the sequences $(J_n(t, x))$ are increasing and bounded and the limits

$$J(t,x) := \lim_{n \to \infty} J_n(t,x), \qquad t \in \mathbb{R}_+, \quad x \in \Omega,$$
(6.3)

exist pointwise. By Beppo Levi's theorem of monotone convergence, we can take the limit $n \to \infty$ in (6.1) and obtain that J is a solution of (5.16) and $J(\cdot, x)$ is increasing for all $x \in \Omega$.

Let \tilde{J} be also a solution of (5.16). By induction, $J_n(t,x) \leq \tilde{J}(t,x)$ for all $n \in \mathbb{N}$ and $J(t,x) \leq \tilde{J}(t,x)$. So J is the minimal solution of (5.16). Of course, there is at most one minimal solution. J_n can be interpreted as the cumulative infective force due to the infected generations from the initial (0^{th}) to the n^{th} generation. This suggests that the minimal solution, which is their limit, is the epidemiologically relevant solution. In summary:

Theorem 6.1. There is a minimal solution J of (5.16) which is the monotone limit of the recursion (6.1). J is the epidemiologically relevant solution of (5.16).

Analogously, as in [47], we could derive conditions for the minimal solution to be the only solution of (5.16).

6.1 The case of a positive minimum latency period

Assume that there is some $a_0 > 0$ such that

$$\eta(x, a, \xi) = 0, \qquad a \in [0, a_0), \quad x, \xi \in \Omega.$$
 (6.4)

Then, for all traits x, there is a latency period with length greater or equal to a_0 .

Theorem 6.2. For $t \leq na_0$, $J(t, x) = J_n(t, x)$ for all $x \in \Omega$.

Proof. This holds for n = 1. Let $n \in \mathbb{N}$ and the assertion is true for n. By (6.1), $J(t,x) = J_{n+1}(t,x)$ for $t \in [0, na_0]$. Let $t \in (na_0, (n+1)a_0)$. By (5.16),

$$J(t,x) - J_0(t,x) = \int_{\Omega} S_0(dx) \int_{a_0}^t A(x,a,\xi) f(J(t-a,x)) da.$$

Since $J(t-a, x) = J_n(t-a, x)$ for $t \le (n+1)a$ and $a \ge a_0$, $J(t, x) = J_{n+1}(t, x)$ for $t \le (n+1)a$, $x \in \Omega$.

7 The final size of the epidemic

Recall that the minimal solution J of (5.16) has the property that J(t, x) is an increasing bounded function of $t \ge 0$ for any $x \in \Omega$. Hence, the limit

$$w(x) = \lim_{t \to \infty} J(t, x), \quad x \in \Omega,$$
(7.1)

exists for all $x \in \Omega$. By (5.17),

$$J_{0}(t,x) \nearrow w_{0}(x) := \int_{0}^{\infty} \int_{\Omega} A_{0}(x,a,\xi) u_{0}(a,d\xi) da,$$

$$A_{0}(x,a,\xi) = \int_{0}^{\infty} \eta(x,r+a,\xi) Q(r+a,a,\xi) dr.$$
(7.2)

By Beppo Levi's theorem of monotone convergence (or the Lebesgue-Fatou lemma [49, p.468]), we can take the limit $t \to \infty$ in (5.16) (cf. [24, (31)]),

$$w(x) = w_0(x) + \int_{\Omega} k(x,\xi) f(w(\xi)) S_0(d\xi), \qquad x \in \Omega,$$
(7.3)

with f in (2.2) and

$$k(x,\xi) = \int_0^\infty A(x,a,\xi)da, \qquad x,\xi \in \Omega.$$
(7.4)

Theorem 7.1. The function $w : \Omega \to \mathbb{R}_+$ given by

$$w(x) = \lim_{t \to \infty} J(t, x), \qquad x \in \Omega, \tag{7.5}$$

is the minimal solution of (7.3) and is obtained as pointwise limit

$$w(x) = \lim_{n \to \infty} w_n(x), \quad x \in \Omega,$$
(7.6)

of the recursion

$$w_{n+1}(x) = \int_{\Omega} k(x,\xi) f(w_n(\xi)) S_0(d\xi) + w_0(x), \qquad n \in \mathbb{Z}_+, \quad x \in \Omega.$$
(7.7)

Further, for all $x \in \Omega$ and $n \in \mathbb{N}$

$$J_n(t,x) \nearrow w_n(x), \qquad t \nearrow \infty.$$
 (7.8)

We mention that the concept of a minimal solution for the final size equation has already been considered in [47, Thm.3.2]. By (7.5) and (5.1),

$$e^{-w(x)} = \lim_{t \to \infty} s(t, x) =: s_{\infty}(x), \qquad x \in \Omega.$$
(7.9)

Proof. Since f is increasing, by induction, for each $n \in \mathbb{N}$, $w_{n+1}(x) \ge w_n(x)$ for all $x \in \Omega$. Further, $\{w_n(x); x \in \Omega, n \in \mathbb{Z}_+\}$ is bounded. So, the pointwise limit $\tilde{w}(x) = \lim_{n \to \infty} w_n(x)$ exists and, as in the proof of Theorem 6.1, \tilde{w} is the minimal solution of (7.3). Recall $J(t, x) = \lim_{n \to \infty} J_n(t, x)$ and the recursion (6.1). By induction,

$$J_n(t,x) \le w_n(x), \qquad n \in \mathbb{N}, \quad t \ge 0, \quad x \in \Omega.$$

So $J(t,x) \leq \tilde{w}(x)$ and, by (7.5), $w(x) \leq \tilde{w}(x)$ for all $x \in \Omega$. Since w is a solution of (7.3) and \tilde{w} is the minimal solution of (7.3), we also have $\tilde{w}(x) \leq w(x)$ for all $x \in \Omega$. The monotone convergence in (7.8) follows inductively from (6.1) and Beppo Levi's theorem of monotone convergence by taking the limit for $t \to \infty$ and by using (7.7).

Using the concavity of the function f, one can derive conditions which make the minimal solution the only solution of (7.3). But since the minimal solution is the epidemiologically relevant solution, we will not go into the technicalities of such a proof yet [48]. See Theorem 10.17.

Proposition 7.2. Assume that $\eta(x, a, \xi) = 0$ for all $a \in [0, a_0]$, $x \in \Omega$. Then $J(t, x) \leq w_n(x)$ for $t \geq na_0$, $x \in \Omega$ where J is the minimal solution of (5.16).

Proof. This follows from Theorem 6.2 and (7.8).

7.1Measure kernels

Recall the concept of a measure kernel, (2.3). Define

$$\kappa(\omega, x) = \int_{\omega} k(x, \xi) S_0(d\xi), \qquad x \in \Omega, \quad \omega \in \mathcal{B},$$
(7.10)

with k from (7.4). To make κ a measure kernel, (2.3), we assume that

$$\int_{\Omega} k(\cdot,\xi) S_0(d\xi) \text{ is bounded on } \Omega.$$

Remark 7.3. Recall the measure kernel

$$\kappa_{\infty}(\omega, x) = \int_{\omega} s_{\infty}(\xi) \kappa(d\xi, x), \qquad \omega \in \mathcal{B}, \quad x \in \Omega,$$

in Remark 3.4. By (7.10),

$$\kappa_{\infty}(\omega, x) = \int_{\omega} k(x, \xi) s_{\infty}(\xi) S_0(d\xi), \qquad \omega \in \mathcal{B}, \quad x \in \Omega,$$

is the measure kernel associated with the final trait distribution of the susceptible hosts,

$$S_{\infty}(\omega) = \int_{\omega} s_{\infty}(\xi) S_0(d\xi), \qquad \omega \in \mathcal{B}.$$

Recall the function f in (2.2), $f(r) = 1 - e^{-r}$, $r \in \mathbb{R}_+$. We define $F : M_+^b(\Omega) \to M_+^b(\Omega)$ by (2.1), $F(w)(x) = \int_{\Omega} f(w(\xi))\kappa(d\xi, x), \ x \in \Omega, \ w \in M_+^b(\Omega).$

Proposition 7.4 ([17, Prop.4.1.5]). If $g : \Omega \to \mathbb{R}_+$ is a measurable function, then there exists an increasing sequence of simple nonnegative functions such that $g = \lim_{n \to \infty} g_n$ pointwise, and g is integrable. If g is bounded, the convergence is uniform.

Since κ is a measure kernel, F maps $M^b_+(\Omega)$ into itself by Proposition 7.4. The recursive equation (7.7) takes the form

 $w_{n+1} = F(w_n) + w^{\circ}, \quad n \in \mathbb{Z}_+, \qquad w_0 = w^{\circ} \in M^b_+(\Omega), \quad x \in \Omega.$ (7.11)

By induction, since κ is a measure kernel and f is increasing, (w_n) is an increasing sequence in $M^b_+(\Omega)$ which is bounded by $\kappa(\Omega, \cdot) + w^\circ$. The final cumulative force of infection, w is the pointwise limit of (w_n) ,

$$w(x) = \lim_{n \to \infty} w_n(x), \qquad x \in \Omega.$$
(7.12)

By Beppo Levi's monotone convergence theorem, $w \in M^b_+(\Omega)$ satisfies

$$w = w^{\circ} + F(w).$$
 (7.13)

Remark 7.5. We introduce the measure kernel notation less to have shorter formulas but to make the connection to the final size consideration in [47, Sec.3].

Not surprisingly, the final size of the cumulative force of infection depends on the final size of the cumulative initial force of infection in an increasing way.

Theorem 7.6. Let $w^{\circ}, \tilde{w}^{\circ} \in M^b_+(\Omega)$ and $w, \tilde{w} \in M^b_+(\Omega)$ be the minimal solutions of

 $w = w^{\circ} + F(w)$ and $\tilde{w} = \tilde{w}^{\circ} + F(\tilde{w}).$

Then, if $w^{\circ} \leq \tilde{w}^{\circ}$ on Ω , also $w \leq \tilde{w}$ on Ω and $w - w^{\circ} \leq \tilde{w} - \tilde{w}^{\circ}$ as well.

Proof. Both $w = \lim_{n \to \infty} w_n$ and $\tilde{w} = \lim_{n \to \infty} \tilde{w}_n$ pointwise on Ω , where w_n is given by the recursion (7.11) and \tilde{w}_n by an analogous recursion. By induction, since f is increasing, $w_n \leq \tilde{w}_n$ on Ω for all $n \in \mathbb{N}$ and so $w \leq \tilde{w}$ on Ω . The last inequality follows from $w - w^\circ = F(w) \leq F(\tilde{w}) = \tilde{w} - \tilde{w}^\circ$. \Box

We continue this section with the following observation which is as trivial as it is fundamental.

Theorem 7.7. $w = w_0$ if and only if $w_1 = w_0$.

Proof. Let $w = w_0$. Since $w \ge w_1 \ge w_0$, this implies $w_1 = w_0$.

Let $w_1 = w_0$. By induction and (7.11), $w_n = w_0$ for all $n \in \mathbb{N}$ and so $w = w_0$ by (7.13).

Corollary 7.8. $w = w_0$ if and only if

$$\int_{\Omega} w_0(\xi) \kappa(d\xi, x) = 0, \qquad x \in \Omega.$$

Let $\mathcal{R}_0 = \mathbf{r}(\kappa) > 0$ and θ be the eigenfunctional of the operator K associated with its spectral radius (Theorem 3.1).

Lemma 7.9. If $\theta(w^{\circ}) = 0$ and (w_n) is provided by the recursion the recursion (7.11), then $\theta(w_n) = 0$ for all $n \in \mathbb{N}$.

Unfortunately, since θ may not be continuous with respect to pointwise convergence, this may not imply that $\theta(w) = 0$ for the pointwise limit w of (w_n) .

If there is a positive minimum latency period, we have the following without an extra assumption. **Theorem 7.10.** Let $a_0 > 0$ and $\eta(x, a, \xi) = 0$ for all $a \in [0, a_0)$. Then, if $\theta(w_0) = 0$, $\theta(J(t, \cdot)) = 0$ for all $t \ge 0$ and the minimal solution J to (5.16).

Proof. By Theorem 7.2,

$$J(t,x) \le w_n(x), \quad t \in [0, na_0), \quad x \in \Omega.$$

Since θ is additive, by Lemma 7.9

$$\theta(J(t,\cdot)) \le \theta(w_n) = 0, \quad t \in [0, na_0), \quad x \in \Omega.$$

Since this holds for all $n \in \mathbb{N}$, $\theta(J(t, \cdot)) = 0$ for all $t \ge 0$.

7.2 Sequences of minimal solutions

Here are first results that relate sequences of minimal solutions to $w = F(w) + w^{\circ}$ to fixed points of F.

Theorem 7.11. Let $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ be a sequence in $M_{+}^{b}(\Omega)$ and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω . For all $\ell \in \mathbb{N}$, let $w_{\ell} \in M_{+}^{b}(\Omega)$ be the minimal solutions of

$$w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}.$$

Then there exist a minimal $\tilde{w} \in M^b_+(\Omega)$ such that $\tilde{w} = F(\tilde{w})$ and

$$\limsup_{n \to \infty} w_{\ell}(x) \le \tilde{w}(x), \qquad x \in \Omega.$$

Proof. Set

$$\breve{w}(x) = \limsup_{n \to \infty} w_{\ell}(x), \qquad x \in \Omega.$$

By Fatou's Lemma, applied for any $x \in \Omega$,

$$\breve{w}(x) \leq \int_{\Omega} \limsup_{n \to \infty} f(w_{\ell}(\xi))\kappa(d\xi, x) + \limsup_{n \to \infty} w_{\ell}^{\circ}(x).$$

Since f is monotone and continuous and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω ,

$$\breve{w}(x) \leq \int_{\Omega} f(\breve{w}_{\ell}(\xi))\kappa(d\xi, x) = F(\breve{w})(x).$$

We define recursively

$$\tilde{w}_{n+1} = F(\tilde{w}_n), \quad n \in \mathbb{Z}_+, \qquad \tilde{w}_0 = \breve{w}.$$

Since F is an increasing map, by induction, (\tilde{w}_n) is an increasing sequence of functions that is also bounded and has a pointwise limit $\tilde{w} \in M^b_+(\Omega)$, $\tilde{w} \geq \tilde{w}$. By Lebesgue's theorem of dominated convergence, $\tilde{w} = F(\tilde{w})$.

We claim that \tilde{w} is the minimal solution of $\check{w} = F(\check{w})$ with $\check{w} \geq \check{w}$. Let \check{w} be such a solution. Then $\check{w} \geq \tilde{w}_0$. If $n \in \mathbb{N}$ and $\check{w} \geq \tilde{w}_n$, then

$$\check{w} = F(\check{w}) \ge F(\tilde{w}_n) = \tilde{w}_{n+1}.$$

By induction, $\check{w} \geq \tilde{w}_n$ for all $n \in \mathbb{Z}_+$ and $\check{w} \geq \tilde{w}$ by taking the pointwise limit.

The next two results give us some vague idea of the final size of the epidemic if the number of initial infectives is small.

Corollary 7.12. Let $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ be a sequence in $M_{+}^{b}(\Omega)$ and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω . For all $\ell \in \mathbb{N}$, let $w_{\ell} \in M_{+}^{b}(\Omega)$ be the minimal solutions of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

Assume that $\tilde{w} = 0$ is the only solution to $F(\tilde{w}) = \tilde{w} \in M^b_+(\Omega)$. Then $w_\ell \to 0$ as $\ell \to \infty$ pointwise on Ω .

Theorem 7.13. Let (w_{ℓ}°) be a decreasing sequence in $M_{+}^{b}(\Omega)$, $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω . Let (w_{ℓ}) be the sequence of minimal solutions of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

Then there exists some fixed point $\tilde{w} = F(\tilde{w})$ in $M^b_+(\Omega)$ such that $w_\ell \searrow \tilde{w}$ and $w_\ell - w^\circ_\ell \searrow \tilde{w}$ as $\ell \to \infty$ pointwise on Ω .

Proof. By Theorem 7.6, the sequences (w_{ℓ}) and $(w_{\ell} - w_{\ell}^{\circ})$ are decreasing. Since they are bounded below by the zero function, they converge pointwise to some $\tilde{w} \in M^b_+(\Omega)$ with

$$w_{\ell} - w_{\ell}^{\circ} \ge \tilde{w}, \qquad \ell \in \mathbb{N}.$$

By Lebesgue's dominated convergence theorem, applied to

$$w_{\ell}(x) = \int_{\Omega} f(w_{\ell}(\xi))\kappa(d\xi, x) + w_{\ell}^{\circ}(x)$$

for each $x \in \Omega$, we take the limit as $\ell \to \infty$, and \tilde{w} satisfies $\tilde{w} = F(\tilde{w})$. \Box

8 Dominated measure kernels

A measure kernel κ is called *dominated* by $0 \neq \nu \in \mathcal{M}_+(\Omega)$ if

$$\kappa(\omega, x) \le \nu(\omega), \qquad \omega \in \mathcal{B}.$$
 (8.1)

The measure kernel defined by (7.10) is dominated by a multiple of S_0 if the function k is bounded on $\Omega \times \Omega$.

Throughout this section, we assume that $\mathcal{R}_0 = \mathbf{r}(\kappa) = \mathbf{r}(K) > 0$ and that θ is the bounded linear eigenfunctional of K associated with \mathcal{R}_0 by Theorem 3.1.

Proposition 8.1. Let κ be a dominated measure kernel. Let w be the minimal solution of $w = F(w) + w^{\circ}$ and (w_n) be the recursion $w_n = F(w_{n-1}) + w^{\circ}$, $n \in \mathbb{N}, w_0 = w^{\circ}$.

Then $||w - w_n||_{\infty} \to 0$ and $\theta(w_n) \to \theta(w)$ as $n \to \infty$.

Proof. Since $w_n \nearrow w$ pointwise on Ω and f is continuous and increasing, $f \circ w_n \nearrow f \circ w$ pointwise on Ω . By Beppo Levi's theorem of monotone convergence,

$$\int_{\Omega} (f \circ w_n) d\nu \to \int_{\Omega} (f \circ w) d\nu, \qquad n \to \infty.$$

Since $f \circ w_n \leq f \circ w$,

$$\int_{\Omega} |(f(w(\xi)) - f(w_n(\xi)))|\nu(d\xi)| = \int_{\Omega} \left((f(w(\xi)) - f(w_n(\xi))) \right) \nu(d\xi) \xrightarrow{n \to \infty} 0.$$

By (8.1),

$$\|w - w_{n+1}\|_{\infty} = \sup_{x \in \Omega} \int_{\Omega} \left| (f(w(\xi)) - f(w_n(\xi))) \middle| \kappa(d\xi, x) \right| \le \int_{\Omega} \left((f(w(\xi)) - f(w_n(\xi))) \nu(d\xi) \xrightarrow{n \to \infty} 0. \right]$$

Lemma 7.9 implies the following result.

Proposition 8.2. Let w be the minimal solution of $w = F(w) + w^{\circ}$. Let the measure kernel κ be dominated. Then $\theta(w) = 0$ if $\theta(w^{\circ}) = 0$.

The next result gives us some better idea than before (Theorem 7.13) about the final size of the epidemic when the number of initial infectives is small.

Theorem 8.3. Let κ be dominated by a measure ν and let (w_{ℓ}°) be a decreasing sequence in $M_{+}^{b}(\Omega)$, $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω . Let (w_{ℓ}) be the sequence of minimal solutions of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

(a) Then $w_{\ell} - w_{\ell}^{\circ} \searrow \tilde{w}$ as $\ell \to \infty$ uniformly on Ω for some solution $\tilde{w} = F(\tilde{w})$ in $M^{b}_{+}(\Omega)$.

(b) Assume in addition that $\mathcal{R}_0 > 1$ and $\theta(w_{\ell}^{\circ}) > 0$ for all $\ell \in \mathbb{N}$. Then $\int_{\Omega} (f \circ \tilde{w}) d\nu \geq \ln \mathcal{R}_0$ and $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$.

Proof. (a) By Theorem 7.6, the sequences (w_{ℓ}) and $(w_{\ell} - w_{\ell}^{\circ})$ are decreasing. By Theorem 7.13, $w_{\ell} - w_{\ell}^{\circ} \to \tilde{w}$ as $\ell \to \infty$ pointwise on Ω for some $w \in M^b_+(\Omega)$ with $\tilde{w} = F(\tilde{w})$. For all $\ell \in \mathbb{N}$ and $x \in \Omega$,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) - \tilde{w}(x) = \int_{\Omega} \left(f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \kappa(d\xi, x) \right) d\xi$$

Since κ is dominated by ν ,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) - \tilde{w}(x) \le \int_{\Omega} \left(f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \right) \nu(d\xi).$$

Since the right hand side does not depend on x, $||w_{\ell} - w_{\ell}^{\circ} - \tilde{w}||_{\infty} \to 0$ as $\ell \to \infty$.

(b) Let $\mathcal{R}_0 > 1$. For all $\ell \in \mathbb{N}$, since κ is dominated by ν ,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) \le \int_{\Omega} (f \circ w_{\ell}) d\nu$$

and, by Theorem 3.2,

$$\ln \mathcal{R}_0 \le \|w_\ell - w_\ell^\circ\|_\infty \le \int_\Omega (f \circ w_\ell) d\nu.$$

By Lebesgue's theorem of dominated convergence,

$$\int_{\Omega} (f \circ w) d\nu \ge \ln \mathcal{R}_0.$$

Further $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$.

Proof of Theorem 3.5. Apply Theorem 8.3 with $w_{\ell}^{\circ} = (1/\ell)w^{\circ}$ for $\ell \in \mathbb{N}$ and $w = w_1$.

8.1 Positivity points of θ

In view of the previous results, it is of interest for which $w^{\circ} \in M^b_+(\Omega)$ we have $\theta(w^{\circ}) > 0$.

Proposition 8.4. Let $\mathcal{R}_0 > 0$ and let there exist $\nu_j, \tilde{\nu}_j \in \mathcal{M}_+(\Omega)$ and $k_j \in \mathcal{M}_+^b(\Omega), j = 1, \ldots, n$, such that

$$\sum_{j=1}^{n} \tilde{\nu}_j(\omega) k_j(x) \le \kappa(\omega, x) \le \sum_{j=1}^{n} \nu_j(\omega) k_j(x), \qquad \omega \in \mathcal{B}, \quad x \in \Omega.$$
(8.2)

Then, for any $w \in M^b_+(\Omega)$, $\theta(w) > 0$ if $\int_{\Omega} w \, d\tilde{\nu}_j > 0$ for j = 1, ..., n, while $\theta(w) = 0$ if $\int_{\Omega} w \, d\nu_j = 0$ for j = 1, ..., n.

Proof. By (8.2), for $w \in M^b_+(\Omega)$,

$$\sum_{j=1}^{n} \left(\int_{\Omega} w \, d\tilde{\nu}_j \right) k_j \le Kw \le \sum_{j=1}^{n} \left(\int_{\Omega} w \, d\nu_j \right) k_j.$$

Then Kw = 0 and $\theta(w) = 0$ if $\int_{\Omega} w d\nu_j = 0$ for $j = 1, \dots, j$.

Recall the constant function u_1 with value 1. Since θ is linear,

$$0 < \mathcal{R}_0 \theta(u_1) = \theta(Ku_1) \le \sum_{j=1}^{\infty} \nu_j(\Omega) \theta(k_j).$$

This implies that $\sum_{j=1}^{n} \theta(k_j) > 0$. Further

$$\mathcal{R}_0 \theta(w) \ge \sum_{j=1}^n \Big(\int_{\Omega} w d\tilde{\nu}_j \Big) \theta(k_j).$$

So $\theta(w) > 0$ if $\int_{\Omega} w \, d\tilde{\nu}_j > 0$ for $j = 1, \dots, n$.

		-
н		н
		-

9 Semi-separable measure kernels

A measure kernel κ is called *semi-separable* if there are nonzero $k_0 \in M^b_+(\Omega)$, $\nu \in \mathcal{M}_+(\Omega)$ and $\delta \in (0, 1]$ such that

$$\delta\nu(\omega)k_0(x) \le \kappa(\omega, x) \le \nu(\omega)k_0(x), \qquad x \in \Omega, \quad \omega \in \mathcal{B}.$$
(9.1)

Proposition 9.1. Assume that κ is semi-separable, (9.1), and let $w \in M^b_+(\Omega)$. Then, $\mathcal{R}_0 > 0$ if and only if $\int_{\Omega} k_0 d\nu > 0$, and $\theta(w) > 0$ if and only if $\int_{\Omega} w d\nu > 0$.

Proof. By (9.1),

$$\delta \Big(\int_{\Omega} k_0 \, d\nu \Big) k_0 \le K k_0 \le \Big(\int_{\Omega} k_0 \, d\nu \Big) k_0.$$

This implies [51, Thm.3.1][54, Thm.5.31] [51, Thm.3.3][54, Thm.6.15] that

$$\delta \int_{\Omega} k_0 d\nu \leq \mathcal{R}_0 = \mathbf{r}(K) \leq \int_{\Omega} k_0 d\nu.$$

The remaining statement is a special case of Proposition 8.4 for n = 1. \Box

Proposition 9.2. Assume that κ is semi-separable, (9.1).

- (a) Then there exists at most one non-zero solution $w \in M^b_+(\Omega)$ of w = F(w).
- (b) If $\mathcal{R}_0 \leq 1$, the zero function is the only $w \in M^b_+(\Omega)$ with w = F(w).

Proof. We follow [29, Sec.6.1]. By (9.1),

$$\delta k_0(x) \int_{\Omega} (f \circ w) d\nu \le F(w)(x) \le k_0(x) \int_{\Omega} (f \circ w) d\nu \le k_0(x)\nu(\Omega).$$
(9.2)

By (9.2), F(w) is not the zero function if and only if

$$\int_{\Omega} (f \circ w) d\nu > 0. \tag{9.3}$$

Since f is concave and f(0) = 0,

$$F(tw) \ge tF(w), \qquad t \in [0,1], \quad w \in M^b_+(\Omega).$$

(a) According to [29, Thm.6.3], it is sufficient to show that, for any $t \in (0, 1)$ and any $w \in M^b_+(\Omega)$ with

$$\int_{\Omega} (f \circ w) d\nu > 0,$$

some $\eta > 0$ can be found such that

$$F(tw) \ge (1+\eta)tF(w). \tag{9.4}$$

Since f is strictly concave, f(tr) > tf(r) for all r > 0, $t \in (0,1)$. Let $t \in (0,1)$. Then $f(tw(\xi)) - tf(w(\xi))$ is nonnegative for all $\xi \in \Omega$ and is positive for $\xi \in \Omega$ if $w(\xi) > 0$.

Suppose that

$$\int_{\Omega} (f(tw(\xi)) - tf(w(\xi)))\nu(d\xi) = 0$$

Then $w(\xi) = 0$ for ν -a.a. $\xi \in \Omega$ and $\int_{\Omega} (f \circ w) d\nu = 0$. By contraposition if $\int_{\Omega} (f \circ w) d\mu > 0$.

By contraposition, if $\int_{\Omega} (f \circ w) d\nu > 0$,

$$\tilde{\eta} = \int_{\Omega} \left(f(tw(\xi)) - tf(w(\xi)) \right) \nu(d\xi) > 0.$$

Now,

$$F(tw)(x) - tF(w)(x) = \int_{\Omega} \left[f(tw(\xi)) - tf(w(\xi)) \right] \kappa(d\xi, x)$$

with the expression in $[\cdot]$ being nonnegative. By (9.1),

$$F(tw)(x) - tF(w)(x) \ge \delta k_0(x) \int_{\Omega} \left[f(tw(\xi)) - tf(w(\xi)) \right] \nu(d\xi) \ge \delta k_0(x) t\hat{\eta}$$

with $\hat{\eta} = \tilde{\eta}/t$. By (9.2) and (9.1),

$$F(tw)(x) - tF(w)(x) \ge \delta k_0(x)t\hat{\eta}\frac{1}{\nu(\Omega)}\int_{\Omega} f(w(\xi))\nu(d\xi) \ge \delta t\frac{\hat{\eta}}{\nu(\Omega)}F(w(x)).$$

We reorganize,

$$F(tw)(x) \ge \left(1 + \frac{\delta\hat{\eta}}{\nu(\Omega)}\right) tFw(x), \qquad x \in \Omega,$$

and we have shown (9.4) with $\eta = \frac{\delta \hat{\eta}}{\nu(\Omega)}$.

(b) Suppose that $0 \neq w \in M^b_+(\Omega)$ satisfies w = F(w). Then (9.3) holds. Let t = 1/2. By the same considerations as before, there exists some $\eta > 0$ such that (9.4) is valid. By (2.7),

$$K(tw) \ge F(tw) \ge (1+\eta)(tw).$$

This implies that $\mathcal{R}_0 = \mathbf{r}(K) \ge (1 + \eta)$ [51, Thm.3.1] [54, Thm.5.31], a contradiction.

We are now in the position to prove the preview results in Section 3.4.

Proof of Theorem 3.7. Part (a) follows from Corollary 7.8 and (9.1).

(b) Existence of \tilde{w} follows from Corollary 3.6 with ν being replaced by $||k_0||_{\infty}\nu$. Uniqueness of \tilde{w} follows from Proposition 9.2.

The remaining statements follow from Proposition 9.1, Theorem 3.2 (b) and Theorem 3.5. $\hfill \Box$

Proof of Theorem 3.8. (a) Pointwise convergence of $w_{\ell} \to 0$ follows from Corollary 7.12 and Proposition 9.2 (b). For all $\ell \in \mathbb{N}$,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) \le k_0(x) \int_{\Omega} f(w_{\ell}(\xi))\nu(d\xi) \to 0$$

as $\ell \to \infty$ by Lebesgue's theorem of dominated convergence because $f(w_{\ell}(\xi)) \to 0$ as $\ell \to \infty$. Since k_0 is bounded, $w_{\ell} - w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ uniformly on Ω .

(b) By Fatou's lemma and the increase and continuity of f,

$$\int_{\Omega} f \circ (\limsup_{\ell \to \infty} w_{\ell}) d\nu \ge \int_{\omega} \limsup_{\ell \to \infty} (f \circ w_{\ell}) d\nu \ge \limsup_{\ell \to \infty} \int_{\Omega} (f \circ w_{\ell}) d\nu.$$

By Theorem 3.7 (b),

$$\int_{\Omega} (f \circ w_{\ell}) d\nu \ge \ln \mathcal{R}_0 / \|k_0\|_{\infty}, \qquad \ell \in \mathbb{N}.$$

So, $\limsup_{\ell \to \infty} w_{\ell}$ is not the zero function. By Theorem 7.11, there exists a solution \hat{w} of $\hat{w} = F(\hat{w})$ with $\limsup_{\ell \to \infty} w_{\ell} \leq \hat{w}$. Then \hat{w} is not the zero function and $\hat{w} = \tilde{w}$ by Theorem 3.7 (b). Since $w_{\ell} \geq \tilde{w}$ for all $\ell \in \mathbb{N}$ by Theorem 3.7 (b), $w_{\ell} \to \tilde{w} = \hat{w}$ as $\ell \to \infty$ pointwise on Ω .

For all $\ell \in \mathbb{N}$,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) - \tilde{w}(x) = \int_{\Omega} \left(f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \right) \kappa(d\xi, x).$$

By (9.1),

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) - \tilde{w}(x) \le k_0(x) \int_{\Omega} \left(f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \right) \nu(d\xi) d\xi$$

Since k_0 is bounded and $f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \to 0$ as $\ell \to \infty$ pointwise for $\xi \in \Omega$, by Lebesgue's theorem of dominated convergence,

$$w_{\ell}(x) - w_{\ell}^{\circ}(x) \to \tilde{w}(x), \quad \ell \to \infty$$

uniformly for $x \in \Omega$.

10 Metric spaces of traits and Feller kernels

To replace semi-separability of the kernel as an assumption, we assume that Ω is a metric space with metric ρ and \mathcal{B} the σ -algebra of Borel sets. Assume that Ω is not just a single point. Let $C^b(\Omega)$ denote the Banach space of bounded continuous functions with the supremum norm which is a closed subspace of $M^b(\Omega)$. We start with a few technical observations.

Remark 10.1. If μ is a finite nonnegative measure on \mathcal{B} , then μ is inner and outer regular and $C^{b}(\Omega)$ is dense in $L^{1}(\Omega, \mu)$.

Proof. See [1, Sec.12.1] for the regularity statement and [18, L.IV.8.19] for the density statement. Notice that μ is regular in [18, L.IV.8.19] if and and only it is inner and outer regular in [1, Sec.12.1].

Lemma 10.2. Let $g \in M^b_+(\Omega)$ be topologically positive (Definition 3.9).

- (a) Then there exists a non-zero Lipschitz continuous function $\tilde{g}: \Omega \to \mathbb{R}_+$ with a Lipschitz constant ≤ 1 such that $\tilde{g} \leq g$ and \tilde{g} is strictly positive on every nonempty open subset U with $\inf_U g > 0$.
- (b) If g is lower semicontinuous, $x \in \Omega$ and g(x) > 0, then $\tilde{g}(x) > 0$ for the function \tilde{g} from (a).

Proof. (a) Recall that ρ denotes the metric on Ω . Let $g \in M^b_+(\Omega)$. Define

$$\tilde{g}(x) = \inf \left\{ \rho(x,\xi) + g(\xi); \ \xi \in \Omega \right\}, \qquad x \in \Omega.$$

Then \tilde{g} is Lipschitz continuous with a Lipschitz constant ≤ 1 and $0 \leq \tilde{g} \leq g$ on Ω [54, Prop.2.78].

Let U be a nonempty open subset of Ω such that $\inf_U g > 0$. Then, for any $x \in U$ there exists some $\delta \in (0, \inf_U g)$ such that $\xi \in U$ and $g(\xi) > \delta$ whenever $\xi \in \Omega$ and $\rho(x, \xi) < \delta$. This implies that

$$\rho(x,\xi) + g(\xi) \ge \delta, \quad \xi \in \Omega,$$

and $\tilde{g}(x) \geq \delta$.

(b) Let g be lower semicontinuous. Suppose that $x \in \Omega$ and $\tilde{g}(x) = 0$. Then there exists a sequence (ξ_n) in Ω such that $\rho(x,\xi_n) + g(\xi_n) \to 0$ as $n \to \infty$. In particular, $\xi_n \to x$ as $n \to \infty$. Since g is lower continuous, $0 = \liminf_{n\to\infty} g(\xi_n) \ge g(x)$ [54, L.A.49].

Conversely, g(x) > 0 implies that $\tilde{g}(x) > 0$.

10.1 Feller kernels

A measure kernel κ is called a *Feller kernel* if the map K on $M^b(\Omega)$ induced by κ maps $C^b(\Omega)$ into itself (Definition 3.9).

Proposition 10.3. Let κ be a Feller kernel and $g : \Omega \to \mathbb{R}_+$ be lower semicontinuous. Then Kg and Fg are lower semicontinuous.

Proof. Let $g : \Omega \to \mathbb{R}_+$ be lower semicontinuous. Then g is the pointwise limit of an increasing sequence of Lipschitz continuous functions [1, Thm.3.13][54, Prop.2.78] and $f \circ g$ is the pointwise limit of an increasing sequence of continuous functions. Since κ is a Feller kernel, Kg and F(g) are pointwise limits of increasing sequences of continuous functions and thus lower semicontinuous [54, L.A.53].

Theorem 10.4. Let κ be a Feller kernel and $w^{\circ} \in M^b_+(\Omega)$ be lower semicontinuous. Then the minimal solution $w \in M^b_+(\Omega)$ of $w = F(w) + w^{\circ}$ is lower semicontinuous.

Proof. Recall that w is given as the pointwise limit $w = \lim_{n \to \infty} w_n$ of the recursion

$$w_n = F(w_{n-1}) + w^\circ, \quad n \in \mathbb{N}, \qquad w_0 = w^\circ.$$
 (10.1)

By induction, (w_n) is a increasing sequence of functions in $M^b_+(\Omega)$ and every w_n is lower semicontinuous by Proposition 10.3 and so is the pointwise limit $w = \sup_{n \in \mathbb{N}} w_n$ [54, L.A.53].

Proposition 10.5. Let κ be a measure kernel such that $\kappa(\Omega, \cdot)$ is continuous on Ω and Kg is lower semicontinuous for any $g \in C^b_+(\Omega)$. Then κ is a Feller kernel.

Proof. Let $g \in C^b_+(\Omega)$. It is sufficient to show that Kg is upper semicontinuous. Define $\tilde{g} \in C^b_+(\Omega)$ by $\tilde{g}(x) = ||g||_{\infty} - g(x), x \in \Omega$. By assumption, $K\tilde{g}$ is lower semicontinuous,

$$K\tilde{g} = \|g\|_{\infty}\kappa(\Omega, \cdot) - Kg.$$

So, Kg is upper semicontinuous. See [54, Rem.A.48].

10.2 Topological irreducibility

We call κ topologically irreducible if for any nonempty open strict subset U of Ω there exist some $x \in \Omega \setminus U$ such that $\kappa(U, x) > 0$. If κ is topologically irreducible, then [54, Rem.13.57]

$$\kappa(\Omega \setminus \{x\}, x) > 0, \qquad x \in \Omega.$$
(10.2)

For the special case (7.10), κ is topologically irreducible if $S_0(\omega) > 0$ for any nonempty open subset of Ω and if for any nonempty open strict subset ω of Ω there exists some $x \in \Omega \setminus \omega$ such that $k(x, \cdot)$ is not zero a.e. on ω . **Theorem 10.6.** Let κ be topologically irreducible, w° be lower semicontinuous and not the zero function and w be the minimal solution of $w = F(w) + w^{\circ}$ (7.6). Then w is lower semicontinuous and $w - w^{\circ}$ is strictly positive on Ω .

Proof. By Proposition 10.4, w is lower semicontinuous and $U = \{w > 0\}$ is open [54, A.51].

Since w° is not the zero function, $\{w > 0\}$ is not the empty set.

Suppose that $U = \{w > 0\} \neq \Omega$. Since κ is topologically irreducible, there exist some $x \in \Omega \setminus U$ such that $\kappa(U, x) > 0$. Recall that

$$w(x) = \int_{\Omega} f(w(\xi))\kappa(d\xi, x) + w^{\circ}(x).$$

Now $U = \bigcup_{n \in \mathbb{N}} \{w > 1/n\}$. Since $\kappa(\cdot, x)$ is a measure, there is some $n \in \mathbb{N}$ such that $\kappa(\{w > 1/n\}, x) > 0$. Then

$$w(x) \ge f(1/n)\kappa(\{w > 1/n\}, x) > 0,$$

and $x \in U$, a contradiction. Now $f \circ w$ is strictly positive and $F(w) = w - w^{\circ}$ is strictly positive by (10.2).

Remark 10.7. Topological irreducibility of the kernel is necessary for the epidemic to always reach all traits.

Proof. Assume that κ is not topologically irreducibility. Then there exists some nonempty open strict subset U of Ω such that $\kappa(U, x) = 0$ for all $x \in \Omega \setminus U$. Since Ω is a metric space, there is some nonzero continuous function w_0 such that $w_0(x) = 0$ for all $x \in \Omega \setminus U$ [1, L.3.20]. By induction and (7.11) $w_n(x) = 0$ for all $n \in \mathbb{N}$ and $x \in \Omega \setminus U$. Then the pointwise limit function w also satisfies w(x) = 0 for all $x \in \Omega \setminus U$.

Proof of Theorem 3.10. Let U be an open nonempty subset of Ω such that $\inf_U w > 0$. By Lemma 10.2, there exists some $\tilde{w}^{\circ} \in C^b_+(\Omega)$ which is strictly positive on U and $0 \leq \tilde{w}^{\circ} \leq w^{\circ}$ on Ω . Let \tilde{w} be the minimal solution of $\tilde{w} = F(\tilde{w}) + \tilde{w}^{\circ}$. By Theorem 7.6, $\tilde{w} \leq w$ on Ω . By Theorem 10.6, $\{\tilde{w} > 0\} = \Omega$ and so $\{w > 0\} = \Omega$. Then $f \circ w$ is strictly positive on Ω and $F(w) = w - w^{\circ}$ is strictly positive by (10.2).

Theorem 10.8. Let $\tilde{w} : \Omega \to \mathbb{R}_+$ be a lower semicontinuous non-zero solution to the inequality

$$\tilde{w}(x) \geq \int_{\Omega} f(\tilde{w}(\xi)) \kappa(d\xi, x), \qquad x \in \Omega,$$

and let κ be a topologically irreducible Feller kernel. Then \tilde{w} is strictly positive on Ω .

Proof. Since \tilde{w} is lower semicontinuous and not the zero function, the set $U = \{\tilde{w} > 0\}$ is an open nonempty set [54, A.51]. If the assertion is false, U is a strict subset of Ω . Since κ is topologically irreducible, there exists some $x \in \Omega \setminus U$ such that $\kappa(U, x) > 0$. Since $\kappa(\cdot, x)$ is a measure, there is some $n \in \mathbb{N}$ such that $\kappa(\{w > 1/n\}, x\} > 0$. Then

$$w(x) \ge f(1/n)\kappa(\{w > 1/n\}, x) > 0,$$

and $x \in U$, a contradiction.

10.3 Tightness

Recall Definition 3.9 and 3.11.

For instance, the Feller kernel κ in (7.10) is tight if S_0 is tight and k is bounded on Ω^2 . Recall that every finite nonnegative measure on \mathcal{B} is tight if Ω is a Polish space, i.e., Ω is separable and complete under a metric that is topologically equivalent to the original one.

Proof of Theorem 3.12. Existence of the tight eigenmeasure μ follows from [54, Thm.13.39] and [54, Thm.13.42].

In addition, let κ be topologically irreducible. Apply [54, Cor.13.60] and Lemma 10.2.

10.4 Strong Feller kernels

While Theorem 3.13 in conjunction with Theorem 3.2 (a) gives a good display of the threshold properties of \mathcal{R}_0 , the relation of minimal solutions of $w = F(w) + w^\circ$ to fixed points $\tilde{w} = F(\tilde{w})$ may be informative. To this end, we strengthen the concept of a Feller kernel (Definition 3.14).

Proposition 10.9. Let κ be a measure kernel. Assume that $\kappa(\Omega, \cdot)$ is continuous on Ω and that $\kappa(\omega, \cdot)$ is lower semicontinuous for any $\omega \in \mathcal{B}$.

Then κ is a strong Feller kernel.

Proof. Let $\omega \in \mathcal{B}$. Then $\kappa(\omega, \cdot)$ is lower semicontinuous. Further, $\Omega \setminus \omega \in \mathcal{B}$. Then $\kappa(\Omega \setminus \omega, \cdot)$ is lower semicontinuous,

$$\kappa(\Omega\setminus\omega,\cdot)=\kappa(\Omega,\cdot)-\kappa(\omega,\cdot).$$

Since $\kappa(\Omega, \cdot)$ is continuous, $\kappa(\omega, \cdot)$ is upper semicontinuous [54, Rem.A.48]. Since $\kappa(\omega, \cdot)$ is both lower and upper semicontinuous, it is continuous. Use Lemma A.49 and A.50 in [54].

In the framework of the model in Section 5 it is difficult to find an example of a Feller kernel that is not a strong Feller kernel. See Proposition 10.11.

In the context of [48], it is easy to give an example of Feller kernel that is not a strong Feller kernel like $\kappa(\omega, x) = \chi_{\omega}(x)$ where χ_{ω} is the characteristic or indicator function of ω , $\chi_{\omega}(x) = 1$ if $x \in \omega$ and 0 otherwise.

Lemma 10.10. If κ is a strong Feller kernel, then K and F map $M^b_+(\Omega)$ into $C^b_+(\Omega)$.

Proof. Let κ be a strong Feller kernel. Since bounded measurable functions are uniform limits of linear combinations of characteristic functions (Proposition 7.4), the operators K and F induced by a strong Feller kernel map $M^b_+(\Omega)$ into $C^b_+(\Omega)$.

In Section 8, we considered dominated measure kernels. For perspective, we mention the following result.

Proposition 10.11. Let κ be a Feller kernel that is dominated by some $\nu \in \mathcal{M}_+(\Omega)$. Then κ is a strong Feller kernel. Actually,

$$Kg \in C^b(\Omega), \qquad g \in L^1(\Omega, \nu).$$

Proof. Since the Feller kernel κ is dominated by the measure ν , for each $x \in \Omega$, the measure $\kappa(\cdot, x)$ is absolutly continuous with respect to ν . By the Radon-Nikodym theorem, there exists some $k_x \in L^1_+(\Omega, \nu)$ such that

$$\nu(\omega) \ge \kappa(\omega, x) = \int_{\omega} k_x(\xi)\nu(d\xi), \qquad \omega \in \mathcal{B}, \quad x \in \Omega.$$

Hence, $k_x(\xi) \leq 1$ for all $\xi \in \Omega$. By Remark 10.1, $C^b(\Omega)$ is dense in $L^1(\Omega, \nu)$. Let $g \in L^1(\Omega, \nu)$. Then there exists a sequence (g_n) in $C^b(\Omega)$ such that $||g - g_n||_1 \to 0$ as $n \to \infty$. For all $x \in \Omega$,

$$|Kg_n(x) - Kg(x)| \le \int_{\Omega} |g_n(\xi) - g(\xi)| k_x(\xi) \nu(d\xi) \le ||g_n - g||_1,$$

and $Kg_n \to Kg$ as $n \to \infty$ uniformly on Ω . Since κ is a Feller kernel, all Kg_n are continuous and so is their uniform limit Kg.

Theorem 10.12. Let κ be a tight strong Feller kernel. Let (w_{ℓ}°) be a decreasing sequence in $M^b_+(\Omega)$ that converges to 0 uniformly on all compact subsets of Ω . For $\ell \in \mathbb{N}$, let w_{ℓ} be the minimal solution to $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$ in $M^b_+(\Omega)$.

Then there is some $\tilde{w} \in C^b_+(\Omega)$ such that $\tilde{w} = F(\tilde{w})$ and $w_\ell - w^\circ_\ell \to \tilde{w}$ uniformly on Ω ,

$$w_{\ell} \ge \tilde{w} + w_{\ell}^{\circ}, \qquad \ell \in \mathbb{N}$$

Further, the following holds:

If $\mathcal{R}_0 > 1$ and κ is topologically irreducible and all functions w_{ℓ}° are topologically positive, then \tilde{w} is strictly positive and $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$.

Proof. By Theorem 7.6, (w_{ℓ}) and $(w_{\ell} - w_{\ell}^{\circ})$ are decreasing sequences that converge to some $\tilde{w} \in M^b_+(\Omega)$ pointwise on Ω . Since κ is a strong Feller kernel, all $w_{\ell} - w_{\ell}^{\circ} = F(w_{\ell})$ are continuous. We have $w_{\ell} \geq \tilde{w} + w_{\ell}^{\circ}$ for all $\ell \in \mathbb{N}$. By Lebesgue's theorem of dominated convergence, $\tilde{w} = F(\tilde{w})$ and \tilde{w} is continuous because κ is a strong Feller kernel.

By Dini's lemma, $(w_{\ell} - w_{\ell}^{\circ})$ converges to \tilde{w} uniformly on every compact subset W of Ω . Since $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ uniformly on every compact subset W of Ω , $w_{\ell} \to \tilde{w}$ as $\ell \to \infty$ uniformly on every compact subset W of Ω . Since f is uniformly continuous,

$$\sup_{w} (f \circ w_{\ell} - f \circ \tilde{w}) \to 0, \qquad \ell \to \infty$$
(10.3)

for every compact subset W of Ω .

Let $\epsilon > 0$. Since κ is tight, there exists some compact subset W of Ω such that $\kappa(\Omega \setminus W, x) \leq \epsilon$ for all $x \in \Omega$. For all $\ell \in \mathbb{N}$ and $x \in \Omega$,

$$0 \le w_{\ell}(x) - w_{\ell}^{\circ}(x) - \tilde{w}(x)$$

$$\le \int_{W} \left(f(w_{\ell}(\xi)) - f(\tilde{w}(\xi)) \right) \kappa(d\xi, x) + \int_{\Omega \setminus W} f(w_{\ell}(\xi)) \kappa(d\xi, x)$$

By the properties of f,

$$\|w_{\ell} - w_{\ell}^{\circ} - \tilde{w}\|_{\infty} \leq \sup_{W} (f \circ w_{\ell} - f \circ \tilde{w}) \sup_{x \in \Omega} \kappa(\Omega, x) + \sup_{x \in \Omega} \kappa(\Omega \setminus W, x).$$

By (10.3),

$$\limsup_{\ell \to \infty} \|w_{\ell} - w_{\ell}^{\circ} - \tilde{w}\|_{\infty} \le \epsilon.$$

Since this hold for any $\epsilon > 0$, $||w_{\ell} - w_{\ell}^{\circ} - \tilde{w}||_{\infty} \to 0$ as $\ell \to \infty$.

Assume that $\mathcal{R}_0 > 1$ and all w_{ℓ}° are topologically positive. By Theorem 3.13

$$\|\tilde{w}\|_{\infty} = \lim_{\ell \to \infty} \|w_{\ell} - w_{\ell}^{\circ}\|_{\infty} \ge \ln \mathcal{R}_0, \qquad \ell \in \mathbb{N}.$$
(10.4)

Since \tilde{w} is continuous and κ is topologically irreducible, by Theorem 10.8, \tilde{w} is strictly positive on Ω .

Corollary 10.13. Let κ be a strong Feller kernel that is tight and topologically irreducible and let $\mathcal{R}_0 > 1$. Then there exists some strictly positive $\tilde{w} \in C^b_+(\Omega)$ such that $\tilde{w} = F(\tilde{w})$ and $\|\tilde{w}\|_{\infty} \geq \ln \mathcal{R}_0$.

Proof. Apply Theorem 10.12 with $w_{\ell}^{\circ} = 1/\ell$.

Corollary 10.14. Let κ be a strong Feller kernel that is tight and topologically irreducible and let $\mathcal{R}_0 > 1$.

Let w° be a topologically positive function in $M^b_+(\Omega)$ and w be the minimal solution of $w = F(w) + w^{\circ}$.

Then there exist strictly positive $\tilde{w} \in C^b_+(\Omega)$ with $\|\tilde{w}\|_{\infty} \ge \ln \mathcal{R}_0$ and

$$\tilde{w} \leq w - w^{\circ}.$$

Proof. Apply Theorem 10.12 with $w_{\ell}^{\circ} = (1/\ell) w^{\circ}$.

10.5 Comparability kernels

In Corollary 10.14, the fixed point $\tilde{w} = F(\tilde{w})$ may depend on w° . The Corollary would send a much stronger message if this were not the case. Further, we do not yet know what happens if $\mathcal{R}_0 = 1$. This leads to the concept of a comparability kernel (Definition 3.16). Cf. [48, (K-2)].

Proposition 10.15. Let κ be a comparability kernel and $g \in M^b_+(\Omega)$ be strictly positive and lower semicontinuous. Then there exists some $\delta > 0$ such that

$$\int_{\Omega} g(\xi) \kappa(d\xi, x) \ge \delta \, \kappa(\Omega, x), \qquad x \in \Omega$$

Proof. Let $g \in M^b_+(\Omega)$ be strictly positive and lower semicontinuous on Ω . By Lemma 10.2 (b), there exists a strictly positive Lipschitz continuous $\tilde{g} \in M^b_+(\Omega)$ such that $0 \leq \tilde{g} \leq g$.

Since κ is a comparability kernel, there exists some $\delta > 0$ such that

$$\delta\,\kappa(\Omega,x) \leq \int_\Omega \tilde{g}(\xi)\kappa(d\xi,x) \leq \int_\Omega g(\xi)\kappa(d\xi,x), \qquad x\in\Omega. \qquad \ \Box$$

Proposition 10.16. Assume that the Feller kernel κ is topologically irreducible and a comparability kernel and $w^{\circ} \in M^b_+(\Omega)$. Then, there exists at most one non-zero continuous solution $\tilde{w} \in M^b_+(\Omega)$ to

$$\tilde{w}(x) = \int_{\Omega} f(\tilde{w}(\xi) + w^{\circ}(\xi))\kappa(d\xi, x), \qquad x \in \Omega.$$

Proof. Assume that there are two, \tilde{w}_1 and \tilde{w}_2 . Since κ is topologically irreducible, by Theorem 10.8, both are strictly positive and so are $f \circ \tilde{w}_i$, i = 1, 2. By Proposition 10.15, there are $\delta_i > 0$ such that

$$\tilde{w}_i(x) \ge \delta_i \kappa(\Omega, x), \qquad x \in \Omega.$$

Since $\tilde{w}_i \leq \kappa(\Omega, \cdot)$, $t = \inf_{\Omega} \tilde{w}_1/\tilde{w}_2 > 0$ and $\tilde{w}_1(x) \geq t\tilde{w}_2(x)$ for all $x \in \Omega$. Suppose that $t \in (0, 1)$. Since f is increasing,

$$0 \ge \tilde{w}_1(x) - t\tilde{w}_2(x) \ge \int_{\Omega} \left[f(t\tilde{w}_2(\xi) + w^{\circ}(\xi)) - tf(\tilde{w}_2(\xi) + w^{\circ}(\xi)) \right] \kappa(d\xi, x).$$

Since \tilde{w}_2 is strictly positive and $t \in (0, 1)$ and f is increasing and strictly sublinear, for all $\xi \in \Omega$,

$$f(t\tilde{w}_{2}(\xi) + w^{\circ}(\xi)) - tf(\tilde{w}_{2}(\xi) + w^{\circ}(\xi))$$

$$\geq f(t(\tilde{w}_{2}(\xi) + w^{\circ}(\xi))) - tf(\tilde{w}_{2}(\xi) + w^{\circ}(\xi)) > 0.$$

Since the left hand side of this inequality is a continuous function of ξ , by Proposition 10.16, there is some $\delta > 0$ such that

$$\tilde{w}_1(x) - t\tilde{w}_2(x) \ge \delta\kappa(\Omega, x) \ge \delta\tilde{w}_2(x), \qquad x \in \Omega.$$

So, $\tilde{w}_1(x) \ge (t+\delta)\tilde{w}_2(x)$ for all $x \in \Omega$, contradicting the definition of t. This proves that $\tilde{w}_1 \ge \tilde{w}_2$. By symmetry, equality holds.

Theorem 10.17. Assume that the Feller kernel κ is topologically irreducible and a comparability kernel and $w^{\circ} \in C^{b}_{+}(\Omega)$. Then there exist at most one non-zero solution $w \in C^{b}_{+}(\Omega)$ to $w = F(w) + w^{\circ}$.

Proof. If w° is the zero function, the statement directly follows from Proposition 10.16. If w° is not the zero function, apply Proposition 10.16 to $\tilde{w} = w - w^{\circ} \ge 0$.

Theorem 10.18. Let κ be a Feller kernel that is topologically irreducible. Assume that κ is a comparability kernel and $\mathcal{R}_0 \leq 1$.

- (a) Then there exists no nonzero solution $\tilde{w} \in C^b_+(\Omega)$ to the equation $\tilde{w} = F(\tilde{w})$.
- (b) If, in addition, κ is a strong Feller kernel, there exists no nonzero solution $\tilde{w} \in M^b_+(\Omega)$ to the equation $\tilde{w} = F(\tilde{w})$.

(c) If κ is a strong Feller kernel and $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ is a sequence in $M^{b}_{+}(\Omega)$ and $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ pointwise on Ω , then $w_{\ell} \to 0$ as $\ell \to \infty$ pointwise on Ω for the minimal solutions w_{ℓ} of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

Proof. (a) Assume that such a solution \tilde{w} exists. Since κ is topologically irreducible, w is strictly positive on Ω by Theorem 10.8. Let t = 1/2. Then $f(t\tilde{w}(x)) - tf(\tilde{w}(x)) > 0$ for all $x \in \Omega$ and this difference is a continuous function of $x \in \Omega$. Since κ is a comparability kernel, there is some $\delta > 0$ such that

$$\begin{split} \delta\kappa(\Omega,x) &\leq \int_{\Omega} [f(t\tilde{w}(\xi)) - tf(\tilde{w}(\xi))]\kappa(d\xi,x) \\ &= F(t\tilde{w})(x) - tF(\tilde{w})(x), \qquad x \in \Omega. \end{split}$$

So there exists some $\epsilon > 0$ such that

$$F(t\tilde{w}) \ge (1+\epsilon)tF(\tilde{w}) = (1+\epsilon)t\tilde{w}.$$

By (2.7), $K\tilde{w} \ge (1+\epsilon)\tilde{w}$. This implies $\mathcal{R}_0 = \mathbf{r}(K) \ge (1+\epsilon)$ [51, Thm.3.1][54, Thm5.31], a contradiction.

(b) If $F(\tilde{w}) = w \in M^b_+(\Omega)$, and κ is a strong Feller kernel, then \tilde{w} is continuous.

(c) This follows from part (b) and Corollary 7.12.

Theorem 10.19. Let κ be a strong Feller kernel that is tight and topologically irreducible. Assume that κ is a comparability kernel and $\mathcal{R}_0 > 1$.

- (a) Then there exists a unique nonzero solution $\tilde{w} \in M^b_+(\Omega)$ to the equation $\tilde{w} = F(\tilde{w}); \tilde{w}$ is continuous, strictly positive and $\|\tilde{w}\| \ge \ln \mathcal{R}_0$.
- (b) For any topologically positive w° in $M^{b}(\Omega)$ and the minimal solution $w \in M^{b}_{+}(\Omega)$ of $w = F(w) + w^{\circ}$, we have $w w^{\circ} \geq \tilde{w}$ with the unique \tilde{w} from (b).
- (c) Finally, if $(w_{\ell}^{\circ})_{\ell \in \mathbb{N}}$ is a decreasing sequence of topologically positive functions in $M^b_+(\Omega)$ with $w_{\ell}^{\circ} \to 0$ as $\ell \to \infty$ uniformly on all compact subsets of Ω , then $w_{\ell} - w_{\ell}^{\circ} \to \tilde{w}$ uniformly on Ω for the solutions w_{ℓ} of $w_{\ell} = F(w_{\ell}) + w_{\ell}^{\circ}$.

Proof. (a) Since κ is a strong Feller kernel, any solution $\tilde{w} = F(\tilde{w})$ in $M^b_+(\Omega)$ is continuous. Uniqueness now follows from Theorem 10.17, existence from Corollary 10.13.

- (b) This follows from part (a) and Corollary 10.14.
- (c) Combine Theorem 10.12 and parts (a) and (b).

10.5.1 Examples of comparability kernels

Proposition 10.20. A Feller κ is a comparability kernel if there exist some compact subset W of Ω and some $\delta > 0$ such that

$$\kappa(W, x) \ge \delta \kappa(\Omega, x), \qquad x \in \Omega.$$

Proof. Let $g : \Omega \to (0, \infty)$ be continuous. Let W and $\delta > 0$ as in the statement of the proposition. Since W is compact, $\inf_W g > 0$. Further, for all $x \in \Omega$,

$$Kg(x) \ge \int_{W} g(\xi)\kappa(d\xi, x) \ge \inf_{w} g \ \kappa(W, x) \ge \inf_{W} g \ \delta \ \kappa(\Omega, x).$$

Proposition 10.21. Let κ be a tight Feller kernel and

$$\inf_{x \in \Omega} \kappa(\Omega, x) > 0. \tag{10.5}$$

Then κ is a comparability kernel.

Proof. By assumption, $\delta = \inf_{x \in \Omega} \kappa(\Omega, x) > 0$. Since κ is tight, there exists a compact subset W of Ω such that $\kappa(\Omega \setminus W, x) \leq \delta/2$ for $x \in \Omega$. Since $\kappa(\Omega, \cdot)$ is bounded on Ω , the assertion follows from Proposition 10.20.

Proposition 10.22. Let $n \in \mathbb{N}$ and $\nu_j, \mu_j \in \mathcal{M}_+(\Omega)$ be nonzero measures and $k_j \in M^b_+(\Omega)$ be nonzero functions, j = 1, ..., n, such that

$$\sum_{i=1}^{n} \mu_i(\omega) k_i(x) \le \kappa(\omega, x) \le \sum_{i=1}^{n} \nu_i(\omega) k_i(x), \qquad x \in \Omega, \quad \omega \in \mathcal{B}.$$
(10.6)

Then κ is a comparability kernel. Further, κ is tight if all ν_i are tight measures, e.g, if Ω is a Polish space (complete and separable).

Compare [48, Exp.1.3b]. It follows that every semi-separable Feller kernel, (3.6), is a comparability kernel.

Proof. By (10.6),

$$\kappa(\Omega, x) \le \sum_{i=1}^{n} \nu_i(\Omega) k_i(x) \le \sup_{i=1}^{n} \nu_i(\Omega) \sum_{i=1}^{n} k_i(x), \qquad x \in \Omega$$

Let $g \in M^b_+(\Omega)$ be strictly positive. By (10.6),

$$\int_{\Omega} g(\xi) \kappa(d\xi, x) \ge \sum_{i=1}^{n} \left(\int_{\Omega} g \, d\mu_i \right) k_i(x)$$

Since g is strictly positive on Ω , $\Omega = \bigcup_{\ell \in \mathbb{N}} \{g \ge 1/\ell\}$. Since $\mu_i \in \mathcal{M}_+(\Omega)$,

$$0 < \mu_i(\Omega) = \lim_{\ell \to \infty} \mu_i \big(\{ g \ge 1/\ell \} \big)$$

For sufficiently large $\ell \in \mathbb{N}$,

$$\int_{\Omega} g d\mu_i \ge \frac{1}{\ell} \mu_i \big(\{g \ge 1/\ell\} \big) > 0.$$

We combine these inequalities: For all $x \in \Omega$,

$$\int_{\Omega} g(\xi)\kappa(d\xi,x) \ge \inf_{i=1}^{n} \int_{\Omega} gd\mu_{i} \sum_{i=1}^{n} k_{i}(x) \ge \frac{\inf_{i=1}^{n} \int_{\Omega} gd\mu_{i}}{\sup_{i=1}^{n} \nu_{i}(\Omega)} \kappa(\Omega,x). \qquad \Box$$

References

- Aliprantis CD, Border KC (1999, 2006) Infinite Dimensional Analysis. A Hitchhiker's Guide, 3rd edition. Springer, Berlin Heidelberg
- [2] Bootsma MCJ, Chan KMD, Diekmann O, Inaba H (2024) The effect of host population heterogeneity on epidemic outbreaks. Math Appl Sci Eng 5:1-35 (https://doi.org/10.5206/mase/16718)
- [3] Bootsma MCJ, Chan KMD, Diekmann O, Inaba H (2023) Separable mixing: the general formulation and a particular example focusing on mask efficiency. Math Biosc Eng 20:17661-17671.
- [4] Busenberg S, Cooke KL (1993) Vertically Transmitted Diseases. Models and Dynamics. Springer, Berlin Heidelberg
- [5] Demongeot J, Griette Q, Maday Y, Magal P (2022) A Kermack-McKendrick model with age of infection starting from a single or multiple cohorts of infected patients. Proceedings of the Royal Society A, 479: 20220381 Supplementary material
- [6] Demongeot J, Magal P (2024) Data-driven mathematical modeling approaches for Covid-19: a survey. Physics of Life Reviews 50:166-208
- [7] Diekmann O (1978) Thresholds and travelling waves for the geographical spread of infection. J Math Biol 6:109-130
- [8] Diekmann O (2020) The 1927 epidemic model of Kermack and McKendrick: a success story or a tragicomedy? SMB Newsletter No. 92:1–4

- [9] Diekmann O, Heesterbeek H, Britton T (2013) Mathematical tools for Understanding Infectious Disease Dynamics. Princeton University Press, Princeton
- [10] Diekmann O, Inaba H (2023) A systematic procedure for incorporating separable static heterogeneity into compartmental epidemic models. *Journal of Mathematical Biology* 86:29
- [11] Diekmann O, Othmer HG, Planqué R, Bootsma MCJ (2021) The discrete-time Kermack–McKendrick model: A versatile and computationally attractive framework for modeling epidemics. PNAS 118 (39):e2106332118d
- [12] d'Onofrio A, Iannelli M, Manfredi P, Marinoschi G (2023) Optimal epidemic control by social distancing and vaccination of an infection structured by time since infection: the COVID-19 case study. Siam J Appl Math 00:S199-S224
- [13] d'Onofrio A, Iannelli M, Manfredi P, Marinoschi G (2024) Epidemic control by social distancing and vaccination: Optimal strategies and remarks on the COVID-19 Italian response policy. Math Biosc Eng 21:6493-6520
- [14] d'Onofrio A, Iannelli M, Marinoschi G, Manfredi P (2024) Multiple pandemic waves vs multi-period/multip-phasic epidemics: Global shape of the Covid-19 pandemic. J Theor Biol 593:111881
- [15] d'Onofrio A, Manfredi P, Iannelli M (2021) Dynamics of partially mitigated multi-phasicepidemics at low susceptible depletion: Phases of covid-19 control in Italy as case study. Math Biosci 340:108671.
- [16] Ducrot A, Griette Q, Liu Z, Magal P (2022) Differential equations and population dynamics I. Introductory approaches. With forewords by Jacques Demongeot and Glenn Webb. Lecture Notes on Mathematical Modelling in the Life Sciences. Springer, Cham.
- [17] Dudley RM (2002) Real Analysis and Probability, sec. ed.. Cambridge University Press, Cambridge.
- [18] Dunford N, Schwartz JT (1988) Linear Operators. Part I. General Theory. John Wiley, Classics Library Edition, New York.

- [19] Eikenberry S, Mancuso M, Iboi E, Phan T, Eikenberry K, Kuang Y, Kostelich E, Gumel A (2020) To mask or not to mask: Modeling the potential for face mask use by the general public to curtail the COVID-19 pandemic. Infect Dis Model 5:293-308
- [20] Farrell AP, Collins JP, Greer AL, Thieme HR (2018) Times from infection to disease-induced death and their influence on final population sizes after epidemic outbreaks. Bull Math Biol 80:1937-1961
- [21] Gomes MGM, Ferreira MU, Corder RM, King JG, Souto-Maior C, Penha-Gonçalves C, Gonçalves G, Chikina M, Pegden W, Aguas R (2022) Individual variation in susceptibility or exposure to SARS-CoV-2 lowers the herd immunity threshold. J Theor Biol 540:111063.
- [22] Iannelli M, Milner F (2017) The Basic Approach to Age-Structured Population Dynamics. Springer, Dordrecht
- [23] Inaba H (2017) Age-structured Population Dynamics in Demography and Epidemiology. Springer, Singapore
- [24] Inaba H (2023) Basic concepts for the Kermack and McKendrick model with static heterogeneity. arXiv:2311.11247, 26 pages.
- [25] Karlin S (1959) Positive operators. J. Math. Mech. 8: 907-937
- [26] Kermack WO, McKendrick AG (1927) A contribution to the mathematical theory of epidemics. Proc Roy Soc A 115:700-721; reprinted in Bull Math Biol 53 (1991):33-55
- [27] Kermack WO, McKendrick AG (1932) A contribution to the mathematical theory of epidemics II. The problem of endemicity. Proc Roy Soc A 138:55-85; reprinted in Bull Math Biol 53 (1991):57-87
- [28] Kermack WO, McKendrick AG (1933) A contribution to the mathematical theory of epidemics III. Further studies of the problem of endemicity. Proc Roy Soc A 141:94-122; reprinted in Bull Math Biol 53 (1991):89-118
- [29] Krasnosel'skij MA (1964) Positive Solutions of Operator Equations. Noordhoff, Groningen
- [30] Kreck M, Scholz E (2022) doi: 10.1007/s11538-022-00994-9. Back to the roots: a discrete Kermack-McKendrick model adapted to Covid-19. Bull Math Biol 84:44.

- [31] Krein MG, Rutman MA (1948) Linear operators leaving invariant a cone in a Banach space (Russian). Uspehi Mat. Nauk (N.S.) 3:3-95, English Translation, Amer Math Soc Trans 26 (1950)
- [32] Li X-Z, Yang J, Martcheva M (2020) Age Structured Epidemic Modeling. Interdisciplinary Applied Mathematics, 52. Springer, Cham
- [33] Liu Z, Magal P, Seydi O, Webb GF (2020), A COVID-19 epidemic model with latency period. Infectious Disease Modelling 5:323-337
- [34] Liu Z, Magal P, Webb GF (2021) Predicting the number of reported and unreported cases for the COVID-19 epidemics in China, South Korea, Italy, France, Germany and United Kingdom. J. Theoret. Biol. 509:Paper No. 110501, 10 pp.
- [35] Luckhaus S, Stevens A (2023) A free boundary problem -in time- for the spread of Covid-19. J Math Biol 86, Paper No. 45, 17 pp.
- [36] Luckhaus S, Stevens A (2023) Kermack and McKendrick models on a two-scale network and connections to the Boltzmann equations. Mathematics Going Forward. Collected Mathematical Brushstrokes (eds. J.-M. Morel and B. Teissier), Lecture Notes in Mathematics 2313:399 – 408.
- [37] Magal P, Ruan S (2018) Theory and Applications of Abstract Semilinear Cauchy Problems. With a foreword by Glenn Webb. Applied Mathematical Sciences, 201. Springer, Cham
- [38] Martcheva M (2015) An Introduction to Mathematical Epidemiology. Springer, New York
- [39] Ponce J, Thieme HR (2023) Can infectious diseases eradicate host species? The effect of infection-age structure. Math Biosci Eng 20:18717-18760.
- [40] Ponce J, Thieme HR (2024) A Kermack–McKendrick type epidemic model with double threshold phenomenon (and a possible application to Covid-19). arXiv:2409.17278
- [41] Rass L, Radcliffe J (2003) Spatial Deterministic Epidemics. AMS, Providence
- [42] Saldaña F, Velasco-Hernández JX (2022) Modeling the COVID-19 pandemic: a primer and overview of mathematical epidemilogy. SeMA J 79:225-251

- [43] Schaefer HH (1966) Topological Vector Spaces. The Macmillan Company, New York
- [44] Smith HL, Thieme HR (2011) Dynamical Systems and Population Persistence. Amer Math Soc, Providence
- [45] Tkachenko AV, Maslov S, Elbanna A, Wong GN, Weiner ZJ, Goldenfeld N (2021) Time-dependent heterogeneity leads to transient suppression of the COVID-19 epidemic, not herd immunity. PNAS Apr 2021, 118 (17) e2015972118; DOI: 10.1073/pnas.2015972118
- [46] Tkachenko AV, Maslov S, Wang T, Elbanna A, Wong GN, Goldenfeld N (2021) Stochastic social behavior coupled to COVID-19 dynamics leads to waves, plateaus, and an endemic state. eLife, https://doi.org/10.7554/eLife.68341
- [47] Thieme HR (1977) A model for the spatial spread of an epidemic. J Math Biol 4:337-351
- [48] Thieme HR (1979) On a class of Hammerstein integral equations. Manuscr math 29:49-84
- [49] Thieme HR (2003) Mathematical Population Biology. Princeton University Press, Princeton
- [50] Thieme HR (2016) Eigenfunctionals of homogeneous order-preserving maps with applications to sexually reproducing populations. J Dynamics Differential Equations 28:1115-1144
- [51] Thieme HR (2017) From homogeneous eigenvalue problems to two-sex population dynamics. J Math Biol 75:783-804
- [52] Thieme HR (2020) Discrete-time population dynamics on the state space of measures. Math Biosc Eng 17:1168-1217
- [53] Thieme HR (2020) Persistent discrete-time dynamics on measures. Progress on Difference Equations and Discrete Dynamical Systems (Stephen Baigent, Saber Elaydi and Martin Bohner, eds.), 59-100, Springer Proceedings in Mathematics & Statistics 341, Springer Nature Switzerland AG
- [54] Thieme HR (2024) Discrete-Time Dynamics of Structured Populations and Homogeneous Order-Preserving Operators. AMS, Providence