

On powers of the Diophantine function $\star : x \mapsto x(x + 1)$

Maxwell Lippmann* Donald Silberger†

February 19, 2025

Dedicated to Jan Mycielski, 1932 February 07 - 2025 January 23

Abstract

We treat the functions $\star^k : \mathbf{N} \rightarrow \mathbf{N}$ where $\star : x \mapsto \star x := x(x + 1)$. The set $\{\star^k x + 1 : \{x, k + 1\} \subset \mathbf{N}\}$ is pairwise coprime; so, the set \mathbf{P} of primes is infinite. Our Theorem 4 resorts to the *mother sequence*, \mathbf{M} , that is obtained by factoring the infinite integer-sequence $2, 3, 4, 5, \dots$ into prime powers.

For each $x \geq 1$ we define the *gross x -sequence*, $\gamma_\star(x) := \langle x + 1; \star x + 1; \star^2 x + 1; \star^3 x + 1; \dots \rangle$, and also the *star sequence*, x^\star , obtained by factoring the terms of $\gamma_\star(x)$ into prime powers. It turns out that $\gamma_\star(1)$ is Sylvester's sequence, A00058 in the *On-Line Encyclopedia of Integer Sequences*, OEIS, and that $\gamma_\star(2)$ is the sequence A082732 in the OEIS.

Theorem 3. For every integer $x \geq 1$ there is a prime $p(x)$ that divides no member of $\{\star^k x + 1 : k \geq 0\}$.

Theorem 4. For each sequence η of powers of primes there are infinitely many subsequences \mathbf{c}_j of \mathbf{M} such that numerically $\eta = \mathbf{c}_j$ but where the term-set family in \mathbf{M} of those \mathbf{c}_j is formally pairwise disjoint.

Theorem 6. $1/x = \sum_{k=0}^{n-1} 1/(\star^k x + 1) + 1/(\star^n x) = \sum_{k=0}^{\infty} 1/(\star^k x + 1)$ for all $\{x, n\} \subseteq \mathbf{N}$.

Theorem 7. For every $x \in \mathbf{N}$, when $x^\star := \langle x_j \rangle_{j=0}^{\infty}$ then $\sum_{j=0}^{\infty} 1/x_j = \infty$.

1 Introduction

Our initial interest in the elementary function $\star : \mathbf{N} \rightarrow \mathbf{N}$ defined by $\star : x \mapsto \star x := x(x + 1)$, where $\mathbf{N} := \{1, 2, 3, \dots\}$, derives from its appearance in what we have called in [4, 7] the Vital Identity

$$\frac{1}{z} = \frac{1}{z+1} + \frac{1}{z(z+1)} \quad \text{This identity holds for every complex number } z \notin \{-1, 0\}.$$

In [4, 7] the Vital Identity has facilitated proofs about segments of the harmonic series and more generally about sums of the reciprocals of positive integers. We define \star^0 to be the identity function and $\star^{k+1} := \star^k \circ \star$.

We recall Theorem 1 in [7] where the fact that two consecutive positive integers are necessarily coprime entails that for every pair $k \geq 0$ and $x \geq 1$ of integers the integer $\star^k x + 1$ has at least k distinct prime factors. This fact immediately yields Euclid's theorem that there exist infinitely many prime integers.¹

We are interested for each $x \in \mathbf{N}$ in two infinite sequences of integers. One is the *star sequence* x^\star ; its terms are powers of primes. Each such x^\star is uniquely determined and is created thus: First we produce the other sequence of interest to us; this is the *gross x -sequence*

$$\gamma_\star(x) := \langle x + 1; \star x + 1; \star^2 x + 1; \star^3 x + 1; \dots; \star^k x + 1; \dots \rangle = \langle \star^k x + 1 \rangle_{k=0}^{\infty}.$$

The terms of $\gamma_\star(x)$ are separated by semicolons. It is easy to see, but it also follows from Theorem 3 of [7], that $\gamma_\star(x)$ is injective, that it very rapidly increases and that the set of its terms is pairwise coprime.

We manufacture x^\star from $\gamma_\star(x)$ by factoring each term of $\gamma_\star(x)$ into a product of the powers of its primes and listing them separated by commas between successive semicolons in ascending order of prime sizes. Change the semicolons into commas. The integers separated by commas are the terms of $x^\star := \langle x_i \rangle_{i=1}^{\infty}$.

*max.lippmann@yahoo.com Roslyn High School Roslyn, NY 11576

†SilbergerD@newpaltz.edu or, preferably, DonaldSilberger@gmail.com State University of New York New Paltz, NY 12561

¹This inescapable observation arising from Sylvester's sequence has been noted by other people; e.g., R. W. K. Odoni mentions it in the second sentence of his 1984 paper [6].

Example One. The gross 1-sequence $\gamma_*(1)$ begins with $\langle 2; 3; 7; 43; 1807; 3263443; \dots \rangle$ and turns out to be Sylvester's sequence, A00058 in the *On-Line Encyclopedia of Integer Sequences - OEIS*. We compute that $1^* = \langle 2, 3, 7, 43, 13, 139, 3263443, \dots \rangle$; i.e., $1_0 = 2, 1_1 = 3; 1_2 = 7, 1_3 = 43, 1_4 = 13, 1_5 = 139, 1_6 = 3263443$. The OEIS says that it is an open question whether every term in $\langle w_k \rangle_{k=1}^\infty$ is square-free.

Example Two. We start the gross 2-sequence $\gamma_*(2) = \langle 3; 7; 43; 1807; 3263443; \dots \rangle$ and then we compute $2^* = \langle 3, 7, 43, 13, 139, 3263443, \dots \rangle$. Our $\gamma_*(2)$ occurs as A082732 in the *OEIS*.

Example Three. The gross 3-sequence begins with $\gamma_*(3) = \langle 4; 13; 157; 24493; 599882557; \dots \rangle$. We compute $3^* = \langle 2^2, 13, 157, 24493, 67, 277, 32323, \dots \rangle$

Notice that $\gamma_*(2)$ is a proper suffix of $\gamma_*(1)$. This implies that 2^* is a proper suffix of 1^* . These observations together with Theorem 3 in [7] give us

Proposition 1. *If $y \neq x$ and $y + 1$ is a term in the sequence $\gamma_*(x)$, then y^* is a proper suffix of x^* , and no prime that is a factor of a term of the prefix of x^* that is complementary to y^* is a factor of any term of y^* .*

Proposition 1 informs us that there are integers $x \geq 1$ for which there are primes that are factors of no member of $\{\star^k x + 1 : k \geq 0\}$. Theorem 3, below, extends this fact.

For $x \in \mathbf{N}$ we define the sequence $\eta(x) := \langle w_{x,j} \rangle_{j=1}^\infty$ recursively by

$$w_{x,k+1} = 1 + \prod_{j=0}^k w_{x,j} \quad \text{and} \quad w_{x,0} := x.$$

According to Odoni [6], whose paper's title contains it, $\eta(1)$ is Sylvester's sequence, A00058 in the OEIS.

Lemma 2. *Let $x \in \mathbf{N}$. The sequence $\eta(x) := \langle w_{x,j} \rangle_{j=1}^\infty$ is identical to the sequence $\gamma_*(x) := \langle \star^k x + 1 \rangle_{k=0}^\infty$.*

Proof. We show that a variant of the recursion defining $\eta(x)$ produces $\gamma_*(x)$. Observe that $\star^0 x + 1 = x + 1 = w_{x,1}$. Suppose for $k > 0$ that $w_{x,k} = \star^{k-1} x + 1$. Then $\star^k x + 1 := (\star^{k-1} x)(\star^{k-1} x + 1) + 1 = (w_{x,k} - 1)w_{x,k} + 1 = (w_{x,1}w_{x,2} \cdots w_{x,k-1})w_{x,k} + 1 = 1 + w_{x,1}w_{x,2} \cdots w_{x,k} = w_{x,k+1}$. The lemma follows by induction. \square

Theorem 3. *For every integer $x \geq 1$ there is a prime $p(x)$ that divides no member of $\{\star^k x + 1 : k \geq 0\}$.*

Proof. In his paper's first paragraph R. W. K. Odoni [6] shows that, if p is a prime that divides a term of $\eta(1)$, then either $p = 3$ or $p \equiv 1 \pmod{6}$. Therefore 11 divides no term w_k of $\eta(1)$, and hence by Lemma 2 we have that $p(1) := 11$ is a factor of no term of the sequence $\gamma_*(1) := \langle \star^j 1 + 1 \rangle_{j=0}^\infty$.

Let $1 < x \in \mathbf{N}$, and let $p(x)$ be any prime factor of x . Observe that x is coprime to every term of the sequence $\gamma_*(x)$ and therefore that $p(x)$ divides no term of $\gamma_*(x) := \langle \star^j x + 1 \rangle_{j=0}^\infty$. \square

2 Parallel embeddings

Our x^* are members of the class of sequences η whose terms are powers of prime integers. Each of these η occurs in an infinite assemblage of subsequences of the *mother sequence* $M := \langle m_0, m_1, m_2, \dots \rangle = \langle m_i \rangle_{i=0}^\infty$ in whose womb each η resides along with infinitely many pairwise formally disjoint identical siblings $c_j = \eta$.

The sequence M is created by listing the natural numbers greater than 1 in their normal ascending order but then replacing each of those integers with its normal-order factorization into powers of primes separated by commas. The following finite prefix of the infinite sequence M ought to eliminate misunderstandings:

$$M = \langle 2, 3, 2^2, 5, 2, 3, 7, 2^3, 3^2, 2, 5, 11, 2^2, 3, 13, 2, 7, 3, 5, 2^4, 17, 2, 3^2, 19, 2^2, 5, 3, 7, 2, 11, 23, 2^3, 3, 5^2, 2, 13, 3^3, \dots \rangle$$

Notice that $2 = m_0 = m_4 = m_9 = m_{15} = m_{21} = \dots$ and that $5 = m_3 = m_{10} = m_{18} = \dots$. That is, not only does M fail to be injective, but every prime power occurs infinitely often as a term in M . We call the symbols m_i and m_j *formally distinct* as terms of M if and only if $i \neq j$. We emphasize that m_i and m_j can be formally distinct even in the event that a numerical equality $m_i = m_j$ obtains. For instance, we say that m_0 "formally distinct" from m_4 although both m_0 and m_4 are numerically identical; i.e., $m_0 = 2 = m_4$.

We call a collection \mathbf{H} of subsequences of M *pairwise formally disjoint* - aka *PFJ* - if and only if, for every pair $\{c, d\} \subset \mathbf{H}$, whenever $\langle m_i, m_j \rangle \in \mathcal{T}(c) \times \mathcal{T}(d)$ then $i \neq j$, where $\mathcal{T}(c)$ is the term set of c .

By a *parallel embedding* of a sequence η of powers of primes we mean an infinite PFJ collection of subsequences of M each of which is term by term a numerical copy of η .

Theorem 4. *Every sequence η of powers of primes has a parallel embedding.*

Proof. We leave to the reader the very easy proof that every finite η has a parallel embedding, and prove the theorem for infinite η using the paradigm argument pertaining to $\eta := x^*$ for an arbitrary $x \in \mathbf{N}$. Recall from Section 1 the terminology $x^* := \langle x_0, x_1, x_2, \dots \rangle$. We will recursively develop the promised parallel embedding $\mathbf{S}(x) := \{\mathbf{c}_0, \mathbf{c}_1, \mathbf{c}_2, \dots\}$ consisting of subsequences \mathbf{c}_j of \mathbf{M} that term-by-term satisfy the numerical identity $\mathbf{c}_j = x^*$ for all $j \geq 0$. In order for our procedure to build all of the $\mathbf{c}_j \in \mathbf{S}(x)$ and not merely a solitary one such infinite sequence, we shall resort to the trick used to prove Theorem 2 in [7] and reminiscent of the corner-slicing gimmick employed by Georg Cantor [1] in his proof that the set \mathbf{Q}^+ is countable.

We intend each \mathbf{c}_i to be a sequence $\mathbf{c}_i := \langle m_{i_0}, m_{i_1}, m_{i_2}, \dots, m_{i_j}, \dots \rangle$ of powers of primes for which $m_{i_j} = x_j$ is a numerical fact for every $j \geq 0$. Let 0_0 be the least index t for which $m_t = x_0$. The term m_{0_0} was just now ‘‘captured from \mathbf{M} .’’ For our purposes in building the set $\mathbf{S}(x)$ of subsequences of \mathbf{M} , our designation of m_{0_0} as ‘‘captured’’ in effect removes m_{0_0} as an available term from the mother sequence \mathbf{M} .

Next, let 0_1 be the smallest integer t such that m_t is not yet captured and also such that $m_t = x_1$. We have thus far chosen the first two terms, m_{0_0} and m_{0_1} , of the subsequence \mathbf{c}_0 . Our third choice will be the first term m_{1_0} of the subsequence \mathbf{c}_1 ; here 1_0 is the smallest integer t for which both m_t is uncaptured and $m_t = x_0$. Three subsequential terms have been designated and thus captured. The order in which we select (and capture) terms of \mathbf{M} for subsequences destined to comprise $\mathbf{S}(x)$ is easily recognized from this pattern:

$$m_{0_0} \rightarrow m_{0_1} \rightarrow m_{1_0} \rightarrow m_{0_2} \rightarrow m_{1_1} \rightarrow m_{2_0} \rightarrow m_{0_3} \rightarrow m_{1_2} \rightarrow m_{2_1} \rightarrow m_{3_0} \rightarrow m_{0_4} \rightarrow m_{1_3} \rightarrow m_{2_2} \rightarrow \dots$$

As a paradigm step in our subsequence term-choosing procedure, we now choose the second term of the subsequence \mathbf{c}_3 . Let 3_1 be the smallest integer t for which $m_t = x_1$ but also for which m_t is not yet captured. We capture m_{3_1} from \mathbf{M} to serve as the the second term, m_{3_1} , in the subsequence \mathbf{c}_3 .

It is clear for every $\langle u, v \rangle \in \omega \times \omega$ that², after a finite number of procedural steps of the sort described in the preceding three paragraphs, the $(v+1)^{\text{st}}$ term, m_{u_v} , of the subsequence \mathbf{c}_u will be captured and put into its place in \mathbf{c}_u and that $m_{u_v} = x_v$. Moreover, for each pair $p \neq q$ of nonnegative integers, our refusal to re-use any previously captured term of \mathbf{M} , that term’s having already been assigned to be a term of some \mathbf{c}_r , guarantees that the set of terms of the subsequence \mathbf{c}_p is formally disjoint from the set of terms of \mathbf{c}_q . \square

Corollary 5. *Every copy of \mathbf{M} has a parallel embedding the family of whose term sets is a partition of \mathbf{M} .*

3 Sequence sums

In [4, 7] the function $\sigma : \mathcal{P}(\mathbf{N}) \setminus \{\emptyset\} \rightarrow \mathbf{Q}^+ \cup \{\infty\}$ is defined by $\sigma : X \mapsto \sigma X := \sum_{y \in X} 1/y$, where $\mathcal{P}(\mathbf{N})$ is the family of sets of positive integers.

Define $\mathbf{T}(x)$ to be the set of terms x_i of x^* . Remember that each such x_i is the power of a prime.

Our next result generalizes a fact about Sylvester’s sequence, i.e., the case $x = 1$.

Theorem 6. *Let $x \geq 1$. Then $1/x = \sigma \bigcup_{k=0}^n \{\star^k x + 1\} + 1/\star^n x$ for all $n \in \mathbf{N}$ and $1/x = \sigma \bigcup_{k=0}^{\infty} \{\star^k x + 1\}$.*

Proof. Since $\star^k x \notin \{-1, 0\}$ when $k \geq 0$, by the Vital Identity we have that

$$\begin{aligned} \frac{1}{x} &= \frac{1}{x+1} + \frac{1}{\star x} =: \frac{1}{\star^0 x + 1} + \frac{1}{\star^1 x} = \frac{1}{\star^0 x + 1} + \frac{1}{\star^1 x + 1} + \frac{1}{\star^2 x} = \frac{1}{\star^0 x + 1} + \frac{1}{\star^1 x + 1} + \frac{1}{\star^2 x + 1} + \frac{1}{\star^3 x} = \\ &\quad \frac{1}{\star^0 x + 1} + \frac{1}{\star^1 x + 1} + \frac{1}{\star^2 x + 1} + \frac{1}{\star^3 x + 1} + \frac{1}{\star^4 x} = \dots = \sum_{k=0}^{n-1} \frac{1}{\star^k x + 1} + \frac{1}{\star^n x}. \end{aligned}$$

The first claim is proved. The second follows from it together with the fact that $\lim_{n \rightarrow \infty} 1/\star^n x = 0$. \square

In the light of Theorem 6, of the paucity of factors that occur in each of the first few terms of the sequences $\gamma_\star(x)$ at which we have looked, and also of the swiftly burgeoning sizes of the primes encountered there, one might wager that $\sigma \mathbf{T}(x) < \infty$. However, there are reasons to hedge this bet.

One such reason is that $1/xy < (x+y)/xy = 1/x + 1/y$ with $x > 1$ and $y > 1$. So, if factors-rich terms of the sort $\star^k x + 1 = p_{i_1}^{e_1} p_{i_2}^{e_2} \dots p_{i_v}^{e_v}$ occur frequently in $\gamma_\star(x)$, then such numbers as $1/(\star^k x + 1)$ that are

²Following the custom of set theoreticians we define $\omega := \{0\} \cup \mathbf{N}$.

summands in the sum treated in Theorem 6 get replaced by significantly larger numbers $1/p_{i_1}^{e_1} + \dots + 1/p_{i_v}^{e_v}$ that are partial sums in $\sigma\mathbb{T}(x)$. This makes the convergence of $\sigma\mathbb{T}(x)$ begin to seem questionable. For, if among the terms of $\gamma_*(x)$ there is an increasing manifestation of factors-rich terms as k increases, then $\sigma \cup \{\star^k x + 1 : k \geq 0\} \ll \sigma\mathbb{T}(x)$. Indeed, $\sigma\mathbb{T}(x) = \infty$ appears possible.

On the other hand, since huge primes do occur early as terms in some sequences x^* , possibly prime-power terms of x^* tend to increase in size swiftly enough to entail rapidly diminishing summands in $\sigma\mathbb{T}(x)$ so as to support convergence, $\sigma\mathbb{T}(x) < \infty$, after all.

So, what actually happens?

Theorem 7. $\sigma\mathbb{T}(x) = \infty$ for every integer $x \geq 1$.

Proof. We rely upon the Second Theorem of Franz Mertens [5] and upon Theorem 2 of R. W. K. Odoni [6] together with Section 8 of Odoni's paper where he observes that the provisions of his Theorem 2 that are met by $\gamma_*(1)$ hold also for the sequences $\gamma_*(x)$ with $x > 1$.

The terms of $x^* := \langle x_j \rangle_{j=0}^\infty$ are powers of primes. We write x_j^- to designate the prime factor of x_j .

The Mertens theorem estimates the size of the sum of the reciprocals of all primes less or equal to n by

$$\lim_{n \rightarrow \infty} \left(\sum_{p \leq n} \frac{1}{p} - \log \log n - M \right) = 0 \quad \text{where } M := 0.26149721 \dots \text{ is the Meissel-Mertens constant.}$$

The Odoni theorem estimates the number $\mathbf{P}_1(n)$ of distinct prime factors less than or equal to n of terms, $\star^k 1 + 1$ of the sequence $\gamma_*(1)$, by asserting that

$$\mathbf{P}_1(n) = O\left(\frac{n}{(\log n)(\log \log \log n)}\right) \quad \text{as } n \rightarrow \infty.$$

Section 8 of the Odoni paper claims that the foregoing assertion about $\gamma_*(1)$ applies to $\gamma_*(x)$ for all $x \geq 1$ and consequently extends our statements here about the sequence 1^* to the sequences x^* as well.

Since the Prime Number Theorem estimates $n/\log n$ to be the number $\pi(n)$ of primes no greater than n as $n \rightarrow \infty$, the Odoni paper implies that the proportion of $\pi(n)$ of those such primes that occur as factors of terms of $\gamma_*(x)$ is asymptotic to $1/(\log \log \log n)$. Using also the Merten theorem we infer that

$$\sum_{p \in \mathbf{P}_x(n)} \frac{1}{p} \approx \frac{\log \log n + M}{\log \log \log n} \quad \text{whence } \sum_{j=0}^{\infty} \frac{1}{x_j^-} = \sum_{p \in \mathbf{P}_x} \frac{1}{p} = \infty \quad \text{since } \lim_{n \rightarrow \infty} \frac{\log \log n + M}{\log \log \log n} = \infty, \quad \text{where}$$

\mathbf{P}_x is the set of all prime factors p of terms of $\gamma_*(x)$ and $\mathbf{P}_x(n)$ is the set of such p no larger than n . Thus although we have obtained the tightening

$$\sum_{j=0}^{\infty} \frac{1}{x_j^-} = \infty$$

of Leonhard Euler's 1937 theorem [2] that $\sum_{p \in \mathbf{P}} 1/p = \infty$, we cannot yet infer as ultimately desired that

$$\sigma\mathbb{T}(x) := \sum_{j=0}^{\infty} \frac{1}{x_j} = \infty.$$

We must dismiss the chance that $x_j = x_j^-$ happens too seldom to compel $\sum_{j \in \omega} 1/x_j$ to diverge. That this might threaten our theorem is suggested by Euler's Basel problem solution [3], $\sum_{n \in \mathbf{N}} 1/n^2 = \pi^2/6$.

We will use probability theory to make an end run around this ostensible hazard we have noted.

It is evident that $\sigma\mathbb{T}(x) = \infty$ if and only if, for every probability $r < 1$ and every real number ℓ , there is an index b such that the probability is greater than or equal to r that $\sum_{j=b}^{\infty} 1/x_j > \ell$.

Since no prime divides more than one term x_j of the sequence x^* , no prime power can recur with an increased exponent. Moreover, among all multiples $z \in \mathbf{N}$ of a prime p , the probability that $p^2|z$ is equal to $1/p$ whence the probability that $\neg(p^2|z)$ is equal to $1 - 1/p = (p - 1)/p$.

For each $v \in \mathbf{N}$ there is an index b such that the smallest prime dividing any term in the suffix subsequence $\langle x_{b+i} \rangle_{i=0}^\infty$ of x^* exceeds v . So, for $r < 1$ a probability, there exists $b \in \mathbf{N}$ with $r < (x_j^- - 1)/x_j^- < 1$ for

every $j \geq b$. Hence, for some $m > b$ we have that the fraction, $f(b, m)$ of the number of terms of the sort $x_j = x_j^-$ divided by the number $m - b + 1$, is greater than r . Thus, $\liminf_{m \rightarrow \infty} f(b, m) \geq r$. It follows, for an arbitrary real number ℓ , that there exists b for which the probability is greater than or equal to r that $\sum_{j=b}^{\infty} 1/x_j > \ell$. We conclude that $\sigma T(x) = \infty$. \square

The proof of Theorem 7 gives us the following fact gratis.

Corollary 8. For all $x \in \mathbf{N}$ the approximation $\sum_{p \in \mathbf{P}_x(n)} \frac{1}{p} \approx \pi(\log \log n)$ obtains as $n \rightarrow \infty$.

Seeking a boundary between convergence and divergence we pose two questions pertaining to each $x \in \mathbf{N}$:

One. If $\mathbf{b}_{x,k}$ is the largest prime with $\mathbf{b}_{x,k}^{e_{x,k}} \parallel \star^k x + 1$ for $e_{x,k} > 0$, then $\sigma\{\mathbf{b}_{x,k}^{e_{x,k}} : k \geq 0\} < \infty$?

Two. If $\mathbf{d}_{x,k}$ is the smallest prime with $\mathbf{d}_{x,k}^{f_{x,k}} \parallel \star^k x + 1$ for $f_{x,k} > 0$, is $\sigma\{\mathbf{d}_{x,k}^{f_{x,k}} : k \geq 0\} < \infty$?

Googling Sylvester's sequence led us to Wikipedia which offers hundreds of related references, and it was Wikipedia that apprised us of the valuable *On-Line Encyclopedia of Integer Sequences* where we found that our $\gamma_*(1)$ had been discovered in another form by James Joseph Sylvester in 1880. Indeed, Euclid may have known about that sequence two and one-third millennia ago.

Both $\gamma_*(1)$ and $\gamma_*(2)$ (respectively A00058 and A082732 in the *OEIS*) have number-theoretic significance. Each term of $\gamma_*(1)$ is 1 plus the product of the terms that precede it and each term of $\gamma_*(2)$ is 1 plus the least common multiple of the terms preceding it. The *OEIS* says also that, starting with the fifth term of the sequence $\gamma_*(2)$, the ultimate two-digit suffixes of its terms alternate perpetually between 57 and 93.

Are there number-theoretic interpretations of $\gamma_*(x)$ for other values of x besides $x = 1$ and $x = 2$? For x a prime power, what patterns emerge for the $\gamma_*(x)$ and the x^* ? What happens when x is square-free?

The set $\{\star^k(1 + \prod \mathbf{P}_{\text{known}}) : k \geq 0\}$ supplements $1 + \prod \mathbf{P}_{\text{known}}$ as a source for undiscovered primes.

References

- [1] Georg Cantor, *Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen*, Crelle's Journal, (1874)
- [2] Leonhard Euler, *Variae observationes circa series infinitas*, Commentarii Academiae Scientiarum Petropolitanae, **9** (1737), 160-188.
- [3] Leonhard Euler, *De summis serierum reciprocarum*, Commentarii Academiae Scientiarum Imperialis Petropolitanae, **7** (1740), 123-134.
- [4] David Hobby, Donald Silberger, Sylvia Silberger, *Sums of finitely many distinct rationals*, arXiv:1702.01316v2[math.NT] 19 Feb 2019, 1-12.
- [5] Franz Mertens, *Ein Beitrag zur analitischen Zahlentheorie*, J. reine angew. Math. **78** (1874) 46-62.
- [6] R. W. K. Odoni, *On the prime divisors of the sequence $w(n+1) = 1 + w_1 + \dots + w_n$* , J. London Math. Soc. (**2**) 32 (1985) 1-11.
- [7] Donald Silberger, *On sums of Egyptian fractions*, arXiv:2412.02049[math.NT] 2 Dec 2024, 1-7.

Acknowledgments. Kira Adaricheva, Evan O'Dorney and Allan J. Silberger offered us helpful counsel.

2020 Mathematics Subject Classification: 01A55, 01A60, 11A25, 11N05, 11N25