Modern Physics Letters A © World Scientific Publishing Company

# Inhomogeneous transformations in a gauged twistor formulation of a massive particle

SHINICHI DEGUCHI

Institute of Quantum Science, College of Science and Technology, Nihon University, Chiyoda-ku, Tokyo 101-8308, Japan deguchi.shinnichi@nihon-u.ac.jp

SATOSHI OKANO\*

Department of Liberal Arts and Basic Sciences, College of Industrial Technology, Nihon University, Narashino, Chiba 275-8576, Japan okano.satoshi@nihon-u.ac.jp

> Received Day Month Year Revised Day Month Year

In this paper, we show that the mass-shell constraints in the gauged twistor formulation of a massive particle given in [Deguchi and Okano, *Phys. Rev. D* **93**, 045016 (2016) [Erratum **93**, 089906(E) (2016)]] are incorporated in an action automatically by extending the local U(2) transformation to its inhomogeneous extension denoted by IU(2). Therefore, it turns out that all the necessary constraints are incorporated into an action by virtue of the local IU(2) symmetry of the system.

Keywords: Twistor; massive particles; gauge symmetries

PACS Nos.: 11.10.Ef, 11.30.Ly, 11.90.+t

# 1. Introduction

In the mid to late 1970's, Penrose, Perjés, and Hughston independently studied the twistor description of a massive spinning particle in four-dimensional Minkowski space.<sup>1–7</sup> In their studies,  $n(\geq 2)$  twistors are introduced and an inhomogeneous extension of SU(n), which is denoted by ISU(n), is found as an internal symmetry group associated with massive particles. (See Ref. 8 for recent topics concerning a twistor description of massive particles.)

In the early 2000's, Lagrange mechanics of a massive spinning particle formulated using two twistors has been studied from both classical and quantum mechanical points of view.<sup>9–17</sup> In earlier formulations, however, all the necessary constraints are

<sup>\*</sup>Corresponding author

incorporated into actions of a massive particle by hand with the use of appropriate Lagrange multipliers.

About 8 years ago from now, the present authors gave a gauged twistor formulation of a massive spinning particle in four dimensions.<sup>18</sup> In this formulation, the necessary constraints except mass-shell constraints are automatically incorporated in an action by virtue of the gauge symmetry under a local U(2) transformation. In contrast, the mass-shell constraints given here are still incorporated into the action by hand.

The purpose of this paper is to show that the mass-shell constraints can also be incorporated in the action automatically by extending the local U(2) transformation to its inhomogeneous extension denoted by IU(2). In this way, it turns out that all the necessary constraints are incorporated in the action on the basis of the local IU(2) symmetry of the system.

This paper is organized as follows. In Sec. 2, we review the gauged twistor model of a massive spinning particle presented in Ref. 18. In Sec. 3, we show that the mass-shell constraints are certainly derived by inhomogeneous extensions of the transformation rules of twistor and some associated variables. Section 4 is devoted to conclusions. In Appendix A, we derive geometric formulas for the coset space SU(2)/U(1).

# 2. Gauged twistor model of a massive spinning particle in four dimensions (A review)

Penrose, Perjés, and Hughston introduced two or more independent twistors to describe a massive particle in four-dimensional Minkowski space, **M**. About 7 years ago from now, it was proven that the  $n(\geq 2)$ -twistor expression of four-momentum vector of a massive particle reduces to the two-twistor expression of it by a unitary transformation.<sup>19</sup> Taking into account this fact, we now introduce two independent twistors  $Z_i^A = (\omega_i^{\alpha}, \pi_{i\dot{\alpha}})$   $(A = 0, 1, 2, 3; \alpha = 0, 1; \dot{\alpha} = \dot{0}, \dot{1})$  distinguished by the index i (i = 1, 2) and their dual twistors  $\bar{Z}_A^i = (\bar{\pi}_{\alpha}^i, \bar{\omega}^{i\dot{\alpha}})$ . Here,  $\bar{\pi}_{\alpha}^i$  and  $\bar{\omega}^{i\dot{\alpha}}$  denote the complex conjugates of  $\pi_{i\dot{\alpha}}$  and  $\omega_i^{\alpha}$ , respectively:  $\bar{\pi}_{\alpha}^i := \bar{\pi}_{i\dot{\alpha}}, \ \bar{\omega}^{i\dot{\alpha}} := \bar{\omega}_i^{\alpha}$ . It is assumed that  $Z_1^A$  and  $Z_2^A$  are not proportional to each other, i.e.,  $Z_1^A \neq cZ_2^A$   $(c \in \mathbb{C})$ , and hence  $\bar{Z}_A^1 \neq \bar{c}Z_A^2$ . The spinors  $\omega_i^{\alpha}$  and  $\pi_{i\dot{\alpha}}$  are related by  $\omega_i^{\alpha} = i z^{\alpha\dot{\alpha}} \pi_{i\dot{\alpha}}$  with  $z^{\alpha\dot{\alpha}}$  being coordinates of a point in complexified Minkowski space,  $\mathbb{C}M$ . In addition to the twistors and their dual twistors, we introduce an inhomogeneous coordinate  $\xi (\in \mathbb{C})$  of a point on the coset space  $\mathcal{C} := SU(2)/U(1)$ . The SU(2) symmetry considered here is linearly realized for the pair of twistors  $(Z_1^A, Z_2^A)$ . (For the nonlinear realization of SU(2), see Appendix A.)

Now, we recall the action for a massive spinning particle found in Ref. 18:

$$S = \int_{\tau_0}^{\tau_1} d\tau \left[ \frac{i}{2} \left( \bar{Z}_A^i D Z_i^A - Z_i^A \bar{D} \bar{Z}_A^i \right) - 2sa - 2t \left( b^r \mathcal{V}_r^3 - \dot{\xi} e_{\xi}^3 - \dot{\bar{\xi}} e_{\bar{\xi}}^3 \right) - \frac{1}{e} g_{\xi\bar{\xi}} D\xi D\bar{\xi} - \frac{k^2}{2} \mathbf{e} + h \left( \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_j^{\dot{\alpha}} - \sqrt{2} m e^{i\varphi} \right) + \bar{h} \left( \epsilon_{ij} \bar{\pi}_{\alpha}^i \bar{\pi}^{j\alpha} - \sqrt{2} m e^{-i\varphi} \right) \right], \qquad (2.1)$$

with

$$DZ_i^A := \dot{Z}_i^A - iaZ_i^A - ib_i{}^j Z_j^A, \qquad (2.2a)$$

$$D\overline{Z}_A^i := \overline{Z}_A^i + ia\overline{Z}_A^i + i\overline{Z}_A^j b_j^{\ i} , \qquad (2.2b)$$

$$D\xi := \xi - b^r K_r^{\xi} , \qquad (2.2c)$$

$$D\bar{\xi} := \bar{\xi} - b^r K_r^{\ \xi} . \tag{2.2d}$$

Here,  $Z_i^A$ ,  $\overline{Z}_A^i$ , and  $\xi$  are understood to be complex scalar fields on the onedimensional parameter space  $\mathcal{T} := \{\tau | \tau_0 \leq \tau \leq \tau_1\}$  of a particle's worldline,  $\varphi$  is a real scalar field on  $\mathcal{T}$ , h is a complex scalar-density field of weight 1 on  $\mathcal{T}$ ,  $\mathbf{e}$  is a positive real scalar-density field of weight 1 on  $\mathcal{T}$ , and a and  $b^r$  (r = 1, 2, 3) are real scalar-density fields of weight 1 on  $\mathcal{T}$ . The field a is regarded as a one-dimensional U(1) gauge field, while the fields  $b^r$  are regarded as one-dimensional SU(2) gauge fields. The  $b_i{}^j$  (i, j = 1, 2) are the matrix elements of the matrix  $b := \sum_{r=1}^3 b^r \sigma_r$ defined using the Pauli matrices  $\sigma_r$ . The field  $\mathbf{e}$  plays the role of an einbein field. The specific forms of  $\mathcal{V}_r{}^3$ ,  $e_{\xi}{}^3$ ,  $e_{\overline{\xi}}{}^3$ , and  $g_{\xi\overline{\xi}}$  are given in Eqs. (A.10c), (A.5a), (A.5b), and (A.6b), respectively (see Appendix A). The three pairs of  $K_r{}^{\xi}$  and  $K_r{}^{\overline{\xi}}$  distinguished by r constitute the SU(2) Killing vectors on  $\mathcal{C}$ . In addition, m is a constant mass parameter, s and t are real constants, and k is a positive real constant. A dot over a field denotes its derivative with respect to  $\tau$ .

The action (2.1) remains invariant under the reparametrization  $\tau \to \tau' = \tau'(\tau)$ . In addition, this action remains invariant under the local U(1) transformation

$$Z_i^A \to Z_i^{\prime A} = e^{i\theta(\tau)} Z_i^A \,, \tag{2.3a}$$

$$\bar{Z}_A^i \to \bar{Z}_A^{\prime i} = e^{-i\theta(\tau)} \bar{Z}_A^i \,, \tag{2.3b}$$

$$\xi \to \xi' = \xi \,, \tag{2.3c}$$

$$\bar{\xi} \to \bar{\xi}' = \bar{\xi},$$
(2.3d)

$$h \to h' = e^{-2i\theta(\tau)}h$$
, (2.3e)

$$\bar{h} \to \bar{h}' = e^{2i\theta(\tau)}\bar{h}\,,\tag{2.3f}$$

$$\varphi \to \varphi' = \varphi + 2\theta(\tau),$$
 (2.3g)

$$a \to a' = a + \theta$$
, (2.3h)

$$b \to b' = b \,, \tag{2.3i}$$

$$\mathbf{e} \to \mathbf{e}' = \mathbf{e} \,, \tag{2.3j}$$

with a real transformation parameter  $\theta(\tau)$  satisfying  $\theta(\tau_1) = \theta(\tau_0)$ . Furthermore, the action (2.1) remains invariant under the local SU(2) transformation

$$Z_i^A \to Z_i^{\prime A} = U_i^{\ j}(\tau) Z_j^A \,, \tag{2.4a}$$

$$\bar{Z}_A^i \to \bar{Z}_A^{\prime i} = \bar{Z}_A^j U^{\dagger}{}_j{}^i(\tau) \,, \tag{2.4b}$$

$$\xi \to \xi' = \xi'(\xi) \,, \tag{2.4c}$$

$$\bar{\xi} \to \bar{\xi}' = \bar{\xi}'(\bar{\xi}),$$
(2.4d)

$$h \to h' = h \,, \tag{2.4e}$$

$$\bar{h} \to \bar{h}' = \bar{h} \,, \tag{2.4f}$$

$$\varphi \to \varphi' = \varphi$$
, (2.4g)

$$a \to a' = a \,, \tag{2.4h}$$

$$b \to b' = UbU^{\dagger} - i\dot{U}U^{\dagger},$$
 (2.4i)

$$\mathbf{e} \to \mathbf{e}' = \mathbf{e} \,, \tag{2.4j}$$

with a transformation matrix  $U(\tau) \in SU(2)$  satisfying  $U(\tau_1) = U(\tau_0)$ .

Now, we take the unitary gauge specified by V = 1, where V denote coset representatives of the coset space SU(2)/U(1). The condition V = 1 leads to  $\xi = \bar{\xi} = 0$  (see Appendix A). Then, as shown in Appendix A,  $\mathcal{V}_r{}^s = \delta_r{}^s$ ,  $e_{\xi}{}^3 = e_{\bar{\xi}}{}^3 = 0$ ,  $g_{\xi\bar{\xi}} = 1/2$ , and Eq. (A.13) hold, and hence we have  $D\xi = -b^-$  and  $D\bar{\xi} = -b^+$ for  $b^{\pm} := b^1 \mp ib^2$ . (Here, note that b can be expanded as  $b = \sum_{r=+,-,3} b^r \sigma_r$  with  $\sigma_{\pm} := (\sigma_1 \pm i\sigma_2)/2$ .) Hence, in the unitary gauge, the action (2.1) becomes

$$S = \int_{\tau_0}^{\tau_1} d\tau \left[ \frac{i}{2} \left( \bar{Z}_A^i D Z_i^A - Z_i^A \bar{D} \bar{Z}_A^i \right) - 2sa - 2tb^3 - \frac{1}{2e} b^{\hat{\imath}} b^{\hat{\imath}} - \frac{k^2}{2} e + h \left( \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_j^{\dot{\alpha}} - \sqrt{2}m e^{i\varphi} \right) + \bar{h} \left( \epsilon_{ij} \bar{\pi}_{\alpha}^i \bar{\pi}^{j\alpha} - \sqrt{2}m e^{-i\varphi} \right) \right],$$
(2.5)

where  $b^{\hat{i}}b^{\hat{i}} := (b^1)^2 + (b^2)^2 = b^+b^-$  ( $\hat{i} = 1, 2$ ). In Ref. 18, this action was referred to as the gauged generalized Shirafuji (GGS) action. In the unitary gauge, the local SU(2) invariance of S is spoiled, while S remains invariant under another local U(1)transformation

$$Z_i^A \to Z_i^{\prime A} = \Theta_i{}^j(\tau) Z_j^A \,, \tag{2.6a}$$

$$\bar{Z}_A^i \to \bar{Z}_A^{\prime i} = \bar{Z}_A^j \Theta^{\dagger}{}_j{}^i(\tau) \,, \qquad (2.6b)$$

$$h \to h' = h \,, \tag{2.6c}$$

$$h \to h' = h \,, \tag{2.6d}$$

$$\varphi \to \varphi' = \varphi \,, \tag{2.6e}$$

$$a \to a' = a \,, \tag{2.6f}$$

$$b \to b' = \Theta b \Theta^{\dagger} + \vartheta \sigma_3 , \qquad (2.6g)$$

$$\mathbf{e} \to \mathbf{e}' = \mathbf{e} \,, \tag{2.6h}$$

with the matrix  $\Theta(\tau) := \exp\{i\vartheta(\tau)\sigma_3\}$ . Here,  $\vartheta(\tau)$  is a real transformation parameter satisfying  $\vartheta(\tau_1) = \vartheta(\tau_0)$ . The local SU(2) invariance of S is thus converted to the invariance under the local U(1) transformation (2.6) by means of the gauge fixing such that V = 1. To avoid confusion, we hereafter refer to the transformation (2.3) as the  $U(1)_a$  transformation and refer to the transformation (2.6) as the  $U(1)_b$ transformation. Their corresponding gauge groups are simply denoted as  $U(1)_a$  and  $U(1)_b$ .

For our purpose, it is convenient to rewrite the action (2.5) as

$$S = \int_{\tau_0}^{\tau_1} d\tau \left[ \frac{i}{2} \left( \bar{Z}_A^i \dot{Z}_i^A - Z_i^A \dot{\bar{Z}}_A^i \right) + a \left( \bar{Z}_A^i Z_i^A - 2s \right) \right. \\ \left. + b^3 \left( \bar{Z}_A^j \sigma_{3j}{}^k Z_k^A - 2t \right) + b^{\hat{\imath}} \bar{Z}_A^j \sigma_{\hat{\imath}j}{}^k Z_k^A - \frac{1}{2e} b^{\hat{\imath}} b^{\hat{\imath}} - \frac{k^2}{2} e \right. \\ \left. + h \left( \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_j^{\dot{\alpha}} - \sqrt{2} m e^{i\varphi} \right) + \bar{h} \left( \epsilon_{ij} \bar{\pi}_{\alpha}^i \bar{\pi}^{j\alpha} - \sqrt{2} m e^{-i\varphi} \right) \right].$$
(2.7)

Variation of the action (2.7) with respect to  $a, b^3$ , and  $b^{\hat{i}}$  yields the constraints

$$T_0 - s = 0$$
, (2.8a)

$$T_3 - t = 0$$
, (2.8b)

$$T_{\hat{i}} - \frac{1}{2e}b^{\hat{i}} = 0,$$
 (2.8c)

respectively, where

$$T_0 := \frac{1}{2} \bar{Z}_A^i Z_i^A, \quad T_r := \frac{1}{2} \bar{Z}_A^j \sigma_{rj}{}^k Z_k^A \quad (r = \hat{\imath}, 3).$$
(2.9)

Additionally, variation of the action (2.7) with respect to e yields  $b^{\hat{i}}b^{\hat{i}} = k^2 e^2$ . Combining this and Eq. (2.8c), we obtain

$$T_i T_i - \frac{1}{4}k^2 = 0. (2.10)$$

In earlier formulations,<sup>9–17</sup> constraints similar to Eqs. (2.8a), (2.8b), and (2.10) have been incorporated into an action by hand with the help of appropriate Lagrange multipliers. In contrast, in our formulation, the constraints (2.8a)–(2.8c) are automatically incorporated in the action S on the basis of the gauge principle. In fact, Eq. (2.8a) is based on the  $U(1)_a$  invariance of S and Eqs. (2.8b) and (2.8c) are based on the  $U(1)_b$  invariance of S. For the constraint (2.10), it is obtained in connection with the invariance of S under the reparametrization  $\tau \to \tau'(\tau)$  and the  $U(1)_b$  transformation.

Variation of S with respect to h and  $\bar{h}$  yields the mass-shell constraints

$$\epsilon^{ij}\pi_{i\dot{\alpha}}\pi^{\dot{\alpha}}_{j} - \sqrt{2}me^{i\varphi} = 0, \qquad (2.11a)$$

$$\epsilon_{ij}\bar{\pi}^i_\alpha\bar{\pi}^{j\alpha} - \sqrt{2}me^{-i\varphi} = 0. \qquad (2.11b)$$

Equation (2.11b) is the complex conjugate of Eq. (2.11a). In our previous paper,<sup>18</sup> unlike the constraints (2.8a)–(2.8c), the constraints (2.11a) and (2.11b) have not

been considered to be that which are derived in relation to some gauge invariance of S. In earlier papers,<sup>9–18</sup> the constraints (2.11a) and (2.11b), or constraints similar to them, have been introduced by hand without taking into account gauge symmetries.

# 3. Inhomogeneous transformations

For convenience, we combine the transformations (2.3) and (2.4) into the following local U(2) transformation:

$$Z_i^A \to Z_i'^A = \mathcal{U}_i^{\ j}(\tau) Z_j^A , \qquad (3.1a)$$

$$\bar{Z}_A^i \to \bar{Z}_A^{\prime i} = \bar{Z}_A^j \mathcal{U}^{\dagger}{}_j{}^i(\tau) \,. \tag{3.1b}$$

$$\xi \to \xi' = \xi'(\xi) , \qquad (3.1c)$$

$$\bar{\xi} \to \bar{\xi}' = \bar{\xi}'(\bar{\xi}),$$
 (3.1d)

$$h \to h' = e^{-2i\theta(\tau)}h, \qquad (3.1e)$$

$$\bar{h} \to \bar{h}' = e^{2i\theta(\tau)}\bar{h},$$
(3.1f)

$$\varphi \to \varphi' = \varphi + 2\theta(\tau),$$
 (3.1g)

$$\mathbf{a} \to \mathbf{a}' = \mathcal{U}\mathbf{a}\,\mathcal{U}^{\dagger} - i\dot{\mathcal{U}}\mathcal{U}^{\dagger},$$
 (3.1h)

$$\mathbf{e} \to \mathbf{e}' = \mathbf{e} \,, \tag{3.1i}$$

where  $\mathcal{U}(\tau)$  is defined by  $\mathcal{U}(\tau) := e^{i\theta(\tau)}U(\tau)$ , being an element of  $U(2) \cong U(1)_a \times SU(2)$  and satisfying  $\mathcal{U}(\tau_1) = \mathcal{U}(\tau_0)$ . In addition, **a** is a one-dimensional U(2) gauge field defined by  $\mathbf{a} := a\sigma_0 + b$ , with  $\sigma_0$  being the 2 by 2 unit matrix. The transformation rule (3.1h) can be verified by using the transformation rules (2.3h) and (2.4i). As pointed out by Perjés and Hughston independently, the linear transformations that preserve both the momentum vector  $p_{\alpha\dot{\alpha}} := \sum_{i=1}^{n} \bar{\pi}^i_{\alpha} \pi_{i\dot{\alpha}} \ (n \geq 2)$  and the angular momentum spinor  $\mu_{\alpha\beta} := (i/2) \sum_{i=1}^{n} (\omega_{i\alpha} \bar{\pi}^i_{\beta} + \omega_{i\beta} \bar{\pi}^i_{\alpha})$  form a Lie group by themselves.<sup>3-7</sup> This group is a group extension of U(n) defined so as to include inhomogeneous transformations, and it is denoted by IU(n).

For a massive particle, the *n*-twistor expression of  $p_{\alpha\dot{\alpha}}$  given above reduces to the two-twistor expression  $p_{\alpha\dot{\alpha}} = \sum_{i=1}^{2} \bar{\pi}_{\alpha}^{i} \pi_{i\dot{\alpha}}$  by a unitary transformation, as was proved in Ref. 19. Therefore it is sufficient to consider only the two-twistor system in which the IU(2) symmetry is realized. Inhomogeneous extensions of Eqs. (3.1a) and (3.1b) are, respectively, found to be<sup>3-7</sup>

$$Z_i^A \to Z_i^{\prime A} = \mathcal{U}_i^{\ j}(\tau) \Big( Z_j^A + \Lambda(\tau) \epsilon_{jk} I^{AB} \bar{Z}_B^k \Big) , \qquad (3.2a)$$

$$\bar{Z}^{i}_{A} \to \bar{Z}^{\prime i}_{A} = \left(\bar{Z}^{j}_{A} + \bar{\Lambda}(\tau)\epsilon^{jk}I_{AB}Z^{B}_{k}\right)\mathcal{U}^{\dagger}{}_{j}{}^{i}(\tau) , \qquad (3.2b)$$

where  $\Lambda(\tau)$  is a complex transformation parameter for the local inhomogeneous transformation, satisfying  $\Lambda(\tau_1) = \Lambda(\tau_0)$ . (In the two-twistor system, a skew-symmetric parameter  $\Lambda_{ij}$  can be written as  $\Lambda\epsilon_{ij}$ .) In addition,  $I^{AB}$  and  $I_{AB}$  denote

the so-called infinity twistors defined by

$$I^{AB} := \begin{pmatrix} \epsilon^{\alpha\beta} & 0\\ 0 & 0 \end{pmatrix}, \qquad I_{AB} := \begin{pmatrix} 0 & 0\\ 0 & \epsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix}.$$
(3.3)

If  $\Lambda = 0$ , Eqs. (3.2a) and (3.2b) become Eqs. (3.1a) and (3.1b), respectively.

Using Eqs. (3.2) and (3.1h), we can derive the transformation behaviors of  $DZ_i^A$ and  $\bar{D}Z_A^i$  under the local IU(2) transformation as follows:

$$DZ_{i}^{A} \rightarrow D'Z_{i}^{\prime A} = \mathcal{U}_{i}^{j} \left\{ DZ_{j}^{A} + \dot{A}\epsilon_{jk}I^{AB}\bar{Z}_{B}^{k} + \Lambda I^{AB} \left( \epsilon_{jk}\dot{\bar{Z}}_{B}^{k} - i\mathbf{a}_{j}^{k}\epsilon_{kl}\bar{Z}_{B}^{l} \right) \right\}, \qquad (3.4a)$$
$$\bar{D}\bar{Z}_{A}^{i} \rightarrow \bar{D}'\bar{Z}_{A}^{\prime i} = \left\{ \bar{D}\bar{Z}_{A}^{j} + \dot{\bar{A}}\epsilon^{jk}I_{AB}Z_{k}^{B} \right\}$$

$$+ \bar{\Lambda} I_{AB} \left( \epsilon^{jk} \dot{Z}_k^B + i \mathsf{a}_k^j \epsilon^{kl} Z_l^B \right) \Big\} \mathcal{U}^{\dagger}{}_j{}^i.$$
(3.4b)

Additionally, using Eqs (3.2) and (3.4) together with the formulas

$$I_{AB}Z_i^A Z_j^B = \frac{1}{2} \epsilon^{kl} \pi_{k\dot{\alpha}} \pi_l^{\dot{\alpha}} \epsilon_{ij} , \quad I^{AB} \bar{Z}_A^i \bar{Z}_B^j = \frac{1}{2} \epsilon_{kl} \bar{\pi}_{\alpha}^k \bar{\pi}^{l\alpha} \epsilon^{ij} , \qquad (3.5)$$

we can show that

$$\frac{i}{2} \left( \bar{Z}_{A}^{\prime i} D^{\prime} Z_{i}^{\prime A} - Z_{i}^{\prime A} \bar{D}^{\prime} \bar{Z}_{A}^{\prime i} \right) = \frac{i}{2} \left( \bar{Z}_{A}^{i} D Z_{i}^{A} - Z_{i}^{A} \bar{D} \bar{Z}_{A}^{i} \right) - \frac{i}{2} \left( \dot{A} + 2ia\bar{A} \right) \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_{j}^{\dot{\alpha}} + \frac{i}{2} \left( \dot{A} - 2ia\bar{A} \right) \epsilon_{ij} \bar{\pi}_{\alpha}^{i} \bar{\pi}^{j\alpha} .$$
(3.6)

Thus we see that the integrand in the first line of Eq. (2.1) is not invariant under the inhomogeneous transformation, even when it is the global transformation specified by  $\dot{\Lambda} = 0$ .

Now, we note that the second and third lines in the right-hand side of Eq. (3.6), which we refer to as extra terms, are proportional to the quantity  $\epsilon^{ij}\pi_{i\dot{\alpha}}\pi_{j}^{\dot{\alpha}}$  or  $\epsilon_{ij}\bar{\pi}_{\alpha}^{i}\bar{\pi}^{j\alpha}$ . Since the same quantities are included in the action S in the form of  $h\epsilon^{ij}\pi_{i\dot{\alpha}}\pi_{j}^{\dot{\alpha}} + \bar{h}\epsilon_{ij}\bar{\pi}_{\alpha}^{i}\bar{\pi}^{j\alpha}$ , it is possible to cancel out the extra terms by carrying out an appropriate modification of the transformation rules (3.1e) and (3.1f). In fact, we can completely cancel out the extra terms by modifying Eqs. (3.1e) and (3.1f) to be

$$h \to h' = e^{-2i\theta(\tau)} \left\{ h + \frac{i}{2} \left( \dot{\bar{A}}(\tau) + 2ia\bar{A}(\tau) \right) \right\}, \qquad (3.7a)$$

$$\bar{h} \to \bar{h}' = e^{2i\theta(\tau)} \left\{ \bar{h} - \frac{i}{2} \left( \dot{A}(\tau) - 2iaA(\tau) \right) \right\}, \qquad (3.7b)$$

respectively. Here, h and  $\bar{h}$  are regarded as gauge fields for the inhomogeneous transformation, because the transformation of h and  $\bar{h}$  given in Eq. (3.7) provides the terms that contribute to canceling out the extra terms in Eq. (3.6).

At the same time, we modify the transformation rule (3.1g) in such a way that the mass term  $-\sqrt{2}m\mathcal{H}$  with  $\mathcal{H} := he^{i\varphi} + \bar{h}e^{-i\varphi}$  included in S remains invariant under the simultaneous transformation that is defined by Eq. (3.7) and a modified version of Eq. (3.1g). The square of the both sides of the invariant condition  $\mathcal{H}' = \mathcal{H}$ leads to  $h'(e^{i\varphi'})^2 - \mathcal{H}e^{i\varphi'} + \bar{h}' = 0$ . This can be solved to yield

$$\varphi \to \varphi' = -i \ln\left(\frac{\mathcal{H} \pm i\sqrt{4|h'|^2 - \mathcal{H}^2}}{2h'}\right).$$
 (3.8)

Here,  $4|h'|^2 \geq \mathcal{H}^2$  holds, because  $h'e^{i\varphi'} - \bar{h}'e^{-i\varphi'} = \pm i |h'e^{i\varphi'} - \bar{h}'e^{-i\varphi'}|$  is satisfied. Equation (3.8) is precisely a modified version of Eq. (3.1g). When  $\Lambda = 0$ , Eqs. (3.7a), (3.7b), and (3.8) reduce to Eqs. (3.1e), (3.1f), and (3.1g), respectively.

The action S turns out to be invariant under the local IU(2) transformation

$$Z_i^A \to Z_i^{\prime A} = \mathcal{U}_i^{\ j}(\tau) \Big( Z_j^A + \Lambda(\tau) \epsilon_{jk} I^{AB} \bar{Z}_B^k \Big) , \qquad (3.9a)$$

$$\bar{Z}_A^i \to \bar{Z}_A^{\prime i} = \left( \bar{Z}_A^j + \bar{\Lambda}(\tau) \epsilon^{jk} I_{AB} Z_k^B \right) \mathcal{U}_j^{\dagger}{}^i(\tau) , \qquad (3.9b)$$

$$\xi \to \xi' = \xi'(\xi) \,, \tag{3.9c}$$

$$\bar{\xi} \to \bar{\xi}' = \bar{\xi}'(\bar{\xi}) , \qquad (3.9d)$$

$$h \to h' = e^{-2i\theta(\tau)} \left\{ h + \frac{i}{2} \left( \dot{\bar{A}}(\tau) + 2ia\bar{A}(\tau) \right) \right\}, \qquad (3.9e)$$

$$\bar{h} \to \bar{h}' = e^{2i\theta(\tau)} \left\{ \bar{h} - \frac{i}{2} \left( \dot{A}(\tau) - 2iaA(\tau) \right) \right\}, \tag{3.9f}$$

$$\varphi \to \varphi' = -i \ln\left(\frac{\mathcal{H} \pm i\sqrt{4|h'|^2 - \mathcal{H}^2}}{2h'}\right),$$
(3.9g)

$$\mathbf{a} \to \mathbf{a}' = \mathcal{U} \mathbf{a} \, \mathcal{U}^{\dagger} - i \dot{\mathcal{U}} \mathcal{U}^{\dagger}, \tag{3.9h}$$

$$\mathbf{e} \to \mathbf{e}' = \mathbf{e} \,, \tag{3.9i}$$

rather than under the local U(2) transformation in Eq. (3.1). In this way, the local U(2) symmetry of the present twistor model can be extended to the local IU(2) symmetry.

## 4. Conclusions

As we have seen, it is necessary for S to include  $\int_{\tau_0}^{\tau_1} d\tau \left[h\epsilon^{ij}\pi_{i\dot{\alpha}}\pi_j^{\dot{\alpha}} + \bar{h}\epsilon_{ij}\bar{\pi}_{\alpha}^i\bar{\pi}^{j\alpha}\right]$ as an additive term in order that S can remain invariant under the local IU(2)transformation as a result of canceling out the extra terms in Eq. (3.6). Here, it is essential that h and  $\bar{h}$  transform inhomogeneously as the gauge fields corresponding to the parameters  $\bar{\Lambda}$  and  $\Lambda$ , respectively, as in Eq. (3.7).

The action S also includes the mass term  $\int_{\tau_0}^{\tau_1} d\tau [-\sqrt{2}m\mathcal{H}]$ , which itself remains

invariant under the local IU(2) transformation. The sum of these two terms reads

$$S_{h} := \int_{\tau_{0}}^{\tau_{1}} d\tau \left[ h \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_{j}^{\dot{\alpha}} + \bar{h} \epsilon_{ij} \bar{\pi}_{\alpha}^{i} \bar{\pi}^{j\alpha} - \sqrt{2} m \mathcal{H} \right]$$
$$= \int_{\tau_{0}}^{\tau_{1}} d\tau \left[ h \left( \epsilon^{ij} \pi_{i\dot{\alpha}} \pi_{j}^{\dot{\alpha}} - \sqrt{2} m e^{i\varphi} \right) + \bar{h} \left( \epsilon_{ij} \bar{\pi}_{\alpha}^{i} \bar{\pi}^{j\alpha} - \sqrt{2} m e^{-i\varphi} \right) \right], \qquad (4.1)$$

which is precisely the mass-shell term of S. The local IU(2) invariance of S holds with the aid of the one-dimensional gauge fields a, b, h, and  $\bar{h}$ .

In our formulation,  $S_h$  has been found on the basis of the local IU(2) symmetry of the system. Therefore the mass-shell constraints (2.11a) and (2.11b) are considered to be outcomes originated in the symmetry under the inhomogeneous transformation. We can say that all the constraints, including the mass-shell constraints, are automatically derived from the local IU(2) symmetry in a self-contained way.

Canonical quantization of the twistor model governed by S was actually carried out in Ref. 18. In the quantization procedure, the first-class constraints (2.8a), (2.8b) and (2.10) are treated as conditions imposed on the physical state vector, after replacing functions in the constraints by the corresponding operators. Another canonical quantization is performed after fixing gauge at the classical level. We expect that the convenient gauge-fixing for quantization can be carried out by using the local IU(2) symmetry.

# Appendix A.

In this paper and Ref. 18, the coset space  $\mathcal{C} := SU(2)/U(1) (\cong \mathbb{C}\mathbf{P}^1)$  is introduced to nonlinearly realize the SU(2) symmetry that is linearly realized for the pair of twistors  $(Z_1^A, Z_2^A)$ . In the appendix, we briefly mention  $\mathcal{C}$  and some formulas used in the gauged twistor formulation of a massive particle.

Let  $\xi \in \mathbb{C}$  be an inhomogeneous coordinate of a point on  $\mathcal{C}$  and let  $V(\xi, \bar{\xi}) \in SU(2)$  be representative elements chosen from each left coset of U(1) labeled by  $\xi$ . The left action of  $U \in SU(2)$  on  $V(\xi, \bar{\xi})$  causes a nonlinear transformation  $\xi \to \xi' = \xi'(\xi)$  in accordance with

$$V(\xi,\bar{\xi}) \to V(\xi',\bar{\xi}') = UV(\xi,\bar{\xi})e^{-i\vartheta\sigma_3}.$$
(A.1)

Here,  $\sigma_3$  is the third component of the Pauli matrices, and  $\vartheta$  is a real parameter for the U(1) transformation generated by  $\sigma_3$ .<sup>20–22</sup>

From  $V(\xi, \bar{\xi})$ , we define  $e_{\xi}^r$  and  $e_{\bar{\xi}}^r$  (r = +, -, 3) by

$$e_{\xi}{}^{r}\sigma_{r} = -iV^{\dagger}\frac{\partial V}{\partial\xi}, \quad e_{\bar{\xi}}{}^{r}\sigma_{r} = -iV^{\dagger}\frac{\partial V}{\partial\bar{\xi}}.$$
 (A.2)

Here,  $\sigma_{\pm}$  are defined from the Pauli matrices  $\sigma_1$  and  $\sigma_2$  as  $\sigma_{\pm} := (\sigma_1 \pm i\sigma_2)/2$ . Choosing V to be an appropriate form

$$V(\xi,\bar{\xi}) = \exp\left[i\left(\bar{\zeta}\sigma_{+} + \zeta\sigma_{-}\right)\right] \tag{A.3}$$

with

$$\zeta := \frac{\xi}{|\xi|} \arctan |\xi|, \quad \bar{\zeta} := \frac{\bar{\xi}}{|\xi|} \arctan |\xi|, \quad (A.4)$$

we obtain

$$e_{\xi}^{+} = 0, \qquad e_{\xi}^{-} = \frac{1}{1 + |\xi|^2}, \quad e_{\xi}^{3} = \frac{-i\xi}{2(1 + |\xi|^2)}, \quad (A.5a)$$

$$e_{\bar{\xi}}^{+} = \frac{1}{1+|\xi|^2}, \quad e_{\bar{\xi}}^{-} = 0, \qquad e_{\bar{\xi}}^{-3} = \frac{i\xi}{2(1+|\xi|^2)}.$$
 (A.5b)

The so-called zweibeins  $(e_{\xi}^+, e_{\bar{\xi}}^+)$  and  $(e_{\xi}^-, e_{\bar{\xi}}^-)$  lead to a proper metric tensor on C:

$$g_{\xi\xi} = e_{\xi}^{+} e_{\xi}^{-} = 0, \qquad (A.6a)$$

$$g_{\xi\bar{\xi}} = g_{\bar{\xi}\xi} = \frac{1}{2} \left( e_{\xi}^{+} e_{\bar{\xi}}^{-} + e_{\bar{\xi}}^{+} e_{\xi}^{-} \right) = \frac{1}{2(1+|\xi|^2)^2} , \qquad (A.6b)$$

$$g_{\bar{\xi}\bar{\xi}} = e_{\bar{\xi}}^+ e_{\bar{\xi}}^- = 0.$$
 (A.6c)

The  $\xi \bar{\xi}$ -component of the metric tensor is precisely the Fubini-Study metric tensor on  $\mathbb{C}\mathbf{P}^1$ .

Now, suppose that U is infinitely near the identity so that  $U = 1 + i\epsilon^r \sigma_r$ (r = +, -, 3) is valid with infinitesimal parameters  $\epsilon^r$ . Accordingly,  $\xi', \bar{\xi}'$ , and  $e^{-i\vartheta\sigma_3}$  can take the following form:  $\xi' = \xi + \epsilon^r K_r \xi, \bar{\xi}' = \bar{\xi} + \epsilon^r K_r \xi$ , and  $e^{-i\vartheta\sigma_3} = 1 - i\epsilon^r \Omega_r \sigma_3$ . Substituting these into Eq. (A.1), we obtain

$$K_r^{\xi} \frac{\partial V}{\partial \xi} + K_r^{\xi} \frac{\partial V}{\partial \bar{\xi}} = i\sigma_r V - i\Omega_r V \sigma_3 \tag{A.7}$$

after removing  $\epsilon_r$ . It turns out that  $(K_r^{\xi}, K_r^{\overline{\xi}})$ , or more precisely  $K_r := K_r^{\xi} \partial/\partial \xi + K_r^{\overline{\xi}} \partial/\partial \overline{\xi}$ , are the so-called SU(2) Killing vectors on  $\mathcal{C}$  and the  $\Omega_r$  are associated compensators. Multiplying Eq.(A.7) by  $-iV^{\dagger}$  from the left and using Eq. (A.2) yield

$$\mathcal{V}_{r}^{\ \hat{i}} = K_{r}^{\ \xi} e_{\xi}^{\ \hat{i}} + K_{r}^{\ \bar{\xi}} e_{\bar{\xi}}^{\ \hat{i}} \quad (\hat{i} = +, -), \tag{A.8a}$$

$$\mathcal{V}_{r}^{\ 3} = K_{r}^{\ \xi} e_{\xi}^{\ 3} + K_{r}^{\ \xi} e_{\bar{\xi}}^{\ 3} + \Omega_{r} \,, \tag{A.8b}$$

where  $\mathcal{V}_r{}^s$   $(s = \hat{\imath}, 3)$  are defined by

$$V^{\dagger}\sigma_r V = \mathcal{V}_r^{\ s}\sigma_s \,. \tag{A.9}$$

The matrix  $\mathcal{V}$  is the adjoint representation of V. From Eq. (A.9), we have

$$\mathcal{V}_r^{+} = \operatorname{tr} \left( V^{\dagger} \sigma_r V \sigma_- \right), \qquad (A.10a)$$

$$\mathcal{V}_r^{-} = \operatorname{tr} \left( V^{\dagger} \sigma_r V \sigma_+ \right), \qquad (A.10b)$$

$$\mathcal{V}_r^{\ 3} = \frac{1}{2} \mathrm{tr} \left( V^{\dagger} \sigma_r V \sigma_3 \right). \tag{A.10c}$$

In the unitary gauge specified by V = 1, it follows that  $\mathcal{V}_r{}^s = \delta_r{}^s$ . When V = 1, we see from Eq. (A.3) that  $\zeta = 0$ , and hence we see from (A.4) that  $\xi = 0$ . By substituting  $\xi = 0$  into Eqs. (A.5a) and (A.5b), they reduce to

$$e_{\xi}^{+} = 0, \quad e_{\xi}^{-} = 1, \quad e_{\xi}^{3} = 0,$$
 (A.11a)

$$e_{\bar{\xi}}^{+} = 1, \quad e_{\bar{\xi}}^{-} = 0, \quad e_{\bar{\xi}}^{3} = 0.$$
 (A.11b)

Accordingly, Eq. (A.6) becomes

$$g_{\xi\xi} = 0, \quad g_{\xi\bar{\xi}} = g_{\bar{\xi}\xi} = \frac{1}{2}, \quad g_{\bar{\xi}\bar{\xi}} = 0.$$
 (A.12)

Using Eqs. (A.8a) and (A.11) and  $\mathcal{V}_r{}^s = \delta_r{}^s$ , we can show that

$$K_{+}^{\xi} = 0, \quad K_{-}^{\xi} = 1, \quad K_{3}^{\xi} = 0,$$
 (A.13a)

$$K_{+}^{\xi} = 1, \quad K_{-}^{\xi} = 0, \quad K_{3}^{\xi} = 0.$$
 (A.13b)

Similarly, using Eq. (A.8b), we can show that

$$\Omega_{+} = 0, \quad \Omega_{-} = 0, \quad \Omega_{3} = 1.$$
(A.14)

# References

- R. Penrose, Twistors and Particles: An Outline, in Feldafing Conference of the Max-Planck Inst. on Quantum Theory and the Structure of Space-time, 1974, pp. 129-145, Carl Hanser Verlag, Munich, 1975.
- 2. R. Penrose, "The twistor programme," Rep. Math. Phys. 12, 65 (1977).
- 3. Z. Perjés, "Twistor variables of relativistic mechanics," Phys. Rev. D 11, 2031 (1975).
- 4. Z. Perjés, "Unitary space of particle internal states," Phys. Rev. D 20, 1857 (1979).
- Z. Perjés, "Perspectives of Penrose theory in particle physics," Rep. Math. Phys. 12, 193 (1977).
- 6. Z. Perjés, "Internal symmetries in twistor theory," Czech. J. Phys. B 32, 540 (1982).
- L. P. Hughston, *Twistors and Particles*, Lecture Notes in Physics Vol. 97 (Springer-Verlag, Berlin, 1979).
- 8. G. Albonico, Y. Geyer, and L. Mason, "From twistor-particle models to massive amplitudes," SIGMA 18, 045 (2022), arXiv:2203.08087 [hep-th].
- S. Fedoruk and V. G. Zima, "Bitwistor formulation of massive spinning particle," J. Kharkiv Univ. 585, 39 (2003), arXiv:hep-th/0308154.
- A. Bette, J. A. de Azcárraga, J. Lukierski, and C. Miquel-Espanya, "Massive relativistic free fields with Lorentz spins and electric charges," Phys. Lett. B 595, 491 (2004), arXiv:hep-th/0405166.
- J. A. de Azcárraga, A. Frydryszak, J. Lukierski, and C. Miquel-Espanya, "Massive relativistic particle model with spin from free two-twistor dynamics and its quantization," Phys. Rev. D 73, 105011 (2006), arXiv:hep-th/0510161.
- S. Fedoruk, A. Frydryszak, J. Lukierski, and C. Miquel-Espanya, "Extension of the Shirafuji model for massive particles with spin," Int. J. Mod. Phys. A 21, 4137 (2006), arXiv:hep-th/0510266.
- 13. J. A. de Azcárraga, J. M. Izquierdo, and J. Lukierski, "Supertwistors, massive superparticles and  $\kappa$ -symmetry," J. High Energy Phys. 01 (2009) 041, arXiv:0808.2155 [hep-th].
- L. Mezincescu, A. J. Routh, and P. K. Townsend, "Supertwistors and massive particles," Ann. Phys. (Amsterdam) 346, 66 (2014), arXiv:1312.2768 [hep-th].

- S. Fedoruk and J. Lukierski, "Massive twistor particle with spin generated by Souriau-Wess-Zumino term and its quantization," Phys. Lett. B 733, 309 (2014), arXiv:1403.4127 [hep-th].
- J. A. de Azcárraga, S. Fedoruk, J. M. Izquierdo, and J. Lukierski, "Two-twistor particle models and free massive higher spin fields," J. High Energy Phys. 04 (2015) 010, arXiv:1409.7169 [hep-th].
- L. Mezincescu, A. J. Routh, and P. K. Townsend, "Twistors and the massive spinning particle," J. Phys. A 49, 025401 (2016), arXiv:1508.05350 [hep-th].
- S. Deguchi and S. Okano, "Gauged twistor formulation of a massive spinning particle in four dimensions," Phys. Rev. D 93, 045016 (2016) [Erratum-ibid. D 93, 089906(E) (2016)], arXiv:1512.07740 [hep-th].
- 19. S. Okano and S. Deguchi, "A no-go theorem for the *n*-twistor description of a massive particle," J. Math. Phys. **58**, 031701 (2017), arXiv:1606.01339 [hep-th].
- S. Coleman, J. Wess, and B. Zumino, "Structure of phenomenological Lagrangians I," Phys. Rev. 177, 2239 (1969).
- A. Salam and J. Strathdee, "Nonlinear realizations I: The role of goldstone bosons," Phys. Rev. 184, 1750 (1969).
- P. van Nieuwenhuizen, "General Theory of Coset Manifolds and Antisymmetric Tensors Applied to Kaluza-Klein Supergravity," in *Supersymmetry and Supergravity '84*, ed. B. de Wit et al. (World Scientific, Singapore, 1984), p. 239.