

On the CP Nature of the ‘95 GeV’ Anomalies

Tanmoy Mondal,^a Stefano Moretti^{b,c} and Prasenjit Sanyal^d

^a*Birla Institute of Technology and Science, Pilani, 333031, Rajasthan, India*

^b*School of Physics and Astronomy, University of Southampton, Southampton SO17 1BJ, United Kingdom*

^c*Department of Physics and Astronomy, Uppsala University, Box 516, SE-751 20 Uppsala, Sweden*

^d*Department of Physics, Konkuk University, Seoul 05029, Republic of Korea*

E-mail: tanmoy.mondal@pilani.bits-pilani.ac.in, s.moretti@soton.ac.uk,
stefano.moretti@physics.uu.se, prasenjit.sanyal01@gmail.com

ABSTRACT: Under the assumption that the various evidences of a ‘95 GeV’ excess, seen in data at the Large Electron Positron (LEP) collider as well as the Large Hadron Collider (LHC), correspond to actual signals of new physics Beyond the Standard Model (BSM), we characterise the underlying particle explaining these in terms of its Charge/Parity (CP) quantum numbers in a model independent way. In doing so, we assume the new object having spin-0 and test the CP-even (scalar) and CP-odd (pseudoscalar) hypotheses as well superpositions of these in its $\tau^+\tau^-$ decays. We prove that the High Luminosity LHC (HL-LHC) will be in a position to disentangle the CP nature of such a new particle.

Contents

1	Introduction	1
2	A Simplified Model to Address the Excesses	3
2.1	Production and Decay Modes	4
2.2	Allowed Parameter Space	4
3	Uncovering the CP Nature of the Resonance	5
3.1	Impact Parameter Method	6
3.2	Rho Decay Plane Method	6
3.3	Combined IP and Rho Decay Plane Methods	7
3.4	Event Simulation and Selection	7
4	Results	7
5	Conclusions	9

1 Introduction

Since the discovery of the 125 GeV Higgs boson at the Large Hadron Collider (LHC) in 2012 [1, 2], significant efforts have been made by the ATLAS and CMS collaborations in ascertaining the nature of this particle, that is, measuring its width, couplings, spin and CP quantum numbers. All corresponding measurements are pointing towards the discovered Higgs boson being the Standard Model (SM) one. Despite all this, though, the search for physics Beyond the SM (BSM) continues, spurred, on the theoretical side by its many flaws and, on the experimental side, by the ever increasing precision of the analyses as well as additional data sets being collected during Run 3 of the LHC.

The 2-Higgs Doublet Model (2HDM) [3, 4], as embedded in various BSM scenarios, including fundamental ones like Supersymmetry [5] and Compositeness [6, 7], is particularly interesting, as it offers in its particle spectrum both a CP-even and a CP-odd (neutral) Higgs boson, alongside the discovered SM-like Higgs state, both of which can in fact be lighter or heavier than the latter¹. Hence, this BSM scenario has been studied extensively since its inception. However, its general version allows for non-diagonal Yukawa couplings, potentially leading to large Flavor Changing Neutral Currents (FCNCs) at tree level, contrary to experimental evidence. To constrain the latter, from the theoretical side, a Z_2 symmetry (which can be softly broken, in fact) is typically imposed to define the coupling structure of the two Higgs doublets to SM fermions. This classification includes the so-called Type-I, Type-II, lepton-specific and flipped scenarios [4], alongside the 2HDM Type-III, which allows direct couplings of both doublets to all SM fermions. The 2HDM Yukawa structure is then refined by both theoretical consistency requirements and experimental measurements of the h mass and couplings or limits on the the same parameters for the H , A and H^\pm states.

Given that a 2HDM, as mentioned, allows (in some Types) for neutral Higgs bosons states lighter than 125 GeV, both ATLAS and CMS have pursued this possibility through dedicated searches in presence of a mass configuration wherein m_h and/or $m_A < m_H = 125$ GeV (with m_{H^\pm}

¹There are also charged Higgs bosons.

also rather light, to comply with various high-precision data). (Hereafter, in our notation, the CP-even H state is the SM-like one already discovered while the h and A states are the (light) CP-even and CP-odd Higgs bosons, respectively, plus H^\pm are the charged ones.)

During searches for a low mass neutral Higgs boson, the CMS collaboration reported an excess near 95 GeV in the di-photon invariant mass already back in 2018 [8]. In March 2023, they confirmed such an excess, eventually claiming a local significance of 2.9σ computed at $m_{\gamma\gamma} = 95.4$ GeV [9, 10]. Similarly, ATLAS observed an excess at about 95 GeV with a local significance of 1.7σ , aligning with CMS findings [11, 12]. Furthermore, the CMS collaboration has reported an excess in the search for a light neutral (pseudo)scalar Higgs boson decaying into $\tau^+\tau^-$ pairs, with local significance of 2.6σ around a mass of 95 GeV. (Yet, other CMS data tend to invalidate attempts to attribute such a di-tau excess to a new Higgs boson because searches for a spin-0 resonance produced in $t\bar{t}$ -associated production and decaying into τ pairs, seem not to support this interpretation [13].) Finally, the Large Electron Positron (LEP) collider collaborations [14] explored the low Higgs mass domain extensively in the $e^+e^- \rightarrow Zh$ production mode, with h decaying via the $\tau^+\tau^-$ and $b\bar{b}$ channels. Interestingly for our purposes is the fact that an excess had been reported in 2006 in the $e^+e^- \rightarrow Zb\bar{b}$ channel for $m_{b\bar{b}}$ around 98 GeV [15]. Given the limited mass resolution of the di-jet invariant mass at LEP, though, this anomaly may well coincide with the discussed excesses seen by CMS and/or ATLAS in the $\gamma\gamma$ and $\tau^+\tau^-$ final states.

Since all the excesses appear in very similar mass regions, several studies [16–49] have explored the possibility of simultaneously explaining these anomalies within BSM frameworks featuring a non-standard Higgs state lighter than 125 GeV, while being in agreement with current measurements of the properties of the SM-like Higgs state observed at the LHC. In the attempt to explain the excesses in the $\gamma\gamma$ and $b\bar{b}$ channels, it was found in Refs. [50–54] that the 2HDM Type-III with a particular Yukawa texture can successfully accommodate both measurements simultaneously with the lightest CP-even Higgs boson of the model, while being consistent with all relevant theoretical and experimental constraints. In fact, also a 2HDM Type-I can afford similar explanations to the excesses [55]. Further recent studies have shown that actually all three aforementioned signatures can be simultaneously explained in the 2HDM plus a real (N2HDM) [29] or a complex (S2HDM) [30, 32] singlet.

The purpose of this paper is to assume that, indeed, a spin-0 particle of yet undefined CP is behind all such anomalies. However, amongst the latter, we focus here on the $\tau^+\tau^-$ one, as this has some diagnostic power in terms of pinning down its CP properties, as shown in Refs. [56–68]. So, let us spend some time on this particular anomaly. The CMS collaboration has reported such an excess in the $\tau^+\tau^-$ final state across all τ decay modes (leptonic and hadronic) for both leptons. The local and global significances of this particular excess are 2.6σ and 2.3σ for $m_{\tau^+\tau^-} = 95$ GeV, respectively. At $m_{\tau^+\tau^-} = 100$ GeV, these values are 3.1σ and 2.7σ . By introducing an additional single neutral narrow resonance h , the best fit values to the signal rate for this excess are [69]:

$$\sigma(gg \rightarrow h) \text{BR}(h \rightarrow \tau^+\tau^-) = 7.7_{-3.1}^{+3.9} \text{ pb} \quad \text{for } m_h = 95 \text{ GeV}, \quad (1.1)$$

$$\sigma(gg \rightarrow h) \text{BR}(h \rightarrow \tau^+\tau^-) = 5.8_{-2.0}^{+2.4} \text{ pb} \quad \text{for } m_h = 100 \text{ GeV}. \quad (1.2)$$

In our work, we assume a 95 GeV resonance. Interpreting Eq. (1.1) in terms of a signal strength, we get

$$\mu_{\tau^+\tau^-}^{\text{exp}} = \frac{\sigma^{\text{exp}}(gg \rightarrow h \rightarrow \tau^+\tau^-)}{\sigma^{\text{SM}}(gg \rightarrow H \rightarrow \tau^+\tau^-)} = 1.2 \pm 0.5, \quad (1.3)$$

where we use a symmetric uncertainty interval for such a signal strength, which is obtained from the lower uncertainty interval of the quoted result for $\sigma(gg \rightarrow h) \text{BR}(h \rightarrow \tau^+\tau^-)$. Also, in the signal strength, H represents a fictitious SM Higgs boson with mass 95 GeV.

For completeness, we also report the signal strengths for the two other aforementioned anomalies, as follows:

$$\mu_{\gamma\gamma}^{\text{exp}} = \frac{\sigma^{\text{exp}}(gg \rightarrow h \rightarrow \gamma\gamma)}{\sigma^{\text{SM}}(gg \rightarrow H \rightarrow \gamma\gamma)} = 0.6 \pm 0.2 \quad (1.4)$$

for $\gamma\gamma$ (see [8]) and

$$\mu_{b\bar{b}}^{\text{exp}} = \frac{\sigma^{\text{exp}}(e^+e^- \rightarrow ZS \rightarrow Zb\bar{b})}{\sigma^{\text{SM}}(e^+e^- \rightarrow Zh \rightarrow Zb\bar{b})} = 0.117 \pm 0.057 \quad (1.5)$$

for $b\bar{b}$ (see [15, 70]).

In this work, in order to ascertain the CP state of the underlying spin-0 resonance (which, for illustrative purposes, we also assume to be a Higgs state), we have adopted a model agnostic explanation to all the excesses. As we particularly focus on the CP nature of this 95 GeV object, we have setup a field with Yukawa couplings to all fermions, i.e., proportional to their mass, scaled by the corresponding SM Yukawa, while allowing in such an interaction for the simultaneous presence of both a γ_5 (i.e., a pseudoscalar interaction) and a unit matrix (i.e., a scalar interaction) structure, in both the $ht\bar{t}$ and $h\tau^+\tau^-$ vertices, entering in production and decay, respectively. After exploring the parameter space of this simplified model by testing it against theoretical and experimental constraints, including fitting the described excesses, we look at unravelling the CP properties of the h state at the LHC in the $\tau^+\tau^-$ channel using as diagnostic observable the signed acoplanarity angle ϕ_{CP} between the two τ lepton decay planes. In order to reconstruct the latter, we have used one-prong hadronic τ decays and Impact Parameter (IP) information [62, 64], wherein for the former we limited ourselves to the case of π and ρ mesons. By then performing a robust detector level simulation, we show that it is indeed possible to unambiguously differentiate a CP-even from a CP-odd as well as a maximally CP-mixed signal at the LHC with 3000 fb^{-1} of data per experiment, which is the expected final dataset size of the High-Luminosity LHC (HL-LHC) [71, 72]. However, we deem this result a conservative one since, as we gather more data at the current LHC (Run 3), the significance of the aforementioned excesses may well grow further, thereby stimulating dedicated analyses aimed at accessing the CP properties of this new resonance which may well prove to offer a better sensitivity than the one worked out in this article.

Our paper is organised as follows. In the next section, we describe our simplified model approach, including delineating its parameter space compatible with current constraints and explaining the $\gamma\gamma$, $\tau^+\tau^-$ and/or $b\bar{b}$ excesses. Then we move on to describe the construction of the mentioned CP-sensitive observable. Numerical results will follow, in turn preceding our conclusions.

2 A Simplified Model to Address the Excesses

We are interested in the study of CP properties of the observed spin-0 resonance, which might come from various UV complete scenarios like the Two Higgs Doublet Model, Next-to Minimal Supersymmetric scenarios (NMSSM) and other scalar extensions of the Standard Model. To remain model agnostic, we assumed a simplified model where, in addition to the SM spectrum, we consider only an additional scalar (S) of mass 95 GeV. The scalar particle S couples with the fermions as follows:

$$\mathcal{L}_{Sf\bar{f}} = -\rho_f^S \frac{m_f}{v} \left(\cos \alpha \bar{f}f + i \sin \alpha \bar{f}\gamma_5 f \right) S. \quad (2.1)$$

We have introduced CP-Violating (CPV) Yukawa coupling with α being the CPV parameter. α will be zero ($\pi/2$) when the scalar S is a CP-even (CP-odd) particle. For simplicity, we have assumed a uniform CP mixing among generations. The dimensionless real parameters ρ_f^S ($f = t, b, \tau$) are the

Yukawa coupling modifiers. The observed SM-like CP -even Higgs boson (h) of mass 125 GeV can couple with S depending on the UV completion. Our analysis will not depend on the coupling of S with the SM Higgs boson.

The coupling of the scalar S with the weak neutral gauge boson, denoted as g_{SVV} , does not affect the LHC excess observed in $\tau^+\tau^-$ and $\gamma\gamma$ channel. However, a non-zero g_{SVV} is necessary to explain the LEP $b\bar{b}$ excess. Such coupling can appear if S is part of a weak multiplet which mixes with the SM Higgs doublet. Since the observed 125 GeV Higgs boson at the LHC closely resembles the SM, the mixing of the Higgs doublet can not be too large, and we have considered $g_{SVV} < 0.3$.

2.1 Production and Decay Modes

Considering the large top mass and presuming ρ_b^h not to be significantly large compared to ρ_t^h , we can obtain the gluon effective coupling described by a dimension five operator

$$\mathcal{L}_{hgg} = \rho_t^S \cos \alpha A_S SG_{\mu\nu}^a G^{a,\mu\nu} + \rho_t^S \sin \alpha A'_S SG_{\mu\nu}^a \tilde{G}^{a,\mu\nu} \quad (2.2)$$

where $G_{\mu\nu}$ is the gluon field strength tensor and $\tilde{G}_{\mu\nu} = \frac{1}{4}\epsilon_{\mu\nu\rho\sigma}G^{\rho\sigma}$ is the corresponding dual field strength tensor and A_S and A'_S are effective couplings. We incorporated Eq. (2.1) in `FeynRules` [73, 74] and used `SusHi` [75, 76] in order to estimate the NNLO production cross-section via gluon fusion. The gluon fusion cross section at the large top mass limit at NNLO is given by

$$\sigma(pp \rightarrow gg \rightarrow S) = \rho_t^{S^2} \left(76.35 \cos^2 \alpha + 176.32 \sin^2 \alpha \right) \text{ pb}, \quad (2.3)$$

where the first term is the contribution from the CP even part, and the second is the CP odd contribution. The numerical coefficients are obtained from `SusHi` [75, 76].

For our analysis we have considered that the scalar S can decay via the following channels [77],

$$S \rightarrow f\bar{f}, VV^*, gg, \gamma\gamma \text{ and } Z\gamma. \quad (2.4)$$

2.2 Allowed Parameter Space

In order to explain all the anomalies, the scalar S must be coupled to third-generation fermions as well as massive gauge bosons. We fix the CPV parameter α and scan the free parameters ρ_f^S to satisfy the excesses defined in Eq. (1.1), Eq. (1.4) and Eq. (1.5). We have shown the parameter space in Fig. 1, which can simultaneously explain all the excesses. For a CP-even scalar, the cross-section is relatively smaller, as shown in Eq. (2.3), and consequently, a large value of ρ_t is required, as depicted in the top left panel of Fig. 1. For a mixed CP state, the required values of ρ_t decreases as the cross-section increases due to a pseudoscalar component, the parameter is shown in the top-right panel of Fig. 1. In the bottom panel, we have considered a pure pseudoscalar model. Note that such a pure pseudoscalar scenario can not explain the LEP $b\bar{b}$ excess. Thus, the g_{SVV} is redundant, and the parameter space does not depend on g_{SVV} . In all cases, the colour represents the ρ_b value. From the figure, it is evident that it is possible to satisfy all three excesses simultaneously.

If we exclude the LEP excess, we can take $g_{SVV} = 0$. Such a choice alters the decay spectrum of S as $S \rightarrow Zff$ is not possible. In addition, it is possible to assume $\rho_b^S = 0$ to explain the LHC excesses. Hence, to explain the LHC anomalies, as given in Eq. (1.1)-(1.4), a lower value of $\rho_{t/\tau}^S$ compared to the parameter space shown in Fig 1 will be enough, which corroborates [22]. Note that in Fig. 1, the parameter space does not depend on the sign of the angle α , and thus, the results are identical for both $\pm\alpha$. As discussed in the next section, the CP observables will distinguish $+\alpha$ from $-\alpha$.




Figure 1. The allowed parameter space for the CP mixing angles $\alpha = 0, \pi/4$ and $\pi/2$. For the case of $\pi/2$, we cannot explain the LEP $b\bar{b}$ excess, and the allowed region is for the CMS $\tau^+\tau^-$ and $\gamma\gamma$ excesses. We can explain all the excesses for $\alpha = 0$ and $\pi/4$.

3 Uncovering the CP Nature of the Resonance

The CP transformation property of a scalar particle can be inferred from its decay to tau pairs since the correlation of the tau lepton polarisation planes is governed by the CP properties of the parent particle. From the reconstruction of the decay planes of the tau leptons, it is possible to infer the CP properties of the scalar particle.

The differential decay width of the scalar boson can be written as [62, 63, 78]

$$d\Gamma_{S \rightarrow \tau^+\tau^-} \propto 1 - \frac{\pi^2}{16} b(E_+) b(E_-) \cos(\phi_{CP} - 2\alpha), \quad (3.1)$$

where $b(E_{\pm})$ are the spectral function describing the spin analyzing power of a given decay mode [63, 79]. For $\tau \rightarrow \pi\nu$ decay, $b(E_{\pm}) = 1$, and for other decay channels, it depends on the energy of the charged hadron at their respective tau-lepton rest frame. There are several methods [56–64] to estimate ϕ_{CP} and in this work we have used Impact Parameter (IP) method [62, 64] and the ρ

decay plane method [56, 64, 65, 67] for hadronic τ decay. For completeness, we have described the methods briefly.

3.1 Impact Parameter Method

In the Impact Parameter (IP) method [62, 64], the τ -lepton decay plane is constructed using the 3-momentum vector (\vec{q}^\pm) and the three-dimensional (3D) impact parameter vector (\vec{n}_\pm) of the charged particles. The method can be used for any one-prong decay of the tau lepton, and in this work we have implemented the method for $S \rightarrow \tau^+ \tau^- \rightarrow (\pi^+ \nu_\tau)(\pi^- \bar{\nu}_\tau)$ decay. The IP vector in the laboratory frame is defined as the distance of the closest approach of the charged particle's (π^\pm) track to the reconstructed primary vertex (PV) of an event. It is possible to reconstruct the 3D IP vector using the distance between the point of closest approach and the primary vertex in the $x-y$ -plane (d_0) and the z -coordinate of the point (z_0) [80]. We have used d_0 and z_0 and charged track momentum (highest p_T charged track inside a τ -tagged jet) provided by `Delphes` to find out the IP vector using the procedure describe in [80]. The IP vector then converted to a four vector $n_\pm^\mu = (0, \vec{n}_\pm)^T$ and boosted to zero-momentum frame (ZMF), the $\pi^+ \pi^-$ rest frame. The boosted IP four-vectors are denoted as $n_\pm^{*\mu}$ and we define the angle,

$$\phi^* = \arccos(\vec{n}_{+\perp}^* \cdot \vec{n}_{-\perp}^*), \quad (3.2)$$

where $n_{\pm\perp}^{*\mu}$ is the perpendicular component of $n_\pm^{*\mu}$ along the charged track h^\pm . To obtain a signed angle, we define the triple product, $\mathcal{O}(= q^{*-} \cdot (\vec{n}_{+\perp}^* \cdot \vec{n}_{-\perp}^*))$ where q^{*-} is the 3-momentum vector of π^- in ZMF. The CP observable is given by,

$$\phi_{CP} = \begin{cases} \tilde{\phi}^* + \pi & \mathcal{O} \geq 0 \\ \tilde{\phi}^* & \mathcal{O} < 0. \end{cases} \quad (3.3)$$

3.2 Rho Decay Plane Method

When the τ lepton decays via the rho meson, $\tau \rightarrow \rho^\pm \nu_\tau \rightarrow \pi^\pm \pi^0 \nu_\tau$, we use the information from the charged and neutral pion to define a CP-odd observable. In the $\pi^+ \pi^-$ rest frame, the acoplanarity angle ϕ^* is defined between the plane spanned by the (π^+, π^0) and (π^-, π^0) vectors. The acoplanarity angle is given by [56, 64, 80],

$$\phi^* = \arccos(\vec{q}_{+\perp}^{*0} \cdot \vec{q}_{-\perp}^{*0}), \quad (3.4)$$

where $\vec{q}_{+(-)\perp}^{*0}$ is the 3-momentum of the π^0 associated with $\pi^{+(-)}$ in the $\pi^+ \pi^-$ rest-frame transverse to the direction of the associated charged pion. The signed angle $\tilde{\phi}^*$ is obtained using the CP-odd triple correlation $\mathcal{O}_{CP}(= q^{*-} \cdot (q_{+\perp}^{*0} \times q_{-\perp}^{*0}))$ and we define

$$\tilde{\phi}^* = \begin{cases} \phi^* & \mathcal{O}_{CP} \geq 0 \\ 2\pi - \phi^* & \mathcal{O}_{CP} < 0. \end{cases} \quad (3.5)$$

In the $S \rightarrow \tau^+ \tau^- \rightarrow (\pi^+ \pi^0 \nu_\tau)(\pi^- \pi^0 \bar{\nu}_\tau)$ decay, the spin analyzing power is proportional to $y_\rho = \frac{(E_{\pi^+} - E_{\pi^0})}{(E_{\pi^+} + E_{\pi^0})} \times \frac{(E_{\pi^-} - E_{\pi^0})}{(E_{\pi^-} + E_{\pi^0})}$ at the respective τ^\pm rest frame [56]. Since y_ρ can be both positive and negative and if we integrate over the charged and neutral pion moments, ϕ_{CP} does not display CP sensitivity. We categorize the events depending on y_ρ and define the final CP sensitive variable,

$$\phi_{CP} = \begin{cases} \tilde{\phi}^* + \pi & y_\rho < 0 \\ \tilde{\phi}^* & \text{otherwise.} \end{cases} \quad (3.6)$$

To estimate y_ρ , we have calculated the energies of the pions in the lab frame, as it is not possible to reconstruct the τ rest frames. The choice of lab frame energy does not significantly reduce the asymmetry in ϕ_{CP} as long as the ρ mesons are energetic ($p_T(\rho^\pm) > 20$ GeV) [64] which is in general true for LHC.

3.3 Combined IP and Rho Decay Plane Methods

In the case of mixed decay, $\tau^+\tau^- \rightarrow (\pi\nu)(\rho\nu)$, we can combine the above-mentioned methods to define a CP-sensitive observable. This is possible if we consider the $\pi^+\pi^-$ rest frame for the ρ Decay plane method.

In the case of $\tau^+\tau^- \rightarrow (\rho^+\nu_\tau)(\pi^-\bar{\nu}_\tau)$ decay, the angle is defined by [64],

$$\phi^* = \arccos(\vec{q}_{+\perp}^{*0} \cdot \vec{n}_{-\perp}^*), \quad (3.7)$$

where, \vec{q}_{\pm}^* is the 3-momentum of π^{\pm} and \vec{n}_{-}^* is the IP vector of π^- and all the vectors are in the $\pi^+\pi^-$ rest frame. As before, we can define a signed angle depending on the value of the triple product $\mathcal{O}_3(= \vec{q}_{-}^* \cdot (\vec{q}_{+\perp}^{*0} \times \vec{n}_{-\perp}^*))$.

On the other hand, if τ^+ decay to $\pi^+\nu_\tau$, we can define ϕ^* and the triple product appropriately. Note that, in both cases of mixed decay, we have considered the event classification based on $y_{\pm}(= (E_{\pi^{\pm}} - E_{\pi^0}))$.

3.4 Event Simulation and Selection

We implemented our simplified model for event simulation in `FeynRules-2.3`[74]. The signal and the background events are generated using `MadGraph5-aMC@NLO-3.5.3`[81, 82]. To handle the τ polarization, we have used the `TauDecay` [83] package in `MadGraph5-aMC@NLO-3.5.3`. For showering and hadronization we used `PYTHIA-8.3` [84, 85], and detector effects were simulated using `Delphes-3.5.0`[86]. We use anti-kt algorithm [87] for jet clustering with radius parameter $R = 0.5$ and $p_T(j) > 20$ GeV. For τ -tagging we have used medium tag point with 70% tagging and 5×10^{-3} mis-tagging efficiency. The signal events are classified as two τ -tagged jets with $p_T > 40$ GeV and $|\eta| < 2.1$. We veto events with any b -tagged jets.

The background for such signal at the LHC is dominated by $Z \rightarrow \tau^+\tau^-$, $W^{\pm} + jets$ and QCD multi-jet events. We have generated sample background data to understand the nature of BG distribution for CP sensitive observable. As we will discuss later, we have used actual data provided by the CMS collaboration through HEPDATA [88] and rescaled it accordingly for different luminosity.

We obtain the charged and neutral pions from the τ -jet constituents. From all the tracks within each τ -jet, the track with the highest p_T is considered to be the charged pion. Neutral pions decay promptly into a pair of photons and therefore deposit their energies in the electromagnetic calorimeter, ECAL. From all the energy deposits, called towers, we estimate the one with the highest E_T and assign the tower to the neutral pion. For the combination method, the charged pion track with higher p_T is associated with the two-body decay of the τ ($\tau \rightarrow \pi\nu_\tau$).

4 Results

Now, we show how to uncover the CP properties of the scalar particle at the LHC. For a realistic representation of signal and background events, we have used the present CMS data of the $\tau^+\tau^-$ excess [8, 10]. The data is taken from HEPDATA[88, 89]. For our analysis, we have selected the events for which the invariant mass $m_{\tau^+\tau^-}$ lies within 60 – 120 GeV and we have selected the events with $p_T^{\tau^+\tau^-} > 200$ GeV obtained by the vectorial p_T sum of the visible decay products of tau lepton. This helps to minimise the background events significantly, as evident from Fig. 9 in ref [9].

To demonstrate the CP nature of the scalar, we plot the ϕ_{CP} for both signal and background events. The distribution peaks at a particular value of ϕ_{CP} for a given CP-admixture, whereas the background remains primarily flat. This enables estimating the CP nature from the ϕ_{CP} distribution alone. We found that with an integrated luminosity of 138 fb^{-1} , the total event distribution remains within statistical uncertainty of the background ($\sqrt{N_B}$). We considered high




Figure 2. (Top panel) Distribution of CP Observable for CP-even and CP-odd scalars with the background events with integrated luminosity of $3000 fb^{-1}$. (Bottom panel) Same distribution for CP-Mix($\alpha = \pm\pi/4$) states.

α	ϕ_{CP} Window	\mathcal{S}_0	$\mathcal{S}_{\sqrt{b}}$	$\mathcal{S}_{1\%}$	$\mathcal{S}_{10\%}$
0	$120^\circ - 240^\circ$	3.10	2.18	2.01	0.26
$\pi/2$	$0^\circ - 80^\circ$ & $280^\circ - 360^\circ$	3.51	2.47	2.09	0.26
$\pi/4$	$200^\circ - 320^\circ$	3.13	2.2	2.03	0.27
$-\pi/4$	$40^\circ - 160^\circ$	3.12	2.2	2.03	0.27

Table 1. Significance of different CP eigenstates with various background uncertainties at HL-LHC.

luminosity LHC (HL-LHC) [71, 72], and appropriately rescaled the existing data to an integrated luminosity of $3000 fb^{-1}$. The ϕ_{CP} distribution for signal and background events for HL-LHC is shown in Fig. 2, where background events are shown in yellow with statistical uncertainty bars, and the signal events are shown in green. As anticipated, the background events exhibit a near-uniform distribution in all the cases. The top panel depicts the distribution for CP-Even (top-left) and CP-odd (top-right) scalar, whereas the bottom panels depict maximally CP-Mixed states. From all the figures, it is evident that for a given CP nature of the resonance, the cumulative number of signal and background events rise beyond the background uncertainty, signalling the CP property of the scalar.

In order to quantify the result, we have chosen a window in ϕ_{CP} distribution and estimated a signal significance using the following formula [90],

$$\mathcal{S}_{\sigma_b} = \sqrt{2} \left\{ (N_S + N_B) \ln \left[\frac{(N_S + N_B)(N_B + \sigma_b^2)}{N_B^2 + (N_S + N_B)\sigma_b^2} \right] - \frac{N_B^2}{\sigma_b^2} \ln \left[1 + \frac{\sigma_B^2 N_S}{N_B(N_B + \sigma_b^2)} \right] \right\}^{1/2}, \quad (4.1)$$

where $N_{S(B)}$ denotes signal(background) events and σ_b denotes the background uncertainty. In Tab. 4 we have tabulated the significance for an optimal ϕ_{CP} window with various background uncertainty to achieve a robust estimation of the Significance. As expected, vanishing BG uncertainty provides a significance of more than 3σ at HL-LHC. As uncertainty increases, Significance reduces but remains large enough if we consider the background uncertainty solely statistical ($\sqrt{N_B}$) or 1%. If σ_b becomes large ($\sim 10\%$), the number of signal events need to be higher to achieve any significance.

5 Conclusions

In summary, under the assumption that several anomalies seen in the invariant mass of $\gamma\gamma$, $\tau^+\tau^-$ (from LHC data) and $b\bar{b}$ (from LEP data) events are due to a single spin-0 object with mass 95 GeV having Yukawa couplings to third generation fermions, we have performed a detector level analysis aimed at extracting the CP properties of such an object. We have done so limitedly to the $\tau^+\tau^-$ final state, as it offers the possibility of accessing the corresponding CP quantum numbers whichever the τ decay, whether leptonic or hadronic, through the relative angle between the two decay planes defined through the visible particles emerging from the τ lepton decays. Our approach has been based on a simplified model, which is characterized by the relevant Yukawa interactions being defined in terms of correcting factors to the corresponding SM vertex strengths while allowing for both a scalar and pseudoscalar interaction as well as any superposition thereof.

We have found that, under the assumption that the $\gamma\gamma$ and $\tau^+\tau^-$ anomalies will persist throughout the LHC era, the HL-LHC, in standard energy and luminosity configurations, will offer the chance to extract the CP properties of such a new spin-0 resonance. Once the latter is combined with cross-section and BR information, it would then be possible to map the results obtained in our model-independent parameterization into well-defined models like, e.g., a 2HDM of whichever Type, so long that it can host a suitable 95 GeV state in its Higgs particle spectrum, as it has indeed been proven to be the case in previous literature.

We therefore advocate experimental analyses by ATLAS and CMS aimed at implementing our approach, or indeed improved versions of it, with upcoming datasets at the CERN machine.

Acknowledgements

T.M. and P.S. would like to thank Artur Gottmann and Ravindra Kumar Verma for some useful discussions. T.M. is supported by the BITS Pilani Grant NFSG/PIL/2023/P3801. S.M. is supported in part through the NExT Institute and STFC Consolidated Grant No. ST/L000296/1.

References

- [1] ATLAS collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [1207.7214].
- [2] CMS collaboration, *Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [1207.7235].
- [3] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *Errata for the Higgs hunter's guide*, [hep-ph/9302272](https://arxiv.org/abs/hep-ph/9302272).

- [4] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, *Theory and phenomenology of two-Higgs-doublet models*, *Phys. Rept.* **516** (2012) 1 [[1106.0034](#)].
- [5] S. Moretti and S. Khalil, *Supersymmetry Beyond Minimality: From Theory to Experiment*. CRC Press, 2019.
- [6] S. De Curtis, L. Delle Rose, S. Moretti and K. Yagyu, *A Concrete Composite 2-Higgs Doublet Model*, *JHEP* **12** (2018) 051 [[1810.06465](#)].
- [7] S. De Curtis, L. Delle Rose, S. Moretti and K. Yagyu, *Supersymmetry versus Compositeness: 2HDMs tell the story*, *Phys. Lett. B* **786** (2018) 189 [[1803.01865](#)].
- [8] CMS collaboration, *Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV*, *Phys. Lett. B* **793** (2019) 320 [[1811.08459](#)].
- [9] CMS collaboration, *Searches for additional Higgs bosons and for vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2023) 073 [[2208.02717](#)].
- [10] CMS collaboration, “Search for a standard model-like Higgs boson in the mass range between 70 and 110 GeV in the diphoton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV.” <https://cds.cern.ch/record/2852907/>, CERN, Geneva, 2023.
- [11] C. Arcangeletti, “ATLAS, LHC Seminar.” <https://indico.cern.ch/event/1281604/>, 2023.
- [12] ATLAS collaboration, “Search for diphoton resonances in the 66 to 110 GeV mass range using 140 fb^{-1} of 13 TeV pp collisions collected with the ATLAS detector.” <http://cds.cern.ch/record/2862024>, CERN, Geneva, 2023.
- [13] CMS collaboration, “Search for dilepton resonances from decays of (pseudo)scalar bosons produced in association with a massive vector boson or top quark anti-top quark pair at $\sqrt{s} = 13$ TeV.” <http://cds.cern.ch/record/2815307>, CERN, Geneva, 2022.
- [14] LEP WORKING GROUP FOR HIGGS BOSON SEARCHES, ALEPH, DELPHI, L3, OPAL collaboration, *Search for the standard model Higgs boson at LEP*, *Phys. Lett. B* **565** (2003) 61 [[hep-ex/0306033](#)].
- [15] ALEPH, DELPHI, L3, OPAL, LEP WORKING GROUP FOR HIGGS BOSON SEARCHES collaboration, *Search for neutral MSSM Higgs bosons at LEP*, *Eur. Phys. J. C* **47** (2006) 547 [[hep-ex/0602042](#)].
- [16] J. Cao, X. Guo, Y. He, P. Wu and Y. Zhang, *Diphoton signal of the light Higgs boson in natural NMSSM*, *Phys. Rev. D* **95** (2017) 116001 [[1612.08522](#)].
- [17] S. Heinemeyer, C. Li, F. Lika, G. Moortgat-Pick and S. Paasch, *Phenomenology of a 96 GeV Higgs boson in the 2HDM with an additional singlet*, *Phys. Rev. D* **106** (2022) 075003 [[2112.11958](#)].
- [18] T. Biekötter, A. Grohsjean, S. Heinemeyer, C. Schwanenberger and G. Weiglein, *Possible indications for new Higgs bosons in the reach of the LHC: N2HDM and NMSSM interpretations*, *Eur. Phys. J. C* **82** (2022) 178 [[2109.01128](#)].
- [19] T. Biekötter, M. Chakraborti and S. Heinemeyer, *A 96 GeV Higgs boson in the N2HDM*, *Eur. Phys. J. C* **80** (2020) 2 [[1903.11661](#)].
- [20] J. Cao, X. Jia, Y. Yue, H. Zhou and P. Zhu, *96 GeV diphoton excess in seesaw extensions of the natural NMSSM*, *Phys. Rev. D* **101** (2020) 055008 [[1908.07206](#)].
- [21] T. Biekötter, S. Heinemeyer and G. Weiglein, *Excesses in the low-mass Higgs-boson search and the W-boson mass measurement*, *Eur. Phys. J. C* **83** (2023) 450 [[2204.05975](#)].
- [22] S. Iguro, T. Kitahara and Y. Omura, *Scrutinizing the 95–100 GeV di-tau excess in the top associated process*, *Eur. Phys. J. C* **82** (2022) 1053 [[2205.03187](#)].
- [23] W. Li, J. Zhu, K. Wang, S. Ma, P. Tian and H. Qiao, *A light Higgs boson in the NMSSM confronted with the CMS di-photon and di-tau excesses*, [2212.11739](#).

- [24] J. M. Cline and T. Toma, *Pseudo-Goldstone dark matter confronts cosmic ray and collider anomalies*, *Phys. Rev. D* **100** (2019) 035023 [[1906.02175](#)].
- [25] T. Biekötter and M. O. Olea-Romacho, *Reconciling Higgs physics and pseudo-Nambu-Goldstone dark matter in the S2HDM using a genetic algorithm*, *JHEP* **10** (2021) 215 [[2108.10864](#)].
- [26] A. Crivellin, J. Heeck and D. Müller, *Large $h \rightarrow bs$ in generic two-Higgs-doublet models*, *Phys. Rev. D* **97** (2018) 035008 [[1710.04663](#)].
- [27] G. Cacciapaglia, A. Deandrea, S. Gascon-Shotkin, S. Le Corre, M. Lethuillier and J. Tao, *Search for a lighter Higgs boson in Two Higgs Doublet Models*, *JHEP* **12** (2016) 068 [[1607.08653](#)].
- [28] A. A. Abdelalim, B. Das, S. Khalil and S. Moretti, *Di-photon decay of a light Higgs state in the BLSSM*, *Nucl. Phys. B* **985** (2022) 116013 [[2012.04952](#)].
- [29] T. Biekötter, S. Heinemeyer and G. Weiglein, *Mounting evidence for a 95 GeV Higgs boson*, *JHEP* **08** (2022) 201 [[2203.13180](#)].
- [30] T. Biekötter, S. Heinemeyer and G. Weiglein, *The CMS di-photon excess at 95 GeV in view of the LHC Run 2 results*, *Phys. Lett. B* **846** (2023) 138217 [[2303.12018](#)].
- [31] D. Azevedo, T. Biekötter and P. M. Ferreira, *2HDM interpretations of the CMS diphoton excess at 95 GeV*, *JHEP* **11** (2023) 017 [[2305.19716](#)].
- [32] T. Biekötter, S. Heinemeyer and G. Weiglein, *95.4 GeV diphoton excess at ATLAS and CMS*, *Phys. Rev. D* **109** (2024) 035005 [[2306.03889](#)].
- [33] J. Cao, X. Jia and J. Lian, *Unified Interpretation of Muon $g-2$ anomaly, 95 GeV Diphoton, and $b\bar{b}$ Excesses in the General Next-to-Minimal Supersymmetric Standard Model*, [2402.15847](#).
- [34] K. Wang and J. Zhu, *A 95 GeV light Higgs in the top-pair-associated diphoton channel at the LHC in the Minimal Dilaton Model*, [2402.11232](#).
- [35] W. Li, H. Qiao, K. Wang and J. Zhu, *Light dark matter confronted with the 95 GeV diphoton excess*, [2312.17599](#).
- [36] P. S. B. Dev, R. N. Mohapatra and Y. Zhang, *Explanation of the 95 GeV $\gamma\gamma$ and $b\bar{b}$ excesses in the minimal left-right symmetric model*, *Phys. Lett. B* **849** (2024) 138481 [[2312.17733](#)].
- [37] D. Borah, S. Mahapatra, P. K. Paul and N. Sahu, *Scotogenic $U(1)_{L\mu-L\tau}$ origin of $(g-2)\mu$, W -mass anomaly and 95 GeV excess*, *Phys. Rev. D* **109** (2024) 055021 [[2310.11953](#)].
- [38] J. Cao, X. Jia, J. Lian and L. Meng, *95 GeV diphoton and $b\bar{b}$ excesses in the general next-to-minimal supersymmetric standard model*, *Phys. Rev. D* **109** (2024) 075001 [[2310.08436](#)].
- [39] U. Ellwanger and C. Hugonie, *Additional Higgs Bosons near 95 and 650 GeV in the NMSSM*, *Eur. Phys. J. C* **83** (2023) 1138 [[2309.07838](#)].
- [40] J. A. Aguilar-Saavedra, H. B. Câmara, F. R. Joaquim and J. F. Seabra, *Confronting the 95 GeV excesses within the $U(1)'$ -extended next-to-minimal 2HDM*, *Phys. Rev. D* **108** (2023) 075020 [[2307.03768](#)].
- [41] S. Ashanujjaman, S. Banik, G. Coloretti, A. Crivellin, B. Mellado and A.-T. Mulaudzi, *$SU(2)_L$ triplet scalar as the origin of the 95 GeV excess?*, *Phys. Rev. D* **108** (2023) L091704 [[2306.15722](#)].
- [42] S. Bhattacharya, G. Coloretti, A. Crivellin, S.-E. Dahbi, Y. Fang, M. Kumar et al., *Growing Excesses of New Scalars at the Electroweak Scale*, [2306.17209](#).
- [43] J. Dutta, J. Lahiri, C. Li, G. Moortgat-Pick, S. F. Tabira and J. A. Ziegler, *Dark Matter Phenomenology in 2HDMs in light of the 95 GeV excess*, [2308.05653](#).
- [44] U. Ellwanger and C. Hugonie, *NMSSM with correct relic density and an additional 95 GeV Higgs boson*, [2403.16884](#).
- [45] M. A. Diaz, G. Cerro, S. Dasmahapatra and S. Moretti, *Bayesian Active Search on Parameter Space: a 95 GeV Spin-0 Resonance in the $(B - L)$ SSM*, [2404.18653](#).

- [46] U. Ellwanger, C. Hugonie, S. F. King and S. Moretti, *NMSSM Explanation for Excesses in the Search for Neutralinos and Charginos and a 95 GeV Higgs Boson*, [2404.19338](#).
- [47] S. Y. Ayazi, M. Hosseini, S. Paktinat Mehdiabadi and R. Rouzbehi, *The Vector Dark Matter, LHC Constraints Including a 95 GeV Light Higgs Boson*, [2405.01132](#).
- [48] A. Arhrib, K. H. Phan, V. Q. Tran and T.-C. Yuan, *When Standard Model Higgs Meets Its Lighter 95 GeV Higgs*, [2405.03127](#).
- [49] J. Gao, J. Ma, L. Wang and H. Xu, *A 95 GeV Higgs boson and spontaneous CP-violation at the finite temperature*, [2408.03705](#).
- [50] R. Benbrik, M. Boukidi, S. Moretti and S. Sehlali, *Explaining the 96 GeV Di-photon anomaly in a generic 2HDM Type-III*, *Phys. Lett. B* **832** (2022) 137245 [[2204.07470](#)].
- [51] R. Benbrik, M. Boukidi, S. Moretti and S. Sehlali, *Probing a 96 GeV Higgs Boson in the Di-Photon Channel at the LHC*, *PoS ICHEP2022* (2022) 547 [[2211.11140](#)].
- [52] A. Belyaev, R. Benbrik, M. Boukidi, M. Chakraborti, S. Moretti and S. Sehlali, *Explanation of the Hints for a 95 GeV Higgs Boson within a 2-Higgs Doublet Model*, [2306.09029](#).
- [53] A. Belyaev, R. Benbrik, M. Boukidi, M. Chakraborti, S. Moretti and S. Sehlali, *Probing 95 GeV Higgs in the 2HDM Type-III*, *PoS LHCP2023* (2024) 299 [[2402.03998](#)].
- [54] R. Benbrik, M. Boukidi and S. Moretti, *Superposition of CP-Even and CP-Odd Higgs Resonances: Explaining the 95 GeV Excesses within a Two-Higgs Doublet Model*, [2405.02899](#).
- [55] A. Khanna, S. Moretti and A. Sarkar, *Explaining 95 (or so) GeV Anomalies in the 2-Higgs Doublet Model Type-I*, [2409.02587](#).
- [56] G. R. Bower, T. Pierzchala, Z. Was and M. Worek, *Measuring the Higgs boson's parity using tau \rightarrow rho nu*, *Phys. Lett. B* **543** (2002) 227 [[hep-ph/0204292](#)].
- [57] K. Desch, A. Imhof, Z. Was and M. Worek, *Probing the CP nature of the Higgs boson at linear colliders with tau spin correlations: The Case of mixed scalar - pseudoscalar couplings*, *Phys. Lett. B* **579** (2004) 157 [[hep-ph/0307331](#)].
- [58] S. Berge, W. Bernreuther and H. Spiesberger, *Higgs CP properties using the τ decay modes at the ILC*, *Phys. Lett. B* **727** (2013) 488 [[1308.2674](#)].
- [59] S. Berge, W. Bernreuther and J. Ziethe, *Determining the CP parity of Higgs bosons at the LHC in their tau decay channels*, *Phys. Rev. Lett.* **100** (2008) 171605 [[0801.2297](#)].
- [60] S. Berge and W. Bernreuther, *Determining the CP parity of Higgs bosons at the LHC in the tau to 1-prong decay channels*, *Phys. Lett. B* **671** (2009) 470 [[0812.1910](#)].
- [61] S. Berge, W. Bernreuther, B. Niepelt and H. Spiesberger, *How to pin down the CP quantum numbers of a Higgs boson in its tau decays at the LHC*, *Phys. Rev. D* **84** (2011) 116003 [[1108.0670](#)].
- [62] S. Berge, W. Bernreuther and S. Kirchner, *Determination of the Higgs CP-mixing angle in the tau decay channels at the LHC including the Drell-Yan background*, *Eur. Phys. J. C* **74** (2014) 3164 [[1408.0798](#)].
- [63] S. Berge, W. Bernreuther and S. Kirchner, *Determination of the Higgs CP-mixing angle in the tau decay channels*, *Nucl. Part. Phys. Proc.* **273-275** (2016) 841 [[1410.6362](#)].
- [64] S. Berge, W. Bernreuther and S. Kirchner, *Prospects of constraining the Higgs boson's CP nature in the tau decay channel at the LHC*, *Phys. Rev. D* **92** (2015) 096012 [[1510.03850](#)].
- [65] T. Han, S. Mukhopadhyay, B. Mukhopadhyaya and Y. Wu, *Measuring the CP property of Higgs coupling to tau leptons in the VBF channel at the LHC*, *JHEP* **05** (2017) 128 [[1612.00413](#)].
- [66] S. Antusch, O. Fischer, A. Hammad and C. Scherb, *Testing CP Properties of Extra Higgs States at the HL-LHC*, *JHEP* **03** (2021) 200 [[2011.10388](#)].

- [67] S. Dutta, A. Goyal, M. Kumar and A. K. Swain, *Measuring CP nature of $h\tau\bar{\tau}$ coupling at e^-p collider*, *Eur. Phys. J. C* **82** (2022) 530 [[2109.00329](#)].
- [68] W. Esmail, A. Hammad, M. Nojiri and C. Scherb, *Testing CP properties of the Higgs boson coupling to τ leptons with heterogeneous graphs*, [2409.06132](#).
- [69] CMS collaboration, *Searches for additional Higgs bosons and vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, tech. rep., CERN, Geneva, 2022. <https://cds.cern.ch/record/2803739>.
- [70] LEP WORKING GROUP FOR HIGGS BOSON SEARCHES, ALEPH, DELPHI, L3, OPAL collaboration, *Search for the standard model Higgs boson at LEP*, *Phys. Lett. B* **565** (2003) 61 [[hep-ex/0306033](#)].
- [71] F. Gianotti et al., *Physics potential and experimental challenges of the LHC luminosity upgrade*, *Eur. Phys. J. C* **39** (2005) 293 [[hep-ph/0204087](#)].
- [72] G. Apollinari, O. Brüning, T. Nakamoto and L. Rossi, *High Luminosity Large Hadron Collider HL-LHC*, *CERN Yellow Rep.* (2015) 1 [[1705.08830](#)].
- [73] N. D. Christensen and C. Duhr, *FeynRules - Feynman rules made easy*, *Comput. Phys. Commun.* **180** (2009) 1614 [[0806.4194](#)].
- [74] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 - A complete toolbox for tree-level phenomenology*, *Comput. Phys. Commun.* **185** (2014) 2250 [[1310.1921](#)].
- [75] R. V. Harlander, S. Liebler and H. Mantler, *SusHi: A program for the calculation of Higgs production in gluon fusion and bottom-quark annihilation in the Standard Model and the MSSM*, *Comput. Phys. Commun.* **184** (2013) 1605 [[1212.3249](#)].
- [76] R. V. Harlander, S. Liebler and H. Mantler, *SusHi Bento: Beyond NNLO and the heavy-top limit*, *Comput. Phys. Commun.* **212** (2017) 239 [[1605.03190](#)].
- [77] S. Y. Choi, J. S. Lee and J. Park, *Decays of Higgs bosons in the Standard Model and beyond*, *Prog. Part. Nucl. Phys.* **120** (2021) 103880 [[2101.12435](#)].
- [78] M. Kramer, J. H. Kuhn, M. L. Stong and P. M. Zerwas, *Prospects of measuring the parity of Higgs particles*, *Z. Phys. C* **64** (1994) 21 [[hep-ph/9404280](#)].
- [79] Y.-S. Tsai, *Decay Correlations of Heavy Leptons in $e^+e^- \rightarrow \text{Lepton}^+ \text{Lepton}^-$* , *Phys. Rev. D* **4** (1971) 2821.
- [80] M. C. Hansen, *Studies into measuring the Higgs CP-state in $H \rightarrow \tau\tau$ decays at ATLAS*, Ph.D. thesis, Bonn U., 2020.
- [81] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, *MadGraph 5 : Going Beyond*, *JHEP* **06** (2011) 128 [[1106.0522](#)].
- [82] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[1405.0301](#)].
- [83] K. Hagiwara, T. Li, K. Mawatari and J. Nakamura, *TauDecay: a library to simulate polarized tau decays via FeynRules and MadGraph5*, *Eur. Phys. J. C* **73** (2013) 2489 [[1212.6247](#)].
- [84] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten et al., *An Introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[1410.3012](#)].
- [85] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*, *SciPost Phys. Codeb.* **2022** (2022) 8 [[2203.11601](#)].
- [86] DELPHES 3 collaboration, *DELPHES 3, A modular framework for fast simulation of a generic collider experiment*, *JHEP* **02** (2014) 057 [[1307.6346](#)].
- [87] M. Cacciari, G. P. Salam and G. Soyez, *The Anti- $k(t)$ jet clustering algorithm*, *JHEP* **04** (2008) 063 [[0802.1189](#)].

- [88] E. Maguire, L. Heinrich and G. Watt, *HEPData: a repository for high energy physics data*, *J. Phys. Conf. Ser.* **898** (2017) 102006 [[1704.05473](#)].
- [89] CMS Collaboration, “Searches for additional higgs bosons and for vector leptoquarks in $\tau\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ tev.” <https://www.hepdata.net/record/ins2132368>, 2022. <https://doi.org/10.17182/hepdata.128147>.
- [90] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [[1007.1727](#)].