

Improvements for lower bounds of mutually orthogonal Latin squares of sizes 54, 96 and 108

R. Julian R. Abel *
 School of Mathematics and Statistics
 UNSW Sydney
 NSW 2052, Australia

Ingo Janiszczak** and Reiner Staszewski
 Faculty of Mathematics
 University of Duisburg-Essen
 45127 Essen, Germany

December 3, 2024

Abstract

We will show that there are at least 8, 10 and 9 mutually orthogonal Latin squares (MOLS) of orders $n = 54, 96$ and 108 . The cases $n = 54$ and 96 are obtained by constructing separable permutation codes consisting of 8×54 and 10×96 codeword respectively; in addition, these codes respectively have lengths 54, 96 and minimum distances 53, 95. Here we will follow exactly the procedure given in [6]. The case $n = 108$ is obtained by constructing a $(108, 10, 1)$ difference matrix. Also, an error in [1] for $n = 45$ will be corrected.

Keywords: bounds, difference matrix, DM, isometry, permutation code, permutation arrays, MOLS, mutually orthogonal Latin squares.

1 Introduction

Let n be a positive integer and let V and W be sets consisting of n elements. A Graeco-Latin square of order n is a $n \times n$ array $M = ((v_{ij}, w_{ij}))$ with entries in the Cartesian product $V \times W$ such that the set of all different entries in M is $V \times W$ and the arrays $L_1 = (v_{ij})$ and $L_2 = (w_{ij})$ are Latin squares, i.e. all rows and all columns of L_1 and L_2 are permutations of V and W respectively. Two such Latin squares are called orthogonal if the array $((v_{ij}, w_{ij}))$ is a Graeco-Latin square. A set of t mutually orthogonal Latin squares (MOLS) of order n (or t MOLS(n)) consists of t Latin squares of order n which are pairwise orthogonal. Let $N(n)$ be size of the largest set of MOLS of order n .

In this paper, we obtain improvements for $N(n)$ when $n \in \{54, 96, 108\}$. These MOLS determined using two different methods. For $n = 54$ and $n = 96$, permutation codes corresponding to the relevant MOLS were determined. In the case of $n = 108$, a difference matrix with 10 rows was constructed; from this difference matrix, 9 MOLS for $n = 108$ can be obtained.

Since parts of the present paper for the cases $n = 54$ and $n = 96$ are to be seen as a supplement to [6], we will simply extract the relevant part for the necessary notations and definitions.

Let X be an arbitrary finite set of order n . Let $S_X := \{\sigma : X \rightarrow X \mid \sigma \text{ is bijective}\}$. This group acts on X from the right, and for $x \in X, \sigma \in S_X$, we denote the image of x under σ by x^σ . If $X = \{1, 2, \dots, n\}$, then we will write S_n instead of S_X . For $\sigma, \tau \in S_X$ the Hamming distance $d_H(\sigma, \tau) = d(\sigma, \tau)$ equals n minus the number of fixed points of $\sigma^{-1}\tau$. A subset C of S_X is called a *permutation code* or *permutation*

*r.j.abel@unsw.edu.au

**ingo.janiszczak@uni-due.de

array of length n and of minimum distance $d = d(C)$ if

$$d(C) := \min \{d_H(\sigma, \tau) \mid \sigma, \tau \in C, \sigma \neq \tau\} .$$

Such a permutation array C is called an (n, d) -PA. If C is an $(n, n-1)$ -PA, then C is called (r, m) -separable if it is the join of m disjoint (n, n) -PAs P_1, \dots, P_m of cardinality r such that for all pairs P_s, P_t , $s \neq t$, the distance $d_H(\sigma, \tau)$ equals $n-1$ for all $\sigma \in P_s$ and all $\tau \in P_t$. When $r = n$, existence of an (n, m) -separable $(n, n-1)$ -PA is equivalent to existence of m MOLS of order n [3]. In this case we order the elements in P_s such that $P_s = \{\sigma_{sk} : 1 \leq k \leq n\}$ for all $1 \leq s \leq m$. From this m MOLS A_1, A_2, \dots, A_m of order n can be obtained as follows: $A_s(i, j) = k$ if $j^{\sigma_{sk}} = i$.

S_n is a metric space via the Hamming distance, and the isometry group $Iso(n)$ is isomorphic to the wreath product $S_n \wr S_2$ [4]. $Iso(n)$ can be described as a subgroup of S_{2n} . Let B_1 and B_2 be the naturally embedded subgroups isomorphic to S_n acting on the sets $\{1, 2, \dots, n\}$ and $\{n+1, n+2, \dots, 2n\}$, respectively. Moreover let $t_n := (1, n+1)(2, n+2) \cdots (n, 2n) \in S_{2n}$. Then $Iso(n) = \langle B_1, B_2, t_n \rangle = (B_1 \times B_2) : \langle t_n \rangle$, and the action of this group on B_1 from right is given by

$$b * x := \begin{cases} x^{-1}b & \text{if } x \in B_1 \\ b\varphi(x) & \text{if } x \in B_2 \\ b^{-1} & \text{if } x = t_n, \end{cases}$$

where $b \in B_1$, $x \in Iso(n)$ and φ denotes the natural isomorphism from B_2 to B_1 . Moreover, if U is a subgroup of $Iso(n)$ and $b \in B_1$, then $b * U := \{b * u \mid u \in U\}$ denotes the U -orbit of b . Our codes C will now be regarded as subsets of B_1 , and a code C will be called U -invariant if C is closed under the action of U . This means that C is a union of U -orbits. For the strategy of constructing (n, m) -separable PAs that are invariant under a given subgroup U of $Iso(n)$ we refer to [5]. But now we will only join separable U -orbits.

In section 2 we will prove the existence of 8 MOLS for $n = 54$ and 10 MOLS for $n = 96$ by constructing a $(54, 8)$ -separable PA and a $(96, 10)$ -separable PA. These are invariant under specific subgroups U of $Iso(n)$ using the described group action of $Iso(n)$ on S_n and will be given by generators of U and representatives of the U -orbits.

As all the data for the constructed PAs is contained in the symmetric group S_{2n} , it can easily be read into a computer algebra system such as MAGMA [2] or GAP [7]. The orbits can then be computed using elementary programs consisting of a few lines of GAP or MAGMA-code. In the appendix we give a GAP code for calculating the orbits, the resulting PA and the MOLS. There is also a routine to check if the PA is (n, m) -separable.

Now we define a difference matrix. Let G be a finite group of order n and let $k \geq 2$ be an integer. An $(n, k, 1)$ -difference matrix (DM) over G , or briefly $(n, k, 1)$ -DM over G , is a $k \times n$ matrix $D = (d_{i,j})$ with entries from G such that for any two distinct rows s and t , the multiset of differences $\{(d_{s,j})(d_{t,j})^{-1} : 1 \leq j \leq n\}$ contains each element of G exactly once. For most known MOLS obtained from difference matrices, the group G is abelian and the expression $d_{s,j} - d_{t,j}$ instead of $(d_{s,j})(d_{t,j})^{-1}$ is more commonly used.

If D is an $(n, k, 1)$ -DM over G , and the first row of D is normalised (that is, for all j , the entries in column j have been multiplied on the right by $d_{1,j}^{-1}$ so that the first entry in each column is 1, the identity element in G) then $k-1$ MOLS(n), L_1, L_2, \dots, L_{k-1} can be obtained as follows. First let $G = \{g_1 = 1, g_2, \dots, g_n\}$. Then for $j = 1, 2, \dots, n$, the j 'th row of L_m is obtained by multiplying the $(m+1)$ 'th row of D on the right by g_j . In section 2 we obtain a $(108, 10, 1)$ -DM which gives 9 MOLS(108).

2 Improvements

Let $n = 54$ and $U := \langle x_1, x_2, x_3, x_4, x_5 \rangle \leq Iso(54)$ of order 243 generated by x_1, x_2, x_3, x_4, x_5 given in cycle structure

$x_1 := (1, 6, 26)(2, 4, 27)(3, 5, 25)(7, 10, 13)(8, 11, 14)(9, 12, 15)(16, 21, 23)(17, 19, 24)(18, 20, 22)(28, 33, 53)$
 $(29, 31, 54)(30, 32, 52)(34, 37, 40)(35, 38, 41)(36, 39, 42)(43, 48, 50)(44, 46, 51)(45, 47, 49)(55, 60, 80)$
 $(56, 58, 81)(57, 59, 79)(61, 64, 67)(62, 65, 68)(63, 66, 69)(70, 75, 77)(71, 73, 78)(72, 74, 76)(82, 87, 107)$
 $(83, 85, 108)(84, 86, 106)(88, 91, 94)(89, 92, 95)(90, 93, 96)(97, 102, 104)(98, 100, 105)(99, 101, 103),$
 $x_2 := (1, 11, 19)(2, 12, 20)(3, 10, 21)(4, 15, 22)(5, 13, 23)(6, 14, 24)(7, 16, 25)(8, 17, 26)(9, 18, 27)$
 $(28, 38, 46)(29, 39, 47)(30, 37, 48)(31, 42, 49)(32, 40, 50)(33, 41, 51)(34, 43, 52)(35, 44, 53)(36, 45, 54),$
 $x_3 := (1, 2, 3)(4, 5, 6)(7, 8, 9)(10, 11, 12)(13, 14, 15)(16, 17, 18)(19, 20, 21)(22, 23, 24)(25, 26, 27)$
 $(28, 29, 30)(31, 32, 33)(34, 35, 36)(37, 38, 39)(40, 41, 42)(43, 44, 45)(46, 47, 48)(49, 50, 51)(52, 53, 54),$
 $x_4 := (55, 65, 73)(56, 66, 74)(57, 64, 75)(58, 69, 76)(59, 67, 77)(60, 68, 78)(61, 70, 79)(62, 71, 80)(63, 72, 81)$
 $(82, 92, 100)(83, 93, 101)(84, 91, 102)(85, 96, 103)(86, 94, 104)(87, 95, 105)(88, 97, 106)(89, 98, 107)$
 $(90, 99, 108),$
 $x_5 := (55, 56, 57)(58, 59, 60)(61, 62, 63)(64, 65, 66)(67, 68, 69)(70, 71, 72)(73, 74, 75)(76, 77, 78)(79, 80, 81)$
 $(82, 83, 84)(85, 86, 87)(88, 89, 90)(91, 92, 93)(94, 95, 96)(97, 98, 99)(100, 101, 102)(103, 104, 105)$
 $(106, 107, 108).$

Furthermore let

$a_1 := (1, 34, 20, 53, 10, 45)(2, 51, 12, 22, 43, 23, 9, 47, 26)(3, 13, 35, 14, 25, 37, 18, 21, 42)$
 $(4, 17, 28, 7, 11, 32, 19, 6, 54)(5, 46, 15, 30, 24, 39)(8, 41)(16, 50)(27, 31)(29, 48, 38)(36, 44, 52),$
 $a_2 := (1, 4, 21, 15, 27, 5, 30, 19, 16, 2, 7, 23, 18, 50, 29, 28, 36, 49, 14, 48, 45, 17, 13, 10, 25, 40, 33, 47)$
 $(3, 43, 42, 39, 54, 44, 53, 31, 41, 24)(6, 9, 12, 22, 37, 46, 38, 11, 34, 35, 26, 20, 52, 8, 32),$
 $a_3 := (1, 38, 26, 35, 6, 41)(2, 39, 27, 36, 4, 42)(3, 37, 25, 34, 5, 40)(16, 30, 51, 18, 29, 50, 17, 28, 49)$
 $(19, 33, 45, 21, 32, 44, 20, 31, 43)(22, 54, 48, 24, 53, 47, 23, 52, 46),$
 $a_4 := (1, 41, 37, 19, 43, 8, 28, 9, 48, 35, 4, 22, 34, 13, 38, 10)(2, 32, 6, 5, 14, 30, 16, 17, 7, 39)$
 $(3, 49, 11, 21, 53, 50)(15, 47, 45, 18, 27, 31)(20, 36, 24, 44, 25, 51)(23, 54, 42, 29, 26, 40, 46, 52, 33),$
 $a_5 := (1, 25, 11, 23, 8, 34, 35, 43, 30, 32, 41, 36, 54)(2, 9, 45, 46, 48, 38, 22, 27)$
 $(3, 17, 5, 33, 50, 7, 53, 20, 47, 29, 51, 18, 15, 12, 4, 49, 26, 21, 28, 42, 44, 39, 6, 40, 52, 10, 14)(13, 19, 37)(16, 24),$
 $a_6 := (1, 54, 48, 40, 20, 34, 6, 38, 51, 41, 52, 36, 53, 15, 4, 19, 12, 21, 16, 14, 10, 44, 43, 28, 17, 25, 18, 33, 3)$
 $(2, 9, 8, 22, 47, 27, 5, 13, 30, 46, 29, 31, 35, 39, 37, 7, 42, 49, 11, 26, 50, 45, 23),$
 $a_7 := (1, 23, 5, 14, 52, 36, 17, 8, 25, 12, 11, 6, 45, 27, 30, 19, 15, 9, 38, 28, 48, 10, 49, 50, 44, 29, 51,$
 $24, 35, 46, 39, 31, 33, 4, 26, 18, 47, 40, 34, 43, 53, 37, 2, 3, 41, 13, 22, 16)(7, 21, 32, 54),$
 $a_8 := (1, 35, 17, 45, 25, 11, 24, 51, 12, 43, 47)(2, 29, 16, 33, 52, 9, 49, 4, 32, 14, 18)(3, 15, 46, 8, 21, 54, 22, 26, 40)$
 $(5, 7, 34, 6, 39, 41, 23, 30, 10, 38, 20)(13, 42, 36, 37, 27, 53, 28, 31, 19, 48, 50, 44),$
 $a_9 := (1, 27, 29, 21, 26, 36, 8, 23, 44, 54, 52, 14, 32, 5, 31)(2, 47, 39, 18, 24, 11, 40)(3, 42, 51)$
 $(4, 12, 25, 22, 33, 50, 49)(6, 45, 13, 10, 48, 16, 35, 15, 43, 9, 34, 53, 7, 30, 37, 19, 46, 20, 41)(17, 28),$
 $a_{10} := (1, 18, 34, 33, 12, 7, 36, 51)(2, 27, 44, 49, 11, 25, 53, 32)(3, 9, 54, 40, 10, 16, 43, 42)$
 $(4, 28, 14, 29, 24, 37, 15, 39)(5, 38, 22, 48, 23, 30, 6, 46)(8, 45, 31, 20, 17, 52, 50, 19)(13, 47)(21, 26, 35, 41),$
 $a_{11} := (1, 32, 13, 53, 10, 24, 40, 22, 18, 29, 52, 41, 28, 19, 51, 37, 14, 33, 47, 44)$
 $(2, 30, 4, 26, 21, 5, 45, 16, 54, 27, 38, 3, 15, 7, 43, 31, 17, 11, 42, 25, 35, 20, 46, 12, 39, 36, 8)$
 $(6, 49, 9, 48, 23, 34, 50),$
 $a_{12} := (1, 17, 38, 45, 12, 25, 37, 54)(2, 8, 47, 53, 11, 7, 29, 43)(3, 26, 30, 34, 10, 16, 46, 35)$
 $(4, 24, 15, 13, 5, 14, 23, 22)(9, 39, 36, 21)(18, 28, 52, 19, 27, 48, 44, 20)(31, 32, 50, 33, 40, 42, 49, 41),$
 $a_{13} := (1, 15, 51, 31, 46, 18, 34, 53, 44, 50, 38, 37, 45, 35, 5, 41, 28, 36, 40, 19, 20, 4, 17, 21, 47, 54,$
 $23, 8, 25, 3, 2, 29, 30, 27, 16, 7, 52, 32, 42, 49)(6, 14, 26, 43, 13, 24, 33, 11, 39, 9, 10, 22),$
 $a_{14} := (1, 38, 21, 11, 7, 22, 6, 32, 46, 49, 34, 27, 44, 30, 54, 41, 15)(2, 37, 19, 10, 8, 24, 4, 31, 47, 51,$
 $35, 26, 45, 29, 52, 40, 13, 3, 39, 20, 12, 9, 23, 5, 33, 48, 50, 36, 25, 43, 28, 53, 42, 14)(16, 18),$
 $a_{15} := (1, 21, 19, 43, 2, 45, 49, 36, 3, 23, 30, 14, 4, 25, 42, 37, 52, 51, 18, 40, 16, 11, 22, 6, 29, 53)$
 $(7, 9, 12, 20, 24, 26, 10, 44, 47, 15, 28, 33, 38, 13, 27)(8, 41, 34, 48, 31, 17)(32, 35, 50, 39)(46, 54),$
 $a_{16} := (1, 42, 5, 44, 18, 48)(2, 34, 28, 6, 47, 50, 25, 33, 36)(3, 17, 4)(7, 41, 45, 12, 43, 38, 14, 29, 32)$
 $(8, 22, 21)(9, 37, 19, 31, 23, 35)(10, 26, 15)(11, 49, 13, 53, 27, 30)(16, 51, 54, 20, 52, 46, 24, 39, 40).$

For $i \in \{1, \dots, 16\}$ the a_i are codewords of B_1 and the orbits $a_i * U$ all have cardinality 27. The $(54, 53)$ - PA $C := \bigcup_{i=1}^{16} a_i * U$ is $(54, 8)$ - separable. This proves

Theorem 1. $N(54) \geq 8$.

Let $n = 96$. In [6] the existence of 8 MOLS has been proven by constructing a $(96, 8)$ - separable code. This implies $N(96) \geq 8$. Now we like to improve this bound using a different isometry group. Let $U := \langle x_1, x_2, x_3, x_4 \rangle \leq Iso(96)$ of order 2304 generated by x_1, x_2, x_3, x_4 given in cycle structure

$x_1 := (1, 4)(2, 3)(5, 16)(6, 15)(7, 13)(8, 14)(9, 11)(10, 12)(17, 20)(18, 19)(21, 32)(22, 31)$
 $(23, 29)(24, 30)(25, 27)(26, 28)(33, 36)(34, 35)(37, 48)(38, 47)(39, 45)(40, 46)(41, 43)$
 $(42, 44)(49, 52)(50, 51)(53, 64)(54, 63)(55, 61)(56, 62)(57, 59)(58, 60)(65, 68)(66, 67)$
 $(69, 80)(70, 79)(71, 77)(72, 78)(73, 75)(74, 76)(81, 84)(82, 83)(85, 96)(86, 95)(87, 93)$
 $(88, 94)(89, 91)(90, 92)(97, 99)(98, 100)(101, 111)(102, 112)(103, 110)(104, 109)(105, 108)$
 $(106, 107)(113, 115)(114, 116)(117, 127)(118, 128)(119, 126)(120, 125)(121, 124)(122, 123)$
 $(129, 131)(130, 132)(133, 143)(134, 144)(135, 142)(136, 141)(137, 140)(138, 139)(145, 147)$
 $(146, 148)(149, 159)(150, 160)(151, 158)(152, 157)(153, 156)(154, 155)(161, 163)(162, 164)$
 $(165, 175)(166, 176)(167, 174)(168, 173)(169, 172)(170, 171)(177, 179)(178, 180)(181, 191)$
 $(182, 192)(183, 190)(184, 189)(185, 188)(186, 187),$
 $x_2 := (1, 20, 33, 4, 17, 36)(2, 19, 34, 3, 18, 35)(5, 32, 37, 16, 21, 48)(6, 31, 38, 15, 22, 47)$
 $(7, 29, 39, 13, 23, 45)(8, 30, 40, 14, 24, 46)(9, 27, 41, 11, 25, 43)(10, 28, 42, 12, 26, 44)$
 $(49, 68, 81, 52, 65, 84)(50, 67, 82, 51, 66, 83)(53, 80, 85, 64, 69, 96)(54, 79, 86, 63, 70, 95)$
 $(55, 77, 87, 61, 71, 93)(56, 78, 88, 62, 72, 94)(57, 75, 89, 59, 73, 91)(58, 76, 90, 60, 74, 92)$
 $(97, 119, 129, 103, 113, 135)(98, 120, 130, 104, 114, 136)(99, 126, 131, 110, 115, 142)$
 $(100, 125, 132, 109, 116, 141)(101, 123, 133, 107, 117, 139)(102, 124, 134, 108, 118, 140)$
 $(105, 128, 137, 112, 121, 144)(106, 127, 138, 111, 122, 143)(145, 167, 177, 151, 161, 183)$
 $(146, 168, 178, 152, 162, 184)(147, 174, 179, 158, 163, 190)(148, 173, 180, 157, 164, 189)$
 $(149, 171, 181, 155, 165, 187)(150, 172, 182, 156, 166, 188)(153, 176, 185, 160, 169, 192)$
 $(154, 175, 186, 159, 170, 191),$
 $x_3 := (1, 96, 17, 64, 33, 80)(2, 87, 29, 63, 41, 75)(3, 86, 19, 54, 35, 70)(4, 94, 24, 53, 44, 74)$
 $(5, 92, 26, 52, 46, 72)(6, 83, 22, 51, 38, 67)(7, 93, 31, 57, 43, 66)(8, 85, 28, 58, 36, 78)$
 $(9, 91, 18, 55, 45, 79)(10, 84, 30, 56, 37, 76)(11, 82, 23, 61, 47, 73)(12, 90, 20, 62, 40, 69)$
 $(13, 95, 25, 59, 34, 71)(14, 88, 21, 60, 42, 68)(15, 89, 27, 50, 39, 77)(16, 81, 32, 49, 48, 65)$
 $(97, 164, 137, 151, 125, 192)(98, 168, 130, 152, 114, 184)(99, 162, 140, 158, 120, 182)$
 $(100, 173, 132, 157, 116, 189)(101, 171, 133, 155, 117, 187)(102, 175, 142, 156, 122, 179)$
 $(103, 169, 139, 145, 128, 181)(104, 166, 131, 146, 124, 190)(105, 167, 141, 160, 113, 180)$
 $(106, 163, 134, 159, 126, 188)(107, 161, 144, 149, 119, 185)(108, 174, 136, 150, 115, 178)$
 $(109, 176, 129, 148, 121, 183)(110, 172, 138, 147, 118, 191)(111, 170, 143, 154, 127, 186)$
 $(112, 165, 135, 153, 123, 177),$
 $x_4 := (1, 86, 17, 54, 33, 70)(2, 94, 29, 53, 41, 74)(3, 96, 19, 64, 35, 80)(4, 87, 24, 63, 44, 75)$
 $(5, 89, 26, 50, 46, 77)(6, 81, 22, 49, 38, 65)(7, 88, 31, 60, 43, 68)(8, 95, 28, 59, 36, 71)$
 $(9, 90, 18, 62, 45, 69)(10, 82, 30, 61, 37, 73)(11, 84, 23, 56, 47, 76)(12, 91, 20, 55, 40, 79)$
 $(13, 85, 25, 58, 34, 78)(14, 93, 21, 57, 42, 66)(15, 92, 27, 52, 39, 72)(16, 83, 32, 51, 48, 67)$
 $(97, 165, 137, 153, 125, 177)(98, 170, 130, 154, 114, 186)(99, 175, 140, 156, 120, 179)$
 $(100, 171, 132, 155, 116, 187)(101, 173, 133, 157, 117, 189)(102, 162, 142, 158, 122, 182)$
 $(103, 167, 139, 160, 128, 180)(104, 163, 131, 159, 124, 188)(105, 169, 141, 145, 113, 181)$
 $(106, 166, 134, 146, 126, 190)(107, 176, 144, 148, 119, 183)(108, 172, 136, 147, 115, 191)$
 $(109, 161, 129, 149, 121, 185)(110, 174, 138, 150, 118, 178)(111, 168, 143, 152, 127, 184)$
 $(112, 164, 135, 151, 123, 192).$

Furthermore let

$a_1 := (2, 9, 30, 43, 34, 77, 51, 76, 54, 70, 17, 3, 58, 15, 16, 7, 19, 87, 72, 39, 61, 49, 44, 42, 59, 55)$
 $(4, 50, 35, 31, 6, 63, 46, 47, 68, 32, 14, 83, 5, 56, 10, 22, 81, 88, 79, 85)$
 $(8, 28, 37, 18, 12, 94, 29, 36, 24, 71, 48, 75, 62, 57)(11, 86)$
 $(13, 92, 20, 96, 82, 95, 90, 74, 33, 69, 25, 73, 41, 52, 67, 23, 80, 93, 21, 89, 66, 91, 27, 45,$
 $40, 53, 78, 60, 64, 38, 26, 65, 84),$
 $a_2 := (2, 11, 8, 70, 61, 24, 15, 41, 95, 30, 85, 19, 67, 31, 92, 46, 80, 52, 49, 53, 84, 45, 4, 5, 48,$
 $14, 47, 74, 25, 38, 3, 7, 42, 66, 81, 21, 86, 63, 93, 64, 17, 78, 33, 65, 26, 77, 62, 83, 27, 68,$
 $87, 29, 10)(6, 75, 32, 16, 69, 34, 96, 58, 54, 23, 91, 28, 44, 90, 22, 9, 12)$
 $(13, 76, 88, 57, 56, 18, 37, 79, 40)(20, 43, 72, 51, 59, 60, 50, 55, 94)(35, 71, 39, 73, 82, 36, 89),$
 $a_3 := (3, 18, 55, 81, 41, 90, 11, 48, 92, 23, 62, 85, 30, 94, 53, 67, 26, 31, 22, 75, 40, 58, 49, 84, 63,$

52, 95, 9, 35, 72, 69, 46, 78, 28, 73, 7, 4, 17, 56, 82, 42, 89, 12, 47, 91, 24, 61, 86, 29, 93, 54,
68, 25, 32, 21, 76, 39, 57, 50, 83, 64, 51, 96, 10, 36, 71, 70, 45, 77, 27, 74, 8)(5, 37)(6, 38)
(13, 20, 87, 43, 16, 34, 60, 65, 79, 14, 19, 88, 44, 15, 33, 59, 66, 80),
 $a_4 := (1, 49, 11, 59)(2, 51, 14, 57, 5, 54, 13, 60, 3, 63, 9, 55, 6, 64, 12, 58, 15, 52, 7, 56, 16, 50,$
 $10, 62, 4, 53, 8, 61)(17, 65, 27, 75)(18, 67, 30, 73, 21, 70, 29, 76, 19, 79, 25, 71, 22, 80, 28,$
 $74, 31, 68, 23, 72, 32, 66, 26, 78, 20, 69, 24, 77)(33, 81, 43, 91)(34, 83, 46, 89, 37, 86, 45,$
 $92, 35, 95, 41, 87, 38, 96, 44, 90, 47, 84, 39, 88, 48, 82, 42, 94, 36, 85, 40, 93),$
 $a_5 := (2, 13, 14)(3, 12, 7, 11, 4, 15)(5, 9, 6)(8, 16)(18, 29, 30)(19, 28, 23, 27, 20, 31)(21, 25, 22)$
 $(24, 32)(34, 45, 46)(35, 44, 39, 43, 36, 47)(37, 41, 38)(40, 48)(49, 54, 50, 58, 61, 57)$
 $(51, 55, 56)(53, 62)(59, 63, 64)(65, 70, 66, 74, 77, 73)(67, 71, 72)(69, 78)(75, 79, 80)$
 $(81, 86, 82, 90, 93, 89)(83, 87, 88)(85, 94)(91, 95, 96),$
 $a_6 := (1, 11, 9, 3, 13, 12, 6, 8, 14, 16, 10, 7)(2, 15, 5, 4)(17, 27, 25, 19, 29, 28, 22, 24, 30,$
 $32, 26, 23)(18, 31, 21, 20)(33, 43, 41, 35, 45, 44, 38, 40, 46, 48, 42, 39)(34, 47, 37, 36)$
 $(49, 63, 58, 51, 50, 59, 54, 52, 61, 64, 53, 56)(55, 62, 60, 57)(65, 79, 74, 67, 66, 75, 70,$
 $68, 77, 80, 69, 72)(71, 78, 76, 73)(81, 95, 90, 83, 82, 91, 86, 84, 93, 96, 85, 88)(87, 94, 92, 89).$

The orbits $a_1 * U, a_2 * U, a_3 * U, a_4 * U, a_5 * U, a_6 * U$ have cardinality 288, 288, 288, 48, 24 and 24 respectively and the $(96, 95)$ - PA $C := a_1 * U \cup a_2 * U \cup a_3 * U \cup a_4 * U \cup a_5 * U \cup a_6 * U$ is $(96, 10)$ - separable. This proves

Theorem 2. $N(96) \geq 10$.

Now let $n = 108$. For the finite fields $\text{GF}(4)$ and $\text{GF}(27)$, let z and x respectively be primitive elements satisfying $z^2 = z + 1$ and $x^3 = x + 2$. Let F_1, F_2 and F_3 be the arrays below. Also let $V = ((0, v_1), (0, v_2), \dots, (0, v_{10}))^T$ where $(v_1, v_2, \dots, v_{10}) = (0, 1, x, x^2, x^3 = x + 2, x^4 = x^2 + 2x, x^5 = 2x^2 + x + 2, x^9 = x + 1, x^{25} = 2x^2 + 1, x^{20} = 2x^2 + x + 1)$. Likewise let $W = ((0, w_1), (0, w_2), \dots, (0, w_{10}))^T$ where $(w_1, w_2, \dots, w_{10}) = x(v_1, v_2, \dots, v_{10}) = (0, x, x^2, x^3, x^4 = x^2 + 2x, x^5 = 2x^2 + x + 2, x^6 = x^2 + x + 1, x^{10} = x^2 + x, x^{26} = 1, x^{21} = x^2 + 1)$.

The columns of our $(108, 10, 1)$ -DM are then obtained by adding the 9 vectors $aV + bW$ ($a, b \in \mathbb{Z}_3$) to each column of $F = [F_1|F_2|F_3]$.

$$F_1 = \begin{pmatrix} (0, 0) & (z + 1, 0) & (z, 0) & (z + 1, 0) \\ (0, 0) & (0, x^2) & (0, x^2) & (1, 2x^2) \\ (0, 0) & (0, 2x + 1) & (0, 2) & (z, 1) \\ (0, 0) & (1, x^2 + 2x) & (z + 1, x^2 + x + 1) & (0, 0) \\ (0, 0) & (1, x^2 + x + 2) & (z + 1, 1) & (z + 1, 2x^2 + x + 1) \\ (0, 0) & (1, 2x^2 + 2x + 2) & (z + 1, x^2 + 2) & (z, 2x^2 + 1) \\ (0, 0) & (z, 2x^2 + 1) & (1, x^2 + 1) & (1, x + 2) \\ (0, 0) & (z, 2x^2 + x + 2) & (1, 2x) & (z, x^2) \\ (0, 0) & (z + 1, x + 1) & (z, x^2 + 2x) & (1, 2x^2 + 2x) \\ (0, 0) & (z + 1, 2x^2 + x) & (z, 2x^2 + 2x + 1) & (0, x^2 + x + 2) \end{pmatrix},$$

$$F_2 = \begin{pmatrix} (1, 0) & (0, 0) & (z + 1, 0) & (z, 0) \\ (1, 2x^2) & (1, 0) & (z, 2x^2) & (z, x^2) \\ (z, 2) & (z, 1) & (z + 1, 2x + 1) & (z + 1, 0) \\ (z, 2x^2 + 2) & (z + 1, 1) & (z, x) & (0, x) \\ (0, 2x + 1) & (z, x^2 + 2x + 2) & (1, x^2 + x + 1) & (z, 2) \\ (z + 1, 2x^2 + x + 1) & (0, x^2 + x + 1) & (0, 2) & (1, 2x^2 + 1) \\ (z + 1, x^2 + 2x + 1) & (z, x + 1) & (0, x^2 + x) & (z + 1, 2x) \\ (1, 2) & (z + 1, 2x^2 + x + 1) & (1, 2) & (0, x^2) \\ (0, x + 2) & (z + 1, x) & (z, 2x^2 + 2x + 2) & (1, x^2) \\ (z + 1, 0) & (1, 2x^2 + 1) & (1, 2x + 2) & (z + 1, x^2 + 2x) \end{pmatrix},$$

$$F_3 = \begin{pmatrix} (1, 0) & (z, 0) & (1, 0) & (0, 0) \\ (z, 0) & (z+1, x^2) & (z+1, 0) & (z+1, 2x^2) \\ (z+1, 0) & (1, x) & (1, 2) & (1, x+2) \\ (1, 2x) & (z, x) & (z+1, 2x^2+2x+1) & (1, 2x+1) \\ (0, 2x^2+2x+1) & (1, 2x^2+2x) & (z, x^2+1) & (z+1, 2x^2+x) \\ (z, 2x^2+2x+2) & (z+1, 0) & (1, x^2+1) & (z, 2x^2+2) \\ (1, 1) & (z, 2x+2) & (0, 2x+1) & (z+1, x^2+x+2) \\ (z+1, 2x^2+1) & (z+1, x^2+1) & (z, 2x^2+x+2) & (0, x^2+x+1) \\ (z+1, x^2+1) & (1, 2x+1) & (0, 2x^2+1) & (z, x+1) \\ (z, x+2) & (0, 2x^2+1) & (z, x) & (1, 2x^2+2x) \end{pmatrix}.$$

This proves

Theorem 3. $N(108) \geq 9$.

Finally we come to the correction of an error in [1] for the case $n=45$. Here we give the corrected $(45, 7, 1)$ difference matrix. Consider the arrays

$$E_1 = \begin{pmatrix} (0, 0, 0) \\ (0, 0, 0) \\ (0, 0, 0) \\ (0, 0, 0) \\ (0, 0, 0) \\ (0, 0, 0) \\ (0, 0, 0) \end{pmatrix}, \quad E_2 = \begin{pmatrix} (2, 2, 1) & (3, 1, 1) & (4, 1, 2) & (4, 0, 1) & (0, 1, 1) & (0, 2, 1) & (3, 2, 2) \\ (1, 2, 1) & (4, 2, 2) & (1, 2, 0) & (4, 1, 0) & (3, 1, 1) & (3, 0, 0) & (2, 1, 2) \\ (4, 1, 1) & (2, 2, 1) & (3, 2, 0) & (1, 2, 0) & (2, 1, 0) & (1, 0, 0) & (3, 2, 1) \\ (0, 1, 0) & (2, 1, 1) & (4, 0, 0) & (0, 0, 2) & (4, 2, 2) & (3, 2, 2) & (1, 2, 2) \\ (3, 1, 2) & (2, 1, 0) & (0, 2, 2) & (4, 2, 1) & (0, 2, 1) & (2, 0, 1) & (1, 1, 2) \\ (2, 1, 1) & (1, 2, 2) & (3, 0, 1) & (2, 0, 1) & (1, 0, 0) & (4, 2, 1) & (1, 1, 0) \\ (0, 0, 0) & (0, 0, 0) & (0, 0, 0) & (0, 0, 0) & (0, 0, 0) & (0, 0, 0) & (0, 0, 0) \end{pmatrix}.$$

Now let E_3 be the array over $Z_5 \times Z_3 \times Z_3$ obtained by interchanging the following pairs of rows in E_2 : 1 and 6, 2 and 5, 3 and 4, while leaving row 7 unaltered. Also, let $V = [(0, 0, 1), (0, 2, 0), (0, 1, 2), (0, 2, 1), (0, 1, 0), (0, 0, 2), (0, 0, 0)]^T$. The columns of our $(45, 7, 1)$ -DM are then obtained by adding the 3 vectors aV ($a \in Z_3$) to each column of $E = [E_1|E_2|E_3]$.

Acknowledgements

The authors would like to thank Alice Miller for several useful suggestions and comments, and for checking a number of results within this paper.

ORCID

R. J. R. Abel: <https://orcid.org/0000-0002-3632-9612>

I. Janiszczak: <https://orcid.org/0000-0002-4892-1189>

References

- [1] R.J.R. Abel, C.J. Colbourn and J.H. Dinitz, Mutually orthogonal Latin squares (MOLS), in: The CRC Handbook of Combinatorial Designs, Second Edition (C. J. Colbourn and J. H. Dinitz, eds.), CRC Press, Boca Raton FL, 2007, 160–193.
- [2] Bosma W., Cannon J., Playoust C.: The Magma algebra system. I. The user language. J. Symbolic Computation **24**, 235-265 (1997). URL: <http://magma.maths.usyd.edu.au/magma/>.

- [3] Colbourn C.J., Kløve T., Ling A.C.H.: Permutation arrays for powerline communication and mutually orthogonal Latin squares. *IEEE Trans. Inform. Theory* **50**, 1289-1291 (2004).
- [4] Farahat H.: The symmetric group as a metric space. *J. London Math. Soc.* **35**, 215-220 (1960).
- [5] Janiszczak, I., Lempken, W., Östergård, P.R.J. et al. Permutation codes invariant under isometries. *Des. Codes Cryptogr.* **75**, 497–507 (2015).
- [6] Janiszczak, I., Staszewski, R.: Isometry invariant permutation codes and mutually orthogonal Latin squares. *J. Combin. Designs.* **27**, 541–551 (2019).
- [7] The GAP Group. GAP – Groups, Algorithms, and Programming, Version 4.13.1, 2024. URL: <http://www.gap-system.org>.

Appendix: GAP code

```
#####
# HERE YOU NEED TO DEFINE n.
n:= ;
#####

a1:=[1..n];
b1:=(1,2);
a2:=[n+1..2*n];
b2:=(n+1,n+2);
t:=Identity(SymmetricGroup([1..2*n]));
for i in [1..n] do
  t:=t*(i,n+i);
od;

# THE GROUPS A,B,B1 AND B2 WILL BE DEFINED
A:=Group(CycleFromList(a1),CycleFromList(a2),b1,b2,t);
B:=Subgroup(A,[CycleFromList(a1),b1,CycleFromList(a2),b2]);
B1:=Subgroup(A,[CycleFromList(a1),b1]);
B2:=Subgroup(A,[CycleFromList(a2),b2]);

#####
# HERE YOU HAVE TO ENTER THE GENERATORS FOR THE ISOMETRY GROUP
# SEPARATED BY ,
UGenerators=[];
#####

#####
# HERE ARE THE ORBITREPRESENTATIVES SEPARATED BY ,
OrbReps=[];
#####

U:=Subgroup(A,UGenerators);

action:=function(x,y)
#THE ACTION OF ELEMENTS OF A ON ELEMENTS OF B1
  local x1,x2,xony;
  if x in B then
    x1:=RestrictedPerm(x,a1);
    x2:=RestrictedPerm(x,a2);
    xony:=x1^-1*y*x2^t;
  else
    x1:=RestrictedPerm(x*t,a1);
    x2:=RestrictedPerm(x*t,a2);
    xony:=(x1^-1*y*x2^t)^-1;
  fi;
  return xony;
end;

grouporbit:=function(G,y)
# RETURNS THE ORBIT GIVEN BY THE ACTION OF G ON y
  local x,orb;
  orb=[];

```



```

    for x in G do
        AddSet(orb,action(x,y));
    od;
    return orb;
end;

dist:=function(x,y)
# RETURNS THE DISTANCE OF x AND y FOR x,y IN B1
    return(Number(MovedPoints(x^-1 * y)));
end;

mindistset:=function(X)
# RETURNS THE MINIMUMDISTANCE OF THE SUBSET X OF B1
    local i,j,d,mindist;
    mindist:=n;
    for i in [1..Number(X)] do
        for j in [i+1..Number(X)] do
            d:=dist(X[i],X[j]);
            if d < mindist then
                mindist:=d;
            fi;
        od;
    od;
    return(mindist);
end;

isdistancebetweensetseq:=function(distance,X,Y)
# RETURNS true IF FOR ALL x IN X AND ALL y IN Y dist(x,y) EQUALS distance
# AND false OTHERWISE
    local i,j;
    for i in [1..Number(X)] do
        for j in [1..Number(Y)] do
            if dist(X[i],Y[j]) = distance then
                continue;
            else
                return(false);
            fi;
        od;
    od;
    return(true);
end;

mkorbitbits:=function(OrbitReps)
# INPUT IS THE SET OF ORBITREPRESENTATIVES
# RETURNS THE CORRESPONDING SET OF ORBITS
    local i, orbs;
    orbs=[];
    for i in [1..Number(OrbitReps)] do
        orbs[i]:=grouporbit(U,OrbitReps[i]);
    od;
    return(orbs);
end;

mkcode:=function(Orbits)
# INPUT IS A SET OF ORBITS, RETURNS THE UNION OF ALL ORBITS

```

```

local i, Code;
Code:=[];
for i in [1..Number(Orbits)] do
    UniteSet(Code,Orbits[i]);
od;
return(Code);
end;

nseparable:=function(Code)
# RETURNS THE Code SEPARATED IN CODES OF LENGTH n WITH MINIMUMDISTANCE n
# IF THE Code IS (n,Number(Code)/n)-SEPARABLE AND [] OTHERWISE
local i,j,r, latinsquare, latinsquares, elementshandled,ls1,ls2;
if Number(Code) mod n > 0 then
    return([]);
fi;
r:=Number(Code)/n;
latinsquares:=[];
elementshandled:=[];
for j in [1..Number(Code)] do
    elementshandled[j]:=0;
od;
for i in [1..r] do
    latinsquare:=[];
    for j in [1..Number(Code)] do
        if elementshandled[j] = 1 then
            continue;
        fi;
        if Number(latinsquare) = 0 then
            AddSet(latinsquare,Code[j]);
            elementshandled[j] := 1;
        else
            if dist(latinsquare[1],Code[j]) = n then
                AddSet(latinsquare,Code[j]);
                elementshandled[j] := 1;
            fi;
        fi;
    od;
    if Number(latinsquare) = n then
        if mindistset(latinsquare) = n then
            AddSet(latinsquares,latinsquare);
            continue;
        else
            return([]);
        fi;
    fi;
od;
for i in [1..Number(latinsquares)-1] do
    ls1:=latinsquares[i];
    for j in [i+1..Number(latinsquares)] do
        ls2:=latinsquares[j];
        if isdistancebetweensetseq(n-1,ls1,ls2) then
            continue;
        else
            return([]);
        fi;
    od;
od;

```

```

        od;
    od;
    return(latinsquares);
end;

makeMOLS:=function(sepcode)
# GIVEN A (n,m)-SEPARABLE CODE sepcode WHICH IS SEPARATED IN m (n,n) PAs
# THE SET OF m MOLS WILL BE RETURNED
    local i,j,k,sepdeg,MOLS;
    MOLS=[];
    sepdeg:=Number(sepcode);
    for i in [1..sepdeg] do
        MOLS[i]:=[];
        for j in [1..n] do
            MOLS[i][j]:=[];
        od;
        for j in [1..n] do
            for k in [1..n] do
                MOLS[i][j^sepcode[i][k]][j]:=k;
            od;
        od;
    od;
    return(MOLS);
end;

isls:=function(ls)
# RETURNS true IF ls IS A LATINSQUARE AND false OTHERWISE
    local i,j;
    for i in [1..n] do
        if Number(Set(ls[i])) = n then
            continue;
        else
            return(false);
        fi;
    od;
    return(true);
end;

lsorth:=function(ls1,ls2)
# RETURNS true IF ls1 AND ls2 ARE ORTHOGONAL AND false OTHERWISE
    local i,j,pairset;
    pairset=[];
    for i in [1..n] do
        for j in [1..n] do
            AddSet(pairset,[ls1[i][j],ls2[i][j]]);
        od;
    od;
    if Number(pairset) = n^2 then
        return(true);
    else
        return(false);
    fi;
end;

orbs:=mkorbits(OrbReps);

```

```
# orbs CONTAINS ALL ORBITS FOR THE REPRESENTATIVES IN OrbReps

MOLSCode:=mkcode(orbs);
# MOLSCode IS THE UNION OF ALL ORBITS

SepCode:=nseparable(MOLSCode);
Print(Number(SepCode),"\n");
# PRINTS Number(MOLSCode)/n IF MOLSCode IS SEPARABLE AND 0 ELSE

if Number(SepCode) = Number(MOLSCode)/n then
  Mols:=makeMOLS(SepCode);
fi;
# Mols ARE THE RESULTING MOLS FROM MOLSCODE
```