

Counting humps and peaks in Motzkin paths with height k

Xiaomei Chen

School of Mathematics and Statistics
Hunan University of Science and Technology
Xiangtan 411201, China
xmchen@hnust.edu.cn

Mathematics Subject Classifications: 05A15

Abstract

In this paper, we construct several bijections among humps of Motzkin paths, standard Young tableaux (SYTs) in a $(2, 1)$ -hook and two special kinds of Motzkin prefixes. As corollaries, we obtain the number of humps and peaks in Motzkin paths with height k , and the number of SYTs with shapes in $\{(\lambda_1, \lambda_2, 1, \dots) \mid \lambda_1 - \lambda_2 = k\}$, which refine Regev's results in 2009. We also give new combinatorial proofs of Regev's results, and reveal some new recurrence relations related to humps, free Motzkin paths and SYTs.

Keywords: Motzkin paths, humps, peaks, standard Young tableaux, Motzkin prefixes, Riordan arrays

Mathematics Subject Classifications: 05A15

1 Introduction

A *Motzkin path* of order n is a lattice path from $(0, 0)$ to $(n, 0)$, using up steps $(1, 1)$ (denoted by U), down steps $(1, -1)$ (denoted by D) and flat steps $(1, 0)$ (denoted by F), and never going below the x -axis. If such a path is allowed to go below the x -axis, it is called a *free Motzkin path*. Let \mathcal{M}_n denote the set

of all Motzkin paths of order n . The cardinality of \mathcal{M}_n is the n th Motzkin number m_n ([11], A001006).

A *hump* in a Motzkin path is an up step followed by zero or more flat steps followed by a down step. Especially, a hump with no flat steps is called a *peak*. See Figure 1 for an example.




Figure 1: An example of hump and peak, where the hump P_1 is colored red, and the peak P_2 is colored blue.

Let HM_n denote the total number of humps in all Motzkin paths of order n , and let SM_n denote the number of free Motzkin paths of order n . By applying the WZ method [8, 14], Regev [9] proved that

$$HM_n = \frac{1}{2}(SM_n - 1) = \frac{1}{2} \left(\sum_{j \geq 1} \binom{n}{j} \binom{n-j}{j} \right). \quad (1.1)$$

A combinatorial proof of (1.1) was given by Ding and Du [1], and generalizations of this study to (k, a) -paths were given in [2, 5, 12].

Regev pointed out that (1.1) is also related to the number of standard Young tableaux (SYT) in a $(2, 1)$ -hook, where a (k, l) -hook is a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ with $\lambda_{k+1} \leq l$. Let $S(k, l; n)$ denote the number of SYTs of order n in a (k, l) -hook. It is obvious that $S(1, 1; n) = 2^{n-1}$. For the “hook” sums of general shapes, only the following result of Regev [9, 10] is known:

$$S(2, 1; n) - 1 = HM_n = \frac{1}{2} \left(\sum_{j \geq 1} \binom{n}{j} \binom{n-j}{j} \right). \quad (1.2)$$

Du and Yu [3] gave a combinatorial proof of (1.2) by proving bijectively that

$$S(2, 1; n) = \frac{1}{2}(SM_n + 1).$$

The goal of this paper is to refine identities (1.1) and (1.2), and reveal some further relations between humps of Motzkin paths and Young tableaux.

The statistic *height* for a hump in a Motzkin path is defined to be the y -coordinate reached by the up step of the hump. Note that the number of peaks in Dyck paths with a given height has been studied in several papers, such as [6] and [7]. We denote

$$\mathcal{HM}_{n,k} = \{(M, P) \mid M \in \mathcal{M}_n, P \text{ is a hump of } M \text{ with height } k\},$$

$$\mathcal{PM}_{n,k} = \{(M, P) \mid M \in \mathcal{M}_n, P \text{ is a peak of } M \text{ with height } k\},$$

and denote $HM_{n,k} = \#\mathcal{HM}_{n,k}$ and $PM_{n,k} = \#\mathcal{PM}_{n,k}$.

To count humps with a given height, the relation between humps and Motzkin prefixes is considered. Here a *Motzkin prefix* is a lattice path that is a prefix of a Motzkin path. We denote $\mathcal{MP}_{n,k}$ the set of Motzkin prefixes from $(0, 0)$ to (n, k) , and denote

$$\mathcal{MP}_{n,k}^{*U} = \{M \mid M \in \mathcal{MP}_{n,k} \text{ whose last non-flat step is } U\},$$

$$\mathcal{MP}_{n,k}^{D*} = \{M \mid M \in \mathcal{MP}_{n,k} \text{ whose first non-up step is } D\}.$$

See [4] and [13] for some further results about Motzkin prefix numbers.

Our first result gives a bijection from $\mathcal{HM}_{n,k}$ to $\mathcal{MP}_{n,2k}^{*U}$. As a corollary, we obtain the following generalization of Equation (1.1):

$$HM_{n,k} = SM_{n,k}, \tag{1.3}$$

where $SM_{n,k}$ denotes the number of free Motzkin paths of order n whose smallest y -coordinate is $-k$ and last non-flat step is U . We also reveal some new recurrence relations between $HM_{n,k}$ and Motzkin prefix numbers. Let $MP_{n,k}$ denote the cardinality of $\mathcal{MP}_{n,k}$ (A064189 in OEIS [11]). Then we show that

$$HM_{n,k} = \sum_{m=2k-1}^{n-1} MP_{m,2k-1}, \tag{1.4}$$

$$MP_{n,2k} = HM_{n,k} + HM_{n,k+1}. \tag{1.5}$$

Moreover, we obtain the following formulas for $HM_{n,k}$ and $PM_{n,k}$.

Theorem 1.1. *Let $HM_{n,k}$ (resp. $PM_{n,k}$) be the total number of humps (resp. peaks) with height k in all Motzkin paths of order n . Then for $n \geq 2$ and*

$k \geq 1$, we have

$$HM_{n,k} = \sum_{j=0}^{n-2k} \chi(j \equiv n \pmod{2}) \frac{4k}{n-j+2k} \binom{n}{j} \binom{n-j-1}{(n-j)/2+k-1},$$

$$PM_{n,k} = \sum_{j=0}^{n-2k} \chi(j \equiv n \pmod{2}) \frac{4k}{n-j+2k} \binom{n-1}{j} \binom{n-j-1}{(n-j)/2+k-1},$$

where χ is the characteristic function. Moreover, the number j in the sum tracks up to the number of flat steps of the Motzkin paths.

We also consider the relation between SYTs contained in a $(2, 1)$ -hook and Motzkin prefixes. As a consequence, we are able to count SYTs in a $(2, 1)$ -hook according to the difference of the first two rows of the hook. Let

$$\mathcal{S}_k(2, 1; n) = \bigcup_{\lambda} \text{SYT}(\lambda),$$

where $\lambda = (\lambda_1, \lambda_2, 1, 1, \dots)$ ranges over all $(2, 1)$ -hooks with $|\lambda| = n$ and $\lambda_1 - \lambda_2 = k$, and $\text{SYT}(\lambda)$ denotes the set of SYTs with shape λ . Let $S_k(2, 1; n)$ denote the cardinality of $\mathcal{S}_k(2, 1; n)$. We firstly show that

$$S_{2k-1}(2, 1; n+1) + S_{2k-1}(2, 1; n) = HM_{n+1,k} - HM_{n,k}. \quad (1.6)$$

Then by constructing several bijections, we obtain a new combinatorial proof of (1.2), and obtain the following result for $S_k(2, 1; n)$.

Theorem 1.2. *Let $S_k(2, 1; n)$ be the number of standard Young tableaux with shapes in $\{(\lambda_1, \lambda_2, 1^{n-\lambda_1-\lambda_2}) \mid \lambda_1 - \lambda_2 = k\}$. For integers n, k with $n - 2 \geq k \geq 0$, we have*

$$S_k(2, 1; n) = (-1)^{n+k} + \sum_{i=0}^{\lfloor \frac{n-k-1}{2} \rfloor} \sum_{j=0}^{n-k-1-2i} \chi(j \equiv n+k-1 \pmod{2}) \cdot \frac{2k+2}{n+k+1-2i-j} \binom{n-2i-2}{j} \binom{n-2i-j-1}{(n+k-j-1)/2-i}.$$

2 The number of humps and peaks with height k

A Motzkin prefix M of order n can be represented as a word $M = w_1w_2 \cdots w_n$ with $w_i \in \{U, D, F\}$. We use \overline{M} to denote the path obtained from M by reading the steps in reverse order and then swapping the U 's and D 's.

Lemma 2.1. *There is a bijection*

$$\psi : \bigcup_{1 \leq k \leq \lfloor \frac{n}{2} \rfloor} \mathcal{HM}_{n,k} \mapsto \bigcup_{1 \leq k \leq \lfloor \frac{n}{2} \rfloor} \mathcal{MP}_{n,2k}^{*U},$$

such that $(M, P) \in \mathcal{HM}_{n,k}$ if and only if $\psi(M, P) \in \mathcal{MP}_{n,2k}^{*U}$. Especially, P is a peak of M , if and only if $\psi(M, P)$ ends with an up step.

Proof. Given $(M, P) \in \mathcal{HM}_{n,k}$, assume that $P = UF^rD$. We decompose M as $M = M_1PM_2$, and define

$$\psi(M, P) = M_1U\overline{M_2}UF^r.$$

It is not difficult to see that $\psi(M, P) \in \mathcal{MP}_{n,2k}^{*U}$. Especially, if $M = M_1PM_2$ with $P = UD$ a peak, then $\psi(M, P) = M_1U\overline{M_2}U$ ends with an up step. See Figure 2 for an example of ψ .




Figure 2: An example of ψ with $(M, P) \in \mathcal{HM}_{9,2}$, where the hump P is colored red.

Conversely, given $M \in \mathcal{MP}_{n,2k}^{*U}$, let $M = M_1UM_2UF^r$ be the decomposition of M , such that the rightmost endpoint of M_1 is the last endpoint of M with y -coordinate $k - 1$. Then we have $\psi^{-1}(M) = (M_1UF^rD\overline{M_2}, P)$, where P is the hump connecting M_1 and $\overline{M_2}$. \square

Similar to the proof of Lemma 2.1, we can give a combinatorial proof of Equation (1.3).

Proof of Equation (1.3). Given $(M, P) \in \mathcal{HM}_{n,k}$, assume that $P = UF^rD$. We decompose M as $M = M_1PM_2$, and define

$$\psi^*(M, P) = M_2DM_1UF^r.$$

It is not difficult to check that ψ^* is a bijection from $\mathcal{HM}_{n,k}$ to the set of free Motzkin paths of order n whose smallest y -coordinate is $-k$ and last non-flat step is U , which implies Equation (1.3). \square

By Lemma 2.1, we can obtain proofs of Equation (1.4) and Equation (1.5).

Corollary 2.2. *For $n \geq 2$ and $k \geq 1$, we have*

$$\begin{aligned} HM_{n,k} &= \sum_{m=2k-1}^{n-1} MP_{m,2k-1}, \\ MP_{n,2k} &= HM_{n,k} + HM_{n,k+1}. \end{aligned}$$

Proof. Given $P \in \mathcal{MP}_{n,2k}^{*U}$, assume that $P = M_1UH^r$, then we define $\rho_1(P) = M_1$. It is not difficult to check that

$$\rho_1 : \mathcal{MP}_{n,2k}^{*U} \longrightarrow \bigcup_{m=2k-1}^{n-1} \mathcal{MP}_{m,2k-1}$$

is a bijection. Thus the first identity can be deduced from Lemma 2.1.

On the other hand, given $P \in \mathcal{MP}_{n,2k}$, we define $\rho_2(P)$ as follows.

- If $P \in \mathcal{MP}_{n,2k}^{*U}$, we define $\rho_2(P) = P$.
- If the last non-flat step of P is D , we define $\rho_2(P)$ to be the path obtained from P by replacing the last D step with U .

Then ρ_2 is a bijection from $\mathcal{MP}_{n,2k}$ to the union of $\mathcal{MP}_{n,2k}^{*U}$ and $\mathcal{MP}_{n,2k+2}^{*U}$, and the second identity follows. \square

A *Dyck prefix* is a Motzkin prefix with no flat steps. It is well known that the number of Dyck paths of semilength n ending with UD^m is counted by $\frac{m}{n} \binom{2n-m-1}{n-1}$, which is also equal to the number of Dyck prefixes from $(0, 0)$ to $(2n - m, m)$ ending with U . (See A33184 in OEIS [11]). Using the above fact and Lemma 2.1, we can give the proof of Theorem 1.1.

Proof of Theorem 1.1. A path $M \in \mathcal{MP}_{n,2k}^{*U}$ with j flat steps can be obtained uniquely by inserting j flat steps into $n-j+1$ positions in a Dyck prefix from $(0,0)$ to $(n-j, 2k)$ ending with U . Especially, M ends with an up step if and only if the above insertion is not allowed at the end of the path. Combining the above fact and Lemma 2.1, we then obtain the required identities in Theorem 1.1. \square

The first few entries of $(HM_{n,k})_{n \geq 2, k \geq 1}$ are as follows:

[2]	1;			
[3]	3;			
[4]	8,	1;		
[5]	20,	5;		
[6]	50,	19,	1;	
[7]	126,	63,	7;	
[8]	322,	196,	34,	1;
[9]	834,	588,	138,	9;
[10]	2187,	1728,	507,	53, 1.

Remark 2.3. $(HM_{n,1})$ is the sequence A140662 in OEIS [11]. In the case of $k \geq 2$, $(HM_{n,k})$ does not appear in [11].

We can also consider the generating function of $\{HM_{n,k}\}$. For fixed k , it is well known that the generating function of $\{MP_{n,k}\}$ is

$$\sum_{n \geq k} MP_{n,k} x^n = x^k M^{k+1}(x), \quad (2.1)$$

where $M(x)$ is the generating function of the Motzkin numbers. See A064189 in OEIS [11] for instance. Combining Equation (2.1) and Corollary 2.2, we obtain the following result.

Corollary 2.4. *The infinite lower triangular array $\{HM_{n,k}\}_{n,k \in \mathbb{N}}$ is the Riordan array $(\frac{1}{1-x}, x^2 M^2(x))$. Equivalently, for $k \geq 1$, we have*

$$\sum_{n \geq 2k} HM_{n,k} x^n = \frac{x^{2k} M^{2k}(x)}{1-x},$$

where $M(x)$ is the generating function of the Motzkin numbers.

3 The number of SYTs in $\mathcal{S}_k(2, 1; n)$

In this section, our first goal is to give a new combinatorial proof of (1.2). Note that there is a simple bijection

$$\phi : \bigcup_{0 \leq k \leq n-2} \mathcal{S}_k(2, 1; n) \mapsto \bigcup_{0 \leq k \leq n-2} \mathcal{MP}_{n,k}^{D*}$$

defined as follows: given $T \in \bigcup_{0 \leq k \leq n-2} \mathcal{S}_k(2, 1; n)$, we define $\phi(T) = w_1 w_2 \dots w_n$, where

$$w_i = \begin{cases} U, & \text{if } i \text{ appears in the first row of } T; \\ D, & \text{if } i \text{ appears in the second row of } T; \\ F, & \text{otherwise.} \end{cases}$$

Therefore, to prove Equation (1.2), it is sufficient to prove the following result.

Lemma 3.1. *There is a bijection*

$$\varphi : \bigcup_{0 \leq k \leq n-2} \mathcal{MP}_{n,k}^{D*} \mapsto \bigcup_{1 \leq k \leq \lfloor \frac{n}{2} \rfloor} \mathcal{MP}_{n,2k}^{*U}$$

Proof. Given $0 \leq k \leq n-2$ and $M \in \mathcal{MP}_{n,k}^{D*}$, we define $\varphi(M)$ as follows.

- If k is odd, we assume that $M = U^r D M_1$, and define $\varphi(M) = U^{r-1} F M_1 U$.
- If k is even, we define $\varphi(M)$ as follows.
 - * If the last non-flat step of M is U , then $\varphi(M) = M$.
 - * If $M = U^{r_1} D M_1 D F^{r_2}$, then $\varphi(M) = U^{r_1-1} F M_1 U F^{r_2+1}$.
 - * If $M = U^{r_1} D F^{r_2}$, then $\varphi(M) = U^{r_1+1} F^{r_2}$.

Conversely, given $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$ and $M \in \mathcal{MP}_{n,2k}^{*U}$, we obtain $\varphi^{-1}(M)$ in the following way.

- If the first non-up step of M is D , then $\varphi^{-1}(M) = M$.
- If the first non-up step of M is F or $M = U^n$, we obtain $\varphi^{-1}(M)$ as follows.

* If $M = U^r FM_1 U$, then $\varphi^{-1}(M) = U^{r+1} DM_1$.

* If $M = U^{r_1} FM_1 U F^{r_2}$ with $r_2 \geq 1$, then

$$\varphi^{-1}(M) = U^{r_1+1} DM_1 D F^{r_2-1}.$$

* If $M = U^{2k} F^{n-2k}$, then $\varphi^{-1}(M) = U^{2k-1} D F^{n-2k}$.

□

Combining the three bijections ψ , φ^{-1} and ϕ^{-1} together, we then obtain the following result, which implies Equation (1.2) immediately. See Figure 3 for an example.

Corollary 3.2. *There is a bijection*

$$\Phi : \bigcup_{1 \leq k \leq \lfloor \frac{n}{2} \rfloor} \mathcal{HM}_{n,k} \mapsto \bigcup_{0 \leq k \leq n-2} \mathcal{S}_k(2, 1; n).$$




Figure 3: An example of Φ with $(M, P) \in \mathcal{HM}_{9,2}$, where the hump P is colored red.

We now calculate the number of SYTs in $\mathcal{S}_k(2, 1; n)$. Let

$$S_{n,k} = S_k(2, 1; n) + (-1)^{n+k+1},$$

where we set $S_{0,0} = 0$. Then we have the following result for the generating function of $\{S_{n,k}\}$ for fixed k .

Lemma 3.3. *The infinite lower triangular array $\{S_{n,k}\}_{n,k \in \mathbb{N}}$ is the Riordan array $(\frac{xM(x)}{1+x}, xM(x))$. Equivalently, we have*

$$S_k(x) = \sum_{n \geq k} S_{n,k} x^n = \frac{1}{1+x} (xM(x))^{k+1},$$

where $M(x)$ is the generating function of the Motzkin numbers.

Proof. For $n \geq k$, we firstly show the following recurrence relation:

$$S_{n+1,k} + S_{n,k} = MP_{n,k}, \quad (3.1)$$

where $MP_{n,k}$ is the cardinality of $\mathcal{MP}_{n,k}$.

The left-hand side counts the number of paths in $\mathcal{MP}_{n+1,k}^{D*} \cup \mathcal{MP}_{n,k}^{D*}$, and the right-hand side counts the number of Motzkin prefixes in $\mathcal{MP}_{n,k}$. It is obvious that $S_{k,k} = 0$ and $S(k+1, k) = 1$ for $k \geq 0$, which implies (3.1) for $n = k$. For $n > k$ and $M \in \mathcal{MP}_{n,k}$, we define $f(M)$ as follows.

- If $M \in \mathcal{MP}_{n,k}^{D*}$, we define $f(M) = M$.
- If $M = U^r FM_1$, then $f(M) = U^{r+1} DM_1$.

It is not difficult to check that f is a bijection between $\mathcal{MP}_{n,k}$ and $\mathcal{MP}_{n+1,k}^{D*} \cup \mathcal{MP}_{n,k}^{D*}$. Therefore, we obtain the recurrence relation (3.1).

Combining Equation (2.1) and Equation (3.1), we obtain Lemma 3.3. \square

Remark 3.4. *For $k = 0$ and $k = 1$, $(S_{n,k})$ are the sequences A187306 and A284778 respectively in OEIS [11].*

The proof of Equation (3.1) also implies a combinatorial proof of the following recurrence relation.

Corollary 3.5. *For $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$,*

$$S_{2k-1}(2, 1; n+1) + S_{2k-1}(2, 1; n) = HM_{n+1,k} - HM_{n,k}.$$

Proof. By the proof of (3.1), the left-hand side counts the number of paths in $\mathcal{MP}_{n,2k-1}$. By Lemma 2.1, the right-hand side counts the number of paths in $\mathcal{MP}_{n+1,2k}^{*U}$ ending with U , which is also equal to the number of paths in $\mathcal{MP}_{n,2k-1}$. In fact, by deleting the last up step of a path in $\mathcal{MP}_{n+1,2k}^{*U}$ ending with U , we obtain a path in $\mathcal{MP}_{n,2k-1}$ immediately. \square

To obtain $S_k(2, 1; n)$, we define the weight w of a Motzkin prefix $M \in \mathcal{MP}_{n,k}^{*U}$ as $w(M) = (-1)^r$, where r is the number of flat steps after the last up step.

Proof of Theorem 1.2. By Lemma 3.3, we have

$$S(n, k) = \sum_{M \in \mathcal{MP}_{n,k+1}^{*U}} w(M).$$

Given $M \in \mathcal{MP}_{n,k+1}^{*U}$ with weight -1 , we define $f(M)$ to be the path obtained from M by moving the last flat step to the beginning. Then f is a bijection from the set of paths in $\mathcal{MP}_{n,k+1}^{*U}$ with weight -1 to those with weight 1 beginning with a flat step.

As a consequence of the above bijection, $S_{n,k}$ equals to the number of paths in $\mathcal{MP}_{n,k+1}^{*U}$ beginning with U and ending with an even number of flat steps. Since those paths can be obtained from the corresponding Dyck prefixes by inserting $2i$ flat steps to the end and inserting j flat steps between the first up step and the last up step, we obtain the required identity for $S_k(2, 1; n)$. \square

The first few entries of $(S_k(2, 1; n))_{n,k \in \mathbb{N}}$ are as follows:

[0]	1;								
[1]	0,	1;							
[2]	1,	0,	1;						
[3]	1,	2,	0,	1;					
[4]	3,	3,	3,	0,	1;				
[5]	6,	9,	6,	4,	0,	1;			
[6]	15,	21,	19,	10,	5,	0,	1;		
[7]	36,	55,	50,	34,	15,	6,	0,	1;	
[8]	91,	141,	139,	99,	55,	21,	7,	0,	1.

Remark 3.6. $(S_0(2, 1; n))$ is the sequence A005043 in OEIS [11], known as Riordan numbers.

Acknowledgements

The author was supported by the Natural Science Foundation of Hunan Province (2021JJ40186). The author is grateful to an anonymous for providing helpful advice to improve some results of the paper.

References

- [1] Y. Ding, R.R.X. Du, Counting humps in Motzkin paths, *Discrete Appl. Math.*, 160 (2012), 187-191.
- [2] R.R.X. Du, Y. Nie, X. Sun, Enumerations of humps and peaks in (k, a) -paths and (n, m) -Dyck paths via bijective proofs, *Discrete Appl. Math.*, 190-191 (2015), 42-49.
- [3] R.R.X. Du, J. Yu, Standard Young tableaux in a $(2, 1)$ -hook and Motzkin paths, *Discrete Math.*, 344 (7) (2021), 112395.
- [4] C. Krattenthaler, D. Yaqubi, Some determinants of path generating functions, II, *Adv. Appl. Math.*, 101 (2018), 232-265.
- [5] T. Mansour, N. Shattuck, Counting humps and peaks in generalized Motzkin paths, *Discrete Appl. Math.*, 161 (2013), 2213-2216.
- [6] T. Mansour, Counting peaks at height k in a Dyck path, *J. Integer Seq.*, 5 (2002), Article 02.1.1.
- [7] P. Peart, W.J. Woan, Dyck paths with no peaks at height k , *J. Integer Seq.*, 4 (2001), Article 01.1.3.
- [8] M. Petkovesk, H.S. Wilf, D. Zeilberger, *A=B*, AK Peters Ltd., 1996.
- [9] A. Regev, Humps for Dyck and for Motzkin paths, 2010, [arXiv: 1002.4504](https://arxiv.org/abs/1002.4504).
- [10] A. Regev, Probabilities in the (k, l) -hook, *Israel J. Math.*, 169 (2009), 61-88.
- [11] N.J.A. Sloane, The on-line encyclopedia of integer sequences, <https://oeis.org/>.
- [12] S.H.F. Yan, Bijective counting of humps and peaks in (k, a) -paths, *Discrete Appl. Math.*, 193 (2015), 208-212.
- [13] D. Yaqubi, M. Farrokhi D.G., H. Ghasemian Zoeram, Lattice paths inside a table, *Math. Commun.*, 28(2023), 181-201.
- [14] D. Zeilberger, The method of creative telescoping, *J. Symbolic Comput.*, 11 (1991), 195-204.