

Omega theorem for fractional sigma function

Yuan Qiu, Alexander B. Kalmynin

Abstract

The research in the subfield of analytic number theory around error term of summation of sigma functions possesses a history which can be dated back to the mid-19th century when Dirichlet provided an $O(\sqrt{n})$ estimation of error term of summation of $d(n)$ [MV07]. Later, G. Voronoi [Ivi12], G. Kolesnik [Kol82], and M.N. Huxley [Hux03] (to name just a few) contributed more on the upper bound on the error term of summation of sigma functions. As for Ω -theorems, G.H. Hardy was the first contributor. Later researchers on this topic include G.H. Hardy [Har29] and T.H. Gronwall [Gro13], but the amount of academic effort is much sparser than O -theorems. This research aims to provide a better Ω -bound for the error term of summation of fractional sigma function $\sigma_\alpha(n)$ on the range $0 < \alpha < \frac{1}{2}$, obtaining the result $\Omega((x \ln x)^{\frac{1}{4} + \frac{\alpha}{2}})$.

1 Introduction

Let $\sigma_\alpha(n)$ denote the fractional sigma function: $\sigma_\alpha(n) = \sum_{d|n} d^\alpha$. When $0 < \alpha < 1$, it can be shown by summation techniques that $\sum_{n \leq x} \sigma_\alpha(n) = \frac{\zeta(\alpha+1)}{\alpha+1} x^{\alpha+1} + O(x)$ [Apo13]. Nevertheless, fractional sigma function has not been a hotbed for research in bounding theorems. In fact, the $O(x)$ estimation has been the best known result.

Most research in this field has been diverted to improving the bounding theorem of $d(n)$, which is also called "Dirichlet divisor problem". The problem can be formulated as the estimation of $\Delta(x)$ in the well-known asymptotic formula $\sum_{n \leq x} d(n) = x \log x + (2\gamma - 1)x + \Delta(x)$.

In 1849, J.P.G. Dirichlet proved that $\Delta(x) = O(\sqrt{x})$ with the Dirichlet hyperbola method [MV07]. In the following years, mathematicians made efforts to improve the error term $O(\sqrt{x})$. In 1904, G. Voronoi improved the bound to $\Delta(x) = O(x^{1/3} \log x)$ [Ivi12]. In 1982, G. Kolesnik made an improvement to $x^{35/108+\varepsilon}$ [Kol82]. The best known bound is $\Delta(x) = O(x^{131/416+\varepsilon})$ due to M.N. Huxley [Hux03]. Omega-theorem of $\Delta(x)$ is also a direction of research. In 1915, G.H. Hardy proved that $\Delta(x) = \Omega(x^{\frac{1}{4}})$ [Har17]. There is a widely propagated conjecture that $\Delta(x) = O(x^{1/4+\varepsilon})$.

Bounding theorem for summation of $\sigma(n) = \sigma_1(n)$ have also been another target of research. From elementary summation techniques a basic O -theorem of summation of $\sigma(n)$ can be obtained: $\sum_{n \leq x} \sigma(n) = \frac{\pi^2 x^2}{12} + O(x \log x)$ [Har29]. The best known O -bound is obtained by A. Walfisz: $\sum_{n \leq x} \sigma(n) = \frac{\pi^2 x^2}{12} + O\left(x(\log x)^{\frac{2}{3}}\right)$.

As for Ω -theorems, In 1913, T.H. Gronwall proved the omega theorem $\Omega(x \log \log x)$ for $\sigma(n)$ [Gro13]. In 1987, Pétermann improved the bound to $\Omega_-(x \log \log x)$ [Pét87].

Our research primarily aims to produce an Ω -theorem for $\sum_{n \leq x} \sigma_\alpha(n)$ when $0 < \alpha < \frac{1}{2}$.

2 Main results

We provide a Ω -bound for the remainder term of summation of $\sigma_\alpha(n)$, which is better than any previously known bounds.

Our main goal is to prove the following theorem:

Theorem 1. *Let $\sigma_\alpha(n)$ denote the fractional sigma function: $\sigma_\alpha(n) = \sum_{d|n} d^\alpha$, $0 < \alpha < \frac{1}{2}$. Then*

$$\sum_{n \leq x} \sigma_\alpha(n) = \frac{x^{1+\alpha}}{1+\alpha} \zeta(1+\alpha) + x \zeta(1-\alpha) + E_\alpha(x) \quad (1)$$

where $E_\alpha(x) = \Omega\left((x \ln x)^{\frac{1}{4} + \frac{\alpha}{2}}\right)$.

Before proving this result, we will show that E_α can be approximated up to $O(x^{\frac{1}{4} + \frac{\alpha}{2}})$ by a certain sum of oscillating terms.

Theorem 2. *For any $x^{\frac{1}{2} + \alpha + \varepsilon} \ll N \ll x^{\frac{1}{2\alpha} - \varepsilon}$,*

$$E_\alpha(x) = \frac{x^{\frac{1}{4} + \frac{\alpha}{2}}}{\sqrt{2\pi}} \sum_{m \leq N} \sigma_\alpha(m) m^{-\frac{3}{4} - \frac{\alpha}{2}} \cos\left(4\pi\sqrt{mx} - \frac{\pi}{4}\right) + O(x^{\frac{1}{4} + \frac{\alpha}{2}}) \quad (2)$$

Note that in the above theorem and throughout the paper, ε indicates a arbitrarily small positive number.

In the process of proving Theorem 2, we also reveal the following bound of $E_\alpha(x)$:

Corollary 1.

$$E_\alpha(x) = O\left(x^{\frac{2\alpha^2 + 3\alpha + 1}{2\alpha + 3} + \varepsilon}\right) \quad (3)$$

This bound is better than $O(x)$ when $\alpha < \frac{\sqrt{17}-1}{4}$.

Note that we are aware of the following "combinatorial" Ω -bound of $E_\alpha(x)$, namely

Theorem 3. For $x \rightarrow +\infty$ we have

$$\sup_{n \leq x} \sigma_\alpha(n) = x^\alpha \exp \left(\left(\frac{1}{1-\alpha} + o(1) \right) \frac{(\ln x)^{1-\alpha}}{\ln \ln x} \right) \quad (4)$$

hence

$$E_\alpha(x) = \Omega \left(x^\alpha \exp \left(\left(\frac{1}{1-\alpha} + o(1) \right) \frac{(\ln x)^{1-\alpha}}{\ln \ln x} \right) \right) \quad (5)$$

and here we give a short proof.

Proof. We have

$$\begin{aligned} \sigma_\alpha(n) &= n^\alpha \sigma_{-\alpha}(n) = n^\alpha \prod_{p|n} (1 + p^{-\alpha} + O(p^{-2\alpha})) \\ &= n^\alpha \exp \left(\sum_{p|n} p^{-\alpha} + O \left(\sum_{p|n} p^{-2\alpha} \right) \right) \end{aligned} \quad (6)$$

Note that

$$\sup_{n \leq x} \sum_{p|n} p^{-\alpha} = \sum_{p|P(Y_x)} p^{-\alpha} \quad (7)$$

where $P(y) = \prod_{p \leq y} p$ and Y_x is the largest positive integer that satisfies $P(Y_x) \leq x$.

Hence

$$\sup_{n \leq x} \sigma_\alpha(n) = x^\alpha \exp \left(\sum_{p|P(Y_x)} p^{-\alpha} + O \left(\sum_{p|P(Y_x)} p^{-2\alpha} \right) \right) \quad (8)$$

If $\beta \geq 1$, $\sum_{p|P(Y_x)} p^{-\beta}$ is bounded.

If $0 < \beta < 1$, by Prime Number Theorem and Abel summation, we have

$$\sum_{p|P(Y_x)} p^{-\beta} \sim \frac{Y_x^{1-\beta}}{(1-\beta) \ln Y_x} \sim \frac{(\ln x)^{1-\alpha}}{(1-\beta) \ln \ln x} \quad (9)$$

Applying the above observations for $\beta = \alpha$ and 2α , we obtain

$$\sup_{n \leq x} \sigma_\alpha(n) = x^\alpha \exp \left(\frac{(\ln x)^{1-\alpha}}{\ln \ln x} \left(\frac{1}{1-\alpha} + o(1) \right) \right) \quad (10)$$

as desired. \square

3 Proof of Theorems

Consider $\sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^s} = \zeta(s)\zeta(s-\alpha)$. By Mellin's Inversion Formula, we have

$$\sum_{n \leq x} \sigma_{\alpha}(n) = \frac{1}{2\pi i} \lim_{T \rightarrow +\infty} \int_{c-iT}^{c+iT} f(s) \frac{x^s}{s} ds \quad (11)$$

where $f(s) = \sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^s} = \zeta(s)\zeta(s-\alpha)$ and we choose $c = 1 + \alpha + \varepsilon$ with $\varepsilon > 0$.

Nevertheless, we wish to remove the infinity in the expression and approximate

$$\sum_{n \leq x} \sigma_{\alpha}(n) \text{ with } I_1(x) = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} f(s) \frac{x^s}{s} ds.$$

Take $T \geq 1$, then

$$\begin{aligned} I_1(x) &= \frac{1}{2\pi i} \int_{c-iT}^{c+iT} f(s) \frac{x^s}{s} ds \\ &= \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \left(\sum_{n=1}^{\infty} \frac{\sigma_{\alpha}(n)}{n^s} \right) \frac{x^s}{s} ds \\ &= \sum_{n=1}^{\infty} \sigma_{\alpha}(n) \frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{x^s}{sn^s} ds \end{aligned} \quad (12)$$

Notice that now we can apply Perron's formula [Ten15] to (12):

$$\frac{1}{2\pi i} \int_{c-iT}^{c+iT} \frac{a^s}{s} ds = \begin{cases} 1 + O\left(\frac{a^c}{T|\ln a|}\right) & \text{if } a > 1 \\ O\left(\frac{a^c}{T|\ln a|}\right) & \text{if } a < 1 \end{cases} \quad (13)$$

and we have

$$\begin{aligned} I_1(x) &= \sum_{n \geq 1} \sigma_{\alpha}(n) \mathbb{1}_{n < x} + O\left(\frac{x^c}{Tn^c |\ln \frac{x}{n}|}\right) \\ &= \sum_{n \leq x} \sigma_{\alpha}(n) + O\left(x^c \sum_{n \geq 1} \frac{\sigma_{\alpha}(n)}{Tn^c |\ln \frac{x}{n}|}\right) \end{aligned} \quad (14)$$

Theorem 3 implies the following bound for $\sigma_{\alpha}(n)$:

Lemma 1.

$$\sigma_{\alpha}(x) = O(x^{\alpha+o(1)}) \quad (15)$$

We have that if $n \leq \frac{x}{2}$ or $n \geq 2x$, then $|\ln \frac{x}{n}| \geq \ln 2$; otherwise, $\ln \frac{x}{n} \geq \frac{|n-x|}{x}$. Plugging in these bounds into (14) and utilizing and Lemma (1), we get

$$\begin{aligned}
O\left(x^c \sum_{n \geq 1} \frac{\sigma_\alpha(n)}{Tn^c |\ln \frac{x}{n}|}\right) &= O\left(x^c \sum_{n \leq \frac{x}{2} \text{ or } n \geq 2x} \frac{\sigma_\alpha(n)}{Tn^c |\ln \frac{x}{n}|}\right) + O\left(x^c \sum_{\frac{x}{2} < n < 2x} \frac{\sigma_\alpha(n)}{Tn^c |\ln \frac{x}{n}|}\right) \\
&= O\left(x^c \sum_{n \geq 1} \frac{n^{\alpha+o(1)}}{Tn^c}\right) + O\left(x^{c+1} \sum_{\frac{x}{2} < n < 2x} \frac{n^{\alpha+o(1)}}{Tn^c |n-x|}\right) \\
&= O\left(\frac{x^c}{T} \sum_{n \geq 1} n^{\alpha-c+o(1)}\right) + O\left(\frac{x^{\alpha+1+o(1)}}{T}\right)
\end{aligned} \tag{16}$$

For the first term, further notice that

$$\sum_{n \geq 1} n^{\alpha-c+o(1)} < 1 + \int_1^\infty n^{\alpha-c+o(1)} dn = 1 - \frac{1}{1+\alpha-c+o(1)} < 1 + \frac{2}{\varepsilon} \tag{17}$$

So we have

$$O\left(x^c \sum_{n \geq 1} \frac{\sigma_\alpha(n)}{Tn^c |\ln \frac{x}{n}|}\right) = O\left(\frac{x^{c+o(1)}}{T}\right) + O\left(\frac{x^{\alpha+1+o(1)}}{T}\right) = O\left(\frac{x^{c+o(1)}}{T}\right) \tag{18}$$

Plugging back to (14), we get our desired estimation:

$$\sum_{n \leq x} \sigma_\alpha(n) = I_1(x) + O\left(\frac{x^{c+o(1)}}{T}\right) \tag{19}$$

Our following work is focused on determining the bound of error term in $I_1(x)$. Now that the integral involved in $I_1(x)$ is along the real axis $\sigma = c = 1 + \alpha + \varepsilon$, we can shift the contour to $\sigma = 1 + \alpha - c = -\varepsilon$ through Cauchy's residue theorem and then apply the functional equation of Riemann zeta function.

$$I_1(x) = -I_2(x) - I_3(x) - I_4(x) + \nu(x) \tag{20}$$

where

$$I_2(x) = \frac{1}{2\pi i} \int_{c+iT}^{1+\alpha-c+iT} f(s) \frac{x^s}{s} ds \tag{21}$$

$$I_3(x) = \frac{1}{2\pi i} \int_{1+\alpha-c+iT}^{1+\alpha-c-iT} f(s) \frac{x^s}{s} ds \tag{22}$$

$$I_4(x) = \frac{1}{2\pi i} \int_{1+\alpha-c-iT}^{c-iT} f(s) \frac{x^s}{s} ds \quad (23)$$

and

$$\begin{aligned} \nu(x) &= \text{Res} \left(f(s) \frac{x^s}{s}, \alpha + 1 \right) + \text{Res} \left(f(s) \frac{x^s}{s}, 0 \right) + \text{Res} \left(f(s) \frac{x^s}{s}, 1 \right) \\ &= \frac{x^{1+\alpha}}{1+\alpha} \zeta(1+\alpha) - \frac{\zeta(-\alpha)}{2} + x\zeta(1-\alpha) \end{aligned} \quad (24)$$

We first estimate I_2 and I_4 .

$$\begin{aligned} I_2(x) &= O \left(\int_{1+\alpha-c+iT}^{c+iT} f(s) \frac{x^s}{s} ds \right) \\ &= O \left(\int_{1+\alpha-c}^c |\zeta(\sigma+iT)\zeta(\sigma-\alpha+iT)| \frac{x^\sigma}{T} d\sigma \right) \\ &= O \left(\int_{1+\alpha-c}^c T^{\mu(\sigma)+\mu(\sigma-\alpha)+\varepsilon-1} x^\sigma d\sigma \right) \end{aligned} \quad (25)$$

where the last step comes from the following bound for $|\zeta(\sigma+it)|$ from [Tit86]:

Lemma 2.

$$|\zeta(\sigma+it)| \ll t^{\mu(\sigma)} \ln t, \mu(\sigma) = \begin{cases} 0 & \text{if } \sigma \geq 1 \\ \frac{1-\sigma}{3} & \text{if } \frac{1}{2} \leq \sigma < 1 \\ \frac{1}{2} - \frac{2\sigma}{3} & \text{if } \sigma < \frac{1}{2} \end{cases} \quad (26)$$

After integrating $T^{\mu(\sigma)+\mu(\sigma-\alpha)+\varepsilon-1} x^\sigma$ by partitioning $[1+\alpha-c, c]$ into separate intervals corresponding to different values of $\mu(\sigma)$ and $\mu(\sigma-\alpha)$, we get

$$I_2(x) = O \left(\frac{x^{1+\alpha+\varepsilon}}{T^{1-\varepsilon}} \right) \quad (27)$$

Similar results can be deduced for $I_4(x)$:

$$I_4(x) = O \left(\frac{x^{1+\alpha+\varepsilon}}{T^{1-\varepsilon}} \right) \quad (28)$$

Now our task is to estimate I_3 . We can apply the functional equation of zeta function on $f(s)$ in the integral.

Lemma 3. (*Functional equation of Riemann zeta function*)

$$\zeta(s) \pi^{-\frac{s}{2}} \Gamma\left(\frac{s}{2}\right) = \zeta(1-s) \pi^{-\frac{1-s}{2}} \Gamma\left(\frac{1-s}{2}\right) \quad (29)$$

Corollary 2.

$$\zeta(s)\zeta(s-\alpha)\Gamma\left(\frac{s}{2}\right)\Gamma\left(\frac{s-\alpha}{2}\right) = \zeta(1-s)\zeta(1+\alpha-s)\pi^{2s-\alpha-1}\Gamma\left(\frac{1-s}{2}\right)\Gamma\left(\frac{1+\alpha-s}{2}\right) \quad (30)$$

We define $\gamma(s)$ to be the factor in the functional equation of $f(s)$ by applying Corollary 2:

$$f(1+\alpha-s) = \gamma(s)f(s), \gamma(s) = \pi^{1+\alpha-2s} \frac{\Gamma\left(\frac{s}{2}\right)\Gamma\left(\frac{s-\alpha}{2}\right)}{\Gamma\left(\frac{1-s}{2}\right)\Gamma\left(\frac{1+\alpha-s}{2}\right)} \quad (31)$$

By utilizing (31) on $I_3(x)$, we get

$$\begin{aligned} I_3(x) &= \frac{1}{2\pi i} \int_{1+\alpha-c+iT}^{1+\alpha-c-iT} f(s) \frac{x^s}{s} ds \\ &= \frac{1}{2\pi i} x^{1+\alpha} \int_{c+iT}^{c-iT} f(s) \gamma(s) \frac{x^{-s}}{1+\alpha-s} ds \\ &= x^{1+\alpha} \sum_{m \geq 1} \sigma_\alpha(m) c_T(mx) \end{aligned} \quad (32)$$

where $c_T(y) = \frac{1}{2\pi i} \int_{c+iT}^{c-iT} \frac{y^{-s} \gamma(s)}{1+\alpha-s} ds$.

Now the most difficult task is to estimate $c_T(y)$.

To estimate $\gamma(s)$, we use the following standard bound for $\Gamma(\sigma + it)$ from Stirling's formula when t is large:

$$\Gamma(\sigma + it) = \sqrt{2\pi} (it)^{\sigma-\frac{1}{2}} e^{-\frac{\pi}{2}t} \left(\frac{t}{e}\right)^{it} (1 + O(t^{-1})) \quad (33)$$

By plugging in this into $\gamma(s)$, we have

$$\gamma(\sigma + it) = \left(\frac{t}{2\pi}\right)^{2\sigma-\alpha-1} e^{4\pi i\theta(t)} \left(1 + O\left(\frac{1}{t}\right)\right), \theta(t) = \frac{t}{2\pi} \ln\left(\frac{t}{2\pi}\right) - \frac{t}{2\pi} - \frac{1}{8} \quad (34)$$

Then

$$\begin{aligned} |c_T(y)| &= \Re \left(\frac{1}{\pi} \int_c^{c+iT} y^{-s} \frac{\gamma(s)}{1+\alpha-s} ds \right) \\ &= \Re \left(\frac{1}{\pi} \left(\int_c^{c+T} y^{-s} \frac{\gamma(s)}{1+\alpha-s} ds + \int_{c+T}^{c+iT} y^{-s} \frac{\gamma(s)}{1+\alpha-s} ds \right) \right) \\ &= O(y^{-c}) + \Re \int_1^T e^{4\pi i\theta(t) - it \ln y} \left(\frac{t}{2\pi}\right)^{2c-\alpha-1} \left(-\frac{1}{t}\right) (1 + O(t^{-1})) dt \end{aligned} \quad (35)$$

From [Ivi12], the following inequality holds:

$$\int_J e^{2\pi i g(t)} h(t) dt \ll \frac{\max |h(t)|}{\delta} \text{ if } |g'(t)| > \delta \text{ for all } t \in J \quad (36)$$

In our case $g(t) = 2\theta(t) - \frac{t \ln y}{2\pi}$, $g'(t) = \frac{1}{\pi} \ln\left(\frac{t}{2\pi}\right) - \frac{\ln y}{2\pi}$. When $y > 2T^2$, we have that $g'(t) < U(T)$ where $U(T)$ is a function of T .

Then $|c_T(y)| = O(y^{-c}) + O(y^{-c} T^{2c-\alpha-2}) = O(y^{-c} T^{2c-\alpha-2})$ when $y > 2T^2$.

When $y \leq 2T^2$, we move the contour of integration to $b = 1 + \frac{\alpha}{2}$.

Since there are no singularities in the contour of integration, we have

$$\begin{aligned} c_T(y) &= \frac{1}{2\pi i} \left(\int_{b-iT}^{b+iT} \frac{\gamma(s)}{\alpha+1-s} y^{-s} ds + \int_{b+iT}^{c+iT} \frac{\gamma(s)}{\alpha+1-s} y^{-s} ds + \int_{c-iT}^{b-iT} \frac{\gamma(s)}{\alpha+1-s} y^{-s} ds \right) \\ &= \frac{1}{2\pi i} \int_{b-iT}^{b+iT} \frac{\gamma(s)}{\alpha+1-s} y^{-s} ds + O\left(\int_b^c T^{2\sigma-\alpha-2} y^{-\sigma} d\sigma \right) \\ &= \frac{1}{2\pi i} \int_{b-iT}^{b+iT} \frac{\gamma(s)}{\alpha+1-s} y^{-s} ds + O\left(T^{-\alpha-2} \left(\left(\frac{T^2}{y}\right)^c + \left(\frac{T^2}{y}\right)^b \right) \right) \\ &= \frac{1}{\pi} \Re \int_1^T \frac{\gamma(\sigma+it)}{\alpha+1-b-it} y^{-b-it} dt + O(T^{\alpha+2\epsilon} y^{-c} + y^{-b}) \end{aligned} \quad (37)$$

Using (34) and $\frac{1}{a+1-b-it} = \frac{a+1-b+it}{(a+1-b)^2+t^2} = O\left(\frac{1}{t}\right)$,

$$\begin{aligned} c_T(y) &= \frac{i}{2\pi^2} y^{-b} \Re \left(\int_1^T e^{4\pi i \theta(t)} y^{-it} O((1+t^{-1})^2) dt \right) + O(T^{\alpha+2\epsilon} y^{-c} + y^{-b}) \\ &= \frac{i}{2\pi^2} y^{-b} \Re \left(\int_1^T e^{4\pi i \theta(t)} y^{-it} dt \right) + O(T^{\alpha+2\epsilon} y^{-c} + y^{-b} \ln T) \end{aligned} \quad (38)$$

From Lemma 2.1 in [FI05]:

Lemma 4. *If $2 \leq z \leq 2T$, then*

$$\int_1^T \left(\frac{t}{ez}\right)^{imt} dt = \sqrt{\frac{2\pi z}{m}} e^{\frac{\pi i}{4} - imz} + O\left(\left(\frac{1}{\sqrt{T}} + \left|\log \frac{z}{T}\right|^{-1}\right)\right) \quad (39)$$

We have

$$\begin{aligned} \int_1^T e^{4\pi i\theta(t)} y^{-it} dt &= -i \int_1^T \left(\frac{t}{2\pi e\sqrt{y}} \right)^{2it} dt \\ &= \begin{cases} -\sqrt{2}\pi i y^{\frac{1}{4}} e^{\frac{\pi i}{4} - 4\pi\sqrt{y}i} + O\left(\left(\frac{1}{\sqrt{T}} + \ln \frac{T}{2\pi\sqrt{y}}\right)^{-1}\right) & y \leq \frac{T^2}{4\pi^2} \\ O\left(\left(\frac{1}{\sqrt{T}} + \ln \frac{2\pi\sqrt{y}}{T}\right)^{-1}\right) & \frac{T^2}{4\pi^2} < y \leq 2T^2 \end{cases} \end{aligned} \quad (40)$$

Hence,

$$c_T(y) = \begin{cases} \frac{1}{\sqrt{2\pi}} y^{-\frac{3}{4} - \frac{\alpha}{2}} \cos(4\pi\sqrt{y} - \frac{\pi}{4}) + O\left(y^{-1 - \frac{\alpha}{2}} \left(\frac{1}{\sqrt{T}} + \ln \frac{T}{2\pi\sqrt{y}}\right)^{-1}\right) \\ + O(T^{\alpha+2\varepsilon} y^{-c} + y^{-1 - \frac{\alpha}{2}} \ln T) & y \leq \frac{T^2}{4\pi^2} \\ O\left(y^{-1 - \frac{\alpha}{2}} \left(\frac{1}{\sqrt{T}} + \ln \frac{2\pi\sqrt{y}}{T}\right)^{-1}\right) + O(T^{\alpha+2\varepsilon} y^{-c} + y^{-1 - \frac{\alpha}{2}} \ln T) & \frac{T^2}{4\pi^2} < y \leq 2T^2 \\ O(y^{-c} T^{2c-\alpha-2}) & y > 2T^2 \end{cases} \quad (41)$$

Let $y = mx$.

$$\begin{aligned} I_3(x) &= x^{1+\alpha} \sum_{m \geq 1} \sigma_\alpha(m) c_T(mx) \\ &= x^{1+\alpha} \left(\sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} y^{-\frac{3}{4} - \frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) \right. \\ &\quad + O\left(\left(\frac{T^2}{x}\right)^{\frac{\alpha}{2} + o(1)} x^{-1 - \frac{\alpha}{2}} + \left(\frac{T^2}{x}\right)^{\frac{\alpha}{2} + o(1) - 1} \sqrt{T} x^{-1 - \frac{\alpha}{2}}\right) \\ &\quad \left. + O\left(\left(\frac{T^2}{x}\right)^{\alpha+1-c+o(1)} T^{\alpha+2\varepsilon} x^{-c}\right) + O\left(\left(\frac{T^2}{x}\right)^{\frac{\alpha}{2} + o(1)} x^{-1 - \frac{\alpha}{2}} \ln T\right) + O(T^{2c-\alpha-2} x^{-c}) \right) \\ &= x^{1+\alpha} \sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} y^{-\frac{3}{4} - \frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) + O(T^{\alpha-\frac{3}{2}+o(1)} x^{1-o(1)}) \\ &\quad + O(T^{\alpha+o(1)} x^{-o(1)}) + O(T^{2c-\alpha-2} x^{1+\alpha-c}) \\ &= x^{1+\alpha} \sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} y^{-\frac{3}{4} - \frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) + O(T^{\alpha-\frac{3}{2}+o(1)} x^{1-o(1)}) \\ &\quad + O(T^{\alpha+o(1)} x^{-o(1)}) + O(T^{\alpha+2\varepsilon} x^{-\varepsilon}) \end{aligned} \quad (42)$$

Summing up (19) (20) (24) (27) (28) (42), we have

$$\begin{aligned}
\sum_{n \leq x} \sigma_\alpha(n) &= \frac{x^{1+\alpha}}{1+\alpha} \zeta(1+\alpha) - \frac{\zeta(-\alpha)}{2} + x\zeta(1-\alpha) \\
&\quad + x^{1+\alpha} \sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} y^{-\frac{3}{4}-\frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) \\
&\quad + O(T^{\varepsilon-1} x^{1+\alpha+\varepsilon}) + O(T^{\alpha-\frac{3}{2}+o(1)} x^{1-o(1)}) + O(T^{\alpha+o(1)} x^{-o(1)}) + O(T^{\alpha+2\varepsilon} x^{-\varepsilon}) \\
&= \frac{x^{1+\alpha}}{1+\alpha} \zeta(1+\alpha) - \frac{\zeta(-\alpha)}{2} + x\zeta(1-\alpha) \\
&\quad + x^{\frac{1}{4}+\frac{\alpha}{2}} \sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} m^{-\frac{3}{4}-\frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) \\
&\quad + O\left(x^{o(1)} \left(\frac{x^{1+\alpha}}{T} + T^\alpha\right)\right)
\end{aligned} \tag{43}$$

This gives

$$\begin{aligned}
E_\alpha(x) &= O\left(x^{\frac{1}{4}+\frac{\alpha}{2}} \left(\frac{T^2}{x}\right)^{\frac{1}{4}+\frac{\alpha}{2}}\right) + O\left(x^{o(1)} \left(\frac{x^{1+\alpha}}{T} + T^\alpha\right)\right) \\
&= O(T^{\frac{1}{2}+\alpha}) + O\left(\frac{x^{1+\alpha+o(1)}}{T}\right)
\end{aligned} \tag{44}$$

Pick $T = x^{\frac{2+2\alpha}{3+2\alpha}}$. Then $E_\alpha(x) = O\left(x^{\frac{2\alpha^2+3\alpha+1}{2\alpha+3}+\varepsilon}\right)$, proving Corollary 1.

Now fix $x^{\frac{3}{4}+\frac{\alpha}{2}+\varepsilon} \ll T \ll x^{\frac{1}{4\alpha}+\frac{1}{2}-\varepsilon}$. Then $O\left(\frac{x^{1+\alpha+O(1)}}{T}\right) = O\left(\frac{x^{1+\alpha}}{x^{\frac{3}{4}+\frac{\alpha}{2}}}\right) = O(x^{\frac{1}{4}+\frac{\alpha}{2}})$, $O(x^{o(1)}T^\alpha) = O((x^{\frac{1}{4\alpha}+\frac{1}{2}})^\alpha) = O(x^{\frac{1}{4}+\frac{\alpha}{2}})$. So

$$E_\alpha(x) = x^{\frac{1}{4}+\frac{\alpha}{2}} \sum_{mx \leq \frac{T^2}{4\pi^2}} \sigma_\alpha(m) \left(\frac{1}{\sqrt{2\pi}} m^{-\frac{3}{4}-\frac{\alpha}{2}} \cos\left(4\pi\sqrt{y} - \frac{\pi}{4}\right) \right) + O(x^{\frac{1}{4}+\frac{\alpha}{2}}) \tag{45}$$

In addition, let $x^{\frac{1}{2}+\alpha+\varepsilon} \ll N \ll x^{\frac{1}{2\alpha}-\varepsilon}$. Define

$$F_\alpha(x, N) = \sum_{n \leq N} \frac{\sigma_\alpha(n)}{n^{\frac{3}{4}+\frac{\alpha}{2}}} \cos(4\pi\sqrt{nx} - \frac{\pi}{4})$$

under this constraint on N . Our goal is to find $x \in (N^{\frac{2}{1+2\alpha}-2\delta}, N^{\frac{2}{1+2\alpha}-\delta})$ ($\delta > 0$) such that $F_\alpha(x, N) = \Omega((\ln x)^{\frac{1}{4}+\frac{\alpha}{2}})$.

Now we apply Soundararajan's Lemma [Sou03] on $F_\alpha(x^2, N)$ to give an Ω -theorem. The lemma is summarized in the following statement:

Lemma 5. (*Soundararajan's Lemma*) Let $(f(n))_{n=1}^{\infty}$ and $(\lambda_n)_{n=1}^{\infty}$ be sequences of nonnegative real numbers, (λ_n) non-decreasing, and $\sum_{n=1}^{\infty} f(n) < \infty$. Let $L \geq 2$ and Y be integers, $\gamma \in \mathbb{R}$ fixed. Let

$$S(t) = \sum_{n=1}^{\infty} f(n) \cos(2\pi\lambda_n t + \gamma) \quad (46)$$

Suppose \mathcal{M} is a finite set of positive integers such that $\{\lambda_m : m \in \mathcal{M}\} \subset [\frac{\lambda_Y}{2}, \frac{3\lambda_Y}{2}]$. Then for any real $X \geq 2$, there exists some $x \in [\frac{X}{2}, (6L)^{|\mathcal{M}|+1}X]$ such that

$$|S(x)| \geq \frac{1}{8} \sum_{m \in \mathcal{M}} f(m) - \frac{1}{L-1} \sum_{\{n|\lambda_n \leq 2\lambda_Y\}} f(n) - \frac{4}{\pi^2 X \lambda_Y} \sum_{n=1}^{\infty} f(n) \quad (47)$$

In our application, $S(x) = F_{\alpha}(x^2, N)$. We have $F_{\alpha}(x^2, N) = \sum_{n=1}^{\infty} f(n) \cos(2\pi\lambda_n x + \beta)$, where $\lambda_n = 2\sqrt{n}$, $\beta = -\frac{\pi}{4}$,

$$f(n) = \begin{cases} \frac{\sigma_{\alpha}(n)}{n^{\frac{\alpha}{2} + \frac{1}{4}}} & n \leq N \\ 0 & \text{otherwise} \end{cases} \quad (48)$$

Take $\mathcal{M} = [\frac{T}{4}, \frac{3T}{4}]$.

We require $\frac{X}{2} > N^{\frac{1}{1+2\alpha} - \delta}$ and $(6L)^{|\mathcal{M}|+1}X < N^{\frac{1}{1+2\alpha} - \frac{\delta}{2}}$. This gives $(6L)^{|\mathcal{M}|+1} = (6L)^{2T+1} < \frac{N^{\frac{\delta}{2}}}{2}$. We take L to be a large constant (say $L = 10^8$). Then $T < \frac{\delta \ln N}{4 \ln(6 \cdot 10^8)}$.

Take $T = \frac{\delta \ln N}{8 \ln(6 \cdot 10^8)}$.

$$\sum_{m \in \mathcal{M}} f(m) \geq \sum_{\frac{T}{4} \leq m \leq \frac{3T}{4}} m^{-\frac{3}{4} + \frac{\alpha}{2}} \gg T^{\frac{1}{4} + \frac{\alpha}{2}} \quad (49)$$

$$\sum_{\{n|\lambda_n \leq 2\lambda_Y\}} f(n) \ll T^{\frac{1}{4} + \frac{\alpha}{2}} \quad (50)$$

$$\frac{1}{X \lambda_T} \ll \frac{1}{X \sqrt{T}} \ll \frac{1}{\sqrt{\ln N}} N^{\delta - \frac{1}{1+2\alpha}} \quad (51)$$

$$\sum_{n=1}^{\infty} f(n) = \sum_{n \leq N} \frac{\sigma_{\alpha}(n)}{n^{\frac{3}{4} + \frac{\alpha}{2}}} \ll N^{\frac{1}{4} + \frac{\alpha}{2}} \quad (52)$$

So

$$\frac{1}{X\lambda_T} \sum_{n=1}^{\infty} f(n) \ll \frac{N^\delta}{\sqrt{\ln N}} N^{\frac{1}{4} + \frac{\alpha}{2} - \frac{1}{1+2\alpha}}. \quad (53)$$

Note that $N \ll x^{\frac{1}{2}}$. In order to confine $\frac{1}{X\lambda_T} \sum_{n=1}^{\infty} f(n)$ at the order of $(\ln x)^{\frac{1}{4} + \frac{\alpha}{2}}$, we need that $\frac{1}{4} + \frac{\alpha}{2} - \frac{1}{1+2\alpha} < 0$. Solving the inequality gives $\alpha < \frac{1}{2}$, which perfectly satisfies the assumption in Theorem 1.

Plugging (49) (50) (53) into Lemma 5, we get $|F_\alpha(x^2, N)| \geq T^{\frac{1}{4} + \frac{\alpha}{2}} = \Omega((\ln N)^{\frac{1}{4} + \frac{\alpha}{2}}) = \Omega((\ln x^2)^{\frac{1}{4} + \frac{\alpha}{2}})$, thus fulfilling our goal.

So $E_\alpha(x) = \Omega((x \ln x)^{\frac{1}{4} + \frac{\alpha}{2}})$.

4 Concluding remarks

It is highly probable that our bound $\Omega((x \ln x)^{1/4 + \alpha/2})$ is not optimal. However, trying to improve it using Soundararajan's method meets substantial obstacles, mainly because the function

$$\omega_\alpha(n) = \sum_{p|n} p^{-\alpha}$$

behaves very differently from $\omega = \omega_0$, which is the number of distinct prime divisors. To be more precise, ω_α is concentrated around its average, which is $O(1)$ in this case, hence the sets $\mathcal{M} \subset [1, N]$ such that $\sigma_\alpha(m)$ is large for all $m \in \mathcal{M}$ are much smaller than the analogous sets for $d = \sigma_0$.

Results like Theorem 2 are a consequence of a summation formula analogous to the Voronoi summation formula. The corresponding kernel of an integral transform takes form

$$W_\alpha(y) = \lim_{T \rightarrow \infty} \frac{1}{2\pi i} \int_{2-iT}^{2+iT} y^{-s} \frac{\Gamma(\frac{s}{2})\Gamma(\frac{s-\alpha}{2})}{(\alpha+1-s)\Gamma(\frac{1-s}{2})\Gamma(\frac{1+\alpha-s}{2})} ds.$$

One can prove that similarly to the original Voronoi summation, the kernel can be expressed in terms of Bessel functions, namely

$$W_\alpha(y) = \frac{1}{2 \sin(\frac{\pi\alpha}{2}) y^{\frac{1+\alpha}{2}}} (I_{-1-\alpha}(4\sqrt{y}) - I_{\alpha+1}(4\sqrt{y}) - J_{-\alpha-1}(4\sqrt{y}) - J_{\alpha+1}(4\sqrt{y})).$$

This function interpolates between the kernels for $d(n)$ and $\sigma(n)$ for $\alpha \in [0, 1]$.

References

[Apo13] Tom M. Apostol. *Introduction to analytic number theory*. Springer Science & Business Media, 2013.

- [FI05] John B Friedlander and Henryk Iwaniec. Summation formulae for coefficients of l-functions. *Canadian Journal of Mathematics*, 57(3):494–505, 2005.
- [Gro13] T.H. Gronwall. Some asymptotic expressions in the theory of numbers. *Transactions of the American Mathematical Society*, 14(1):113–122, 1913.
- [Har17] Gr H Hardy. On dirichlet’s divisor problem. *Proceedings of the London Mathematical Society*, 2(1):1–25, 1917.
- [Har29] G.H. Hardy. An introduction to the theory of numbers. 1929.
- [Hux03] Martin N Huxley. Exponential sums and lattice points iii. *Proceedings of the London Mathematical Society*, 87(3):591–609, 2003.
- [Ivi12] Aleksandar Ivic. *The Riemann zeta-function: theory and applications*. Courier Corporation, 2012.
- [Kol82] Grigori Kolesnik. On the order of $\zeta(1/2 + it)$ and $\delta(r)$. *Pacific Journal of Mathematics*, 98(1):107–122, 1982.
- [MV07] Hugh L Montgomery and Robert C Vaughan. *Multiplicative number theory I: Classical theory*. Number 97. Cambridge university press, 2007.
- [Pét87] Y-FS Pétermann. An ω -theorem for an error term related to the sum-of-divisors function. *Monatshefte für Mathematik*, 103(2):145–157, 1987.
- [Sou03] Kannan Soundararajan. Omega results for the divisor and circle problems. *arXiv preprint math/0302010*, 2003.
- [Ten15] Gérald Tenenbaum. *Introduction to analytic and probabilistic number theory*, volume 163. American Mathematical Soc., 2015.
- [Tit86] EC Titchmarsh. *The theory of the Riemann zeta-function*. The Clarendon Press Oxford University Press, 1986.