Numerical approach to compressible shallow-water dynamics of neutron-star spreading layers

ALEKSANDR RUSAKOV, PAVEL ABOLMASOV, AND OMER BROMBERG²

¹Saint Petersburg State University, 7/9 Universitetskaya nab., St. Petersburg, 199034, Russia; rusakov.sasha@outlook.com ²The Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel

ABSTRACT

A weakly magnetized neutron star (NS) undergoing disk accretion should release about a half of its power in a compact region known as the accretion boundary layer. Latitudinal spread of the accreted matter and efficient radiative cooling justify the approach to this flow as a two-dimensional *spreading layer* (SL) on the surface of the star. Numerical simulations of SLs are challenging because of the curved geometry and supersonic nature of the problem. We develop a new two-dimensional hydrodynamics code that uses the multislope second-order MUSCL scheme in combination with an HLLC+ Riemann solver on an arbitrary irregular mesh on a spherical surface. The code is suitable and accurate for Mach numbers at least up to 5-10. Adding sinks and sources to the conserved variables, we simulate constant-rate accretion onto a spherical NS. During the early stages of accretion, heating in the equatorial region triggers convective instability that causes rapid mixing in latitudinal direction. One of the outcomes of the instability is the development of a two-armed 'tennis ball' pattern rotating as a rigid body. From the point of view of a high-inclination observer, its contribution to the light curve is seen as a high-quality-factor quasi-periodic oscillation mode with a frequency considerably smaller than the rotation frequency of the matter in the SL. Other variability modes seen in the simulated light curves are probably associated with low-azimuthal-number Rossby waves.

Keywords: Hydrodynamics (1963), Astrophysical fluid dynamics (101), Stellar accretion (1578), Neutron stars (1108), Computational astronomy (293), Low-mass X-ray binary stars (939)

1. INTRODUCTION

A typical neutron star (NS) is born in the end of the evolution of a massive star and initially possesses a strong magnetic field and a large angular momentum. Ohmic losses in the crust (Bransgrove et al. 2018) and field burial during accretion (Payne & Melatos 2004) destroy the magnetic field, creating a population of old, weakly magnetized, rapidly rotating, relatively massive NSs observed as accretors in low-mass X-ray binaries (LMXBs, see Bahramian & Degenaar 2023).

The absence of strong magnetic fields allows LMXBs to have accretion disks extending all the way toward the NS surface (or the innermost stable circular orbit that might be slightly above the surface; this scenario is known as the accretion-gap case, see Kluzniak et al. 1990). The structure of the innermost parts of the flow, where accreting matter slows down from approximately Keplerian rotation in the disk to the much slower rotation of the NS, is understood in very little detail. This part of the flow is called accretion boundary layer, or simply boundary layer (BL).

Weakly magnetized NSs in LMXBs are among the brightest and the most numerous X-ray sources in the sky, and their

timing and spectral properties are being studied since the 70-s. The spectral and timing properties of the BL allow to separation of its contribution to the observed flux from that of the disk (Gilfanov et al. 2003). With advances in X-ray polarization measurement techniques, timing and spectral data are now complemented by polarization measurements for several sources (Ursini et al. 2024). Some of them demonstrate behavior unexpected from LMXBs, such as considerable variations of the polarization angles. All this creates a demand for new theoretical work and numerical simulations of the accretion flows onto weakly-magnetized NSs.

The rich phenomenology of NS LMXBs includes different quasi-periodic oscillations (QPOs), in particular kilohertz-timescale QPOs, kHz QPO (van der Klis 2000). Their periods are of the same order ar the free-fall time near the surface. Besides, the properties of kHz QPOs in NS systems are profoundly different from the short-timescale QPOs in the systems containing black holes (Ingram & Motta 2019). Their observational properties suggest that this specific oscillation mode must be related to the processes near the surface of the accreting star. Obviously, some version of a BL is a good candidate for the origin site of any fast variability of a weakly magnetized accreting NS.

First models of the BL treated it as a part of the disk (Papaloizou & Stanley 1986; Shakura & Sunyaev 1988) where the Keplerian rotation of the thin disk decreases to the much smaller value matching the spin of the star. One of the insights of such an approach is the result that the radial extent of the layer should be small compared to the thickness of the disk, making it essentially a three-dimensional flow with a complicated geometry. Modeling such a flow must take into account the latitudinal components of the velocities within it, that are usually ignored in accretion disk models.

An important step in understanding the BL dynamics is the concept of spreading layer (SL), introduced by Inogamov & Sunyaev (1999) and further developed in Suleimanov & Poutanen (2006) and Inogamov & Sunyaev (2010). In this approach, the latitudinal motions dominate over radial, and efficient radiative cooling from the surface makes the flow effectively two-dimensional, except for a narrow injection region near the equator. Existing analytic models of the SL are one-dimensional and steady-state.

One would expect the SL to have two- and threedimensional instability and variability modes. An analytical study of such oscillation modes and their stability was done by Watts et al. (2003). An important step in applying these results to real flows and making a self-consistent model of an SL would be two-dimensional numerical simulations. An attempt to consider supersonic two-dimensional dynamics on a sphere was done by Abolmasov et al. (2020), who have also found an inertial oscillation mode potentially responsible for the kHz OPO mentioned above. The method used in this paper to address the flow dynamics was by solving the system of equations in terms of spherical harmonics. For subsonic motions, this is an extremely reliable technique, usually applied to shallow-water equations (Jakob-Chien et al. 1995). Deriving the system of compressible shallow-water equations is straightforward.

The real NS SLs are expected to be moderately supersonic, with the rotational Mach number spanning the range between several and several tens in different parts of the flow. For the spectral approach, supersonic flows are hard to simulate because of the Gibbs phenomenon (Gottlieb & Shu 1997) caused by shock fronts and strong density and pressure contrasts. To be able to handle correctly the dynamics of an SL, a hydrodynamic code should work reliably at high Mach numbers. Besides, correct description of the flow close to the poles of the sphere requires that the code should not have a singularity in the polar regions. A solution to all these numerical problems is to develop a numerical code for solving compressible shallow-water equations using a mesh without singular points and is able to handle large Mach number and large density and pressure contrasts and discontinuities.

In this paper, we present a new numerical code SPLASH (SPreading LAyer Simulation for Hydrodynamics) that

solves the compressible shallow-water equations on a sphere using an arbitrary irregular mesh with help of a second-order MUSCL scheme. The paper is organized as follows. In Section 2, we introduce the main equations and physical quantities. In Section 3, the numerical algorithm is described. In Section 4, we give the results of stability tests, and in Section 5 the results of our simulations of an SL on a spherical star. We discuss the results in Section 6 and make conclusions in Section 7.

2. PHYSICAL PROBLEM

2.1. Typical values and basic equations

We will consider a spherical NS with typical global parameter values: mass $M=1.5 \rm M_{\odot}$ and radius $R=10 \rm km$. Its rotation will be considered negligibly slow, but we focus on the dynamics of an envelope, that may rotate at a considerable fraction of the Keplerian velocity. We will consider a fixed mass accretion rate of $\dot{M}=10^{-8} \rm M_{\odot} \, yr^{-1}$. In terms of emitted luminosity, this is typical for a Z-source similar to Sco X-1 (Hasinger & van der Klis 1989). The source of accreting matter has net angular momentum about 80% of Keplerian and is limited to an equatorial band as one would expect for a source accreting through a disk.

For a mass accretion rate close to the Eddington limit, the expected effective temperature of the source is ~ 1keV. On long enough time scales, the matter in the layer slows down due to some kind of (very weak) interaction with the crust of the NS, cools down and joins the crust. The time needed to lose the excess angular momentum may be very long. According to Inogamov & Sunyaev (1999), it should be of the order of thousands of Keplerian times. The correlation times related to the parallel-tracks phenomenon (van der Klis 2001; Abolmasov & Poutanen 2021) suggest even longer times of the order millions of Keplerian scales. Slow dissipation in the SL allow to estimate its surface density as $\Sigma \sim \dot{M}t_{\rm corr}/4\pi R^2 \sim 10^8 {\rm g \, cm^{-2}}$. Below, we will consider an initial configuration with a mean surface density of $\sim 10^7 \mathrm{g \, cm^{-2}}$. The internal temperature in the layer is $T \sim 100 \text{keV}$, implying that radiation pressure is important and matter is mostly ionized (see also estimates in Inogamov & Sunyaev 1999 and Abolmasov et al. 2020). The speed of sound is $c_s \sim 0.01c$, where c is the speed of light. The velocity of the accreting matter reaches about 0.5c, hence the expected Mach number is of the order of several tens.

The equations we are solving are vertically integrated equations of ideal hydrodynamics, including the continuity equation

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho v) = 0, \tag{1}$$

Euler equations

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{1}{\rho}\nabla P - \mathbf{g},\tag{2}$$

and energy conservation equation

$$\frac{\partial}{\partial t} \left(e + \frac{v^2}{2} \rho - \Phi \rho \right) + \nabla \cdot \left[\left(e + P + \frac{v^2}{2} \rho - \Phi \rho \right) v + q \right] = 0, (3)$$

where e is internal energy density, P is pressure, $\Phi = GM/(R+h)$ is gravitational potential, $g = -\nabla \Phi$, and h is elevation above the surface. We use a spherical coordinate system where the position is given by a radius vector

$$r = (R + h) \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}, \tag{4}$$

and $h \ll R$. For the vertically integrated equations, we will only take into account tangential velocity components v_{θ} and v_{φ} along the coordinate vectors of the spherical coordinate system.

We treat these equations as conservation laws of the form

$$\frac{\partial}{\partial t}(u)(\mathbf{r},t) + \nabla \cdot \mathbf{F}_{u}(\mathbf{r},t) = S_{u}(\mathbf{r},t), \tag{5}$$

where u is the density of the particular conserved quantity, F_u is the corresponding flux vector, and S_u is the corresponding source. All the quantities are expressed as functions of the position on the sphere r and time t.

The quantities used in the calculations are surface density

$$\Sigma = \int \rho \mathrm{d}h,\tag{6}$$

angular momentum surface density

$$l = \int \rho(r \times v) dh, \tag{7}$$

and surface density of energy

$$E = \int \left(e + \frac{v^2}{2} \rho + \Phi \rho \right) dh. \tag{8}$$

The local energy density e is related to pressure P by the equations for a fluid with adiabatic index $\gamma = 4/3$,

$$e = \frac{P}{\gamma - 1}. (9)$$

For the integration scheme, we will use the values of the above five vertically integrated quantities in the centers of the cells of an irregular grid (see Section 3.2 for details). Mass flux is calculated as

$$\mathbf{F}_{\Sigma} = \Sigma \mathbf{v}.\tag{10}$$

For the other quantities, the fluxes and sources are introduced in the following subsections.

2.2. Velocity and angular momenta

A fluid element on the surface of the sphere tends to conserve its angular momentum. Gravity directed along the radius vector gives zero contribution to the angular momentum change, as the torque it creates is proportional to $r \times g = 0$. Including surface deformation due to rotation or tangential stresses would introduce force components along the surface that we will ignore in this paper. All the changes in angular momentum come from the forces not aligned with the radius vector, and they may be included in the right-hand side of the conservation laws. Angular momentum per unit surface area is

$$\boldsymbol{l} = \Sigma \boldsymbol{r} \times \boldsymbol{v}.\tag{11}$$

To convert the primitives using the conservatives, we need to divide the mass by the surface area of the cell K_i . To get velocities from l, one can take a vector product of (11) by r

$$r \times l = \Sigma ((r \cdot v) - vR^2) = -\Sigma R^2 v,$$
 (12)

where we neglect the next-order terms in h/R, assuming the layer is thin in vertical direction. The last step is possible because all the motions are restricted to the surface of the sphere, and thus are orthogonal to the radius vector. This means that the angular momentum vector contains all the necessary information to restore the velocity components, and

$$v = -\frac{r \times l}{\Sigma R^2}. (13)$$

Vector \boldsymbol{l} is a vector in 3D space, and thus needs to be defined in a global coordinate system unrelated to the coordinates on the sphere. We will use the three Cartesian components of the angular momenta.

Angular momentum flux is obtained by integrating the stress $T_{\alpha\beta} = \rho v_{\alpha}v_{\beta} + \delta_{\alpha\beta}P$ over the radial coordinate and taking a vector product with \mathbf{R} . Here, the Greek indices correspond to the spatial coordinates in the 3D space. Tensor formalism is applied to Cartesian 3D coordinates only and thus does not require distinction between covariant and contravariant components. Given \mathbf{n} is a unit normal vector to a cell edge tangential to the surface of the sphere, the flux of the α component of \mathbf{l} along the normal is

$$F_{l,\alpha} = l_{\alpha} \mathbf{v} - (\mathbf{n} \times \mathbf{R})_{\alpha} \Pi. \tag{14}$$

2.3. Vertical structure and energy

In the absence of vertical velocities, the vertical component of Euler equation becomes

$$\frac{\mathrm{d}P}{dr} = -g_{\mathrm{eff}}\rho,\tag{15}$$

where

$$g_{\text{eff}} = \frac{GM}{R^2} - \frac{v^2}{R}.\tag{16}$$

Velocity v here has two tangential components, v_{θ} (along the polar angle direction) and v_{φ} (azimuthal). If we assume that, for given coordinates on the sphere θ and φ , $P \propto \rho^{\gamma_{\gamma}}$ with changing h. We will assume this exponent is the same as the global adiabatic index, $\gamma_{\gamma} = \gamma$, but it is not necessarily the case in general. If the vertical structure is affected by radiation transfer, $\gamma_{\gamma} \simeq 4/3$. Vertical integration results in

$$\rho = \rho_0 \left(1 - \frac{h}{H} \right)^{\frac{1}{\gamma_V - 1}},\tag{17}$$

$$P = P_0 \left(1 - \frac{h}{H} \right)^{\frac{\gamma_V}{\gamma_V - 1}},\tag{18}$$

where H is the radius where density and pressure become zero,

$$H = \frac{\gamma_{\rm v}}{\gamma_{\rm v} - 1} \frac{P_0}{q_{\rm eff} \rho_0}.$$
 (19)

By direct integration, one can prove that

$$P_0 = q_{\text{eff}} \Sigma \tag{20}$$

and

$$H = \frac{2\gamma_{\rm v} - 1}{\gamma_{\rm v} - 1} \frac{\Pi}{q_{\rm eff} \Sigma}.$$
 (21)

Total surface energy density

$$E = \int_0^H \left(\frac{P}{\gamma - 1} + \rho \Phi + \rho \frac{v^2}{2}\right) dh =$$

$$= \frac{\gamma}{\gamma - 1} \Pi + \Sigma \frac{v^2}{2} + g_{\text{eff}} \Sigma \Delta R,$$
(22)

where ΔR is the deformation of the stellar surface related to rotation (the term is analogous to $\Delta \Phi$ used in Abolmasov et al. 2020). If we want the deformation to correspond to a certain equilibrium rotation figure, we can replace in the first order in ΔR ,

$$\frac{GM}{R_*^2} \Delta R \simeq -v_0^2/2 = -\frac{1}{2} \Omega_0^2 R_*^2 \sin^2 \theta.$$
 (23)

In such an approximation, total energy is

$$E = \frac{\gamma}{\gamma - 1} \Pi + \frac{1}{2} \Sigma \left(v_{\theta}^2 + v_{\varphi}^2 - \Omega_0^2 R_*^2 \sin^2 \theta \right). \tag{24}$$

In this paper, we restrict ourselves to the case of a spherical star, setting $\Omega_0 = 0$. Energy flux is found as

$$F_{E} = \int_{0}^{H} \left(\frac{\gamma P}{\gamma - 1} + \rho \Phi + \rho \frac{v^{2}}{2} \right) v dr =$$

$$= \left[\frac{2\gamma - 1}{\gamma - 1} \Pi + \frac{1}{2} \left(v_{\theta}^{2} + v_{\varphi}^{2} - \Omega_{0}^{2} R_{*}^{2} \sin^{2} \theta \right) \Sigma \right] v.$$
(25)

Thermal contribution to vertically integrated energy density contains coefficients that differ from their three-dimensional analogues. However, defining two-dimensional effective adiabatic exponent $\Gamma = 2 - 1/\gamma$, we can restore the original form of the expressions

$$E = \frac{1}{\Gamma - 1} \Pi + \frac{1}{2} \Sigma \left(v_{\theta}^2 + v_{\varphi}^2 - \Omega_0^2 R_*^2 \sin^2 \theta \right), \tag{26}$$

$$\boldsymbol{F}_{E} = \left[\frac{\Gamma}{\Gamma - 1} \Pi + \frac{1}{2} \left(v_{\theta}^{2} + v_{\varphi}^{2} - \Omega_{0}^{2} R_{*}^{2} \sin^{2} \theta \right) \Sigma \right] \boldsymbol{v}. \tag{27}$$

As a consequence, adiabatic relationship between vertically integrated pressure and density is $\Pi \propto \Sigma^{\Gamma}$. Hence, the speed of sound is also $c_s = \sqrt{\Gamma \Pi/\Sigma}$.

2.4. Sources and sinks

Deformation of the NS surface is included in energy equation through the potential terms in equations (24) and (27), hence there is no need to include the associated work as a source of energy. However, the deformation should be included in the angular momentum source as

$$\dot{l}_{\Phi} = \Sigma \mathbf{r} \times \mathbf{g} = \mathbf{e}_{\theta} \times \mathbf{r} \ \Sigma \Omega_0^2 R_* \sin \theta \cos \theta. \tag{28}$$

In order to properly simulate accretion and other physical effects relevant for the SL problem, we include additional source and sink terms. Accretion is included as a mass source dependent on the coordinates as

$$\dot{\Sigma}_{\rm acc} = \frac{\dot{M}}{4\pi R^2} \frac{1}{\sqrt{2\pi}\sigma_{\alpha}} e^{-\frac{\cos^2 \alpha}{2\sigma_{\alpha}^2}},\tag{29}$$

where α is the polar angle with the axis of the disk that we assume here to be inclined by $\alpha_{\rm tilt} = 6^{\circ}$ with respect to the rotation axis of the NS, and σ_{α} determines the width of the band within which accretion occurs.

$$\cos \alpha = \frac{r \cdot \Omega_{\text{orb}}}{|\Omega_{\text{orb}}|} = \cos \theta \cos \alpha_{\text{tilt}} + \sin \theta \sin \alpha_{\text{tilt}} \sin \varphi, \quad (30)$$

where Ω_{orb} is the orbital rotation frequency of the accreting matter. This mass source distribution and the associated injection velocity profile (see below) correspond to a thin accretion disk with the relative thickness $\sim 10\%$.

We assume that the accreting matter has net angular momentum (aligned with the symmetry axis of the source) and kinetic and internal energy, that allows to write down the accretion-associated sources as

$$\dot{\boldsymbol{l}}_{\rm acc} = \dot{\Sigma}_{\rm acc} \left(\boldsymbol{r} \times \boldsymbol{v}_{\rm orb} \right) \tag{31}$$

and

$$\dot{E}_{\rm acc} = \dot{\Sigma}_{\rm acc} \left(\left(\frac{E}{\Sigma} \right)_{\rm d} + \frac{1}{2} (\boldsymbol{v}_{\rm orb} - \boldsymbol{v})^2 \right). \tag{32}$$

We assume $v_{\rm orb}=r \times \Omega_{\rm orb}$ to be slightly lower than the Keplerian velocity at the surface. We consider $\Omega_{\rm orb}$ to be constant.

The first sink term we introduce is the loss of mass from the bottom of the spreading layer. Mass sink is locally proportional to surface density as

$$\dot{\Sigma}_{\text{fall}} = -\frac{1}{t_{\text{fall}}} \Sigma. \tag{33}$$

We define the constant t_{fall} from the assumption of mass balance in the spreading layer. It has a physical meaning of the e-folding time for the surface density.

$$t_{\text{fall}} = \frac{M_{\text{tot}}}{\dot{M}}.$$
 (34)

Here, \dot{M} is the accreted mass, $M_{\rm tot}$ is the initial mass of the SL.

If mass is lost from the bottom of the layer at a rate $\dot{\Sigma}$ (for mass surface density), this loss should be accompanied with the loss in angular momentum and energy, proportional to $\dot{\Sigma}$. Associated momentum loss is

$$\dot{l}_{\text{fall}} = \frac{l}{\Sigma} \dot{\Sigma}_{\text{fall}},\tag{35}$$

meaning that the net angular momentum of the matter locally lost from the system is the local vertically average value. The amount of energy lost from the cell is defined by the vertical energy flux (Landau & Lifshitz 1987, section I.6)

$$F_{E,r} = v_r \rho \left(\frac{v^2}{2} + \frac{\gamma_v}{\gamma_v - 1} \frac{P_0}{\rho_0} \right). \tag{36}$$

We do not know the vertical velocity $v_r \ll v$, but $\rho v_r = \dot{\Sigma}_{\rm fall}$ (mass flux through the surface), that allows (taking into account the results of Section 2.3) to re-write the expression as

$$\dot{E}_{\text{fall}} = \dot{\Sigma}_{\text{fall}} \left(\frac{v^2}{2} + \frac{\Gamma}{\Gamma - 1} \frac{\Pi}{\Sigma} \right). \tag{37}$$

Another sink term we introduce is the radiation energy loss in a way it is described in Abolmasov et al. (2020):

$$\dot{E}_{\rm rad} = -\frac{cg_{\rm eff}}{\varkappa} (1 - \beta). \tag{38}$$

Here, κ is the Rosseland average opacity (we assume Thomson scattering the main opacity source, that implies $\kappa \simeq \kappa_{\rm T} \simeq 0.34~{\rm cm^2~g^{-1}}$), $g_{\rm eff}$ is given by equation (16), and β is the ratio of gas pressure to the total pressure of gas and radiation. β can be computed by solving

$$\frac{\beta}{(1-\beta)^{1/4}(1-\beta/2)} = \frac{12}{5} \frac{k}{m} \left(\frac{3}{4} \frac{c}{\sigma_{\rm SB}} g_{\rm eff} \Sigma \right)^{1/4} \frac{\Sigma}{E}.$$
 (39)

Here, m is the mass of a hydrogen atom, k is the Boltzmann constant, and σ_{SB} is the Stefan–Boltzmann constant.

In order to avoid numerically solving this algebraic equation at every time step, we introduce an approximation for the left-hand side of equation (39)

$$\frac{\beta}{(1-\beta)^{1/4}(1-\beta/2)} \approx \begin{cases} \frac{\beta}{1-\beta} & \beta \le \beta_{\text{switch}} \\ \frac{2}{\sqrt[4]{1-\beta}} & \beta > \beta_{\text{switch}}, \end{cases}$$
(40)

where $\beta_{\text{switch}} \simeq 0.736$. The precise value of β_{switch} is chosen to match the values given by the two approximations given above

Thus, the general expressions for the source terms we use are

$$\dot{\Sigma} = \dot{\Sigma}_{acc} + \dot{\Sigma}_{fall},
\dot{I} = \dot{I}_{\Phi} + \dot{I}_{acc} + \dot{I}_{fall},
\dot{E} = \dot{E}_{acc} + \dot{E}_{fall} + \dot{E}_{rad},$$
(41)

where the individual source/sink terms are described by the equations given earlier this section.

3. NUMERICAL ALGORITHM

3.1. MUSCL scheme for general unstructured meshes

The conservative scheme we present here is based on the multislope MUSCL (Monotonic Upstream-centered Scheme for Conservation Laws van Leer (1979)) method for general unstructured meshes described in Touze et al. (2015). The main advantage of this scheme is its second order of error, that does not come at the cost of stability. It demonstrates lower accuracy compared to the ENO and WENO (Shu 2020); however, it is possible to implement on an unstructured mesh. We adapt the 2D method for use on an arbitrary spherical mesh. We tested several different meshes (Figure 1) in order to find which of them worked best for the given problem.

MUSCL scheme works for any hyperbolic conservation equation system with initial condition. The general form for each of the equations of the system is

$$\partial_t u(\mathbf{x}, t) + \nabla F(\mathbf{x}, t) = \dot{u} \quad u(\mathbf{x}, t = 0) = u_0, \tag{42}$$

where $u = \Sigma$, l, and E, fluxes are calculated using equations (10), (14), and (27), and the right-hand side contains the sources given by equation (41).

As we are dealing with a sphere, the boundary condition is not necessary. Let $\Omega \subset R^2$ be a domain, $x \in \Omega$. Let us look at a discretization of Ω made up of polygons K_i . We can define V(i) as the vicinity of the element K_i , defined as the set of neighboring elements K_j that share with K_i a common edge $S_{ij} = K_i \cap K_j$. U_i^n defines the discrete value of a conservative variable u in the center of the face K_i at the moment $t_n = t_0 + n\Delta t$. Then the numeric scheme can be written as

$$U_i^{n+1} = U_i^n - \Delta t \sum_{V(i)} \frac{S_{ij}}{K_i} \phi(U_{ij}^n, U_{ji}^n, \mathbf{n}_{ij}, \mathbf{F}),$$
 (43)

where S_{ij} is the length of the edge between K_i and K_j , K_i is the surface area of a cell, ϕ is the Riemann solver that is dependent on the physical flux F, U_{ij}^n , U_{ji}^n are the reconstructed values, n_{ij} is the unit normal vector of the edge directed outwards.

In order to find a reconstructed value of a conservative variable at the edge center M_{ij} (see Figure 2) we need to find forward and backward slopes p_{ij}^+ and p_{ij}^- . With $r=p_{ij}^-/p_{ij}^+$ we can introduce a flux limiter function $\varphi(r)$ that turns two slopes into one limited slope in order to avoid spurious oscillations. The flux limiter should also be dependent on the local geometry. Dropping the temporal index n, the reconstructed values can be defined as

$$U_{ij} = U_i + p_{ij}^+ \varphi(r) || \mathbf{B}_i \mathbf{M}_{ij} ||.$$
 (44)

We can find slopes by finding 2 points on our mesh: H_{ij}^+ and H_{ij}^- . If we know the values of the conservative variables in this point, we can find the slopes using

$$p_{ij}^{+} = \frac{U_{H_{ij}^{+}} - U_{i}}{\|B_{i}H_{ij}^{+}\|}, \quad p_{ij}^{-} = \frac{U_{i} - U_{H_{ij}^{-}}}{\|B_{i}H_{ij}^{-}\|}.$$
 (45)

The process of finding these "special" points depends on the mesh. An example is given on the Figure 2. Let us denote the set of all neighboring faces of a cell K_i as W(i). It is different from V(i) because W(i) also contains neighboring faces of K_i that share only one common vertice. Connecting the face center B_i with the point M_{ij} results in a line. We have to find $K_{ij_1}^-$, the most backward neighboring element of K_i . (in Fig 2 it is a face with a center $B_{ij_1}^-$) We can define it formally as

$$\cos(B_{ij_1}^- B_i, B_i M_{ij}) = \min_{k \in W(i)} \cos(B_k B_i, B_i M_{ij}).$$
 (46)

Then we also have to define K_{ij_2} , the second most backward neighboring element of K_i that is located on the other side of the line B_iM_{ij} .

$$\cos(\boldsymbol{B}_{ij_2}^{-}\boldsymbol{B}_i,\boldsymbol{B}_i\boldsymbol{M}_{ij}) = \min_{\boldsymbol{k} \in \overline{\boldsymbol{W}(i)}} \cos(\boldsymbol{B}_{\boldsymbol{k}}\boldsymbol{B}_i,\boldsymbol{B}_i\boldsymbol{M}_{ij}). \tag{47}$$

$$\overline{\mathcal{W}}(i) = \{k \in \{\mathcal{W}(i) \setminus K_{ij_1}^-\} | \sin(B_k B_i, B_i M_{ij}^\perp) \cdot \\ \cdot \sin(B_{ij_1}^- B_i, B_i M_{ij}^\perp) \le 0\}.$$

$$(48)$$

Using a similar algorithm, we can find $B_{ij_1}^+$ and $B_{ij_2}^+$. Then, by finding the intersections between lines $B_i M_{ij}$ and $B_{ij_1}^+ B_{ij_2}^+$ or $B_{ij_1}^- B_{ij_2}^-$, we are able to find points H_{ij}^+ and H_{ij}^- . Finally, in order to find our conservative values in these points, we have to introduce barycentric coordinates

$$\beta_{ij_1}^- = \frac{\|B_{ij_2}^- H_{ij}^-\|}{\|B_{ij_1}^- B_{ij_2}^-\|}, \quad \beta_{ij_2}^- = \frac{\|B_{ij_1}^- H_{ij}^-\|}{\|B_{ij_1}^- B_{ij_2}^-\|}, \tag{49}$$

$$\beta_{ij_1}^+ = \frac{\|B_{ij_2}^+ H_{ij}^+\|}{\|B_{ij_1}^+ B_{ij_2}^+\|}, \quad \beta_{ij_2}^+ = \frac{\|B_{ij_1}^+ H_{ij}^+\|}{\|B_{ij_1}^+ B_{ij_2}^+\|}. \tag{50}$$

Then the variable values at points H_{ij}^+ and H_{ij}^- can be found as

$$U_{H_{ij}^{-}} = \beta_{ij_{1}}^{-} U_{ij_{1}}^{-} + \beta_{ij_{2}}^{-} U_{ij_{2}}^{-}, \quad U_{H_{ij}^{+}} = \beta_{ij_{1}}^{+} U_{ij_{1}}^{+} + \beta_{ij_{2}}^{+} U_{ij_{2}}^{+}.$$
 (51)

3.2. Spherical adaptation and meshes

We use meshes created using an algorithm from Sieger (2021) shown in Figure 1. Further, we will use cubic, triangular, and hexagonal mesh. We abstain from using UV-mesh, as it is less symmetrical than the other options; moreover, its polar area is not optimal for the reconstruction algorithm due to having a large number of faces sharing the same vertex.

Cubic, triangular, and hexagonal meshes are based on cube, icosahedron, and dodecahedron, respectively. The refinement of cubic and triangular meshes is achieved by subdividing each face into 4 parts and then projecting the resulting vertices onto the sphere. This results in a fast and consistent way of creating a fine mesh. One issue with this approach is the impossibility to directly set the desired number of faces. The hexagonal mesh is generated by projecting the centers of a triangular mesh onto a sphere and using them as new vertices.

Different types of mesh have different defining traits. The triangular mesh has a large number of similarly sized faces, so it is more computationally expensive. The cubic mesh has fewer faces on the same level of subdivision; however, the shape and size of these faces differ. The hexagonal mesh shares an edge with every neighbor, although some of the faces are pentagons that are considerably smaller than their neighboring faces.

In order to apply the algorithm to a spherical mesh, several adaptations have to be made. We can treat meshes either like polyhedra or as spherical tessellations. The first case requires us to work with broken straight lines, while the second one turns straight lines into great circle arcs and requires spherical trigonometry. After multiple tests, we decided to use spherical tessellation as it is more physically correct and no 'mountains' or 'trenches' are introduced into geometry.

Every face is treated as a spherical polygon with vertices $\{r_k\}_{k=1}^n$. The center of a cell R_i is therefore defined as

$$\boldsymbol{R}_{i} = \frac{\sum_{k=1}^{n} \boldsymbol{r}_{k}}{\left|\sum_{k=1}^{n} \boldsymbol{r}_{k}\right|}.$$
 (52)

 r_c is located on the same unit sphere as R_i .

One of the main differences lies in the choice of edge normals that are needed for a Riemann solver $\phi(U_{ij}^n, U_{ji}^n, n_{ij}, F)$. As we were going to reconstruct the velocities on the edges, we need to ensure that if the velocity vector is orthogonal to




Figure 1. Different meshes used from Sieger (2021).




Figure 2. Geometric reconstruction for 2D configuration.

the edge then $(n_{ij} \cdot v) = ||v||$. The edge normals are defined as follows: if r_1, r_2 are vertices that define an edge, then

$$n_{ij} = \frac{(r_1 + r_2) \times (r_1 - r_2)}{\|(r_1 + r_2) \times (r_1 - r_2)\|}.$$
 (53)

This gives a normal unit vector that is tangent to the surface of the sphere in the center of the edge.

3.3. Choice of the Riemann solver and limiter

One of the biggest challenges of the problem we are solving is the presence of large velocity differences. Because our initial state includes rigid body rotation, the velocity around the poles is close to zero. On the other hand, the accreted matter that falls onto the surface as a source term can reach Mach numbers $\mathcal{M} \sim 10$ and higher, up to 50 (see Section 2). Being able to solve the Riemann problem in both cases required us to choose an all-speed Riemann solver. We chose HLLC+ from Chen et al. (2020), that is based on the original approximate Riemann solver of Harten–Lax–van Leer with contact restoration and includes multiple wave fixes for

both low and high Mach numbers. We also tested the regular HLLC Riemann solver, as well as the HLLE for isothermal system of equations.

We introduce several changes into HLLC+ in order to adapt it to our choice of conservatives. HLLC (Harten-Laxvan Leer with Contact restoration) based Riemann solvers use notations of F(U) = vU + pD. In our case $v = v \cdot n$. Replacing p with Π we can set $D = [0, -(n \times R), S_*]$, where S_* is the speed of the middle wave in HLLC Riemann solver. Additionally, we replace normals n with $-(n \times R)$ in fixes that were introduced in HLLC+.

As for the limiter, we used the hybrid limiter introduced in Touze et al. (2015). It is proven to be stable for multislope MUSCL reconstruction, and it was the one that worked best in most of the tests.

3.4. Time-step stability conditions and temporal integration

In Touze et al. (2015), the authors provide sufficient stability condition and time-step limitations. Unfortunately, this condition results in unreasonably small Δt , which required us to choose a time step from other considerations. The sufficient condition allows that larger time steps without making the scheme unstable. We choose a CFL-based time step that accounts for the highest speed on the mesh in a given snapshot. If h_0 is the length of the shortest edge,

$$\Delta t = \frac{h_0 \cdot \text{CFL}}{\max\left(\max_{i \in \Omega} |v_i|, \max_{i \in \Omega} c_i\right)}.$$
 (54)

After series of different tests, the CFL value of 0.3 proved to provide stability in most cases and was used to generate all of the test results presented further.

As for the temporal integration, we use explicit secondorder Runge-Kutta integrator. Since the reconstruction al-




Figure 3. Difference between a 2D mesh, a spherical polyhedron mesh and a spherical tessellation.

gorithm has a second order of numerical error, we needed a corresponding temporal integrator.

4. CODE TESTS

Based on the algorithm described above, a numerical code has been written in C++ programming language. It is publicly available in the online repository ¹. The code supports OpenMP parallelism on one node.

In order to ensure the robustness of the algorithm as well as to test its capabilities, a set of tests has been conducted. These tests were selected in a way that allows to confirm the applicability of the code to different scenarios that may occur in real physical problems. Below is a list of the results of applying the code to different physical problems (steady-state and dynamic).

In this section and in the following Section 5, we will present two-dimensional maps of vertically-integrated pressure and density in Gall–Peters cylindrical equal-area projection from Gall (1885) defined as

$$x = \frac{\varphi}{\sqrt{2}}, \quad y = \sqrt{2}\cos\theta. \tag{55}$$

4.1. Stationary rotating atmosphere solution and mesh comparison

Two stationary analytical solutions are described in Appendix A. The scheme allows for the preservation of these profiles with minimal, yet still non-vanishing numerical error. The application of Riemann solver ensures that the expected analytical solution is recovered even with high Mach numbers of the flow.

We present the results of isentropic solution tests performed on a hexagonal, triangular, and cubic meshes with comparable number of faces. The initial pressure and density profiles have been selected as described in the isentropic solution described in Appendix A. The rotation is rigid-body with $\mathcal{M}_0 = 5$. Full simulation time is 10 full rotations. The results show conservation of full mass and energy to the ac-

curacy $\sim 10^{-14}$, proving that there are no numerical errors that lead to leaks or sources of conservative values (see Figure 4 for results on a hexagonal mesh with 10242 faces). An important property of the method is its conservation of static profiles. As shown in Figures 5, 6, and 7, the profile experiences only minor deviations after 10 rotations. Here we compare hexagonal, triangular, and cubic meshes with 10242, 20480, and 24576 faces, respectively. Due to the subdivision algorithm, we cannot set arbitrary numbers of faces while keeping them approximately the same size. With this in mind, we can only compare meshes with relatively similar numbers of faces. The results show that hexagonal mesh yields a bigger error near the 45th parallels, yet the general profile is preserved quite well. In further tests that required isentropic configuration, we used triangular or cubic meshes. These results allowed us to use this static profile as a baseline for further tests where we can safely add sink or source terms. The Mach number in this solution is ≤ 2.5 , as rigid-body rotating isentropic configuration always retains moderate Mach numbers (see Appendix A).

We have performed similar tests for the equal-density solution; however, it was not stable due to a convective instability. Over time this instability led to equalization of entropy, thus creating previously discussed isentropic solution.

4.2. Shock test

A simple shock test that has been conducted. The initial conditions in the test consist of a static constant-density, constant-pressure configuration with a small size of R/3 region overpressured by a factor of 50. In order to avoid symmetry effects, the high-pressure zone was created at non-zero latitude of 17.2° and longitude of 28.6°. The mach number of this shock is $\mathcal{M} \approx 7$. The results demonstrated in Figure 8 show how the spot exploded and how the front of the shock moved over time.

Over time, the shock just bounces back and forward from the start point to the opposite side of a sphere. As we do not introduce any friction yet, it persists for some time. We continued simulation for 3 total rotations and observed the

 $^{^{1}\} https://github.com/TURBOLOSE/MUSCL-scheme-on-spherical-mesh-WIP$




Figure 4. Relative error in conservation of total mass and total energy.




Figure 5. Conservation of Σ and Π profiles after 10 full rotations on a hexagonal mesh with 10242 faces.




Figure 6. Conservation of Σ and Π profiles after 10 full rotations on a triangular mesh with 20480 faces.




Figure 7. Conservation of Σ and Π profiles after 10 full rotations on a cubic mesh with 24576 faces.

initial spot compressing even further into a smaller area on a sphere before exploding once again.

4.3. *Split-sphere test*

The initial conditions for this test consist of an atmosphere that rotates in one direction above the equator and in the opposite direction below it. The cells on the equator are assumed to be initially static. Such a setup ensures that the transition is as sharp as possible, and the grid itself introduces perturbations to the axisymmetric velocity field. In each of the hemispheres, the rotation frequency is constant. The discontinuity in the rotation velocity triggers Kelvin–Helmholtz instability near the equator and forms multiple small vortices that merge over time. In Figure 9, we show the maps of the radial component of vorticity calculated as

$$\boldsymbol{\omega}_r = [\nabla \times \boldsymbol{v}]_r. \tag{56}$$

This quantity is sensitive to rotations and shears in the velocity field and has the same units as Ω , that allows to use the initial rotation frequency as normalization. Individual vortices are seen in the maps as local vorticity maxima.

The observed evolution towards larger spatial scales is typical for two-dimensional incompressible turbulence on a sphere. In particular, similar behavior was reproduced in Nättilä et al. (2024) for a more complex setup with several streams rotating in alternating directions.

5. ACCRETION SPREADING LAYER

Here we combine the initial equal-entropy atmosphere described in the Section 4.1 with additional source terms representing accretion and other physical effects from Section 2.4. The initial state is an equal-entropy atmosphere (see Appendix A) with the density at the poles $\Sigma_0 = 7.7 \times 10^5 \mathrm{g \ cm^{-2}}$, $\mathcal{M}_0 = 5$, and rotation frequency $\Omega = 300 s^{-1}$. Equatorial rotation velocity is thus $\sim 0.01 \mathrm{c}$. The average surface density for this solution is $\langle \Sigma \rangle = 10^7 \mathrm{g \ cm^{-2}}$. We also assume the speed of sound to be equal to $2 \times 10^{-3} \mathrm{c}$. We do not incorporate any NS deformation into our computation for now, so $\Omega_0 = 0$.

We set the accretion rate to a $10^{-8}M_{\odot}$ yr⁻¹, with the equatorial velocity $|v_{\rm orb}| = 0.4$ c, which is close to the Kepler velocity of 0.47c. $\left(\frac{E}{\Sigma}\right)_{\rm d}$ is set to 6×10^{-3} in code units. Accreted mass is distributed over the sphere according to the Gaussian law (equation 29), where we set $\alpha_{\rm tilt} = 6^{\circ}$ and $\sigma = 6^{\circ}$. Tilting the source allows to break the axial symmetry of the problem. From the physical point of view, this distribution corresponds to a thin disk with the thickness $\sim 0.1R$. Expected standard disk thickness at about the last stable orbit is (Shakura & Sunyaev 1973, equation 2.8)

$$\left(\frac{H}{R}\right)_{\rm st} = \frac{3}{8\pi} \frac{\varkappa \dot{M}}{cR} \left(1 - \sqrt{\frac{R_{\rm in}}{R}}\right),\tag{57}$$

that varies from ~ 0.5 away from the inner radius $R_{\rm in} \simeq 6GM/c^2$ to essentially zero near this radius. Making more accurate estimates of the disk thickness in this region would require a more accurate model of its transonic part near the last stable orbit.

We used a cubic mesh with 98304 faces. The simulation covers a time period of ~ 1 s. Results of the simulations are stored as snapshots of all the primitives in all the cells saved several times per rotation period and as high-cadence 'light curves' representing the properties of the object from the point of view of a distant observer (see later Section 5.1).

The simulation passes several stages shown in Figure 10. The accretion breaks the initial state by heating the equatorial regions and creates conditions for convective instability. The latter develops on the time scales of $\sim 1/\Omega$ and by $t \sim 0.05$ s enters a non-linear stage. As we expect for a Rayleigh-Taylor instability, the small-scale modes develop faster but are overrun by longer-wavelength modes, that finally merge to form the global 'tennis ball' pattern clearly visible in the last panel of Figure 10. The pattern is rotating around the z axis at a frequency close to the average angular frequency. During the later stages of evolution, this pattern is surprisingly stable and rotating at a frequency close to the initial rotation frequency of the layer. The low-density, low-pressure regions located within the meanders of the pattern are long-living cyclones with its own local velocity fields. In Figure 12, we show two density maps for the last snapshot of the simulation together with the velocity streamlines with rotation and without it. In order to subtract rotation for the second picture in Figure 12 we compute average linear rotation frequency as

$$v = \frac{1}{2\pi} \sum_{i} (\mathbf{R}_i \times \mathbf{v}_i)_z \csc^2 \theta_i.$$
 (58)

The linear frequency can be turned into a rotational frequency by multiplying it by 2π . After that, we compute rotational velocities and subtract them from the overall velocity field.

This subtraction allows to resolve the circular pattern of the strongest cyclone located in the southern hemisphere at $\theta \sim -60^{\circ}$ and $\varphi \sim 200^{\circ}$. After $t \sim 0.35$ s, this cyclone coincides with the global minimum of pressure, that allows to accurately trace its position and measure the rotation velocity of the 'tennis-ball' pattern.

It is important to note that rotation at later stages of simulation is not uniform and varies with latitude. For example, averaging out rotations in northern and southern hemispheres leads to a noticeable split in frequencies, as demonstrated in Figure 11.

5.1. Timing analysis

We compute light curves as follows: let \vec{R}_{obs} be the unit vector setting the direction towards the observer. Let ζ be a




Figure 8. Pressure maps in the shock test.

set of faces with radius vectors \vec{R}_i , such that $\vec{R}_{obs} \cdot \vec{R}_i > 0$. Then we find the light curves like:

$$L_{\rm c} = \sum_{i \in \zeta} \Pi_i K_i (\mathbf{R}_{\rm obs} \cdot \mathbf{R}_i), \qquad (59)$$

where K_i is the surface area of the i th cell, and \vec{R}_i is the normalized radius vector of its center given by equation (52). Calculating such a time series for weighed pressure allows to outline the most dynamically violent processes in the layer, that do not necessarily affect the radiative energy loss term (equation 38). We leave a more elaborate study of spectral variability within our model for a separate study. The series are calculated for four different observers with the inclination angles i = 0 (along the initial rotation axis, North pole), 45°, 90° (in the equatorial plane), and 180° (South pole).

As a result, we have four slightly inhomogeneous time series of $\sim 5 \times 10^5$ data points each. The inhomogeneity is caused by the variable time-step in equation (54) taking effect during some parts of the simulation. This led to a loss of even spacing between points of the time series.

During the simulation, the spreading layer is out of thermal balance, and all the variability curves are dominated by accretion-related heating. As a result, all the light curves are demonstrating a strong growing trend that needs to be

removed to study the variability on smaller time scales (of the order sound propagation time and rotation period). We subtract the trends using multiple connected splines of third order, split our observations into 10 even (in terms of the amount of data points) segments, and then compute a periodogram for each of them. We use the Lomb-Scargle algorithm (Lomb 1976; Scargle 1982) to account for the inhomogeneity of the timing data. This split allows us to see changes in the power density spectrum over time.

The results are presented in Figure 13. For the 90° observer, we can see entropic modes caused by rotation. Around 0.2s into simulation, one can see a signal at 2Ω that may be explained by the visibility of the 'tennis ball' structure.

The low-frequency mode is more complicated, as it is close to ν but not equal to it. After a certain point in the simulation, rotation starts to differ between the southern and northern hemispheres, which can be seen in Figure 11. The southern hemisphere, where the cyclone is located, rotates slower than the northern one. Here, instead of averaging ν on a sphere, we can assess the $\nu_{\rm cycl}$ —rotational frequency of a cyclone. We can do it by tracing its center over multiple snapshots. At its center, it creates a local minimum in pressure and maxi-




Figure 9. Vorticity plots for the split-sphere test. The merge of multiple smaller vortices into a bigger one can be observed.

mum in vorticity. Plotting $v_{\rm cycl}$ over the power spectra reveals that it is indeed the source of low-frequency oscillations for equatorial observer, as seen in Figure 13.

6. DISCUSSION

6.1. Oscillation modes and instabilities

Since rotation is supersonic, one may expect the variability modes to be dominated by Rossby modes. Their frequencies are for given latitude equal to $v_R(\theta) = v_e(\theta) + mv(\theta)$, where m is a whole number,

$$v_{\rm e} = \sqrt{2\nu \frac{\partial}{\partial \theta} \left(\nu \sin^2 \theta\right)},\tag{60}$$

and $v = v(\theta)$ is the local rotation frequency (see discussion section in Abolmasov et al. 2020). The observed power density spectrum depends on the contribution of different latitudes to the observed variability. One possibility is that the oscillations are excited at the latitude or latitudes where the rotation frequency is equal to the rotation rate of the double-wave pattern. For most of the time span, however, the pattern rotation frequency is outside the rotation frequency range within the flow. More importantly, Rossby waves may be excited at the location of the cyclone in the Southern hemisphere, and the strongest peaks seen in the lower panels of

Figure 13 correspond to the local v_c . The lower-frequency peaks in the polar periodograms are likely of different origin, as their frequencies do not depend on the rotation frequency during the most of the simulation time span.

Gravity modes are unlikely to be present in the simulated variability curves, as the configuration starts with neutral convective stability, and the heat released near the equator causes latitudinal mixing that restores an approximately flat entropy profile. Pressure modes might be present but confined to latitudinal strips $\sim 1/\mathcal{M}$ wide. Their frequencies may be estimated as

$$v_{\rm p} \simeq \frac{c_{\rm s}}{2\pi R} \simeq \frac{v}{\mathcal{M}}.$$
 (61)

In our setup, the observed temporal evolution is an outcome of the large velocity difference between the initial rigidly rotating configuration and the newly accreting matter from the disk. Depending on the mass accretion rate and cooling rate, this likely causes either Kelvin-Helmholtz (if accretion is fast enough to create a discontinuity in the velocity profile) or Rayleigh-Taylor instability. In the simulations we present in Section 5, heating in the equatorial region is much more important. As a result, the excess of entropy near the equator creates favorable conditions for Rayleigh-Taylor instability




Figure 10. Density maps of the accretion simulation. Six snapshots: initial state, instability start, instability evolution, and the final state. Full video of simulation is available on YouTube: https://youtu.be/9_TPXbtqDNY.




Figure 11. Average rotational frequency of northern and southern hemispheres.

with the wave numbers $k \lesssim 1/R\sigma_{\alpha}$. For a wave number larger than this value, the instability analysis is done in Appendix B. However, for the long-wavelength modes that dominate the evolution at $t \gtrsim 0.05$ s, the instability is classical Rayleigh-Taylor with the effective gravity equal to

$$q_{\theta, \text{eff}} \simeq \Omega^2 R \sin \theta \cos \theta,$$
 (62)

and the growth increment

$$\sigma_{\rm RT} \simeq \sqrt{gk} \simeq \Omega \sqrt{kR \sin \theta \cos \theta}.$$
 (63)

The value varies from $\sim 700 \text{Hz}$ for $k=1/R\sigma_{\alpha}$ to $\sim 200 \text{Hz}$ if $k\sim 1/R$, that ensures a rapid growth stage during the first several rotation periods in the SL simulation, that stops growing after entering non-linear regime at $t\sim 0.2 \text{s}$. After that, the density, pressure, and vorticity maps show remarkable stability and rigid-body-like rotation. Apparently, the longest instability wave number allowed by the topology of the sphere corresponds to the azimuthal number of m=2, that determines the azimuthal structure of the pattern. The rotation velocity of the pattern is close to the rotation velocity in the initial conditions, and grows much slower than the average velocity of the flow.

Starting from $t \sim 0.07$ s, there is a measurable difference in the rotation velocities and stream patterns between the Northern and the Southern hemispheres, reaching about 20% by the end of the simulation (see Figure 11). One of the most striking differences between the two hemispheres (probably related to the angular frequency asymmetry) is the formation of a single large cyclone in the Southern hemisphere. We suggest that this symmetry breaking is spontaneous and related to the rapid growth of the instability at earlier times. It may also be thought as a manifestation of a

more global heating instability discussed in Abolmasov et al. (2020): larger velocity difference in one of the hemispheres leads to larger energy dissipation, the latter creates latitudinal pressure asymmetry that pushes the accreting matter towards the hemisphere already rotating faster. Asymmetry in accreted mass would ultimately stop this instability, but the relevant time scale is difficult to estimate because of the large number of factors such as cooling rate, mass sources and sinks, and velocity profiles, are at play at the same time. More simulations are needed to check this hypothesis.

A valid question is if the initial conditions we use correspond to some real stage in the evolution of an accreting NS. We start with a gaseous atmosphere rotating as a rigid body at a rate several times less than the Keplerian rate. This is an expected outcome of a low-accretion-rate stage when the flow efficiently exchanges angular momentum in latitudinal direction (otherwise, rotation is going to be differential) and slows down due to some kind of friction with the NS crust, presumably rotating even slower. Hence, our setup may correspond to an episode of rapidly growing mass accretion rate. In future simulations, we are planning to include a friction force and evolve the simulation toward a steady state.

7. CONCLUSIONS

This code presented in this work allows to address any possible two-dimensional hydrodynamical problems on a spherical surface up to reasonably high Mach numbers of $\sim 5-10$. Both steady-state and dynamical problems set on a sphere show good consistency with the analytical solutions as well as internal consistency between different resolutions and grid types.

The capabilities of the code allow us for the first time to simulate the dynamics of an accretion spreading layer in the realistic parameter range appropriate for a real weakly magnetized accreting NS. Applying the code to the spreading layer problem reveals some interesting properties of this flow. In particular, its early evolution is dominated by the heating of the equatorial regions, that leads to Rayleigh-Taylor instability and efficient mixing of the accreting matter in latitudinal direction. As a by-product, it produces a global twoarmed 'tennis-ball' pattern that causes flux variations for an observer at a large inclination. The pattern retains its shape and changes its rotation velocity at a much slower rate than the spin-up of the spreading layer in general. Other observed oscillation modes may be connected to pressure modes propagating in longitudinal direction or to inertial Rossby modes, presumably with the azimuthal numbers m = 0 and 2.

Somehow, the development of the instabilities and rotating patterns leads to a considerable asymmetry between the two hemispheres. In terms of the average rotation velocities, the asymmetry reaches about 20% by the end of the simulation run, and is likely to grow further. To answer the question if




Figure 12. Final state of accretion simulation with streamlines. First plot contains streamlines with rotation, the second is without average rotation.




Figure 13. Dynamic power spectra of the accretion simulation for different observers. Overplotted are the *z*-component of the average frequency, its first harmonic, and cyclone frequency.

this asymmetry is likely to survive in the steady-state regime, we are planning more simulations with a similar setup but longer duration. This will also allow to test the validity of steady-state spreading-layer models.

- This work was supported by a grant from the Simons Foun-
- ² dation (00001470, PA). The authors would like to thank
- Joonas Nättilä and Sergey Khaibrakhmanov for valuable dis-
- 4 cussions.

A. ANALYTIC SOLUTIONS FOR RIGID-BODY ROTATION

Hydrostatic balance in vector form on the surface of a sphere, assuming rigid-body rotation with angular frequency Ω

$$\nabla \Pi = -\frac{v^2}{R\sin\theta} \Sigma e_{\varpi},\tag{A1}$$

where θ is polar angle, e_{ϖ} is cylindrical radial unit vector (directed away from the axis of rotation), and $v = \Omega R \sin \theta$ is rotation velocity. Projecting the equation onto the polar angle unit vector e_{θ} yields

$$\partial_{\theta}\Pi = \Omega^2 R \Sigma \sin \theta \cos \theta. \tag{A2}$$

As there is no other equation to constrain the pressure and density profile, it is possible to set an arbitrary profile of density, pressure, or entropy. In particular, for Σ =const, pressure is found as

$$\Pi_{\Sigma=\text{const}} = \Pi_{\text{poles}} + \frac{\Omega^2 R^2 \Sigma}{2} \sin^2 \theta. \tag{A3}$$

If globally $\Pi = \Pi_0 (\Sigma/\Sigma_0)^{\Gamma}$, solving equation (A2) results in

$$\Sigma_{\text{isentropic}} = \Sigma_0 \left[1 + \frac{\Gamma - 1}{2} \mathcal{M}_0^2 \sin^2 \theta \right]^{\frac{1}{\Gamma - 1}},\tag{A4}$$

$$\Pi_{\text{isentropic}} = \Pi_0 \left[1 + \frac{\Gamma - 1}{2} \mathcal{M}_0^2 \sin^2 \theta \right]^{\frac{\Gamma}{\Gamma - 1}},\tag{A5}$$

where the index '0' corresponds to the poles, and $\mathcal{M}_0 = \frac{\Omega R}{a_0}$, $c_{s,0} = \sqrt{\frac{\Gamma \Pi_0}{\Sigma_0}}$. Pressure normalization may be expressed using other quantities as

$$\Pi_0 = \frac{\Sigma_0}{\Gamma \mathcal{M}_0^2} \Omega^2 R^2. \tag{A6}$$

The Mach number of the isentropic flow is

$$\mathcal{M}(\theta) = \mathcal{M}_0 \frac{\sin \theta}{\sqrt{1 + \frac{\Gamma - 1}{2} \mathcal{M}_0^2 \sin^2 \theta}}.$$
 (A7)

It is maximal at the equator, where

$$\mathcal{M}_{\text{eq}} = \frac{\mathcal{M}_0}{\sqrt{1 + \frac{\Gamma - 1}{2} \mathcal{M}_0^2 \sin^2 \theta}}.$$
 (A8)

The upper limit for the Mach number for a rigidly rotating isentropic solution is $\mathcal{M}_{eq, max} = \sqrt{2/(\Gamma - 1)} = \sqrt{6} \approx 2.5$.

Depending on the latitudinal entropy gradient, a rigidly rotating configuration may be stable or unstable to buoyancy modes (Chandrasekhar 1961). The derivation in Boussinesq approximation is given in App. B and implies that the constant-density case should show convective instability with the increment $\sim M\Omega/2$.

B. STABILITY OF RIGID-BODY ROTATION

Let us consider steady-state rigid-body rotation flow from App. A and perform its linear analysis in Boussinesq approximation (Chandrasekhar 1961, chapter II). The equation of state is considered adiabatic with the index of Γ , hence we can introduce entropy as

$$S = \ln \frac{\Pi}{\Sigma^{\Gamma}}.$$
 (B9)

The unperturbed solution is described by equation (A2). In the first order, we will need to linearize the continuity equation

$$\partial_t \Sigma + \nabla(\Sigma v) = 0, (B10)$$

Euler equation (2 tangential components)

$$\partial_t \mathbf{v} + (\mathbf{v}\nabla)\mathbf{v} = \mathbf{g} - \frac{1}{\Sigma}\nabla\Pi,$$
 (B11)

and equation of state

$$\partial_t S + (v\nabla)S = 0. ag{B12}$$

The background solution is determined only by the θ component of the Euler equation, as the flow is stationary and incompressible, and equations (B10) and (B12) are fulfilled independently of the pressure and density profile.

Let us consider linear perturbations of Π , Σ , and $v_{\theta,\,\varphi}$ in the co-rotating frame. For every quantity f, $\delta f \propto \exp\left(\mathrm{i}\left(\omega t - m\varphi\right)\right) f_{\theta}(\theta)$. As we consider the problem in the co-rotating frame, both velocity components are zero in zeroth order, and v_{θ} and v_{φ} will be used without deltas. Thermodynamical quantities Π and Σ without deltas in this section will refer to the background solution.

Linearized continuity equation

$$i\omega R \sin \theta \frac{\delta \Sigma}{\Sigma} - imv_{\varphi} + \partial_{\theta} (v_{\theta} \sin \theta) = 0.$$
 (B13)

Linearized entropy conservation equation

$$i\omega R \left(\frac{\delta \Pi}{\Pi} - \Gamma \frac{\delta \Sigma}{\Sigma} \right) + v_{\theta} \partial_{\theta} S = 0.$$
 (B14)

Linearized θ component of Euler

$$i\omega R v_{\theta} = \frac{\delta \Sigma}{\Sigma} \Omega^{2} R^{2} \sin \theta \cos \theta - \frac{1}{\Sigma} \partial_{\theta} (\delta \Pi). \tag{B15}$$

If we ignore pressure variations (that corresponds to the Boussinesq approximation, valid for the processes slow enough for pressure to be leveled by sonic waves), equations (B14) and (B15) may be combined to reproduce the local Brunt-Väisälä frequency N as

$$\frac{N^2}{\Omega^2} = -\partial_{\theta} S \sin \theta \cos \theta. \tag{B16}$$

The right-hand side of the equation is always negative if entropy increases towards the equator. In particular, for Σ =const,

$$\partial_{\theta} S = \frac{\partial_{\theta} \Pi}{\Pi} = \frac{\Omega^2 R^2 \Sigma}{\Pi} \sin \theta \cos \theta = \Gamma \mathcal{M}^2 \sin \theta \cos \theta \left[1 + \frac{\Gamma - 1}{2} \mathcal{M}^2 \sin^2 \theta \right]^{-\frac{\Gamma}{\Gamma - 1}}, \tag{B17}$$

where $\mathcal{M} = \sqrt{\Gamma \Pi(\theta = \pi/2)/\Sigma}$, hence

$$N = i\Omega \sqrt{\Gamma} \mathcal{M} \sin \theta \cos \theta \left[1 + \frac{\Gamma - 1}{2} \mathcal{M}^2 \sin^2 \theta \right]^{-\frac{\Gamma}{2(\Gamma - 1)}}.$$
 (B18)

The frequency is always imaginary, meaning convective stability on all the latitudes. The increment itself is, however, a strong function of latitude. For $\mathcal{M} \ll 1$, the maximal value is achieved at $\theta \simeq \pi/4$ and equals $\text{Im }\omega \simeq \sqrt{\Gamma}\mathcal{M}\Omega/2$. In the general case (for arbitrary \mathcal{M}), the maximal increment is at

$$\sin \theta = \frac{1}{\sqrt{1 + \frac{M^2}{4} + \sqrt{1 + \frac{\Gamma - 1}{2}M^2 + \frac{M^4}{16}}}},$$
(B19)

that implies

$$\operatorname{Im} \omega = \Omega \sqrt{\Gamma} \mathcal{M} \left[1 + \frac{M^2}{4} + D \right]^{\frac{2-\Gamma}{2(\Gamma-1)}} \left(1 + \frac{2\Gamma - 1}{4} \mathcal{M}^2 + D \right) \sqrt{\frac{\mathcal{M}^2}{4} + D}, \tag{B20}$$

where

$$D = \sqrt{1 + \frac{\Gamma - 1}{2} \mathcal{M}^2 + \frac{\mathcal{M}^4}{16}}.$$
 (B21)

For large Mach numbers, however, the estimate is useless, as one of the assumptions was the negligibility of pressure variations, that is questionable for a supersonic flow.

REFERENCES

Abolmasov, P., Nättilä, J., & Poutanen, J. 2020, A&A, 638, A142, doi: 10.1051/0004-6361/201936958 Abolmasov, P., & Poutanen, J. 2021, A&A, 647, A45, doi: 10.1051/0004-6361/202039485

- Bahramian, A., & Degenaar, N. 2023, in Handbook of X-ray and Gamma-ray Astrophysics, 120,
 - doi: 10.1007/978-981-16-4544-0_94-1
- Bransgrove, A., Levin, Y., & Beloborodov, A. 2018, MNRAS, 473, 2771, doi: 10.1093/mnras/stx2508
- Chandrasekhar, S. 1961, Hydrodynamic and hydromagnetic stability
- Chen, S., Lin, B., Li, Y., & Yan, C. 2020, SIAM Journal on Scientific Computing, 42, B921, doi: 10.1137/18M119032X
- Gall, J. 1885, Scottish Geographical Magazine, 1, 119
- Gilfanov, M., Revnivtsev, M., & Molkov, S. 2003, A&A, 410, 217, doi: 10.1051/0004-6361:20031141
- Gottlieb, D., & Shu, C.-W. 1997, SIAM Review, 39, 644, doi: 10.1137/S0036144596301390
- Hasinger, G., & van der Klis, M. 1989, A&A, 225, 79
- Ingram, A. R., & Motta, S. E. 2019, NewAR, 85, 101524, doi: 10.1016/j.newar.2020.101524
- Inogamov, N. A., & Sunyaev, R. A. 1999, Astronomy Letters, 25, 269, doi: 10.48550/arXiv.astro-ph/9904333
- —. 2010, Astronomy Letters, 36, 848, doi: 10.1134/S1063773710120029
- Jakob-Chien, R., Hack, J. J., & Williamson, D. L. 1995, Journal of Computational Physics, 119, 164, doi: 10.1006/jcph.1995.1125
- Kluzniak, W., Michelson, P., & Wagoner, R. V. 1990, ApJ, 358, 538, doi: 10.1086/169006
- Landau, L. D., & Lifshitz, E. M. 1987, Fluid Mechanics Lomb, N. R. 1976, Ap&SS, 39, 447, doi: 10.1007/BF00648343

- Nättilä, J., Cho, J. Y. K., Skinner, J. W., Most, E. R., & Ripperda, B. 2024, ApJ, 971, 37, doi: 10.3847/1538-4357/ad54c2
- Papaloizou, J. C. B., & Stanley, G. Q. G. 1986, MNRAS, 220, 593, doi: 10.1093/mnras/220.3.593
- Payne, D. J. B., & Melatos, A. 2004, MNRAS, 351, 569, doi: 10.1111/j.1365-2966.2004.07798.x
- Scargle, J. D. 1982, ApJ, 263, 835, doi: 10.1086/160554
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337
- —. 1988, Advances in Space Research, 8, 135, doi: 10.1016/0273-1177(88)90396-1
- Shu, C.-W. 2020, Acta Numerica, 29, 701–762, doi: 10.1017/S0962492920000057
- Sieger, D. 2021, Generating Meshes of a Sphere. https: //danielsieger.com/blog/2021/03/27/generating-spheres.html
- Suleimanov, V., & Poutanen, J. 2006, MNRAS, 369, 2036, doi: 10.1111/j.1365-2966.2006.10454.x
- Touze, C. L., Murrone, A., & Guillard, H. 2015, Journal of Computational Physics, 284, 389, doi: 10.1016/j.jcp.2014.12.032
- Ursini, F., Gnarini, A., Capitanio, F., et al. 2024, Galaxies, 12, 43, doi: 10.3390/galaxies12040043
- van der Klis, M. 2000, ARA&A, 38, 717, doi: 10.1146/annurev.astro.38.1.717
- —. 2001, ApJ, 561, 943, doi: 10.1086/323378
- van Leer, B. 1979, Journal of Computational Physics, 32, 101, doi: https://doi.org/10.1016/0021-9991(79)90145-1
- Watts, A. L., Andersson, N., Beyer, H., & Schutz, B. F. 2003, MNRAS, 342, 1156, doi: 10.1046/j.1365-8711.2003.06612.x