

Transversal Structures in Graph Systems: A Survey

Wanting Sun *

Guanghai Wang †

Lan Wei ‡

Abstract

Given a system $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ of graphs/digraphs/hypergraphs on the common vertex set V of size n , an m -edge graph/digraph/hypergraph H on V is *transversal* in \mathcal{G} if there exists a bijection $\phi : E(H) \rightarrow [m]$ such that $e \in E(G_{\phi(e)})$ for all $e \in E(H)$. In this survey, we consider extremal problems for transversal structures in graph systems. More precisely, we summarize some sufficient conditions that ensure the existence of transversal structures in graph/digraph/hypergraph systems, which generalize several classical theorems in extremal graph theory to transversal version. We also include a number of conjectures and open problems.

1 Introduction

How global parameters of a graph, such as its degree or chromatic number, can influence its local substructures? Is there some sufficiently large average degree or chromatic number that forces a given substructure to appear? Questions of this type are one of the most natural problems in graph theory, and there are many profound and interesting results. These are collectively referred to as *extremal graph theory*. It is a branch of combinatorial mathematics that studies the maximal or minimal properties of graphs under certain conditions.

Many important problems in extremal graph theory can be framed as subgraph containment problems, which ask for conditions on a graph G that ensure it contains a copy of a general graph H . For example, Mantel's theorem states that every n -vertex graph with more than $\lfloor n^2/4 \rfloor$ edges contains a triangle; Turán [132] generalized it by determining the maximum number of edges in a graph that forbids K_r , where K_r denotes a complete graph on r vertices; Dirac [39] proved that every n -vertex graph with minimum degree at least $n/2$ contains a Hamilton cycle. Such a graph is called a *Dirac graph*. For more results, the reader is referred to the book of Bollobás [15].

In this paper, we survey some recent developments on transversal problems in graph systems. The notion of transversal appears in diverse branches of mathematics. A *transversal* X over a system $\mathcal{F} = \{F_1, \dots, F_m\}$ of objects refers to an object that intersects each F_i exactly once. Here, the object can be any mathematical object, such as sets, spaces, set systems, and matroids, etc. Several classical theorems, which guarantee the existence of certain objects, have been reiterated in the context of transversals, demonstrating the possibility and impossibility of obtaining such an object as a transversal over a certain system. To name a few, transversals are extensively studied for Carathéodory's theorem [14], Helly's theorem [75], Erdős–Ko–Rado theorem [7], and Rota's basis conjecture [71, 118], etc.

Building on this trend, there have been numerous studies dedicated to generalizing classical results in extremal graph theory to transversal version. In addition to being interesting in their own right, they often provide a strengthening of the original results, which also reflects the robustness of graph properties. A transversal in a graph system is also referred to as a “rainbow subgraph” in much literature, and it was formally introduced in the recent work of Joos and Kim [74]. For a positive integer n , we write $[n] := \{1, 2, \dots, n\}$.

*Data Science Institute, Shandong University, Jinan, 250100, China. Email: wtsun@sdu.edu.cn. Supported by the China Postdoctoral Science Foundation (2023M742092).


†School of Mathematics, Shandong University, Jinan, 250100, China. Email: ghwang@sdu.edu.cn. Supported by the National Key Research and Development Program (2023YFA1009603) and the Natural Science Foundation of China (12231018).

‡School of Mathematics, Shandong University, Jinan, 250100, China. Email: lanwei@mail.sdu.edu.cn.


Definition 1.1. For a given system $\mathcal{G} = \{G_1, \dots, G_m\}$ of graphs/digraphs/hypergraphs with the same vertex set V , an m -edge graph/digraph/hypergraph H on the vertex set V is a \mathcal{G} -*transversal* (or a transversal inside \mathcal{G}) if there exists a bijection $\phi : E(H) \rightarrow [m]$ such that $e \in E(G_{\phi(e)})$ for all $e \in E(H)$.

By interpreting each G_i as the set of edges colored with the color i , the function ϕ in the above definition is referred to as a *rainbow coloring* and H is a *rainbow subgraph* inside \mathcal{G} . We shall emphasize that in what follows, when we say that \mathcal{G} is a graph system with order n , then all graphs in \mathcal{G} are not necessarily distinct and they have the same vertex set V of size n . Many classical results in extremal graph theory have been extended to transversal settings, exhibiting interesting phenomena.

Perhaps the most famous transversals are those related to Latin squares, which were studied by Euler. In 1782, Euler [48] considered a Latin square of order n , which is an $n \times n$ array filled with symbols $1, \dots, n$, where each symbol appears exactly once in each row and column. A *partial transversal* of a Latin square is a set of cells such that every pair of cells shares no row, column or symbol, in particular a *transversal* is a partial transversal with n cells. The famous Ryser–Brualdi–Stein conjecture [124, 22, 126] is one of the most important problems on transversals in Latin squares.



(a) Latin square of order 4.



(b) Edge-colored bipartite graph.

Conjecture 1.2 (Ryser–Brualdi–Stein conjecture, [124, 22, 126]). *Every Latin square of order n has a partial transversal with $n - 1$ cells, and a transversal with n cells if n is odd.*

In 2023, Montgomery [113] introduced the first techniques to identify and exploit the possible algebraic properties behind the entries in a Latin square, and proved the following.

Theorem 1.3 ([113]). *There is some $n_0 \in \mathbb{N}$ such that every Latin square of order $n \geq n_0$ contains a partial transversal with $n - 1$ cells.*

For more relevant results of transversals in Latin squares, see the recent survey by Montgomery [114], which collected results from the past decade.

Considering the rows and columns of the Latin square as a complete bipartite graph $K_{n,n}$, where each symbol represents a color and each cell represents a colored edge, the Latin square naturally corresponds to a properly edge-colored $K_{n,n}$. The edge-colored $K_{n,n}$ can be viewed as a set of graphs $\mathcal{G} = \{G_1, \dots, G_n\}$, where each G_i consists of edges with color i , then a rainbow matching of \mathcal{G} corresponds to a transversal of the Latin square. We refer the reader to see the nice survey [133]. These studies have inspired research on the existence of transversal structures in a system of graphs. The central question in the study of transversals can be generalized and depicted as follows.

Problem 1.4 ([74]). Let H be a graph/digraph/hypergraph with m edges, $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ be a system of graphs/digraphs/hypergraphs with order n . Which property imposed on \mathcal{G} will ensure a transversal copy of H ?

Since we do not require the objects in the family \mathcal{G} to be distinct, a necessary condition for a positive answer to the above problem is that every n -vertex graph/digraph/hypergraph satisfying the property contains a copy of H . In the following of this survey, it will be shown by some examples that this necessary condition is not sufficient in some cases.

In this paper, we aim to survey known results for transversal problems in graph/random graph/digraph/hypergraph systems. Throughout the survey, we include a number of conjectures and open problems. It is of course impossible to cover all the known results for graph systems in a single survey; thus, the choice of results we present is inevitably subjective. Nevertheless, we hope to describe enough results from this fascinating area to engage many researchers in extremal and probabilistic combinatorics, and motivate further study of this subject.

2 Turán-type problem in graph systems

One of the central topics in extremal combinatorics is to determine the maximum number of edges of a graph on n vertices that does not contain a copy of a given graph H as a subgraph, such a maximum number is denoted by $\text{ex}(n, H)$. The most notable result is Turán's theorem [132], which states that $\text{ex}(n, K_{r+1}) = t_r(n)$, where $t_r(n)$ is the number of edges of a balanced complete r -partite graph on n vertices (denoted by $T_r(n)$). Erdős, Stone and Simonovits [46, 47] found a surprising connection between $\text{ex}(n, H)$ and the chromatic number $\chi(H)$ of H .

In addition to the Turán problem itself, there are also many variants involving colorings. For example, Diwan and Mubayi [40] considered a generalization of Turán's theorem in a graph system containing two graphs. It is of course possible to consider similar question with more than two graphs. In this section, we survey results on the Turán-type problems in graph systems. In other words, we are to determine the "largest" n -vertex graph systems that do not contain a copy of H as a rainbow subgraph.

Define the k -rainbow extremal number $\text{ex}_k^*(n, H)$ of a graph H to be the maximum $\min_{i \in [k]} |E(G_i)|$ among all rainbow H -free graph systems $\mathcal{G} = \{G_1, \dots, G_k\}$ with order n . The simplest case here is a rainbow triangle. Diwan and Mubayi [40] proposed a problem: determine the value of $\text{ex}_3^*(n, K_3)$. Magnant [111] answered this problem under the assumption that the multigraphs (i.e., graphs formed by taking the union of the edge sets $E(G_i)$) in question contain at least one edge between every pair of vertices. In 2020, Aharoni, DeVos, de la Maza, Montejano and Šámal [3] solved the above problem completely.

Theorem 2.1 ([3]). $\text{ex}_3^*(n, K_3) = \lfloor \frac{26-2\sqrt{7}}{81} \rfloor n^2$.

Interestingly, the constant $\frac{26-2\sqrt{7}}{81}$ is an irrational number and larger than $\frac{1}{4}$. Therefore, compared to Mantel's theorem, the transversal triangle problem requires a stronger condition. This signifies the difference between the transversal setting and the classical setting, and renders the rainbow extremal number more intriguing. Aharoni, DeVos, de la Maza, Montejano and Šámal [3] proposed a potential interesting direction to proceed, that is,

Problem 2.2 ([3]). For what real numbers $\alpha_1, \alpha_2, \alpha_3 > 0$, every graph system $\{G_1, G_2, G_3\}$ satisfying $|E(G_i)| > \alpha_i n^2$ must have a transversal triangle?

Falgas-Ravry, Markström and Rätty [50, Theorem 1.13] solved this problem when n is sufficiently large. Recall that Turán's theorem [132] generalizes Mantel's theorem by determining the maximum number of edges in n -vertex K_r -free ($r \geq 3$) graphs. Analogously, one may consider the following.

Problem 2.3 ([3]). Determine the value of $\text{ex}_{(2)}^*(n, K_r)$ for $r \geq 4$.

In general cases, the problem becomes more complicated and there are only few known results related to the k -rainbow extremal number $\text{ex}_k^*(n, H)$. As a corollary of Keevash, Saks, Sudakov and Verstraëte [83], we have $\text{ex}_k^*(n, K_3) \leq n^2/4 + o(n^2)$ when $k \geq 4$. Frankl [55] gave a new proof

of this result. Babiński and Grzesik [12] determined the value of $\text{ex}_k^*(n, P_4)$ up to additive $o(n^2)$ error term for every $k \geq 3$, where P_4 denotes the path on 4 vertices.

The k -rainbow Turán density of a graph H is defined as follows:

$$\pi_k^*(H) = \lim_{n \rightarrow \infty} \frac{\text{ex}_k^*(n, H)}{\binom{n}{2}}.$$

Im, Kim, Lee and Seo [72] showed that the limit exists for every graph H . They [72] conjectured that the rainbow Turán density is always an algebraic number, and they proved the conjecture for all trees. Furthermore, they [72] gave an answer to an analogue of Problem 2.2 for trees.

Theorem 2.4 ([72]). *Let $\alpha_1, \dots, \alpha_k$ be nonnegative reals such that $\sum_{i=1}^k (1 - \sqrt{\alpha_i}) < 1$. For every $\epsilon > 0$, there exists an integer n_0 such that for every $n \geq n_0$ and every k -edge tree T , every graph system $\mathcal{G} = \{G_1, \dots, G_k\}$ of order n with $|E(G_i)| \geq (\alpha_i + \epsilon) \binom{n}{2}$ contains a transversal copy of T .*

Except the rainbow Turán number of a graph, depending on the contexts, different measures for the “size” of graph systems have been introduced, for example, the sum $\sum_i |E(G_i)|$ [83, 24], the product $\prod_i |E(G_i)|$ [55, 68] and other measures.

The k -color Turán number, denoted by $\text{ex}_k(n, H)$, is the maximum value of $\sum_{i \in [k]} |E(G_i)|$ among all graph systems $\mathcal{G} = \{G_1, \dots, G_k\}$ of order n and without rainbow copy of H . We call such a graph system \mathcal{G} satisfying $\sum_{i \in [k]} |E(G_i)| = \text{ex}_k(n, H)$ the k -color extremal graph system of H . To determine $\text{ex}_k(n, H)$, there are two natural lower bounds that arise from the following constructions. One is that each G_i is the n -vertex H -free graph with $\text{ex}(n, H)$ edges; the other one is that there are $|E(H)| - 1$ copies of the complete graph K_n and all the other graphs are empty.

A graph is called r -color-critical if it has chromatic number r , and it has an edge whose removal reduces the chromatic number to $r - 1$. In 2004, Keevash, Saks, Sudakov and Verstraëte [83] proposed the following conjecture.

Conjecture 2.5 ([83]). *Suppose $r \geq 3$ and $k \geq h$. Let H be an r -color-critical graph with h edges. Then, there exists an $n_0 = n_0(H) > 0$ such that for all $n \geq n_0$, the k -color extremal graph system of H either consists of exactly $h - 1$ nonempty graphs where each of them is a copy of K_n or consists of k graphs where all of them are identical copies of $T_{r-1}(n)$. In particular,*

$$\text{ex}_k(n, H) = \begin{cases} (h - 1) \binom{n}{2}, & \text{if } h \leq k < \frac{r-1}{r-2}(h - 1), \\ k \cdot t_{r-1}(n), & \text{if } k \geq \frac{r-1}{r-2}(h - 1). \end{cases}$$

They provided evidence for this conjecture by proving it holds when H is a complete graph or $r = 3$. Recently, Chakraborti, Kim, Lee, Liu and Seo [24] showed that the conjecture holds for every 4-color-critical graph H and almost all r -color-critical graphs H with $r \geq 5$. Furthermore, they [24] proved that if H is not r -color-critical for some $r \geq 3$ and n is sufficiently large, then neither of the constructions in Conjecture 2.5 is extremal. Li, Ma and Zheng [103] proved Conjecture 2.5 holds when $k \geq 2(r - 1)(h - 1)/r$, significantly improving earlier results. Furthermore, their proof combined the stability argument of Chakraborti, Kim, Lee, Liu and Seo [24] with a novel graph packing technique for embedding multigraphs.

Theorem 2.6 ([103]). *Let $r \geq 5$ and H be an r -color-critical graph with h edges. If n is sufficiently large and $k \geq \frac{2(r-1)(h-1)}{r}$, then $\text{ex}_k(n, H) = k \cdot t_{r-1}(n)$, and the unique n -vertex k -color extremal graph system of H consists of k colors all of which are identical copies of $T_{r-1}(n)$.*

There is also some research focusing on maximizing the product $\prod_i |E(G_i)|$ of the number of edges of a graph system without a rainbow copy of H . Frankl [55] conjectured that a graph system $\{G_1, G_2, G_3\}$ on a common vertex set of order n with $\prod_{i \in [3]} |E(G_i)| > \lfloor n^2/4 \rfloor^3$ contains a transversal triangle. This conjecture was disproved by Frankl, Györi, He, Lv, Salia, Tompkins, Varga and Zhu [68], and they further conjectured that:

Conjecture 2.7 ([68]). Let $\mathcal{G} = \{G_1, G_2, G_3\}$ be a graph system of order n . If \mathcal{G} contains no transversal triangle, then

$$\prod_{i \in [3]} |E(G_i)| \leq \gamma n^6 (1 + o(1)),$$

where γ is the maximum of $\frac{x^2}{2} \left(\frac{x^2}{2} + \frac{(1-x)^2}{2} \right) \left(x(1-x) + \frac{(1-x)^2}{2} \right)$ on $[0, 1]$.

In fact, this conjecture is a direct consequence of Problem 2.2, which has been solved by Falgas-Ravry, Markström and Rätty [50] when n is sufficiently large.

3 Dirac-type problem in graph systems

Let $G = (V, E)$ be a graph. For a vertex $v \in V$, denote by $N_G(v)$ the neighborhood of v and let $d_G(v) = |N_G(v)|$ be the degree of v . The subscripts are omitted when G is clear from the context. Let $\delta(G)$ and $\Delta(G)$ be the minimum degree and maximum degree of G respectively. Let $\mathcal{G} = \{G_1, G_2, \dots, G_m\}$ be a graph system. Denote by $\delta(\mathcal{G}) = \min\{\delta(G_i) : i \in [m]\}$ the minimum degree of \mathcal{G} .

In this section, we investigate the minimum degree condition that ensure the existence of transversal spanning structures in graph systems. When one is interested in embedding a spanning graph, it makes sense to consider minimum degree conditions rather than size conditions, as, for example, a graph could be very dense but contain an isolated vertex, and therefore not contain a copy of any given spanning graph with minimum degree at least 1.

Given the difficulty of obtaining exact results in (rainbow) graph embedding problems, many results have focused on determining the minimum d such that for all $\epsilon > 0$ and sufficiently large integer n , each graph (system) of order n with minimum degree at least $(d + \epsilon)n$ contains a (transversal) copy of H . Note that when we consider this problem in graph systems, d is at least the value required to guarantee a copy of H in a single graph, since all graphs in the system could be identical.

Definition 3.1. Let H be a graph with m edges. Denote by δ_H (if it exists) the smallest real number δ such that for all $\alpha > 0$ and sufficiently large n , every n -vertex graph G with $\delta(G) \geq (\delta + \alpha)n$ contains a copy of H .

Denote by δ_H^{rb} (if it exists) the smallest real number δ such that for all $\alpha > 0$ and sufficiently large n , every graph system $\mathcal{G} = \{G_1, \dots, G_m\}$ of order n with $\delta(\mathcal{G}) \geq (\delta + \alpha)n$ contains a transversal copy of H .

For a given graph H , if the above two values exist, then it is clear that $\delta_H^{rb} \geq \delta_H$. Indeed, if G is H -free, then the graph system \mathcal{G} consisting of $|E(H)|$ copies of G does not contain a transversal copy of H . However, Montgomery, Müyesser and Pehova [115] showed that δ_H^{rb} can be much larger than δ_H : taking H to be the disjoint union of copies of $K_{2,3} \cup C_4$, then $\delta_H^{rb} \geq 1/2$, whereas Kühn and Osthus [95] showed that $\delta_H \leq 4/9$.

On the other hand, there are many natural instances with $\delta_H^{rb} = \delta_H$, for example, Hamilton cycle and perfect matching (see [74]). When this equality holds, we always say that the corresponding H is *color-blind*.

3.1 Hamilton path and Hamilton cycle

A *Hamilton path (cycle)* in a graph is a path (cycle) that visits every vertex exactly once. Hamiltonicity is one of the most well-studied properties in graph theory. Dirac [39] showed that every n -vertex graph with minimum degree at least $n/2$ contains a Hamilton cycle. Later on, Ore [117] relaxed this condition and proved that every n -vertex graph G with $\sigma_2(G) \geq n$ contains a Hamilton cycle, where $\sigma_2(G) = \min\{d(x) + d(y) : xy \notin E(G)\}$. Moon and Moser [116] established the bipartite analogues of Dirac's theorem.

In 2020, Aharoni [3] conjectured that Dirac's theorem can be extended to a transversal version. Cheng, Wang and Zhao [33] proved an asymptotic version of this conjecture; Joos and Kim [74] confirmed the conjecture completely.

Theorem 3.2 ([74]). *Let $n \in \mathbb{N}$ and $n \geq 3$. Suppose that $\mathcal{G} = \{G_1, \dots, G_n\}$ is a graph system of order n and $\delta(\mathcal{G}) \geq \frac{n}{2}$. Then \mathcal{G} contains a transversal Hamilton cycle.*

Bradshaw [20] gave a degree condition to guarantee the existence of transversal Hamilton cycles in bipartite graph systems, which generalizes the result of Moon and Moser [116].

The *distance* of two distinct vertices in a graph is the length of a shortest path connecting them. The k -th power G^k of a graph G is the graph obtained from G by joining every pair of vertices with distance at most k in G . Pósa (for $r = 3$, [43]) and Seymour (for $r > 3$, [125]) conjectured that for all $r \geq 3$ and all $n \geq r$, every n -vertex graph G with $\delta(G) \geq (r-1)n/r$ must contain the $(r-1)$ -th power of a Hamilton cycle. This conjecture has been confirmed by Komlós, Sárközy, and Szemerédi [90, 92]. Recently, Gupta, Hamann, Müyesser, Parczyk and Sgueglia [62] determined the asymptotically tight bound on $\delta(\mathcal{G})$ for the existence of transversal powers of Hamilton cycles, and they proposed a problem for the exact version of transversal Pósa–Seymour conjecture.

Problem 3.3 ([62]). *Let $\mathcal{G} = \{G_1, \dots, G_{rn}\}$ be a graph system of order n . If $\delta(\mathcal{G}) \geq (1 - \frac{1}{r})n$, is already sufficient for a transversal copy of the r -th power of a Hamilton cycle?*

Bowtell, Morris, Pehova and Staden [19] proposed a further generalization of the transversal problem, where one seeks a copy of H with a given coloring.

Problem 3.4 ([19]). *Given a graph system $\mathcal{G} = \{G_1, \dots, G_m\}$ of order n , what conditions imposed on \mathcal{G} will guarantee that for every coloring $\chi : E(H) \rightarrow \{1, \dots, m\}$, there is a copy of H such that the image of e is an edge of $G_{\chi(e)}$ for all $e \in E(H)$?*

When H is a Hamilton cycle, Bowtell, Morris, Pehova and Staden [19] gave an asymptotically optimal minimum degree condition for Problem 3.4.

Theorem 3.5 ([19]). *For every $\alpha > 0$ there exists n_0 such that for every $n \geq n_0$ the following holds. Let $m \in \mathbb{N}$ and let $\mathcal{G} = \{G_1, \dots, G_m\}$ be a graph system of order n . If $\delta(\mathcal{G}) \geq (\frac{1}{2} + \alpha)n$, then for every $\chi : [n] \rightarrow [m]$, there is a Hamilton cycle C such that the image of e is an edge of $G_{\chi(e)}$ for all $e \in E(C)$.*

Determining the exact minimum degree condition to guarantee all Hamilton cycle patterns remains an intriguing open problem. Bowtell, Morris, Pehova and Staden [19] provided a construction demonstrating that $n/2 + 1$ is a necessary lower bound and is conjectured as an exact threshold. In light of this, it seems more pertinent to either obtain results that improve this $o(n)$ term to an explicit and relatively small constant, or find counterexamples to this being possible.

It is known that one can measure the robustness of Dirac graphs with respect to Hamiltonicity by counting the number of distinct Hamilton cycles that a Dirac graph must contain. For more details about the robustness, we refer the reader to see the nice survey of Sudakov [127]. This method can be extended to measure the robustness of Dirac graph systems.

It follows from Theorem 3.5 that there exist at least $n! \sim \sqrt{2\pi n}(n/e)^n$ different transversal Hamilton cycles in a graph system consisting of n asymptotically Dirac graphs. Bradshaw, Halasz, and Stacho [21] showed that given a system of n Dirac graphs with order n , there are at least $(cn/e)^{cn}$ different transversal Hamilton cycles for some constant $c \geq 1/68$. Anastos and Chakraborti [11] improved the result of [21] by showing that there exists a constant $C > 0$ such that there are $(Cn)^{2n}$ different transversal Hamilton cycles in a graph system consisting of n Dirac graphs, which is tight up to the choice of C . They [11] also showed that every graph system consisting of n Dirac graphs with order n spans $n/2$ pairwise edge-disjoint transversal Hamilton cycles, and this is best possible.

Cheng and Staden [31] established a stability result for transversal Hamilton cycles. Given integers $a, b \geq 0$, let \mathcal{H}_a^b be a graph system obtained by taking a copies of $K_{\frac{n}{2}} \cup K_{\frac{n}{2}}$ and b copies of $K_{\frac{n}{2}, \frac{n}{2}}$ where they are defined on the same equitable partition $A \cup B$. For two vertex subsets $X, Y \subseteq V$, let $G[X]$ denote the subgraph of G induced by X , let $G[X, Y]$ be the subgraph of G with vertex set $X \cup Y$ and edge set $\{xy : x \in X, y \in Y \text{ and } xy \in E(G)\}$. We say that a graph

system \mathcal{J} of order n is *half-split* if there is $A \subseteq V$ with $|A| = \lfloor n/2 \rfloor + 1$ such that $J[A] = \emptyset$ and $J[A, V \setminus A]$ is complete for all $J \in \mathcal{J}$. Given $\kappa > 0$, we say that two graph systems \mathcal{G} and \mathcal{G}' on a common vertex set of size n are κ -close if \mathcal{G}' can be obtained from \mathcal{G} by adding or deleting at most κn^3 edges.

Theorem 3.6 ([31]). *For all $\kappa > 0$, there exist $\alpha > 0$ and n_0 such that the following holds for all integers $n \geq n_0$. Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a graph system of order n and $\delta(\mathcal{G}) \geq \frac{1-\alpha}{2}n$. If \mathcal{G} contains no transversal Hamilton cycles, then either \mathcal{G} is κ -close to \mathcal{H}_{n-t}^t for some odd integer $t \in [n]$, or \mathcal{G} is κ -close to a half-split graph system.*

The proof of Theorem 3.6 was a combination of their newly developed transversal regularity-blow-up method [30] (see Theorem 4.4 in the next section) and the absorption method. As a continuation of the transversal version of Dirac's theorem (Theorem 3.2) and the stability result for transversal Hamilton cycles (Theorem 3.6), Cheng, Sun, Wang and Wei [32] characterized all graph systems with order n and minimum degree at least $n/2 - 1$, each of which does not contain transversal Hamilton cycles.

Li, Li and Li [101] studied the existence of rainbow Hamilton paths in a graph system $\mathcal{G} = \{G_i : i \in [n]\}$ under Ore-type conditions. However, the edges of a rainbow Hamilton path only come from $n - 1$ graphs of \mathcal{G} . Thus, it is worth studying whether $n - 1$ graphs can guarantee the existence of a rainbow Hamilton path in a graph system $\mathcal{G} = \{G_i : i \in [n - 1]\}$. It is well-known that Dirac's theorem implies that every n -vertex graph with minimum degree at least $(n - 1)/2$ contains a Hamilton path. Cheng, Sun, Wang and Wei [32] proved a transversal version of this result.

Theorem 3.7 ([32]). *Let $n \in \mathbb{N}$ and $n \geq 3$. Suppose $\mathcal{G} = \{G_1, \dots, G_{n-1}\}$ is a graph system of order n . If $\delta(\mathcal{G}) \geq \frac{n-1}{2}$, then \mathcal{G} contains a transversal Hamilton path.*

Similar to the case of Hamilton cycles, Cheng and Staden [31] established a stability result for transversal Hamilton paths; Cheng, Sun, Wang and Wei [32] characterized all graph systems with order n and minimum degree at least $(n - 3)/2$, each of which does not contain transversal Hamilton paths.

In 1971, Bondy [16] posed a meta-conjecture which states that almost all nontrivial sufficient conditions on a graph which implies that the graph is Hamiltonian also implies that the graph is pancyclic, or there might be a simple family of exceptional graphs. Some concepts similar to pancyclic have also attracted extensive attention, such as, vertex-pancyclic and panconnected. Here, we only illustrate those concepts in graph systems. Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a system of graphs on a common vertex set V of order n . We say

- \mathcal{G} is *transversal pancyclic* if there is a rainbow cycle of length ℓ in \mathcal{G} for each integer $\ell \in [3, n]$;
- \mathcal{G} is *transversal vertex-pancyclic* if each vertex of V is contained in a rainbow cycle of length ℓ for all $\ell \in [3, n]$;
- \mathcal{G} is *rainbow panconnected* if for any two vertices $u, v \in V$, \mathcal{G} contains a rainbow path of length ℓ joining u and v for all $\ell \in [d_{\mathcal{G}}(u, v) + 1, n]$, where $d_{\mathcal{G}}(x, y)$ is the length of the shortest rainbow path (if it exists) connecting u and v in \mathcal{G} .

Li, Li and Li [101, 102] obtained Dirac-type conditions for transversal pancyclicity, transversal vertex-pancyclicity and rainbow panconnectedness respectively. The analogue results in bipartite graph systems were obtained by Bradshaw [20] and Hu, Li, Li and Xu [69].

In addition to the Dirac-type condition, there are many other sufficient conditions to guarantee the Hamiltonicity of a graph, such as Ore's condition [117] and Pósa's condition [120]. Bradshaw [20] and Li, Li and Li [101] proposed that it is also interesting to consider those conditions in graph systems. We list one of them here.

Question 3.8 ([20, 101]). *Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a graph system of order n . If $\sigma_2(G_i) \geq n$ for all $i \in [n]$, does \mathcal{G} contain a transversal Hamilton cycle?*

3.2 F -factor

Given two graphs F and G , an F -tiling is a set of vertex-disjoint copies of F in G . A *perfect F -tiling* (or an F -factor) in G is an F -tiling covering all vertices of G . Characterizing sufficient conditions for the existence of F -factors is one of the central areas of extremal graph theory. The most celebrated one is the Hajnal–Szemerédi theorem.

Theorem 3.9 (Hajnal–Szemerédi [63], Corrádi–Hajnal [34] for $k = 3$). *Every n -vertex graph G with $n \in k\mathbb{N}$ and $\delta(G) \geq (1 - \frac{1}{k})n$ has a K_k -factor. Moreover, the minimum degree condition is sharp.*

For any r -vertex graph F , Kühn and Osthus [95] determined the smallest value of δ such that there is an $n_0 \in \mathbb{N}$ and a constant $c = c(F)$ such that any graph with $n \geq n_0$ vertices and minimum degree at least $\delta n + c$ contains an F -factor if $r|n$. For more results, please see the survey of Kühn and Osthus [94].

Now, we survey some results on minimum degree conditions for the existence of transversal F -factors in graph systems. Joos and Kim [74] established an exact minimum degree condition to guarantee the existence transversal perfect matching (i.e., K_2 -factor) in graph systems.

Theorem 3.10 ([74]). *Let $n \in \mathbb{N}$ and $n \geq 2$ even. Suppose $\mathcal{G} = \{G_1, \dots, G_{\frac{n}{2}}\}$ is a graph system of order n and $\delta(\mathcal{G}) \geq \frac{n}{2}$. Then \mathcal{G} contains a transversal perfect matching.*

Bradshaw [20] proved the analogue result in bipartite graph systems, which is equivalent to a special case of a matching theorem for r -partite r -uniform hypergraphs obtained by Aharoni, Georgakopoulos and Sprüssel [4], for the case $r = 3$.

Cheng, Han, Wang and Wang [27] gave an asymptotical version of the transversal Hajnal–Szemerédi theorem. Furthermore, they proposed a conjecture for the exact version of transversal Hajnal–Szemerédi theorem.

Conjecture 3.11 ([27]). *Let $\mathcal{G} = \{G_1, \dots, G_{\frac{n}{k} \binom{k}{2}}\}$ be a graph system of order n . If $\delta(\mathcal{G}) \geq (1 - \frac{1}{k})n$, then \mathcal{G} contains transversal K_k -factor.*

Montgomery, Müyesser and Pehova [115] showed that, in most cases, the same minimum degree bound is asymptotically sufficient for the existence of transversal F -factors, which generalized the Kühn–Osthus theorem [95] on factors.

Theorem 3.12 ([115]). *Let $\varepsilon > 0$ and let F be a graph on r vertices with t edges. If $\delta_{F\text{-factor}} \geq \frac{1}{2}$ or F has a bridge, then let $\delta_F^T = \delta_{F\text{-factor}}$, and otherwise let $\delta_F^T = \frac{1}{2}$. Then, there is some n_0 such that the following holds for all $n \geq n_0$.*

Assume that $\mathcal{G} = \{G_1, \dots, G_{tn}\}$ is a graph system of order rn and $\delta(\mathcal{G}) \geq (\delta_F^T + \varepsilon)rn$. Then, \mathcal{G} contains a transversal F -factor.

Suppose V_1, V_2, \dots, V_k are disjoint vertex sets each of order n , and G is a k -partite graph on vertex classes V_1, V_2, \dots, V_k . Denote the *partite minimum degree* of G by

$$\delta'(G) := \min_{i \in [k]} \min_{v \in V_i} \min_{j \in [k] \setminus \{i\}} |N(v) \cap V_j|.$$

Fischer [52] conjectured that the k -partite graph G contains a K_k -factor if $\delta'(G) \geq (1 - 1/k)n + 1$ (actually his original conjecture did not include the term $+1$, but this stronger conjecture is known to be false). An approximate version of this conjecture was demonstrated independently by Keevash and Mycroft [81], and by Lo and Markström [104]. Later, Keevash and Mycroft [82] showed that the conjecture holds for any sufficiently large graph (in fact, they obtained a more general result for r -partite graphs with $r \geq k$). Cheng, Han, Wang and Wang [27] extended the approximate result of Fischer’s conjecture to the transversal version.

Theorem 3.13 ([27]). *For every $\varepsilon > 0$ and integer k , there exists $n_0 \in \mathbb{N}$ such that the following holds for all integers $n \geq n_0$. If $\mathcal{G} = \{G_1, \dots, G_{n \binom{k}{2}}\}$ is a system of k -partite graphs with a common vertex partition V_1, \dots, V_k each of size n such that $\delta'(G_i) \geq (1 - \frac{1}{k} + \varepsilon)n$, then \mathcal{G} contains a transversal K_k -factor.*

3.3 Graphs with bounded bandwidth

Montgomery, Müyesser and Pehova [115] determined asymptotically tight bound on $\delta(\mathcal{G})$ for the existence of a transversal spanning tree with maximum degree at most $cn/\log n$, generalizing the tree-embedding theorem of Komlós, Sárközy and Szemerédi [93].

It is noteworthy that many graphs, including F -factors, powers of Hamilton cycles, and trees, exhibit a somewhat bounded maximum degree and low connectivity. This property of low connectivity can be captured by the following notion of bandwidth. A graph H has a *bandwidth* at most b if there exists an ordering x_1, \dots, x_n of $V(H)$ such that all edges $x_i x_j \in E(H)$ satisfies $|i - j| \leq b$. Böttcher, Schacht and Taraz [18] proved the celebrated bandwidth theorem, which determines the asymptotically optimal minimum degree condition for a graph to contain a spanning subgraph H with bounded maximum degree and low bandwidth.

Since the bandwidth theorem provides a common generalization of many classical results, it is natural to characterize such a generalization for the above transversal version results. Chakraborti, Im, Kim and Liu [23] proved that such a generalization holds by showing the following bandwidth theorem for graph transversals.

Theorem 3.14 ([23]). *For every $\epsilon > 0$ and positive integers Δ, k , there exist $\alpha > 0$ and $h_0 > 0$ satisfying the following for every $h \geq h_0$. Let H be an n -vertex graph with h edges and bandwidth at most αn such that $\Delta(H) \leq \Delta$ and $\chi(H) \leq k$. If $\mathcal{G} = \{G_1, \dots, G_h\}$ is a graph system of order n and $\delta(\mathcal{G}) \geq (1 - \frac{1}{k} + \epsilon)n$, then \mathcal{G} contains a transversal copy of H .*

This covers several of the above results, and is asymptotically best possible for many graphs H . Note that by taking each G_i in Theorem 3.14 to be a same graph, the bandwidth theorem of Böttcher, Schacht and Taraz [18] can be deduced immediately. Therefore, Theorem 3.14 generalizes and strengthens the original bandwidth theorem.

4 Transversal via regularity

The classical “regularity-blow-up method” has been used to prove many results concerning the embedding of a given spanning subgraph H in a large graph G . Such proofs are typically developed using two important tools: Szemerédi’s regularity lemma [128] and blow-up lemma of Komlós, Sárközy and Szemerédi [91]. Cheng and Staden [30] provided a tool needed to obtain transversal embeddings by applying the regularity-blow-up method. More precisely, they proved a transversal blow-up lemma that embeds separable graphs into graph systems.

Definition 4.1 (Regularity and superregularity). Suppose that $\mathcal{G} = \{G_c : c \in \mathcal{C}\}$ is a system of bipartite graphs on a common vertex partition $V_1 \cup V_2$ with color set \mathcal{C} . We say that

- \mathcal{G} is (ϵ, d) -regular if whenever $V'_i \subseteq V_i$ with $|V'_i| \geq \epsilon|V_i|$ for $i = 1, 2$ and $\mathcal{C}' \subseteq \mathcal{C}$ with $|\mathcal{C}'| \geq \epsilon|\mathcal{C}|$, we have

$$\left| \frac{\sum_{c \in \mathcal{C}'} |E(G_c[V'_1, V'_2])|}{|\mathcal{C}'||V'_1||V'_2|} - \frac{\sum_{c \in \mathcal{C}} |E(G_c)|}{|\mathcal{C}||V_1||V_2|} \right| < \epsilon$$

and $\sum_{c \in \mathcal{C}} |E(G_c)| \geq d|\mathcal{C}||V_1||V_2|$.

- \mathcal{G} is (ϵ, d) -superregular if it is (ϵ, d) -regular and $\sum_{c \in \mathcal{C}} d_{G_c}(x) \geq d|\mathcal{C}||V_{3-i}|$ for all $x \in V_i$ with $i \in [2]$, and $|E(G_c)| \geq d|V_1||V_2|$ for all $c \in \mathcal{C}$.

Note that if every G_c with $c \in \mathcal{C}$ is the same, then \mathcal{G} is (ϵ, d) -regular if and only if G_c is (ϵ, d) -regular; and \mathcal{G} is (ϵ, d) -superregular if and only if G_c is (ϵ, d) -superregular.

Theorem 4.2 (Regularity lemma for graph systems, [30]). *For all integers $L_0 \geq 1$ and every $\epsilon, \delta > 0$, there is an $n_0 = n_0(\epsilon, \delta, L_0)$ such that for every $d \in [0, 1)$ and every graph system $\mathcal{G} = \{G_c : c \in \mathcal{C}\}$ on vertex set V of size $n \geq n_0$ with $\delta n \leq |\mathcal{C}| \leq \frac{n}{\delta}$, there exists a partition of*

V into V_0, V_1, \dots, V_L , of \mathcal{C} into $\mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_M$ and a spanning subgraph G'_c of G_c for each $c \in \mathcal{C}$ such that the following properties hold:

- (i) $L_0 \leq L, M \leq n_0$ and $|V_0| + |\mathcal{C}_0| \leq \varepsilon n$;
- (ii) $|V_1| = \dots = |V_L| = |\mathcal{C}_1| = \dots = |\mathcal{C}_M|$;
- (iii) $\sum_{c \in \mathcal{C}} d_{G'_c}(v) > \sum_{c \in \mathcal{C}} d_{G_c}(v) - (\frac{3d}{8^2} + \varepsilon)n^2$ for all $v \in V$ and $|E(G'_c)| > |E(G_c)| - (\frac{3d}{8^2} + \varepsilon)n^2$ for all $c \in \mathcal{C}$;
- (iv) if, for $c \in \mathcal{C}$, the graph G'_c has an edge with both vertices in a single vertex cluster V_i for some $i \in [L]$, then $c \in \mathcal{C}_0$;
- (v) for all triples $(\{h, i\}, j) \in \binom{[L]}{2} \times [M]$, we have that either $G'_c[V_h, V_i] = \emptyset$ for all $c \in \mathcal{C}_j$, or $\mathcal{G}'_{hi,j} := \{G'_c[V_h, V_i] : c \in \mathcal{C}_j\}$ is (ε, d) -regular.

The sets V_i are called vertex clusters and the sets \mathcal{C}_j are called color clusters, while V_0 and \mathcal{C}_0 are the exceptional vertices and colors sets respectively.

Definition 4.3 (Reduced graph system). Given a graph system $\mathcal{G} = \{G_c : c \in \mathcal{C}\}$ on V and parameters $\varepsilon > 0$, $d \in [0, 1)$ and $L_0 \geq 1$, the *reduced graph system* $\mathcal{R} = \mathcal{R}(\varepsilon, d, L_0)$ is defined as follows. Apply Theorem 4.2 to \mathcal{G} with parameters ε, d, L_0 to obtain a graph system $\mathcal{G}' = \{G'_c : c \in \mathcal{C}\}$, a partition V_0, \dots, V_L of V and $\mathcal{C}_0, \dots, \mathcal{C}_M$ of \mathcal{C} , where V_0, \mathcal{C}_0 are the exceptional sets, V_1, \dots, V_L are the vertex clusters and $\mathcal{C}_1, \dots, \mathcal{C}_M$ are the color clusters. Then $\mathcal{R} = \{R_1, \dots, R_M\}$ is a graph system consisting of M graphs each on the same vertex set $[L]$, where $hi \in R_j$ if and only if $\mathcal{G}'_{hi,j}$ is (ε, d) -regular for each $(\{h, i\}, j) \in \binom{[L]}{2} \times [M]$.

An n -vertex graph H is μ -separable if there is $X \subseteq V(H)$ of size at most μn such that $H - X$ consists of components of size at most μn . With suitable small μ , separable graphs includes many natural graphs, for example, F -factors for a fixed F , 2-regular graphs, trees, powers of Hamilton cycles and graphs of small bandwidth.

Theorem 4.4 (Transversal blow-up lemma, [30]). For all $\nu, d, \Delta, r > 0$, where $r \geq 2$ is an integer, there exist $\varepsilon, \mu, \alpha > 0$ and $m_0 \in \mathbb{N}$ such that the following holds for all integers $m \geq m_0$. Assume $\mathcal{G} = \{G_i : i \in \mathcal{C}\}$ is a graph system with the following properties.

- There is a graph R with vertex set $[r]$ and a partition $\mathcal{C} = \cup_{e \in E(R)} \mathcal{C}_e$ where $|\mathcal{C}_e| \geq \delta m$ for all $e \in E(R)$;
- for all $ij \in E(R)$ and $c \in \mathcal{C}_{ij}$, G_c is bipartite with parts V_i, V_j , and V is a vertex set of size n with partition $V = V_1 \cup V_2 \cup \dots \cup V_r$, where $m \leq |V_i| \leq \frac{m}{8}$ for each $i \in [r]$;
- for all $ij \in E(R)$,
 - for all $V_h' \subseteq V_h$ ($h = i, j$) and $\mathcal{C}'_{ij} \subseteq \mathcal{C}_{ij}$ with $|V_h'| \geq \varepsilon |V_h|$ ($h = i, j$) and $|\mathcal{C}'_{ij}| \geq \varepsilon |\mathcal{C}_{ij}|$, we have that
$$\sum_{c \in \mathcal{C}'_{ij}} |E(G_c[V_i', V_j'])| \geq d |\mathcal{C}'_{ij}| |V_i'| |V_j'|;$$
 - for every $v \in V_h$ with $h = i, j$ we have $\sum_{c \in \mathcal{C}'_{ij}} d_{G_c}(v) \geq d |\mathcal{C}'_{ij}| m$, and for every $c \in \mathcal{C}_{ij}$, we have $|E(G_c)| \geq dm^2$.

Suppose that H is a graph with the following properties.

- $\Delta(H) \leq \Delta$;
- H is μ -separable;

- H has vertex partition $A_1 \cup A_2 \cup \dots \cup A_r$ such that $|A_i| = |V_i|$ for all $i \in [r]$, and for every $xy \in E(H)$ there is $ij \in E(R)$ such that $x \in A_i$ and $y \in A_j$;
- $|E(H[A_i, A_j])| = |\mathcal{C}_{ij}|$ for all $ij \in E(R)$;
- for $i \in [r]$, there is a set $U_i \subseteq A_i$ with $|U_i| \leq \alpha|A_i|$ and for each $x \in U_i$, there exists a target set $T_x \subseteq V_i$ with $|T_x| \geq \nu m$.

Then \mathcal{G} contains a transversal copy of H such that every $x \in U_i$ is embedded inside T_x for $i \in [r]$.

By utilizing the transversal blow-up lemma, Cheng and Staden [30] proved a stability result for transversal Hamilton cycles, and characterized a large class of spanning 3-uniform linear hypergraphs H such that any sufficiently large uniformly dense n -vertex 3-uniform hypergraph with minimum vertex degree $\Omega(n^2)$ contains H as a subhypergraph.

5 Random graph systems

In this section, we survey some results on the existence of a transversal structures in a system of random graphs in various settings. Let $0 \leq p \leq 1$ and $G(n, p)$ be the binomial *random graph*. For graphs $G = (V, E)$ and $G' = (V', E')$, we set $G \cap G' = (V \cap V', E \cap E')$. Given an n -vertex graph G , we call $G(p) := G \cap G(n, p)$ the *random subgraph* of G . In particular, $G(p) = G(n, p)$ when G is the complete graph K_n . We say that $G(p)$ has a graph property \mathcal{P} with high probability, or whp for brevity, if the probability that $G(p)$ has \mathcal{P} approaches 1 as n tends to infinity. Sudakov [127] pointed out that random subgraph $G(p)$ can be used to measure the robustness of graph properties.

A natural question concerns the existence of transversal spanning structures in random graph systems. A graph property is called *monotone increasing* if it is closed under the addition of edges. For a monotone increasing graph property, the *resilience* quantifies the robustness in terms of the number of edges one must delete from a graph, locally or globally, in order to destroy the property. Ferber, Han and Mao [51] proved a resilient result for a system of n graphs with order n where each graph is an independent copy of $G(n, p)$ provided that $np = \omega(\log n)$.

Theorem 5.1 ([51]). *For every $\epsilon > 0$, there exists an absolute constant $c = c(\epsilon)$ such that for any sufficiently large n and $p \geq \frac{c \log n}{n}$ the following holds. Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a system of n independent copies of $G(n, p)$ on the same vertex set V of size n . For all $i \in [n]$, let $H_i \subseteq G_i$ be such that $d_{H_i}(v) \geq (\frac{1}{2} + \epsilon)d_{G_i}(v)$ for all $v \in V$. Then, whp $\{H_1, \dots, H_n\}$ contains a transversal Hamilton cycle.*

Similar result for transversal perfect matching was also obtained in [51]. Note that the above theorem implies that if $\mathcal{G} = \{G_1, \dots, G_n\}$ is a graph system of order n with $\delta(\mathcal{G}) \geq (1/2 + \epsilon)n$, and $\mathcal{F} = \{F_1, \dots, F_n\}$ where $F_i = G_i(p)$ for some $p \geq c \log n/n$, then whp \mathcal{F} contains a transversal Hamilton cycle. Anastos and Chakraborti [11] improved this by removing the error term in the minimum degree condition with spread techniques.

Theorem 5.2 ([11]). *There exists an absolute c such that the following holds. Suppose that $\mathcal{G} = \{G_1, \dots, G_n\}$ is a graph system of order n with $\delta(\mathcal{G}) \geq \frac{n}{2}$, and $p \geq \frac{c \log n}{n}$. Let $\mathcal{F} = \{F_1, \dots, F_n\}$ where $F_i = G_i(p)$. Then, whp \mathcal{F} contains a transversal Hamilton cycle.*

In the above theorem, each graph in the given system intersect with the same random graph $G(n, p)$. They [11] also considered a system that consists of n independent random subgraphs of a Dirac graph.

Theorem 5.3 ([11]). *There exists an absolute constant c such that the following holds. Suppose G is a graph on n vertices with $\delta(G) \geq \frac{n}{2}$, and $p \geq \frac{c \log n}{n^2}$. Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a family of graphs where each G_i is independently distributed as $G(p)$, for $i \in [n]$. Then, whp \mathcal{G} contains a transversal Hamilton cycle.*

Naturally, they [11] hoped for a combined generalization of the above two theorems.

Problem 5.4 ([11]). Does there exist a constant c such that the following holds? Let $\mathcal{G} = \{G_1, \dots, G_n\}$ be a graph system of order n with $\delta(\mathcal{G}) \geq \frac{n}{2}$, and $p \geq \frac{c \log n}{n^2}$. For each $i \in [n]$, let $F_i = G_i(p)$ and let $\mathcal{F} = \{F_1, \dots, F_n\}$. Then, whp \mathcal{F} contains a transversal Hamilton cycle.

Anastos and Chakraborti [11] showed that Problem 5.4 holds when n is odd, whereas it turns out to be false when n is even.

Han, Hu and Yang [65] obtained a transversal robust version of the Hajnal–Szemerédi theorem [63].

Theorem 5.5 ([65]). *Let $r \geq 3$ be an integer and let $\gamma > 0$. There exists a constant $c = c(r, \gamma)$ such that for any sufficiently large $n \in r\mathbb{N}$ and $p \geq cn^{-\frac{2}{r}}(\log n)^{1/\binom{r}{2}}$ the following holds. Suppose that $\mathcal{G} = \{G_1, \dots, G_{\frac{n}{r}\binom{r}{2}}\}$ is a graph system of order n with $\delta(\mathcal{G}) \geq (1 - \frac{1}{r} + \gamma)n$. Let $\mathcal{F} = \{F_1, \dots, F_{\frac{n}{r}\binom{r}{2}}\}$ where $F_i = G_i(p)$. Then whp \mathcal{F} contains a transversal K_r -factor.*

The transversal robust version of the multipartite Hajnal–Szemerédi theorem [63] was also obtained in [65].

6 Digraph systems

In this section, we consider transversal problems in digraph systems. Let D be a *digraph* with no loops and at most one edge in each direction between every pair of vertices. An *oriented graph* is a digraph with no cycle of length two. Let $\delta^+(D)$ and $\delta^-(D)$ be the *minimum out-degree* and *minimum in-degree*, respectively. Let $\delta^0(D) := \min\{\delta^+(D), \delta^-(D)\}$ be the *minimum semi-degree* of D . Let $\mathcal{D} = \{D_1, D_2, \dots, D_m\}$ be a system of not necessarily distinct digraphs with common vertex set V . Define $\delta^+(\mathcal{D}) := \min\{\delta^+(D_i) : i \in [m]\}$ and $\delta^0(\mathcal{D}) := \min\{\delta^0(D_i) : i \in [m]\}$.

A *tournament* is an oriented graph where every pair of distinct vertices are adjacent. We call a tournament T *strongly connected* if for every pair of vertices x, y in T , there is a directed path from x to y . We say that a digraph D is *transitive* if xy and yz are arcs in D with $x \neq z$, then the arc xz is also in D .

Let T_k be the transitive tournament on k vertices and \mathcal{T}_k be the family of all tournaments on k vertices. Czygrinow, DeBiasio, Kierstead and Molla [35] proved that every n -vertex digraph D with $\delta^+(D) \geq (1 - 1/k)n$ contains a T_k -factor for integers n, k with $k|n$. Treglown [131] proved that for $T \in \mathcal{T}_k$ with $k \geq 3$ and sufficiently large integer n with $k|n$, every digraph D on n vertices with $\delta^0(D) \geq (1 - 1/k)n$ contains a T -factor.

In contrast to Theorem 3.13, it is natural to seek an analogue of the transversal Hajnal–Szemerédi theorem [63] in the digraph systems. Cheng, Han, Wang and Wang [27] established the minimum out-degree and semi-degree condition for the existence of transversal tournament factors in digraph systems.

Theorem 6.1 ([27]). *For every integer $k \geq 3$, $T \in \mathcal{T}_k$ and real $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that the following holds for all integers $n \geq n_0$ and $n \in k\mathbb{N}$. If $\mathcal{D} = \{D_1, \dots, D_m\}$, $m = \frac{n}{k}\binom{k}{2}$, is a system of n -vertex digraphs on the same vertex set such that $\delta^+(\mathcal{D}) \geq (1 - \frac{1}{k} + \varepsilon)n$ (resp. $\delta^0(\mathcal{D}) \geq (1 - \frac{1}{k} + \varepsilon)n$), then \mathcal{D} contains a transversal T_k -factor (resp. T -factor).*

It is well-known that every tournament contains a Hamilton path, and every strongly connected tournament contains a Hamilton cycle. Chakraborti, Kim, Lee and Seo [25] proved the following result concerning the existence of transversal directed Hamilton paths and cycles in a system of tournaments.

Theorem 6.2 ([25]). *For every sufficiently large n ,*

- (i) *every system of $(n - 1)$ tournaments with order n contains a transversal directed Hamilton path;*

- (ii) every system \mathcal{T} of n tournaments with order n satisfies the following. If all tournaments in \mathcal{T} possibly except one are strongly connected, then \mathcal{T} contains a transversal directed Hamilton cycle.

Thomason [130] proved that every sufficiently large tournament contains Hamilton paths and cycles with all possible orientations, except possibly the consistently oriented Hamilton cycle. Chakraborti, Kim, Lee and Seo [26] established the transversal generalization of this result.

Theorem 6.3 ([26]). *For every sufficiently large n ,*

- (i) every system of $(n - 1)$ tournaments with order n contains a transversal of every orientation of the Hamilton path;
- (ii) every system of n tournaments with order n contains a transversal of every orientation of the Hamilton cycle except possibly the directed one.

Babiński, Grzesik and Prorok [13] studied the transversal version of Mantel's theorem in digraph systems and oriented graph systems. More precisely, they determined the minimum number of edges in each graph that ensure the existence of transversal transitive triangles or transversal directed triangles in digraph systems and oriented graph systems. Gerbner, Grzesik, Palmer and Prorok [58] determined the minimum number of edges of digraphs in a digraph system which guarantees the existence of any rainbow directed star.

Ghouila-Houri [59] showed that any n -vertex digraph with minimum semi-degree at least $n/2$ contains a directed Hamilton cycle. Cheng, Li, Sun and Wang [29] provided a transversal generalization of Ghouila-Houri's theorem, which solved a problem proposed by Chakraborti, Kim, Lee and Seo [26].

Theorem 6.4 ([29]). *Let $\mathcal{D} = \{D_1, \dots, D_n\}$ be a system of digraphs on a common vertex set V of sufficiently large order n . If $\delta^0(\mathcal{D}) \geq \frac{n}{2}$, then \mathcal{D} contain a transversal directed Hamilton cycle.*

DeBiasio and Molla [38] proved the the anti-directed Hamilton cycle (an orientation in which consecutive edges alternate direction) is guaranteed to appear if $\delta^0(D) \geq n/2 + 1$. DeBiasio, Kühn, Molla, Osthus and Taylor [37] showed that for sufficiently large n , every n -vertex digraph D with $\delta^0(D) \geq n/2$ contains every orientation of a Hamilton cycle except, possibly, the anti-directed one. It is interesting to generalize the above two results to transversal version.

Keeshash, Kühn and Osthus [77] proved the oriented version of Dirac's theorem, which states that every n -vertex digraph with minimum semi-degree at least $(3n - 4)/8$ contains a directed Hamilton cycle. Considering a transversal version of this theorem would be interesting.

Question 6.5. *Let $\mathcal{D} = \{D_1, \dots, D_n\}$ be a system of oriented graphs with common vertex set V of size n . If $\delta^0(\mathcal{D}) \geq \lceil \frac{3n-4}{8} \rceil$, does \mathcal{D} contain a transversal directed Hamilton cycle?*

7 Hypergraph systems

In this section, we will survey results in k -uniform hypergraph systems. For a set V , we denote by $\binom{V}{k}$ the set of subsets of size k of V . A k -uniform hypergraph (or k -graph) $H = (V, E)$ consists of a vertex set V and an edge set E which is a family of k -element subsets of V , i.e., $E \subseteq \binom{V}{k}$. For any $S \subseteq V$, the degree of S in H , denoted by $\deg_H(S)$, is the number of edges containing S . For any integer $s \geq 0$, define the *minimum s -degree* $\delta_s(H)$ to be $\min\{\deg_H(S) : S \in \binom{V(H)}{s}\}$. Let $\mathcal{H} = \{H_1, H_2, \dots, H_m\}$ be a system of not necessarily distinct k -graphs with common vertex set V . Define $\delta_s(\mathcal{H}) = \min\{\delta_s(H_i) : i \in [m]\}$.

The problem of characterizing k -graphs which contain a Hamilton cycle or a perfect matching is perhaps the most prominent one in hypergraph theory. In the remainder of this section, we focus on those two most well studied structures.

7.1 Transversal Hamilton cycle in hypergraph systems

We say that a k -graph C is an ℓ -cycle if there exists a cyclic ordering of the vertices of C such that every edge of C consists of k consecutive vertices and every two consecutive edges (in the natural ordering of the edges) intersects in precisely ℓ vertices. We say that C is a *Hamilton ℓ -cycle* of a k -graph H if C is a spanning sub- k -graph of H . We call C a *tight Hamilton cycle* if $\ell = k - 1$, a *loose Hamilton cycle* if $\ell = 1$. A Hamilton cycle of a k -graph H is a cycle which is a spanning subhypergraph of H . Note that if a k -graph H on n vertices contains a Hamilton ℓ -cycle then $(k - \ell)|n$, since every edge of the cycle contains exactly $k - \ell$ vertices which were not contained in the previous edge.

In 1999, Katona and Kierstead [76] proved a Dirac-type theorem: every n -vertex k -graph H with $\delta_{k-1}(H) > (1 - \frac{1}{2k})n + 4 - k - \frac{5}{2k}$ contains a tight Hamilton cycle. They anticipated that a much stronger result is true.

Conjecture 7.1 ([76]). *Let H be a k -graph on $n \geq k + 1 \geq 4$ vertices. If $\delta_{k-1}(H) \geq \lfloor \frac{n-k+3}{2} \rfloor$, then H has a tight Hamilton cycle.*

Rödl, Ruciński and Szemerédi [121, 122] confirmed this asymptotically, and they [123] also gave the exact version for $k = 3$.

Theorem 7.2 ([123, 121, 122]). *Let $k \geq 3$, $\gamma > 0$ and H be an n -vertex k -graph, where n is sufficiently large. If $\delta_{k-1}(H) \geq (\frac{1}{2} + \gamma)n$, then H contains a tight Hamilton cycle. Furthermore, when $k = 3$ it is enough to have $\delta_2(H) \geq \lfloor \frac{n}{2} \rfloor$.*

However, Kühn and Osthus [96] and Zhao [135] noted that it is much more difficult to determine the minimum d -degree condition for tight Hamilton cycle for $d \in [k - 2]$. The best general bound for such a problem was given by Lang and Sanhueza-Matamala [99], Polcyn, Reiher, Rödl and Schülke [119] independently.

Theorem 7.3 ([99, 119]). *Let $k \geq 3$, $\gamma > 0$ and H be an n -vertex k -graph, where n is sufficiently large. If $\delta_{k-2}(H) \geq (\frac{5}{9} + \gamma)\binom{n}{2}$, then H contains a tight Hamilton cycle.*

Let $1 \leq d, \ell \leq k - 1$. For $n \in (k - \ell)\mathbb{N}$, define $h_d^\ell(k, n)$ to be the smallest integer h such that every n -vertex k -graph H satisfying $\delta_d(H) \geq h$ contains a Hamilton ℓ -cycle. Han and Zhao [66] proved that

$$h_d^{k-1}(k, n) \geq \left(1 - \binom{t}{\lfloor \frac{t}{2} \rfloor} \frac{\lceil \frac{t}{2} \rceil \lceil \frac{t}{2} \rceil (\lfloor \frac{t}{2} \rfloor + 1)^{\lfloor \frac{t}{2} \rfloor}}{(t+1)^t + o(1)} \right) \binom{n}{t}, \quad (1)$$

where $d \in [k - 1]$ and $t = k - d$. In particular, $h_{k-3}^{k-1}(k, n) \geq (5/8 + o(1))\binom{n}{3}$. Lang and Sanhueza-Matamala [99] proved $h_d^{k-1}(k, n) \leq 2^{-1/(k-d)}$ and they conjectured that the minimum d -degree threshold for k -uniform tight Hamilton cycles coincides with the lower bounds in (1).

Recently, Cheng, Han, Wang, Wang and Yang [28] and Tang, Wang, Wang and Yan [129] extended Theorem 7.2 and Theorem 7.3 to transversal version, respectively.

Theorem 7.4 ([28, 129]). *For every $k \geq 3, \gamma > 0$, there exists n_0 such that the following holds for $n \geq n_0$. Let $\mathcal{H} = \{H_1, \dots, H_n\}$ be a k -graph system of order n .*

- (i) *If $\delta_{k-1}(\mathcal{H}) \geq (\frac{1}{2} + \gamma)n$, then \mathcal{H} admits a transversal tight Hamilton cycle.*
- (ii) *If $\delta_{k-2}(\mathcal{H}) \geq (\frac{5}{9} + \gamma)\binom{n}{2}$, then \mathcal{H} admits a transversal tight Hamilton cycle.*

Cheng, Han, Wang, Wang and Yang [28] suspected that the threshold for transversal tight Hamilton cycles in a k -graph system is the same with the threshold conjectured for tight Hamilton cycles in a single k -graph (i.e., Conjecture 7.1) and provided the following conjecture.

Conjecture 7.5 ([28]). *Suppose $\mathcal{H} = \{H_1, \dots, H_n\}$ is a k -graph system of order n ($n \geq k + 1 \geq 4$) and $\delta_{k-1}(\mathcal{H}) \geq \lfloor \frac{n-k+3}{2} \rfloor$, then \mathcal{H} contains a transversal tight Hamilton cycle.*

Motivated by (1), Tang, Wang, Wang and Yan [129] proposed the following conjecture, which concerns the minimum d -degree condition for transversal tight Hamilton cycles.

Conjecture 7.6 ([129]). *For every $k \geq d + 1$, $\mu > 0$, there exists n_0 such that the following holds for $n \geq n_0$. Let $\mathcal{H} = \{H_1, \dots, H_n\}$ be a k -graph system of order n .*

- (i) *If $k \geq 4$ and $\delta_{k-3}(\mathcal{H}) \geq (\frac{5}{8} + \mu)\binom{n}{3}$, then \mathcal{H} admits a transversal tight Hamilton cycle.*
- (ii) *If $\delta_d(\mathcal{H}) \geq h_d^{k-1}(k, n) + \mu\binom{n}{d}$, then \mathcal{H} admits a transversal tight Hamilton cycle.*

Furthermore, the problem of giving sufficient conditions for transversal Hamilton ℓ -cycles with $\ell \in [k - 2]$ is still open. It would be very nice to settle this problem.

Theorem 7.4 implies that if we consider $(k - 1)$ -degree and $(k - 2)$ -degree conditions, then tight Hamilton cycle is color-blind. Gupta, Hamann, Müyesser, Parczyk and Sgueglia [62] provided a unified approach to find color-blind graphs, which is general enough to recover many previous results and obtain novel transversal variants of several classical Dirac-type results. The precise statement of their main result is quite technical, we refer the reader to see [62, Theorem 2.6]. The following theorem lists some applications in their paper and they believed that their setting can capture even more families of hypergraphs.

Theorem 7.7 ([62]). *Under d -degree condition, the following families of hypergraphs are all color-blind.*

- (i) *The family of the r -th powers of Hamilton cycles for $d = 1$ and fixed $r \geq 2$.*
- (ii) *The family of k -uniform Hamilton ℓ -cycles for the following ranges of k , ℓ and d .*
 - $1 < \ell < \frac{k}{2}$ and $d = k - 2$;
 - $1 \leq \ell < \frac{k}{2}$ or $\ell = k - 1$, and $d = k - 1$;
 - $\ell = \frac{k}{2}$ and $\frac{k}{2} < d \leq k - 1$ with k even.

7.2 Transversal matching in hypergraph systems

A t -matching M in a hypergraph H is a subset of $E(H)$ consisting of t pairwise disjoint edges, which is *perfect* if $V(M) = V(H)$. We denote by $\binom{[n]}{r}$ the set of subsets (also called edges) of size r of $[n]$. For a set X , let 2^X be the set of all subsets of X . Let F be a subfamily of 2^X , denote by $\nu(F)$ its *matching number*, that is, the largest m such that F contains m pairwise disjoint sets. A classical problem in extremal set theory is to determine the maximum size of F with fixed $\nu(F)$. In 1965, Erdős [44] proposed the following conjecture, which is known as Erdős matching conjecture.

Conjecture 7.8 (Erdős matching conjecture, [44]). *If $n \geq (s + 1)k$ and a family $F \subset \binom{[n]}{k}$ satisfies $\nu(F) \leq s$, then $|F| \leq \max\{\binom{n}{k} - \binom{n-s}{k}, \binom{(s+1)k-1}{k}\}$.*

Notice that the bound in Conjecture 7.8 is tight. In fact, one may consider the edge set of the complete k -graph on $(s + 1)k - 1$ vertices or the k -graph on n vertices in which every edge intersects a fixed set of s vertices. Setting $s = 1$ in Conjecture 7.8, then it corresponds to the well-known Erdős–Ko–Rado theorem [45]. There are numerous papers devoted to the Erdős matching conjecture. For recent development on this conjecture, see [10, 54, 64, 89].

Many scholars have attempted to generalize the Erdős matching conjecture to transversal version in hypergraph systems. Matsumoto and Tokushige [112] generalized the Erdős–Ko–Rado theorem to two families, that is,

Theorem 7.9 ([112]). *Let n, k, ℓ be integers such that $n \geq \max\{2k, 2\ell\}$. Assume that $\mathcal{H}_1 \subseteq \binom{[n]}{k}$ and $\mathcal{H}_2 \subseteq \binom{[n]}{\ell}$. If $|\mathcal{H}_1||\mathcal{H}_2| > \binom{n-1}{k-1}\binom{n-1}{\ell-1}$, then there exist two disjoint subsets $X_1 \in \mathcal{H}_1, X_2 \in \mathcal{H}_2$.*

It is natural to extend Erdős–Ko–Rado theorem to more than two hypergraphs. For this, Aharoni and Howard [6] and Huang, Loh and Sudakov [70] proposed the following conjecture independently.

Conjecture 7.10 ([6, 70]). *Let $\mathcal{H} = \{H_1, \dots, H_t\}$ be a system of k -graphs on a common vertex set V of size n . If $|E(H_i)| > \max \left\{ \binom{n}{k} - \binom{n-t+1}{k}, \binom{kt-1}{k} \right\}$ for all $1 \leq i \leq t$, then \mathcal{H} has a transversal t -matching.*

When $k = 2$, this conjecture was confirmed by Meshulam (see [6]). Indeed, the case $k = 2$ of this conjecture follows directly from an earlier result established by Akiyama and Frankl [8], as restated in [53]. Huang, Loh, and Sudakov [70] proved that Conjecture 7.10 holds for $n > 3k^2t$. Keller and Lifshitz [84] proved that Conjecture 7.10 holds when $n \geq f(t)k$ for some large constant $f(t)$ which only depends on t , and this was further improved the bound on n to $12kt \log(e^2t)$ by Frankl and Kupavskii [56]. Lu, Wang and Yu [108] improved this bound to $n > 2kt$ for sufficiently large t . By using the sharp threshold techniques developed in [79], Keevash, Lifshitz, Long and Minzer [78] proved a more general result with $n = \Omega(kt)$, which is the optimal order of magnitude. Kupavskii [98] gave the concrete dependencies on the parameters by showing the conjecture holds for $n > 3ekt$ with $t > 10^7$. Recently, Gao, Lu, Ma and Yu [57] showed that Conjecture 7.10 is true when $k = 3$ and n is sufficiently large.

Aharoni and Howard [6] proposed the following conjecture in k -partite k -graph systems.

Conjecture 7.11 ([6]). *Let $\mathcal{H} = \{H_1, \dots, H_t\}$ be a k -partite k -graph systems with n vertices in each partition class. If $|E(H_i)| \geq (t-1)n^{k-1}$ for all $i \in [t]$, then \mathcal{H} admits a transversal t -matching.*

Aharoni and Howard proved that Conjecture 7.11 holds when $t = 2$ [6] or $k \leq 3$ [7]. Lu and Yu [109] proved Conjecture 7.11 for $n > 3(k-1)(t-1)$. Kiselev and Kupavskii [88] showed that this conjecture is true for $t \geq t_0$ where t_0 is a constant.

The degree conditions for the existence of large matchings in uniform hypergraphs have been extensively studied. Kühn, Osthus and Treglown [97] (also see Khan [85]) and Khan [86] determined the minimum 1-degree condition for an n -vertex 3-graph and 4-graph containing a perfect matching, respectively. Lu, Yu and Yuan [110] and Lu, Wang and Yu [107] proved the transversal version of the above results.

Theorem 7.12 ([110, 107]). *Let n be a sufficiently large integer and $k \in \{3, 4\}$ with $k|n$. If $\mathcal{H} = \{H_1, \dots, H_{\frac{n}{k}}\}$ is a k -graph system of order n with $\delta_1(\mathcal{H}) > \binom{n-1}{k} - \binom{(k-1)n/k}{k}$, then \mathcal{H} contains a transversal perfect matching.*

Recently, Lu and Wang [106] determined the minimum 1-degree condition that guarantees a 3-partite 3-graph system admits transversal perfect matchings. This generalized a result of Lo and Markström [105] on the 1-degree threshold for the existence of perfect matchings in 3-partite 3-graphs. Moreover, in their proof, they used fractional transversal matching theory obtained by Aharoni, Holzman and Jiang [5].

It is well-known that perfect matchings are closely related to its fractional counterpart. Given a k -graph H , a *fractional matching* is a function $f : E(H) \rightarrow [0, 1]$, subject to the requirement that $\sum_{e:v \in e} f(e) \leq 1$, for every $v \in V(H)$. Furthermore, if equality holds for every $v \in V(H)$, then we call the fractional matching *perfect*. Denote the maximum size of a fractional matching of H by $\nu^*(H) = \max_f \sum_{e \in E(H)} f(e)$. Let $c_{k,d}$ be the minimum d -degree threshold for perfect fractional matchings in k -graphs, namely, for every $\varepsilon > 0$ and sufficiently large $n \in \mathbb{N}$, every n -vertex k -graph H with $\delta_d(H) \geq (c_{k,d} + \varepsilon) \binom{n-d}{k-d}$ contains a perfect fractional matching.

Aharoni, Holzman and Jiang [5] studied the existence of rainbow fractional matching in a k -graph system.

Theorem 7.13 ([5]). *Let $k \geq 2$ be an integer, and let n be a positive rational number. Assume that $\mathcal{H} = \{H_1, \dots, H_{\lceil kn \rceil}\}$ is a k -graph system with $\nu^*(H_i) \geq n$ for all $i \in [\lceil kn \rceil]$. Then \mathcal{H} has a rainbow fractional matching of size n .*

When the k -graph is k -partite and n is an integer, the number of graphs in \mathcal{H} required in Theorem 7.13 goes down from kn to $kn - k + 1$. Theorem 7.13 provided a fractional version of the corresponding problem concerning rainbow matchings, which was solved by Drisko [42] and by Aharoni and Berger [1] for bipartite graphs (the rainbow matching problem remains open for general graphs as well as for k -partite hypergraphs when $k > 2$).

Alon, Frankl, Huang, Rödl, Ruciński and Sudakov [9] established the relationship between perfect matching and perfect fractional matching by showing that every n -vertex k -graph H with $\delta_d(H) \geq (\max\{c_{k,d}, 1/2\} + o(1))\binom{n-d}{k-d}$ has a perfect matching, and this condition is asymptotically best possible. Cheng, Han, Wang, and Wang [27] proved that the d -degree condition that ensures the existence a perfect matching in a k -graph also guarantees the existence of a transversal perfect matching in a k -graph system for $d \in [k - 1]$.

Theorem 7.14 ([27]). *For every $\varepsilon > 0$ and integer $d \in [k - 1]$, there exists $n_0 \in \mathbb{N}$, such that the following holds for all integers $n \geq n_0$ and $n \in k\mathbb{N}$. Every n -vertex k -graph system $\mathcal{H} = \{H_1, \dots, H_{\frac{n}{k}}\}$ with $\delta_d(\mathcal{H}) \geq (\max\{c_{k,d}, \frac{1}{2}\} + \varepsilon)\binom{n-d}{k-d}$ contains a transversal perfect matching.*

8 Further results

There are some additional results on transversal problems in graph systems whose details we will not discuss in this survey. For convenience of the reader we give a concise list of such results together with relevant references.

- The rainbow Turán problem can be viewed as a special case of Turán problems in graph systems. Let H be a graph. The *rainbow Turán number* of H , denoted by $\text{ex}^*(n, H)$, is defined to be the the maximum number of edges in a properly edge-colored n -vertex graph G with no rainbow copies of H . Notice that this properly edge-colored graph G can be viewed as system $\mathcal{G} = \{G_1, \dots, G_{\chi(G)}\}$ of graphs where each G_i is the graph consisting of edges with color i . Then find a rainbow copy of H in G is equivalent to find a transversal copy of H inside \mathcal{G} . The rainbow Turán number of some classical families of graphs were studied by Keevash, Mubayi, Sudakov and Verstraëte [80]. See more results in [17, 36, 73, 87].
- In section 3, we only survey results on the minimum degree condition for transversal spanning structures in graph systems. In fact, some studies have considered the problem of embedding transversal non-spanning structures in a graph system. Li, Li and Li [101] studied the existence of transversal cliques in graph systems under Dirac-type conditions. Falgas-Ravry, Markström and Rätty [49] studied the minimum degree version of Problem 2.2, and they found that the extremal behavior in the minimum degree condition differs strikingly from that seen in the density condition, with discrete jumps as opposed to continuous transitions.
- The *adjacency matrix* of G is $A(G) = (a_{ij})$, where $a_{xy} = 1$ if $xy \in E(G)$ and 0 otherwise. The largest eigenvalue of $A(G)$ is called the *spectral radius* of G . The maximum spectral radius of graphs in a graph system that forbids transversal perfect matching, Hamilton path or Hamilton cycle was studied in [61, 67, 134].
- One may consider the problem of embedding a rainbow graph H in a graph system \mathcal{G} provided that H and all graphs in \mathcal{G} belong to the same class. For example, Aharoni, Briggs, Holzman and Jiang [2] proved that every family of $2\lfloor n/2 \rfloor - 1$ odd cycles of order n contains a rainbow odd cycle. For more results, we refer the reader to [41, 60, 100].

9 Conclusion

Although we have made an effort to provide a systematic coverage of recent classical results in graph systems, it is inevitable that some topics were left out of this survey due to constraints of space and time. Nevertheless, we would like to believe that we have presented enough results demonstrating

how one can generalize known results in extremal graph theory to transversal version. Therefore we hope this survey will inspire further research in this area.

10 Acknowledgments

The authors would like to thank Yangyang Cheng and Jie Hu for helpful comments and valuable suggestions.

References

- [1] R. Aharoni and E. Berger. Rainbow matchings in r -partite r -graphs. *Electron. J. Combin.*, 16(1):Research Paper 119, 9, 2009.
- [2] R. Aharoni, J. Briggs, R. Holzman, and Z. Jiang. Rainbow odd cycles. *SIAM J. Discrete Math.*, 35(4):2293–2303, 2021.
- [3] R. Aharoni, M. DeVos, S. González Hermosillo de la Maza, A. Montejano, and R. Šámal. A rainbow version of Mantel’s theorem. *Adv. Comb.*, pages Paper No. 2, 12, 2020.
- [4] R. Aharoni, A. Georgakopoulos, and P. Sprüssel. Perfect matchings in r -partite r -graphs. *European J. Combin.*, 30(1):39–42, 2009.
- [5] R. Aharoni, R. Holzman, and Z. Jiang. Rainbow fractional matchings. *Combinatorica*, 39(6):1191–1202, 2019.
- [6] R. Aharoni and D. Howard. Size conditions for the existence of rainbow matchings. 2011. <https://math.colgate.edu/dmhoward/rsc.pdf>.
- [7] R. Aharoni and D. Howard. A rainbow r -partite version of the Erdős-Ko-Rado theorem. *Combin. Probab. Comput.*, 26(3):321–337, 2017.
- [8] J. Akiyama and P. Frankl. On the size of graphs with complete factors. *J. Graph Theory*, 9(1):197–201, 1985.
- [9] N. Alon, P. Frankl, H. Huang, V. Rödl, A. Ruciński, and B. Sudakov. Large matchings in uniform hypergraphs and the conjecture of Erdős and Samuels. *J. Combin. Theory Ser. A*, 119(6):1200–1215, 2012.
- [10] N. Alon, H. Huang, and B. Sudakov. Nonnegative k -sums, fractional covers, and probability of small deviations. *J. Combin. Theory Ser. B*, 102(3):784–796, 2012.
- [11] M. Anastos and D. Chakraborti. Robust Hamiltonicity in families of Dirac graphs. *arXiv preprint arXiv:2309.12607*, 2023.
- [12] S. Babiński and A. Grzesik. Graphs without a rainbow path of length 3. *SIAM J. Discrete Math.*, 38(1):629–644, 2024.
- [13] S. Babiński, A. Grzesik, and M. Prorok. Directed graphs without rainbow triangles. *arXiv preprint arXiv:2308.01461*, 2023.
- [14] I. Bárány. A generalization of Carathéodory’s theorem. *Discrete Math.*, 40(2-3):141–152, 1982.
- [15] B. Bollobás. *Extremal graph theory*, volume 11 of *London Mathematical Society Monographs*. Academic Press, Inc., London-New York, 1978.
- [16] J. A. Bondy. Pancyclic graphs. I. *J. Combinatorial Theory Ser. B*, 11:80–84, 1971.

- [17] J. A. Bondy and M. Simonovits. Cycles of even length in graphs. *J. Combinatorial Theory Ser. B*, 16:97–105, 1974.
- [18] J. Böttcher, M. Schacht, and A. Taraz. Proof of the bandwidth conjecture of Bollobás and Komlós. *Math. Ann.*, 343(1):175–205, 2009.
- [19] C. Bowtell, P. Morris, Y. Pehova, and K. Staden. Universality for transversal Hamilton cycles. *arXiv preprint arXiv:2310.04138*, 2023.
- [20] P. Bradshaw. Transversals and bipancyclicity in bipartite graph families. *Electron. J. Combin.*, 28(4):Paper No. 4.25, 20, 2021.
- [21] P. Bradshaw, K. Halasz, and L. Stacho. From one to many rainbow Hamiltonian cycles. *Graphs Combin.*, 38(6):Paper No. 188, 21, 2022.
- [22] R. A. Brualdi and H. J. Ryser. *Combinatorial matrix theory*, volume 39 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1991.
- [23] D. Chakraborti, S. Im, J. Kim, and H. Liu. A bandwidth theorem for graph transversals. *arXiv preprint arXiv:2302.09637*, 2023.
- [24] D. Chakraborti, J. Kim, H. Lee, H. Liu, and J. Seo. On a rainbow extremal problem for color-critical graphs. *Random Structures Algorithms*, 64(2):460–489, 2024.
- [25] D. Chakraborti, J. Kim, H. Lee, and J. Seo. Hamilton transversals in tournaments. *Combinatorica*, 2024. <https://doi.org/10.1007/s00493-024-00123-1>.
- [26] D. Chakraborti, J. Kim, H. Lee, and J. Seo. Transversal cycles and paths in tournaments. *arXiv preprint arXiv:2407.14300*, 2024.
- [27] Y. Cheng, J. Han, B. Wang, and G. Wang. Rainbow spanning structures in graph and hypergraph systems. *Forum Math. Sigma*, 11:Paper No. e95, 20, 2023.
- [28] Y. Cheng, J. Han, B. Wang, G. Wang, and D. Yang. Transversal Hamilton cycle in hypergraph systems. *arXiv preprint arXiv:2111.07079*, 2021.
- [29] Y. Cheng, H. Li, W. Sun, and G. Wang. Transversal Hamilton cycles in digraph collections. *In preparation*, 2024.
- [30] Y. Cheng and K. Staden. Transversals via regularity. *arXiv preprint arXiv:2306.03595*, 2023.
- [31] Y. Cheng and K. Staden. Stability of transversal Hamilton cycles and paths. *arXiv preprint arXiv:2403.09913*, 2024.
- [32] Y. Cheng, W. Sun, G. Wang, and L. Wei. Transversal Hamilton paths and cycles. *arXiv preprint arXiv:2406.13998*, 2024.
- [33] Y. Cheng, G. Wang, and Y. Zhao. Rainbow pancyclicity in graph systems. *Electron. J. Combin.*, 28(3):Paper No. 3.24, 9, 2021.
- [34] K. Corrádi and A. Hajnal. On the maximal number of independent circuits in a graph. *Acta Mathematica Hungarica*, 14(3-4):423–439, 1963.
- [35] A. Czygrinow, L. DeBiasio, H. A. Kierstead, and T. Molla. An extension of the Hajnal-Szemerédi theorem to directed graphs. *Combin. Probab. Comput.*, 24(5):754–773, 2015.
- [36] S. Das, C. Lee, and B. Sudakov. Rainbow Turán problem for even cycles. *European J. Combin.*, 34(5):905–915, 2013.

- [37] L. DeBiasio, D. Kühn, T. Molla, D. Osthus, and A. Taylor. Arbitrary orientations of Hamilton cycles in digraphs. *SIAM J. Discrete Math.*, 29(3):1553–1584, 2015.
- [38] L. DeBiasio and T. Molla. Semi-degree threshold for anti-directed Hamiltonian cycles. *Electron. J. Combin.*, 22(4):Paper 4.34, 23, 2015.
- [39] G. A. Dirac. Some theorems on abstract graphs. *Proc. London Math. Soc. (3)*, 2:69–81, 1952.
- [40] A. A. Diwan and D. Mubayi. Turán’s theorem with colors. 2006. <https://homepages.math.uic.edu/mubayi/papers/coloredturan.pdf>.
- [41] Z. Dong and Z. Xu. Rainbow even cycles. *SIAM J. Discrete Math.*, 38(2):1269–1284, 2024.
- [42] A. A. Drisko. Transversals in row-Latin rectangles. *J. Combin. Theory Ser. A*, 84(2):181–195, 1998.
- [43] P. Erdős. Problem 9. *Theory of Graphs and its Applications (Pro. Sympo. Smolenice, 1963)*, 13:85–90, 1964.
- [44] P. Erdős. A problem on independent r -tuples. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, 8:93–95, 1965.
- [45] P. Erdős, C. Ko, and R. Rado. Intersection theorems for systems of finite sets. *Quart. J. Math. Oxford Ser. (2)*, 12:313–320, 1961.
- [46] P. Erdős and M. Simonovits. A limit theorem in graph theory. *Studia Sci. Math. Hungar.*, 1:51–57, 1966.
- [47] P. Erdős and A. H. Stone. On the structure of linear graphs. *Bull. Amer. Math. Soc.*, 52:1087–1091, 1946.
- [48] L. Euler. Recherches sur un nouvelle espèce de quarrés magiques. *Verh. Zeeuw. Genoot. Weten. Vliss 9*, pages 85–239, 1782.
- [49] V. Falgas-Ravry, K. Markström, and E. Rätty. Minimum-degree conditions for rainbow triangles. *J. Graph Theory*, 107(2):298–329, 2024.
- [50] V. Falgas-Ravry, K. Markström, and E. Rätty. Rainbow variations on a theme by Mantel: extremal problems for Gallai colouring templates. *Combinatorica*, 44(5):977–1010, 2024.
- [51] A. Ferber, J. Han, and D. Mao. Dirac-type problem of rainbow matchings and Hamilton cycles in random graphs. *arXiv preprint arXiv:2211.05477*, 2022.
- [52] E. Fischer. Variants of the Hajnal-Szemerédi theorem. *J. Graph Theory*, 31(4):275–282, 1999.
- [53] P. Frankl. The shifting technique in extremal set theory. In *Surveys in combinatorics 1987 (New Cross, 1987)*, volume 123 of *London Math. Soc. Lecture Note Ser.*, pages 81–110. Cambridge Univ. Press, Cambridge, 1987.
- [54] P. Frankl. Improved bounds for Erdős’ matching conjecture. *J. Combin. Theory Ser. A*, 120(5):1068–1072, 2013.
- [55] P. Frankl. Graphs without rainbow triangles. *arXiv preprint arXiv:2203.07768*, 2022.
- [56] P. Frankl and A. Kupavskii. Simple juntas for shifted families. *Discrete Anal.*, pages Paper No. 14, 18, 2020.
- [57] J. Gao, H. Lu, J. Ma, and X. Yu. On the rainbow matching conjecture for 3-uniform hypergraphs. *Sci. China Math.*, 65(11):2423–2440, 2022.

- [58] D. Gerbner, A. Grzesik, C. Palmer, and M. Prorok. Directed graphs without rainbow stars. *arXiv preprint arXiv:2402.01028*, 2024.
- [59] A. Ghouila-Houri. Une condition suffisante d'existence d'un circuit hamiltonien. *C. R. Acad. Sci. Paris*, 251:495–497, 1960.
- [60] I. Goorevitch and R. Holzman. Rainbow triangles in families of triangles. *arXiv preprint arXiv:2209.15493*, 2022.
- [61] M. Guo, H. Lu, X. Ma, and X. Ma. Spectral radius and rainbow matchings of graphs. *Linear Algebra Appl.*, 679:30–37, 2023.
- [62] P. Gupta, F. Hamann, A. Müyesser, O. Parczyk, and A. Sgueglia. A general approach to transversal versions of Dirac-type theorems. *Bull. Lond. Math. Soc.*, 55(6):2817–2839, 2023.
- [63] A. Hajnal and E. Szemerédi. Proof of a conjecture of P. Erdős. In *Combinatorial theory and its applications, I-III (Proc. Colloq., Balatonfüred, 1969)*, volume 4 of *Colloq. Math. Soc. János Bolyai*, pages 601–623. North-Holland, Amsterdam-London, 1970.
- [64] J. Han. Perfect matchings in hypergraphs and the Erdős matching conjecture. *SIAM J. Discrete Math.*, 30(3):1351–1357, 2016.
- [65] J. Han, J. Hu, and D. Yang. A robust version of the multipartite Hajnal–Szemerédi theorem. *arXiv preprint arXiv:2311.00950*, 2023.
- [66] J. Han and Y. Zhao. Forbidding Hamilton cycles in uniform hypergraphs. *J. Combin. Theory Ser. A*, 143:107–115, 2016.
- [67] X. He, Y. Li, and L. Feng. Spectral radius and rainbow Hamilton paths of a graph. *Discrete Math.*, 347(10):Paper No. 114128, 11, 2024.
- [68] Z. He, P. Frankl, E. Győri, Z. Lv, N. Salia, C. Tompkins, K. Varga, and X. Zhu. Extremal results for graphs avoiding a rainbow subgraph. *Electron. J. Combin.*, 31(1):Paper No. 1.28, 11, 2024.
- [69] J. Hu, L. Li, X. Li, and N. Xu. Vertex-bipancyclicity in a bipartite graph collection. *Discrete Math.*, 347(7):Paper No. 113980, 10, 2024.
- [70] H. Huang, P.-S. Loh, and B. Sudakov. The size of a hypergraph and its matching number. *Combin. Probab. Comput.*, 21(3):442–450, 2012.
- [71] R. Huang and G.-C. Rota. On the relations of various conjectures on Latin squares and straightening coefficients. *Discrete Math.*, 128(1-3):225–236, 1994.
- [72] S. Im, J. Kim, H. Lee, and H. Seo. On rainbow Turán densities of trees. *arXiv preprint arXiv:2312.15956*, 2023.
- [73] D. Johnston, C. Palmer, and A. Sarkar. Rainbow Turán problems for paths and forests of stars. *Electron. J. Combin.*, 24(1):Paper No. 1.34, 15, 2017.
- [74] F. Joos and J. Kim. On a rainbow version of Dirac's theorem. *Bull. Lond. Math. Soc.*, 52(3):498–504, 2020.
- [75] G. Kalai and R. Meshulam. A topological colorful Helly theorem. *Adv. Math.*, 191(2):305–311, 2005.
- [76] G. Y. Katona and H. A. Kierstead. Hamiltonian chains in hypergraphs. *J. Graph Theory*, 30(3):205–212, 1999.

- [77] P. Keevash, D. Kühn, and D. Osthus. An exact minimum degree condition for Hamilton cycles in oriented graphs. *J. Lond. Math. Soc. (2)*, 79(1):144–166, 2009.
- [78] P. Keevash, N. Lifshitz, E. Long, and D. Minzer. Global hypercontractivity and its applications. *arXiv preprint arXiv:2103.04604*, 2021.
- [79] P. Keevash, N. Lifshitz, E. Long, and D. Minzer. Hypercontractivity for global functions and sharp thresholds. *J. Amer. Math. Soc.*, 37(1):245–279, 2024.
- [80] P. Keevash, D. Mubayi, B. Sudakov, and J. Verstraëte. Rainbow Turán problems. *Combin. Probab. Comput.*, 16(1):109–126, 2007.
- [81] P. Keevash and R. Mycroft. A geometric theory for hypergraph matching. *Mem. Amer. Math. Soc.*, 233(1098):vi+95, 2015.
- [82] P. Keevash and R. Mycroft. A multipartite Hajnal-Szemerédi theorem. *J. Combin. Theory Ser. B*, 114:187–236, 2015.
- [83] P. Keevash, M. Saks, B. Sudakov, and J. Verstraëte. Multicolour Turán problems. *Adv. in Appl. Math.*, 33(2):238–262, 2004.
- [84] N. Keller and N. Lifshitz. The junta method for hypergraphs and the Erdős-Chvátal simplex conjecture. *Adv. Math.*, 392:Paper No. 107991, 95, 2021.
- [85] I. Khan. Perfect matchings in 3-uniform hypergraphs with large vertex degree. *SIAM J. Discrete Math.*, 27(2):1021–1039, 2013.
- [86] I. Khan. Perfect matchings in 4-uniform hypergraphs. *J. Combin. Theory Ser. B*, 116:333–366, 2016.
- [87] J. Kim, J. Lee, H. Liu, and T. Tran. Rainbow cycles in properly edge-colored graphs. *Combinatorica*, 44(4):909–919, 2024.
- [88] S. Kiselev and A. Kupavskii. Rainbow matchings in k -partite hypergraphs. *Bull. Lond. Math. Soc.*, 53(2):360–369, 2021.
- [89] D. Kolupaev and A. Kupavskii. Erdős matching conjecture for almost perfect matchings. *Discrete Math.*, 346(4):Paper No. 113304, 9, 2023.
- [90] J. Komlós, G. N. Sárközy, and E. Szemerédi. On the square of a Hamiltonian cycle in dense graphs. In *Proceedings of the Seventh International Conference on Random Structures and Algorithms (Atlanta, GA, 1995)*, volume 9, pages 193–211, 1996.
- [91] J. Komlós, G. N. Sárközy, and E. Szemerédi. Blow-up lemma. *Combinatorica*, 17(1):109–123, 1997.
- [92] J. Komlós, G. N. Sárközy, and E. Szemerédi. Proof of the Seymour conjecture for large graphs. *Ann. Comb.*, 2(1):43–60, 1998.
- [93] J. Komlós, G. N. Sárközy, and E. Szemerédi. Spanning trees in dense graphs. *Combin. Probab. Comput.*, 10(5):397–416, 2001.
- [94] D. Kühn and D. Osthus. Embedding large subgraphs into dense graphs. In *Surveys in combinatorics 2009*, volume 365 of *London Math. Soc. Lecture Note Ser.*, pages 137–167. Cambridge Univ. Press, Cambridge, 2009.
- [95] D. Kühn and D. Osthus. The minimum degree threshold for perfect graph packings. *Combinatorica*, 29(1):65–107, 2009.

- [96] D. Kühn and D. Osthus. Hamilton cycles in graphs and hypergraphs: an extremal perspective. In *Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. IV*, pages 381–406. Kyung Moon Sa, Seoul, 2014.
- [97] D. Kühn, D. Osthus, and A. Treglown. Matchings in 3-uniform hypergraphs. *J. Combin. Theory Ser. B*, 103(2):291–305, 2013.
- [98] A. Kupavskii. Rainbow version of the Erdős matching conjecture via concentration. *Comb. Theory*, 3(1):Paper No. 1, 19, 2023.
- [99] R. Lang and N. Sanhueza-Matamala. Minimum degree conditions for tight Hamilton cycles. *J. Lond. Math. Soc. (2)*, 105(4):2249–2323, 2022.
- [100] E. Y. H. Li, L. Li, and P. Li. Transversals in a collections of trees. *arXiv preprint arXiv:2310.06354*, 2023.
- [101] L. Li, P. Li, and X. Li. Rainbow structures in a collection of graphs with degree conditions. *J. Graph Theory*, 104(2):341–359, 2023.
- [102] L. Li, P. Li, and X. Li. Rainbow pancyclicity in a collection of graphs under the Dirac-type condition. *Acta Math. Appl. Sin. Engl. Ser.*, 40(2):269–274, 2024.
- [103] X. Li, J. Ma, and Z. Zheng. On the multicolor Turán conjecture for color-critical graphs. *arXiv preprint arXiv:2407.14905*, 2024.
- [104] A. Lo and K. Markström. A multipartite version of the Hajnal-Szemerédi theorem for graphs and hypergraphs. *Combin. Probab. Comput.*, 22(1):97–111, 2013.
- [105] A. Lo and K. Markström. Perfect matchings in 3-partite 3-uniform hypergraphs. *J. Combin. Theory Ser. A*, 127:22–57, 2014.
- [106] H. Lu and Y. Wang. Rainbow perfect matchings in 3-partite 3-uniform hypergraphs. *arXiv preprint arXiv:2408.08523*, 2024.
- [107] H. Lu, Y. Wang, and X. Yu. Rainbow perfect matchings for 4-uniform hypergraphs. *SIAM J. Discrete Math.*, 36(3):1645–1662, 2022.
- [108] H. Lu, Y. Wang, and X. Yu. A better bound on the size of rainbow matchings. *J. Combin. Theory Ser. A*, 195:Paper No. 105700, 19, 2023.
- [109] H. Lu and X. Yu. On rainbow matchings for hypergraphs. *SIAM J. Discrete Math.*, 32(1):382–393, 2018.
- [110] H. Lu, X. Yu, and X. Yuan. Rainbow matchings for 3-uniform hypergraphs. *J. Combin. Theory Ser. A*, 183:Paper No. 105489, 21, 2021.
- [111] C. Magnant. Density of Gallai multigraphs. *Electron. J. Combin.*, 22(1):Paper 1.28, 6, 2015.
- [112] M. Matsumoto and N. Tokushige. The exact bound in the Erdős-Ko-Rado theorem for cross-intersecting families. *J. Combin. Theory Ser. A*, 52(1):90–97, 1989.
- [113] R. Montgomery. A proof of the Ryser-Brualdi-Stein conjecture for large even n . *arXiv preprint arXiv:2310.19779*, 2023.
- [114] R. Montgomery. Transversals in Latin squares. *arXiv preprint arXiv:2406.19873*, 2024.
- [115] R. Montgomery, A. Müyesser, and Y. Pehova. Transversal factors and spanning trees. *Adv. Comb.*, pages Paper No. 3, 25, 2022.
- [116] J. Moon and L. Moser. On Hamiltonian bipartite graphs. *Israel J. Math.*, 1:163–165, 1963.

- [117] O. Ore. Note on Hamilton circuits. *Amer. Math. Monthly*, 67:55, 1960.
- [118] A. Pokrovskiy. Rota’s basis conjecture holds asymptotically. *arXiv preprint arXiv:2008.06045*, 2020.
- [119] J. Polcyn, C. Reiher, V. Rödl, and B. Schülke. On Hamiltonian cycles in hypergraphs with dense link graphs. *J. Combin. Theory Ser. B*, 150:17–75, 2021.
- [120] L. Pósa. A theorem concerning Hamilton lines. *Magyar Tud. Akad. Mat. Kutató Int. Közl.*, 7:225–226, 1962.
- [121] V. Rödl, A. Ruciński, and E. Szemerédi. A Dirac-type theorem for 3-uniform hypergraphs. *Combin. Probab. Comput.*, 15(1-2):229–251, 2006.
- [122] V. Rödl, A. Ruciński, and E. Szemerédi. An approximate Dirac-type theorem for k -uniform hypergraphs. *Combinatorica*, 28(2):229–260, 2008.
- [123] V. Rödl, A. Ruciński, and E. Szemerédi. Dirac-type conditions for Hamiltonian paths and cycles in 3-uniform hypergraphs. *Adv. Math.*, 227(3):1225–1299, 2011.
- [124] H. J. Ryser. Neuere probleme der kombinatorik. *Vorträge über Kombinatorik, Oberwolfach*, 69(91):35, 1967.
- [125] P. D. Seymour. Problem section, problem 3, combinatorics. *London Math. Soc. Lecture Note Ser. (Proc. British Combinatorial Conf., Univ. Coll. Wales, Aberystwyth, 1973)*, pages 201–202, 1974.
- [126] S. K. Stein. Transversals of Latin squares and their generalizations. *Pacific J. Math.*, 59(2):567–575, 1975.
- [127] B. Sudakov. Robustness of graph properties. In *Surveys in combinatorics 2017*, volume 440 of *London Math. Soc. Lecture Note Ser.*, pages 372–408. Cambridge Univ. Press, Cambridge, 2017.
- [128] E. Szemerédi. Regular partitions of graphs. In *Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976)*, volume 260 of *Colloq. Internat. CNRS*, pages 399–401. CNRS, Paris, 1978.
- [129] Y. Tang, B. Wang, G. Wang, and G. Yan. Rainbow Hamilton cycle in hypergraph system. *arXiv preprint arXiv:2302.00080*, 2023.
- [130] A. Thomason. Paths and cycles in tournaments. *Trans. Amer. Math. Soc.*, 296(1):167–180, 1986.
- [131] A. Treglown. On directed versions of the Hajnal-Szemerédi theorem. *Combin. Probab. Comput.*, 24(6):873–928, 2015.
- [132] P. Turán. Eine Extremalaufgabe aus der Graphentheorie. *Mat. Fiz. Lapok*, 48:436–452, 1941.
- [133] I. M. Wanless. Transversals in Latin squares: a survey. In *Surveys in combinatorics*, volume 392 of *London Math. Soc. Lecture Note Ser.*, pages 403–437. Cambridge Univ. Press, Cambridge, 2011.
- [134] Y. Zhang and E. R. van Dam. Rainbow Hamiltonicity and the spectral radius. *arXiv preprint arXiv:2401.17845*, 2024.
- [135] Y. Zhao. Recent advances on Dirac-type problems for hypergraphs. In *Recent trends in combinatorics*, volume 159 of *IMA Vol. Math. Appl.*, pages 145–165. Springer, 2016.