REPRESENTATION STABILITY IN THE (CO)HOMOLOGY OF VERTICAL CONFIGURATION SPACES

DAVID BARON, URSHITA PAL, CHENGLU WANG, JENNIFER C. H. WILSON, AND CHUNYE YANG

ABSTRACT. In this paper, we study sequences of topological spaces called "vertical configuration spaces" of points in Euclidean space. We apply the theory of FI_G -modules, and results of Bianchi–Kranhold, to show that their (co)homology groups are *representation stable* with respect to natural actions of wreath products $S_k \wr S_n$. In particular, we show that in each (co)homological degree, the (co)homology groups (viewed as $S_k \wr S_n$ -representations) can be expressed as induced representations of a specific form. Consequently, the characters of their rational (co)homology groups, and the patterns of irreducible $S_k \wr S_n$ -representation constituents in these groups, stabilize in a strong sense. In addition, we give a new proof of rational (co)homological stability for unordered vertical configuration spaces, with an improved stable range.

Contents

1. Introduction	2
1.1. Classical (co)homological stability	2
1.2. Representation stability	2
1.3. Vertical configuration spaces	3
1.4. Representation stability for vertical configuration spaces, and its consequences	5
1.5. Overview of the paper	6
2. Irreducible representations of wreath products	6
2.1. The structure of a wreath product $G \wr H$	6
2.2. Conjugacy classes in $G \wr S_n$	7
2.3. The classification of irreducible $G \wr H$ -representations	8
2.4. The branching rule for $G \wr S_n$	9
3. The category FI_G and FI_G -modules	10
3.1. The category FI_G	10
3.2. FI_G -modules	12
3.3. An FI_G -module criterion	12
3.4. Generation of FI_G -modules and the Noetherian property	13
3.5. Representable and induced FI_G -modules	13
4. Representation stability	15
5. $\mathrm{FI}_G \sharp$ -modules	16
5.1. The structure theorem for $\operatorname{FI}_G \sharp$ -modules	17
6. Character polynomials for FI_G -modules	18
7. Representation stability for vertical configuration spaces	19
7.1. Bianchi–Kranhold's computation of the cohomology groups $H^*\left(\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})\right)$	19
7.2. The ranks of the (co)homology groups grow polynomially in n	20
7.3. The (co)homology groups of vertical configuration spaces form $\mathrm{Fl}_G \sharp$ -modules	21
7.4. The (co)homology of vertical configuration space is representation stable	23
7.5. The transfer map and classical (co)homological stability	24
References	27

Date: December 3, 2024.

1. INTRODUCTION

1.1. Classical (co)homological stability. A classical theme in algebraic topology is the study of (co)homological stability: a sequence of spaces with maps $(X_n \xrightarrow{\varphi_n} X_{n+1})_{n \ge 0}$ satisfies homologically stability with stable range $n \ge N_d$ if for all $n \ge N_d$, the map

$$H_d(X_n) \xrightarrow{(\varphi_n)_*} H_d(X_{n+1})$$

induced on degree-d homology is an isomorphism.

One well-studied example of this phenomenon is the families of unordered configuration spaces of points in open manifolds. Given a manifold M, we define the ordered configuration space of M on n points to be

$$\operatorname{Conf}_n(M) := \{ (x_1, \dots, x_n) \in M^n \mid x_i \neq x_j \text{ if } i \neq j \}$$

topologized as a subspace of M^n . The manifold M is called the *background manifold*. We may alternatively view $\operatorname{Conf}_n(M)$ as the space of embeddings of the discrete set $\{1, \ldots, n\}$ into M, and interpret an element of the space as a collection of n distinct points in M, "labelled" by elements of the set $\{1, \ldots, n\}$. The symmetric group S_n acts on $\operatorname{Conf}_n(M)$ by permuting the labels. The *unordered configuration space* of M on n points is the orbit space

$$\operatorname{UConf}_n(M) := \operatorname{Conf}_n(M)/S_n,$$

endowed with the quotient topology. An element of $\operatorname{UConf}_n(M)$ may be interpreted as a set of n distinct unlabelled points in M.

Arnol'd [Arn70a] proved homological stability of the unordered configuration spaces $\mathrm{UConf}_n(\mathbb{R}^2)$ with stable range $n \ge 2d$. McDuff [McD75, Theorem 1.2] generalized this theorem to any connected open background manifold M of dimension at least 2. Segal [Seg79, Appendix to §5] established a stable range for these families of $n \ge 2d$.

The ordered configuration spaces of these manifolds fail to exhibit homological stability, but Church–Farb [CF13] formalized a sense in which a sequence of S_n -representation may stabilize "up to the S_n action." They named this condition *representation stability*. Church [Chu12] and later Church–Ellenberg–Farb [CEF15] proved representation stability for the (co)homology of the families $(Conf_n(M))_n$.

1.2. Representation stability. Let $(V_n)_n$ be a sequence of rational S_n -representations, and let $\phi_n : V_n \to V_{n+1}$ be maps that are compatible with the S_n -actions in an appropriate sense (made precise in Definition 3.13). Church-Farb proposed the following notion of stability for the sequence $(\phi_n : V_n \to V_{n+1})_n$.

Definition 1.1 (Church–Farb [CF13, Definition 1.1]). The sequence $(\phi_n : V_n \to V_{n+1})_n$ is representation stable if, for sufficiently large n, the following conditions hold:

- (i) **Injectivity**: The maps $\phi_n : V_n \to V_{n+1}$ are injective.
- (ii) **Surjectivity**: The span of the S_{n+1} -orbit of $\phi_n(V_n)$ equals all of V_{n+1} .
- (iii) Multiplicity Stability: Decompose V_n into irreducible S_n -representations,

$$V_n = \bigoplus_{\lambda} c_{\lambda,n} V(\lambda)_n$$

with multiplicities $0 \leq c_{\lambda,n} \leq \infty$. For each λ , the multiplicities $c_{\lambda,n}$ are independent of n.

Rational representations of S_n are semisimple, and the irreducible representations are classified: they are in canonical bijection with partitions of n. Definition 1.1 depends on a certain (combinatorially natural) naming scheme for these irreducible representations that is uniform in n: for any partition λ of an integer, there is an associated sequence of irreducible S_n -representations $V(\lambda)_n$ that is defined for all n sufficiently large. This naming scheme is described in Section 4 as a special case of Definition 4.2. With this notation, significantly, Definition 1.1 states that for a representation stable sequence V_n of S_n -representations, the decomposition of each module V_n into irreducible subrepresentations is (for n sufficiently large) independent of n.

To illustrate these patterns, consider the degree-1 cohomology of $\operatorname{Conf}_n(\mathbb{R}^2)$. As S_n -representations, the cohomology groups decompose in the following ways, using the standard indexing of irreducible S_n -representations by Young diagrams of size n.



Once $n \ge 4$, we obtain the decomposition at level (n + 1) from that at level n by adding a single box to the top row of each Young diagram. Church–Farb express the stable decomposition as follows, by recording the portion of each Young diagram below its top row. This description holds for all $n \ge 4$.

 $H^1(\operatorname{Conf}_n(\mathbb{R}^2);\mathbb{Q}) \cong V(\emptyset)_n \oplus V(\Box)_n \oplus V(\Box)_n$

Church [Chu12] proves that the same pattern holds in every (co)homological degree for every family of spaces $\operatorname{Conf}_n(M)$, with connected open orientable background manifold M of dimension at least 2.

Church-Ellenberg-Farb further formalized the theory of representation stability using an algebraic construct they called an FI-module [CEF15, Definition 1.1]. They proved that if a sequence of S_n -representations has the structure of a "finitely generated FI-module"—a combinatorial condition that is easy to verify in many cases—then representation stability in the sense of Definition 1.1 follows, in addition to other structural consequences.

1.3. Vertical configuration spaces. Our paper focuses on a variant of configuration space of \mathbb{R}^s , called *vertical configuration space*, that arise from the theory of certain coloured operads (see Kranhold [Kra21]). These spaces were introduced by Herberz [Her14] and Rosner [Ros14] in their respective Bachelors' theses, and further studied by Latifi [Lat17], Kranhold [Kra21], and and Bianchi–Kranhold [BK22]. The following definitions are stated as in Bianchi–Kranhold [BK22].

Definition 1.2 (Bianchi–Kranhold [BK22, Section 1]). For $k \ge 1$, we say a *cluster* of size k in \mathbb{R}^s is a k-tuple of k distinct points of \mathbb{R}^s . Let $\mathbb{R}^s = \mathbb{R}^p \times \mathbb{R}^q$, and consider the projection map of the first p coordinates, $pr : \mathbb{R}^s \longrightarrow \mathbb{R}^p$. We say a cluster $z = (z^1, \ldots, z^k)$ of k points in \mathbb{R}^s is *vertical* if $pr(z^1) = \cdots = pr(z^k)$, i.e. the k points in the cluster share their first p coordinates.

Notation 1.3. For an *r*-tuple $K = (k_1, \ldots, k_r)$ of positive integers, write $|K| = k_1 + \cdots + k_r$. Let $\operatorname{Conf}_K(\mathbb{R}^s)$ denote the space (canonically isomorphic to $\operatorname{Conf}_{|K|}(\mathbb{R}^s)$) of *r*-tuples $(z_1, \ldots, z_r) \in (\mathbb{R}^s)^{k_1} \times \cdots \times (\mathbb{R}^s)^{k_r}$ of *r* ordered, pairwise disjoint clusters $z_i = (z_i^1, \ldots, z_i^{k_i})$.

In other words, $\operatorname{Conf}_K(\mathbb{R}^s)$ is the space $\operatorname{Conf}_{|K|}(\mathbb{R}^s)$ with the additional data of a partition of the set of labels into r parts of sizes k_1, \ldots, k_r . An element in $\operatorname{Conf}_K(\mathbb{R}^s)$ is a collection of |K| distinct labelled points in \mathbb{R}^s , correspondingly partitioned into r clusters.

Definition 1.4 (Bianchi–Kranhold [BK22, Definition 1.1]). Fix s = p + q. Let $r \ge 0$ and $K = (k_1, \ldots, k_r)$ be an ordered tuple of positive integers. Define the *vertical configuration space*, denoted $\tilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$, to be the following subspace of $\operatorname{Conf}_K(\mathbb{R}^s)$. An *r*-tuple of clusters (z_1, \ldots, z_r) belongs to $\tilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ if and only if each cluster $z_i = (z_i^1, \ldots, z_i^{k_i})$ is vertical in the sense of Definition 1.2.

Again, we may view elements of $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ as collections of |K| distinct labelled points in \mathbb{R}^{p+q} , as illustrated in Figures 1 and 2.

In general the full symmetric group does not act on the vertical configuration space by permuting the labels, since arbitrary permutations may not respect the verticality conditions imposed on each cluster. Instead, if we fix $K = (k_1, \ldots, k_r)$ and let r(k) denote the number of occurrences of k in K, then the product of wreath products $\prod_{k\geq 1} S_k \wr S_{r(k)}$ acts freely on $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ by permuting clusters of the same size and labels

```
3,2•
•2,3
1,3••2,1
3,1•
•2,2
1,1•
•2,4
1,2•
```

FIGURE 1. An element of $\tilde{\mathcal{V}}_{(3,4,2)}(\mathbb{R}^{1,1})$. The 9 points are partitioned into three clusters of 3, 4, and 2 points, respectively. Points in each cluster are subject to a colinearity condition. The case q = 1 is the origin of the name 'vertical' configuration space. Figure adapted from Bianchi–Kranhold [BK22, Figure 1].



FIGURE 2. An element of $\tilde{\mathcal{V}}_{(3,4,3)}(\mathbb{R}^{1,2})$. The 10 points are partitioned into three clusters of 3, 4, and 3 points, respectively. Points in each cluster are subject to a coplanarity condition. Figure adapted from Bianchi–Kranhold [BK22, Figure 3].

on points within each cluster. Analogous to UConf, we define the *unordered* vertical configuration space as the quotient space under this action.

Definition 1.5. Fix $p, q, r \ge 0$, and let $K = (k_1, \ldots, k_r)$ be an ordered tuple of positive integers. The unordered vertical configuration space $\mathcal{V}_K(\mathbb{R}^{p,q})$ is defined to be orbit space

$$\mathcal{V}_K(\mathbb{R}^{p,q}) := \widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q}) / \prod_{k \ge 1} S_k \wr S_{r(k)}.$$

We may view an element of $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ as |K| distinct unlabelled points in \mathbb{R}^{p+q} , partitioned into k_1, k_2, \ldots, k_r unlabelled clusters, as shown in Figure 3. Each cluster must be *vertical* in the sense that the points in a given cluster agree in their first p coordinates.



FIGURE 3. An element of $\mathcal{V}_{(2,2,2,2,2)}(\mathbb{R}^{1,1})$. The 10 points in \mathbb{R}^2 are partitioned into 5 unlabelled "vertical clusters" of 2 unlabelled points each.

We are particularly interested in the case where K is the n-tuple K = (k, ..., k) for some $k \ge 0$. In this case we write $\tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$ for $\tilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ and $\mathcal{V}_n^k(\mathbb{R}^{p,q})$ for $\mathcal{V}_K(\mathbb{R}^{p,q})$. The wreath product $S_k \wr S_n$ acts freely on the space.

Latifi [Lat17] proves homological stability for the families $(\mathcal{V}_n^k(\mathbb{R}^{1,2}))_n$ for fixed $k \ge 1$. In fact, as pointed out by Bianchi–Kranhold [BK22], Latifi's proof applies to any p, q with $p + q \ge 3$. Bianchi–Kranhold [BK22] compute the cohomology groups of the ordered vertical configuration spaces, and extended the homological stability result for the unordered vertical configuration space for all $p \ge 0, q \ge 1$, with $(p,q) \ne (0,1)$. We review the results of Bianchi–Kranhold and Latifi in Sections 7.1 and 7.5.

1.4. Representation stability for vertical configuration spaces, and its consequences. Given the homological stability result for the unordered vertical configuration spaces, it is natural to ask: does the ordered vertical configuration space satisfy representation stability, generalizing the behaviour of $\operatorname{Conf}_n(\mathbb{R}^s)$ and $\operatorname{UConf}_n(\mathbb{R}^s)$? Our project resolves this question in the affirmative for $p \ge 0$ and $q \ge 2$.

To answer this question, we apply the machinery of FI_G -modules and FI_G #-modules, as developed by Sam–Snowden [SS19], Gan–Li [GL15], Ramos [Ram18], and Casto [Cas16]. This machinery generalizes the work of Church–Ellenberg–Farb to sequences of representations of wreath products $G \wr S_n$ for G a fixed finite group. The following is our main theorem.

Theorem 7.24. Let $p \ge 0$ and $q \ge 2$. Fix $k \ge 1$ and (co)homological degree $d \ge 0$.

- Let R be a commutative ring. The sequences $\left(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_n$ and $\left(H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_n$ of $S_k \wr S_n$ representations are representation stable in the sense that they are $\operatorname{FI}_{S_k} \sharp$ -modules finitely generated
 in degree at most $\left\lfloor \frac{2d}{q-1} \right\rfloor$.
- The sequences $\left(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})\right)_n$ and $\left(H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})\right)_n$ of rational $S_k \wr S_n$ -representations are representation stable in the sense of Definition 4.4, with stable range $n \ge \left\lfloor \frac{4d}{a-1} \right\rfloor$.

The theory of $\operatorname{FI}_{S_k} \sharp$ -modules implies, as a consequence, that in each (co)homological degree, the sequence of $S_k \wr S_n$ -representations $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$ (respectively, $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$) has a highly constrained form: it can be expressed as certain induced representations, as described in Corollary 7.25 below. Heuristically, this gives a concrete sense in which, up to the action of the wreath product $S_k \wr S_n$, all the degree-d (co)homology classes "come from" the spaces $\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$ for $n \leq \left\lfloor \frac{2d}{q-1} \right\rfloor$.

Corollary 7.25. For $i = 0, \ldots \lfloor \frac{2d}{q-1} \rfloor$, there exist $S_k \wr S_i$ -representations U_i such that, as a $S_k \wr S_n$ -representations,

$$H^{d}(\widetilde{\mathcal{V}}_{n}^{k}(\mathbb{R}^{p,q});R) \cong \bigoplus_{i=0}^{\left\lfloor\frac{2d}{q-1}\right\rfloor} \operatorname{Ind}_{(S_{k}\wr S_{i})\times(S_{k}\wr S_{n-i})}^{S_{k}\wr S_{n}}(U_{i}\boxtimes R) \quad \text{for all } n$$

The same statement holds for $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$.

The structure of these induced representations implies that, when we take coefficients $R = \mathbb{Q}$, the patterns of irreducible representations stabilize, in a sense analogous to Definition 1.1. The classification of irreducible representations of wreath products $S_k \wr S_n$ is reviewed in Section 2.3 (see Corollary 2.16). Notions of stability for sequences of $S_k \wr S_n$ -representations are described in Section 4.

Corollary 7.26. Fix $d \ge 0$. The decomposition of the $S_k \wr S_n$ -representation $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ into irreducible components $L(\underline{\lambda})_n$ stabilizes in n in the sense of Definition 4.4, with stable range $n \ge \left\lfloor \frac{4d}{q-1} \right\rfloor$. The same statement holds for $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$.

The description of the (co)homology groups as induced $S_k \wr S_n$ -representations in Corollary 7.25 also has implications for the sequence of characters of these representations. Specifically, for each (co)homological degree d, the sequence of characters is equal to a *character polynomial* (defined in Section 6) that is independent of n. **Corollary 7.27.** Fix $d \ge 0$. The sequence of characters of the $S_k \wr S_n$ -representations $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ can be expressed as a single character polynomial in the coloured cycle-counting functions

$$P^{H^{a}(\mathcal{V}_{n}^{\kappa}(\mathbb{R}^{p,q}))} \in \mathbb{Q}[X_{r}^{C_{i}} \mid r \geq 1, C_{i} \text{ a conjugacy class of } S_{k}]$$

of degree at most $\left\lfloor \frac{2d}{q-1} \right\rfloor$, independent of n. The same statement holds for the $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$.

Representation stability for the $S_k \wr S_n$ -representations $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ and $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ means in particular that $S_k \wr S_n$ -coinvariants of these representations stabilize for all *n* sufficiently large. In Section 7.5, we explain why this implies classical stability for the (co)homology of the *unordered* vertical configuration spaces $\mathcal{V}_n^k(\mathbb{R}^{p,q})$. We deduce a new proof of rational cohomological stability for these spaces. Although—in contrast to the work of Latifi [Lat17] and Bianchi–Kranhold [BK22]—our results only hold with rational coefficients, in the rational case we improve the stable range. In Corollary 7.34, we prove stability with respect to stabilization maps $\phi_n : \mathcal{V}_n^k(\mathbb{R}^{p,q}) \to \mathcal{V}_{n+1}^k(\mathbb{R}^{p,q})$ defined in Section 7.5.

Corollary 7.34. Fix $p \ge 0$, $q \ge 2$, and (co)homological degree d. The (co)homology of the unordered vertical configuration spaces, $(H_d(\mathcal{V}_n^k(\mathbb{R}^{p,q});\mathbb{Q}))_n$ and $(H^d(\mathcal{V}_n^k(\mathbb{R}^{p,q});\mathbb{Q}))_n$, stabilize. Specifically, the maps on (co)homology induced by the stabilization maps ϕ_n are always injective, and are isomorphisms for $n \ge \left\lfloor \frac{2d}{q-1} \right\rfloor$.

1.5. Overview of the paper. In Section 2 we review the structure of wreath products and their representation theory. In Sections 3, 4, 5, and 6, we present the theory of FI_G -modules, FI_G #-modules, and their connection to representation stability. In Section 7, we apply this theory to prove the main theorems of the paper.

Acknowledgments. Jennifer Wilson is grateful for the support of the NSF CAREER grant DMS-2142709. David Baron, Chenglu Wang, and Chunye Yang thank the NSF and the University of Michigan for the support of their REU program ("Research Experience for Undergraduates"). David Baron was funded by NSF grant DMS-2142709. Chenglu Wang and Chunye Yang were funded by the University of Michigan. Urshita Pal is grateful for a Research Assistantship from the University of Michigan to support her work on the project and mentorship role in the REU.

The authors thank Gahl Shemy for her mentorship during the REU. They thank Nir Gadish for helpful conversations.

2. IRREDUCIBLE REPRESENTATIONS OF WREATH PRODUCTS

In this section we review the definition and representation theory of wreath products, as it is developed in James–Kerber [JK81, Chapter 4].

2.1. The structure of a wreath product $G \wr H$. Suppose that X is a set with a faithful, transitive action by a group G. For $n \in \mathbb{Z}_{\geq 1}$ let $[n] = \{1, 2, ..., n\}$ and $[0] = \emptyset$. Consider the product $[n] \times X$, which we may interpret as the disjoint union of n (labelled) copies of X. The product G^n acts on $[n] \times X$, with the *i*th factor G of G^n acting on $\{i\} \times X$ and fixing all other copies of X pointwise. Any fixed subgroup H of the symmetric group S_n acts on $[n] \times X$ by acting on [n] by permutations and fixing X pointwise; it permutes the n copies of X. These two actions typically do not commute, but generate a group of symmetries of $[n] \times X$ called the *wreath product* of G by H and denoted $G \wr H$. In the following definition, we give an algebraic characterization of $G \wr H$.

Definition 2.1. Let G be a group and H a subgroup of the symmetric group S_n . We identify the *n*-fold Cartesian product G^n with the set of functions from $[n] = \{1, ..., n\}$ into G:

$$G^n = \{ \alpha \mid \alpha : [n] \longrightarrow G \}.$$

The group S_n acts on [n] by its standard action by permutations, and acts on G^n by precomposition,

$$\pi: G^n \longrightarrow G^n$$
$$\alpha \longmapsto \alpha \circ \pi^{-1}$$

We define the wreath product of G by H, denoted $G \wr H$, to be the group with underlying set

$$G^n \times H = \{ (\alpha, \pi) \mid \alpha \in G^n, \pi \in H \}$$

and group operation

$$(\alpha, \pi)(\alpha', \pi') = (\alpha \cdot \alpha'_{\pi}, \pi\pi')$$

where $(\alpha \cdot \alpha')(i)$ denotes the pointwise product $\alpha(i)\alpha'(i)$ with respect to the group operation on G, $\pi\pi'$ is the product as elements of H, and $\alpha'_{\pi} = \alpha' \circ \pi^{-1}$.

An element (α, π) of $G \wr H$ has inverse $(\bar{\alpha} \circ \pi, \pi^{-1})$, where $\bar{\alpha}$ is the pointwise inverse of α , with $\bar{\alpha}(i) = \alpha(i)^{-1}$.

Remark 2.2. An alternate way to model the group $G \wr H$ is via *monomial matrices*. Realize the group $H \subseteq S_n$ as a subgroup of $n \times n$ permutation matrices. Consider the set of all matrices obtained by replacing each of the 1's in these permutation matrices by an element of G. Then the group $G \wr H$ consists of these matrices, with the group product defined by matrix multiplication and the group operation on G.

We introduce notation for the distinguished subgroups of $G \wr H$ isomorphic to G^n and H. With this notation, we state Lemma 2.4 on the basic structure of the wreath product.

Definition 2.3. Let G, H be as in Definition 2.1. Let 1_H denote the identity permutation in H. Let e denote the identity element of G^n , the constant function at the identity 1_G of G. The base group G^* of $G \wr H$ is the subgroup

$$G^* = \{ (\alpha, 1_H) \mid \alpha \in G^n \}$$

isomorphic to G^n . The *complement* of G^* is the subgroup

$$H' = \{(e,\pi) \mid \pi \in H\}$$

isomorphic to H.

Lemma 2.4 (James-Kerber [JK81, 4.1.15]). Let G, H be as in Definition 2.1. Then,

- (i) As a set $G \wr H$ is equal to the product of subsets $G^* \cdot H' = \{(\alpha, 1_H)(e, \pi) \mid \alpha \in G^n, \pi \in H\}$.
- (ii) G^* is a normal subgroup of $G \wr H$
- (iii) $G^* \cap H'$ is the trivial subgroup $\{(e, 1_H)\}$.

2.2. Conjugacy classes in $G \wr S_n$. To study characters and character polynomials of the wreath product $G \wr S_n$, we review the classification of its conjugacy classes. Throughout this section, we assume that G has countably many conjugacy classes C^1, C^2, \ldots .

Notation 2.5. Fix an element $\pi \in S_n$. If the cycle type of π has $c = c(\pi)$ cycles of lengths ℓ_1, \ldots, ℓ_c , then we can uniquely express π in cycle notation

$$(j_1 \pi(j_1) \cdots \pi^{\ell_1 - 1}(j_1)) (j_2 \pi(j_2) \cdots \pi^{\ell_2 - 1}(j_2)) \cdots (j_c \pi(j_c) \cdots \pi^{\ell_c - 1}(j_c))$$

by imposing the conditions that $j_{\nu} \leq \pi^{i}(j_{\nu})$ for all i, ν , and $j_{1} \leq j_{2} \leq \cdots \leq j_{c}$.

Definition 2.6 (James–Kerber [JK81, 4.2.1]). Let $\pi \in S_n$ be written in cycle notation as in Notation 2.5 and let $(\alpha, \pi) \in G \wr S_n$. Then, to the ν th cyclic factor $(j_{\nu}\pi(j_{\nu})\dots\pi(j_{\nu})^{l_{\nu}-1})$, we associate the element of G

$$g_{\nu}(\alpha,\pi) = \alpha(j_{\nu})\alpha(\pi^{-1}(j_{\nu}))\cdots\alpha(\pi^{-l_{\nu}+1}(j_{\nu})))$$

which we call the ν th cycle product of (α, π) .

James–Kerber show that the conjugacy class of (α, π) in $G \wr S_n$ is indexed by the data of the lengths ℓ_{ν} of each cycle of π , along with the conjugacy class of its cycle product $g_{\nu}(\alpha, \pi)$.

Definition 2.7 (James–Kerber [JK81, 4.2.2]). For $(\alpha, \pi) \in G \wr S_n$, let $a_{ik}(\alpha, \pi)$ be the number of cycles of length k of π for which the associated cycle product $g_{\nu}(\alpha, \pi)$ belongs to the conjugacy class C^i of G. Then, the *type* of (α, π) is the matrix

$$a(\alpha,\pi) = \left[a_{ik}(\alpha,\pi)\right].$$

Theorem 2.8 (James–Kerber [JK81, 4.2.8]). Two elements (α, π) and (α', π') of $G \wr S_n$ are conjugate if and only if $a(\alpha, \pi) = a(\alpha', \pi')$.

Concretely, the conjugacy class of an element of $G \wr S_n$ is indexed by

- a cycle type, that is, an (unordered) partition of n,
- for each cycle, a corresponding conjugacy class of G.

Remark 2.9. Note that the numbers $a_{ik}(\alpha, \pi)$ satisfy

$$\sum_{i,k} ka_{ik}(\alpha,\pi) = n$$

. Conversely, given a matrix with nonnegative entries satisfying the above equation, it is not hard to construct an element of $G \wr S_n$ whose type is given by that matrix.

Corollary 2.10 (James–Kerber [JK81, 4.2.9]). Suppose G has $s \in \mathbb{Z}_{\geq 0}$ conjugacy classes and p(m) denotes the number of partitions of m. Then the number of conjugacy classes of $G \wr S_n$ is equal to

$$\sum_{(n)} p(n_1) \cdots p(n_s)$$

where the sum is taken over all s-tuples (n_1, \ldots, n_s) such that $\sum_{i=1}^s n_i = n$.

2.3. The classification of irreducible $G \wr H$ -representations. In this paper we consider representations of wreath products of a finite group G with the full symmetric group S_n . For the remainder of the discussion, we assume that G is a finite group and that \mathbb{K} is a splitting field for G of characteristic zero. We will now review the classification of irreducible representations of $G \wr H$ over \mathbb{K} .

Notation 2.11. Given an *n*-tuple of representations D_1, \ldots, D_n of G over \mathbb{K} , we use the notation $D_1 \boxtimes D_2 \boxtimes \cdots \boxtimes D_n$ to denote the representation of G^n obtained as the *external* tensor product of the representations D_i ; this is the \mathbb{K} -vector space $D_1 \otimes_{\mathbb{K}} D_2 \otimes_{\mathbb{K}} \cdots \otimes_{\mathbb{K}} D_n$ with action of $\alpha \in G^n$ defined by

 $\alpha \cdot (d_1 \otimes d_2 \otimes \cdots \otimes d_n) = (\alpha(1) \cdot d_1) \otimes (\alpha(2) \cdot d_2) \otimes \cdots \otimes (\alpha(n) \cdot d_n) \quad \text{for } d_1 \otimes \cdots \otimes d_n \in D_1 \otimes_{\mathbb{K}} D_2 \otimes_{\mathbb{K}} \cdots \otimes_{\mathbb{K}} D_n.$

In contrast, the *internal* tensor product $U_1 \otimes U_2$ of representations of a group G' is the tensor product $U_1 \otimes_{\mathbb{K}} U_2$ equipped with the diagonal action of G',

 $g \cdot (u_1 \otimes u_2) = (g \cdot u_1) \otimes (g \cdot u_2)$ for $g \in G$ and $u_1 \otimes u_2 \in U_1 \otimes_{\mathbb{K}} U_2$.

Let $\{D^1, \ldots, D^r\}$ be a complete list of inequivalent irreducible representations of G. A standard result from the representation theory of finite groups (see for example Fulton-Harris [FH13, Exercise 2.36]) states that the irreducible representations of the finite product $G^* \cong G^n$ have the form

$$D_1 \boxtimes D_2 \boxtimes \cdots \boxtimes D_n$$
, where $D_i \in \{D^1, \dots, D^r\}$ for all $i \in \{1, \dots, n\}$.

These r^n representations are irreducible, pairwise inequivalent, and constitute a complete list of irreducible representations of G^n .

Definition 2.12 (James–Kerber [JK81, Section 4.3]). Let G be a finite group and H a subgroup of S_n . Fix an irreducible representation $D^* = D_1 \boxtimes D_2 \boxtimes \cdots \boxtimes D_n$ of G^* . The type of D^* is the r-tuple $\mu = (n_1, \ldots, n_r)$, where n_i denotes the number of factors isomorphic to D^i in D^* . Let $S_{\mu} \cong S_{n_1} \times \cdots \times S_{n_r}$ be the subgroup of S_n that stabilizes the partition of the index set [n] of D^* into the n_1 indices i corresponding to factors D_i equal to D^1 , the n_2 indices i corresponding to factors D_i equal to D^2 , etc. Let H_{D*} denote $H \cap S_{\mu}$. Let $H'_{D*} \cong H_{D*}$ denote the corresponding subgroup of $G \wr H$,

$$H'_{D*} = \{ (e, \pi) \mid \pi \in H_{D*} \} \subseteq H' \subseteq G \wr H.$$

The *inertia subgroup* of D^* is the subgroup $G^* \cdot H'_{D^*}$ of $G \wr H$. The inertia subgroup is isomorphic to the wreath product of G by H_{D^*} , and we denote it by $G \wr H_{D^*}$.

Definition 2.13. The *conjugate* of the G^* -representation D^* by $(\alpha, \pi) \in G \wr H$ is the G^* -representation obtained by pulling back the representation along the group homomorphism $G^* \to G^*$ given by conjugation by (α, π) . Two G^* -representations are *conjugate* if one is conjugate to the other via some $(\alpha, \pi) \in G \wr H$.

James–Kerber characterize the inertia subgroup $G \wr H_{D*}$ as the set of elements (α, π) in $G \wr H$ for which the conjugate of D^* by (α, π) is isomorphic to D^* as a G^* -representation: it is the stabilizer of D^* with respect to the action of $G \wr H$ by conjugation on the set of irreducible G^* -representations.

With these definitions, James–Kerber [JK81] use Clifford Theory [Cli37] to classify the irreducible representations of $G \wr H$. **Theorem 2.14** (James–Kerber [JK81, Theorem 4.3.34]). Let G be a finite group and H a subgroup of S_n . We take coefficients in a splitting field \mathbb{K} for G of characteristic zero. Fix an irreducible representation $D^* = D_1 \boxtimes D_2 \boxtimes \cdots \boxtimes D_n$ of G^* . The action of G^* on $D_1 \boxtimes D_2 \boxtimes \cdots \boxtimes D_n$ extends to an action of the inertia subgroup $G \wr H_{D^*}$, where

$$(\alpha,\pi)\cdot(d_1\otimes d_2\otimes\cdots\otimes d_n)=(\alpha(1)\cdot d_{\pi^{-1}(1)})\otimes(\alpha(2)\cdot d_{\pi^{-1}(2)})\otimes\cdots\otimes(\alpha(n)\cdot d_{\pi^{-1}(n)})$$

for $(\alpha, \pi) \in G \wr H_{D*}$ and $d_1 \otimes \cdots \otimes d_n \in D_1 \boxtimes \cdots \boxtimes D_n$. We denote this $G \wr H_{D*}$ -representation by D^* .

Let D' be an irreducible representation of H_{D^*} . We let \widetilde{D}' denote the representation of $G \wr H_{D^*}$ pulled back from D' along the quotient map

$$G \wr H_{D*} \longrightarrow H_{D*}$$
$$(\alpha, \pi) \longmapsto \pi.$$

Let $\widetilde{D}^* \otimes \widetilde{D}'$ denote the internal tensor product of these $G \wr H_{D^*}$ -representations. We write D for the induced representation

$$D = \operatorname{Ind}_{G \wr H_{D^*}}^{G \wr H} \widetilde{D}^* \otimes \widetilde{D}'.$$

Then,

- (i) D is irreducible and every irreducible representation of $G \wr H$ over \mathbb{K} has this form.
- (ii) D runs through a complete system of pairwise inequivalent and irreducible representations of $G \wr H$ if D^* runs through a complete system of pairwise non-conjugate, irreducible representations of G^* , and, while D^* remains fixed, D' runs through a complete system of pairwise inequivalent irreducible representations of H_{D^*} .

We remark that when $H = S_n$, two G^* -representations are conjugate if and only if they have the same type μ [JK81, Section 4.4]. In this case, their inertia subgroup S_{μ} is a product of symmetric groups. The irreducible representations of the symmetric group S_N are in canonical bijection with partitions of N (see, for example, Fulton-Harris [FH13, Chapter 4]). Thus, when $H = S_n$, we can specialize Theorem 2.14 as follows.

Corollary 2.15. Let χ_1, \ldots, χ_r be a complete list of irreducible representations of a finite group G over a splitting field \mathbb{K} of G of characteristic zero. Then the irreducible representations of $G \wr S_n$ are classified by partition-valued functions $\underline{\lambda}$ from $\{\chi_1, \ldots, \chi_r\}$ to partitions of nonnegative numbers, such that the sum $\sum_{i=1}^r |\underline{\lambda}(\chi_i)|$ is n.

The functions $\underline{\lambda}$ are sometimes called *multipartitions of n with r components*.

When $G = S_k$, we can further specialize the theorem. We note, in this case, that every irreducible representation of $S_k \wr S_n$ is defined over the rational numbers (see James–Kerber [JK81, Theorem 4.4.8 and Corollary 4.4.9]).

Corollary 2.16. Let \mathbb{K} be \mathbb{Q} (or any field of characteristic zero). Then the irreducible representations of $S_k \wr S_n$ over \mathbb{K} are in canonical bijection with functions $\underline{\lambda}$ from the set of partitions ρ of k to the set of all partitions, such that $\sum_{\rho} |\underline{\lambda}(\rho)| = n$. Explicitly, if V_{ρ} denotes the S_k -representation associated to ρ , then (in the notation of Theorem 2.14) the function $\underline{\lambda}$ corresponds to the irreducible $S_k \wr S_n$ -representation

$$\operatorname{Ind}_{S_k\wr (\prod_{\rho} S_{n_{\rho}})}^{S_k\wr S_n} \left(\underbrace{\boxtimes V_{\rho}^{\boxtimes n_{\rho}}}_{\rho} \right) \otimes \left(\underbrace{\boxtimes V_{\underline{\lambda}(\rho)}}_{\rho} \right)$$

where n_{ρ} denotes the size $|\underline{\lambda}(\rho)|$ of the partition $\underline{\lambda}(\rho)$.

2.4. The branching rule for $G \wr S_n$. Fix a finite group G. From the construction of the irreducible representations of $G \wr S_n$ and the Littlewood–Richardson rule for the symmetric groups, we can describe the structure of certain induced representations of wreath products. See, for example, Stein [Ste17, Theorem 4.5], or Ingram–Jing–Stitzinger [IJS08, Theorem 4.3].

Theorem 2.17 (E.g. [Ste17, Theorem 4.5])). Let χ_1, \ldots, χ_r be a complete list of the *r* irreducible representations of a finite group *G* over a splitting field \mathbb{K} of *G* of characteristic zero.

Fix integers $0 \leq d \leq n$. Let $\underline{\delta} = (\delta_1, \dots, \delta_r)$ be a multipartition of d with r components, and let $\underline{\rho} = (\rho_1, \dots, \rho_r)$ be a multipartition of (n - d) of r components. Let $W_{\underline{\delta}}$ and $W_{\underline{\rho}}$ denote the corresponding irreducible representations of $G \wr S_d$ and $G \wr S_{n-d}$, respectively.

$$\operatorname{Ind}_{G\wr S_d \times G\wr S_{n-d}}^{G\wr S_d} W_{\underline{\delta}} \boxtimes W_{\underline{\rho}} \cong \bigoplus_{\underline{\lambda}} \left(\prod_{i=1}^r C_{\rho_i,\delta_i}^{\lambda_i} \right) W_{\underline{\lambda}}$$

Here, the sum is taken over all multipartitions $\underline{\lambda}$ of n with r components satisfying $|\lambda_i| = |\rho_i| + |\delta_i|$ for $i = 1, \ldots, r$. The constants $C_{\rho_i, \delta_i}^{\lambda_i}$ are the Littlewood–Richardson coefficients.

To obtain an analogue of Pieri's rule, we specialize to the case that $W_{\underline{\rho}}$ is the trivial $G \wr S_{n-d}$ -representation. We let χ_1 denote the trivial G-representation, so $W_{\underline{\rho}}$ corresponds to the multipartition $\underline{\rho} = ((n-d), \emptyset, \emptyset, \dots, \emptyset)$. We deduce the following.

Corollary 2.18. Let G be a finite group. Let χ_1, \ldots, χ_r be a complete list of the r irreducible representations of G over a splitting field K of G of characteristic zero, with χ_1 the trivial representation. Let $\underline{\delta} = (\delta_1, \ldots, \delta_r)$ be a multipartition of d with r components, and let $W_{\underline{\delta}}$ denote the corresponding irreducible representations of $G \wr S_d$. Then,

$$\begin{aligned} \operatorname{Ind}_{G\wr S_d}^{G\wr S_d} &\cong \bigoplus_{\substack{\text{partitions } \lambda_1 \text{ of } \\ |\delta_1| + (n-d)}} C_{\delta_1,(n-d)}^{\lambda_1} W_{(\lambda_1,\delta_2,\delta_3,\ldots,\delta_r)} \\ &\cong \bigoplus_{\substack{\text{Young diagrams } \lambda_1 \text{ obtained} \\ \text{from } \delta_1 \text{ by adding } (n-d) \\ \text{boxes in distinct columns}}} W_{(\lambda_1,\delta_2,\delta_3,\ldots,\delta_r)}. \end{aligned}$$

3. The category FI_G and FI_G -modules

The field of *representation stability* was developed by Church, Ellenberg, Farb, and Nagpal [CF13, CEF15, CEFN14, CE17], Putman, Sam, and Snowden [Sno13, Put15, SS15, SS17, PS17], Gan and Li [GL15, GL19], and many others. See [CE15] for some additional references and, for example, [SS12, Far14, Wil18, Sam17, Sno19, Sam20, JRW22] for surveys and notes.

Church-Ellenberg-Farb developed the theory of FI-modules as a formal categorical framework for studying sequences of representations of the symmetric groups. The foundations of this theory are developed in their paper [CEF15]. In this section, we describe a generalization of this framework to sequences of representations of wreath products $G \wr S_n$, for a fixed finite group G. This generalization has been developed by Sam–Snowden [SS19], Gan–Li [GL15], Ramos [Ram18], Casto [Cas16], and others. The special case of $G = S_2$ was studied in earlier work [CF13, Wil12, Wil14, Wil15], in part because $S_2 \wr S_n$ is isomorphic to the Weyl group of type B_n/C_n .

In the case that G is the trivial group, then FI_G is canonically isomorphic to the category FI, and the results here specialize to the results of Church–Ellenberg–Farb.

In Sections 3, 4, 5, and 6 we review the theory of representation stability for wreath products, the tools we will use to prove it, and some combinatorial consequences for the structure of the representations and their characters. We do not claim originality for this theory—credit is due to Sam–Snowden [SS19], Gan–Li [GL15], Ramos [Ram18], and Casto [Cas16]—and many results closely parallel those of Church–Ellenberg– Farb [CEF15] for FI-modules. We do review the material in detail, and establish versions of some results that are implicit in the literature but not readily citable (such as Propositions 3.14 and 5.6). We use a novel combinatorial characterization of the categories FI_G and $FI_G \sharp$, which we hope the reader may find conceptually or computationally useful.

3.1. The category FI_G .

Definition 3.1. Let G be a finite group. Let X denote the underlying set of G, viewed as a faithful, transitive G-set with respect to the action of G by right multiplication. We define a category FI_G as follows. The objects are products $A \times X$ of a finite set A with the set X. The morphisms are all injective maps of the form $f: A \times X \to B \times X$ such that if $f(a, 1_G) = (b, g)$, then f(a, g') = (b, gg') for all $a \in A$ and $g' \in G$.

We may view the object $A \times X$ as the disjoint union of |A| copies of X, each labelled by an element of A. A morphism $A \times X \to B \times X$, then, is an injective map $\phi : A \to B$ on label sets, and on each copy $\{a\} \times X$ of X a G-equivariant map to $\{\phi(a)\} \times X$.

We can therefore specify the data of an FI_G morphism by an injective map $\phi : A \to B$ and a map $\alpha : A \to G$ that determines the values of the points $A \times \{1_G\}$. Then the pair (α, ϕ) corresponds to the injective map

$$A \times X \longrightarrow B \times X$$
$$(a,g) \longmapsto (\phi(a), \alpha(a)g)$$

This observation implies the following proposition, which reconciles Definition 3.1 with the definition of FI_G used by Sam–Snowden [SS19, Section 1.1], Gan–Li [GL15, Definition 1.1], Casto [Cas16], and Ramos [Ram18, Section 1].

Proposition 3.2. Let G be a finite group. The objects of FI_G are indexed by finite sets A, and the morphisms from A to B correspond precisely to the set of pairs (α, ϕ) consisting of an injective map $\phi : A \to B$ and an arbitrary map $\alpha : A \to G$, subject to the composition rule

$$(\alpha, \phi)(\alpha', \phi') = ((\alpha \circ \phi') \cdot \alpha', \phi \circ \phi')$$

where \cdot denotes the pointwise product with respect to the group operation on G.

For the purposes of the rest of the paper, we take the endomorphisms of the finite set $[n] \times X$ as the definition of the wreath product $G \wr S_n$. We observe that it is isomorphic to the group described in Definition 2.1.

Remark 3.3. Fix $k \in \mathbb{N}$. In the case that G is the symmetric group S_k , we may replace X with the faithful, transitive S_k -set $[k] = \{1, 2, ..., k\}$ equipped with the standard action of S_k by permutations. We obtain the same category FI_{S_k} as in Definition 3.1.

Remark 3.4. To study functors with domain FI_G , it suffices to consider the objects of the form $[n] \times X$ for natural numbers $n \ge 0$. Thus we sometimes restrict to the full subcategory of FI_G (equivalent to FI_G) with objects $[n] \times X$ indexed by the natural numbers. By abuse of notation, some authors also denote this subcategory by FI_G .

Notation 3.5. For brevity, for $n \in \mathbb{Z}_{\geq 0}$, we will sometimes write \mathbf{n}_X to denote the set $[n] \times X$. We similarly write \mathbf{d}_X , \mathbf{k}_X , \mathbf{m}_X , ..., and $\mathbf{0}_X$, $\mathbf{1}_X$,

Notation 3.6. Given $m, n \ge 0$, let $\iota_{m,n} : \mathbf{m}_X \to \mathbf{n}_X$ denote the natural inclusion map

$$\iota_{m,n} = (\iota \times id_X) : [m] \times X \longrightarrow [n] \times X$$

for $\iota : [m] \hookrightarrow [n]$ the inclusion. Notably, we can factor $\iota_{m,n} = \iota_{n-1,n} \circ \cdots \circ \iota_{m,m+1}$.

As in the case of FI, it is convenient to observe that we can factor morphisms of FI_G as a composite of a natural inclusion map and an automorphism.

Proposition 3.7. Every morphism in $f \in \text{Hom}_{\text{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$ factors as a composite

$$\mathbf{m}_X \xrightarrow{\iota_{m,n}} \mathbf{n}_X \xrightarrow{\omega} \mathbf{n}_X$$

for some (non-canonical) $\omega \in \operatorname{End}_{\operatorname{FI}_G}(\mathbf{n}_X) = G \wr S_n$.

In light of Proposition 3.7, as in the theory of FI-modules, we will see that the key to understanding the combinatorics of the FI_G morphisms is to calculate the stabilizer of the morphism $\iota_{m,n}$ under the action of End_{FI_G} (\mathbf{n}_X) = $G \wr S_n$ by postcomposition.

Proposition 3.8. Fix $m, n \ge 0$. Consider the action of $\operatorname{End}_{\operatorname{FI}_G}(\mathbf{n}_X) = G \wr S_n$ by postcomposition on the set of morphisms $\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$. Then the stabilizer $\operatorname{stab}(\iota_{m,n})$ of $\iota_{m,n}$ is precisely the set of morphisms

$$\mathbf{n}_X \longrightarrow \mathbf{n}_X$$

that restrict to the identity on $(\{1, 2, ..., m\} \times X) \subseteq ([n] \times X) = \mathbf{n}_X$. By considering the restriction of these maps to $(\{m + 1, ..., n\} \times X) \subseteq ([n] \times X) = \mathbf{n}_X$, we can canonically identify the stabilizer

 $\operatorname{stab}(\iota_{m,n}) \cong \operatorname{End}_{\operatorname{FI}_G}(\{m+1,\ldots,n\} \times X),$

a group isomorphic to the wreath product $G \wr S_{n-m}$.

3.2. FI_G -modules.

Definition 3.9. An FI_G-module V over a commutative ring R is a functor V from FI_G to the category R-Mod of left R-modules. Given a finite set A, for simplicity we write V_A for $V(A \times X)$. We let V_n denote $V(\mathbf{n}_X)$.

Definition 3.10. Let R be a commutative ring and let V and W be FI_G -modules over R. We define a map

$$F: V \longrightarrow W$$

to be map of FI_G -modules if F is a natural transformation of functors V and W. More concretely, F is a family of R-linear maps

$$F_A: V_A \longrightarrow W_A$$

for each finite set A such that the following diagram commutes for all FI_G morphisms $f: A \times X \to B \times X$,



where f_* denotes the map induced by f. We say that U and V are isomorphic as FI_G -modules if there is a natural isomorphism $F: U \to V$ of FI_G -modules, that is, a natural transformation F such that F_A is an R-linear isomorphism for every finite set A.

Definition 3.11. Let V be an FI_G-module. We say that U is an FI_G-submodule of V if there is a map of FI_G-modules $F: U \to V$ such that F_A is the inclusion of an R-submodule of V_A for each finite set A.

Definition 3.12. Suppose that $F: V \to U$ is a map of Fl_G -modules. Then the family of kernels $\ker(F_A) \subseteq V_A$ for each finite set A form an Fl_G -submodule of V which we denote by $\ker(F)$ and call the *kernel* of F. The family of images $\operatorname{im}(F_A) \subseteq U_A$ form an Fl_G -submodule of U, which we denote by $\operatorname{im}(F_A)$ and call the *image* of F.

Sam–Snowden [SS19, Theorem 1.2.4] show that, in a certain strong sense, the theory of FI_G -modules reduces to the theory of FI-modules.

3.3. An FI_G -module criterion.

Definition 3.13. Let G be a finite group, and R a commutative ring. For each integer $n \ge 0$, let V_n be a $G \wr S_n$ -representation over R, and let $\varphi_n : V_n \to V_{n+1}$ be an R-linear map. View $G \wr S_n$ as the subgroup of $G \wr S_{n+1}$ of symmetries of $([n] \times X) \subseteq ([n+1] \times X)$. We call the sequence

$$V_0 \xrightarrow{\varphi_0} V_1 \xrightarrow{\varphi_1} V_2 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_{n-1}} V_n \xrightarrow{\varphi_n} V_{n+1} \xrightarrow{\varphi_{n+1}} \dots$$

a consistent sequence of $G \wr S_n$ -representations if the maps φ_n are $G \wr S_n$ -equivariant with respect to the action of $G \wr S_n$ on V_n and the restriction of the action to $G \wr S_n \subseteq G \wr S_{n+1}$ on V_{n+1} .

An FI_G-module V defines in particular a consistent sequence of $G \wr S_n$ -representations V_n with equivariant maps $(\iota_{n,n+1})_* : V_n \to V_{n+1}$. However, not every consistent sequence of $G \wr S_n$ -representations arises in this way. The following results show when a sequence of $G \wr S_n$ -representations can be given the structure of a functor from (the full subcategory on objects \mathbf{n}_X of) the category FI_G. This result generalizes Church– Ellenberg–Farb [CEF15, Remark 3.31].

Lemma 3.14. Let G be a finite group, and R a commutative ring. Suppose that

$$V_0 \xrightarrow{\varphi_0} V_1 \xrightarrow{\varphi_1} V_2 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_{n-1}} V_n \xrightarrow{\varphi_n} V_{n+1} \xrightarrow{\varphi_{n+1}} \dots$$

is a consistent sequence of $G \wr S_n$ -representations. Let $\varphi_{m,n} : V_m \to V_n$ denote the composite $\varphi_{n-1} \circ \cdots \circ \varphi_m$.

Let $\iota_{m,n} \in \operatorname{Hom}_{\mathrm{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$ be the natural inclusion map as defined in Notation 3.6. Let $\operatorname{stab}(\iota_{m,n})$ denote

 $\operatorname{stab}(\iota_{m,n}) = \{ \omega \in \operatorname{End}_{\operatorname{FI}_G}(\mathbf{n}_X) = G \wr S_n \mid \omega \circ \iota_{m,n} = \iota_{m,n} \},\$

as in Proposition 3.8. Then there exists a functor $V : \operatorname{FI}_G \to R$ -Mod satisfying $V(\mathbf{n}_X) = V_n$ and $(\iota_{m,n})_* = \varphi_{m,n}$ if and only if the following Condition (*) is satisfied:

(*) for all
$$m < n$$
, $\omega \cdot v = v$, for all $\omega \in \operatorname{stab}(\iota_{m,n})$ and $v \in \varphi_{m,n}(V_m)$

Proof. Given an FI_G -module $V : \operatorname{FI}_G \to R$ -Mod, the sequence $(\iota_{n,n+1})_* : V_n \to V_{n+1}$ satisfies Condition (*) by functoriality. Conversely, suppose that $\varphi_n : V_n \to V_{n+1}$ is a consistent sequence of $G \wr S_n$ -representations that satisfies (*). We will construct the functor V and verify that it is functorial.

It suffices to construct V on the full subcategory of FI_G of objects $\mathbf{n}_X = [n] \times X$ for integers $n \ge 0$. Let $V(\mathbf{n}_X) = V_n$, and let $(\iota_{m,n})_* = \varphi_{m,n}$. Let $f \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$. By Proposition 3.7, we can factor f (non-uniquely) as a composite $f = \sigma \circ \iota_{m,n}$, where $\sigma \in \operatorname{End}_{\operatorname{FI}_G}(\mathbf{n}_X) = G \wr S_n$. Thus we define f_* to be the function $\sigma \circ \varphi_{m,n} : V_m \to V_n$.

We first check that f_* is well-defined. Suppose that $f = \sigma \circ \iota_{m,n} = \tau \circ \iota_{m,n}$ in $\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$, for $\sigma, \tau \in G \wr S_n$. Then $\tau^{-1}\sigma \in \operatorname{stab}(\iota_{m,n})$, so $\sigma \circ \varphi_{m,n} = \tau \circ \varphi_{m,n}$ by Condition (*).

We finally check that V is functorial. Let $h \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{k}_X, \mathbf{m}_X)$ and $f \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_X, \mathbf{n}_X)$. Factor $h = \rho \circ \iota_{k,m}$ and $f = \sigma \circ \iota_{m,n}$. Observe that the map $\iota_{m,n}$ defines an inclusion of $G \wr S_m$ into $G \wr S_n$ as the symmetry group of $\mathbf{m}_X = ([m] \times X) \subseteq ([n] \times X) \subseteq \mathbf{n}_X$. For $\rho \in G \wr S_m$, write $\rho' \in G \wr S_n$ for its image; by definition ρ' satisfies $\iota_{m,n} \circ \rho = \rho' \circ \iota_{m,n}$ as FI_G morphisms. Then,

$$\begin{aligned} f_* \circ h_* &= (\sigma \circ \iota_{m,n})_* \circ (\rho \circ \iota_{k,m})_* \\ &= (\sigma \circ \varphi_{m,n}) \circ (\rho \circ \varphi_{k,m}) \\ &= \sigma \circ \rho' \circ \varphi_{m,n} \circ \varphi_{k,m} \qquad \text{(by the equivariance assumption on the maps } \varphi_i) \\ &= (\sigma \circ \rho' \circ \iota_{m,n} \circ \iota_{k,m})_* \\ &= (\sigma \circ \iota_{m,n} \circ \rho \circ \iota_{k,m})_* \\ &= (f \circ h)_* \qquad \text{which concludes the proof.} \end{aligned}$$

3.4. Generation of FI_G -modules and the Noetherian property. In this section we recall the Noetherian property of FI_G -modules over Noetherian rings. The following definitions appear in Casto [Cas16, Section 2] and Sam–Snowden [SS19, Section 2.1], and are direct analogues of the corresponding definitions for FI-modules [CEF15, Section 2.3].

Definition 3.15. An FI_G-module V is generated by a set $S \subseteq \coprod_{n \ge 0} V_n$ if V is itself the smallest FI_G-submodule of V containing S.

Definition 3.16. An FI_G-module V is *finitely generated* if it is generated by a finite set. We say V is generated in degree $\leq d$ whenever V is generated by $\coprod_{0 \leq n \leq d} V_n$. If V is generated in degree $d < \infty$, then we say V has *finite generation degree*.

Definition 3.17. A finitely generated FI_G -module V is *Noetherian* if every FI_G -submodule of V is finitely generated.

Sam–Snowden [SS19] proved a Noetherianity result for FI_G -modules for a class of groups G called *virtually* polycyclic groups; it includes in particular the case that G is finite.

Theorem 3.18 ([SS19, Corollary 1.2.2 and Theorem 1.2.3]). Let G be a finite group and let R be a left-Noetherian ring. Then an FI_G -module V over R is Noetherian.

3.5. Representable and induced FI_G -modules. In this section, we define a class of "induced" modules with highly controlled combinatorial behaviour. These modules play an essential role in Section 5. The theory developed here, and our notation, closely mirrors Church–Ellenberg–Farb [CEF15, Section 2.2] in the case of FI-modules. The definitions below for general FI_G -modules appear in Casto [Cas16, Section 2], and Ramos [Ram18, Definition 2.4].

Definition 3.19. Fix $d \ge 0$ and a commutative ring R. We denote by M(d) the representable or free FI_G-module over R, defined to be

$$M(d)_n = R[\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{n}_X)]$$

with the action of FI_G morphisms by post composition,

$$f_*: R[\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{m}_X)] \longrightarrow R[\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{n}_X)]$$
$$h \longmapsto f \circ h$$

for $f \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_X, \mathbf{n}_X), h \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{m}_X).$

We observe that, for each n, $M(d)_n$ admits an action of $G \wr S_d = \operatorname{End}_{\operatorname{FI}_G}(\mathbf{d}_X)$ on the right by precomposition. With this observation we can make the following definition.

Definition 3.20. Let R be a commutative ring. Suppose that T is a $R[G \wr S_d]$ -module. Define the *induced* FI_G-module by

$$M(T)_n = M(d)_n \otimes_{R[G \wr S_d]} T$$

and the FI_G morphisms act on the left tensor factor M(d).

Remark 3.21. Sam–Snowden [SS17, Section 1.1] refer to modules of the form M(d) as principal projective modules. Gadish [Gad17a, Definition 3.1] calls the modules M(T) induction FI_G-modules. Ramos [Ram18, Definition 2.4] refers to sums of modules of the form M(T) as relatively projective modules, and Gadish [Gad17a, Section 3] calls all such modules free.

We collect some facts about the FI_G -modules M(d) and M(T). These are direct analogues of the corresponding results for FI-modules (see, for example, Church-Ellenberg-Farb [CEF15, Section 2.2]), and we omit the details of the proofs. Recall the definition of the external tensor product \boxtimes from Notation 2.11.

Proposition 3.22. Let R be a commutative ring. Fix a natural number $d \ge 0$, and let T be a $G \wr S_d$ -representation over R.

- (i) $M(d)_n = 0$ if n < d.
- (ii) $M(d)_d \cong R[G \wr S_d]$, the regular representation of $G \wr S_d$.
- (iii) By an orbit-stabilizer argument, $M(d)_n$ is (as a $G \wr S_n$ -representation) the permutation representation on the cosets

$$\operatorname{End}_{\operatorname{FI}_G}(\mathbf{n}_X)/\operatorname{stab}(\iota_{d,n}) \cong (G \wr S_n)/(G \wr S_{n-d})$$

by Proposition 3.8. Consequently, there are isomorphisms of $G \wr S_n$ -representations

$$M(d)_n \cong \operatorname{Ind}_{G\wr S_{n-d}}^{G\wr S_n} R$$

where R has trivial $G \wr S_{n-d}$ action.

- (iv) M(d) is finitely generated in degree $\leq d$ as an FI_G-module by the identity morphism id in End_{FI_G}(\mathbf{d}_X) $\subseteq M(d)_d$.
- (v) Given any FI_G-module V and element $v \in V_d$, the assignment $id \mapsto v$ extends uniquely to a map of FI_G-modules $M(d) \to V$,

$$\begin{aligned} M(d)_n &\longrightarrow V_n \\ f &\longmapsto f_*(v) \end{aligned}$$

for $f \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{n}_X)$.

(vi) Let V be a FI_G -module and $S \subseteq \coprod_n V_n$ a subset with $s \in V_{d_s}$. Then V is generated by S if and only if the corresponding map of FI_G -modules

$$\bigoplus_{s \in S} M(d_s) \to V$$

is surjective. In particular, V is finitely generated if and only if it admits a surjective map of FI_{G} -modules of the form

$$\bigoplus_{1 \leq i \leq N} M(d_i) \to V$$

and generated in degree $\leq d$ if and only if it admits a surjective map of FI_G-modules of the form

$$\bigoplus_{0 \le m \le d} M(m)^{\oplus c_m} \to V.$$

(vii) $M(d) \cong M(R[G \wr S_d]).$

- (viii) $M(T)_n \cong 0$ if n < d.
- (ix) $M(T)_d \cong T$ as a $G \wr S_d$ -representation.
- (x) There are isomorphisms of $G \wr S_n$ -representations

$$M(T)_n \cong \operatorname{Ind}_{(G \wr S_d) \times (G \wr S_{n-d})}^{G \wr S_n}(T \boxtimes R).$$

Here, $(G \wr S_d) \times (G \wr S_{n-d})$ is the subgroup of $G \wr S_n$ that stabilizes the partition of $\mathbf{n}_X = ([n] \times X)$ into $\{1, \ldots, d\} \times X$ and $\{d + 1, \ldots, n\} \times X$. Again R has trivial $G \wr S_{n-d}$ action. (xi) Let T be a $G \wr S_d$ -module with finite rank over R. Then, as an R-module, $M(T)_n$ has rank

$$\operatorname{rank}_{R}(M(T)_{n}) = \left(\frac{|G \wr S_{n}|}{|(G \wr S_{d}) \times (G \wr S_{n-d})|}\right) \operatorname{rank}_{R}(T) = \binom{n}{d} \operatorname{rank}_{R}(T).$$

Notably, for $T \neq 0$, the rank of $M(T)_n$ is a polynomial in n of degree d.

- (xii) M(T) is generated as an FI_G-module in degree $\leq d$ by $M(T)_d \cong T$.
- (xiii) Given an FI_G-module V, any $G \wr S_d$ -equivariant map $T \to V_d$ extends uniquely to a map of FI_G-modules $M(T) \to V$.

Remark 3.23. We call an FI_G-module projective if it is a projective object in the category of FI_G-modules. As with FI-modules [CEF15, Remark 2.2.A], the projective FI_G-modules are precisely the direct sums of induced modules M(T) for T a projective $R[G \wr S_d]$ -module (Casto [Cas16, Section 2.2], Ramos [Ram18, Proposition 2.13]).

4. Representation stability

Church-Ellenberg-Farb prove that, if V is a finitely generated FI-module over \mathbb{Q} , for all n sufficiently large the decomposition of V_n into irreducible S_n -representations stabilizes in a strong sense [CEF15, Theorem 1.13]. Gan-Li generalize this result to FI_G-modules [GL15, Theorem 1.12]. In this section we summarize their results.

Let G be a finite group. Throughout the section, assume that R is a subfield of \mathbb{C} over which all complex irreducible representations of G are defined. For $G = S_d$ we may take $R = \mathbb{Q}$.

Notation 4.1. For a finite group G, let $Irr(G) = \{\chi_1, \ldots, \chi_r\}$ denote the set of its r distinct isomorphism classes of irreducible representations over R. Let $\chi_1 \in Irr(G)$ denote the 1-dimensional trivial representation.

To make sense of what it means for the irreducible constituents of a sequence V_n of $G \wr S_n$ -representations to stabilize, we need to associate to each irreducible $G \wr S_n$ -representation an irreducible representation of $G \wr S_{n+1}$, of $G \wr S_{n+2}$, of $G \wr S_{n+3}$, The following definition gives a uniform notation for these "stable" sequences of irreducible representations. Recall from Corollary 2.15 that the irreducible representations of $G \wr S_n$ are classified by *n*-multipartitions $\underline{\lambda}$ with *r* components, that is, partition-valued functions

$$\underline{\lambda}: \{\chi_1, \dots, \chi_r\} \to \{ \text{ partitions of nonnegative numbers } \} \qquad \text{satisfying } \sum_{i=1}^r |\underline{\lambda}(\chi_i)| = n$$

For a general multipartition $\underline{\lambda}$ with r components, we write $|\underline{\lambda}|$ for the sum $\sum_{i=1}^{r} |\underline{\lambda}(\chi_i)|$.

Definition 4.2 (Gan–Li [GL15, Section 1.4]). Let $\underline{\lambda}$ be any partition-valued function on $\operatorname{Irr}(G)$. Let $n \in \mathbb{Z}_{\geq 0}$ such that $n \geq |\underline{\lambda}| + \underline{\lambda}(\chi_1)_1$ and $\underline{\lambda}(\chi_1)_1$ is the largest part of the partition $\underline{\lambda}(\chi_1)$. Define

$$\underline{\lambda}[n](\chi) = \begin{cases} (n - |\underline{\lambda}|, \ \underline{\lambda}(\chi_1)) & \chi = \chi_1 \\ \underline{\lambda}(\chi) & \chi \neq \chi_1 \end{cases}$$

Let $L(\underline{\lambda})_n$ denote the irreducible $R[G \wr S_n]$ -module corresponding to the multipartition $\underline{\lambda}[n]$ when $n \ge |\underline{\lambda}| + \underline{\lambda}(\chi_1)_1$, and let $L(\underline{\lambda})_n = 0$ for $n < |\underline{\lambda}| + \underline{\lambda}(\chi_1)_1$.

Remark 4.3. We may visualize a multipartition $\underline{\lambda}$ as an *r*-tuple of Young diagrams. The multipartition $\underline{\lambda}[n]$, then, is constructed from $\underline{\lambda}$ by adding a row of length $n - |\underline{\lambda}|$ to the top of the Young diagram $\underline{\lambda}(\chi_1)$ in the first component. As *n* increases, the tuples $\underline{\lambda}[n]$ and $\underline{\lambda}[n+1]$ differ only in the length of the top row of the first Young diagram.

We illustrate this pattern in the case $G = \mathbb{Z}/3\mathbb{Z}$ (so r = 3) and the multipartition $\underline{\lambda} = (\Box, \Box, \Box)$. In this case, we have



Using this notation, the following definition states the criteria for stability for a consistent sequence of $G \wr S_n$ -representations. This generalizes the notion of representation stability stated for symmetric groups in the introduction in Definition 1.1.

Definition 4.4 (Church–Farb [CF13, Definition 1.1], Gan–Li [GL15, Definition 1.10]). Let $\{\varphi_n : V_n \to V_{n+1}\}$ be a consistent sequence of $R[G \wr S_n]$ -modules V_n in the sense of Definition 3.13. This sequence is representation stable if there exists N > 0 such that for each $n \ge N$, the following three conditions hold:

- (i) **Injectivity:** The map $\varphi_n : V_n \to V_{n+1}$ is injective.
- (ii) Surjectivity: The span of the $G \wr S_{n+1}$ -orbit of $\varphi_n(V_n)$ is all of V_{n+1} .
- (iii) Multiplicities: There is a decomposition

$$V_n = \bigoplus_{\underline{\lambda}} L(\underline{\lambda})_n^{\bigoplus c(\underline{\lambda})}$$

for each n and where the multiplicities $0 \leq c(\underline{\lambda}) \leq \infty$ do not depend on n and with $|\underline{\lambda}| < N$ for each $\underline{\lambda}$ with positive multiplicity $c(\underline{\lambda})$.

Condition (iii) of Definition 4.4 implies that, for all n large enough, we can express the decomposition of V_n into irreducible subrepresentations uniformly in n using finitely many multipartitions $\underline{\lambda}$. By Remark 4.3, for all n large enough, the multipartitions $\underline{\lambda}[n]$ indexing the irreducible constituents of V_n only grow in n in the first row of the first component.

Following the work of Church–Ellenberg–Farb on FI-modules, Gan and Li [GL15] related finite generation of FI_G-modules V to representation stability for sequences $(\iota_{n,n+1})_* : V_n \to V_{n+1}$.

Theorem 4.5 (Church-Ellenberg-Farb [CEF15, Theorem 1.13], Gan-Li [GL15, Theorem 1.12]). Suppose that R is a splitting field for G of characteristic 0. Let V be an FI_G-module over R. Then V is finitely generated as an FI_G-module if and only if the consistent sequence $\{(\iota_{n,n+1})_* : V_n \to V_{n+1}\}$ is a representation stable sequence of $R[G \wr S_n]$ -modules with dim_R $V_n < \infty$ for each n.

5. FI_G #-MODULES

Church-Ellenberg-Farb [CEF15, Section 4.1] study a class of FI-modules V with additional structure which, they proved, places even stronger constraints on the combinatorics of the representations V_n and their characters. Roughly speaking, these are the FI-modules for which the FI morphisms act by split injections, and admit one-sided inverses (satisfying appropriate compatibility conditions). More precisely, these are FI-modules for which the action of FI extends to an action of a larger category—denoted by Church-Ellenberg-Farb by FI \sharp —containing as subcategories both FI and its opposite category.

Ramos [Ram18, Section 2.3] and Casto [Cas16, Section 2.5] generalize the theory FI \sharp -modules to the setting of FI_G. In this section we review the main results. We use a different characterization of the category FI_G \sharp that was suggested to the fourth author by Peter May during her thesis work.

Definition 5.1. Let G be a finite group. Let X be a faithful transitive G-set as in Definition 3.1. For a finite set A, let $(A \times X)_{\clubsuit}$ denote the disjoint union of the set $A \times X$ with a basepoint that we denote by \clubsuit .

Define the category $\operatorname{FI}_G \sharp$ as follows. The objects are, for each finite set A, the based set $(A \times X)_{\clubsuit}$ with basepoint \clubsuit . The morphisms are based maps of sets $f : (A \times X)_{\clubsuit} \to (B \times X)_{\clubsuit}$ satisfying the following conditions.

- $f(\clubsuit) = \clubsuit$
- For any point (b, x) in the codomain distinct from the basepoint, its preimage contains at most 1 element of $A \times X$. We say f is "injective away from the basepoint."
- For each $a \in A$, the image of $(\{a\} \times X)$ is either the basepoint $\{\clubsuit\}$ or the set $(\{b\} \times X)$ for some $b \in B$. In the latter case, the map

$$f|_{(\{a\}\times X)}:(\{a\}\times X)\to (\{b\}\times X)$$

is G-equivariant.

For brevity we denote the object $([n] \times X)_{\bullet}$ by $\mathbf{n}_{X\bullet}$, for all $n \in \mathbb{Z}_{\geq 0}$.

 FI_G embeds in $\operatorname{FI}_G \sharp$ as the wide subcategory of injective morphisms. Each injective morphism in $\operatorname{FI}_G \sharp$ admits a left inverse in $\operatorname{FI}_G \sharp$ that maps the complement of its image to the basepoint. These left inverses form a subcategory isomorphic to FI_G^{op} . The subcategories $\operatorname{FI}_G, \operatorname{FI}_G^{op} \subseteq \operatorname{FI}_G \sharp$ together generate $\operatorname{FI}_G \sharp$.

Remark 5.2. It is not too difficult to verify that the definition of $\operatorname{Fl}_G \sharp$ given in Definition 5.1 is equivalent to the definition used by Casto [Cas16, Section 2.5]. The objects are indexed by finite sets. A morphism from A to B is a pair (Z, f), where $Z \subseteq A$ is a subset, and $f : Z \to B$ is an Fl_G morphism. We may identify (Z, f) with the morphism (in the category of Definition 5.1) that is defined by f on $Z \times X$ and that maps $(A \setminus Z) \times X$ to the basepoint.

As in the case of FI-modules [CEF15, Example 4.1.4], the action of FI_G on M(d) and M(T) extends to an action of FI_G[#]. For $f \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{m}_{X, \mathbf{A}}, \mathbf{n}_{X, \mathbf{A}}), h \in \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{m}_X) \subseteq \operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_{X, \mathbf{A}}, \mathbf{m}_{X, \mathbf{A}})$, we define

$$\begin{aligned} f_* : R[\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{m}_X)] &\longrightarrow R[\operatorname{Hom}_{\operatorname{FI}_G}(\mathbf{d}_X, \mathbf{n}_X)] \\ h &\longmapsto \begin{cases} (f \circ h) \big|_{\mathbf{d}_X}, & \text{if } \clubsuit \notin (f \circ h)(\mathbf{d}_X) \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

This in turn defines an action of FI_G on M(T) for any d and W_d -representation T.

Definition 5.3. For a commutative ring R, an $\operatorname{FI}_G \sharp$ -module is a functor V from $\operatorname{FI}_G \sharp$ to the category of left R-modules. A map of $\operatorname{FI}_G \sharp$ -modules over R is a natural transformation of functors.

We define kernels, images, etc, of $\operatorname{FI}_G \#$ -module maps pointwise, as with FI_G -modules. Any $\operatorname{FI}_G \#$ -module map will also be a map of FI_G -modules under the restriction of the actions to $\operatorname{FI}_G \subseteq \operatorname{FI}_G \#$.

5.1. The structure theorem for $FI_G \#$ -modules. Church–Ellenberg–Farb [CEF15, Theorem 4.1.5] give a complete classification of FI #-modules. Their argument adapts readily to $FI_G \#$ -modules; see Ramos [Ram18, Theorem 2.19] or Casto [Cas16, Proposition 2.7].

Theorem 5.4 (Church–Ellenberg–Farb [CEF15, Theorem 4.1.5]; Casto [Cas16, Proposition 2.7]). Any $\operatorname{FI}_G \sharp$ -module, up to isomorphism of $\operatorname{FI}_G \sharp$ -modules, has the form $\bigoplus_{d \ge 0} M(T_d)$ for T_d a $G \wr S_d$ -representation (possibly 0).

Concretely, in light of the description of M(T) in Proposition 3.22, Theorem 5.4 implies that any $\operatorname{Fl}_G \sharp$ module has the following structure.

Corollary 5.5. Let V be a $\operatorname{FI}_G \sharp$ -module over a commutative ring R. Then there is some sequence of $G \wr S_d$ -representations T_d such that, as a $G \wr S_n$ -representations,

$$V_n \cong \bigoplus_{d \ge 0} \operatorname{Ind}_{(G \wr S_d) \times (G \wr S_{n-d})}^{G \wr S_n} (T_d \boxtimes R) \quad \text{for all } n$$

The underlying FI_G-module of V is finitely generated in degree $\leq m$ if and only if $T_d = 0$ for all $d \geq m$.

If each representation T_d has finite rank as an R-module, then

$$\operatorname{rank}_{R}(V_{n}) = \sum_{d \ge 0} \binom{n}{d} \operatorname{rank}_{R}(T_{d}).$$

If some representation T_d has infinite rank, then V_d will have infinite rank.

As observed implicitly by Gan–Li [GL15, Section 8.1] and Casto [Cas16, Corollary 2.8], Pieri's rule for wreath products (stated in Corollary 2.18) therefore implies that finitely generated $FI_G \#$ -modules are representation stable in the sense of Definition 4.4, with stable range depending only on the degree of generation. In the following corollary, we collect this and other consequences of the structure theorem for $FI_G \#$ -modules.

Corollary 5.6. Let V be an FI_G #-module over a commutative ring R. The following hold.

- (i) Suppose, as an FI_G-module, V is generated in degree $\leq m$. Then $V \cong \bigoplus_{0 \leq m \leq d} M(T_m)$ for some $G \wr S_m$ representations T_m for $0 \leq m \leq d$.
- (ii) $V \cong \bigoplus_m M(T_m)$ is finitely generated as an FI_G-module if and only if $\bigoplus_m T_m$ is finite dimensional, that is, each representation T_m is finite dimensional, and nonzero for at most finitely many m.
- (iii) Let $R = \mathbb{K}$ be a splitting field for G of characteristic 0. Suppose, as an FI_G-module, V is finitely generated in degree $\leq d$. Then V is representation stable in the sense of Definition 4.4, stabilizing for $n \geq 2d$.
- (iv) Suppose V is finitely generated as an FI_G -module in degree $\leq d$. Then the sequence $\operatorname{rank}_R(V_n)$ is polynomial in n of degree $\leq d$.
- (v) Conversely, suppose $\operatorname{rank}_R(V_n)$ is a polynomial in n of degree $\leq d$. Then $V \cong \bigoplus_{0 \leq m \leq d} M(T_m)$. Consequently, as an FI_G-module, V is finitely generated in degree $\leq d$. If $\operatorname{rank}_R(V_n)$ is a polynomial of degree exactly d, then $T_d \neq 0$, and the bound on degree of generation is sharp, in the sense that V is not generated in degree $\leq (d-1)$.

6. Character polynomials for FI_G -modules

Casto shows that, as with FI \sharp -modules (Church–Ellenberg–Farb [CEF15, Theorem 4.1.7]), the characters of FI_G \sharp -modules are highly constrained and admit a description that is uniform in n.

Recall from Section 2.2 that the conjugacy classes of $G \wr S_n$ are classified by the cycle type of a permutation in S_n , along with (associated to each unordered cycle) a conjugacy class of an element of G.

Notation 6.1. Let G be a finite group, and let C_1, \ldots, C_r be a complete list of its distinct conjugacy classes. Fix an integer $d \ge 1$. For any n, let $X_d^{C_i}$ denote the class function on $G \wr S_n$ that outputs the number of d-cycles in its cycle type labelled by the conjugacy class C_i . We call the functions $X_d^{C_i}$ the coloured cyclecounting functions, as they count the number of cycles in the cycle type of an element of $G \wr S_n$ that are "coloured" by a specified conjugacy class of G.

In the notation of James–Kerber (see Definition 2.7), the function $X_d^{C_i}$ is the function a_{id} .

As Casto calculates explicitly [Cas16, Theorem 2.2], the formula for M(T) as an induced representation (Theorem 3.22 part (x)) allows us to express the character of $M(T)_n$ in terms of the coloured cycle-counting functions. The result is that the sequence of characters of $M(T)_n$ are given by a unique polynomial in the coloured cycle-counting functions, independent of n.

The coloured cycle-counting functions $X_d^{C_i}$ (viewed as functions on the set $\coprod_n G \wr S_n$) are algebraically independent, and we consider the polynomial ring

$$\mathbb{K}[X_d^{C_i} \mid d \ge 1, C_i \text{ a conjugacy class of } G]$$

as a graded ring for which the element $X_d^{C_i}$ has degree d.

Proposition 6.2 ([Cas16, proof of Theorem 2.2]). Let G be a finite group with conjugacy classes C_1, \ldots, C_r . Let K be a splitting field for G of characteristic zero, and let χ^1, \ldots, χ^r be the irreducible G-representations over K. Let T be the $G \wr S_d$ -representation with character χ^T . Then, for every n, the $G \wr S_n$ -representation $M(T)_n$ has character equal to the degree-d character polynomial

$$\chi^{M(T)_n} = \sum_{\substack{\text{conjugacy classes}\\\sigma \text{ of } G \wr S_d}} \chi^T(\sigma) \left(\prod_i \prod_{0 \leqslant m \leqslant d} \begin{pmatrix} X_m^{C_i} \\ X_m^{C_i}(\sigma) \end{pmatrix} \right).$$

In light of the structure theorem for $FI_G \sharp$ -modules and its consequences (see Theorem 5.4 and Corollary 5.5), Proposition 6.2 has the following corollary.

Corollary 6.3. Let V be an $\operatorname{FI}_G \sharp$ -module over a splitting field \mathbb{K} of G of characteristic zero. Suppose $\dim_{\mathbb{K}}(V_n)$ is finite for each n, and V is generated as an FI_G -module in degree $\leq m$. Then the sequence of characters χ^{V_n} is given by a single polynomial

$$P^V \in \mathbb{K}[X_d^{C_i} \mid d \ge 1, C_i \text{ a conjugacy class of } G],$$

independent of n, in the coloured cycle-counting functions $X_d^{C_i}$, and P^V has degree $\leq m$.

Casto [Cas16, Theorem 2.2], generalizing Church–Ellenberg–Farb, in fact proves that the characters of all finitely generated FI_G -modules are equal to a character polynomial for n sufficiently large, with the stable range depending on an invariant of V called the *relation degree*.

7. Representation stability for vertical configuration spaces

Recall the vertical configuration spaces $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$ of points in \mathbb{R}^{p+q} defined in Definition 1.4. In this section, we prove the main theorems of our paper: representation stability for the (co)homology these spaces, and classical (co)homological stability for the unordered vertical configuration spaces.

In Section 7.1, we summarize the computation of the cohomology groups $H^*\left(\tilde{\mathcal{V}}_K(\mathbb{R}^{p,q})\right)$ due to Bianchi– Kranhold [BK22]. We use this calculation in Section 7.2 to prove that, in each cohomological degree, the Betti numbers of these groups grow polynomially. Then, in Section 7.3, we verify that these cohomology groups assemble to form an FI_G-module. This enables us to prove our main theorem, Theorem 7.24. Finally, in Section 7.5, we review the theory of the transfer homomorphism of a regular covering space, and use it to deduce classical (co)homological stability for the unordered vertical configuration spaces.

Remark 7.1. In his beautiful work [Gad17b], Gadish gave broad general topological criteria for a family of complements of linear subspace arrangements to satisfy (co)homological representation stability. We believe that these results would give an alternate approach to our main theorem, which does not rely on the calculations done by Bianchi–Kranhold.

7.1. Bianchi–Kranhold's computation of the cohomology groups $H^*\left(\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})\right)$. Bianchi–Kranhold index a basis for the integral cohomology groups $H^*\left(\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})\right)$ using combinatorial objects they call *ray partitions* [BK22, Section 3.1 and Theorem 3.7]. We now review their definitions.

Throughout this subsection, we fix a partition $K = (k_1, \ldots, k_r)$ with $k_i \ge 1$.

Definition 7.2 ([BK22, Definition 3.1]). The table associated with the partition K is the set

 $T_K := \{(i,j) : 1 \leq i \leq r, 1 \leq j \leq k_i\}$

We order T_K lexicographically, which means we write (i, j) < (i', j') if either i < i', or i = i' and j < j'.

For the following two definitions, let Q be a partition of T_K into non-empty subsets Q_1, \ldots, Q_ℓ .

Definition 7.3 ([BK22], Notation 3.2). Let $\ell(Q) = \ell$ denote the *length* of the partition. Note that

$$1 \leq \ell(Q) \leq |K|.$$

Definition 7.4 ([BK22, Notation 3.2]). Let ~ be the equivalence relation on the index set $\{1, \ldots, \ell\}$ generated by the identifications

$$\beta \sim \beta' \iff \exists (i,j), (i,j') \in T_K \text{ with } j \neq j' \text{ such that } (i,j) \in Q_\beta \text{ and } (i,j') \in Q_{\beta'}$$

We define the *agility* of Q, denoted by a(Q), to be the number of equivalence classes under \sim . Note that

$$1 \le a(Q) \le \min(\ell(Q), r).$$

Definition 7.5 ([BK22, Definition 3.3]). A ray partition Q of type K is a partition of non-empty subsets Q_1, \ldots, Q_ℓ of T_K , with a total order $<_\beta$ on each piece Q_β (called a ray), such that the following hold

(i) if < denotes the global order, i.e, the order in T_K , then the parts Q_i are indexed according to their global order,

$$\min(Q_1, <) < \cdots < \min(Q_\ell, <)$$

(ii) for each $1 \leq \beta \leq \ell$,

$$\min(Q_{\beta}, <_{\beta}) = \min(Q_{\beta}, <)$$

Bianchi–Kranhold [BK22, Theorem 3.7] provide the following explicit description of the cohomology groups of $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$.

Theorem 7.6 ([BK22, Theorem 3.7]). Let $p \ge 0$, $q \ge 1$ and $K = (k_1, \ldots, k_r)$ with $k_i \ge 1$. The integral cohomology $H^*(\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q}))$ is freely generated as an abelian group by classes u_Q for each ray partition Q, and the cohomological degree of u_Q is

$$|u_Q| = p(r - a(Q)) + (q - 1)(|K| - \ell(Q)).$$

7.2. The ranks of the (co)homology groups grow polynomially in *n*. Throughout this section, we let $k, n \in \mathbb{Z}_{>0}$ and define a partition K = (k, k, ..., k) such that |K| = nk. We set $p \ge 0$ and $q \ge 2$, and write $\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$ for $\widetilde{\mathcal{V}}_K(\mathbb{R}^{p,q})$.

In this section, we show that for a fixed k and d, the rank of $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}))$ grows polynomially in n as $n \to \infty$. According to Theorem 7.6, the degree-d cohomology group $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}))$ has a basis indexed by ray partitions Q that satisfy

(1)
$$d = p(n - a(Q)) + (q - 1)(nk - \ell(Q))$$

We first collect two general combinatorial lemmas. We then observe how Equation 1 constrains the possible lengths of the ray partitions in fixed cohomological degree d. We use the combinatorial lemmas to bound the number of possible ray partitions of a given length. We deduce that $\operatorname{rank}(H^d(\tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})))$ is bounded by a polynomial of degree at most $\left|\frac{2d}{q-1}\right|$ when $q \ge 2$.

Notation 7.7. Let S(n,t) denote the Stirling number of the second kind, that is, S(n,t) is the number of ways to partition a set of n objects into t nonempty unordered subsets.

See (for example) [Aig07, Chapter 1.3] for the combinatorial theory of these Stirling numbers. Crucially, they satisfy the following defining combinatorial identity.

Proposition 7.8. For $n, t \in \mathbb{Z}_{>0}$,

$$S(n,t) = tS(n-1,t) + S(n-1,t-1)$$
 for $n \ge 1$.

Proposition 7.9. For fixed t, the sequence $(S(n, n-t))_{n>t}$ is polynomial in n of degree 2t.

Proof. We proceed by induction on t. When t = 0 and $n \ge 1$, S(n, n) = 1. Suppose now for some fixed $t \ge 1$ that $(S(n, n-t))_{n>t}$ is polynomial in n of degree 2t. We want to show that the sequence $s_n := S(n, n-(t+1))$ is polynomial of degree 2t + 2. Proposition 7.8 states

$$S(n, n-t-1) = (n-t-1)S(n-1, n-t-1) + S(n-1, n-t-2)$$

that is, the differences $s_n - s_{n-1}$ satisfy

$$s_n - s_{n-1} = (n - t - 1)S(n - 1, n - t - 1).$$

Let p(n) = (n-t-1)S(n-1, n-t-1). Since p(n) is polynomial of degree 2t+1 by the inductive hypothesis, $s_n = \sum_{i=1}^n p(n-i)$ is a polynomial of degree 2t+2.

We also make the following elementary observation:

Lemma 7.10. For nonnegative integers a_1, a_2, \ldots, a_p ,

$$(a_1 + a_2 + \dots + a_p)! \ge a_1!a_2!\dots a_p!$$

The following lemma gives a bound on the lengths of a ray partition Q associated to a cohomology generator in cohomological degree d.

Lemma 7.11. Suppose $q \ge 2$. Then

$$\left(nk - \frac{d}{q-1}\right) \leqslant \ell(Q) \leqslant nk$$

In particular, ray partitions Q that index a cohomology basis element in fixed cohomological degree d admit at most $\left(1 + \frac{d}{q-1}\right)$ possible distinct values of $\ell(Q)$, a number that does not depend on n.

Proof. The right hand side is clear by the definition of $\ell(Q)$. The left hand side follows from the inequality

$$= (q-1)(nk-\ell(Q)) + p(n-a(Q)) \geq (q-1)(nk-\ell(Q))$$

since the term p(n - a(Q)) is nonnegative.

d

We now bound the number of ray partitions of a given length.

Lemma 7.12. Fix an integer t with $0 \le t \le \left\lfloor \frac{d}{q-1} \right\rfloor$. As n tends to infinity, the number of ray partitions Q of length $\ell(Q) = nk - t$ is polynomial in n of degree at most 2t.

Proof. There are S(nk, nk-t) possible partitions of the table T_K (Definition 7.2) of length $\ell = nk-t$. Denote the parts Q_1, \ldots, Q_ℓ according to indexing convention (i) in the definition of a ray partition (Definition 7.5). By Definition 7.5, then, there is an associated ray partition Q for each choice of total order on each part Q_i , each subject to a specified minimal element. The number of such choices of total orderings is

$$(Q_1 - 1)!(Q_2 - 1)! \dots (Q_\ell - 1)!$$

$$\leq \left(\sum_i Q_i - 1\right)! \quad \text{by Lemma 7.10}$$

$$= (nk - \ell)!$$

$$= t!$$

Since t! is constant in n, the result follows from Lemma 7.9.

We can now prove the main theorem of this section.

Theorem 7.13. Let $p \ge 0$ and $q \ge 2$. For a fixed degree d, as n tends to infinity, the rank of $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Z})$ is polynomial in n of degree at most $\left|\frac{2d}{a-1}\right|$.

Proof. The result follows from Lemmas 7.11 and 7.12.

The universal coefficient theorems imply the following corollary.

Corollary 7.14. Let $p \ge 0$ and $q \ge 2$. For a fixed degree d, rank $(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q}))$ is polynomial in n of degree at most $\left|\frac{2d}{q-1}\right|$.

7.3. The (co)homology groups of vertical configuration spaces form $\operatorname{FI}_G \sharp$ -modules. Again fix $p \ge 0, q \ge 2, k \in \mathbb{Z}^+$, and $K = (k, \ldots, k)$ with |K| = nk. Again we let $\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$ denote the ordered vertical configuration space of n clusters with k points in each.

Let $G = S_k$. We work over the category FI_{S_k} . Our goal is to verify that, for a fixed (co)homology degree d, the (co)homology groups of $\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$ admit an $\operatorname{FI}_G \sharp$ -module structure. Our construction closely parallels Church-Ellenberg-Farb [CEF15, Proposition 6.4.2]. We will view the objects of FI_G as sets of the form $A \times [k]$ for A a finite set, as discussed in Remark 3.3, and analogously view objects of $\operatorname{FI}_G \sharp$ as based sets $(A \times [k]) \cup \{\clubsuit\}$. We first adapt some definitions introduced in [CEF15] to FI_G -modules.

Definition 7.15. An FI_G -space is a covariant functor X from FI_G to the category Top of topological spaces. A co- FI_G -space is a contravariant functor X from FI_G to Top. $\operatorname{FI}_G \sharp$ -spaces and co- $\operatorname{FI}_G \sharp$ -spaces are defined analogously.

Definition 7.16. A homotopy Fl_G -space is a functor X from $\operatorname{Fl}_G \sharp$ to hTop, the category of topological spaces and homotopy classes of continuous maps. Homotopy $\operatorname{Fl}_G \sharp$ -spaces are defined analogously.

Since a homotopy class of maps induces a well-defined map on homology and cohomology, the sequences of singular homology and cohomology groups of a homotopy $FI_G \sharp$ -space form $FI_G \sharp$ -modules.

Notation 7.17. Let B be a finite set. Write

 $T_B^k := \{(b, j) \mid b \in B \text{ and } 1 \leq j \leq k\} = B \times [k]$

In other words, T_B^k is the object of FI_G associated to the finite set *B*. This notation is suggestive of Definition 7.2, as $T_{(k,...,k)}$ is the table T_B^k for B = [n].

Definition 7.18. Let π_p be the projection of $\mathbb{R}^{p,q}$ onto the first p coordinates. Define the space

$$\operatorname{Ver}_{B}^{k}(\mathbb{R}^{p,q}) := \{ \rho \mid \rho : T_{S}^{k} \hookrightarrow \mathbb{R}^{p,q} \text{ satisfying } \pi_{p}(\rho(b,i)) = \pi_{p}(\rho(b,j)) \text{ for } (b,i), (b,j) \in T_{B}^{k} \}$$

A choice of bijection of sets $B \cong [|B|]$ defines an $S_k \wr S_{|B|}$ -equivariant homeomorphism $\operatorname{Ver}_B^k(\mathbb{R}^{p,q}) \cong \widetilde{\mathcal{V}}_{|B|}^k(\mathbb{R}^{p,q})$. The "categorical" description of the vertical configuration spaces of Definition 7.18 enables us to directly define a contravariant action of the category FI_G , with morphisms acting by precomposition.

Proposition 7.19. Ver^k ($\mathbb{R}^{p,q}$) is a co-FI_G-space, where an FI_G morphism $f: T_A^k \to T_B^k$ acts by

$${}^{\mathfrak{k}*}: \operatorname{Ver}_B^k(\mathbb{R}^{p,q}) \longrightarrow \operatorname{Ver}_A^k(\mathbb{R}^{p,q})$$

$$\rho \longmapsto \rho \circ f.$$

To extend this co-FI_G-action to an action of $FI_G \sharp$, we define stabilization maps on that introduce a new cluster of points "far away."

Definition 7.20. Fix $p \ge 0, q \ge 1$, and an ordered partition $(k_1, \ldots, k_n, k_{n+1})$. Let e_1, \ldots, e_{p+q} denote the standard basis for \mathbb{R}^{p+q} , and pr_{e_1} the projection onto the first component. Define a map

$$\tilde{\phi}: \tilde{\mathcal{V}}_{(k_1,\dots,k_n)}(\mathbb{R}^{p,q}) \to \tilde{\mathcal{V}}_{(k_1,\dots,k_n,k_{n+1})}(\mathbb{R}^{p,q})$$

as follows. We map an element $(z_1, \ldots, z_n) \in \tilde{\mathcal{V}}_{(k_1, \ldots, k_n)}(\mathbb{R}^{p,q})$ with clusters $z_i = (z_i^1, \ldots, z_i^{k_i})$ to the element $\tilde{\phi}(z_1, \ldots, z_n) = (z_1, \ldots, z_n, z_{n+1})$, where the new cluster $z_{n+1} = (z_{n+1}^1, \ldots, z_{n+1}^{k_{n+1}})$ consists of the points

$$z_{n+1}^{t} = \left(\max_{\substack{1 \le j \le n \\ 1 \le i \le k_j}} |pr_{e_1}(z_i^j)| + 1\right) e_1 + t e_{p+q}$$

as illustrated in Figure 4. For the partition (k, \ldots, k) , we denote the stabilization map

$$\tilde{\phi}_n: \tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}) \longrightarrow \tilde{\mathcal{V}}_{n+1}^k(\mathbb{R}^{p,q})$$

3,2•			3,2•		•4,1
1,3•	• 2, 3		1,3•	• 2, 3	
3,1•	• 2, 1	\mapsto	3,1•	• 2, 1	
1, 1 • 1, 2 •	• 2, 2 • 2, 4		1,1• 1,2•	• 2, 2 • 2, 4	• 4, 2

FIGURE 4. The stabilization map $\tilde{\mathcal{V}}_{(3,2,4)}(\mathbb{R}^{1,1}) \to \tilde{\mathcal{V}}_{(3,2,4,2)}(\mathbb{R}^{1,1})$. Figure adapted from Bianchi–Kranhold [BK22, Figure 1].

We observe that, because $q \ge 2$ by assumption, if we permute the labels on the points in the new cluster z_{n+1} , the resulting map would be homotopic (continuously in $\tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q})$) to $\tilde{\phi}_n$. Similarly, up to homotopy, we can place the cluster further away in the e_1 direction.

Proposition 7.21. Ver^k_{\blacksquare </sup>($\mathbb{R}^{p,q}$) is a homotopy FI_G-space, where the action of (the skeleton of) FI_G is generated by the action of the groups $S_k \wr S_n$ and the stabilization maps $\tilde{\phi}_n : \tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}) \to \tilde{\mathcal{V}}_{n+1}^k(\mathbb{R}^{p,q})$ of Definition 7.20.}

Proof. By construction, the sequence $\left(\tilde{\phi}_n : \tilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}) \to \tilde{\mathcal{V}}_{n+1}^k(\mathbb{R}^{p,q})\right)_n$ is a consistent sequence in the sense of Definition 3.13. By Proposition 3.14, it suffices to check that, for all $\omega \in \operatorname{stab}(\iota_{m,n}) \cong S_k \wr S_{n-m}$, there is equality between $\omega(\iota_{m,n})_* = \omega \tilde{\phi}_{n-1} \cdots \tilde{\phi}_m$ and $(\iota_{m,n})_* = \tilde{\phi}_{n-1} \cdots \tilde{\phi}_m$. These are not equal as functions—we have potentially permuted the labels on the points in the new clusters introduced "far away"-but because $q \ge 2$, they are homotopic maps. This concludes the proof.

The co-FI_G-space structure of Proposition 7.19 and the homotopy FI_G-space structure of Proposition 7.21 combine to give the spaces $\operatorname{Ver}_{\blacksquare}^{k}(\mathbb{R}^{p,q})$ the structure of a homotopy FI_G \sharp -space. We can verify functoriality directly using the arguments from Church–Ellenberg–Farb [CEF15, Proof of Proposition 6.4.2].

Theorem 7.22. $\operatorname{Ver}_{\blacksquare}^{k}(\mathbb{R}^{p,q})$ admits a homotopy FI_{G} space structure with respect to the maps defined in Proposition 7.19 and Proposition 7.21.

Corollary 7.23. Fix $d \ge 0$ and a commutative ring R. The sequence of degree-d cohomology groups $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$, and the sequence of degree-d homology groups $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$, form FI_G \sharp -modules.

7.4. The (co)homology of vertical configuration space is representation stable. We now prove the main theorem.

Theorem 7.24. Let $p \ge 0$ and $q \ge 2$. Fix $k \ge 1$ and (co)homological degree $d \ge 0$.

- Let R be a commutative ring. The sequences $\left(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_n$ and $\left(H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_n$ of $S_k \wr S_n$ representations are representation stable in the sense that they are $\operatorname{FI}_{S_k} \sharp$ -modules finitely generated
 in degree at most $\left|\frac{2d}{q-1}\right|$.
- The sequences $\left(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})\right)_n$ and $\left(H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})\right)_n$ of rational $S_k \wr S_n$ -representations are representation stable in the sense of Definition 4.4, with stable range $n \ge \left|\frac{4d}{a-1}\right|$.

Proof. By Corollary 7.23, for any fixed d, the sequences of representations $\left(H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_{n \ge 1}$ and $\left(H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)\right)_{n \ge 1}$ each have the structure of an $\operatorname{FI}_{S_k} \sharp$ -module. By Theorem 7.13 and Corollary 7.14, their ranks are polynomial in n of degree at most $\left\lfloor \frac{2d}{q-1} \right\rfloor$. By the universal coefficient theorem, the sequences $\left(\dim_{\mathbb{Q}} H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); \mathbb{Q})\right)_n$ and $\left(\dim_{\mathbb{Q}} H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); \mathbb{Q})\right)_n$ are bounded by a polynomial in n of degree $\left\lfloor \frac{2d}{q-1} \right\rfloor$. The result now follows from Corollary 5.6.

Corollary 5.5 implies the following structural results for the (co)homology groups over R.

Corollary 7.25. For $i = 0, \ldots \lfloor \frac{2d}{q-1} \rfloor$, there exist $S_k \wr S_i$ -representations U_i such that, as a $S_k \wr S_n$ -representations,

$$H^{d}(\widetilde{\mathcal{V}}_{n}^{k}(\mathbb{R}^{p,q});R) \cong \bigoplus_{i=0}^{\left\lfloor\frac{2d}{q-1}\right\rfloor} \operatorname{Ind}_{(S_{k}\wr S_{i})\times(S_{k}\wr S_{n-i})}^{S_{k}\wr S_{n}}(U_{i}\boxtimes R) \quad \text{for all } n.$$

The same statement holds for $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q}); R)$.

The definition of representation stability implies the following for the rational (co)homology groups.

Corollary 7.26. Fix $d \ge 0$. The decomposition of the $S_k \wr S_n$ -representation $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ into irreducible components $L(\underline{\lambda})_n$ stabilizes in n in the sense of Definition 4.4, with stable range $n \ge \left\lfloor \frac{4d}{q-1} \right\rfloor$. The same statement holds for $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$.

Corollary 6.3 implies the following.

Corollary 7.27. Fix $d \ge 0$. The sequence of characters of the $S_k \wr S_n$ -representations $H^d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$ can be expressed as a single character polynomial in the coloured cycle-counting functions

$$\mathcal{P}^{H^d(\mathcal{V}_n^k(\mathbb{R}^{p,q}))} \in \mathbb{Q}[X_r^{C_i} \mid r \ge 1, C_i \text{ a conjugacy class of } S_k]$$

of degree at most $\left\lfloor \frac{2d}{q-1} \right\rfloor$, independent of n. The same statement holds for the $H_d(\widetilde{\mathcal{V}}_n^k(\mathbb{R}^{p,q});\mathbb{Q})$.

Remark 7.28. The range in Corollary 7.26 may not be sharp in general. In the case p = 0 and k = 1, which is the ordinary ordered configuration space $\operatorname{Conf}_n(\mathbb{R}^q)$, Theorem 7.24 gives us the stable range $n \ge \left\lfloor \frac{4d}{q-1} \right\rfloor$ for $q \ge 2$. In this case the cohomology groups are nonzero only in degrees d divisible by (q-1), and for the nonzero groups the sharp range is given by Hersh–Reiner [HR17, Theorem 1.1]:

$$n \ge \begin{cases} \frac{3d}{q-1} & \text{if } q \ge 3 \text{ odd} \\ \frac{3d}{q-1} + 1 & \text{if } q \ge 2 \text{ even}. \end{cases}$$

7.5. The transfer map and classical (co)homological stability. In this section we give another proof of the stability of the rational (co)homology group of the unordered vertical configuration spaces $\mathcal{V}_n^k(\mathbb{R}^{p,q})$. Our proof uses representation stability and the transfer map.

Analogous to the stabilization maps we defined on the ordered vertical configuration spaces in Definition 7.20, we define stabilization maps $\phi_n : \mathcal{V}_n^k(\mathbb{R}^{p,q}) \to \mathcal{V}_{n+1}^k(\mathbb{R}^{p,q})$ on the unordered spaces by introducing a $(n+1)^{st}$ cluster of k points "far away," as in Figure 5. See Latifi [Lat17, Section 4.1] or Bianchi–Kranhold [BK22, Construction 4.1].



FIGURE 5. The stabilization map $\phi_5 : \mathcal{V}_5^2(\mathbb{R}^{1,1}) \to \mathcal{V}_6^2(\mathbb{R}^{1,1})$. Figure adapted from Bianchi–Kranhold [BK22, Figure 6].

Latifi and Bianchi–Kranhold proved the following stability result with respect to the maps ϕ_n . Their results generalize the work of Arnol'd [Arn70b] and others on homological stability for classical configuration spaces of points in \mathbb{R}^s , and more recent work of Palmer [Pal21] on unlabelled clusters of points in \mathbb{R}^s (without the "verticality" condition).

Theorem 7.29 (Latifi [Lat17, Theorem 4.1.4] and Bianchi–Kranhold [BK22, Theorem 4.3]). For all $p \ge 0$ and $q \ge 1$ with $(p,q) \ne (0,1)$, the induced maps

$$H_d(V_n^k(\mathbb{R}^{p,q})) \longrightarrow H_d(V_{n+1}^k(\mathbb{R}^{p,q}))$$

are isomorphisms for all $n \ge 2d$.

They note that stability fails when (p,q) = (0,1).

We give a new proof of rational (co)homological stability in the cases $p \ge 0$ and $q \ge 2$. Although our results only hold rationally, for $q \ge 3$ they improve the stable range.

To deduce (co)homological stability from our main representation stability result Theorem 7.24, we follow a standard line of reasoning in the representation stability literature (see, for example, Church–Ellenberg– Farb [CEF15, Corollary 6.3.4]). We will, however, give a more detailed justification of the approach, as these details may not be easily found in the literature. We first review the theory of transfer homomorphisms of covering spaces.

7.5.1. The transfer map. Recall that, given a representation U of a finite group \mathcal{G} over \mathbb{Q} , the "averaging map" over the action of \mathcal{G}

$$\begin{array}{c} \alpha_{\mathcal{G}} \colon U \longrightarrow U \\ \\ u \longmapsto \frac{1}{|\mathcal{G}|} \sum_{g \in \mathcal{G}} g \cdot u \end{array}$$

defines a projection from U to its \mathcal{G} -invariant subspace $U^{\mathcal{G}}$, which factors through an isomorphism between the \mathcal{G} -coinvariants and \mathcal{G} -invariants $U_{\mathcal{G}} \cong U^{\mathcal{G}}$.

We now recall some facts about transfer maps (see, for example, Hatcher [Hat02, Section 3.G]).

Definition 7.30. Let \widetilde{X} be an N-sheeted covering space $\pi : \widetilde{X} \to X$ of a space X for some $N \in \mathbb{N}$. Then π induces a map $\pi_{\#} : C_d(\widetilde{X}; \mathbb{Q}) \to C_d(X; \mathbb{Q})$ on singular chains. Define a map

$$\tau: C_d(X; \mathbb{Q}) \longrightarrow C_d(\widetilde{X}; \mathbb{Q})$$
$$[\sigma: \Delta^k \to X] \longmapsto \left[\sum_{\text{lifts } \tilde{\sigma_i} \text{ of } \sigma_i} \frac{(\tilde{\sigma_i}: \Delta^k \to \widetilde{X})}{N}\right]$$

The map τ is a chain map, and so it induces transfer homomorphisms $\tau_* : H_d(X; \mathbb{Q}) \to H_d(\tilde{X}; \mathbb{Q})$ on homology and $\tau^* : H^d(X; \mathbb{Q}) \to H^d(\tilde{X}; \mathbb{Q})$ on cohomology, satisfying $\pi_* \tau_* = id$ and $\tau^* \pi^* = id$. When π is a regular cover with deck group \mathcal{G} , then the map $\tau \pi_{\#}$ is the 'averaging map' $\alpha_{\mathcal{G}}$,

$$C_d(\widetilde{X};\mathbb{Q}) \xrightarrow{\pi_{\#}} C_d(X;\mathbb{Q}) \xrightarrow{\tau} C_d(\widetilde{X};\mathbb{Q})$$
$$\widetilde{\sigma} \longmapsto \pi_{\#}(\widetilde{\sigma}) \longmapsto \frac{1}{|\mathcal{G}|} \sum_{g \in G} g \cdot \widetilde{\sigma}.$$

The induced map $\tau_*\pi_*$ is the averaging map on homology $H_d(\widetilde{X};\mathbb{Q})$, and the induced map $\pi^*\tau^*$ is the averaging map on cohomology $H^d(\widetilde{X};\mathbb{Q})$. This implies the following (standard) result.

Proposition 7.31. Let X be a topological space, \widetilde{X} be the regular covering space with finite deck group \mathcal{G} . Then the map $\pi^* : H^d(X; \mathbb{Q}) \to H^d(\widetilde{X}; \mathbb{Q})$ induces an isomorphism between $H^d(X; \mathbb{Q})$ and the \mathcal{G} -invariant subspace of $H^d(\widetilde{X}; \mathbb{Q})$

$$H^d(X; \mathbb{Q}) \xrightarrow{\cong} H^d(\widetilde{X}; \mathbb{Q})^{\mathcal{G}}$$

The map $\tau^* : H^d(\widetilde{X}; \mathbb{Q}) \to H^d(X; \mathbb{Q})$ factors through an isomorphism between the \mathcal{G} -coinvariants of $H^d(\widetilde{X}; \mathbb{Q})$ and $H^d(X; \mathbb{Q})$,

$$H^d(\widetilde{X};\mathbb{Q})_{\mathcal{G}} \xrightarrow{\cong} H^d(X;\mathbb{Q})$$

Similarly, the maps π_* and τ_* , respectively, factor through isomorphisms

$$H_d(\widetilde{X};\mathbb{Q})_{\mathcal{G}} \xrightarrow{\cong} H_d(X;\mathbb{Q}) \quad \text{and} \quad H_d(X;\mathbb{Q}) \xrightarrow{\cong} H_d(\widetilde{X};\mathbb{Q})^{\mathcal{G}}, \text{ respectively.}$$

7.5.2. Classical (co)homological stability as a consequence of representation stability. We will apply this theory to the situation where we have a sequence of covers \tilde{X}_n equipped with maps $\tilde{\varphi}_n : \tilde{X}_n \to \tilde{X}_{n+1}$ and $\tilde{\varphi}^n : \tilde{X}_{n+1} \to \tilde{X}_n$ as shown

satisfying the relations $\varphi^n \pi_{\#} = \pi_{\#} \tilde{\varphi}^n$ and $\varphi_n \pi_{\#} = \pi_{\#} \tilde{\varphi}_n$. We assume that the induced maps

$$\left((\tilde{\varphi}^n)^* : H^d(\tilde{X}_n; \mathbb{Q}) \longrightarrow H^d(\tilde{X}_{n+1}; \mathbb{Q}) \right)_n \text{ and } \left((\tilde{\varphi}_n)_* : H_d(\tilde{X}_n; \mathbb{Q}) \longrightarrow H_d(\tilde{X}_{n+1}; \mathbb{Q}) \right)_n$$

form consistent sequences in the sense of Definition 3.13, (or assume at a minimum that the composites

$$H^{d}(\widetilde{X}_{n};\mathbb{Q}) \xrightarrow{(\tilde{\varphi}^{n})^{*}} H^{d}(\widetilde{X}_{n+1};\mathbb{Q}) \twoheadrightarrow H^{d}(\widetilde{X}_{n+1};\mathbb{Q})_{\mathcal{G}_{n+1}}$$
$$H_{d}(\widetilde{X}_{n};\mathbb{Q}) \xrightarrow{(\tilde{\varphi}_{n})_{*}} H_{d}(\widetilde{X}_{n+1};\mathbb{Q}) \twoheadrightarrow H_{d}(\widetilde{X}_{n+1};\mathbb{Q})_{\mathcal{G}_{n+1}}$$

factor through the coinvariants $H^d(\widetilde{X}_{n+1}; \mathbb{Q})_{\mathcal{G}_n}$ and $H_d(\widetilde{X}_n; \mathbb{Q})_{\mathcal{G}_n}$, respectively.)

The relations $\varphi^n \pi_{\#} = \pi_{\#} \tilde{\varphi}^n$, $\varphi_n \pi_{\#} = \pi_{\#} \tilde{\varphi}_n$ and $\pi_{\#} \tau = id$, imply

$$\varphi^n = \varphi^n(\pi_{\#}\tau) = (\varphi^n\pi_{\#})\tau = (\pi_{\#}\tilde{\varphi}^n)\tau$$

$$\varphi_n = \varphi_n(\pi_{\#}\tau) = (\varphi_n\pi_{\#})\tau = (\pi_{\#}\tilde{\varphi}_n)\tau.$$

Thus we can identify the induced maps $(\varphi^n)^*$ and $(\varphi_n)_*$ on (co)homology with the composites





In particular, we can express the map $(\varphi^n)^*$ as a composite of the induced map $(\tilde{\varphi}^n)^*_{\mathcal{G}}$ on coinvariants with several other maps that are all isomorphisms. The analogous statement holds for $(\varphi_n)^*$. We conclude the following result.

Proposition 7.32. With notation as in the previous paragraphs, the map

$$(\varphi^n)^* : H^d(X_n; \mathbb{Q}) \longrightarrow H^d(X_{n+1}; \mathbb{Q})$$

is an isomorphism (or injective, surjective, respectively) on cohomology precisely when the induced map

$$(\tilde{\varphi}^n)^*_{\mathcal{G}} : H^d(X_n; \mathbb{Q})_{\mathcal{G}_n} \longrightarrow H^d(X_{n+1}; \mathbb{Q})_{\mathcal{G}_{n+1}}$$

is an isomorphism (or injective, surjective, respectively) on cohomology coinvariants. The map

$$(\varphi_n)_* : H_d(X_n; \mathbb{Q}) \longrightarrow H_d(X_{n+1}; \mathbb{Q})$$

is an isomorphism (or injective, surjective, respectively) on homology precisely when the induced map

$$((\tilde{\varphi}_n)_*)_{\mathcal{G}} : H_d(X_n; \mathbb{Q})_{\mathcal{G}_n} \longrightarrow H_d(X_{n+1}; \mathbb{Q})_{\mathcal{G}_{n+1}},$$

is an isomorphism (or injective, surjective, respectively) on homology coinvariants.

In our setting, we will see that representation stability for the sequences

$$\left((\tilde{\varphi}^n)^* : H^d(\widetilde{X}_n; \mathbb{Q}) \to H^d(\widetilde{X}_{n+1}; \mathbb{Q}) \right)_n \text{ and } \left((\tilde{\varphi}_n)_* : H_d(\widetilde{X}_n; \mathbb{Q}) \to H_d(\widetilde{X}_{n+1}; \mathbb{Q}) \right)$$

will imply that the induced maps on coinvariants $(\tilde{\varphi}^n)^*_{\mathcal{G}}$, $((\tilde{\varphi}_n)_*)_{\mathcal{G}}$ are isomorphisms in a stable range, and hence by Proposition 7.32 deduce (co)homological stability for the sequences

$$\left((\varphi^n)^*: H^d(X_n; \mathbb{Q}) \to H^d(X_{n+1}; \mathbb{Q})\right)_n \text{ and } \left((\varphi_n)_*: H_d(X_n; \mathbb{Q}) \to H_d(X_{n+1}; \mathbb{Q})\right)_n.$$

The following result is a special case of, for example, Gadish [Gad17a, Theorem 4.5]. In the case of FI_{G} modules it is also possible to deduce the statement directly, using the description of the induced modules (Proposition 3.22 part (x)), and the formula for their decomposition into irreducible constituents (stated in Corollary 2.18).

Proposition 7.33 (Gadish [Gad17a, Lemma 4.3 and Theorem 4.5]). Let G be a finite group. Let V be an induced FI_G-module (in the sense of Definition 3.20) generated in degree $\leq r$. Then the FI_G-module structure on V induces a FI_G-module structure on its sequence of coinvariants $(V_n)_{G \wr S_n}$. All morphisms act on the coinvariants by injective maps, and for $n \geq r$ the induced maps

$$(V_n)_{G\wr S_n} \longrightarrow (V_{n+1})_{G\wr S_{n+1}}$$

are isomorphisms.

7.5.3. Rational (co)homological stability for unordered vertical configuration spaces. Consider $\tilde{\mathcal{V}}_{n}^{k}(\mathbb{R}^{p,q})$ as a $((k!)^{n} \cdot n!)$ -sheeted cover of $\mathcal{V}_{n}^{k}(\mathbb{R}^{p,q})$ defined by the covering space action of the wreath product $S_{k} \wr S_{n}$. We observe that these covering spaces and our stabilization maps satisfy the framework of Section 7.5.2. We obtain the following corollary, our classical (co)homological stability result for the rational (co)homology of the unordered vertical configuration spaces.

Corollary 7.34. Fix $p \ge 0$, $q \ge 2$, and (co)homological degree d. The (co)homology of the unordered vertical configuration spaces, $(H_d(\mathcal{V}_n^k(\mathbb{R}^{p,q});\mathbb{Q}))_n$ and $(H^d(\mathcal{V}_n^k(\mathbb{R}^{p,q});\mathbb{Q}))_n$, stabilize. Specifically, the maps on (co)homology induced by the stabilization maps ϕ_n are always injective, and are isomorphisms for $n \ge \left\lfloor \frac{2d}{q-1} \right\rfloor$.

Proof. This result follows by combining our main result Theorem 7.24 with Propositions 7.32 and 7.33. \Box

References

- [Aig07] Martin Aigner, A course in enumeration, Springer, 2007.
- [Arn70a] V. I. Arnold, Certain topological invariants of algebraic functions, Trudy Moskov. Mat. Obšč. 21 (1970), 27–46. MR 274462
- [Arn70b] Vladimir Igorevich Arnol'd, Certain topological invariants of algebraic functions, Trudy Moskovskogo Matematicheskogo Obshchestva 21 (1970), 27–46.
- [BK22] Andrea Bianchi and Florian Kranhold, Vertical configuration spaces and their homology, Q. J. Math. 73 (2022), no. 4, 1279–1306. MR 4520220
- [Cas16] Kevin Casto, FI_G-modules, orbit configuration spaces, and complex reflection groups, 2016.
- [CE15] Thomas Church and Jordan S Ellenberg, *Representation stability references*, arXiv preprint arXiv:1506.01022 (2015).
- [CE17] Thomas Church and Jordan Ellenberg, *Homology of* FI-modules, Geometry & Topology **21** (2017), no. 4, 2373–2418.
- [CEF15] Thomas Church, Jordan S. Ellenberg, and Benson Farb, FI-modules and stability for representations of symmetric groups, Duke Math. J. 164 (2015), no. 9, 1833–1910. MR 3357185
- [CEFN14] Thomas Church, Jordan S Ellenberg, Benson Farb, and Rohit Nagpal, FI-modules over Noetherian rings, Geometry & Topology 18 (2014), no. 5, 2951–2984.
- [CF13] Thomas Church and Benson Farb, Representation theory and homological stability, Adv. Math. 245 (2013), 250–314. MR 3084430
- [Chu12] Thomas Church, Homological stability for configuration spaces of manifolds, Inventiones mathematicae 188 (2012), no. 2, 465–504.
- [Cli37] Alfred H Clifford, Representations induced in an invariant subgroup, Proceedings of the National Academy of Sciences 23 (1937), no. 2, 89–90.
- [Far14] Benson Farb, *Representation stability*, arXiv preprint arXiv:1404.4065 (2014).
- [FH13] William Fulton and Joe Harris, Representation theory: a first course, vol. 129, Springer Science & Business Media, 2013.
- [Gad17a] Nir Gadish, Categories of FI type: a unified approach to generalizing representation stability and character polynomials, Journal of Algebra 480 (2017), 450–486.
- [Gad17b] _____, Representation stability for families of linear subspace arrangements, Advances in Mathematics **322** (2017), 341–377.
- [GL15] Wee Liang Gan and Liping Li, Coinduction functor in representation stability theory, J. Lond. Math. Soc. (2) 92 (2015), no. 3, 689–711. MR 3431657
- [GL19] _____, Linear stable range for homology of congruence subgroups via FI-modules, Selecta Math. (N.S.) 25 (2019), no. 4, Paper No. 55, 11. MR 3997138
- [Hat02] Allen Hatcher, Algebraic topology, Cambridge University Press, 2002.
- [Her14] Sarah Herberz, Vertikale konfigurationsraume-homotopie-gruppen fur partitionen der lange zwei, Bachelor's thesis, Rheinische Friedrich-Wilhelms-Universitat Bonn, 2014.
- [HR17] Patricia Hersh and Victor Reiner, Representation stability for cohomology of configuration spaces in \mathbb{R}^d , Int. Math. Res. Not. IMRN (2017), no. 5, 1433–1486, With an appendix written jointly with Steven Sam. MR 3658170
- [IJS08] Frank Ingram, Naihuan Jing, and Ernie Stitzinger, Wreath product symmetric functions, arXiv preprint arXiv:0809.2439 (2008).
- [JK81] Gordon James and Adalbert Kerber, The representation theory of the symmetric group, Encyclopedia of Mathematics and its Applications, vol. 16, Addison-Wesley Publishing Co., Reading, MA, 1981, With a foreword by P. M. Cohn, With an introduction by Gilbert de B. Robinson. MR 644144
- [JRW22] Rita Jiménez Rolland and Jennifer C. H. Wilson, Stability properties of moduli spaces, Notices Amer. Math. Soc. 69 (2022), no. 4, 522–533. MR 4398059
- [Kra21] Florian Kranhold, Configuration spaces of clusters as E_d-algebras, arXiv preprint arXiv:2104.02729 (2021).
- [Lat17] Genta Latifi, Vertical configuration spaces and homological stability, Master's thesis, Rheinische Friedrich-Wilhelms-Universität Bonn, 2017.
- [McD75] Dusa McDuff, Configuration spaces of positive and negative particles, Topology 14 (1975), 91-107. MR 358766
- [Pal21] Martin Palmer, Homological stability for moduli spaces of disconnected submanifolds, i, Algebraic & Geometric Topology 21 (2021), no. 3, 1371–1444.
- [PS17] Andrew Putman and Steven V Sam, Representation stability and finite linear groups, Duke Mathematical Journal 166 (2017), no. 13, 2521 – 2598.
- [Put15] Andrew Putman, Stability in the homology of congruence subgroups, Inventiones mathematicae 202 (2015), 987– 1027.
- [Ram18] Eric Ramos, Homological invariants of FI-modules and FI_G-modules, Journal of Algebra 502 (2018), 163–195.
- [Ros14] Christian Martin Wilhelm Rosner, Vertikale konfigurationsraume-fadell-neuwirth fasserung und eilenberg-maclane, Bachelor's thesis, Rheinische Friedrich-Wilhelms-Universitat Bonn, 2014.
- [Sam17] Steven V Sam, Notes for Math 847: Representation Stability.
- $[Sam20] \qquad \underline{\qquad}, Structures in representation stability, Notices of the American Mathematical Society 67 (2020), no. 1, 38-44.$
- [Seg79] Graeme Segal, The topology of spaces of rational functions, Acta Math. 143 (1979), no. 1-2, 39–72. MR 533892
- [Sno13] Andrew Snowden, Syzygies of Segre embeddings and Δ -modules, Duke Math. J. **162** (2013), no. 2, 225–277. MR 3018955
- [Sno19] _____, Algebraic structures in representation stability, 2019.

- [SS12] Steven V Sam and Andrew Snowden, Introduction to twisted commutative algebras, arXiv preprint arXiv:1209.5122 (2012).
- [SS15] _____, Stability patterns in representation theory, Forum of Mathematics, Sigma, vol. 3, Cambridge University Press, 2015, p. e11.
- [SS17] Steven V. Sam and Andrew Snowden, Gröbner methods for representations of combinatorial categories, J. Amer. Math. Soc. 30 (2017), no. 1, 159–203. MR 3556290
- [SS19] _____, Representations of categories of G-maps, J. Reine Angew. Math. 750 (2019), 197–226. MR 3943321
- [Ste17] Itamar Stein, The Littlewood-Richardson rule for wreath products with symmetric groups and the quiver of the category $F \wr FI_n$, Communications in Algebra 45 (2017), no. 5, 2105–2126.
- [Wil12] Jennifer Wilson, Representation stability for the cohomology of the pure string motion groups, Algebraic & Geometric Topology **12** (2012), no. 2, 909–931.
- [Wil14] Jennifer CH Wilson, FI_W-modules and stability criteria for representations of classical Weyl groups, Journal of Algebra **420** (2014), 269–332.
- [Wil15] _____, FI_W-modules and constraints on classical Weyl group characters, Mathematische Zeitschrift **281** (2015), no. 1, 1–42.
- [Wil18] Jennifer C. H. Wilson, An introduction to FI-modules and their generalizations, http://www.math.lsa.umich.edu/ jchw/FILectures.pdf, 2018, Summer School Lecture Notes.

Email address, David Baron: jdf5@williams.edu

WILLIAMS COLLEGE, MATHEMATICS & STATISTICS, WACHENHEIM SCIENCE CENTER, 18 HOXSEY STREET, WILLIAMSTOWN, MA $01267~\mathrm{USA}$

Email address, Urshita Pal: urshita@umich.edu

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, 530 CHURCH ST, ANN ARBOR MI, 48109, USA

Email address, Chenglu Wang: chengluw@sas.upenn.edu

UNIVERSITY OF PENNSYLVANIA, DEPARTMENT OF MATHEMATICS, DAVID RITTENHOUSE LAB., 209 SOUTH 33rd Street, Philadelphia, PA 19104-6395

Email address, Jennifer Wilson: jchw@umich.edu

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, 530 CHURCH ST, ANN ARBOR MI, 48109, USA

Email address, Chunye Yang: chunye@umich.edu

UNIVERSITY OF MICHIGAN, DEPARTMENT OF MATHEMATICS, 530 CHURCH ST, ANN ARBOR MI, 48109, USA

28