LENGTH OF CLOSED GEODESICS ON RIEMANNIAN MANIFOLDS WITH GOOD COVERS

ZHIFEI ZHU

ABSTRACT. In this article, we prove a generalization of our previous result in [12]. In particular, we show that for an *n*-dimensional, simply-connected Riemannian manifold with diameter D and volume V. Suppose that Madmits a good cover consisting of N elements. Then the length of a shortest closed geodesic on M is bounded by some function which only depends on V, D and N.

1. INTRODUCTION

In this paper, we extend the results established in [12]. Specifically, we prove the following:

Theorem 1.1. Let M be a complete, n-dimensional, simply-connected Riemannian manifold with diameter D and volume V. Suppose that M admits a good cover consisting of N elements. Then the length of the shortest closed *geodesic* on M satisfies:

length $\leq A(n, V, D, N)$,

where A is a function that only depends on n, V, D and N.

The concept of a good cover used in Theorem 1.1 is defined as follows:

Definition 1.2. Let M be a closed, n-dimensional, simply-connected Riemannian manifold with diameter D and volume V. An open cover $\mathcal{O} = \{B_i\}_{i \in I}$ of M is said to be a good cover if there exists a refinement $\tilde{\mathbb{O}} = \{A_i\}_{i \in J}$ that satisfies the following conditions:

- Any two points in A_i can be joined by a curve of length $\leq F_A(V, D)$ within A_i .
- Any two points in B_j can be joined by a curve of length $\leq F_B(V, D)$ within B_j .
- If the intersection $A_i \cap A_j \neq \emptyset$, then the union $A_i \cup A_j$ is contained in some $B_k \in \mathcal{O}$.
- For any closed curve $\gamma : S^1 \to B_j \in \mathcal{O}$, there is a contraction $H : S^1 \times [0,1] \to B_j$ of γ , i.e., $H(t,0) = \gamma(t)$ and $H(t,1) = \text{point} \in B_j$,

such that

$$v_H \le G(V, D).$$

• The number of elements in \tilde{O} and O are both bounded by some N.

Remark 1.3. Sometimes, we take $F(V, D) = \max\{F_A(V, D), F_B(V, D)\}$ as a universal upper bound for an arc connecting points within these open sets. If, in particular, N = N(V, D), then, obviously, the function A = A(V, D) in Theorem 1.1.

In Section 2.2, we provide some examples of manifolds that admit a good cover, especially the cases of four-dimensional non-collapsed manifolds with bounded diameter and Ricci curvature considered in [12]. In particular, in Definition 1.2, we do not have restrictions on the lower bound of diameters of the sets A_i or B_j . In other words, it is possible that for some A_i or B_j , their diameters are arbitrarily close to zero.

While the elements A_i or B_j may not be geodesically convex, it is helpful to consider these elements as the $C^{1,\alpha}$ -images of some convex domains or geodesic balls. In fact, if they are constructed to be geodesically convex, then one can take the obvious bound F = D for arcs connecting two points in these sets.

Intuitively, the refinement " A_i " describes how the elements " B_i " intersect each other. Therefore, one may ask about the relation between the number of diffeomorphic types in some collection of manifolds

 $\mathcal{M} = \{ M \text{ admits a good cover} \},\$

and the given parameters V, D and N. Currently, we do not have an affirmative answer for this relation, which can be summarized in the following question.

Question 1. Is it true that if a collection of simply-connected Riemannian manifolds with diameter D and volume V only consists of N diffeomorphic types, then the length of a shortest closed geodesic can be bounded by some function of D, V, and N?

The first study of the length of a shortest closed geodesic on Riemannian manifolds (which are homeomorphic to T^2 or $\mathbb{R}P^2$) is due to C. Leowner and P. Pu, appears in the work [9] of Pu. For simply-connected manifolds, C. Croke [2] obtained the first upperbound for Riemannian S^2 . M. Gromov asked in [3] whether the length of a shortest periodic geodesic in a *n*-dimensional Riemannian manifold M^n can be bounded by $c(n) \operatorname{vol}(M)^{1/n}$. Similar questions can be asked for the diameter D of the manifold. There are various theories developed to study this question, one may refer to [5] for an overview of this topic.

It is a theorem of Lusternik and Fet that on every compact simply-connected manifold, there exists at least one closed geodesic. (See, for example, [6],

or [7]). The outline of the approach for obtaining an upper bound for the length of this closed geodesic is the following. Let us consider the loop space $\Omega_p M$, which consists of all loops based at a fixed point $p \in M$. Suppose the smallest integer m satisfies $\pi_{m+1}(M) \neq 0$. If one can construct a "small" non-contractible m-dimensional sphere in $\Omega_p M$, that is, a non-contractible map $S^m \to \Omega_p^L M$, where $\Omega_p^L M$ denotes the subspace of $\Omega_p M$ consisting of loops of length $\leq L$, then, by a Morse-type argument, there exists a closed geodesic of length $\leq L$. This geodesic arises as a critical point of the length functional on the free loop space ΛM .

A. Nabutovsky and R. Rotman demonstrated in [8] that the primary obstruction to constructing these "small" non-contractible spheres is the existence of certain "short" closed geodesics on M (see [8, Corollary 5.4]). To formalize these ideas, they introduced the notions of the depth of a loop ([8, Definition 7.1 & 7.4]) and the width of a homotopy.

Definition 1.4 (Depth of a loop). Let M be a closed *n*-dimensional simplyconnected Riemannian manifold with diameter D, and let $\gamma : S^1 \to M$ be a loop in M based at p. We define the depth $S(\gamma)$ of γ as the infimum of positive numbers S such that γ is contractible by a path homotopy through loops of length $\leq \text{length}(\gamma) + S$.

We also define $S_p(M, L)$ as

$$S_p(M, L) = \sup_{\text{length}(\gamma) \le L} S(\gamma),$$

where the supremum is taken over all loops γ of length $\leq L$ based at p.

Definition 1.5 (Width of a homotopy). Let M be a Riemannian manifold, and let $\gamma_i : [0,1] \to M$, i = 1, 2, be two curves in M. Suppose γ_1 and γ_2 are homotopic, and let $H : [0,1] \times [0,1] \to M$ be a homotopy between γ_1 and γ_2 . For each fixed $s \in [0,1]$, denote

$$H_s = H(s, \cdot) : [0, 1] \to M,$$

the curve describing the trajectory of a point H(s, 0) during the homotopy. We define the width ω_H of the homotopy H as

$$w_H = \max_{s \in [0,1]} \text{ length of } H_s$$

And in particular,

Theorem 1.6. If for any closed curve γ of length bounded by L, there exists a contraction of γ with width bounded by some constant W, then

$$S_p(M, L) \le \max_{3} \{2L, 2W + 2D\}.$$

The proof of this inequality can be found in [8, Section 8]. In [8], by taking the base point p = q = x in Theorem 7.3 and applying Corollary 5.4, Nabutovsky and Rotman proved the following result:

Theorem 1.7. Let M^n be a closed Riemannian manifold of diameter D, and let p be a point in M. Suppose $S \ge 0$, and there exists $k \in \mathbb{N}$ such that there is no geodesic loop of length in ((2k-1)D, 2kD] based at p, which is a local minimum of the length functional on $\Omega_p M$ of depth > S. Then for every positive integer m, any map $f : S^m \to \Omega_p M$ is homotopic to a map $\tilde{f}: S^m \to \Omega_p^{L+o(1)}M$, where

$$L = ((4k+2)m + (2k-3))D + (2m-1)S.$$

In this case, the length of the shortest closed geodesic on M does not exceed

$$L = ((4k+2)m + (2k-3))D + (2m-1)S.$$

Therefore, it is enough to show that

Theorem 1.8. Let M be a complete, n-dimensional, simply-connected Riemannian manifold with diameter D and volume V that admits a good cover consisting of N elements. Suppose that there is no closed geodesic on M with length bounded by 3D, then the for every closed curve $\gamma : S^1 \to M$, there is a homotopy H that contracts γ with

$$w_H \le B(V, D, N),$$

where B is a function that only depends on V, D and N.

2. Nerve of the cover and simplicial approximation

If a manifold admits a good cover, we may then approximate every contraction of the closed curves in the manifold with homotopies through some particular loops, i.e., cycles in the 1-skeleton of the nerve of the covering $\tilde{\mathcal{O}}$. More precisely, let us consider the following.

Definition 2.1. Let M be a closed Riemannian manifold and $\tilde{O} = \{A_i\}_{i=1}^N$ the refinement of a good covering as in Definition 1.2. We associate to (M, \tilde{O}) a graph Σ , which is essentially the nerve of the covering \tilde{O} , constructed in the following way.

- (1) For each element $A_i \in \mathcal{O}$, we pick a point $x_i \in A_i$ as a vertex in Σ .
- (2) If for some $1 \le i < j \le N$, the open sets $A_i \cap A_j \ne \emptyset$, then we connect the corresponding vertices x_i and x_j with curve γ_{ij} of length bounded by $2F_A(V, D)$ whose image is contained in $A_i \cup A_j$.

In fact, if $A_i \cap A_j \neq \emptyset$, one may pick a point z in $A_i \cap A_j$ and then the vertices x_i and x_j can be connected by arcs from x_i to z and from z to x_j .

Remark 2.2. Note that the number of the vertices in Σ equals to N, which is the number of the elements in \mathcal{O} . And the number of distinct edges in Σ is at most $\binom{2}{N}$.

Definition 2.3. A simplicial curve α in Σ is a simplicial map $\alpha : [0, 1]_{\Delta} \rightarrow \Sigma$, where $[0, 1]_{\Delta}$ is constructed by partitioning the interval [0, 1] into L subintervals with $0 = t_0 < \cdots < t_L = 1$, and $L \ge 1$ is an integer. We define the simplicial length $m(\alpha)$ of α to be the number of edges in α . In other words, $m(\alpha) = L$. When $\alpha(0) = \alpha(1)$, we refer to α as a loop within Σ .

Note that since Σ consists of edges being curves in M, one may always view $\alpha : [0,1] \to \Sigma$ as a map $\alpha : [0,1] \to M$ through the natural embedding $\Sigma \hookrightarrow M$. Through out this section, we assume that our manifold (M,g) admits a good cover in the sense of Definition 1.2 with diameter D and volume V.

Given a smooth curve $\gamma : [0,1] \to M$, we first show that there exists a simplicial curve α in $\Sigma \hookrightarrow M$ which is homotopic to γ though a homotopy with bounded width. The curve α is called a simplicial approximation of the curve γ . The construction in Lemma 2.4 were originally used in the work [10] of R. Rotman.

Lemma 2.4. For any curve $\gamma : [0,1] \to M$, there exists a simplicial curve $\alpha : [0,1] \to \Sigma$ such that γ is homotopic to α through a homotopy H of width $w_H \leq G_{\Sigma}(V,D)$.

Proof. Let us denote by $\mathcal{O} = \{A_i\}$ the refined cover of the manifold as before. Since the image of γ is path-connected, we may choose a sufficiently fine subdivision of [0,1], say, $\mathcal{O} = t_0 < t_1 < \cdots < t_T = 1$, such that if $\gamma(t_i) \in A_k$ and $\gamma(t_{i+1}) \in A_l$, then the intersection $A_k \cap A_l$ is nonempty, and the segment $\gamma([t_i, t_{i+1}]) \subset A_k \cup A_l$.

Now we may construct the simplicial approximation in the following way. Let us connect $\gamma(t_i)$ with x_k by an arc σ_{ik} of length $\leq F_A$, where x_k is the vertex in Σ corresponding to A_k . (and connect $\gamma(t_{i+1}), x_l$ by $\sigma_{i+1,k}$, resp.) Then $\gamma([t_k, t_{k+1}]) \cup -\sigma_{i+1,k} \cup -\gamma_{kl} \cup \sigma_{ik}$ forms a loop in $A_k \cup A_l$. By the assumption of the good cover, there is a contraction of this loop in some B_s with width $\leq G$, which, in particular, induces a homotopy between $\gamma([t_i, t_{i+1}])$ and $\sigma_{i+1,k} \cup \gamma_{kl} \cup -\sigma_{ik}$ with width $\leq G + 2F_A$.

By applying this to all the intervals in the subdivision, and observe that for consecutive intervals, the arc $\sigma_{ik} \cup -\sigma_{ik}$ can be contracted to x_k through a homotopy with width $\leq 2F_A$. Therefore, we conclude that γ is homotopic to $\alpha := \cup \gamma_{ik}$ though a homotopy with width $\leq G_{\Sigma}(V, D) := G + 4F_A$. \Box

We will present several essential technical lemmas, apart from the above Simplicial Approximation Lemma 2.4, that will be used in the proof of our main results. It's worth noting that these results aligns with those outlined in our paper [12]. While we are not providing detailed proofs here, we will present these statements for the sake of completeness.

Lemma 2.5. Let $\gamma : [0,1] \to M$ be a minimizing geosdesic, and α its approximation as in Lemma 2.4. Then the simplicial length of α is bounded by Z(V, D).

Lemma 2.6. Let $\gamma : [0,1] \to M$ be a closed curve, and α its approximation as in Lemma 2.4. Then there are explicit functions X(V,D), W(V,D) and $\delta(M)$, such that if the simplicial length $m(\alpha) > X(V,D)$, then γ is homotopic to $\gamma_1 \cup \gamma_2 \cup \cdots \cup \gamma_k$ through a homopoty $H : [0,1] \times [0,1] \to M$ which satisfies the following.

- (1) The width of the homotopy $w_H \leq W(V, D)$.
- (2) If α_i is the simplicial approximation of each γ_i , then $m(\alpha_i) \leq X(V, D)$.
- (3) The length of each $\gamma_i \leq \text{length}(\gamma) \delta$.

Lemma 2.7. Let $\gamma : [0,1] \to M$ be a closed curve. Suppose that $\gamma = \bigcup_{i=1}^{n} \gamma_i$, where each γ_i is a closed curve with base point p. If each γ_i can be contracted to a point in M through a homotopy with width bounded by W_i , then there exists a homotopy $H : [0,1] \times [0,1] \to M$ such that $H(t,0) = \gamma(t)$ and H(t,1) = p. And the width ω_H of this homotopy is bounded by $2 \cdot \max_i W_i$.

Remark 2.8. The statement of the above lemmas does not depend on the number of the elements N in the good cover.

Our main theorem states that if a Riemannian manifold admits a good cover in the above sense, then the length of a shortest closed geodesic, can be bounded by some function depending on F, G and N. In fact, there are several classes of manifolds that admit such type of cover.

2.1. Manifolds with injectivity radius lower bound. If we consider manifolds with injectivity radius bounded from below by some constant r_0 , for example, manifolds with sectional curvature $\leq k$, vol $\leq V$, then in this case, we may construct a good cover with $F = r_0$ and $G = G(r_0(V, D, k))$. In this case, it is showed by R. Rotman in [10], that if M has a non-trivial second homology group, then the length of a shortest closed geodesic is bounded above. Our Theorem 1.1 does not improve this result. It is worth mentioning that the assumption on the second homology group can be removed by A. Nabutovsky and R. Rotman in [8].

2.2. Non-collapsed manifolds with Ricci curvature bounds. In [12], we have constructed a good cover for four-dimensional manifolds whose diameter $\leq D$, volume $\geq v$ and |Ric| < 3. Let us briefly describe the process here. In [1], Cheeger and Naber showed that for any non-collapsed manifold of dimension 4,

with bounded Ricci curvature, there is a structural theorem called bubble-tree decomposition.

Theorem 2.9 ([1], Theorem 8.64). Let M be a 4-dimensional Riemannian manifold with $|Ric| \leq 3$, vol(M) > v > 0 and $diam(M) \leq D$. Then M admits a decomposition into bodies and necks

$$M = \mathcal{B}^1 \cup \bigcup_{j_2=1}^{N_2} \mathcal{N}_{j_2}^2 \cup \bigcup_{j_2=1}^{N_2} \mathcal{B}_{j_2}^2 \cup \dots \cup \bigcup_{j_k=1}^{N_k} \mathcal{N}_{j_k}^k \cup \bigcup_{j_k=1}^{N_k} \mathcal{B}_{j_k}^k$$

such that the following conditions are satisfied:

- (1) If $x \in \mathcal{B}_i^j$, then $r_h(x) \geq r_0(v, D) \cdot \operatorname{diam}(\mathcal{B}_i^j)$, where r_h is the harmonic radius and r_0 is a constant that only depends on v and D.
- (2) Each \mathcal{N}_i^j is diffeomorphic to $\mathbb{R} \times S^3 / \Gamma_i^j$ for some $\Gamma_i^j \subset O(4)$ with the order $|\Gamma_i^j| < c(v, D).$
- (3) $\mathcal{N}_{i}^{j} \cap \mathcal{B}_{i}^{j}$ is diffeomorphic to $\mathbb{R} \times S^{3}/\Gamma_{i}^{j}$. (4) $\mathcal{N}_{i}^{j} \cap \mathcal{B}_{i'}^{j-1}$ is either empty or diffeomorphic to $\mathbb{R} \times S^{3}/\Gamma_{i}^{j}$.
- (5) Each $N_i \leq n(v, D)$ and $k \leq k(v, D)$.

It is constructed in [12] that, in this case, the manifold M admits a good cover in the sense of Definition 1.2. Namely,

Lemma 2.10. Let M be a 4-dimensional Riemannian manifold which satisfies the conditions of Theorem 2.9. Then there is an open cover \mathfrak{O} of the manifold and its refinement \mathfrak{O} in the sense of Definition 1.2 and Remark 1.3 with F =3D and G = 21D. In other words,

- Any x, y in $A \in \tilde{O}$ can be connected by a curve with length less than 3D.
- For any closed curve $\gamma: [0,1] \to B \in \mathcal{O}$ with $\gamma(0) = \gamma(1) = p$, there is a contraction $H: [0,1] \times [0,1] \rightarrow B$ with $H(t,0) = \gamma(t)$ and H(t,1) = psuch that $w_H \leq 21D$.

Moreover, the total number of elements in \mathfrak{O} is bounded by some N(V, D) which can be computed by the constants in Theorem 2.9.

The construction of this cover can be found in Section 2 of [12]. Here we omit the detailed proof but instead we describe the idea. By Theorem 2.9, Madmits a cover with the body regions \mathcal{B}_i^j and the neck regions \mathcal{N}_i^j .

In each body \mathcal{B}_i^j , for any $x \in \mathcal{B}_i^j$, the harmonic radius $r_h(x) \ge r_0 \cdot \operatorname{diam}(\mathcal{B}_i^j)$ is in proportional to the diameter of the body. Let $R(x) = \frac{r_h(x)}{8 \cdot (1+\varepsilon)}$. It is show in [, Lemma 2.2] that every closed curve in $B_R(x)$ can be contracted within $B_R(x)$ through a homotopy of width $\leq D$. We may pick a cover $\cup_k B_{R(x_k)/4}(x_k) \supseteq \mathcal{B}_i^j$ such that $B_{R(x_k)/16}(x_k)$ are pairwise disjoint. This guarantees the number of the elements in this collection is bounded by some $N_{\mathcal{B}_{i}^{j}}(V, D)$. Note that the collection $\{B_{R(x_k)}\}$ still form a cover of \mathcal{B}_i^j . Therefore, we simply choose $B_{R(x_k)/4}(x_k)$ to be an element in $\tilde{\mathcal{O}}$ and $B_{R(x_k)}(x_k)$ an element in \mathcal{O} .

The neck regions in general do not admit a lower bound on harmonic radius. However, the geometry of \mathcal{N}_i^j is completely understood. In this case, we cover the neck by the image $\{\tilde{T}_{i,k}^j\}$ and $\{T_{i,k}^j\}$ of finitely many convex regions $\{\tilde{K}_{i,k}^j\}$, and their refinements $\{K_{i,k}^j\}$, respectively. Here $k \leq N_{\mathcal{N}_i^j}(V,D)$; $\tilde{K}_{i,k}^j \subset \mathbb{R} \times S^3/\Gamma_i^j$ is convex; and $N_{\mathcal{N}_i^j}(V,D)$ only depends on the order of $|\Gamma_i^j| < c(V,D)$. It is then showed in [, Lemma 2.8] that the cover satisfies Definition 1.2 with F = 3D and G = 21D.

Remark 2.11. If the neck region is empty, e.g., the manifold admits an injectivity radius lower bound which only depends on V and D, then one can pick the cover in the same way as the body region, with G = D or finer estimates depending on the injectivity radius.

3. Proof of the main Theorems

In this section, we will prove our main Theorem 1.1 and Theorem 1.8. The idea of the proof is based on our paper [12]. We first proof Theorem 1.8.

Proof of Theorem 1.8. Let $\gamma : [0,1] \to M$ be a closed curve with $\gamma(0) = \gamma(1) = p$ in M. By Lemma 2.4, there is a loop $\alpha : [0,1]_{\triangle} \to \Sigma$ such that γ is homotopic to α through a homotopy H_1 with $w_{H_1} \leq G_{\Sigma}(V,D)$.

Without lost of generality, we may assume that the simplicial length $m(\alpha)$ does not exceed X(V, D). Otherwise, by Lemma 2.6, the curve γ is homotopic to $\gamma_1 \cup \cdots \cup \gamma_k$, such that if α_i is the approximation of γ_i , then $m(\alpha_i) \leq X(V, D)$. Furthermore, the length of each length $(\gamma_i) \leq \text{length}(\gamma) - \delta(V, D)$, for some $\delta > 0$. Then, by Lemma 2.7, if each γ_i can be contracted through a homotopy of width $\leq W$, then γ can be contracted through a homotopy of width $\leq 2W$. Therefore, it is suffices to contract each γ_i .

Below we will show that γ can be contracted to a point through a homotopy of width bounded by some B(V, D, N).

(0) Birkhoff curve shortening process for free loops (BPFL).

Note that because the length of $\gamma \leq 3D$, under the assumption that there is no closed geodesic whose length is $\leq 3D$, if we apply the Birkhoff curve shortening process for free loops (BPFL) to γ , then there is a length non-increasing contraction of γ .

Let us denote this contraction by $H : [0,1] \times [0,1] \to M$, where $H(t,0) = \gamma(t)$ and H(t,1) = point. We consider a partition of the second interval $[0,1] = s_0 < s_1 < \cdots < s_l = 1$ such that $w_H(s_j, s_{j+1}) \leq D$. And let $\gamma^j(t) := H(t,s_j)$, for every $0 \leq j \leq l$.

(1) First step.

By Lemma 2.4, each γ^j is homotopic to some simplicial $\alpha^j \subset \Sigma$. In particular, by assumption, for the first curve $\gamma^0 := \gamma$, we have $m(\alpha^0) \leq X(V, D)$.

Let γ^{j_0} be the first $j_0 \geq 1$ such that $m(\alpha^{j_0}) > X(V, D)$. Then γ is homotopic to γ^{j_0} through curves $\gamma^1, \gamma^2, \ldots, \gamma^{j_0-1}$. The width of homotopy bewteen each γ^j and γ^{j+1} is less than or equal to D, and the simplicial approximation of each α^j has simplicial length $m(\alpha^j) \leq X(V, D)$, for $j = 0, 1, \ldots, j_0 - 1$.

We now apply Lemma 2.6 to γ^{j_0} . In this case, γ^{j_0} is homotopic to some $\gamma_1^{j_0} \cup \cdots \cup \gamma_k^{j_0}$. Let $\alpha_i^{j_0}$ be the simplicial approximation of $\gamma_i^{j_0}$. By Lemma 2.6, the length of each $\gamma_i^{j_0}$ satisfies

$$\operatorname{length}(\gamma_i^{j_0}) \le \operatorname{length}(\gamma) - \delta$$

And the simplicial length $m(\alpha_i^{j_0}) \leq X(V, D)$.

(2) A generic step.

Now we repeat this construction to the curves $\gamma_i^{j_0}$, which will generate a finite family of curves. (We describe in (3) the parametrization of this family.) Denote by \mathbf{i}_l the indices i_1, i_2, \ldots, i_l . During this process, a generic curve $\gamma_{\mathbf{i}_l} := \gamma_{\mathbf{i}_l}^0$ is homotopic to some $\gamma_{\mathbf{i}_l}^{j_0}$ through curves $\gamma_{\mathbf{i}_l}^1, \ldots, \gamma_{\mathbf{i}_l}^{j_0-1}$ constructed using BPFL, such that if $\alpha_{\mathbf{i}_l}^j$ is the approximation of $\gamma_{\mathbf{i}_l}^j$ as in Lemma 2.4, then $m(\alpha_{\mathbf{i}_l}^j) \leq X(V, D)$, for $j = 0, 1, \ldots, j_0 - 1$ and $m(\alpha_{\mathbf{i}_l}^{j_0}) > X(V, D)$. The width of homotopy between $\gamma_{\mathbf{i}_l}^j$ and $\gamma_{\mathbf{i}_l}^{j+1}$ is bounded by D.

And by Lemma 2.6, the curve $\gamma_{\mathbf{i}_l}^{j_0}$ is homotopic to $\gamma_{\mathbf{i}_l,1} \cup \cdots \cup \gamma_{\mathbf{i}_l,k}$ with the simplicial length of the approximation $\alpha_{\mathbf{i}_l,j}$ of each $\gamma_{\mathbf{i}_l,j}$ is bounded by X(V, D).

This process will terminate if no such j_0 exists, i.e., $\gamma_{\mathbf{i}_l}$ can be contracted to a point through a family of curves $\gamma_{\mathbf{i}_l}^1, \ldots, \gamma_{\mathbf{i}_l}^{j_0-1} = \text{point}$, with simplicial length of the approximation $\leq X(V, D)$.

On the other hand, each time we apply Lemma 2.6,

 $\operatorname{length}(\gamma_{\mathbf{i}_l,j}) \leq \operatorname{length}(\gamma_{\mathbf{i}_l}) - \delta \leq \operatorname{length}(\gamma) - (l+1) \cdot \delta$

Note that the above inequality also implies that

$$l \le \frac{\operatorname{length}(\gamma)}{\delta} - 1,$$

where the right-hand side is finite (but may not be bounded by any functions of V and D). Hence for each "branch" of the curve γ_i , we may apply Lemma 2.6 for at most $\frac{\text{length}(\gamma)}{\delta}$ times before we get a point during the contraction.

(3) Parametrization of the family.

In previous steps, we generate a finite family of curves denoted as $\{\gamma_I\}_{I \in \mathcal{V}(\mathcal{T})}$. This family is parameterized by a finite tree $\mathcal{T} = \{v_i, e_{ij}\}$, where $\mathcal{V}(\mathcal{T}) = \{v_i\}$ is the set of vertices and $\mathcal{E}(\mathcal{T}) = \{e_{ij}\}$ the set of edges. We denote by $h(\mathcal{T})$ the height of a finite tree \mathcal{T} . The tree \mathcal{T} is constructed as follows.

- (a) The root v_0 of the tree is defined to be the curve γ_i , so that the simplicial length of the approximation $m(\alpha_i) \leq X(V, D)$.
- (b) Each curve $\gamma_{\mathbf{i}_l}^j$, including the point curves, occurred during the generic step (2) whose approximation has simplicial length bounded by X(V, D) is identified with a vertex $v_t \in \mathcal{V}$. Since the total number of the curve is finite, we can choose the indices such that if $\gamma_{\mathbf{i}_l}^j$ is a subsequent curve of $\gamma_{\mathbf{i}_l}^k$ during the BPFL, then the corresponding vertices v_{t_j} and v_{t_k} satisfy $t_j > t_k$.

In addition, suppose that a curve $\gamma_{\mathbf{i}_l}^{j_0}$ is homotopic to $\gamma_{\mathbf{i}_l,1} \cup \cdots \cup \gamma_{\mathbf{i}_l,k}$. Note that the curve $\gamma_{\mathbf{i}_l}^{j_0}$ is not identified with a vertex. In this case, we require that if the curves $\gamma_{\mathbf{i}_l,i}$, where $1 \leq i \leq k$, and $\gamma_{\mathbf{i}_l}^{j_0-1}$ are identified with the vertices v_{s_1}, \ldots, v_{s_k} and $v_{t_{j_0-1}}$, then $s_i > t_{j_0-1}$.

(c) Suppose that $\gamma_{\mathbf{i}_l}^{j+1}$ is a subsequent curve of $\gamma_{\mathbf{i}_l}^j$ during BPFL process and they are identified with vertices v_{t_j} , $v_{t_{j+1}}$, respectively. We define an edge $e_{t_j,t_{j+1}} \in \mathcal{E}$ joining these two vertices.

Similarly, if $\gamma_{\mathbf{i}_l}^{j_0}$ is homotopic to $\gamma_{\mathbf{i}_l,1} \cup \cdots \cup \gamma_{\mathbf{i}_l,k}$, and the curves $\gamma_{\mathbf{i}_l,1}, \ldots, \gamma_{\mathbf{i}_l,k}$ and $\gamma_{\mathbf{i}_l}^{j_0-1}$ are identified with the vertices v_{s_1}, \ldots, v_{s_k} and $v_{t_{j_0-1}}$, respectively. Then we define an edge $e_{t_{j_0-1},s_i} \in \mathcal{E}$ joining $v_{t_{j_0-1}}$ with each v_{s_i} , for $1 \leq i \leq k$.

As a consequence of the construction, the family of curves $\{\gamma_I\}_{I \in \mathcal{V}}$ satisfies the following property.

Lemma 3.1. Let $\{\gamma_I\}_{I \in \mathcal{V}(\mathcal{T})}$ be a finite family of the curves constructed from above. Then the simplicial length of the approximation of each γ_I is bounded by X(V, D), and in particular, we have the following.

- (a) If a vertex $v_j \in \mathcal{V}$ is a leaf, i.e., it has no children, then the corresponding curve γ_{v_j} is a point curve in M.
- (b) If a vertex v_j has a single child v_s , then the corresponding curve γ_{v_j} is homotopic to γ_{v_s} through a homotopy of width bounded by D.
- (c) If a vertex v_j has k children v_{s_1}, \ldots, v_{s_k} , then the corresponding curve γ_{v_j} is homotopic to $\gamma_{v_{s_1}} \cup \cdots \cup \gamma_{v_{s_k}}$ through a homotopy of width bound by D+W(V,D), where W(V,D) is the same function as in Lemma 2.6.

(4) Homotopy with marked points.

Given a closed curve γ and its associated family of curves $\{\gamma_I\}_{I \in \mathcal{V}(\mathcal{T})}$ as constructed above, we now develop a new homotopy based on this family which contracts γ to a marked point $p \in \gamma$. This construction can be viewed as a generalization of Lemma 2.7. The construction will be done inductively as the following.

Without lost of generality, we assume a curve $\gamma_i \in {\gamma_I}_{I \in \mathcal{V}(\mathcal{T})}$ corresponds to a vertex $v_i \in \mathcal{V}$. And we denote by p_i a marked point in γ_i . We first discuss in two cases according to the cases in Lemma 3.1. The choice of the marked points on subsequent curves will also be based on this construction.

(a),(b) If v_i has a single child v_j , let γ_i, γ_j be the corresponding curves, and $p_i \in \gamma_i$ is the marked point. We denote by H(t, s) the homotopy between γ_i and γ_j , where $H(t, 0) = \gamma_i(t)$ and $H(t, 1) = \gamma_j(t)$. By assumption, the width of this homotopy $w_H \leq D$.

Let $\sigma_i : [0, 1] \to M$ be the trajectory of the marked point p_i under H. In other words, if $p_i = \gamma(t_0)$, then $\sigma_i(s) = H(t_0, s)$. We define the marked point p_j on γ_j to be the other endpoint of σ_i , i.e., $p_j := H(t_0, 1)$.

Then γ_i is homotopic to $\sigma_i \cup \gamma_j \cup -\sigma_i$ through $\sigma_i([0, t]) \cup H([0, 1], t) \cup -\sigma_i([0, t]), 0 \le t \le 1$. The width of this homotopy is bounded by $\max\{2 \operatorname{length}(\sigma_i), w_H\} \le 2D$.

(c) Similarly, if v_i has k children v_{s_1}, \ldots, v_{s_k} . Let H(s, t) be the homotopy between the corresponding curves γ_i and $\gamma_{v_{s_1}} \cup \cdots \cup \gamma_{v_{s_k}}$. Let $p_i \in \gamma_i$ be the marked point. By Lemma 2.6, the curves $\{\gamma_{v_{s_1}}, \ldots, \gamma_{v_{s_k}}\}$ has a common base point p_s . In this case, we choose the marked point $p_{s_j} = p_s$ on each γ_{s_j} . However, p_s may not be the destination of p_i under H. Therefore, if $\gamma_i(t_0) = p_i$ and $H(t_1, 1) = p_s$, for some t_0, t_1 , w.l.o.g, assume $t_1 \geq t_0$, we first apply a rotation R(s, t) = H(t + s, 0) of γ_i , for $s \in [0, s_0]$, so that $R(t_0, 0) = p_i$ and $R(t_0, s_0) = H(t_1, 0)$. The width $w_R \leq \frac{1}{2} \operatorname{length}(\gamma) \leq \frac{3}{2}D$.

Let $\sigma_i : [0, 1 + s_0] \to M$ be the concatenation of the trajectory of p_i under the rotation R and homotopy H, where $\sigma_i(0) = p_i$ and $\sigma_i(1 + s_0) = p_s$. Then γ_i is homotopic to $\sigma_i \cup (\bigcup_{j=1}^k \gamma_{s_j}) \cup -\sigma_i$, through the curves $\sigma([0, s_0 + t]) \cup H([0, 1], t) \cup -\sigma_i([0, s_0 + t])$. In this case the length of σ_i is bounded by $\frac{3}{2}D + D + W(V, D)$ and hence the width of this homotopy is bounded by 5D + 2W(V, D).

Now, inductively, the root curve γ_0 is homotopic to $\sigma_1 \cup \gamma_1 \cup -\sigma_1$, then $\sigma_1 \cup \sigma_2 (\cup \gamma_2 \dots) \cup -\sigma_2 \cup -\sigma_1, \dots$, and $\cup_{i \in \mathcal{V}} (\sigma_i \cup -\sigma_i)$. The width of this homotopy is bounded by $2D \cdot h_1 + (5D + 2W) \cdot h_2$, where h_1, h_2 are the number of steps (a)(b), (c) performed in the above construction, respectively. Note that $h_1 + h_2 = h(\mathfrak{T})$ which is the height of the tree \mathfrak{T} . Finally, one may contract $\bigcup_{i \in \mathcal{V}} (\sigma_i \cup -\sigma_i)$ to the point p. Because each leaf $\sigma_i \cup -\sigma_i$ can be contracted at the same time, therefore, the width of this contraction is again bounded by $2D \cdot h_1 + (5D + 2W) \cdot h_2$, where $h_1 + h_2 = h(\mathfrak{T})$.

(5) Bounded height construction.

To complete the proof of our main theorem, we show that if γ is a closed curve of length $\leq 3D$ and $\{\gamma_{v_I}\}_{v_I \in \mathcal{V}(\mathfrak{T})}$ is the associated family in the above construction, then there is a new family $\{\gamma_{v_I}\}_{v_I \in \tilde{\mathcal{V}}(\tilde{\mathfrak{T}})}$, which still satisfies the conclusion of Lemma 3.1 with the height of $\tilde{\mathfrak{T}}$ bounded in terms of V, D.

Indeed, for each γ_{v_I} , where $v_I \in \mathcal{V}$, let us denote by \mathcal{L}_I the subtree of \mathcal{T} whose root is v_I . First note the following.

Lemma 3.2. Let γ_1 , γ_2 be two closed curves of length $\leq 3D$. And let α_1 , α_2 be their approximation as in Lemma 2.4. If $\alpha_1 = \alpha_2$ up to a reparametrization, then γ_1 is homotopic to γ_2 through a homotopy H with $w_H \leq Y(V, D)$.

Proof. The proof is the same as in Lemma 2.4. Observe that the curve $\gamma([t_k, t_{k+1}]) \cup -\sigma_{i+1,k} \cup -\gamma_{kl} \cup \sigma_{ik}$ can also be contracted to the point $\gamma(t_k)$ with in B_s , and therefore inducing a homotopy between α_2 and γ_2 with width bounded by G_{Σ} . Therefore, one may take $Y(V, D) = 2G_{\Sigma}(V, D)$.

Let us denote by $h(\mathcal{L})$ the height of a subtree $\mathcal{L} \subset \mathcal{T}$. Since Σ is a finite simplicial complex with N vertices and at most N^2 edges, the number N_0 of closed curves, up to reparametrization, whose simplicial length is less than or equal to X(V, D) is bounded by $(N^2 + 1)^X$.

Now if the height $h(\mathfrak{T}) \geq N_0+1$, then there exists $v_1, v_2 \in \mathcal{V}$ such that the approximation of γ_{v_1} and γ_{v_2} are the same, up to reparametrization. And $v_2 \in \mathcal{V}(\mathcal{L}_1)$, i.e. v_2 is in the subtree with root v_1 .

By Lemma 3.2, γ_{v_1} and γ_{v_2} are homotopic with width $\leq Y(V, D)$. In this case, we construct a new tree \mathcal{T}_1 by replacing \mathcal{L}_1 with \mathcal{L}_2 . Note that \mathcal{T}_1 is indeed a tree, i.e., contains no loops, since $v_2 \in \mathcal{V}(\mathcal{L}_1)$.

Denote by v_0 the parent of v_1 . Again, we discuss by cases as in Lemma 3.1.

- (a)(b) If v_1 is the single child of v_0 , then in \mathcal{T}_1 , v_2 is the single child of v_0 and γ_{v_0} is homotopic to γ_{v_2} with width bounded by D + Y(V, D).
 - (c) If v_0 has k children $v_{s_1}, \ldots, v_1, \ldots, v_{s_{k-1}}$, then in $\mathfrak{T}_1, \gamma_{v_0}$ is homotopic to $\gamma_{v_{s_1}} \cup \cdots \cup \gamma_{v_2} \cup \cdots \cup \gamma_{v_{s_{k-1}}}$ through a homotopy of width bounded by D + Y(V, D) + W(V, D).

Observe that even though we reduced the height of a branch of v_0 , in general, the height $h(\mathfrak{T}_1) \leq h(\mathfrak{T})$, where the equality may happen. However, if the height $h(\mathfrak{T}_1) \geq N_0 + 1$, then we may still apply the above construction to replace a subtree until $h(\mathfrak{T}_k) \leq N_0 \leq (N^2 + 1)^X$.

On the other hand, if in \mathcal{T}_1 , there is some $v_3 \in \mathcal{L}_{v_2}$ such that γ_{v_3} and γ_{v_2} have the same approximation, then in particular, γ_{v_1} and γ_{v_3} also have the same approximation. Hence, in this case v_0 can be connected to v_3 with homotopies of width bounded by D + Y or D + Y + W, respectively for the cases (a)(b) and (c).

And therefore by (4) and (5), we have proved that the total width of contracting the curve γ to a point is bounded by

$$B(V, D, N) \le (N^2 + 1)^X \cdot (7D + 2Y + W).$$

We may now finish the proof of Theorem 1.1.

Proof of Theorem 1.1. Let M be a manifold satisfies the condition in Theorem 1.1. Suppose that there is no closed geodesic of length $\leq 3D$. Then Theorem 1.8 implies that every loop in M may be contracted via a homotopy with width bounded by B(V, D, N). This further implies by Theorem 1.6 that the depth $S_p(M, 3D) \leq \max\{3D, 2B + 2D\}$.

As before, let $\Omega_p(M)$ be the space of continuous maps $\{S^1 \to M\}$ based at $p \in M$ and $\Omega_p^L M$ the subspace where every loop is of length $\leq L$. Now by taking the integer k = 2 in Theorem 1.7, we conclude that for every positive integer m, every map $f: S^m \to \Omega_p M$ is homotopic to a map $\tilde{f}: S^m \to \Omega_p^F M$, where

$$F = F(m, V, D, N) = 10 \cdot m + D + (2m - 1) \cdot S_p.$$

And in particular, the length of a closed periodic geodesic does not exceed F(m, V, D, N).

Because our manifold is simply-connected, for $l \geq 2$, suppose it is (l - 1)-connected but not l-connected, the above argument shows that there is a closed geodesic of length $\leq F(l, V, D)$. Note that F is increasing in l. If M is (n-1)-connected, then by Hurewicz theorem (see [4, Theorem 4.32]), then $\pi_n(M) \cong H^n(M) \neq 0$, and hence we can take A = F(n, V, D, N) in the above argument.

References

 J. Cheeger and A. Naber, Regularity of Einstein manifolds and the codimension 4 conjecture, Ann. Math. (2) 182, No. 3, 1093–1165 (2015).

- [2] C. Croke, Area and the length of the shortest closed geodesic, J. Differential Geom. 27, No. 1, 1–21 (1988).
- [3] M. Gromov, Filling Riemannian manifolds, J. Differential Geom. 18, 1–147 (1983).
- [4] A. Hatcher, *Algebraic topology*. Cambridge: Cambridge University Press (2002).
- [5] M. Katz, Systolic geometry and topology. With an appendix by Jake P. Solomon. Providence, RI: American Mathematical Society (AMS) (2007).
- [6] W. Klingenberg, Lectures on closed geodesics. Grundlehren der Mathematischen Wissenschaften, Vol. 230. Springer-Verlag, Berlin-New York, (1978).
- [7] J. Milnor. *Morse Theory*. Princeton University Press, (1969).
- [8] A. Nabutovsky and R. Rotman, Length of geodesics and quantitative Morse theory on loop spaces, Geom. Funct. Anal. 23, No. 1, 367–414 (2013).
- [9] Pu, Pao Ming, Some inequalities in certain nonorientable Riemannian manifolds, Pacific J. Math. 2(1), 55–71 (1952).
- [10] R. Rotman, Upper bounds on the length of the shortest closed geodesic on simply connected manifolds, Math. Z. 233, No. 2, 365–398 (2000).
- [11] R. Rotman, The length of a shortest closed geodesic and the area of a 2dimensional sphere, Proc. Am. Math. Soc. 134, No. 10, 3041–3047 (2006)
- [12] N. Wu and Z. Zhu, Length of a shortest closed geodesic in manifolds of dimension four, J. Differential Geom. 122, No. 3, 519–564 (2022).