

# ON POLYNOMIALLY HYPO-EP OPERATORS.

R. SEMMAMI AND H. EZZAHRAOUI

**ABSTRACT.** The purpose of this paper is to present a new class of operators known as polynomially hypo-EP operators, extending the notation of hypo-EP,  $n$ -hypo-EP, and polynomially EP. The paper explores numerous properties and characterizations of polynomially hypo-EP operators and polynomially EP operators. In addition, we characterize polynomially hypo-EP operators applying the adequate operator matrix representations. By employing these findings, we derive new characterizations of EP, hypo-EP,  $n$ -EP and  $n$ -hypo-EP.

## 1. INTRODUCTION

Let  $\mathcal{H}$  be a complex Hilbert space and let  $\mathcal{L}(\mathcal{H})$  represent the space of all bounded linear operators defined on  $\mathcal{H}$  and  $\mathcal{R}(\mathcal{H}) \subset \mathcal{L}(\mathcal{H})$  be the set of operators with closed range. For  $T \in \mathcal{L}(\mathcal{H})$ ,  $T^*$ ,  $N(T)$  and  $R(T)$  stand for its adjoint, kernel and range, respectively. For  $T, S \in \mathcal{L}(\mathcal{H})$ , we write  $[T, S] := TS - ST$  for the commutator of  $S$  and  $T$ .

A closed subspace  $E \subset \mathcal{H}$  is said to be  $T$ -invariant if  $T(E) \subset E$ . We define  $T_E : E \rightarrow E$  as  $T_E x = Tx$ ,  $x \in E$ . Also  $E$  is called reducing for  $T$  if it is invariant for both  $T$  and  $T^*$ , or equivalently, if  $T(E) \subset E$  and  $T(E^\perp) \subset E^\perp$ , where  $E^\perp$  is the orthogonal complement of  $E$ .

For  $T \in \mathcal{R}(\mathcal{H})$ , the Moore-Penrose inverse  $T^\dagger$  of  $T$ , is defined as the unique solution of the following four operator equations:

$$TT^\dagger T = T, \quad T^\dagger TT^\dagger = T^\dagger, \quad (T^\dagger T)^* = T^\dagger T \quad \text{and} \quad (TT^\dagger)^* = TT^\dagger.$$

It follows immediately that

$$TT^\dagger = P_{R(T)}, \quad T^\dagger T = P_{R(T^*)}, \quad N(T^\dagger) = N(T^*) \quad \text{and} \quad N(T^{\dagger*}) = N(T).$$

The Cauchy dual of an operator  $T \in \mathcal{L}(\mathcal{H})$  with closed range is defined as  $\omega(T) = T(T^*T)^\dagger$ . For equivalent definitions and further properties of  $\omega(T)$ , see [8].

The concept of EP matrices was defined by Schwerdtfeger [19] and generalized to Hilbert space operators with closed range in [2]. An operator  $T \in \mathcal{L}(\mathcal{H})$  is called EP if  $R(T)$  is closed and  $R(T) = R(T^*)$ . EP operators are significant because they commute with their Moore–Penrose inverse. Normal operators with a closed range and nonsingular operators are

---

2020 *Mathematics Subject Classification.* 47B15; 47B20; 47A05.

*Key words and phrases.* Moore-Penrose inverse of an operator.  $n$ -EP,  $n$ -hypo-EP,  $n$ -normal, polynomially EP and polynomially hypo-EP operator.

particular cases of EP operators. Several authors have thoroughly examined different established characterizations of EP matrices and EP operators. See for example [1, 2, 3, 7, 20, 9, 12].

An operator  $T \in \mathcal{R}(\mathcal{H})$  is SD (star-dagger), if  $[T^*, T^\dagger] = 0$ . This class been studied in detail by several authors. See [20, 9, 10].

The class of  $n$ -EP matrices was introduced in [18] as a generalization of EP matrices. The notion of  $n$ -EP operators was defined in [21] extending the definition of  $n$ -EP matrices. An operator  $T \in \mathcal{L}(\mathcal{H})$  is called  $n$ -EP and we write  $T \in (n\text{-EP})$  if  $R(T)$  is closed and  $T^n T^\dagger = T^\dagger T^n$  for some  $n \in \mathbb{N}$ . For further properties of  $n$ -EP operators, we refer to [8, 11]. Generalizing the notion of  $n$ -EP operators, polynomially EP operators were presented in [12].

M. Itoh introduced in [13] the class of hypo-EP operators as follows. An operator  $T \in \mathcal{R}(\mathcal{H})$  is an hypo-EP operator if  $[T^\dagger, T] \geq 0$ . Let (HEP) be the class of hypo-EP operators. Several authors have thoroughly examined different established characterizations of hypo-EP matrices and hypo-EP operators. See [1, 20, 17, 22] for detailed study. Some research characterized hypo-EP operators through factorizations and when the product of two hypo-EP matrices or hypo-EP operators is also hypo-EP, was studied. See for example [1, 15, 17].

As an extension of hypo-EP operators,  $n$ -hypo-EP operators were introduced in [20] as follows:  $T \in \mathcal{R}(\mathcal{H})$  is  $n$ -hypo-EP ( $n \in \mathbb{N}$ ) if  $R(T^n) \subseteq R(T^*)$ . Let ( $n$ -HEP) be the class of  $n$ -hypo-EP operators.

The operator  $T$  is called normal if  $[T^*, T] = 0$ . As an extension of normal operators,  $n$ -normal operators were introduced in [14] as follows:  $T \in \mathcal{L}(\mathcal{H})$  is  $n$ -normal ( $n \in \mathbb{N}$ ) if  $[T^n, T^*] = 0$ . Generalizing the notion of  $n$ -normal operators, polynomially normal operators were presented in [5].

We use  $Poly_1$  to denote the set of all complex polynomials in one variable. Taking conjugate coefficients of  $p \in Poly_1$ , we obtain  $\bar{p} \in Poly_1$ :  $\bar{p}(z) := \overline{p(\bar{z})}$ ,  $z \in \mathbb{C}$ . Let  $T \in \mathcal{L}(\mathcal{H})$  and  $p \in Poly_1$  be nontrivial. If  $[p(T), T^*] = 0$ , then we say that  $T$  is  $p$ -normal.  $T \in \mathcal{L}(\mathcal{H})$  is polynomially normal, if  $T$  is  $q$ -normal for some  $q \in Poly_1$ . Observe that, by [5, Lemma 1],  $T \in \mathcal{L}(\mathcal{H})$  is  $p$ -normal if and only if  $p(T)$  is normal. Recent results related to polynomially normal operators can be found in [4].

Let  $T \in \mathcal{L}(\mathcal{H})$  and  $p \in Poly_1$  be nontrivial. Following [12], we say that  $T$  is  $p$ -EP, if  $p(T)T^\dagger = T^\dagger p(T)$ . An operator  $T \in \mathcal{L}(\mathcal{H})$  is polynomially EP, if  $T$  is  $q$ -EP for some  $q \in Poly_1$ . It is proved in [12, Lemma 2.4] that  $T$  is  $p$ -EP if and only if  $T^*$  is  $\bar{p}$ -EP where  $(p(T))^* = \bar{p}(T^*)$ . It has been shown in [12] that every polynomially normal operator with a closed range is a polynomially EP operator and under additional assumptions. It has been verified that the converse is valid.

Motivated by recent research and useful properties of EP, hypo-EP,  $n$ -EP,  $n$ -hypo-EP, polynomially normal operators and polynomially EP operators, we define a new class of closed range operators called polynomially hypo-EP

operators. The class of polynomially hypo-EP operators is a wider class than the classes of all hypo-EP,  $n$ -hypo-EP and polynomially EP operators. We prove many properties and characterizations of polynomially hypo-EP operators and we give some additional properties of polynomially EP operators. We show that every polynomially EP operator is a polynomially hypo-EP operator and under additional assumptions, we verify that the converse is valid. What is more, we characterize polynomially hypo-EP operators by the adequate operator matrix representations (Theorems 3 and 4) and we obtain new characterizations of  $n$ -hypo-EP and hypo-EP operators.

## 2. POLYNOMIALLY EP OPERATORS

It's easily seen that if  $T$  is EP, then  $T$  is  $p$ -EP for an arbitrary  $p \in Poly_1$ . So, we proposed a wider class of operators which includes EP and  $n$ -EP operators. Obviously, when  $T$  is  $p$ -EP for  $p(t) = t$  (or  $p(t) = t^n$ ),  $t \in \mathbb{C}$ , then  $T$  is EP ( $n$ -EP). From [12, Exemple 2.2], we observe that a polynomially EP operator is not EP in general. In the following proposition, we give a necessary condition of  $p \in Poly_1$  such that an operator  $m$ -EP is  $p$ -EP.

**Proposition 1.** *Let  $T \in (m\text{-EP})$  where  $m \geq 1$ . Then  $T \in (p\text{-EP})$  for all  $p(t) = \sum_{j=0}^n a_j t^j \in Poly_1$  such that  $m \leq k = \min\{j \geq 1 : a_j \neq 0\}$ .*

*Proof.* Suppose that  $T$  is  $m$ -EP and  $p(t) = \sum_{j=0}^n a_j t^j$ , where  $a_0, a_1, \dots, a_n \in \mathbb{C}$ .

We have

$$p(T) = a_n T^n + a_{n-1} T^{n-1} + \dots + a_k T^k = T^m (a_n T^{n-m} + \dots + a_k T^{k-m}).$$

Thus

$$R(p(T)) = R(T^m (a_n T^{n-m} + \dots + a_k T^{k-m})) \subseteq R(T^m) \subseteq R(T^*).$$

And

$$R(p(T)^*) = R(T^{*m} (a_n T^{*n-m} + \dots + a_k T^{*k-m})) \subseteq R(T^{*m}) \subseteq R(T).$$

From [12, Theorem 2.5],  $T$  is  $p$ -EP.  $\square$

In the special case that  $p(t) = t^n$  in Proposition 1, we obtain one well-known property of  $n$ -EP operators (see for example [21, Theorem 3.3]).

**Corollary 1.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and let  $n \geq 1$  be an integer. If  $T$  is  $n$ -EP, then  $T$  is  $(n+k)$ -EP for an arbitrary integer  $k \geq 0$ .*

It well known that  $T$  is EP operator if and only if  $TT^\dagger T = T^\dagger T$  and  $T^\dagger T^2 T^\dagger = TT^\dagger$ , or equivalently  $T$  and  $T^\dagger$  are both hypo-EP operators. For class ( $n$ -EP) we have  $T \in (n\text{-EP})$  if and only if  $TT^\dagger (T^\dagger T^n)^* = (T^\dagger T^n)^*$  and  $T^\dagger T^{n+1} T^\dagger = T^n T^\dagger$ . For class of polynomially EP operators we have the following general setting.

**Theorem 1.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in Poly_1$  such that  $p(0) = 0$ . The following assertions are equivalent:*

- (1)  $T$  is  $p$ -EP;
- (2)  $TT^\dagger(T^\dagger p(T))^* = (T^\dagger p(T))^*$  and  $T^\dagger T p(T) T^\dagger = p(T) T^\dagger$ ;
- (3)

$$p(T)TT^\dagger + \left(p(T)TT^\dagger\right)^* = p(T) + \bar{p}(T^*) \text{ and } T^\dagger p(T)T + \left(T^\dagger p(T)T\right)^* = p(T) + \bar{p}(T^*).$$

*Proof.* (1) $\implies$ (2) By assumption and from [12, Theorem 2.5] we have,

$$\begin{aligned} R((T^\dagger p(T))^*) &\subset R(T) \text{ and } R(p(T)T^\dagger) \subset R(T^*) \\ &\implies P_{R(T)}(T^\dagger p(T))^* = (T^\dagger p(T))^* \text{ and } P_{R(T^*)}p(T)T^\dagger = p(T)T^\dagger \\ &\implies TT^\dagger(T^\dagger p(T))^* = (T^\dagger p(T))^* \text{ and } T^\dagger T p(T)T^\dagger = p(T)T^\dagger. \end{aligned}$$

(2) $\implies$ (1) We have

$$TT^\dagger(T^\dagger p(T))^* = (T^\dagger p(T))^* \text{ and } T^\dagger T p(T)T^\dagger = p(T)T^\dagger.$$

Multiplying right by  $T^*$  and by  $T$  respectively, we get

$$P_{R(T)}\bar{p}(T^*) = TT^\dagger\bar{p}(T^*) = \bar{p}(T^*) \text{ and } P_{R(T^*)}p(T) = T^\dagger T p(T) = p(T).$$

Thus,  $R(p(T)) \subset R(T^*)$  and  $R(\bar{p}(T^*)) \subset R(T)$ . Therefore by [12, Theorem 2.5],  $T$  is  $p$ -EP.

(1) $\implies$ (3) Is derived from [12, Theorem 2.5(iv)].

(3) $\implies$ (1) Multiplying  $p(T)TT^\dagger + (p(T)TT^\dagger)^* = p(T) + \bar{p}(T^*)$  by  $TT^\dagger$  from the right side, we get

$$p(T)TT^\dagger + TT^\dagger\bar{p}(T^*)TT^\dagger = p(T)TT^\dagger + \bar{p}(T^*)TT^\dagger,$$

which gives  $TT^\dagger\bar{p}(T^*) = \bar{p}(T^*)TT^\dagger = \bar{p}(T^*)$ . Thus,  $p(T) = p(T)TT^\dagger$ . In the same way, we show that  $T^\dagger p(T)T + (T^\dagger p(T)T)^* = p(T) + \bar{p}(T^*)$  implies  $p(T) = T^\dagger T p(T)$ . Therefore by [12, Theorem 2.5(iv)],  $T$  is  $p$ -EP.  $\square$

**Proposition 2.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$  such that  $p(0) = 0$  and  $R(p(T))$  is closed.*

- (1) *If  $p(T) \in (EP)$  then  $T \in (p\text{-EP})$ .*
- (2) *If  $T \in (SD)$  then  $T \in (p\text{-EP})$  if and only if  $p(T) \in (EP)$ .*

*Proof.* (1) If  $p(T) \in (EP)$ , then  $R(p(T)) = R(p(T)^*) \subseteq R(T^*)$  and  $R(p(T)^*) \subset R(T)$ , using [20, Theorem 2] and [12, Theorem 2.5] we get  $T \in (p\text{-EP})$ .

(2) Suppose that  $T \in (p\text{-EP}) \cap (SD)$  and let  $A = p(T)$ ,  $B = T^\dagger$ . We have  $A^*AN(B^*) = p(T)^*p(T)N(T^{\dagger*}) = p(T)^*p(T)N(T) = 0 \subseteq N(B^*)$ . On the other hand, for  $x \in N(A) = N(p(T))$  and since  $T$  is SD, we have  $ABB^*x = p(T)T^\dagger T^{\dagger*}x = T^\dagger p(T)T^{\dagger*}x = T^\dagger T^{\dagger*}p(T)x = 0$ . Thus  $BB^*N(A) \subseteq N(A)$ . So, by [20, Remark 1], we derive that  $(p(T)T^\dagger)^\dagger = T^{\dagger\dagger}p(T)^\dagger = Tp(T)^\dagger$ . Similarly, for  $A = T^\dagger$  and  $B = p(T)$ , we have  $A^*AN(B^*) \subseteq N(B^*)$  and  $BB^*N(A) \subseteq N(A)$ . Thus,  $(T^\dagger p(T))^\dagger = (p(T))^\dagger T$ . Since by assumption  $T$  is  $p$ -EP, we get  $Tp(T)^\dagger = p(T)^\dagger T$  and hence  $p(T)p(T)^\dagger = p(T)^\dagger p(T)$  or equivalently  $p(T)$  is EP.  $\square$

In the special case where  $p(t) = t^n$  in Proposition 2, we obtain one well-known property of  $n$ -EP operators (see [20, Theorem 3.6]).

### 3. POLYNOMIALLY HYPO-EP OPERATORS

At the start of this section, we introduce and study the concept of polynomially hypo-EP operators, which extends the notion of  $n$ -hypo-EP operators.

**Definition 1.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$  be nontrivial. If  $T^\dagger T p(T) = p(T) T^\dagger T$ , then we say that  $T$  is  $p$ -hypo-EP (or  $p$ -HEP, for short). An operator  $S \in \mathcal{L}(\mathcal{H})$  is polynomially hypo-EP, if  $S$  is  $q$ -hypo-EP for some  $q \in \text{Poly}_1$ .*

It is clear that both  $T$  and  $T^*$  are in hypo-EP if and only if  $T$  is EP. Also, for  $n$ -EP operators,  $T$  and  $T^*$  are simultaneously in ( $n$ -HEP) if and only if  $T$  is in ( $n$ -EP). For  $p$ -EP operators, by [12, Theorem 2.5] and Theorem 2 below, we have for  $p \in \text{Poly}_1$  such that  $p(0) = 0$ ,  $T$  is in ( $p$ -EP) if and only if  $T$  is in ( $p$ -HEP) and  $T^*$  is in ( $\bar{p}$ -HEP).

From [13, Theorem 2.4],  $T \in (\text{HEP})$  if and only if  $T = T^\dagger T^2$ , or equivalently,  $R(T) \subset R(T^*)$  ([17]). Obviously, when  $T$  is  $p$ -HEP for  $p(t) = t$  ( or  $p(t) = t^n$ ), then  $T$  is hypo-EP ( $n$ -hypo-EP).

The following proposition, we prove that if  $T$  is hypo-EP, then  $T$  is  $p$ -HEP for an arbitrary  $p \in \text{Poly}_1$ .

**Proposition 3.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and let  $p \in \text{Poly}_1$ . If  $T \in (\text{HEP})$ , then  $T \in (p\text{-HEP})$ .*

*Proof.* Suppose that  $T \in (\text{HEP})$  and  $p(t) = \sum_{i=0}^n a_i t^i$ . We have

$$\begin{aligned} p(T)T^\dagger T &= (a_n T^n + \cdots + a_1 T + a_0 I)T^\dagger T \\ &= a_n T^n T^\dagger T + \cdots + a_1 T T^\dagger T + a_0 T^\dagger T \\ &= a_n T^n + \cdots + a_1 T + a_0 T^\dagger T \\ &= a_n T^\dagger T T^n + \cdots + a_1 T^\dagger T T + a_0 T^\dagger T \\ &= T^\dagger T p(T). \end{aligned}$$

Therefore,  $T$  is  $p$ -HEP. □

The converse of this implication is generally false. In the following examples, we observe that a polynomially hypo-EP operator is not hypo-EP in general.

**Example 1.** Let  $T$  be the following matrix defined by  $T = \begin{pmatrix} 1 & 0 & 2 & 3 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

on  $\mathcal{H} = \mathbb{C}^4$ . Since

$$T^\dagger = \begin{pmatrix} \frac{1}{5} & 0 & -\frac{1}{5} & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ \frac{2}{5} & 0 & -\frac{2}{5} & 0 \\ 0 & 0 & \frac{1}{3} & 0 \end{pmatrix}, \quad T^2 = \begin{pmatrix} 1 & 0 & 2 & 9 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad T^3 = \begin{pmatrix} 1 & 0 & 2 & 9 \\ 0 & 8 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

we get,  $T^3 - T^2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  and

$$(T^3 - T^2)T^\dagger T = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} = T^\dagger T(T^3 - T^2).$$

Thus,  $T$  is  $p$ -HEP where  $p(t) = t^3 - t^2$ . Since

$$TT^\dagger - T^\dagger T = \begin{pmatrix} \frac{4}{5} & 0 & -\frac{2}{5} & 0 \\ 0 & 0 & 0 & 0 \\ -\frac{2}{5} & 0 & \frac{1}{5} & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}$$

is not negative operator, we conclude that  $T$  is not hypo-EP. Therefore  $T$  is  $p$ -HEP but is not hypo-EP.

**Example 2.** Let  $p(t) = t^3 + 4it^2 - (1 + 4i)t + 2$ ,  $t \in \mathbb{C}$ , and

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$$

on  $\mathcal{H} = \mathbb{C}^2$ . Since  $T^2 = T$ , we obtain  $p(T) = T^3 + 4iT^2 - (1 + 4i)T + 2I = 2I$ . Thus,  $T$  is  $p$ -HEP. Using

$$T^\dagger = \begin{pmatrix} \frac{1}{2} & 0 \\ \frac{1}{2} & 0 \end{pmatrix}$$

notice that

$$T^\dagger T - TT^\dagger = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & \frac{1}{2} \\ 0 & 0 \end{pmatrix}$$

is not positive operator. Hence  $T$  is not hypo-EP.

The next theorem provides characterizations of polynomially hypo-EP operators.

**Theorem 2.** Let  $T \in \mathcal{R}(\mathcal{H})$ ,  $p \in \text{Poly}_1$  such that  $p(0)=0$ . The following statements are equivalent:

- (i)  $T \in (p\text{-HEP})$ ;
- (ii)  $p(T) = T^\dagger T p(T)$ ;
- (iii)  $R(p(T)) \subset R(T^*)$ ;
- (iv)  $T^\dagger T p(T) T^\dagger = p(T) T^\dagger$ ;
- (v)  $T^\dagger T p(T) + (T^\dagger T p(T))^* = p(T) + p(T)^*$ .

*Proof.* (i) $\iff$ (ii) Suppose that  $T$  is  $p$ -HEP and  $p(t) = \sum_{i=1}^n a_i t^i$ . Then,

$$p(T) T^\dagger T = (a_n T^n + \cdots + a_1 T) T^\dagger T = (a_n T^{n-1} + \cdots + a_1) T T^\dagger T = p(T).$$

(ii) $\iff$ (iii) Evident because  $p(T) = P_{R(T^*)} p(T) \iff R(p(T)) \subset R(T^*)$ .

(iii) $\implies$ (iv) By the previous equivalence, we get  $T^\dagger T p(T) = p(T)$ , thus,  $T^\dagger T p(T) T^\dagger = p(T) T^\dagger$ .

(iv) $\implies$ (iii) We have

$$T^\dagger T p(T) T^\dagger = p(T) T^\dagger.$$

Multiplying right by  $T$ , we get

$$P_{R(T^*)} p(T) = T^\dagger T p(T) = p(T).$$

Therefore,  $R(p(T)) \subset R(T^*)$ .

(ii) $\implies$ (v) is obvious.

(v) $\implies$ (ii) Multiplying  $T^\dagger T p(T) + (T^\dagger T p(T))^* = p(T) + p(T)^*$  by  $T^\dagger T$  from the right side, we obtain that

$$T^\dagger T p(T) + p(T)^* T^\dagger T T^\dagger T = p(T) T^\dagger T + (p(T))^* T^\dagger T,$$

which gives  $T^\dagger T p(T) = p(T)$ .  $\square$

In the particular case that  $p(t) = t^n$  in Theorem 2, we obtain new characterizations of  $n$ -hypo-EP operators with a closed range.

**Corollary 2.** *Let  $T \in \mathcal{R}(\mathcal{H})$ . The following statements are equivalent:*

- (i)  $T \in (n\text{-HEP})$ ;
- (ii)  $R(T^n) \subset R(T^*)$ ;
- (iii)  $T^\dagger T^{n+1} T^\dagger = T^n T^\dagger$ ;
- (iv)  $T^\dagger T^{n+1} + (T^\dagger T^{n+1})^* = T^n + (T^n)^*$ .

**Proposition 4.** *Let  $T \in \mathcal{R}(\mathcal{H})$ ,  $p \in \text{Poly}_1$  such that  $p(0)=0$ . The following statements are equivalent:*

- (i)  $T \in (p\text{-HEP})$ ;
- (ii)  $p(T)^* = p(T)^* T^\dagger T$ ;
- (iii)  $\|p(T)^* x\| \leq k \|Tx\|$  for some  $k \geq 0$  and all  $x \in \mathcal{H}$ ;
- (iv)  $\|p(T)^* x\| \leq k \|\omega(T)x\|$  for some  $k \geq 0$  and all  $x \in \mathcal{H}$ .

*Proof.* The equivalences (i)  $\iff$  (iii) and (i)  $\iff$  (iv) follow from the fact that  $R(T^\dagger) = R(T^*)$  and the results of Douglas [6].  $\square$

The class of hypo-EP operators is closed under unitary equivalence and restrictions to reducing subspaces, see [22]. For  $p$ -hypo-EP class, we have the following properties.

**Proposition 5.** *Let  $T \in \mathcal{R}(\mathcal{H})$  be unitarily equivalent to some  $S \in \mathcal{R}(\mathcal{H})$  and let  $p \in \text{Poly}_1$ .*

- (1) *If  $M$  is a reducing subspace for  $T$ , then,  $T|_M$  is  $p$ -HEP;*
- (2)  *$S$  is  $p$ -HEP if and only if  $T$  is  $p$ -HEP.*

*Proof.* (1) Evident.

- (2) Suppose that  $S = UTU^*$  for some unitary operator  $U$ . Since  $S^\dagger = UT^\dagger U$  and  $p(S) = Up(T)U^*$ , we get

$$\begin{aligned} p(S) = S^\dagger Sp(S) &\iff Up(T)U^* = UT^\dagger U^* UTU^* Up(T)U^* \\ &\iff Up(T)U^* = UT^\dagger T p(T)U^* \\ &\iff p(T) = T^\dagger T p(T). \end{aligned}$$

Thus,  $S$  is  $p$ -HEP if and only if  $T$  is  $p$ -HEP. □

In the following theorem, we show that a  $p$ -HEP operator is  $p_k$ -HEP, where  $p_k(t) = t^k(p(t) - p(0))$ , for arbitrary  $k \in \mathbb{N}$ .

**Theorem 3.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$ . If  $T$  is  $p$ -HEP, then  $T$  is  $p_k$ -HEP, where  $p_k(t) = t^k(p(t) - p(0))$ , for every  $k \in \mathbb{N}$ .*

*Proof.* Notice that for every  $k \in \mathbb{N}$ ,  $p_k(0) = 0$ . Suppose that  $T$  is  $p$ -HEP and  $p(t) = \sum_{i=0}^n a_i t^i$ , where  $a_0, a_1, \dots, a_n \in \mathbb{C}$ .

For  $k = 1$ ,  $p_1(t) = t(p(t) - p(0)) = a_n t^{n+1} + \dots + a_1 t^2$ . By assumption

$$a_n T^n + \dots + a_1 T + a_0 I = T^\dagger T (a_n T^n + \dots + a_1 T + a_0 I).$$

Multiplying by  $T$  from the right hand side, we get

$$a_n T^{n+1} + \dots + a_1 T^2 + a_0 T = T^\dagger T (a_n T^{n+1} + \dots + a_1 T^2 + a_0 T).$$

Therefore,

$$p_1(T) = T^\dagger T p_1(T).$$

So, by Theorem 2,  $T$  is  $p_1$ -HEP.

Assume that  $T$  is  $p_k$ -HEP, where

$$p_k(t) = t^k(p_k(t) - p(0)) = a_n t^{n+k} + \dots + a_1 t^{k+1}.$$

Then

$$a_n T^{n+k} + \dots + a_1 T^{k+1} = T^\dagger T (a_n T^{n+k} + \dots + a_1 T^{k+1}),$$

multiplying by  $T$  from the right hand side, we get

$$p_{k+1}(T) = T^\dagger T (a_n T^{n+k+1} + \dots + a_1 T^{k+2}) = T^\dagger T p_{k+1}(T).$$

Hence,  $T$  is  $p_{k+1}$ -HEP. Therefore,  $T$  is  $p_k$ -HEP for all  $k \in \mathbb{N}$ . □

**Corollary 3.**  *$(n\text{-HEP}) \subseteq (n+k\text{-HEP})$  for all  $k \in \mathbb{N}$ .*



The next proposition is a generalization of [20, Proposition 13] for  $n$ -hypo-EP operator.

**Proposition 6.** *Let  $T$  be in  $\mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$  such that  $p(0) = 0$ . If one of the following conditions is satisfied:*

- (1)  $[T^\dagger T, p(T) + T^\dagger] = 0$ ,
- (2)  $[T^\dagger T, p(T) + T^*] = 0$ ,

then,  $T \in (p\text{-HEP})$ .

*Proof.* Let us first note that since  $p(0) = 0$ , then,

$$p(T)T^\dagger T = TT^\dagger p(T) = p(T) \text{ and } p(T)^*TT^\dagger = T^\dagger Tp(T)^* = p(T)^*.$$

Suppose that (1) is verified and  $p(t) = \sum_{i=0}^n a_i t^i$ , where  $a_0, a_1, \dots, a_n \in \mathbb{C}$ . Then,

$$\begin{aligned} 0 &= [T^\dagger T, p(T) + T^\dagger] \\ &= T^\dagger Tp(T) + T^\dagger TT^\dagger - p(T)T^\dagger T - T^{\dagger 2}T \\ &= T^\dagger Tp(T) + T^\dagger - p(T) - T^{\dagger 2}T. \end{aligned}$$

Multiplying by  $T^\dagger$  from the right hand side, we get

$$T^\dagger Tp(T)T^\dagger = p(T)T^\dagger.$$

Thus, by Theorem 2, we conclude that  $T \in (p\text{-HEP})$ .

Suppose that (2) is satisfied, then

$$\begin{aligned} 0 &= [T^\dagger T, p(T) + T^*] \\ &= T^\dagger Tp(T) + T^\dagger TT^* - p(T)T^\dagger T - T^*T^\dagger T \\ &= T^\dagger Tp(T) + T^* - p(T) - T^*T^\dagger T. \end{aligned}$$

Multiplying right by  $T^\dagger$ ,

$$T^\dagger Tp(T)T^\dagger = p(T)T^\dagger.$$

Thus, by Theorem 2, we derive that  $T \in (p\text{-HEP})$ . □

In the following proposition, we provide sufficient conditions for a polynomially hypo-EP operator to be a polynomially EP operator.

**Proposition 7.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$  such that  $p(0) = 0$ . Suppose that  $T \in (p\text{-HEP})$ . If  $T$  satisfies one of the following statements,*

- (1)  $[T, p(T)T^\dagger] = 0$ ;
- (2)  $[TT^\dagger, p(T) + T^\dagger] = 0$ ;
- (3)  $[TT^\dagger, p(T) + T^*] = 0$ ;

then  $T \in (p\text{-EP})$ .

*Proof.* (1) We have

$$\begin{aligned}
[T, p(T)T^\dagger] = 0 &\iff Tp(T)T^\dagger - p(T)T^\dagger T = 0 \\
&\iff p(T)TT^\dagger = p(T) \\
&\iff TT^\dagger(p(T))^* = (p(T))^* \\
&\iff R(p(T))^* \subset R(T) \\
&\iff T^* \in (p\text{-HEP}).
\end{aligned}$$

Since by assumption  $T \in (p\text{-HEP})$ , we derive that  $T \in (p\text{-EP})$ .

(2) We have

$$\begin{aligned}
0 &= [TT^\dagger, p(T) + T^\dagger] \\
&= TT^\dagger p(T) + TT^\dagger T^\dagger - p(T)TT^\dagger - T^\dagger TT^\dagger \\
&= p(T) + TT^{\dagger 2} - p(T)TT^\dagger - T^\dagger
\end{aligned}$$

Multiplying by  $TT^\dagger$  from the left hand side, we get,

$$p(T) = p(T)TT^\dagger.$$

Since  $T$  is assumed to be  $p\text{-HEP}$ , the result derives from [12, Theorem 2.5].

(3) Similarly, we get this assertion.  $\square$

The next theorem is generalization of [22, Theorem 3.1] for hypo-EP operator. We use Riesz representation to give a necessary and sufficient condition for an operator to be polynomially HEP operator.

**Theorem 2.** *If  $T \in \mathcal{L}(\mathcal{H})$  and  $p \in \text{Poly}_1$  is such that  $p(0) = 0$ , then  $T$  is  $p\text{-HEP}$  if and only if for each  $x \in \mathcal{H}$ , there exists  $c > 0$  such that*

$$(1) \quad |\langle p(T)x, y \rangle| \leq c\|Ty\|, \text{ for all } y \in \mathcal{H}.$$

*Proof.* Assume  $T$  is  $p\text{-HEP}$ . If  $x \in N(p(T))$ , then the result is trivial. Let  $x \in \mathcal{H}$  such that  $x \notin N(p(T))$ . Then  $p(T)x \in R(p(T)) \subset R(T^*)$ . Therefore there exists a non-zero  $s \in \mathcal{H}$  such that  $T^*s = p(T)x$ . Then for all  $y \in \mathcal{H}$ ,

$$|\langle p(T)x, y \rangle| = |\langle T^*s, y \rangle| = |\langle s, Ty \rangle| \leq \|s\|\|Ty\|.$$

Taking  $c = \|s\|$ , we get  $|\langle p(T)x, y \rangle| \leq c\|Ty\|$ .

Conversely, assume that for each  $x \in \mathcal{H}$ , there exists  $c > 0$  such that

$$|\langle p(T)x, y \rangle| \leq c\|Ty\|.$$

for all  $y \in \mathcal{H}$ . Let  $x \in \mathcal{H}$  be fixed.

Then, for all  $y \in \mathcal{H}$ , we have

$$c\|Ty\| \geq |\langle x, p(T)^*y \rangle| = |f_x(p(T)^*y)|$$

where we set  $f_x(p(T)^*y) = \langle p(T)^*y, x \rangle$ . Hence

$$|(p(T)f_x^*)y| \leq c\|(T^*)^*y\|,$$

for some  $c > 0$ , for all  $y \in \mathcal{H}$ . By Douglas's theorem,

$$p(T)f_x^* = T^*D,$$

for some  $D \in \mathcal{L}(\mathbb{C}, \mathcal{H})$ . Taking adjoint on both sides gives

$$f_x p(T)^* = g_x T,$$

where  $g_x = D^* \in \mathcal{L}(\mathcal{H}, \mathbb{C})$ . By Riesz representation theorem, there exists  $x' \in \mathcal{H}$  such that

$$g_x(Ts) = \langle Ts, x' \rangle$$

for all  $s \in \mathcal{H}$ . Hence for  $s \in \mathcal{H}$ ,

$$f_x p(T)^* s = g_x Ts$$

implies that

$$\langle p(T)^* s, x \rangle = \langle Ts, x' \rangle.$$

Hence for each  $x \in \mathcal{H}$ , there exists  $x' \in \mathcal{H}$  such that  $p(T)x = T^*x'$ . Thus,  $R(p(T)) \subset R(T^*)$ , i.e,  $T$  is  $p$ -HEP.  $\square$

Applying Theorem 2, for  $p(t) = t^n$  we get the following consequence which involve characterization of  $n$ -hypo-EP operators.

**Corollary 4.** *Let  $T \in \mathcal{R}(\mathcal{H})$ . Then  $T$  is  $n$ -hypo-EP if and only if for each  $x \in \mathcal{H}$ , there exists  $c > 0$  such that*

$$(2) \quad |\langle T^n x, y \rangle| \leq c \|Ty\|, \text{ for all } y \in \mathcal{H}.$$

Recall that the closed range operator  $T \in \mathcal{L}(\mathcal{H})$  has the following matrix representation

$$(3) \quad T = \begin{bmatrix} T_1 & 0 \\ T_2 & 0 \end{bmatrix} : \begin{bmatrix} R(T^*) \\ N(T) \end{bmatrix} \rightarrow \begin{bmatrix} R(T^*) \\ N(T) \end{bmatrix},$$

with respect to the orthogonal decomposition

$$\mathcal{H} = R(T^*) \oplus N(T)$$

and  $D = T_1^* T_1 + T_2^* T_2 \in \mathcal{L}(R(T^*))$  is an invertible operator. An easy computation shows that the Moore-Penrose inverse of  $T$  is represented by

$$(4) \quad T^\dagger = \begin{bmatrix} D^{-1} T_1^* & D^{-1} T_2^* \\ 0 & 0 \end{bmatrix}.$$

Using the representation (3), we obtain the following characterizations of  $p$ -HEP operators.

**Theorem 3.** *Let  $T \in \mathcal{L}(\mathcal{H})$  be with a closed range. Suppose that  $T$  is represented by (3),  $p(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_0$  and  $q(z) = a_n z^{n-1} + a_{n-1} z^{n-2} + \dots + a_1$ , where  $a_0, a_1, \dots, a_n \in \mathbb{C}$ . Then the following statements are equivalent:*

- (i)  $T$  is  $p$ -HEP;
- (ii)  $T_2 q(T_1) = 0$ .

In addition, if  $T$  is  $p$ -HEP then

$$p(T) = \begin{bmatrix} p(T_1) & 0 \\ 0 & a_0 I \end{bmatrix}.$$

*Proof.* Since  $T$  is assumed to be represented as in (3), using the equality (4) and the following representation

$$p(T) = \begin{bmatrix} p(T_1) & 0 \\ T_2q(T_1) & a_0I \end{bmatrix},$$

we derive that

$$\begin{aligned} T^\dagger T p(T) &= \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} p(T_1) & 0 \\ T_2q(T_1) & a_0I \end{bmatrix} \\ &= \begin{bmatrix} p(T_1) & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} p(T) T^\dagger T &= \begin{bmatrix} p(T_1) & 0 \\ T_2q(T_1) & a_0I \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} p(T_1) & 0 \\ T_2q(T_1) & 0 \end{bmatrix}. \end{aligned}$$

Hence,  $T^\dagger T p(T) = p(T) T^\dagger T$  if and only if  $T_2q(T_1) = 0$ .  $\square$

**Theorem 4.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and polynomials  $p, q$  be as in Theorem 3. The following statements are equivalent:*

- (i)  $T$  is  $p$ -HEP;
- (ii)  $T^\dagger T p(T) T^* T = p(T) T^* T$ ;
- (iii)  $T^\dagger T p(T) T^* = p(T) T^*$ .

*Proof.* (i)  $\iff$  (ii): Let the polynomial  $q$  be as in Theorem 3. Since

$$\begin{aligned} p(T) T^* T &= \begin{bmatrix} p(T_1) & 0 \\ T_2q(T_1) & a_0I \end{bmatrix} \begin{bmatrix} T_1^* & T_2^* \\ 0 & 0 \end{bmatrix} \begin{bmatrix} T_1 & 0 \\ T_2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} p(T_1)T_1^* & p(T_1)T_2^* \\ T_2q(T_1)T_1^* & T_2q(T_1)T_2^* \end{bmatrix} \begin{bmatrix} T_1 & 0 \\ T_2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} p(T_1)T_1^*T_1 + p(T_1)T_2^*T_2 & 0 \\ T_2q(T_1)T_1^*T_1 + T_2q(T_1)T_2^*T_2 & 0 \end{bmatrix} \\ &= \begin{bmatrix} p(T_1)D & 0 \\ T_2q(T_1)D & 0 \end{bmatrix}. \end{aligned}$$

And

$$T^\dagger T p(T) T^* T = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} p(T_1)D & 0 \\ T_2q(T_1)D & 0 \end{bmatrix} = \begin{bmatrix} p(T_1)D & 0 \\ 0 & 0 \end{bmatrix}.$$

We deduce that  $T^\dagger T p(T) T^* T = p(T) T^* T$  if and only if  $T_2q(T_1) = 0$ . Therefore by Theorem 3 we get equivalent.

(ii)  $\iff$  (iii): Multiplying the identities of assertions (ii) and (iii) from the right hand sides by  $T$  and  $T^\dagger$  respectively allows us to find this equivalence.  $\square$

Applying Theorem 4, we get the following consequences which involve characterizations of  $p$ -EP operators

**Corollary 5.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $p \in \text{Poly}_1$ . The following statements are equivalent:*

- (i)  $T$  is  $p$ -EP;
- (ii)  $T^\dagger T p(T) T^* T = p(T) T^* T$  and  $T T^\dagger \bar{p}(T^*) T T^* = \bar{p}(T^*) T T^*$ ;
- (iii)  $T^\dagger T p(T) T^* = p(T) T^*$  and  $T T^\dagger \bar{p}(T^*) T = \bar{p}(T^*) T$ .

Theorem 4 implies new characterizations of  $n$ -hypo-EP and hypo-EP operators.

**Corollary 6.** *Let  $T \in \mathcal{R}(\mathcal{H})$  and  $n \geq 1$ . The following statements are equivalent:*

- (i)  $T$  is  $n$ -hypo-EP;
- (ii)  $T^\dagger T^{n+1} T^* T = T^n T^* T$ ;
- (iii)  $T^\dagger T^{n+1} T^* = T^n T^*$ .

*In particular, the following conditions are equivalent:*

- (i)  $T$  is hypo-EP;
- (ii)  $T^\dagger T^2 T^* T = T T^* T$ ;
- (iii)  $T^\dagger T^2 T^* = T T^*$ .

## REFERENCES

- [1] Bourdon, Paul, et al, *Closed-range posinormal operators and their products* Linear Algebra and its Applications, **671** (2023), 38-58.
- [2] S. L. Campbell and C. D. Meyer, *EP Operators and Generalized Inverses*, Canadian Mathematical Bulletin, **18** (1975), 327-333.
- [3] S. Cheng, Y.Tian, *Two sets of new characterizations for normal and EP matrices*. Linear Algebr. Appl.**375**(2003), 181–195.
- [4] M.D. Cvetković, D. Mosić, *Drazin invertibility, characterizations and structure of polynomially normal operators.*, Linear Multilinear Algebr. (2021).
- [5] D.S. Djordjević, M. Chō, D. Mosić, *Polynomially normal operators*, Ann. Funct. Anal. **11** (2020), 493–504.
- [6] R.G. Douglas, *On majorization, factorization and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc. **17.2** (1966), 413-415.
- [7] D. Drivaliaris, S. Karanasios, D. Pappas, *Factorizations of EP operators*, Linear Algebr. Appl. **429(7)** (2008), 1555–1567.
- [8] H. Ezzahraoui, M. Mbekhta and E. H. Zerouali, *On the Cauchy dual of closed range operators*, Acta Sci.Math, **85** (2019), 231–248.
- [9] D.E. Ferreyra, F.E. Levis, Saroj B. Malik, A.N. Priori, *On star-dagger matrices and the core-EP decomposition*, Journal of Computational and Applied Mathematics, **438** (2024), 0377-0427.
- [10] R. Hartwig and K. Spindelböck, *Matrices for which  $A^*$  and  $A^\dagger$  commute*, Linear and Multilinear Algebra, **14** (1983), 241-256.
- [11] S. Menkad, and E. Anissa . *Some Results of  $n$ -EP Operators on Hilbert Spaces*. International Journal of Analysis and Applications, **22** (2024), 78-78.
- [12] D.Mosić, and Miloš D. Cvetković, *Polynomially EP operators*, Aequationes mathematicae **96.5** (2022): 1075-1087.
- [13] M. Itoh. *On some EP operators*, Nihonkai Mathematical Journal, **16** (2005), 49-56.

- [14] A. A. S. Jibril, *On  $n$ -Power Normal Operators*, The Journal for Science and Engineering, **33** (2008), 247–253.
- [15] Johnson, P. Sam and Vinoth, A., *Product and factorization of hypo-EP operators* Special Matrices, vol. 6, no. 1, 2018, pp. 376-382.
- [16] M. H. Pearl, *On generalized inverses of matrices*, Mathematical Proceedings of the Cambridge Philosophical Society, **62** (1966), 673-677.
- [17] A. B. Patel and M. P. Shekhawat, *Hypo-EP Operators*, Indian Journal of Pure and Applied Mathematics, **47** (2016), 73-84.
- [18] Saroj B. Malik, Laura Rueda and Néstor Thome *The class of  $m$ -EP and  $m$ -normal matrices*. Linear and Multilinear Algebra, **64.11** (2016) 2119-2132.
- [19] H. Schwerdtfeger, *Introduction to Linear Algebra and the Theory of Matrices*, P. Noordhoff N.V., Groningen-Holland, 1961.
- [20] R. Semmami, H. Ezzahraoui and E. H. Zerouali, *on some properties of  $n$ -EP and  $n$ -hypo-EP operators*, arXiv, arXiv:2410.20920.
- [21] Wang Xiunan and Chunyuan Deng, *Properties of  $m$ -EP operators*. Linear and Multilinear Algebra, **65.7** (2017) 1349-1361.
- [22] A. Vinoth and P. Sam Johnson, *On Sum and Restriction of Hypo-EP Operators*, Functional Analysis, Approximation and Computation **9.1** (2017), 37–41.

R. SEMMAMI, LABORATORY OF MATHEMATICAL ANALYSIS AND APPLICATIONS, FACULTY OF SCIENCES, MOHAMMED V UNIVERSITY IN RABAT, RABAT, MOROCCO.

*Email address:* `semmami.rachi@gmail.com`; `rachid_semmami@um5.ac.ma`

H. EZZAHRAOUI, LABORATORY OF MATHEMATICAL ANALYSIS AND APPLICATIONS, FACULTY OF SCIENCES, MOHAMMED V UNIVERSITY IN RABAT, RABAT, MOROCCO.

*Email address:* `hamid.ezzahraoui@fsr.um5.ac.ma`; `h.ezzahraoui@um5r.ac.ma`