KOŁODZIEJ-TOSATTI'S CONJECTURE ON COMPACT HERMITIAN MANIFOLD WITH BOUNDED MASS PROPERTY

LEI ZHANG

ABSTRACT. In this note, we show a conjecture of Kołodziej-Tosatti about Morse-type integrals in nef (1, 1) classes on compact Hermitian manifold with bounded mass property. As a consequence, we give positive answers to Demailly-Păun's conjecture and Tosatti-Weinkove's conjecture when compact Hermitian manifold with bounded mass property.

1. INTRODUCTION

Let (X^n, ω) be a compact Hermitian manifold and α a closed real (1, 1) form on X. A classes $[\alpha] \in H^{1,1}_{BC}(X; \mathbb{R})$ is called nef (numerical effective) if for every $\varepsilon > 0$ there exists $u_{\varepsilon} \in C^{\infty}(X, \mathbb{R})$ such that

$$\alpha + \sqrt{-1}\partial\bar{\partial}u_{\varepsilon} \ge -\varepsilon\omega.$$

A classes $[\alpha] \in H^{1,1}_{BC}(X;\mathbb{R})$ is called big if for every $\varepsilon > 0$ there exists a quasi-psh function u on X (locally the sum of psh plus smooth) such that

$$T := \alpha + \sqrt{-1} \partial \bar{\partial} u \ge \varepsilon \omega$$

holds in the current sense. In this case, Demailly-Păun [14, Theorem 2.12] show that X must be bimeromorphic to a compact Kähler manifold.

Our main interest is in a version of conjecture of Demailly [11] for nef classes on non-Kähler manifolds, which also encompasses a question proposed by Tosatti in [23, Remark 3.2]. More precisely, Kołodziej-Tosatti [18] proposed the following conjecture.

Conjecture 1:(Kołodziej-Tosatti, [18, Conjecture 1.2]) Let (X^n, ω) be a compact Hermitian manifold and α a closed real (1, 1) form such that $[\alpha]$ is nef. Then we have

$$\int_{X} \alpha^{n} = \inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha + \sqrt{-1}\partial\bar{\partial}u,0)} \left(\alpha + \sqrt{-1}\partial\bar{\partial}u\right)^{n}$$
(1.1)

where $X(\alpha + \sqrt{-1}\partial \bar{\partial} u, 0)$ denotes the set of all points $x \in X$ such that $(\alpha + \sqrt{-1}\partial \bar{\partial} u) \ge 0$.

In [11] Demailly shows that the inequality

$$\int_X \alpha^n \le \inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha + \sqrt{-1}\partial\bar{\partial}u,0)} \left(\alpha + \sqrt{-1}\partial\bar{\partial}u\right)^n.$$

holds (see Step 2 in the proof of Theorem 1.1) and the inequality holds if X is projective surface and $\alpha \in NS_{\mathbb{R}}(X)$ ([11, Theorem 1.18]). The Kähler version of conjecture 1 was shown by Demailly [12, Sec 3.5] whenever the orthogonality conjecture of Zariski decompositions of BDPP [3] holds and has been proved by Kołodziej-Tosatti [18, Proposition 2.2] when α is big or more generally in Fujiki class. In the non-Kähler case, the conjecture 1 holds [18, Proposition 2.1] if either the class $[\alpha]$ is semi-positive or the manifold X admits a Hermitian metric ω with

$$\partial\bar{\partial}\omega = 0 = \partial\bar{\partial}\omega^2. \tag{1.2}$$

Boucksom [2] defines the volume of $[\alpha]$ to be

$$vol([\alpha]) := \sup_{T} \int_{X} T_{ac}^{n} > 0,$$

where the supermum is over all Kähler currents in the class $[\alpha]$ and T_{ac} is the absolutely continuous part of the Lebesgue decomposition with respect to the Lebesgue measure on X. If $[\alpha]$ is not big, then one can define $vol([\alpha]) = 0$. Boucksom [2, Conjecture] conjectured that there exists a Kähler current in $[\alpha] \in H^{1,1}_{BC}$ on a compact Hermitian manifold (X, ω) provided that $\int_X \alpha^n > 0$. Boucksom [2, Theorem 4.7] has proved his conjecture when ω is Kähler. In the non-Kähler manifold, Wang [25, 26] has produced partial results on Boucksom's conjecture when $[\alpha]$ is pseudo-effective real (1,1)-class. Recently, Boucksom-Guedj-Lu [4] shown that the lower volume function vanishes outside the big cone ([4, Theorem 3.20]). As a simple consequence, they proved that X is Fujiki if X has the bounded mass property and it admits a closed positive (1,1)-current T such that $\int_X T^n > 0$ (that is, $\int_X \alpha^n > 0$, see[2, 4] for more details).

As was observed by Tosatti [23], conjecture 1 also has applications to complex Monge-Ampère equations on non-Kähler manifolds. Inspired by the work of Demailly-Păun [14], we see that the following conjecture of Demailly-Păun can be obtained by applying the Demailly-Păun's methods. Based on the work of Demailly [10] and Tosatti-Weinkove [28], it is interesting to attack Tosatti-Weinkove's conjecture.

Conjecture 2: (Demailly-Păun, [14, Conjecture 0.8]) Let (X^n, ω) be a compact Hermitian manifold and α a closed real (1, 1) form such that $[\alpha]$ is nef and $\int_X \alpha^n > 0$. Then $[\alpha]$ is big.

The Kähler version of conjecture 2 was initially proved by Demailly-Păun through Demailly's mass concentration technique [10] and Yau's solution to Calabi conjecture [30]. In the case when n = 2 and α is semi-positive, conjecture 2 was proved by the work of Buchdahl [5, 6] and Lamari [19]. For the case when n = 3, Chiose [8, Theorem 4.1] has proved conjecture 2 under the assumption that there exists a pluriclosed Hermitian metric ω ($\partial \partial \omega = 0$). Later on, Popovici [22, Theorem 1.1] made an observation (based on Xiao's approach [29, Theorem 1.4) that can help to simplify the some arguments in [8] and proved conjecture 2 under the assumption (1.2). Nguyen [21, Theorem 0.7] used Popovici's observation and proved conjecture 2 assuming that α is semi-positive and that there exists a pluriclosed Hermitian metric. Recently, Li-Wang-Zhou [20, Theorem 1.11] relaxed the semi-positivity assumption of α to the assumption that there exists a bounded quasi-psh function ρ such that $\alpha + \sqrt{-1}\partial\bar{\partial}\rho \geq 0$ in the current sense. Guedj-Lu [15, 16] made a breakthrough by introducing bound mass property for the Hermitian metric ([4]), then conjecture is valid. It is worth noting that all the aforementioned results can also be regarded as partial answers to a conjecture of Boucksom [2]. Recent advances towards a conjecture of Boucksom were recently made by Boucksom-Guedj-Lu [4, Theorem C], they proved the Boucksom's conjecture if it has the bounded mass property.

Conjecture 3:(Tosatti-Weinkove, [28]) Let (X^n, ω) be a compact Hermitian manifold and α a closed real (1, 1) form such that $[\alpha]$ is nef and $\int_X \alpha^n > 0$. Fix $x_1, \dots, x_m \in X$ and choose positive real numbers τ_i, \dots, τ_m such that

$$\sum_{i=1}^{m} \tau_i^n < \int_X \alpha^n.$$

Then there exists a α -psh function ϕ with logarithmic poles at x_1, \dots, x_m :

$$\phi(z) \le \tau_i \log_2 |z| + O(1)$$

in a coordinate (z_1, \dots, z_n) at x_i .

In his seminal paper [10], Demailly used a mass concentration technique for a degenerating family of complex Monge-Ampère equations to produce α -psh in a nef and big class $[\alpha]$ on a compact Kähler manifold which have nontrivial Lelong numbers at finitely many given points. This was later extended by Tosatti-Weinkove [28, Main Theorem] to non-Kähler manifolds and proved their conjecture for n = 2, 3 and they proved for general $n \geq 4$ under the conditions that X is Moishezon and α is rational. Significant advancements have been made more recently in this field. Tosatti [23, Theorem 1.3] proved conjecture 3 when α is nef and big. Nguyen [21, Theorem 0.6] proved conjecture 3 under the assumption that α is semi-positive and $\int_X \alpha^n > 0$. Recently, Li-Wang-Zhou [20, Theorem 1.10] relaxed the semi-positivity assumption of α to the assumption that there exists a bounded quasi-psh function ρ such that $\alpha + \sqrt{-1}\partial \bar{\partial} \rho \geq 0$ in the current sense.

Now, we recall that a compact Hermitian manifold (X, ω) has bounded mass property if it satisfies

$$\overline{vol}(\omega) := \sup\left\{\int_X (\omega + \sqrt{-1}\partial\bar{\partial}f)^n; \ f \in C^\infty(X;\mathbb{R}) \ with \ \omega + \sqrt{-1}\partial\bar{\partial}f > 0\right\} < \infty.$$

The condition $\overline{vol}(\omega) < \infty$ is independent of the chioce of ω ; it is moreover invariant under bimeromorphic change of coordinates [15, Theoerm A]. In particular, $\overline{vol}(\omega) < \infty$ if X belongs to the Fujiki class. In recent years, the bounded mass property has been intensively studied by many authors, in relation to quasi-psh envelopes (see [4, 15, 16] and the reference therein).

In this note, we show the following result.

Theorem 1.1. The conjecture 1 is true if it has the bounded mass property.

Theorem 1.2. The conjecture 2 is true if it has the bounded mass property. In particular, the conjecture 3 is true if it has the bounded mass property.

2. Proof of the main result

2.1. **Proof of Theorem 1.1.** We now present a proof of this result following the argument of Kołodziej-Tosatti [18]. The proof is divided into two steps. Let $(n, k) = \frac{n!}{k!(n-k)!}$. **Step 1.** Since $[\alpha]$ is nef, then for every $\varepsilon > 0$ there exists $u_{\varepsilon} \in C^{\infty}(X, \mathbb{R})$ such that

$$\alpha + \sqrt{-1}\partial\bar{\partial}u_{\varepsilon} \ge -\varepsilon\omega$$

So,

$$X(\alpha + \sqrt{-1}\partial\bar{\partial}u_{\varepsilon} + \varepsilon\omega, 0) = X.$$

By rescaling, we can assume that $\alpha \leq \omega$. Therefore, for any smooth function

$$u \in psh(X, \alpha + \varepsilon\omega) \subset psh(X, 2\omega)$$

we apply the bounded mass property and Stokes' theorem to have

$$\inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha+\varepsilon\omega+\sqrt{-1}\partial\bar{\partial}u,0)} \left(\alpha+\varepsilon\omega+\sqrt{-1}\partial\bar{\partial}u\right)^{n} \\
\leq \int_{X} \left(\alpha+\varepsilon\omega+\sqrt{-1}\partial\bar{\partial}u_{\varepsilon}\right)^{n} \\
= \int_{X} \left(\alpha+\sqrt{-1}\partial\bar{\partial}u_{\varepsilon}\right)^{n} + \sum_{k=1}^{n} (n,k)\varepsilon^{k} \int_{X} \left(\alpha+\sqrt{-1}\partial\bar{\partial}u_{\varepsilon}\right)^{n-k} \wedge \omega^{k}$$

$$= \int_{X} \alpha^{n} + \sum_{k=1}^{n} (n,k) \varepsilon^{k} \int_{X} \left(\alpha - 2\omega + 2\omega + \sqrt{-1} \partial \bar{\partial} u_{\varepsilon} \right)^{n-k} \wedge \omega^{k}$$

$$\leq \int_{X} \alpha^{n} + \sum_{k=1}^{n} (n,k) \varepsilon^{k} \int_{X} \left(2\omega + \sqrt{-1} \partial \bar{\partial} u_{\varepsilon} \right)^{n-k} \wedge \omega^{k}$$

$$= \int_{X} \alpha^{n} + O(\varepsilon).$$
(2.1)

On the other hand, given any $u \in C^{\infty}(X, \mathbb{R})$, we have

$$X(\alpha + \sqrt{-1}\partial\bar{\partial}u, 0) \subset X(\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}u, 0)$$

and for any $x \in X(\alpha + \sqrt{-1}\partial \bar{\partial} u, 0)$

$$\left(\alpha + \sqrt{-1}\partial\bar{\partial}u\right)^n(x) \le \left(\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}u\right)^n(x).$$

So, combining (2.1) with above, we have

$$\inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha+\sqrt{-1}\partial\bar{\partial}u,0)} (\alpha + \sqrt{-1}\partial\bar{\partial}u)^{n} \\
\leq \inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha+\varepsilon\omega+\sqrt{-1}\partial\bar{\partial}u,0)} (\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}u)^{n} \\
\leq \int_{X} \alpha^{n} + O(\varepsilon).$$
(2.2)

Letting $\varepsilon \to 0$, we have

$$\int_{X} \alpha^{n} \ge \inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha + \sqrt{-1}\partial\bar{\partial}u,0)} \left(\alpha + \sqrt{-1}\partial\bar{\partial}u\right)^{n}.$$
(2.3)

Step 2. To prove the converse inequality:

$$\int_{X} \alpha^{n} \leq \inf_{u \in C^{\infty}(X,\mathbb{R})} \int_{X(\alpha + \sqrt{-1}\partial\bar{\partial}u,0)} \left(\alpha + \sqrt{-1}\partial\bar{\partial}u\right)^{n}.$$
(2.4)

Fix $u \in C^{\infty}(X, \mathbb{R})$, let $\beta = \alpha + \sqrt{-1}\partial \bar{\partial} u$. We consider the envelope

$$h_{\varepsilon}(x) := \sup\{\varphi(x) \in psh(X, \beta + \varepsilon\omega), \varphi \le 0\}$$

= $-u + u_{\varepsilon} + \sup\{\varphi(x) \in psh(X, \alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}u_{\varepsilon}), \varphi \le u - u_{\varepsilon}\}$

which satisfies $h_{\varepsilon} \in C^{1,1}(X)$ (See [9, 24]). In particular,

$$\left(\beta + \varepsilon \omega + \sqrt{-1}\partial \bar{\partial} h_{\varepsilon}\right)^n = 0, \text{ on } \{h_{\varepsilon} < 0\}.$$

Hence, applying above inequality and Proposition 3.1 (iii) in [1], we have

$$\int_{X} \left(\beta + \varepsilon \omega + \sqrt{-1} \partial \bar{\partial} h_{\varepsilon}\right)^{n} = \int_{h_{\varepsilon}=0} \left(\beta + \varepsilon \omega\right)^{n} \\ \leq \int_{X(\beta + \varepsilon \omega, 0)} \left(\beta + \varepsilon \omega\right)^{n}.$$
(2.5)

Noting that

$$\lim_{\varepsilon \to 0} \int_{X(\beta + \varepsilon\omega, 0)} \left(\beta + \varepsilon\omega\right)^n = \int_{X(\beta, 0)} \beta^n \tag{2.6}$$

Next, we show that

$$\limsup_{\varepsilon \to 0} \int_X \left(\beta + \varepsilon \omega + \sqrt{-1} \partial \bar{\partial} h_\varepsilon \right)^n \ge \int_X \alpha^n.$$
(2.7)

Indeed, similar calculation of (2.1), we have

$$\int_X \left(\beta + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}h_\varepsilon\right)^n = \int_X \left(\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}(h_\varepsilon + u)\right)^n$$
$$= \int_X \alpha^n + O(\varepsilon).$$

Combining (2.5), (2.6) with (2.7), by letting $\varepsilon \to 0$, the inequality (2.4) is proved. From above, this completes the proof.

2.2. **Proof of Theorem 1.2.** We can now give the proof of Theorem 1.2. We will follow closely Popovici's arguments [22] (see [23, p.398-399]). The starting point for our discussion is the following lemma by Lamari in [19, Lemma 3.3](see [23, Lemma 3.3] simple proof).

Lemma 2.1 (Lamari, Lemma 3.3 in [19]). Let (X^n, ω) be a compact Hermitian manifold and α a real (1,1) form on X. Given $\varepsilon \geq 0$, there exists a current of the form $T = \alpha + \sqrt{-1}\partial \bar{\partial}u \geq \varepsilon \omega$ if and only if

$$\int_X \alpha \wedge \omega_G^{n-1} \ge \varepsilon \int_X \omega \wedge \omega_G^{n-1}$$

holds for all Gauduchon metrics ω_G on X.

Since $[\alpha]$ is nef, then for every $\varepsilon > 0$ there exists $u_{\varepsilon} \in C^{\infty}(X, \mathbb{R})$ such that

$$\alpha_{\varepsilon} := \alpha + \varepsilon \omega + \sqrt{-1} \partial \bar{\partial} u_{\varepsilon} > 0.$$

By Lamari's Lemma 2.1, it suffices to show that there exist $\varepsilon, \delta > 0$ such that for any Gauduchon metric ω_G on X we have

$$\int_{X} \alpha_{\varepsilon} \wedge \omega_{G}^{n-1} \ge \varepsilon (1+\delta) \int_{X} \omega \wedge \omega_{G}^{n-1}.$$
(2.8)

In the following, we divide the proof of (2.8) into three steps. **Step 1.** Thanks to Tosatti-Weinkove [27], we can find $\psi_{\varepsilon} \in C^{\infty}(X; \mathbb{R})$ solving

$$\begin{cases} \left(\alpha_{\varepsilon} + \sqrt{-1}\partial\bar{\partial}\psi_{\varepsilon}\right)^{n} &= e^{b_{\varepsilon}} \frac{\omega \wedge \omega_{G}^{n-1}}{\int_{X} \omega \wedge \omega_{G}^{n-1}}, \\ \widetilde{\alpha_{\varepsilon}} &= \alpha_{\varepsilon} + \sqrt{-1}\partial\bar{\partial}\psi_{\varepsilon} \end{cases}$$
(2.9)

where $b_{\varepsilon} \in \mathbb{R}$. Noting that

$$e^{b_{\varepsilon}} = \int_{X} \widetilde{\alpha_{\varepsilon}}^{n} = e^{\frac{b_{\varepsilon}^{n}}{2}} \Big(\int_{X} \Big(\frac{\widetilde{\alpha_{\varepsilon}}^{n}}{\omega_{G}^{n}} \Big)^{\frac{1}{2}} \Big(\frac{\omega \wedge \omega_{G}^{n-1}}{\omega_{G}^{n}} \Big)^{\frac{1}{2}} \omega_{G}^{n} \Big) \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-\frac{1}{2}} \\ = \frac{e^{\frac{b_{\varepsilon}}{2}}}{\sqrt{n}} \Big(\int_{X} \Big(\frac{\widetilde{\alpha_{\varepsilon}}^{n}}{\omega_{G}^{n}} \Big)^{\frac{1}{2}} \Big(tr_{\omega_{G}} \omega \Big)^{\frac{1}{2}} \omega_{G}^{n} \Big) \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-\frac{1}{2}}.$$

Hence,

$$e^{b_{\varepsilon}} = \frac{1}{n} \Big(\int_{X} \Big(\frac{\widetilde{\alpha_{\varepsilon}}^{n}}{\omega_{G}^{n}} \Big)^{\frac{1}{2}} \Big(tr_{\omega_{G}} \omega \Big)^{\frac{1}{2}} \omega_{G}^{n} \Big)^{2} \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-1} \\ \leq \frac{1}{n} \Big(\int_{X} \Big(\frac{\widetilde{\alpha_{\varepsilon}}^{n}}{\omega_{G}^{n}} tr_{\widetilde{\alpha_{\varepsilon}}} \omega \Big)^{\frac{1}{2}} \Big(tr_{\omega_{G}} \widetilde{\alpha_{\varepsilon}} \Big)^{\frac{1}{2}} \omega_{G}^{n} \Big)^{2} \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-1} \\ \leq \frac{1}{n} \Big(\int_{X} \Big(tr_{\widetilde{\alpha_{\varepsilon}}} \omega \Big) \widetilde{\alpha_{\varepsilon}}^{n} \Big) \Big(\int_{X} \Big(tr_{\omega_{G}} \widetilde{\alpha_{\varepsilon}} \Big) \omega_{G}^{n} \Big) \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-1} \\ = n \Big(\int_{X} \omega \wedge \widetilde{\alpha_{\varepsilon}}^{n-1} \Big) \Big(\int_{X} \widetilde{\alpha_{\varepsilon}} \wedge \omega_{G}^{n-1} \Big) \Big(\int_{X} \omega \wedge \omega_{G}^{n-1} \Big)^{-1}$$

that is,

$$\frac{\int_X \alpha_{\varepsilon} \wedge \omega_G^{n-1}}{\varepsilon \int_X \omega \wedge \omega_G^{n-1}} = \frac{\int_X \widetilde{\alpha_{\varepsilon}} \wedge \omega_G^{n-1}}{\varepsilon \int_X \omega \wedge \omega_G^{n-1}} \ge \frac{\int_X \widetilde{\alpha_{\varepsilon}}^n}{n\varepsilon \int_X \omega \wedge \widetilde{\alpha_{\varepsilon}}^{n-1}}$$
(2.10)

Step 2. There exist $\varepsilon, \delta' > 0$ independent of ω_G such that

$$n\varepsilon \int_X \omega \wedge \widetilde{\alpha_\varepsilon}^{n-1} \le (1-\delta') \int_X \widetilde{\alpha_\varepsilon}^n.$$
(2.11)

Indeed, similar calculation of (2.1), we have

$$\int_{X} \widetilde{\alpha_{\varepsilon}}^{n} = \int_{X} \left(\alpha + \sqrt{-1} \partial \overline{\partial} (u_{\varepsilon} + \psi_{\varepsilon}) + \varepsilon \omega \right)^{n}$$
$$= \dots = \int_{X} \alpha^{n} + O(\varepsilon). \tag{2.12}$$

On the other hand, by the bounded mass property and Stokes' formula again, we have

$$n\varepsilon \int_{X} \omega \wedge \widetilde{\alpha_{\varepsilon}}^{n-1} = n\varepsilon \int_{X} \omega \wedge \left(\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}(u_{\varepsilon} + \psi_{\varepsilon})\right)^{n-1}$$
$$= n\varepsilon \int_{X} \omega_{X} \wedge \alpha^{n-1} + n\varepsilon \sum_{k=1}^{n-2} \int_{X} \omega \wedge^{n-k-1} \wedge \left(\varepsilon\omega + \sqrt{-1}\partial\bar{\partial}(u_{\varepsilon} + \psi_{\varepsilon})\right)^{k}$$
$$= n\varepsilon \int_{X} \omega_{X} \wedge \alpha^{n-1} + O(\varepsilon^{2}).$$
(2.13)

Case 1: $\int_X \omega_X \wedge \alpha^{n-1} \leq 0$. By (2.12) and (2.13), (2.11) is obvious. Case 2: $\int_X \omega_X \wedge \alpha^{n-1} \geq 0$. If we choose $0 < \varepsilon \ll \frac{\int_X \alpha^n}{2n \int_X \omega \wedge \alpha^{n-1}}$, then by (2.12) and (2.13), (2.11) is valid. From above, the inequality (2.8) is proved.

Step 3. To prove conjecture 3. Indeed, by above, we see that if α is nef and $\int_X \alpha^n > 0$, then α is big. Hence, applying [23, Theorem 1.3], we finish the conjecture 3. For more details, we refer the reader's see [10, 28, 23] and the reference therein.

From above, this completes the proof of conjecture 2 and conjecture 3.

3. Further Discussion

As remarked above, we will follow closely Demailly-Păun's arguments [14] and Tosatti-Weinkove's arguments [28] (cf.[23]) to give an alternative proof of Theorem 1.2. For reader's convenience, we recall Demailly-Păun's concentration of mass for nef classes of positive self-intersection.

Lemma 3.1. (Demailly-Păun, [14], Boucksom, [2]) Let (X^n, ω) be a compact Hermitian manifold and let $Y \subset X$ be an analytic subset of X. Then there exist globally defined quasi-psh function ψ with analytic singularities along Y and a sequence of smooth functions $(\psi_{\varepsilon})_{\varepsilon \in (0,1]}$ on X decreasing pointwise to ψ such that

$$\omega_{\varepsilon} := \omega + \sqrt{-1}\partial\bar{\partial}\psi_{\varepsilon} \ge \frac{1}{2}\omega \tag{3.1}$$

and

$$\int_{U \cap V_{\varepsilon}} \omega_{\varepsilon}^p \wedge \omega^{n-p} \ge \delta_p(U) > 0 \tag{3.2}$$

in any neighborhood U of a regular point $x_0 \in Y$, where $V_{\varepsilon} = \{z \in X; \psi(z) < \log \varepsilon\}$ is the "tubular neighborhood" of radius ε around Y. In particular, for every there exists a closed

positive current $T \in \{\omega_{\varepsilon}^{p}\}$ (any weak limit T of ω_{ε}^{p} as $\varepsilon \to 0$) satisfies $T \ge \delta'[Y]$ for some $\delta' > 0$, that is, $\int_V T \ge \delta'$.

Proposition 3.2. (=Conjecture 2) Let (X^n, ω) be a compact Hermitian manifold with bounded mass property and α a closed real (1,1) form such that $[\alpha]$ is nef and $\int_X \alpha^n > 0$. Then $[\alpha]$ is big. In particular, the conjecture 3 is true.

Proof. The proof of Proposition 3.2 is divided into three steps.

Step 1. In the same setup as above, consider the following complex Monge-Ampère equations

$$\left(\alpha + \varepsilon\omega + \sqrt{-1}\partial\bar{\partial}(u_{\varepsilon} + \phi_{\varepsilon})\right)^n = C_{\varepsilon}\omega_{\varepsilon}^n \tag{3.3}$$

with

$$\widetilde{\alpha_{\varepsilon}} := \alpha + \varepsilon \omega + \sqrt{-1} \partial \bar{\partial} u_{\varepsilon} + \sqrt{-1} \partial \bar{\partial} \phi_{\varepsilon} > 0, \ \sup_{X} (u_{\varepsilon} + \phi_{\varepsilon}) = 0,$$

where for every $\varepsilon > 0$, and some uniquely determined positive constants C_{ε} and ω_{ε} is a family metrics construct by Demailly-Păun [14]. In the Kähler case, we have

$$C_{\varepsilon} = \frac{\int_X \widetilde{\alpha_{\varepsilon}}^n}{\int_X \omega_{\varepsilon}^n} = \frac{\int_X (\alpha + \varepsilon \omega)^n}{\int_X \omega^n} \ge C_0 = \frac{\int_X \alpha^n}{\int_X \omega^n} > 0.$$

In the non-Kähler manifolds, by the bounded mass property again,

$$C_{\varepsilon} = \frac{\int_X \widetilde{\alpha_{\varepsilon}}^n}{\int_X \omega_{\varepsilon}^n} = \frac{\int_X \alpha^n + O(\varepsilon)}{\int_X \omega^n + O(\varepsilon)} = C_0 > 0.$$

Step 2. Let us denote by

$$\lambda_1(z) \leq \cdots \leq \lambda_n(z)$$

the eigenvalues of $\widetilde{\alpha_{\varepsilon}}(z)$ with respect to $\omega_{\varepsilon}(z)$, at every point $z \in X$ (these functions are continuous with respect to z and of course depend also on ε). The equation (3.3) is equivalent to the fact that

$$\lambda_1(z)\cdots\lambda_n(z)=C_\varepsilon$$

is constant (which is bounded away from 0).

Fix a regular point $x_0 \in Y$ and a small neighborhood U (meeting only the irreducible component of x_0 in Y). By Lemma 3.1, we have a uniform lower bound

$$\int_{U \cap V_{\varepsilon}} \omega_{\varepsilon}^p \wedge \omega^{n-p} \ge \delta_p(U) > 0.$$
(3.4)

Now, by looking at the p smallest (resp. (n-p) largest) eigenvalues λ_i of $\widetilde{\alpha_{\varepsilon}}(z)$ with respect to $\omega_{\varepsilon}(z)$,

$$\widetilde{\alpha_{\varepsilon}}^p \ge \lambda_1 \cdots \lambda_p \omega_{\varepsilon}^p, \tag{3.5}$$

$$\widetilde{\alpha_{\varepsilon}}^{n-p} \wedge \omega_{\varepsilon}^{p} \ge \frac{1}{n!} \lambda_{p+1} \cdots \lambda_{n} \omega_{\varepsilon}^{n}.$$
(3.6)

The inequality (3.6) and bound mass property implies

$$\int_X \lambda_{p+1} \cdots \lambda_n \omega_{\varepsilon}^n \le n! \int_X \widetilde{\alpha_{\varepsilon}}^{n-p} \wedge \omega_{\varepsilon}^p$$
$$\le n! \int_X \alpha^{n-p} \wedge \omega^p + O(\varepsilon) \le M$$

for some constant M when $\varepsilon \ll 1$. In particular, for every $\delta > 0$ the subset $E_{\delta} \subset X$ of points $z \in X$ such that . .

$$\lambda_{p+1}(z)\cdots\lambda_n(z) > \frac{M}{\delta}.$$

Hence, combining above inequality with (3.1)

$$\int_{E_{\delta}} \omega_{\varepsilon}^{p} \wedge \omega^{n-p} \le 2^{n-p} \int_{E_{\delta}} \omega_{\varepsilon}^{n} \le 2^{n-p} \delta.$$
(3.7)

The combination of (3.4) and (3.7) yields

$$\int_{(U \cap V_{\varepsilon}) \setminus E_{\delta}} \omega_{\varepsilon}^{p} \wedge \omega^{n-p} \ge \delta_{p}(U) - 2^{n-p} \delta$$

On the other hand, by (3.5), we have

$$\widetilde{\alpha_{\varepsilon}}^p \geq \frac{C_{\varepsilon}}{\lambda_{p+1}\cdots\lambda_n} \omega_{\varepsilon}^p \geq \frac{C_{\varepsilon}}{M/\delta} \omega_{\varepsilon}^p, \text{ on } (U \cap V_{\varepsilon}) \setminus E_{\delta}.$$

From this we infer

$$\int_{U \cap V_{\varepsilon}} \widetilde{\alpha_{\varepsilon}}^{p} \wedge \omega^{n-p} \geq \frac{C_{\varepsilon}}{M/\delta} \int_{(U \cap V_{\varepsilon}) \setminus E_{\delta}} \omega_{\varepsilon}^{p} \wedge \omega^{n-p} \\ \geq \frac{C_{\varepsilon}}{M/\delta} (\delta_{p}(U) - 2^{n-p}\delta) > 0.$$
(3.8)

provided that δ is taken small enough, e.g., $\delta = 2^{-(n-p+1)}\delta_p(U)$. **Step 3.** The family of (p, p)-forms is uniformly bounded in mass since

$$\int_X \widetilde{\alpha_\varepsilon}^p \wedge \omega^{n-p} = \int_X \alpha^p \wedge \omega^{n-p} + O(\varepsilon) \le M$$

by the bound mass property. Let T be any weak limit of α_{ε}^{p} . By (3.8), T carries nonzero mass on every p-codimensional component of Y (near regular point). By Skoda's extension theorem, $1_Y T$ is a closed positive current with support in Y, hence $1_Y T = \sum c_j [Y_j]$ is a combination of the various components Y_j of Y with coefficients $c_j > 0$. This shows that T belongs to the cohomology class $\{\alpha\}^p$. Argue as the proof of Theorem 4.7 in [2] (cf. [2, Lemma 4.10]), we can finish the proof of Proposition 3.2.

Acknowledgments: The author would like to thank Professor Zhiwei Wang and Zhuo Liu for useful discussions and insightful suggestions, which helped greatly to improve the exposition of the present article. The author is greatly indebted to Professor Zhenlei Zhang for his stimulating conversations and encouragements towards this problem.

References

- Robert J Berman. Bergman kernels and equilibrium measures for line bundles over projective manifolds. Amer. J. Math. 131 (2009), no. 5, 1485-1524.
- [2] Sèbastien Boucksom. On the volume of a line bundle. Internat. J. Math. 13 (2002), no. 10, 1043-1063.
- [3] Sèbastien Boucksom, Jean-Pierre Demailly, Mihai Păun, Thomas Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. J. Algebraic Geom. 22 (2013), no. 2, 201-248.
- [4] Sèbastien Boucksom, Vincent Guedj, Chinh H. Lu Volumes of Bott-Chern classes. priprint, arXiv:2406.01090.
- [5] Nicholas Buchdahl. On compact Kähler surfaces. Ann. Inst. Fourier (Grenoble) 49 (1999), no. 1, vii, xi, 287-302.
- [6] Nicholas Buchdahl. A Nakai-Moishezon criterion for non-Kähler surfaces. Ann. Inst. Fourier (Grenoble) 50 (2000), no. 5, 1533-1538.
- [7] Ionuţ Chiose. Obstructions to the existence of Kähler structures on compact complex manifolds. Proc. Amer. Math. Soc. 142 (2014), no. 10, 3561-3568.
- [8] Ionuţ Chiose. The Kähler rank of compact complex manifolds. J. Geom. Anal. 26 (2016), no. 1, 603-615. bibitemChu21 Jianchun Chu. C^{1,1} regularity of degenerate complex Monge-Ampère equations and some applications. Anal. PDE 14 (2021), no. 6, 1671-1700.

- [9] Jianchun Chu, Bin Zhou. Optimal regularity of plurisubharmonic envelopes on compact Hermitian manifolds. Sci. China Math. 62 (2019), no. 2, 371-380.
- [10] Jean-Pierre Demailly. A numerical criterion for very ample line bundles. J. Differential Geom. 37 (1993), no. 2, 323-374.
- [11] Jean-Pierre Demailly. Holomorphic Morse inequalities and asymptotic cohomology groups: a tribute to Bernhard Riemann. Milan J. Math. 78 (2010), no. 1, 265-277.
- [12] Jean-Pierre Demailly. A converse to the Andreotti-Grauert theorem. Ann. Fac. Sci. Toulouse Math. (6) 20 (2011), Fascicule Spècial, 123-135.
- [13] Jean-Pierre Demailly. Complex analytic and differential geometry. freely accessible book. https://wwwfourier.ujf-grenoble.fr/ demailly/manuscripts/agbook.pdf
- [14] Jean-Pierre Demailly, Mihai Păun. Numerical characterization of the Kähler cone of a compact Kähler manifold. Ann. of Math. (2) 159 (2004), no. 3, 1247-1274.
- [15] Vincent Guedj, Chinh H Lu. Quasi-plurisubharmonic envelopes 2: Bounds on Monge-Ampère volumes. Algebr. Geom. 9 (2022), no. 6, 688-713.
- [16] Vincent Guedj, Chinh H Lu. Quasi-plurisubharmonic envelopes 3: Solving Monge-Ampère equations on hermitian manifolds. J. Reine Angew. Math. 800 (2023), 259-298.
- [17] Slawomir Kolodziej. Hölder continuity of solutions to the complex Monge-Ampère equation with the right-hand side in L^p: the case of compact Kähler manifolds. Math. Ann. 342 (2008), no. 2,379-386.
- [18] Slawomir Kolodziej, Valentino Tosatti. Morse-type integrals on non-Kähler manifolds. Pure Appl. Math. Q. 17 (2021), no. 3, 991-1004.
- [19] Ahche Lamari. Courants kählériens et surfaces compactes. Ann. Inst. Fourier (Grenoble) 49 (1999), no. 1, vii, x, 263-285.
- [20] Yinji Li, Zhiwei Wang, Xiangyu Zhou Degenerate complex Monge-Ampère type equations on compact Hermitian manifolds and applications. Trans. Amer. Math. Soc. 377 (2024), no. 8, 5947-5992.
- [21] Ngoc Cuong Nguyen. The complex Monge-Ampère type equation on compact Hermitian manifolds and applications. Adv. Math. 286 (2016), 240-285.
- [22] Dan Popovici. Sufficient bigness criterion for differences of two nef classes. Math. Ann. 364 (2016), no. 1-2, 649-655.
- [23] Valentino Tosatti. The Calabi-Yau theorem and Kähler currents. Adv. Theor. Math. Phys. 20 (2016), no. 2, 381-404.
- [24] Valentino Tosatti. Regularity of envelopes in Kähler classes. Math. Res. Lett. 25 (2018), no. 1, 281-289.
- [25] Zhiwei Wang. On the volume of a pseudo-effective class and semi-positive properties of the Harder-Narasimhan filtration on a compact Hermitian manifold. Ann. Polon. Math. 117 (2016), no. 1, 41-58.
- [26] Zhiwei Wang. On Grauert-Riemenschneider type criteria. Acta Math. Sin. (Engl. Ser.) 35 (2019), no. 5, 609-618.
- [27] Valentino Tosatti, Ben Weinkove. The complex Monge-Ampère equation on compact Hermitian manifolds. J. Amer. Math. Soc. 23 (2010), no. 4, 1187-1195.
- [28] Valentino Tosatti, Ben Weinkove. Plurisubharmonic functions and nef classes on complex manifolds. Proc. Amer. Math. Soc. 140 (2012), no. 11, 4003-4010.
- [29] Jian Xiao. Weak transcendental holomorphic Morse inequalities on compact Kähler manifolds. Ann. Inst. Fourier (Grenoble) 65 (2015), no. 3, 1367-1379.
- [30] ShingTung Yau. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I. Comm. Pure Appl. Math. 31 (1978), 339-411.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, 100084, P.R.CHINA

Email address: leizhang92@mail.tsinghua.edu.cn