

MINIMUM DILATATIONS OF PSEUDO-ANOSOV BRAIDS

CHI CHEUK TSANG AND XIANGZHUO ZENG

ABSTRACT. We determine the minimum dilatation δ_n among pseudo-Anosov braids with n strands, for large enough values of n . This confirms a conjecture of Venzke that $\lim_{n \rightarrow \infty} \delta_n^n = (2 + \sqrt{3})^2 \approx 13.928$. Together with previous work, this also solves the minimum dilatation problem on the n -punctured sphere, for all but 6 values of n .

1. INTRODUCTION

Let S be an orientable finite-type surface. An orientation-preserving homeomorphism $f : S \rightarrow S$ is a **pseudo-Anosov map** if there exists a pair of transverse measured singular foliations ℓ^s and ℓ^u on S such that f contracts the leaves of ℓ^s and expands the leaves of ℓ^u by a factor of $\lambda(f) > 1$. The number $\lambda(f)$ is called the **dilatation** of f , and gives a measure of the dynamical complexity of f .

A theorem of Ivanov [Iva88] states that, with the surface S fixed, there are only finitely many conjugacy classes of pseudo-Anosov maps with dilatation bounded above by a given number. This allows one to ask the **minimum dilatation problem**: What is the minimum value of the dilatation among pseudo-Anosov maps on S ? In this paper, we address this question in the case when S has genus zero.

Let D be a disc with n marked points. A **braid** (with n strands) is a union of n embedded arcs in $D \times [0, 1]$ that meet $D \times \{0, 1\}$ in the marked points and which project to monotone paths in $[0, 1]$. A braid determines a homeomorphism on D by sweeping from $D \times \{0\}$ to $D \times \{1\}$ and letting the braid dictate the movement of the marked points. By puncturing out the n marked points and forgetting about the boundary component, we obtain a homeomorphism on the sphere with $n + 1$ punctures.

Conversely, given a homeomorphism f on a punctured sphere, one can deduce from the Lefschetz fixed point theorem that f must have a fixed puncture or a fixed interior point. The latter case can be reduced to the former case by puncturing out a fixed point. We can then obtain a braid by completing a fixed puncture into a boundary component and suspending f .

These constructions determine a correspondence between braids and homeomorphisms of punctured spheres. We refer to [Bir74] for more details. We say that a braid is **pseudo-Anosov** if its corresponding homeomorphism is a pseudo-Anosov map. The main theorem of this paper is the following.

Theorem 1.1. *Let f be a pseudo-Anosov braid with n strands. Then the dilatation of f satisfies*

$$\lambda(f) \geq \min\{14.5^{\frac{1}{n}}, \begin{cases} |x^{2k+1} - 2x^{k+1} - 2x^k + 1| & \text{if } n = 2k + 1 \\ |x^{4k} - 2x^{2k+1} - 2x^{2k-1} + 1| & \text{if } n = 4k \\ |x^{4k+2} - 2x^{2k+3} - 2x^{2k-1} + 1| & \text{if } n = 4k + 2 \end{cases}\}.$$

Here we denote by $|P(x)|$ the largest positive root of a polynomial $P(x)$.

For the rest of this paper, we denote the second item in the minimum in [Theorem 1.1](#) by $\underline{\delta}_n$. We also denote

$$\begin{aligned} \underline{N} &= \{n \mid \underline{\delta}_n^n \leq 14.5\} \\ &= \{n \mid n = 2k + 1 \geq 9\} \cup \{n \mid n = 4k \geq 16\} \cup \{n \mid n = 4k + 2 \geq 30\}. \end{aligned}$$

This is the set of n for which the right hand side in [Theorem 1.1](#) equals $\underline{\delta}_n$.

In [\[HK06\]](#), Hironaka and Kin constructed a sequence of braids $\sigma_{k-1,k+1}$, where each $\sigma_{k-1,k+1}$ has $n = 2k + 1$ strands and dilatation $\underline{\delta}_n$. In [\[Ven08\]](#), Venzke extended their work by constructing sequences of braids ψ_n with n strands and dilatation $\underline{\delta}_n$ for $n = 4k$ and $n = 4k + 2$ as well. Together with [Theorem 1.1](#), this implies the following corollary.

Corollary 1.2. *For $n \in \underline{N}$, the minimum dilatation δ_n among pseudo-Anosov braids with n strands is $\underline{\delta}_n$. In particular, $\lim_{n \rightarrow \infty} \delta_n^n = (2 + \sqrt{3})^2 \approx 13.928$.*

[Corollary 1.2](#) answers [\[HK06, Question 4.2\]](#) in the negative, confirms [\[Ven08, Conjectures 5.3 and 5.4\]](#) for all $n \in \underline{N}$ and confirms [\[Ven08, Conjecture 5.10\]](#). On the other hand, note that [\[Ven08, Conjectures 5.4\]](#) is not true for $n = 10$. See [Section 13.3](#).

Under the correspondence between braids and homeomorphisms of punctured spheres, [Theorem 1.1](#) and the examples of Hironaka-Kin and Venzke also solve the minimum dilatation problem on n -punctured spheres for $n \in \underline{N}$.

Corollary 1.3. *For $n \in \underline{N}$, the minimum dilatation $\lambda_{0,n}$ among pseudo-Anosov maps on the n -punctured sphere is $\underline{\delta}_n$. In particular, $\lim_{n \rightarrow \infty} \lambda_{0,n}^n = (2 + \sqrt{3})^2 \approx 13.928$.*

We explain the proof of [Corollary 1.3](#) in more detail in [Section 2.2](#).

A pseudo-Anosov map on a closed orientable surface is **hyperelliptic** if it commutes with a hyperelliptic involution. Via taking the quotient under the involution, hyperelliptic pseudo-Anosov maps on the closed genus g surface are in correspondence with pseudo-Anosov maps on the $2g + 2$ punctured sphere. (See [\[FM12, Chapter 9.4\]](#) for more details.) Consequently, we also have the following corollary.

Corollary 1.4. *For $g \in \{g \mid g = 2k + 1 \geq 7\} \cup \{g \mid g = 2k \geq 14\}$, the minimum dilatation η_g among hyperelliptic pseudo-Anosov maps on the closed orientable genus g surface is $\underline{\delta}_{2g+2}$. In particular, $\lim_{g \rightarrow \infty} \eta_g^g = 2 + \sqrt{3} \approx 3.732$.*

For the rest of this introduction, we review some previous work, discuss the ideas used in proving [Theorem 1.1](#), and state an addendum to [Theorem 1.1](#) which gives information about the braids that attain the minimum dilatation.

1.1. Previous work. Recall that we denote by δ_n the minimum dilatation among pseudo-Anosov braids with n strands. It is a classical fact that $\delta_3 = \underline{\delta}_3 = \frac{3+\sqrt{5}}{2}$.

Ko, Los, and Song showed in [\[SKL02\]](#) that $\delta_4 = \underline{\delta}_4$. Ham and Song showed in [\[HS07\]](#) that $\delta_5 = \underline{\delta}_5$. These results use the tool of train track automata. A train track map is a combinatorial way of encoding the dynamics of a pseudo-Anosov map. We explain this in more detail in [Section 1.2](#). The crucial fact here is that every pseudo-Anosov map is encoded by a closed path in some train track automaton, with maps of smaller dilatation giving shorter paths.

In both papers, the authors first construct the suitable train track automata. Then by searching through all paths in the automata that have length bounded by some a priori bound, they recover all the pseudo-Anosov maps with small dilatation and in particular are able to locate the minimum dilatation.

Lanneau and Thiffeault showed in [\[LT11\]](#) that $\delta_6 = \underline{\delta}_5$, $\delta_7 = \underline{\delta}_7$, and $\delta_8 = \underline{\delta}_8$. They do this by considering the orientable double cover of a pseudo-Anosov braid, for which the dilatation can be read off from the action on the first homology. The characteristic polynomial of this action must be reciprocal, and the Lefschetz fixed point theorem imposes additional conditions on its coefficients. Using these restrictions, they were able to list out the possible characteristic polynomials and locate the minimum dilatation.

Note that these previous results concern the values of δ_n for small n , while [Corollary 1.2](#) gives the values of δ_n for large n . We compile all these known values of δ_n in [Table 1](#). The shaded cells are results from previous work. The unshaded cells that contain values are [Corollary 1.2](#). The remaining cells are labelled ‘???’.

A similar table can be compiled for the minimum dilatation $\lambda_{0,n}$ among pseudo-Anosov maps on the n -punctured sphere: Reasoning as in the proof of [Corollary 1.3](#), one can deduce from the aforementioned results of [\[SKL02\]](#), [\[HS07\]](#), [\[LT11\]](#) that $\lambda_{0,4} = \underline{\delta}_3$, $\lambda_{0,5} = \lambda_{0,6} = \underline{\delta}_5$, $\lambda_{0,7} = \underline{\delta}_7$, and $\lambda_{0,8} = \underline{\delta}_8$. With [Corollary 1.3](#), the remaining unknown values of $\lambda_{0,n}$ are for $n = 10, 12, 14, 18, 22, 26$.

See [Section 13.3](#) and [Section 13.5](#) for more discussion.

1.2. Train tracks, curve complexes, and clique polynomials. A train track τ on a finite-type surface S is an embedded graph where the half-edges incident to each vertex are tangent to some tangent line. See [Figure 4](#) top. A **train track map** is a map $g : \tau \rightarrow \tau'$ that sends vertices to vertices and smooth edge paths to smooth edge paths. See [Figure 5](#) bottom. The **transition matrix** of a train track map is the matrix

$$g_* \in \text{Hom}(\mathbb{R}^{E(\tau)}, \mathbb{R}^{E(\tau')})$$

whose (e', e) -entry is the number of times $g(e)$ passes through e' .

k	$n = 4k - 1$	$n = 4k$	$n = 4k + 1$	$n = 4k + 2$
1	$\underline{\delta}_3 \approx 2.61803$	$\underline{\delta}_4 \approx 2.29663$	$\underline{\delta}_5 \approx 1.72208$	$\underline{\delta}_5 \approx 1.72208$
2	$\underline{\delta}_7 \approx 1.46557$	$\underline{\delta}_8 \approx 1.41345$	$\underline{\delta}_9 \approx 1.34372$???
3	$\underline{\delta}_{11} \approx 1.27248$???	$\underline{\delta}_{13} \approx 1.22572$???
4	$\underline{\delta}_{15} \approx 1.19267$	$\underline{\delta}_{16} \approx 1.18129$	\vdots	???
5	\vdots	$\underline{\delta}_{20} \approx 1.14192$???
6		\vdots		???
7				$\underline{\delta}_{30} \approx 1.09309$
8				$\underline{\delta}_{34} \approx 1.08144$
\vdots				\vdots

TABLE 1. The minimum dilatation among pseudo-Anosov braids with n strands. The shaded cells are results from previous work [SKL02], [HS07], [LT11]. The entry for $n = 6$ is *not* a typo.

Bestvina and Handel [BH95] showed that every pseudo-Anosov map $f : S \rightarrow S$ can be homotoped into some train track map $g : \tau \rightarrow \tau$ on a train track τ on S . In this context, we say that g **carries** f .

The train track map g encodes the dynamics of f in the following ways:

- The spectral radius of the transition matrix g_* equals the dilatation of f .
- There is a correspondence between the periodic orbits of g and the periodic orbits of f .

In [HT24], Hironaka and the first author showed that if f has at least two singularity orbits, then one can choose $g : \tau \rightarrow \tau$ to be a **standardly embedded train track map**. In this case, by separating the edges of τ into the **infinitesimal** and **real** edges, the transition matrix g_* admits a block decomposition

$$g_* = \begin{bmatrix} P & * \\ 0 & g_*^{\text{real}} \end{bmatrix}$$

where P is a permutation matrix and g_*^{real} is a Perron-Frobenius matrix. In particular, the spectral radius of g_*^{real} equal the dilatation of f . Morally, the **real transition matrix** g_*^{real} packages the dynamics of f more efficiently than the entire transition matrix.

The fact that g_*^{real} is Perron-Frobenius also means that we can compute its spectral radius as the growth rate of a curve complex. This theory is explained in detail in [McM15]. We briefly recap the key points here.

Let Γ be the directed graph whose adjacency matrix equals g_*^{real} . The fact that g_*^{real} is Perron-Frobenius implies that Γ is strongly connected.

A **curve** of Γ is an embedded directed edge cycle. The **curve complex** of Γ is the graph G whose vertices are the curves of Γ , and where two vertices are connected by an edge if and only if the corresponding curves of Γ are disjoint. For a vertex $v \in V(G)$, we define its **weight** $w(v)$ to be the length of v as a curve.

The **right-angled Artin semi-group** associated to G is defined to be

$$A_+(G) = \langle v \in V(G) \mid [v_1, v_2] = 1 \text{ if there is an edge between } v_1 \text{ and } v_2 \rangle.$$

The function $w : V(G) \rightarrow \mathbb{R}_+$ extends to a semi-group homomorphism $w : A_+(G) \rightarrow \mathbb{R}_+$. The **growth rate** of (G, w) is defined to be

$$\lambda(G, w) = \lim_{T \rightarrow \infty} (\text{number of elements } g \in A_+(G) \text{ with } w(g) \leq T)^{\frac{1}{T}}.$$

[McM15, Theorem 1.4] implies that the spectral radius of g_*^{real} equals the growth rate of (G, w) . In turn, the growth rate of (G, w) can be computed by its clique polynomial. We defer an explanation of this part of the theory to [Section 3.1](#).

The advantage of this perspective is that one can estimate the spectral radius of g_*^{real} with partial information. For example, if we know that f has a periodic point with small period, then g will also have a periodic point with small period, which means that Γ has a short curve. In general, the more intersecting short curves there are in Γ , the larger the growth rate will be, since the curves are non-commuting generators with small weight in $A_+(G, w)$. This gives us a way of finding lower bounds for the dilatation of f .

1.3. Floral train tracks. The main innovation in this paper is the observation that, when f is a pseudo-Anosov braid, one can further arrange the train track τ to be **floral**, i.e. consisting of

- one central infinitesimal polygon, which we call the **pistil**,
- several 1-pronged infinitesimal polygons, which we call the **anthers**,
- some real edges that each connect one anther to the pistil, which we call the **filaments**, and
- some real edges each of whose endpoints lie on the same vertex of the pistil, which we call the **petals**.

See [Figure 1](#) for an example.

Once this is the case, the directed graph Γ associated to the real transition matrix g_*^{real} satisfies some nice properties. For example:

- There is a collection of **filament curves** such that each vertex of Γ that is a filament in τ passes through one and only one filament curve ([Proposition 5.4\(2\)](#)).
- Edges that enter filament curves occur in pairs ([Proposition 5.4\(3\)](#)).
- There is a collection of **petal curves** such that each vertex of Γ that is a petal in τ passes through one and only one petal curve ([Proposition 5.4\(2\)](#)).




FIGURE 1. An example of a floral train track. Here the pistil is in red, the anthers are in yellow, the filaments are in green, and the petals are in blue.

- Edges that exit petal curves occur in pairs ([Proposition 5.9](#)).

Intuitively, all these properties say that there are more edges in Γ than one would expect in a general setting. Along with the existence of certain short curves which come from periodic points that are guaranteed by the Lefschetz fixed point theorem, this gives us enough control on the curve complex in order to show [Theorem 1.1](#).

In reality, the analysis is much more tedious, since we have to divide into many cases depending on how the short curves, the filament curves, and the petal curves are positioned with respect to each other.

For example, if the short curves intersect many filament/petal curves, then this already gives a large growth rate. If the short curves intersect few filament/petal curves, then since Γ is strongly connected, they must be connected to the remaining filament/petal curves via other curves. These connective curves may be long, but by the aforementioned properties, they are plentiful, hence we get large growth rate in those cases as well.

We have chosen not to include all the details of our computations in this paper. Instead, we will only include details in one of the cases, namely the case where the values $\underline{\delta}_n$ arise, in [Section 7](#).

In the rest of the cases, which spans [Section 8](#) to [Section 12](#), we will only describe our casework and present the results of our computations. In each of these cases, we will show that $\lambda^n \geq 14.5$.

1.4. Braids that attain the minimum dilatation. After knowing the values of the minimum dilatations δ_n , a natural problem is to find the set of braids that attain the minimum dilatation. We do not answer this problem in this paper, but we make some progress towards it by showing that the braids in question must satisfy certain topological and dynamical properties.

Theorem 1.5. *Let f be a pseudo-Anosov braid with n strands, where $n \in \underline{N}$. If $\lambda(f) = \underline{\delta}_n$, then:*

- *The action of f on the strands has a single orbit, i.e. the closure of the braid is a knot.*
- *Every strand is a punctured 1-pronged singular point of f .*
- *Aside from the strands, there are exactly two other singular points of f , which are*

$$\begin{cases} (k+1)\text{- and }k\text{-pronged} & \text{if } n = 2k + 1, \\ (2k+1)\text{- and } (2k-1)\text{-pronged} & \text{if } n = 4k, \\ (2k+3)\text{- and } (2k-1)\text{-pronged} & \text{if } n = 4k + 2. \end{cases}$$
- *The action of f on the collection of half-leaves at each singularity has a single orbit.*

Theorem 1.5 will drop out of our proof: In our computations we will see that the value $\underline{\delta}_n$ is attained when the directed graph Γ is of a very specific form, where in particular the lengths and quantities of the short curves, filament curves, and petal curves mentioned in **Section 1.3** are determined. From these, one will be able to deduce the items in **Theorem 1.5** immediately.

In fact, we expect that with more work, the knowledge of the directed graph Γ will allow one to exactly recover the list of pseudo-Anosov braids that attain equality. See **Section 13.4** for more discussion on this.

Organization. In **Section 2** we recall some background material on pseudo-Anosov maps. In **Section 3** we recall the theory of clique polynomials and curve complexes. In **Section 4**, we recall the theory of standardly embedded train tracks, expositing in parallel an upgrade which plays a role in defining floral train tracks.

In **Section 5**, we develop the theory of floral train tracks. We show that every pseudo-Anosov braid (up to a slight modification) is carried by a floral train track map, and we record some properties of floral train track maps.

The proof of **Theorem 1.1** spans the next few sections. **Section 6** explains the casework for our computations while **Section 7** to **Section 12** record the computations that go into proving **Theorem 1.1**. The casual reader is suggested to just look at **Section 6** and **Section 7** to get an idea of these computations.

Finally, in **Section 13**, we discuss some future directions.

Acknowledgements. The first author thanks Eriko Hironaka for introducing him to the subject of pseudo-Anosov maps with small dilatations. We thank Ian Agol for pointing out **Corollary 1.4**. We also thank Michael Landry and Curtis McMullen for comments on an earlier version of the paper.

This project started under the CRM-ISM undergraduate summer research program, and while the first author is a CRM-ISM postdoctoral fellow based at CIRGET. We thank the center and the institute for their support. The second author was supported by a Science Undergraduate Research Award at McGill University.




FIGURE 2. Left: An unpunctured 3-pronged singular point. Right: A punctured 1-pronged singular point.

2. PSEUDO-ANOSOV MAPS

In this section, we recall some facts about pseudo-Anosov maps. We refer to [FLP12, Exposés 9-13] for more details.

2.1. General definitions. In this paper, a **finite-type surface** is an oriented closed surface with finitely many points, which we call the **punctures**, removed.

A orientation-preserving homeomorphism f on a finite-type surface S is said to be a **pseudo-Anosov map** if there exists a pair of singular measured foliations (ℓ^s, μ^s) and (ℓ^u, μ^u) such that:

- (1) Away from a finite set of **singular points**, which includes the punctures, ℓ^s and ℓ^u are locally conjugate to the foliations of \mathbb{R}^2 by vertical and horizontal lines respectively.
- (2) Near a singular point x , ℓ^s and ℓ^u are locally conjugate to either
 - the pull-back of the foliations of \mathbb{R}^2 by vertical and horizontal lines by the map $z \mapsto z^{\frac{n}{2}}$ respectively, for some $n \geq 3$, or
 - the pull-back of the foliations of $\mathbb{R}^2 \setminus \{0\}$ by vertical and horizontal lines by the map $z \mapsto z^{\frac{n}{2}}$ respectively, for some $n \geq 1$.

See [Figure 2](#) for examples of singular points.

- (3) $f_*(\ell^s, \mu^s) = (\ell^s, \lambda^{-1}\mu^s)$ and $f_*(\ell^u, \mu^u) = (\ell^u, \lambda\mu^u)$ for some $\lambda = \lambda(f) > 1$.

We refer to (ℓ^s, μ^s) and (ℓ^u, μ^u) as the **stable** and **unstable** measured foliations respectively. We refer to $\lambda(f)$ as the **dilatation** of f .

The **Poincaré-Hopf index** of a singular point x is defined to be

$$(2.1) \quad \text{ind}_{\text{PH}}(x) = \begin{cases} 1 - \frac{n}{2} & \text{if } x \text{ is } n\text{-pronged and is unpunctured,} \\ -\frac{n}{2} & \text{if } x \text{ is } n\text{-pronged and is punctured.} \end{cases}$$

Here we add the letters PH to avoid confusion with the Lefschetz fixed point index, which we define later in this section.

Theorem 2.1 (Poincaré-Hopf). *Let $f : S \rightarrow S$ be a pseudo-Anosov map. Then*

$$\chi(S) = \sum_x \text{ind}_{\text{PH}}(x)$$

where the sum is taken over all singular points.

Having certain singular points lie in S while others being punctures makes it awkward to phrase some definitions cleanly. To overcome this, we consider the **completion** of f , which we define to be the homeomorphism $\bar{f} : \bar{S} \rightarrow \bar{S}$ obtained by filling in every puncture of S .

The advantage of this definition is that each point of \bar{S} is either a point of S or a puncture of S , so we can, for example, make statements about both types of singular points at once.

On the other hand, note that \bar{f} is in general not a pseudo-Anosov map, since we might have filled in some 1-pronged punctures. Nevertheless, the stable and unstable foliations on S can be extended into singular measured foliations $(\bar{\ell}^s, \bar{\mu}^s)$ and $(\bar{\ell}^u, \bar{\mu}^u)$ on \bar{S} respectively, and we still have $f_*(\bar{\ell}^s, \bar{\mu}^s) = (\bar{\ell}^s, \lambda^{-1}\bar{\mu}^s)$ and $f_*(\bar{\ell}^u, \bar{\mu}^u) = (\bar{\ell}^u, \lambda\bar{\mu}^u)$ where λ is the dilatation of f .

A **stable/unstable half-leaf** is the image of a proper embedding of $[0, \infty)$ into a leaf of $\bar{\ell}^{s/u}$, respectively.

Proposition 2.2 ([FLP12, Proposition 9.6]). *Every stable/unstable half-leaf is dense.*

A point $x \in \bar{S}$ is **periodic** if it is fixed by \bar{f}^p for some $p \geq 1$. The smallest possible value for p is the **period** of x . The set $\{f^k(x) \mid k \in \mathbb{Z}/p\}$ is the **periodic orbit** containing x . The **period** of a periodic orbit is the period of any of its elements.

For example, since \bar{f} acts as a permutation on the finite set of singular points, each singular point is periodic. We define a **singularity orbit** to be a periodic orbit that consists of singular points.

Suppose the stable leaf containing x is the union of n half-leaves $\bar{\ell}_1^s, \dots, \bar{\ell}_n^s$ meeting at x . In this case, we say that x is **n -pronged**. This agrees with our previous terminology when x is singular; if x is nonsingular, it is 2-pronged. The smallest possible value of P such that \bar{f}^P fixes x and each of the half-leaves $\bar{\ell}_i^s$ is the **rotationless period** of x . We always have $p \mid P$ and $P \mid pn$.

We say that a periodic orbit is **n -pronged** if every element in it is n -pronged. The **rotationless period** of a periodic orbit is the rotationless period of any of its elements.

Proposition 2.3. *A leaf of $\bar{\ell}^{s/u}$ contains at most one periodic point.*

Proof. Suppose $\bar{\ell}$ is a leaf of $\bar{\ell}^s$ that contains two periodic points x_1 and x_2 of period p_1 and p_2 respectively. Take a path α within $\bar{\ell}$ between x_1 and x_2 . Then for every $n \geq 0$, $\bar{f}^{np_1p_2}$ fixes x_1 and x_2 , thus $\bar{f}^{np_1p_2}(\alpha)$ is a path between x_1 and x_2 contained in a leaf of $\bar{\ell}^s$. However, $\bar{\mu}^u(\bar{f}^{np_1p_2}(\alpha)) = \lambda^{-np_1p_2}\bar{\mu}^u(\alpha) \rightarrow 0$ as $n \rightarrow \infty$, so the $\bar{\mu}^u$ -length of $\bar{f}^{np_1p_2}(\alpha)$ converges to 0, and $x_1 = x_2$. \square

2.2. Lefschetz fixed point theorem. Let \bar{f} be a homeomorphism on a closed surface. Let x be an isolated fixed point of \bar{f} . Let ν be a small closed neighborhood of x that contains no other fixed points. Choose an orientation-preserving homeomorphism between ν and a neighborhood in \mathbb{R}^2 , so that it makes sense to talk about the difference between two points in ν as an element in \mathbb{R}^2 . The **Lefschetz fixed point index** of \bar{f} at x , which we denote by $\text{ind}_{\text{Lef}}(\bar{f}, x)$, is defined to be the degree of the map $\partial\nu \rightarrow S^1$ taking y to $\frac{f(y)-y}{\|f(y)-y\|}$.

Theorem 2.4 (Lefschetz). *Let \bar{f} be a homeomorphism on a closed surface \bar{S} where every fixed point is isolated. Then*

$$\sum_{k=0}^2 (-1)^k \text{tr}(\bar{f}_* : H_k(\bar{S}) \rightarrow H_k(\bar{S})) = \sum_x \text{ind}_{\text{Lef}}(\bar{f}, x)$$

where the sum on the right hand side is taken over all fixed points of \bar{f} .

We specialize to the setting when \bar{f} is the completion of a pseudo-Anosov map $f : S \rightarrow S$. In this case, every fixed point of \bar{f} is isolated, and the Lefschetz fixed point index can be computed from the action of \bar{f} on the stable and unstable foliations as follows:

Suppose x is a n -pronged fixed point. Then the stable leaf containing x is the union of n half-leaves $\bar{\ell}_1^s, \dots, \bar{\ell}_n^s$ meeting at x . We say that x is **unrotated** if \bar{f} preserves each $\bar{\ell}_i^s$, otherwise we say that x is **rotated**. In particular if $n = 1$ then x must be unrotated.

The Lefschetz fixed point index of \bar{f} at x is given by

$$(2.2) \quad \text{ind}_{\text{Lef}}(\bar{f}, x) = \begin{cases} 1 - n & \text{if } x \text{ is } n\text{-pronged and is unrotated,} \\ 1 & \text{if } x \text{ is } n\text{-pronged and is rotated.} \end{cases}$$

2.3. Pseudo-Anosov braids. We define a **pseudo-Anosov braid** with n strands to be a pseudo-Anosov map $f : S \rightarrow S$ where S is a sphere with $n + 1$ punctures and where f fixes one of the punctures b . Under the usual way of representing braids, one thinks of S as the interior of a disc with n punctures, with the boundary of the disc corresponding to the fixed puncture b .

Theorem 2.4 simplifies for pseudo-Anosov braids.

Proposition 2.5. *Let \bar{f} be the completion of a pseudo-Anosov braid. Then*

$$\sum_x \text{ind}_{\text{Lef}}(\bar{f}, x) = 2$$

where the sum is taken over all fixed points of \bar{f} .

Proof. Since \bar{f} is an orientation-preserving homeomorphism on the sphere S^2 , we have

$$\sum_{k=0}^2 (-1)^k \text{tr}(\bar{f}_* : H_k(\bar{S}) \rightarrow H_k(\bar{S})) = 1 - 0 + 1 = 2$$

on the left hand side of [Theorem 2.4](#). □

We illustrate one application of [Proposition 2.5](#) by showing how [Corollary 1.3](#) follows from [Theorem 1.1](#) and the examples of Hironaka-Kin and Venzke, as promised in the introduction.

Proof of [Corollary 1.3](#) assuming [Theorem 1.1](#). Let f be a pseudo-Anosov map on the n -punctured sphere S , and let \bar{f} be the completion of f .

From [Proposition 2.5](#) we know that $\sum_x \text{ind}_{\text{Lef}}(\bar{f}, x) = 2$. From [Equation \(2.2\)](#), we observe that the contribution of every fixed point is at most 1, thus there are at least two fixed points. Let b be one of them.

If b is a puncture of S , then f is a pseudo-Anosov braid with $n - 1$ strands, thus $\lambda(f) \geq \delta_{n-1}$. If b is not a puncture of S , we let f° be the map obtained from f by puncturing out b . Then f° is a pseudo-Anosov braid with n strands, thus $\lambda(f) = \lambda(f^\circ) \geq \delta_n$.

Combining these two cases, we have $\lambda(f) \geq \min\{\delta_{n-1}, \delta_n\} \geq \min\{14.5^{\frac{1}{n}}, \underline{\delta}_{n-1}, \underline{\delta}_n\}$ by [Theorem 1.1](#). A straightforward computation shows that $\underline{\delta}_{n-1} > \underline{\delta}_n$ for $n \neq 6, 10$. Thus we conclude that $\lambda(f) \geq \underline{\delta}_n$ for $n \in N$.

Meanwhile, the braids $\sigma_{k-1, k+1}$ with $n = 2k + 1$ strands in Hironaka-Kin [[HK06](#)] that have dilatation $\underline{\delta}_n$ have a k -pronged singularity at the fixed puncture. For $n \in N$, we have $k \geq 4$, thus we can fill in the puncture to get pseudo-Anosov maps on the n -punctured sphere that also have dilatation $\underline{\delta}_n$.

Similarly, the braids ψ_n with $n = 4k$ or $n = 4k + 2$ strands in Venzke [[Ven08](#)] that have dilatation $\underline{\delta}_n$ have a $2k - 1$ -pronged singularity at the fixed puncture. For $n \in N$, we have $k \geq 4$, thus we can fill in the puncture to get pseudo-Anosov maps on the n -punctured sphere that also has dilatation $\underline{\delta}_n$. □

3. CLIQUE POLYNOMIALS AND CURVE COMPLEXES

In this section, we recall the theory of clique polynomials and curve complexes as described in [[McM15](#)].

3.1. Clique polynomials. A **weighted graph** is a (simple, undirected) graph G together with a function $w : V(G) \rightarrow \mathbb{R}_+$, which we refer to as the **weight** of the vertices.

Given a weighted graph (G, w) , the **right-angled Artin semi-group** associated to G is defined to be

$$A_+(G) = \langle v \in V(G) \mid [v_1, v_2] = 1 \text{ if there is an edge between } v_1 \text{ and } v_2 \rangle.$$

The function $w : V(G) \rightarrow \mathbb{R}_+$ extends to a semi-group homomorphism $w : A_+(G) \rightarrow \mathbb{R}_+$. The **growth rate** of (G, w) is defined to be

$$\lambda(G, w) = \lim_{T \rightarrow \infty} (\text{number of elements } g \in A_+(G) \text{ with } w(g) \leq T)^{\frac{1}{T}}.$$

A **clique** of G is a subset $K \subset V(G)$ where every pair of vertices in K is connected by an edge. In particular the empty set is always a clique. The **size** of K , which we denote by $|K|$, is the number of vertices in K . The **weight** of K , which we denote by $w(K)$, is the sum of the weights of the vertices in K .

The **clique polynomial** of (G, w) is defined to be

$$Q_{(G,w)}(t) = \sum_K (-1)^{|K|} t^{w(K)}$$

where the sum is taken over all cliques.

Theorem 3.1 ([McM15, Theorem 4.2]). *The smallest positive zero of $Q_{(G,w)}(t)$ equals $\frac{1}{\lambda(G,w)}$.*

We record some properties of the growth rate $\lambda(G, w)$ as a function of the graph G and the weight function w .

Recall that H is an **induced subgraph** of G if $V(H) \subset V(G)$ and there is an edge between v_1 and v_2 in H if and only if there is an edge between v_1 and v_2 in G .

We say that H is a **wide subgraph** of G if $V(H) = V(G)$ and there is an edge between v_1 and v_2 in H only if there is an edge between v_1 and v_2 in G .

Proposition 3.2.

- (1) For any $\alpha \in \mathbb{R}_+$, we have $\lambda(G, \alpha w) = \lambda(G, w)^{\frac{1}{\alpha}}$.
- (2) If $w \leq w'$, then $\lambda(G, w) \geq \lambda(G, w')$.
- (3) If H is an induced subgraph of G , then $\lambda(G, w) \geq \lambda(H, w|_H)$.
- (4) If H is a wide subgraph of G , then $\lambda(H, w) \geq \lambda(G, w)$.
- (5) For any $w_1, w_2 : V(G) \rightarrow \mathbb{R}_+$ and $t \in [0, 1]$, we have $\lambda(G, tw_1 + (1-t)w_2) \leq \lambda(G, w_1)^t \lambda(G, w_2)^{1-t}$.




FIGURE 3. Summing the vertices v_1 and v_2 in a weighted graph (G, w) to get a weighted graph (H, z) .

Proof. (1)-(3) can be found in [McM15, P.196 ‘Basic properties and examples’ 1 and 2].

To see (4), note that there is a surjective semi-group homomorphism $A_+(H) \twoheadrightarrow A_+(G)$. The weight function $w : A_+(H) \rightarrow \mathbb{R}_+$ factors through this homomorphism, so we have an induced surjection

$$\{g \in A_+(H) \mid w(g) \leq T\} \twoheadrightarrow \{g \in A_+(G) \mid w(g) \leq T\}$$

for each T , which implies that $\lambda(H, w) \geq \lambda(G, w)$.

(5) follows from [McM15, Theorems 3.3 and 4.2]. \square

We introduce an operation on weighted graphs that will be handy in our analysis.

Definition 3.3. Let (G, w) be a weighted graph. Let v_1 and v_2 be two vertices of G . We define a weighted graph (H, z) as follows:

- The vertices of H are the same as those of G but with v_1 and v_2 identified into a single vertex v_3 .
- $z(u) = w(u)$ for any vertex $u \neq v_3$ in H .
- $z(v_3) = w(v_1) + w(v_2)$.
- The edges of H between vertices other than v_3 are the same as those of G .
- There is an edge between v_3 and another vertex u in H if and only if there are edges between v_1 and u and between v_2 and u in G .

We say that (H, z) is obtained from (G, w) by **summing** v_1 and v_2 . See Figure 3 for an example.

Proposition 3.4. *Suppose (H, z) is obtained from (G, w) by summing two vertices. Then $\lambda(H, z) \leq \lambda(G, w)$.*

Proof. We use the tool of minimal words as described in [McM15, Section 4], see also [GS00]: Given a total order on $V(G)$, a word in $A_+(G)$ is **minimal** if it does not

contain a subword of the form $c_1 \dots c_k d$ where d commutes with c_1, \dots, c_k and $d < c_1$. Each element of $A_+(G)$ is represented by a unique minimal word.

We choose a total order on $V(G)$ with $v_1 < v_2$ being the lowest ordered two elements. This induces a total order on $V(H)$ by letting v_3 be the lowest ordered element and retaining the ordering on the rest of the elements. Under these orders, given a minimal word in $A_+(H)$, we can replace every maximal string of the form v_3^k by $v_1^k v_2^k$ to obtain a minimal word in $A_+(G)$ with the same weight.

This defines an injection

$$\{g \in A_+(H) \mid z(g) \leq T\} \hookrightarrow \{g \in A_+(G) \mid w(g) \leq T\}$$

for each T , which implies that $\lambda(H, z) \leq \lambda(G, w)$. \square

More generally, we can sum together a collection of vertices. The growth rate is non-increasing under this operation.

3.2. Curve complexes. Let $A \in M_{n \times n}(\mathbb{Z}_{\geq 0})$ be a square matrix with non-negative integer entries. The **directed graph associated to** A , which we denote by Γ_A , is the graph whose vertex set is $\{1, \dots, n\}$ and with A_{ji} edges from i to j . Equivalently, Γ_A is the directed graph whose adjacency matrix is A . More generally, for $k \geq 1$, the (j, i) -entry of A^k is the number of directed edge paths from i to j .

Suppose that A is Perron-Frobenius. This implies that Γ_A is **strongly connected**, i.e. for any pair of vertices (v_1, v_2) there is a directed edge path from v_1 to v_2 .

We use the following terminology for directed graphs: A **cycle** of Γ_A is a cyclic sequence of directed edges $(e_i)_{i \in \mathbb{Z}/p}$ where the terminal point of e_i equals the initial point of e_{i+1} for every i . In particular, a cycle is allowed to have repeated vertices. The **length** of a cycle is the number of edges that it consists of. A cycle is **primitive** if it is not a multiple of another cycle. A **curve** of Γ_A is an embedded cycle, i.e. a cycle that does not have repeated vertices. Every cycle is the concatenation of a collection of curves.

The **curve complex** of Γ_A , which we denote by (G_A, w_A) , is the weighted graph whose vertices are the curves of Γ_A , the weight of a vertex being the length of the curve, and where two vertices are connected by an edge if and only if the corresponding curves of Γ_A are disjoint, i.e. do not share any vertex.

The following theorem provides the connection between Perron-Frobenius matrices and the clique polynomial.

Theorem 3.5 ([McM15, Theorem 1.4]). *The characteristic polynomial of A equals the reciprocal of the clique polynomial of (G_A, w_A) . In particular the spectral radius of A equals the growth rate of (G_A, w_A) .*




FIGURE 4. Top: A train track. Bottom: The tie neighborhood of a train track. The switch ties are bolded.

4. STANDARDLY EMBEDDED TRAIN TRACKS

In this section, we set up the machinery of standardly embedded train tracks. The main new result is [Theorem 4.12](#). Most ideas in this section can be found in [\[HT24\]](#), and we refer to that paper for some of the proofs.

4.1. Train track terminology. A **train track** τ on a finite-type surface S is an embedded graph with a choice of a tangent line at each vertex, such that the half-edges incident to each vertex are tangent to the tangent line and there is at least one half-edge tangent to either side of the tangent line. See [Figure 4 top](#) for a local picture of a train track. We refer to the vertices of a train track as its **switches**.

A **tie neighborhood** of τ is a closed subset N of S that can be written as a union of compact intervals, called the **ties**, such that the map $S \rightarrow S$ defined by quotienting each tie sends N to τ . The ties that map to the switches of τ are the **switch ties**. See [Figure 4 bottom](#).

A train track τ is said to **carry** a lamination ℓ on S if there is a tie neighborhood N of τ such that the leaves of ℓ are contained in N and are transverse to the ties. More generally, τ is said to **carry** a singular foliation ℓ if τ carries the lamination obtained by blowing air into the singular leaves of ℓ .

In this paper, all train tracks will carry the unstable foliation of some pseudo-Anosov map. The laminations obtained by blowing air into the singular leaves of such a foliation have the property that every complementary region is a disc D^2 or a half-open annulus $S^1 \times [0, 1)$ with $n \geq 1$ points on its boundary removed, where n is the number of prongs of the singular point that the leaf contained. Correspondingly, the complementary regions of our train tracks will always be discs or half-open annuli with $n \geq 1$ cusps on its boundary. More precisely, by a **cusps**, we mean a non-smooth point on the boundary of the complementary region.

We define a **boundary component** of a train track τ to be a boundary component of a complementary region. We say that a boundary component is **n -pronged** if it contains n cusps. We denote the (abstract) union of boundary components of τ by $\partial\tau$.




FIGURE 5. Top: A homeomorphism that sends the tie neighborhood N into the tie neighborhood N' . Bottom: The induced map on the train tracks $\tau \rightarrow \tau'$. For clarity, we have drawn the image of τ lying slightly off τ' .

Let τ and τ' be train tracks on a surface S . Let N and N' be tie neighborhoods of τ and τ' respectively. A **train track map** from τ to τ' is a homeomorphism $g : S \rightarrow S$ that sends N into N' , mapping the ties of N into ties of N' and mapping the switch ties of N into the switch ties of N' . See Figure 5 top.

By collapsing along the ties, we get an induced map $\tau \rightarrow \tau'$ that sends the switches of τ to switches of τ' and smooth edge paths of τ to smooth edge paths of τ' . See Figure 5 bottom. In this paper, we take the customary approach of referring to this map as the train track map instead, while implicitly remembering the data of the tie neighborhoods.

The **transition matrix** of a train track map $g : \tau \rightarrow \tau'$ is the matrix

$$g_* \in \text{Hom}(\mathbb{R}^{E(\tau)}, \mathbb{R}^{E(\tau')})$$

whose (e', e) -entry is the number of times $g(e)$ passes through e' .

A **train track isomorphism** is a train track map that is a graph isomorphism. The transition matrix of a train track isomorphism is always a permutation matrix.

4.2. (Based) standardly embedded train tracks.

Definition 4.1 (Standardly embedded train tracks). Let $\partial\tau = \partial_I\tau \sqcup \partial_O\tau$ be a partition of the boundary components of τ into a nonempty set of **inner boundary components** and a nonempty set of **outer boundary components** respectively.

A train track τ is said to be **standardly embedded** with respect to $(\partial_I\tau, \partial_O\tau)$ if its set of edges $E(\tau)$ can be partitioned into a set of **infinitesimal edges** $E_{\text{inf}}(\tau)$ and a set of **real edges** $E_{\text{real}}(\tau)$, such that:

- At each switch, the infinitesimal edges are tangent to one side of the tangent line while the real edges are tangent to the other side.




FIGURE 6. An example of a standardly embedded train track. The infinitesimal edges are in red and the real edges are in blue.

- The union of infinitesimal edges is a disjoint union of cycles, which we call the **infinitesimal polygons**.
- The infinitesimal polygons are exactly the inner boundary components of τ .

We show one example of a standardly embedded train track in [Figure 6](#).

A **standardly embedded train track map** is a train track map $g : \tau \rightarrow \tau'$ between standardly embedded train tracks that send infinitesimal edges to infinitesimal edges. By listing the infinitesimal edges in front of the real edges, the transition matrix of a standardly embedded train track map admits a block decomposition

$$g_* = \begin{bmatrix} P & * \\ 0 & g_*^{\text{real}} \end{bmatrix}$$

where P is a permutation matrix. We call g_*^{real} the **real transition matrix** of g .

The dimension of the real transition matrix is well-controlled.


Proposition 4.2 ([\[HT24, Proposition 3.2\]](#)). *Let τ be a standardly embedded train track. The number of real edges of τ is equal to $-\chi(\tau)$.*

The standardly embedded train track maps that we will deal with in this paper can be decomposed into elementary moves, which we now describe.

Construction 4.3. Let τ be a standardly embedded train track. Let c be an outer boundary component, and let s be a cusp of c . Let e_0 and e_1 be the two edges of τ adjacent to s along c . These must be real edges. Suppose e_0 is not a complete side of c , i.e. suppose there is an infinitesimal edge e'_0 adjacent to e_0 along c such that the concatenation $e_0 * e'_0$ is a smooth edge path lying along c .

We say that the map g that folds a proper subpath of e_1 onto $e_0 * e'_0$ is an **elementary fold** of e_1 onto e_0 at s . See [Figure 7](#).

Note that g is a standardly embedded train track map from τ to another standardly embedded train track τ' that has the same infinitesimal polygons and the same number

FIGURE 7. An elementary fold of e_1 onto e_0 at s .

of real edges as τ . In fact, there is a canonical identification between the edges of τ and that of τ' : We denote the image of e_0 as e_0 , the image of e'_0 as e'_0 , and the subpath of e_1 that was not folded as e_1 , and retain the label on the remaining edges. Under this labelling, the real transition matrix g_*^{real} is the matrix $I + N$ where I is the identity matrix and N has a 1 in the (e_0, e_1) -entry and 0 elsewhere.

In [HT24], Hironaka and the first author showed that the dynamics of any pseudo-Anosov map with at least two singularity orbits can be encoded by a standardly embedded train track map, in the following sense.

Theorem 4.4 ([HT24, Theorem 1.15]). *Let f be a pseudo-Anosov map. Given a partition of the singular points of S into two nonempty f -invariant subsets $\mathcal{X} = \mathcal{X}_I \sqcup \mathcal{X}_O$, there exists a standardly embedded train track τ and a standardly embedded train track map $g : \tau \rightarrow \tau$ such that:*

- (1) *g is induced by a map that sends a tie neighborhood of τ into itself and that is homotopic to f .*
- (2) *τ carries the unstable foliation of f . Each boundary component in $\partial_{I/O}\tau$ corresponds to one element of $\mathcal{X}_{I/O}$ respectively.*
- (3) *g can be factored into elementary folds followed by a train track isomorphism.*
- (4) *The real transition matrix g_*^{real} is Perron-Frobenius and reciprocal.*
- (5) *The spectral radius of g_*^{real} is the dilatation of f .*

Technically, (3) is not recorded in [HT24, Theorem 1.15], but it follows from the proof of [HT24, Lemma 3.20]. In any case, Theorem 4.4 will be subsumed by Theorem 4.12, which is a mild upgrade saying that the cusps of the outer boundary components can be chosen to lie on prescribed inner boundary components.

To work towards Theorem 4.12, we introduce some specializations of our definitions so far.

Definition 4.5 (Based standardly embedded train tracks). Let $\partial\tau = \partial_P\tau \sqcup \partial_A\tau \sqcup \partial_O\tau$ be a partition of the boundary components of τ into a nonempty set of **pistil boundary**

components, a (possibly empty) set of **another boundary components**, and a nonempty set of **outer boundary components** respectively.

A train track that is standardly embedded with respect to $(\partial_P\tau \sqcup \partial_A\tau, \partial_O\tau)$ is said to be **based at** $\partial_P\tau$ if each cusp of an outer boundary component lies on a pistil boundary component. When $\partial_P\tau$ is understood, we simply say that τ is a **based standardly embedded train track**.

The motivation for our terminology comes from the special case of floral train tracks, which we mentioned in the introduction (see [Figure 1](#)) and will define precisely in [Section 5.2](#).

Performing an elementary folding move on a based standardly embedded train track τ returns a standardly embedded train track τ' , but τ' may no longer be based at $\partial_P\tau$. Indeed, in the notation of [Construction 4.3](#), the cusp s now lies on the inner boundary component that contains e'_0 , which may or may not have been an element of $\partial_P\tau$.

To make sure we stay within the category of based standardly embedded train tracks, we define a **based fold** to be a composition of elementary folds at the same cusp s , $\tau = \tau_0 \xrightarrow{g_1} \dots \xrightarrow{g_n} \tau_n$ where the intermediate train tracks $\tau_1, \dots, \tau_{n-1}$ are not based at $\partial_P\tau$, but the final train track τ_n is based at $\partial_P\tau$.

We compose the edge identifications before and after each elementary fold to identify the edges before and after a based fold.

4.3. Train track partitions. The goal of this subsection is to retrace the constructions in [\[HT24, Section 3\]](#) in order to state and prove [Theorem 4.12](#).

Let $\bar{f} : \bar{S} \rightarrow \bar{S}$ be the completion of a pseudo-Anosov map $f : S \rightarrow S$.

A **rectangle** is the image of a map $R : [0, 1] \times [0, 1] \rightarrow \bar{S}$ where

- R is an embedding in the interior $(0, 1) \times (0, 1)$,
- the image of $\{u_0\} \times [0, 1]$ lies within a stable leaf for every u_0 , and
- the image of $[0, 1] \times \{s_0\}$ lies within an unstable leaf for every s_0 .

We refer to $R(\{0\} \times [0, 1])$ and $R(\{1\} \times [0, 1])$ as the **stable sides** and $R([0, 1] \times \{0\})$ and $R([0, 1] \times \{1\})$ as the **unstable sides** of the rectangle.

Let x be a singular point. Suppose x is n -pronged. Then the stable/unstable leaf containing x is the union of n half-leaves $\bar{\ell}_1^{s/u}, \dots, \bar{\ell}_n^{s/u}$ meeting at x . A **stable/unstable prong** at x is a compact segment P_i of some $\bar{\ell}_i^{s/u}$ that contains x . A **stable/unstable star** at x is a union of prongs $\bigcup_{i=1}^n P_i$, where each P_i is contained in $\bar{\ell}_i^{s/u}$. A **side** of a stable star is the union of two adjacent prongs.

Definition 4.6 ((Based) train track partitions). Let $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ be a partition of the set of singular points \mathcal{X} . We write $\mathcal{X}_I = \mathcal{X}_P \sqcup \mathcal{X}_A$.

A **train track partition** with respect to $(\mathcal{X}_I, \mathcal{X}_O)$ consists of

- a stable star σ_x^s at each $x \in \mathcal{X}_I$, and
- an unstable star σ_x^u at each $x \in \mathcal{X}_O$

such that:

- The complementary regions of $\bigcup_{x \in \mathcal{X}_I} \sigma_x^s \cup \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$ are rectangles.
- The interior of each stable star is disjoint from the unstable stars.
- The interior of each unstable star is disjoint from the stable stars.

A train track partition $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is **Markov** if

- $\bar{f}(\bigcup_{x \in \mathcal{X}_I} \sigma_x^s) \subset \bigcup_{x \in \mathcal{X}_I} \sigma_x^s$, and
- $\bar{f}(\bigcup_{x \in \mathcal{X}_O} \sigma_x^u) \supset \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$.

A train track partition $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is **based at \mathcal{X}_P** if each endpoint of an unstable star σ_x^u lies on a stable star σ_x^s where $x \in \mathcal{X}_P$.

Given a f -invariant partition of \mathcal{X} , we demonstrate a way of constructing based Markov train track partitions.

Construction 4.7. Let $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ be a partition of \mathcal{X} into f -invariant subsets, where \mathcal{X}_P and \mathcal{X}_O are nonempty. We write $\mathcal{X}_I = \mathcal{X}_P \sqcup \mathcal{X}_A$.

We construct a stable star σ_x^s for every $x \in \mathcal{X}_I$ and an unstable star σ_x^u for every $x \in \mathcal{X}_O$ as follows:

- (1) For each $x \in \mathcal{X}_P$, let σ_x^s be the stable star at x for which each of its prongs has μ^u -length 1.
- (2) For each $x \in \mathcal{X}_O$, let σ_x^u be the unstable star at x that is maximal with respect to the property that its interior is disjoint from $\bigcup_{x \in \mathcal{X}_P} \sigma_x^s$. That is, each σ_x^u is defined by extending the unstable prongs at x until it bumps into some σ_x^s .
- (3) For each $x \in \mathcal{X}_I$, let σ_x^s be the stable star at x that is maximal with respect to the property that its interior is disjoint from $\bigcup_{x \in \mathcal{X}_O} \sigma_x^u$. That is, each σ_x^s is defined by extending the stable prongs at x until it bumps into some σ_x^u .

Here we have implicitly applied [Proposition 2.2](#) to make sure that we can extend σ_x^u and σ_x^s until they bump into the required stars.

The same argument as in [[HT24](#), Construction 3.6] shows that $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is a train track partition with respect to $(\mathcal{X}_I, \mathcal{X}_O)$: By construction the interior of each stable star is disjoint from the unstable stars, and vice versa. The complementary regions are foliated by the restriction of the stable leaves, and their corners are all convex, so they must be rectangles.

The same argument as in [HT24, Proposition 3.7] shows that $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is Markov: One checks that $\bar{f}(\bigcup_{x \in \mathcal{X}_I} \sigma_x^s) \subset \bigcup_{x \in \mathcal{X}_I} \sigma_x^s$ and $\bar{f}(\bigcup_{x \in \mathcal{X}_O} \sigma_x^u) \supset \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$ are true at each stage of the construction (taking the yet-to-be-defined stars to be empty). At stage (1), we have $\bar{f}(\bigcup_{x \in \mathcal{X}_P} \sigma_x^s) \subset \bigcup_{x \in \mathcal{X}_P} \sigma_x^s$ since \bar{f} contracts the stable stars by a factor of $\lambda(f)$. This implies that $\bar{f}(\bigcup_{x \in \mathcal{X}_O} \sigma_x^u) \supset \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$ in stage (2). This in turn implies that $\bar{f}(\bigcup_{x \in \mathcal{X}_I} \sigma_x^s) \subset \bigcup_{x \in \mathcal{X}_I} \sigma_x^s$ in stage (3).

Finally, $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is based at \mathcal{X}_P by construction.

Construction 4.7 is in a sense the most natural way to build a based train track partition, given the argument in the following proposition.

Proposition 4.8. *A based train track partition $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is uniquely determined by its stable stars σ_x^s for $x \in \mathcal{X}_P$.*

Proof. If two based train track partitions have the same collection of stable stars σ_x^s for $x \in \mathcal{X}_P$, then their unstable stars σ_x^u for $x \in \mathcal{X}_O$ must coincide since these must be the maximal unstable stars whose interiors are disjoint from $\bigcup_{x \in \mathcal{X}_P} \sigma_x^s$. In turn, their stable stars σ_x^s for $x \in \mathcal{X}_A$ must coincide since these must be the maximal stable stars whose interiors are disjoint from $\bigcup_{x \in \mathcal{X}_O} \sigma_x^u$. \square

We also record the following construction, which turns a based Markov train track partition with respect to $(\mathcal{X}'_I, \mathcal{X}'_O)$ into a based Markov train track partition with respect to $(\mathcal{X}_I, \mathcal{X}_O)$ after we move some elements from \mathcal{X}'_I to \mathcal{X}'_O .

Construction 4.9. Let $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}'_A \sqcup \mathcal{X}'_O$ be a partition of \mathcal{X} into f -invariant subsets, where \mathcal{X}_P and \mathcal{X}'_O are nonempty. We write $\mathcal{X}'_I = \mathcal{X}_P \sqcup \mathcal{X}'_A$. Suppose $((\sigma_x^s)_{x \in \mathcal{X}'_I}, (\sigma_x^u)_{x \in \mathcal{X}'_O})$ is a based Markov train track partition.

Let $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ be another partition of \mathcal{X} into f -invariant subsets, where $\mathcal{X}_A \subset \mathcal{X}'_A$ and $\mathcal{X}_O \supset \mathcal{X}'_O$. We write $\mathcal{X}_I = \mathcal{X}_P \sqcup \mathcal{X}_A$. Note that the set of pistil singularities has not been changed.

We construct an unstable star σ_x^u for every $x \in \mathcal{X}'_A \setminus \mathcal{X}_A = \mathcal{X}_O \setminus \mathcal{X}'_O$ by extending the unstable prongs at x until it bumps into some σ_x^s .

Here again we have implicitly applied **Proposition 2.2** to make sure that we can extend σ_x^u until they bump into the stable stars.

We claim that $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is a train track partition with respect to $(\mathcal{X}_I, \mathcal{X}_O)$: By construction the interior of each stable star is disjoint from the unstable stars, and vice versa. Since no endpoint of an unstable star lay on a stable star σ_x^s for $x \in \mathcal{X}'_A$, the complementary regions of $\bigcup_{x \in \mathcal{X}_I} \sigma_x^s \cup \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$ have convex corners. Also, the regions are foliated by the restriction of the stable leaves, so they must be rectangles.

Since $\bar{f}(\bigcup_{x \in \mathcal{X}_P} \sigma_x^s) \subset \bigcup_{x \in \mathcal{X}_P} \sigma_x^s$, we have $\bar{f}(\bigcup_{x \in \mathcal{X}_O \setminus \mathcal{X}'_O} \sigma_x^u) \supset \bigcup_{x \in \mathcal{X}_O \setminus \mathcal{X}'_O} \sigma_x^u$. This implies that $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is Markov.

Finally, $((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ is based at \mathcal{X}_P by construction.

The crucial property about (based) train track partitions is that they give rise to (based) standardly embedded train tracks.

Construction 4.10 ([HT24, Construction 3.8]). Let $\mathcal{M} = ((\sigma_x^s)_{x \in \mathcal{X}_I}, (\sigma_x^u)_{x \in \mathcal{X}_O})$ be a train track partition with respect to $(\mathcal{X}_I, \mathcal{X}_O)$.

Define a graph $\tau_{\mathcal{M}}$ by taking a set of vertices in one-to-one correspondence with the sides of the stable stars σ_x^s . The edges of $\tau_{\mathcal{M}}$ will come in two types: infinitesimal and real. The **infinitesimal** edges correspond to the prongs of σ_x^s . The endpoints of the edge corresponding to a prong P lies at the two vertices corresponding to the two sides that P belongs to. The **real** edges are in one-to-one correspondence with the rectangles R in the complement of $\bigcup_{x \in \mathcal{X}_I} \sigma_x^s \cup \bigcup_{x \in \mathcal{X}_O} \sigma_x^u$. The endpoints of the edge corresponding to a rectangle R lies at the two vertices corresponding to the two sides the stable sides of R lie along.

We embed $\tau_{\mathcal{M}}$ into the surface S by placing the infinitesimal edges corresponding to the prongs of a single star σ_x^s around the singular point x in the cyclic order of the prongs, and placing the real edge along their corresponding rectangle.

See each of the two columns of [Figure 8](#) for local examples of the construction.

Observe that the cusps of the outer boundary components of $\tau_{\mathcal{M}}$ are in one-to-one correspondence with the endpoints of the unstable stars of \mathcal{M} . A cusp lies on the inner boundary component of $\tau_{\mathcal{M}}$ that corresponds to the stable star on which the corresponding endpoint lies on. In particular, if \mathcal{M} is based at \mathcal{X}_P , then $\tau_{\mathcal{M}}$ is based at $\partial_P \tau$.

Construction 4.11 ([HT24, Construction 3.9]). Continuing from [Construction 4.10](#), suppose in addition that \mathcal{M} is Markov. We define a standardly embedded train track map $g_{\mathcal{M}} : \tau_{\mathcal{M}} \rightarrow \tau_{\mathcal{M}}$ by mapping the vertex corresponding to a side s of a stable star to the vertex corresponding to the side containing $\bar{f}(s)$, mapping the infinitesimal edge corresponding to a stable prong P to the infinitesimal edge corresponding to the prong containing $\bar{f}(P)$, and mapping the real edge corresponding to a rectangle R to the edge path corresponding to the sequence of rectangles and prongs passed through by $\bar{f}(R)$.

Observe that the (e', e) -entry of the real transition matrix $(g_{\mathcal{M}})_*^{\text{real}}$ is the number of appearances of e' as an edge in the edge path $g(e)$. In other words, the edges from e to e' in the directed graph Γ associated to $(g_{\mathcal{M}})_*^{\text{real}}$ are in one-to-one correspondence with these appearances, which in turn correspond to the sub-rectangles in $\bar{f}(R) \cap R'$, where R and R' are the rectangles corresponding to e and e' respectively.

Theorem 4.12. *Let f be a pseudo-Anosov map. Let $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ be a partition of \mathcal{X} into f -invariant subsets, where \mathcal{X}_P and \mathcal{X}_O are nonempty. Suppose \mathcal{M} is a Markov train track partition with respect to $(\mathcal{X}_P \sqcup \mathcal{X}_A, \mathcal{X}_O)$ and based at \mathcal{X}_P . Then the based*




FIGURE 8. Each of the two columns: Local examples of [Construction 4.10](#). From the left column to the right column: Retracting an unstable prong and extending a stable prong translates to an elementary fold.

standardly embedded train track $\tau = \tau_{\mathcal{M}}$ and the standardly embedded train track map $g = g_{\mathcal{M}}$ satisfy:

- (1) g is induced by a map that sends a tie neighborhood of τ into itself and that is homotopic to f .
- (2) τ carries the unstable foliation of f . Each boundary component in $\partial_{P/A/O}\tau$ corresponds to one element of $\mathcal{X}_{P/A/O}$ respectively.
- (3) g can be factored into based folds followed by a train track isomorphism.
- (4) The real transition matrix g_*^{real} is Perron-Frobenius and reciprocal.
- (5) The spectral radius of g_*^{real} is the dilatation of f .

Proof. (1) and (2) are clear from construction.

To show (3), we will define a sequence of based train track partitions

$$((\sigma_x^s(k))_{x \in \mathcal{X}_I}, (\sigma_x^u(k))_{x \in \mathcal{X}_O})$$

that interpolates between \mathcal{M} and $\overline{f}^{-1}(\mathcal{M})$.

We start by defining $((\sigma_x^s(0))_{x \in \mathcal{X}_I}, (\sigma_x^u(0))_{x \in \mathcal{X}_O}) = \mathcal{M}$. Inductively, suppose

$$((\sigma_x^s(k))_{x \in \mathcal{X}_I}, (\sigma_x^u(k))_{x \in \mathcal{X}_O})$$

has been defined. If every stable prong P^s in $\sigma_x^s(k)$ at a point $x \in \mathcal{X}_P$ is a stable prong in $\bar{f}^{-1}(\mathcal{M})$, then by [Proposition 4.8](#), $\bar{f}^{-1}(\mathcal{M}) = ((\sigma_x^s(k))_{x \in \mathcal{X}_I}, (\sigma_x^u(k))_{x \in \mathcal{X}_O})$ in which case we are done with the interpolation.

Otherwise, there exists a stable prong P^s in $\sigma_x^s(k)$ at a point $x \in \mathcal{X}_P$ that is not a stable prong in $\bar{f}^{-1}(\mathcal{M})$, i.e. P^s is a strict subset of a stable prong in $\bar{f}^{-1}(\mathcal{M})$. The endpoint of P^s lies on some unstable prong P^u . We define $\sigma_x^{s/u}(k+1)$ by retracting P^u until its endpoint coincides with the endpoint of P^s . We then extend all stable prongs until their endpoints lie on the interior of some unstable star. Using our arguments as above, one can check that $((\sigma_x^s(k+1))_{x \in \mathcal{X}_I}, (\sigma_x^u(k+1))_{x \in \mathcal{X}_O})$ is a based train track partition.

We claim that the train tracks $\tau_{\mathcal{M}(k)}$ and $\tau_{\mathcal{M}(k+1)}$ are related by a sequence of based folds. Indeed, we can further interpolate between $\mathcal{M}(k)$ and $\mathcal{M}(k+1)$ by, in our construction, performing the retraction of P^u step-by-step by pausing whenever the endpoint of P^u coincides with the endpoint of *some* stable prong, and extending that stable prong until its endpoint lies on the interior of some unstable star. The intermediate train tracks are then related by elementary folds at a single cusp s , corresponding to the endpoint of P^u . See [Figure 8](#), and compare with [[HT24](#), Proposition 3.18].

We also observe that the initial and final train tracks $\tau_{\mathcal{M}(k)}$ and $\tau_{\mathcal{M}(k+1)}$ are based at \mathcal{X}_P . Thus by grouping together the elementary folds that go through non-based train tracks, we have our desired sequence of based folds.

Finally, \bar{f} induces a train track isomorphism from $\tau_{\bar{f}^{-1}(\mathcal{M})}$ to $\tau_{\mathcal{M}}$.

(4) and (5) can be shown as in [[HT24](#)]. We sketch an outline of the argument here.

To show that g_*^{real} is Perron-Frobenius, one applies [Proposition 2.2](#) to see that for every rectangle R , $\bar{f}^N(R)$ passes through every rectangle for large N . Correspondingly, for every real edge e , $(g_*^{\text{real}})^N(e)$ passes through every real edge.

To show that g_*^{real} is reciprocal, one defines the weight space $\mathcal{W}(\tau) \subset \mathbb{R}^{E(\tau)}$ ([\[HT24](#), Definition 4.1]). One defines an antisymmetric bilinear form ω on $\mathcal{W}(\tau)$, known as the **Thurston symplectic form** ([\[HT24](#), Definition 4.3]), and shows that g preserves ω ([\[HT24](#), Proposition 4.6]). One then shows that the actions of g on

- the quotient $\mathbb{R}^{E(\tau)}/\mathcal{W}(\tau)$ ([\[HT24](#), Proposition 4.2]),
- the radical $\text{rad}(\omega)$ is reciprocal ([\[HT24](#), Propositions 5.4 and 5.5]), and
- the quotient $\mathcal{W}(\tau)/\text{rad}(\omega)$ ([\[HT24](#), Proposition 2.13(1)])

are reciprocal, and deduce that the action of g on $\mathbb{R}^{E(\tau)}$, i.e. the transition matrix g_* , is reciprocal ([\[HT24](#), Proposition 2.13(3)]). Since g acts by a permutation on the infinitesimal edges, the real transition matrix g_*^{real} is also reciprocal ([\[HT24](#), Proposition 2.13(2) and (3)]).

To show that the spectral radius of g_*^{real} is the dilatation $\lambda(f)$, note that the μ^u -measures of the rectangles provide a positive $\lambda(f)$ -eigenvector, so we can conclude by the Perron-Frobenius theorem. \square

In the sequel, we will refer to a standardly embedded train track map g as in [Theorem 4.12](#) as a train track map that **carries** f with respect to the partition $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$. In view of item (2) we will often identify a boundary component of τ with its corresponding singular point.

4.4. Periodic orbits and cycles. In this subsection, we supplement [Theorem 4.12](#) by showing that if g is a train track map that carries f , then there is a correspondence between the periodic orbits of \bar{f} and the cycles in the directed graph Γ associated to the real transition matrix of g .

Let $\mathbf{a} = \{\bar{f}^k(a) \mid k \in \mathbb{Z}/p\}$ be a periodic orbit of period p . First suppose that \mathbf{a} is not a singularity orbit, i.e. $\mathbf{a} \not\subset \mathcal{X}$. Then by [Proposition 2.3](#), each $\bar{f}^k(a)$ does not lie on a stable or unstable star, hence must lie in the interior of a unique complementary rectangle R_k .

Let e_k be the real edge corresponding to R_k . Recall that the edges from e_k to e_{k+1} in Γ correspond to the sub-rectangles in $\bar{f}(R_k) \cap R_{k+1}$. Let ε_k be the edge that corresponds to the sub-rectangle containing $\bar{f}^{k+1}(x)$. We refer to the cycle $(\varepsilon_k)_{k \in \mathbb{Z}/p}$ in Γ as the cycle **determined by \mathbf{a}** .

Now suppose instead that $\mathbf{a} \subset \mathcal{X}_I$, and suppose \mathbf{a} is n -pronged. Let P be the rotationless period of \mathbf{a} . Recall that $P \mid pn$. The action of \bar{f} on the set of the pn unstable half-leaves at the p points of \mathbf{a} has $\frac{pn}{P}$ orbits, each of size P . We label these orbits as $\{\bar{f}^k(\bar{\ell}_i^u) \mid k \in \mathbb{Z}/P\}$, where $i = 1, \dots, \frac{pn}{P}$. Since each $\bar{f}^k(a)$ lies at the center of a stable star, the initial portion of each $\bar{f}^k(\bar{\ell}_i^u)$ is contained in a unique rectangle $R_{i,k}$.

Let $e_{i,k}$ be the real edge corresponding to $R_{i,k}$. Let $\varepsilon_{i,k}$ be the edge from $e_{i,k}$ to $e_{i,k+1}$ that corresponds to the sub-rectangle containing the initial portion of $\bar{f}^{k+1}(\bar{\ell}_i^u)$. We refer to the cycles $(\varepsilon_{1,k})_{k \in \mathbb{Z}/P}, \dots, (\varepsilon_{\frac{pn}{P},k})_{k \in \mathbb{Z}/P}$ in Γ as the cycles **determined by \mathbf{a}** .

Finally, the case when $\mathbf{a} \subset \mathcal{X}_O$ is similar to the case when $\mathbf{a} \subset \mathcal{X}_I$, but with the unstable half-leaves replaced by stable half-leaves: The action of \bar{f} on the set of the pn stable half-leaves at the p points of \mathbf{a} has $\frac{pn}{P}$ orbits $\{\bar{f}^k(\bar{\ell}_i^s) \mid k \in \mathbb{Z}/P\}$, where $i = 1, \dots, \frac{pn}{P}$. The initial portion of each $\bar{f}^k(\bar{\ell}_i^s)$ is contained in a unique rectangle $R_{i,k}$.

Let $e_{i,k}$ be the real edge corresponding to $R_{i,k}$, and let $\varepsilon_{i,k}$ be the edge from $e_{i,k}$ to $e_{i,k+1}$ that corresponds to the sub-rectangle containing the initial portion of $\bar{f}^{k+1}(\bar{\ell}_i^s)$. We refer to the cycles $(\varepsilon_{1,k})_{k \in \mathbb{Z}/P}, \dots, (\varepsilon_{\frac{pn}{P},k})_{k \in \mathbb{Z}/P}$ in Γ as the cycles **determined by \mathbf{a}** .

Proposition 4.13. *Let g be a train track map that carries f . Then:*

- (1) *Every cycle determined by a periodic orbit is primitive.*
- (2) *The cycles determined by distinct periodic orbits are distinct.*
- (3) *The cycles determined by a singularity orbit are distinct.*
- (4) *Every primitive cycle is determined by some periodic orbit.*

Proof. Suppose $(\varepsilon_k : e_k \rightarrow e_{k+1})_{k \in \mathbb{Z}/P}$ is a cycle determined by a periodic orbit $\mathbf{a} = \{f^k(a)\}$. Let R_k be the rectangle corresponding to e_k . Up to relabeling, we can assume that $f^k(a) \in R_k$ for all k .

For the duration of this proof, we take the intersection $\bar{f}^{-1}(R_{k+1}) \cap R_k$ to mean only the sub-rectangle corresponding to the given edge ε_k . More generally, we interpret $\bar{f}^{-k}(R_{j+k}) \cap R_j$ to mean only the sub-rectangle corresponding to the sub-path $\varepsilon_j \cdots \varepsilon_{j+k}$.

Consider the intersection $\bar{R}_j = \bigcap_{k=-\infty}^{\infty} \bar{f}^{-k}(R_{j+k})$. By definition, we have $f^j(a) \in \bar{R}_j$. The $\bar{\mu}^s$ -measure of $\bar{f}^{-kP}(R_{j+kP}) = \bar{f}^{-kP}(R_j)$ decreases to 0 as $k \rightarrow \infty$, while the $\bar{\mu}^u$ -measure of $\bar{f}^{-kP}(R_{j+kP}) = \bar{f}^{-kP}(R_j)$ decreases to 0 as $k \rightarrow -\infty$, so \bar{R}_j must only consist of the point $f^j(a)$.

If \mathbf{a} is nonsingular, we can conclude that the edge cycle (ε_k) must be primitive, since otherwise $f^j(a) = a$ for j equal to the length of the primitive cycle for which (ε_k) is a multiple of.

We can show primitivity in the case when \mathbf{a} is singular by a similar argument. Suppose $(\varepsilon_k)_{k \in \mathbb{Z}/P}$ is determined by an orbit of an unstable half-leaf $\{\bar{f}^k(\bar{\ell}^u)\}$. Up to relabeling, we can assume that the initial portion of $\bar{f}^k(\bar{\ell}^u)$ lies in R_k for all k .

Consider the intersection $\bar{R}_j = \bigcap_{k=-\infty}^0 \bar{f}^{-k}(R_{j+k})$. By definition, the initial portion of $\bar{f}^j(\bar{\ell}^u)$ lies in \bar{R}_j , and since the $\bar{\mu}^u$ -measure of $\bar{f}^{-kP}(R_{j+kP})$ decreases to 0 as $k \rightarrow -\infty$, \bar{R}_j must be equal to this initial portion.

These arguments also show that we can recover the periodic orbit \mathbf{a} and, if \mathbf{a} is singular, the orbit of the stable/unstable half-leaf, from the edge cycle (ε_k) . This implies (2) and (3).

Finally, to show (4), suppose we are given a cycle $(\varepsilon_k : e_k \rightarrow e_{k+1})_{k \in \mathbb{Z}/P}$. Let R_k be the rectangle corresponding to e_k . Then the intersection $\bigcap_{k=-\infty}^{\infty} \bar{f}^{-k}(R_k)$ is a point a in R_0 . It is straightforward to check that (ε_k) is a cycle determined by the periodic orbit of a . \square

Next, we explain how the notion of a rotated/unrotated fixed point translates under this correspondence. We will not need the full knowledge of this in this paper, but we leave in the generality in anticipation for future applications.

Suppose we have chosen orientations on e and e' . These induce orientations on the unstable leaves in R and R' . Let ε be an edge in the directed graph Γ from e to e' . Let R_ε be the subrectangle in $\bar{f}(R) \cap R'$ corresponding to ε . We say that ε is **orientation-preserving** if \bar{f} restricted to a map from $\bar{f}^{-1}(R_\varepsilon) \subset R$ to $R_\varepsilon \subset R'$ preserves the orientations on the unstable leaves. Otherwise we say that ε is **orientation-reversing**. Equivalently, ε is orientation-preserving if and only if the corresponding appearance of e' in the edge path $g(e)$ is positive.

We say that a cycle in Γ is **orientation-preserving/reversing** if it consists of an even/odd number of orientation-reversing edges, respectively. It is straightforward to deduce the following proposition from our definitions.

Proposition 4.14. *Suppose (ε_k) is a primitive cycle of length p . If (ε_k) is determined by a nonsingular periodic orbit \mathbf{a} , then (ε_k) is orientation-preserving if and only if the elements of \mathbf{a} are unrotated fixed points of f^p . If (ε_k) is determined by a singular periodic orbit, then (ε_k) is always orientation-preserving.*

With this terminology, we can state the following proposition.

Proposition 4.15. *Let g be a train track map carrying f . Let $\alpha = (\varepsilon_k : e_k \rightarrow e_{k+1})_{k \in \mathbb{Z}/P}$ be an orientation-preserving cycle determined by a periodic orbit \mathbf{a} .*

- If $\mathbf{a} \notin \mathcal{X}_O$, then there are at least two edges entering α , i.e.

$$\sum_k \sum_e ((g_*^{\text{real}})_{e_k, e} - 1) \geq 2.$$

- If $\mathbf{a} \notin \mathcal{X}_I$, then there are at least two edges exiting α , i.e.

$$\sum_k \sum_e ((g_*^{\text{real}})_{e, e_k} - 1) \geq 2.$$

Here the inner sums are taken over all vertices e of Γ .

Proof. Let R_k be the rectangle corresponding to e_k , and write $\mathbf{a} = \{f^k(a)\}$ where $f^k(a) \in R_k$. As in the proof of [Proposition 4.13](#), we will write $\bar{f}^k(R_{j+k}) \cap R_j$ to denote the subrectangle corresponding to the path $\varepsilon_j \cdots \varepsilon_{j+k}$. Also, the orientation on the real edges e_k induces orientations on the unstable leaves in R_k , thus we can coherently refer to the two unstable sides of R_k as the top and bottom unstable sides.

Suppose $\mathbf{a} \notin \mathcal{X}_O$. Then no unstable side of $\bar{f}^p(R_0) \cap R_0$ can lie on an unstable side of R_0 , since otherwise a would lie on an unstable side of R_0 . Note that the assumption that α is orientation-preserving is used here.

Let k_1 be the minimum number where the top unstable side of $\bar{f}^{k_1}(R_0) \cap R_{k_1}$ does not lie on an unstable side of R_{k_1} . Then \bar{f} must map some rectangle other than R_{k_1-1} through R_{k_1} , i.e. $\sum_e ((g_*^{\text{real}})_{e_{k_1}, e} - 1) \geq 1$.

Similarly, let k_2 be the minimum number where the bottom unstable side of $\bar{f}^{k_2}(R_0) \cap R_{k_2}$ does not lie on an unstable side of R_{k_2} . Then \bar{f} must map some rectangle other than R_{k_2-1} through R_{k_2} . Hence $\sum_e ((g_*^{\text{real}})_{e_{k_2},e} - 1) \geq 1$ and $\sum_k \sum_e ((g_*^{\text{real}})_{e_k,e} - 1) \geq 2$.

A similar argument using stable sides proves the second statement. \square

The cycles that are determined by singularity orbits will play an outsized role in the proof of [Theorem 1.1](#). We show the following additional properties regarding these cycles.

Proposition 4.16. *Each vertex of Γ can appear at most twice in the collection of cycles determined by periodic orbits in \mathcal{X}_I . The same statement holds with \mathcal{X}_I replaced by \mathcal{X}_O .*

Proof. Each time a vertex e appears in a cycle determined by $\mathbf{a} \subset \mathcal{X}_I$, the rectangle R corresponding to e contains a point in \mathbf{a} in one of its stable side. Since R has only two stable sides, and no two periodic points can lie along the same stable leaf, e cannot appear more than twice in the collection of cycles determined by periodic points in \mathcal{X}_I .

A similar argument using unstable sides proves the second statement. \square

Corollary 4.17. *Each curve Γ can appear at most once in the collection of cycles determined by periodic orbits in \mathcal{X}_I . The same statement holds with \mathcal{X}_I replaced by \mathcal{X}_O .*

Proof. If a curve γ is contained in a cycle α determined by a periodic orbit in \mathcal{X}_I , but $\gamma \neq \alpha$, then there must be a vertex on γ that appears twice in α . For example, the vertex where α enters γ is such a vertex. In this case, γ cannot appear in another cycle determined by a periodic orbit in \mathcal{X}_I .

On the other hand, the cycles determined by periodic orbits in \mathcal{X}_I are all distinct, thus in particular those cycles that are embedded are also all distinct. \square

5. FLORAL TRAIN TRACKS

In this section, we develop the theory of floral train tracks. We will show that every pseudo-Anosov braid (up to a slight modification) is carried by a floral train track map. We also establish some properties regarding the real transition matrix of a floral train track map.

5.1. Jointless train track maps. Let f be a pseudo-Anosov braid and let \bar{f} be the completion of f . By definition, f has a fixed puncture, which we denote by b . Note that b may or may not be 1-pronged. We denote the union of the 1-pronged singularities of f other than (possibly) b by \mathbf{a} .

Lemma 5.1. *Aside from b and (possibly) \mathbf{a} , \bar{f} has at least one other fixed point c .*

Proof. From [Proposition 2.5](#) we know that $\sum_x \text{ind}_{\text{Lef}}(\bar{f}, x) = 2$. From [Equation \(2.2\)](#), we observe that every unrotated fixed point contributes a nonpositive number, while every rotated fixed point contributes a 1. Thus there are at least two rotated fixed points. Every 1-pronged singularity is unrotated, so there is at least one rotated fixed point other than b . \square

Note that a priori c may be nonsingular. If this is the case, we puncture it out to make it into a 2-pronged punctured singularity. (The only reason for doing this is to stay consistent with our convention that boundary components of train tracks correspond to singularities.)

We first discuss a family of train tracks that is more general than floral train tracks.

Definition 5.2. A train track map $g : \tau \rightarrow \tau$ that carries f is **jointless** if it is based with respect to a partition $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ where $\mathbf{a} \subset \mathcal{X}_A$.

We refer to the train track on which a jointless train track map is defined on to be a **jointless** train track. Our terminology is inspired by the fact that a 1-pronged infinitesimal polygon that meets more than one real edge is called a **joint** in [\[FRW23\]](#).

Let τ be a jointless train track. We refer to a real edge of τ that has one endpoint lying on a 1-pronged infinitesimal polygon as a **filament**. Since no cusps of outer boundary components can lie on 1-pronged infinitesimal polygon, the other endpoint of a filament cannot lie on a 1-pronged infinitesimal polygon, otherwise τ just consists of the filament between two 1-pronged infinitesimal polygons, contradicting nonemptiness of \mathcal{X}_P .

We will always orient a filament from the endpoint that lies on a 1-pronged infinitesimal polygon to the endpoint that does not do so.

We refer to the rest of the real edges of τ as the **petals**. We will not place any canonical orientation on the petals.

[Theorem 4.12\(3\)](#) states that any jointless train track map g can be factorized into $g = hg_k \dots g_1$ where g_i are based folds and h is a train track isomorphism. The following lemma classifies each of the based folds.

Lemma 5.3. *Each based fold in g is of one of the following forms:*

- (1) *An elementary fold of a filament e_1 onto another filament e_0 , followed by $k \geq 0$ elementary folds of e_0 onto e_1 , followed by an elementary fold of e_1 onto e_0 . The real transition matrix of this based fold is the identity matrix except with a block of $\begin{bmatrix} k+1 & k+2 \\ k & k+1 \end{bmatrix}$ in the $\{e_0, e_1\} \times \{e_0, e_1\}$ entries. After the based fold, e_0 and e_1 remain as filaments.*
- (2) *An elementary fold of a petal e_1 onto a filament e_0 , followed by another elementary fold of e_1 onto e_0 . The real transition matrix of this based fold is the*

identity matrix except with a block of $\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ in the $\{e_0, e_1\} \times \{e_0, e_1\}$ entries. After the based fold, e_0 remains as a filament and e_1 remains as a petal.

(3) An elementary fold of a petal e_1 onto a filament e_0 , followed by an elementary fold of e_0 onto e_1 . The real transition matrix of this based fold is the identity matrix except with a block of $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix}$ in the $\{e_0, e_1\} \times \{e_0, e_1\}$ entries. After the based fold, e_0 becomes a petal while e_1 becomes a filament.

(4) An elementary fold of a filament or a petal e_1 onto a petal e_0 . The real transition matrix of this based fold is the identity matrix except with a block of $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ in the $\{e_0, e_1\} \times \{e_0, e_1\}$ entries. After the based fold, e_0 remains as a petal and e_1 remains as a filament or a petal respectively.

Proof. Consider the first elementary fold of a based fold. Using the notation of [Construction 4.3](#), e_0 and e_1 are each either a filament or a petal.

If e_0 and e_1 are filaments that meet 1-pronged infinitesimal polygons a_0 and a_1 respectively, then after the first elementary fold, e_1 now connects a_0 to a_1 . The next elementary fold must fold e_1 onto e_0 or fold e_0 onto e_1 . In the former case, the based fold is concluded. In the latter case, we return to the same setting, and we repeat the argument. Eventually, the based fold must conclude. See [Figure 9](#) top row.

If e_0 is a filament meeting a 1-pronged infinitesimal polygon a_0 while e_1 is a petal, then after the first elementary fold e_1 connects $\partial_P\tau$ to a_0 . The next elementary fold must fold e_1 onto e_0 or fold e_0 onto e_1 . In either case the based fold concludes afterwards. See [Figure 9](#) second row.

If e_0 is a petal, then after the first elementary fold the train track is already based, so the based fold just consists of one elementary fold. See [Figure 9](#) bottom two rows. \square

We associate to each based fold g_i a permutation ζ_i on the labels of the real edges:

- If g_i is of type (1) where k is odd, then ζ_i switches e_0 and e_1 .
- If g_i is of type (3) then ζ_i switches e_0 and e_1 .
- In every other case, ζ_i is the identity.

Each ζ_i satisfies the following properties:

- (1) A real edge e is a filament or a petal before the based fold g_i if and only if the real edge $\zeta_i(e)$ is a filament or a petal after the based fold respectively.
- (2) The $(\zeta_i(e), e)$ -entry of the real transition matrix $(g_i)_*^{\text{real}}$ is always positive. In fact, there is a canonical positive appearance of $\zeta_i(e)$ in the edge path $g_i(e)$: If e is a filament, then this is the appearance of $\zeta_i(e)$ as the initial edge of $g_i(e)$. If e is a petal, then this is the unique appearance of $\zeta_i(e)$ in $g_i(e)$.




FIGURE 9. The types of based folds for jointless train track maps, as classified by [Lemma 5.3](#).

- (3) The (e', e) -entry of $(g_i)_*^{\text{real}} - \zeta_i$ is even if e' is a filament. In fact, if $e = \zeta_i^{-1}(e')$, then the number of positive appearances of e' in $g_i(e)$ is exactly one more than the number of negative appearances of e' in $g_i(e)$. Otherwise the number of positive appearances of e' in $g_i(e)$ equals the number of negative appearances of e' in $g_i(e)$.

We let ζ be the composition of the ζ_i and the permutation induced by the final train track isomorphism h in the factorization of g . Then the previous properties imply the following proposition.

Proposition 5.4. *Let g be a jointless train track map that carries f .*

- (1) *A real edge e is a filament or a petal if and only if $\zeta(e)$ is a filament or a petal respectively.*
- (2) *The $(\zeta(e), e)$ -entry of the real transition matrix g_*^{real} is always positive. In fact, there is a canonical positive appearance of $\zeta(e)$ in the edge path $g(e)$.*
- (3) *The (e', e) -entry of $g_*^{\text{real}} - \zeta$ is even if e' is a filament. In fact, if $e = \zeta^{-1}(e')$, then the number of positive appearances of e' in $g(e)$ is exactly one more than the*




FIGURE 10. Since $g(e_k)$ is incident to $\bar{f}^{k+1}(a)$ at its initial point, the initial portion of $\bar{f}^{k+1}(\bar{\ell}^u)$ lies in the subrectangle of $\bar{f}(R_k) \cap R_{k+1}$ that corresponds to the appearance of e_{k+1} as the initial edge of $g(e_k)$.

number of negative appearances of e' in $g(e)$. Otherwise the number of positive appearances of e' in $g(e)$ equals the number of negative appearances of e' in $g(e)$.

Proposition 5.4(2) gives us a canonical edge ε_e in the directed graph Γ associated to g_*^{real} from each e to $\zeta(e)$.

The concatenation of ε_e over the filaments e is a union of curves, which we refer to as the **filament curves**.

Lemma 5.5. *The filament curves are exactly the cycles determined by the 1-pronged puncture orbits in \mathbf{a} .*

Proof. For each $a \in \mathbf{a}$, the initial portion of the unstable leaf $\bar{\ell}^u$ at a must lie in the rectangle R corresponding to the filament e that meets the 1-pronged infinitesimal polygon a , since e is the only real edge meeting a . This implies that the filaments are exactly the vertices in the cycles determined by the 1-pronged puncture orbits.

In fact, suppose $\{\bar{f}^k(a)\}_{k \in \mathbb{Z}/p}$ is the puncture orbit containing a , and suppose R_k are the rectangles containing the initial portions of the unstable leaves of $\bar{f}^k(a)$, with corresponding real edges e_k . Then since $g(e_k)$ is incident to $\bar{f}^{k+1}(a)$ at its initial point, the initial portion of $\bar{f}^{k+1}(\bar{\ell}^u)$ lies in the subrectangle of $\bar{f}(R_k) \cap R_{k+1}$ that corresponds to the appearance of e_{k+1} as the initial edge of $g(e_k)$. See **Figure 10**.

This implies that $e_{k+1} = \zeta(e_k)$, and that this appearance of e_{k+1} in $g(e_k)$ is exactly the one corresponding to the edge ε_{e_k} from e_k to e_{k+1} . Hence the filament curves are exactly the cycles determined by the 1-pronged puncture orbits. \square

On the other hand, we define a **system of petal curves** to be a maximal nonempty collection of disjoint curves that each pass through petals only. We refer to the elements of a system of petal curves as the **petal curves**. We say that a curve that passes through petals only is **persistent** if it is contained in every system of petal curves.

Lemma 5.6. *A curve α that passes through petals only is non-persistent if and only if there is another curve δ that passes through petals only and which intersects α .*

Proof. For the forward direction, we can pick δ to be a petal curve in a system that does not contain α . For the backward direction, we can build a system of petal curves by starting with δ and adding in disjoint curves that pass through petals only until we reach a maximal collection. Such a collection will not contain α since it intersects δ . \square

The concatenation of the canonical edges ε_e over the petals e determines a canonical system of petal curves. This system has the special property that every petal lies in some petal curve. Unlike the case for the filament curves, it is unclear to us whether the petal curves in this system depend on the choice of factorization of g into based folds. See [Section 13.1](#) for more discussion.

[Proposition 5.4](#)(4) says that the edges that enter filament curves always occur in pairs. This implies that if α is a curve that passes through filaments and is not a filament curve, then one can define another curve α' by following α except for an edge where α enters a filament curve, where α' instead takes the other edge in the pair of edges. We note that α and α' have the same length, but have opposite parities in terms of orientation-preserving/reversing. In this context, we say that α' is a **double** of α .

We define $G_{e',e} = \begin{cases} \frac{1}{2}(g_*^{\text{real}} - \zeta)_{e',e} & \text{if } e' \text{ is a filament} \\ (g_*^{\text{real}} - \zeta)_{e',e} & \text{if } e' \text{ is a petal} \end{cases}$ as a measure of the number of edges with this doubling phenomenon excluded. This notation will come in handy in the next subsection.

5.2. Floral train track maps. We are now ready to discuss floral train track maps.

Definition 5.7. A train track map $g : \tau \rightarrow \tau$ that carries f is **floral** if it is based with respect to a partition $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ where $\mathcal{X}_P = \{c\}$ and $\mathcal{X}_A = \mathbf{a}$ (so that g is in particular jointless), and if the two endpoints of every petal in τ coincide.

We refer to the train track on which a floral train track map is defined on to be a **floral train track**. The reason for us in discussing jointless train tracks before defining floral train tracks is so that we can make sense of the intermediate train tracks in [Lemma 5.3](#), since these might not be floral even if τ is floral.

Proposition 5.8. *The pseudo-Anosov braid f is carried by some floral train track map.*

Proof. We first consider the partition $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}'_A \sqcup \mathcal{X}'_O$ where $\mathcal{X}_P = \{c\}$ and $\mathcal{X}'_O = \{b\}$ and take some based train track partition \mathcal{M}' with respect to this partition.

We then apply [Construction 4.9](#) to convert \mathcal{M}' into a based train track partition \mathcal{M} with respect to the partition $\mathcal{X} = \mathcal{X}_P \sqcup \mathcal{X}_A \sqcup \mathcal{X}_O$ where $\mathcal{X}_P = \{c\}$ and $\mathcal{X}_A = \mathbf{a}$.




FIGURE 11. Constructing a floral train track.

We argue that $\tau = \tau_{\mathcal{M}}$ is floral. We first observe that $\tau' = \tau_{\mathcal{M}'}$ is of a specific form: We define a **limb** of τ' to be a maximal connected subset that is disjoint from the infinitesimal polygon c . If we collapse each infinitesimal polygon in a limb to a point, then we get a tree (with one vertex corresponding to where the limb is attached to c removed). The infinitesimal polygons that correspond to elements in $\mathcal{X}_O \setminus \mathcal{X}'_O$ are exactly the infinitesimal polygons in the limbs that get collapsed down to non-leaf vertices of these trees. Moreover, each limb is disjoint from the cusps of b .

To obtain \mathcal{M} from \mathcal{M}' , we split the rectangles along the unstable stars of elements in $\mathcal{X}_O \setminus \mathcal{X}'_O$ until we hit the stable star of c . Correspondingly, we split the cusps of the infinitesimal polygons in τ' that correspond to elements in $\mathcal{X}_O \setminus \mathcal{X}'_O$ until we hit the infinitesimal polygon c .

This procedure turns each limb into a union of filaments and petals (and 1-pronged infinitesimal polygons), see Figure 11. Since each limb is attached to b at one vertex, the endpoints of each petal have to lie at the same vertex. \square

Recall that, by Proposition 5.4(4), the edges that enter filaments always occur in pairs. The following proposition can be thought of as a dual property where the edges that exit petals always occur in ‘pairs’.

Proposition 5.9. *Let $g : \tau \rightarrow \tau$ be a floral train track map carrying f . For every petal p , $\sum_e G_{e,p}$ is even, where the sum is taken over all vertices e of Γ .*

Proof. A smooth edge path E in τ that starts and ends at vertices of the infinitesimal polygon c must be a concatenation $g_1 c_1 g_2 c_2 \cdots g_L c_L g_{L+1}$, where each g_i is of the form

- $f_i^{-1} a_i f_i$, where f_i is a filament meeting a 1-pronged infinitesimal polygon a_i , or
- p_i , where p_i is a petal,

and each c_i is an edge in c . In particular, the endpoints of each g_i lie on the same vertex, while the endpoints of each c_i lie on adjacent vertices.

We apply this observation to the smooth edge path $g(e)$. Here we have $\sum_{e'} G_{e',e} = L$. But since the two endpoints of e lie on the same vertex, the same is true for $g(e)$, so L must be even. \square

However, notice that [Proposition 5.9](#) is not a perfect dual of [Proposition 5.4\(4\)](#) since the ‘pair’ of edges may not have the same terminal points, i.e. we do not claim that each individual $G_{e,p}$ is even.

Regardless, [Proposition 5.9](#) is very helpful for ensuring that many edges exit petals. For example, suppose $\alpha = (\varepsilon_k : e_k \rightarrow e_{k+1})_{k \in \mathbb{Z}/p}$ is an orientation-preserving cycle determined by a periodic orbit \mathbf{a} , where $\mathbf{a} \notin \mathcal{X}_I$, and suppose α passes through petals only. Then the second item of [Proposition 4.15](#) implies that $\sum_k \sum_e G_{e,e_k} \geq 1$. With [Proposition 5.9](#) we can then upgrade this to $\sum_k \sum_e G_{e,e_k} \geq 2$.

In fact, we can often do even better: If some $\sum_e G_{e,e_k} > 2$ for some value of k , or if $\sum_e G_{e,e_k} > 0$ for two values of k , then by [Proposition 5.9](#), $\sum_k \sum_e G_{e,e_k} \geq 4$. The following proposition says that if this is not true, then we can also extract some information. We will need this proposition to wrap up one of the cases in our analysis.

Proposition 5.10. *Suppose $\alpha = (\varepsilon_k : e_k \rightarrow e_{k+1})_{k \in \mathbb{Z}/p}$ is an orientation-preserving cycle determined by a periodic orbit \mathbf{a} , where $\mathbf{a} \notin \mathcal{X}_I$, and suppose α passes through petals only. If $\sum_k \sum_e G_{e,e_k} = 2$, then either*

- $G_{p,e_k} > 0$ for some petal p and some (unique) k , or
- $G_{f,e_k} = 2$ for some filament f and some (unique) k , and there exists some filament f' such that $G_{f,f'} \geq 1$.

Proof. By [Proposition 5.9](#), if $\sum_k \sum_e G_{e,e_k} = 2$ then $\sum_e G_{e,e_k} = 2$ for some unique k . Moreover, in the notation of the proof of that proposition, the smooth edge path $g(e_k)$ must be of the form $g_1 c_1 e_{k+1} c_2 g_3$, where each g_i is of the form

- $f_i^{-1} a_i f_i$, where f_i is a filament, or
- p_i , where p_i is a petal,

If g_1 or g_3 is of the second form, then we belong to the first case. If g_1 and g_3 are both of the first form, we claim that $f_1 = f_3$.

To see this, let R be the rectangle corresponding to e_k . Let u be the unstable side of R that lies closer to b . Let v be the vertex of c that is the common endpoint of e_k . There is a segment s of the side of the stable star at c corresponding to v such that $u \cup s$ bounds a disc D not containing b in the sphere \bar{S} . See [Figure 12](#) top.

Taking the image under \bar{f} , $\bar{f}(u) \cup \bar{f}(s)$ bounds a disc $\bar{f}(D)$ that does not contain b as well.




FIGURE 12. If $f_1 \neq f_3$, then either $\bar{f}(u)$ intersects $\bar{f}(s)$ (bottom left and middle) or $\bar{f}(D)$ contains c (bottom right).




FIGURE 13. Let f' be a filament enclosed by e_k . Then $G_{f,f'} \geq 1$.

Now assume $f_1 \neq f_3$. Then depending on the relative positions of the subpaths of $\bar{f}(u)$ in the rectangles corresponding to f_1 and f_3 , either $\bar{f}(u)$ intersects $\bar{f}(s)$ as in [Figure 12](#) bottom left and middle, or $\bar{f}(D)$ contains c as in [Figure 12](#) bottom right. In both cases we arrive at a contradiction, thus f_1 and f_3 is some common filament f .

Meanwhile, e_k must enclose some 1-pronged puncture, i.e. D must contain some 1-pronged puncture. Indeed, consider the innermost non-1-pronged boundary component d that is enclosed by e_k . If d is n -pronged then it must enclose $n - 1$ 1-pronged infinitesimal polygons. Let f' be a filament connecting to one of these 1-pronged infinitesimal polygons. Then we must have $G_{f,f'} \geq 1$. See [Figure 13](#). \square

Finally, we record the following fact.

Lemma 5.11. *The cycles $\beta_1, \dots, \beta_{\frac{p_0}{p}}$ determined by b are each embedded and are mutually disjoint. Similarly, the cycles $\gamma_1, \dots, \gamma_{\frac{q_0}{q}}$ determined by c are each embedded and are mutually disjoint.*

Proof. For some β_i to be non-embedded, there must be some real edge of τ that lies on two sides of the boundary component b . But each filament, despite possibly appearing twice on b , lies on at most a single side, and each petal appears at most once on b , so this is impossible. The same reason implies that the cycles determined by b are mutually disjoint.

For some γ_i to be non-embedded, there must be some real edge of τ that meets two vertices of the infinitesimal polygon c , but this is not the case by the definition of floral. The same reason implies that the cycles determined by c are mutually disjoint. \square

6. SETUP FOR COMPUTATIONS

The proof of [Theorem 1.1](#) will occupy the rest of this paper. In this section, we set up some notation and explain the casework for our computations.

6.1. Inequalities. Let f be a pseudo-Anosov braid with n strands. Let \bar{f} be the completion of f . Let λ be the dilatation of f .

We define the singular orbits \mathbf{a} , $\{b\}$, $\{c\}$ as in [Section 5.1](#). Recall that in defining c , we might have created an additional puncture. When we do so, we do not change the value of n ; we simply allow f to become a braid with $n + 1$ strands.

We denote by m the cardinality of \mathbf{a} . We denote by p_0 and q_0 the number of prongs of b and c , respectively.

We refer to the singular points in $\mathbf{a} \cup \{b, c\}$ as the **primary** singular points, and refer to the remaining singular points as the **secondary** singular points. We denote by r_0 the number of secondary singular points. This number may be zero or positive.

Lemma 6.1. *If $g : \tau \rightarrow \tau$ is a floral train track map that carries f . Then the number of filaments in τ equals m and the number of petals in τ equals r_0 .*

Proof. The number of filaments equals the cardinality of \mathbf{a} , which is m . Meanwhile, the number of boundary components of τ equals the number of singularities, which is $m + r_0 + 2$. By [Proposition 4.2](#), the number of real edges equals $(m + r_0 + 2) - 2 = m + r_0$. Hence the number of petals equals r_0 . \square

Lemma 6.2. *The numbers m, n, p_0, q_0, r_0 satisfy $m \leq n$ and $p_0 + q_0 + r_0 \leq n$.*

Proof. That $m \leq n$ follows from the definition of m and n .

For the second inequality, let us first suppose that c is unpunctured. The secondary singular points can be separated into the punctured ones and the unpunctured ones. Since we are assuming that c is unpunctured, there are $n - m$ such punctured singularities.

Each of these must be ≥ 2 -pronged, hence have Poincaré-Hopf index ≤ -1 . Meanwhile, there are $r_0 - (n - m)$ such unpunctured singularities. Each of these must be ≥ 3 -pronged, hence have Poincaré-Hopf index $\leq -\frac{1}{2}$.

Applying this observation to [Theorem 2.1](#), we have

$$\chi(S) = \sum_{a \in \mathbf{a}} \text{ind}_{\text{PH}}(a) + \text{ind}_{\text{PH}}(b) + \text{ind}_{\text{PH}}(c) + \sum_{\text{punc.}} \text{ind}_{\text{PH}} + \sum_{\text{unpunc.}} \text{ind}_{\text{PH}}$$

$$(6.1) \quad 1 - n \leq -\frac{m}{2} - \frac{p_0}{2} + \left(1 - \frac{q_0}{2}\right) - (n - m) - \frac{r_0 - (n - m)}{2}$$

Hence $p_0 + q_0 + r_0 \leq n$.

If c is punctured and this is not because of an additional puncture we did in order to make c singular, then in [Equation \(6.1\)](#), one replaces $1 - \frac{p_0}{2}$ with $-\frac{p_0}{2}$ and replaces $n - m$ with $n - m - 1$. In particular the right hand side does not change, so one ends up with the same inequality.

Finally, if c is punctured and this is because of an additional puncture we did in order to make c singular, then in [Equation \(6.1\)](#), one replaces $1 - n$ with $-n$ and replaces $1 - \frac{p_0}{2}$ with $-\frac{p_0}{2}$. In particular the right hand side again does not change, so one ends up with the same inequality. \square

We denote by p and q the rotationless periods of b , and c respectively. We have $p \mid p_0$ and $q \mid q_0$. In particular from [Lemma 6.2](#), we have $p + q + r_0 \leq n$.

Also, up to switching b and c , we can assume that either

$$(6.2) \quad \begin{cases} q \neq r_0, \text{ or} \\ p = q = r_0. \end{cases}$$

The following lemma states that b and c in turn induce another pair of periodic points.

Lemma 6.3. *There exists periodic points b' and c' such that*

- b' is a rotated fixed point of f^p and c' is a rotated fixed point of f^q , and
- the orbits of b, b', c, c' are distinct.

Proof. From [Proposition 2.5](#) we know that $\sum_x \text{ind}_{\text{Lef}}(\bar{f}^p, x) = 2$, where the sum is taken over all fixed points of \bar{f}^p . From [Equation \(2.2\)](#) we observe that b contributes $1 - p_0$, c contributes at most 1, and elements in \mathbf{a} (possibly) contributes 0.

Thus there are at least p_0 points other than b and c that are rotated fixed points of \bar{f}^p . We pick one such point b' . Let p'_0 be the period of b' .

We follow the same strategy for c' . From [Proposition 2.5](#) we know that $\sum_x \text{ind}_{\text{Lef}}(\bar{f}^q, x) = 2$, where the sum is taken over all fixed points of \bar{f}^q . From [Equation \(2.2\)](#) we observe

that b contributes at most 1, c contributes $1 - q_0$, points in the orbit of b' (possibly) contribute at most p'_0 , and elements in \mathbf{a} (possibly) contribute 0.

If p'_0 does not divide q , i.e. points in the orbit of b' do not contribute to the sum, then there are at least q_0 points other than b and c that are rotated fixed points of \bar{f}^q , and we can pick c' to be one such point.

If p'_0 does divide q , i.e. points in the orbit of b' do contribute to the sum, then unless $p'_0 = q$, there are at least $q_0 - p'_0 \geq q - p'_0 > 0$ points other than b , c , and (possibly) the points in the orbit of b' that is a rotated fixed point of \bar{f}^q , and we can pick c' to be one such point.

The remaining case is if $p'_0 = q$. This can only happen if $p \geq q$. Similarly, the symmetric strategy of first picking c' then b' works unless $q \geq p$.

Thus it remains to tackle the case when $p = q$. In this case, we have to use a slightly different argument. From [Proposition 2.5](#) we know that $\sum_x \text{ind}_{\text{Lef}}(\bar{f}^p, x) = 2$, where the sum is taken over all fixed points of \bar{f}^p . From [Equation \(2.2\)](#) we observe that b contributes $1 - p_0$, c contributes $1 - q_0$, and elements in \mathbf{a} (possibly) contribute 0.

Thus there are at least $p_0 + q_0 \geq 2p$ points other than b and c that are rotated fixed points of \bar{f}^p . There must be at least two orbits of such points. We pick b' to be one such point, and pick c' to be a point in some other orbit. \square

We have the following upgrade of [Lemma 6.2](#).

Lemma 6.4.

(1) *Suppose b' is singular. We denote by p' the rotationless period of b' . Then*

(a) $p_0 + q_0 + \frac{1}{3}p' \leq n$, and

(b) for every $\kappa \in [-1, 2]$, we have $(\kappa - 1)p_0 + q_0 + r_0 + \frac{2-\kappa}{3}p' \leq n$

(2) *Suppose b' and c' are singular. We denote by p' and q' the rotationless period of b' and c' respectively. Then*

(a) $p_0 + q_0 + \frac{1}{3}p' + \frac{1}{3}q' \leq n$, and

(b) for every $\kappa_1, \kappa_2 \in [-1, 2]$, we have $(\kappa_1 - 1)p_0 + (\kappa_2 - 1)q_0 + r_0 + \frac{2-\kappa_1}{3}p' + \frac{2-\kappa_2}{3}q' \leq n$

Proof. Let us first suppose that b' is singular and unpunctured. Let p'_0 be the period of b' and $k \geq 3$ be the number of prongs at b' . By splitting off the orbit of b' from the last term of [Equation \(6.1\)](#), we have

$$1 - n \leq -\frac{m}{2} - \frac{p_0}{2} + \left(1 - \frac{q_0}{2}\right) - (n - m) - \frac{r_0 - (n - m) - p'_0}{2} + p'_0\left(1 - \frac{k}{2}\right).$$

Here we are assuming that c is unpunctured. If this is not the case then one modifies the terms to arrive at the same inequality as in the proof of [Lemma 6.2](#).

To show (1a), we bound the fourth term by $-\frac{1}{2}(n-m)$ and bound the fifth term by 0. This gives us $p_0 + q_0 + (k-2)p'_0 \leq n$. Since $k \geq 3$ and $p' \leq kp_0$, we have

$$\begin{aligned} p_0 + q_0 + \frac{1}{3}p' &\leq p_0 + q_0 + \frac{k-2}{k}p' \\ &\leq p_0 + q_0 + (k-2)p'_0 \\ &\leq n. \end{aligned}$$

To show (1b), we retain all the terms, which gives us $p_0 + q_0 + r_0 + (k-3)p'_0 \leq n$. Since $p'_0 \leq p \leq p_0$ and $p' \leq kp_0$, for every $\kappa \in [-1, 2]$ we have

$$\begin{aligned} (\kappa-1)p_0 + q_0 + r_0 + \frac{2-\kappa}{3}p' &\leq (\kappa-1)p_0 + q_0 + r_0 + \frac{(2-\kappa)k}{3}p'_0 \\ &= (\kappa-1)p_0 + (2-\kappa)p'_0 + q_0 + r_0 + \frac{(2-\kappa)(k-3)}{3}p'_0 \\ &\leq (\kappa-1)p_0 + (2-\kappa)p_0 + q_0 + r_0 + (k-3)p'_0 \\ &= p_0 + q_0 + r_0 + (k-3)p'_0 \\ &\leq n. \end{aligned}$$

Now suppose b' is singular and punctured. Let p'_0 be the period of b' and $k \geq 2$ be the number of prongs at b' . By splitting off the orbit of b' from the second-to-last term of Equation (6.1), we have

$$1 - n \leq -\frac{m}{2} - \frac{p_0}{2} + \left(1 - \frac{q_0}{2}\right) - (n - m - p'_0) - \frac{r - (n - m)}{2} + p'_0\left(-\frac{k}{2}\right).$$

Here, as above, we are assuming that c is unpunctured.

To show (1a), we bound the fourth term by $-\frac{1}{2}(n-m-p'_0)$ and bound the fifth term by 0. This gives us $p_0 + q_0 + (k-1)p'_0 \leq n$. Since $k \geq 2$ and $p' \leq kp_0$, we have

$$\begin{aligned} p_0 + q_0 + \frac{1}{3}p' &\leq p_0 + q_0 + \frac{k-1}{k}p' \\ &\leq p_0 + q_0 + (k-1)p'_0 \\ &\leq n. \end{aligned}$$

For (1b), if $k \geq 3$, then the computation in the previous case carries through: For every $\kappa \in [-1, 2]$ we have

$$\begin{aligned} (\kappa-1)p_0 + q_0 + r_0 + \frac{2-\kappa}{3}p' &\leq (\kappa-1)p_0 + (2-\kappa)p'_0 + q_0 + r_0 + \frac{(2-\kappa)(k-3)}{3}p'_0 \\ &\leq (\kappa-1)p_0 + (2-\kappa)p_0 + q_0 + r_0 + \frac{k-3}{2}p'_0 \\ &= p_0 + q_0 + r_0 + \frac{k-3}{2}p'_0 \\ &\leq p_0 + q_0 + r_0 + \frac{k-2}{2}p'_0 \end{aligned}$$

$$\leq n.$$

If $k = 2$, then the second inequality above is incorrect, so we have to use a different chain of inequalities:

$$\begin{aligned} (\kappa - 1)p_0 + q_0 + r_0 + \frac{2 - \kappa}{3}p' &\leq (\kappa - 1)p_0 + (2 - \kappa)p'_0 + q_0 + r_0 - \frac{2 - \kappa}{3}p'_0 \\ &\leq (\kappa - 1)p_0 + (2 - \kappa)p_0 + q_0 + r_0 \\ &= p_0 + q_0 + r_0 \\ &\leq n. \end{aligned}$$

The inequalities in (2) can be proved using similar computations. \square

6.2. Division of cases. Let $g : \tau \rightarrow \tau$ be a floral train track that carries f . Let Γ be the directed graph associated to the real transition matrix g_*^{real} . Let (G, w) be the curve complex of Γ .

Let β be some cycle determined by b and γ be some cycle determined by c . Our analysis is divided into the following six cases.

(I) Both β and γ pass through filaments.

This case includes the situation when $r = 0$, i.e. when there are no secondary singular points. We will encounter the values $\underline{\delta}_n$ in this case.

(II) One of β and γ passes through both filaments and petals while the other passes through petals only.

(III) One of β and γ passes through filaments only while the other passes through petals only and is not persistent.

(IV) One of β and γ passes through filaments only while the other passes through petals only and is persistent.

(V) Both β and γ pass through petals only, and at least one of them is not persistent.

(VI) Both β and γ pass through petals only, and both are persistent.

7. CASE I: BOTH β AND γ PASS THROUGH FILAMENTS

The goal of this section is to prove the following proposition. The assumptions in the proposition are more general than the stated hypothesis for this case. The added generality will come into play in the later cases.

Proposition 7.1. *Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β and γ are orientation-preserving, and
- β and γ pass through filaments and are not filament curves.


Then $\lambda^n \geq \min\{14.5, \delta_n^n\}$.

We first argue that we can assume β and γ are embedded.

Lemma 7.2. *If there is a cycle μ in Γ with length $\leq \frac{n}{2}$ and which consists of at least two distinct curves, then $\lambda^n \geq 16$.*

Proof. If μ contains at least two distinct curves, then we can find two curves μ_1, μ_2 in μ such that μ_1 intersects μ_2 .

Thus the following subgraph G_1 must be an induced subgraph of the curve complex G :



where $u_i = \text{length}(\mu_i)$ have sum $\leq \text{length}(\mu) \leq \frac{n}{2}$.

By convexity and monotonicity, the minimum growth rate of G_1 subject to the condition $u_1 + u_2 \leq \frac{n}{2}$ is attained when $u_1 = u_2 = \frac{n}{4}$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= 1 - 2t^{\frac{n}{4}} \\ &= 1 - 2x^{\frac{1}{4}} \end{aligned}$$

where we changed variables $x = t^n$.

The reciprocal of the smallest positive root of this polynomial in x is 16. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 16$. \square

Lemma 7.3. *If there is a cycle μ in Γ with length $\leq n$ and which consists of at least three distinct curves, then $\lambda^n \geq 16$.*

Proof. If μ contains at least three distinct curves, then we can find three curves μ_1, μ_2, μ_3 in μ such that μ_2 intersects μ_1 and μ_3 .

Thus, depending on whether μ_1 intersects μ_3 , one of the following two subgraphs must be an induced subgraph of the curve complex G :



where $u_i = \text{length}(\mu_i)$ have sum $\leq \text{length}(\mu) \leq n$.

The graph on the left is a wide subgraph of the one on the right, so the growth rate of G is always bounded from below by the growth rate of the graph G_1 on the right.

By convexity and monotonicity, the minimum growth rate of G_1 subject to the condition $u_1 + u_2 + u_3 \leq n$ is attained when $u_1 = u_3$ and $u_2 = n - 2u_1$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= 1 - 2t^{u_1} - t^{n-2u_1} + t^{2u_1} \\ &= 1 - 2x^a - x^{1-2a} + x^{2a} \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{u_1}{n}$.

Via a standard calculus exercise, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is 16, attained when $a = \frac{1}{4}$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 16$. \square

Since β passes through filaments and is not a filament curve, it contains at least one curve β_1 that passes through filaments and is not a filament curve. Similarly, γ contains at least one curve γ_1 that passes through filaments and is not a filament curve.

We let β'_1 be a double of β_1 and γ'_1 be a double of γ_1 . If $\beta_1, \beta'_1, \gamma_1, \gamma'_1$ are distinct, then, depending on which one of β_1, β'_1 and which one of γ_1, γ'_1 is orientation-preserving, we can replace β by β_1 or β'_1 and γ by γ_1 or γ'_1 to arrange it so that β and γ are embedded.

Otherwise, suppose $\beta_1 = \gamma_1$. Then since $\beta \neq \gamma$, we have $\beta \neq \beta_1$ or $\gamma \neq \gamma_1$. Without loss of generality suppose $\beta \neq \beta_1$. Then β contains another curve β_2 .

If β contains a third curve β_3 , then [Lemma 7.3](#) implies that $\lambda^n \geq 16$.

If β does not contain a third curve, but $\beta_2 = \beta'_1$, then β_1 and β_2 would have opposite parity, and β must traverse β_1 or β_2 at least twice, for otherwise it would be orientation-reversing. Thus $4 \text{length}(\beta_1) = 3 \text{length}(\beta_1) + \text{length}(\gamma_1) \leq \text{length}(\beta) + \text{length}(\gamma) \leq n$. The curve obtained by traversing β_1 and β'_1 has length $2 \text{length}(\beta_1) \leq \frac{n}{2}$, so by [Lemma 7.2](#) we have $\lambda^n \geq 16$.

If β does not contain a third curve and $\beta_2 \neq \beta'_1$, then the curve obtained by traversing $\beta_1, \beta'_1, \beta_2$ has length $(\text{length}(\beta_1) + \text{length}(\beta_2)) + \text{length}(\gamma_1) \leq \text{length}(\beta) + \text{length}(\gamma) \leq n$, so by [Lemma 7.3](#) we have $\lambda^n \geq 16$.

Thus we can assume from this point onwards that β and γ are embedded. We denote the lengths of β and γ by p and q respectively.

Next, we argue that we can assume β and γ each enter filament curves only once, i.e. each of β and γ contains exactly one edge that is not part of a filament curve.

Lemma 7.4. *If β enters filament curves at least twice, then $\lambda^n \geq 14.6$.*

Proof. If β enters filament curves at least twice, then by following β but taking the other edge in each of the two pairs of edges, we get a quadruple of curves $\beta = \beta_1, \beta_2, \beta_3, \beta_4$ that are pairwise doubles (in two ways).


If γ is not one of the β_i then we can pick a double γ' of γ that is also not one of the β_i . Thus, depending on whether β and γ intersect, the curve complex must contain an induced subgraph of one of the following two forms:



where

- $\times N$ denotes N vertices of the denoted weight,
- an edge between two vertices labelled $\times N_1$ and $\times N_2$ denotes an edge going between each of the N_1 vertices and each of the N_2 vertices,
- the vertices in the top row are the filament curves, and
- the edges between the vertices in top row and those in the bottom row depend on which ones of the filament curves β and γ intersect.

The graph on the left is a wide subgraph of the one on the right, so the growth rate of G is always bounded from below by the growth rate of the graph on the right. Furthermore, the growth rate of the graph on the right is bounded from below by that of the following graph G_1 :



obtained by summing together the vertices in the top row.


By monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$ and $p + q \leq n$ is attained when $m = n$ and $q = n - p$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= 1 - 4t^p - 2t^{n-p} + 7t^n \\ &= 1 - 4x^a - 2x^{1-a} + 7x \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{p}{n}$.

Via a standard calculus exercise, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is $9 + 4\sqrt{2} \geq 14.6$, attained near $a \approx 0.63$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.6$.

If instead γ is one of the β_i , then $p = q$. Reasoning as above, the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$ and $2p \leq n$ is attained when $m = n$ and $p = \frac{n}{2}$. In this case, the clique polynomial of G_1 is

$$\begin{aligned}
 Q_1(t) &= 1 - 4t^{\frac{n}{2}} - t^n \\
 &= 1 - 4x^{\frac{1}{2}} - x
 \end{aligned}$$


where we changed variables $x = t^n$.

The reciprocal of the smallest positive root of this polynomial in x is $9 + 4\sqrt{5} \geq 17.9$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.6$. \square


Thus we can assume from this point onwards that β and γ each enter filament curves only once. In particular, each of them only intersect one filament curve. We define β' and γ' to be the (unique) doubles of β and γ respectively.

Lemma 7.5. *If β and γ intersect, then $\lambda^n \geq 17.9$*

Proof. If β and γ intersect, then so do the doubles β' and γ' , so the curve complex G must contain an induced subgraph of the following form:



where vertices in the top row are the filament curves as in [Lemma 7.4](#). By summing these vertices, the growth rate of G is bounded from below by that of the following graph G_1 :



By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$ and $p + q \leq n$ is attained when $m = n$ and $p = q = \frac{n}{2}$. In this case, the clique polynomial of G_1 is

$$Q_1(t) = 1 - 4t^{\frac{n}{2}} - t^n$$

$$= 1 - 4x^{\frac{1}{2}} - x$$

where we changed variables $x = t^n$.


The reciprocal of the smallest positive root of this polynomial in x is $9 + 4\sqrt{5} \geq 17.9$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 17.9$. \square

Thus we can assume from this point onwards that β and γ are disjoint.

Next, we rule out the existence of petals.

Lemma 7.6. *If β passes through petals, then $\lambda^n \geq 25$.*

Proof. We pick the canonical system of petal curves given by [Proposition 5.4\(2\)](#). The growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



where the top vertex is the sum of the filament curves and the bottom vertex is the sum of the petal curves.

By monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$ and $p + q + r_0 \leq n$ is attained when $m = n$ and $r_0 = n - p - q$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= (1 - 2t^p)(1 - 2t^q) - t^n - t^{n-p-q}(1 - 2t^q - t^n) \\ &= (1 - 2x^a)(1 - 2x^b) - x - x^{1-a-b}(1 - 2x^b - x) \end{aligned}$$

where we changed variables $x = t^n$, $a = \frac{p}{n}$, and $b = \frac{q}{n}$.


Via a standard calculus exercise, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is 25, attained near $a = 0.5$ and $b \approx 0.28$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 25$. \square

Similarly, if γ passes through petals, then $\lambda^n \geq 25$ as well.

Lemma 7.7. *If τ has any petals, but β and γ do not pass through petals, then $\lambda^n \geq 19.9$.*

Proof. We pick the canonical system of petal curves given by [Proposition 5.4\(2\)](#). Since the directed graph Γ is strongly connected, there must exist a curve μ intersecting some filament curve and some petal curve. In particular, since μ enters a filament curve, it has a double μ' .

If μ intersects β and γ , then the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



where $u = \text{length}(\mu) \leq m + r_0 \leq n + r_0$.


By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$, $p + q + r_0 \leq n$, and $u \leq n + r_0$ is attained when $m = n$, $p = q$, $r_0 = n - 2p$, and $u = n + r_0 = 2n - 2p$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= ((1 - 2t^p)^2 - t^n)(1 - t^{n-2p}) - 2t^{2n-2p} \\ &= ((1 - 2x^a)^2 - x)(1 - x^{1-2a}) - 2x^{2-2a} \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{p}{n}$.

Via a standard calculus exercise, and with computational aid, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is ≥ 20 , attained near $a \approx 0.40$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 19.9$.

If μ intersects β but not γ , then the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



where $u = \text{length}(\mu)$ satisfies $q + u \leq m + r_0 \leq n + r_0$.


By monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$, $p + q + r_0 \leq n$, and $q + u \leq n + r_0$ is attained when $m = n$, $r_0 = n - p - q$, and $u = n + r_0 - q = 2n - p - 2q$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= ((1 - 2t^p)(1 - 2t^q) - t^n)(1 - t^{n-p-q}) - 2t^{2n-p-2q}(1 - 2t^q) \\ &= ((1 - 2x^a)(1 - 2x^b) - x)(1 - x^{1-a-b}) - 2x^{2-a-2b}(1 - 2x^b) \end{aligned}$$

where we changed variables $x = t^n$, $a = \frac{p}{n}$ and $b = \frac{q}{n}$.

Via a standard calculus exercise, and with computational aid, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is $13 + 4\sqrt{3} \geq 19.9$, attained near $a \approx 0.45$ and $b \approx 0.32$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 19.9$.

If μ does not intersect β and γ , then the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



where $u = \text{length}(\mu)$ satisfies $p + q + u \leq m + r_0 \leq n + r_0$.

By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$, $p + q + r_0 \leq n$, and $p + q + u \leq n + r_0$ is attained when $m = n$, $p = q$, $r_0 = n - 2p$, and $u = n + r_0 - p - q = 2n - 4p$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= ((1 - 2t^p)^2 - t^n)(1 - t^{n-2p}) - 2t^{2n-4p}(1 - 2t^p)^2 \\ &= ((1 - 2x^a)^2 - x)(1 - x^{1-2a}) - 2x^{2-4a}(1 - 2x^a)^2 \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{p}{n}$.

Via a standard calculus exercise, and with computational aid, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is ≥ 21 , attained near $a \approx 0.34$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 19.9$. \square


Thus we can assume from this point onwards that there are no petals.

Next, we rule out the existence of more than one filament curve. Recall that we have reduced to the case when β and γ each intersect a unique filament curve.

Lemma 7.8. *If β and γ intersect distinct filament curves, then $\lambda^n \geq 16$.*

Proof. Let α_1 be the filament curve that β intersects. Since Γ is strongly connected, and since we have reduced to the case when there are no petals, there is a curve μ intersecting α_1 and some other filament curve. Since μ enters filament curves at least twice, we can find a quadruple of curves $\mu = \mu_1, \mu_2, \mu_3, \mu_4$ by following μ .

Thus the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



where the top left vertex is α_1 while the top right vertex is the sum of the rest of the filament curves, so $m_1 + m_2 = m \leq n$.

By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m_1 + m_2 \leq n$, $p + q \leq n$, and $u \leq n$ is attained when $m_1 = m_2 = p = q = \frac{n}{2}$ and $u = n$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= (1 - 3t^{\frac{n}{2}})^2 - 4t^n(1 - 2t^{\frac{n}{2}})^2 \\ &= (1 - 3x^{\frac{1}{2}})^2 - 4x(1 - 2x^{\frac{1}{2}})^2 \end{aligned}$$


where we changed variables $x = t^n$.

The reciprocal of the smallest positive root of this polynomial in x is 16. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 16$. \square

Lemma 7.9. *If β and γ intersect the same filament curve but there are other filament curves, then $\lambda^n \geq 68$.*

Proof. Let α_1 be the filament curve that β and γ intersect. Since Γ is strongly connected, and since we have reduced to the case when there are no petals, there is a curve μ intersecting α_1 and some other filament curve. Since μ enters filament curves at least twice, we can find a quadruple of curves $\mu = \mu_1, \mu_2, \mu_3, \mu_4$ by following μ .

Thus the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m_1 + m_2 \leq n$, $p + q \leq m_1$, and $u \leq n$ is attained when $m_2 = n - m_1$, $p = q = \frac{m_1}{2}$, and $u = n$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= (1 - 2t^{\frac{m_1}{2}})^2(1 - t^{m_1} - 4t^n) - t^{n-m_1}(1 - t^{m_1}) \\ &= (1 - 2x^a)^2(1 - x^{2a} - 4x) - x^{1-2a}(1 - x^{2a}) \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{m_1}{2n}$.

Via a standard calculus exercise, and with computational aid, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is ≥ 68 , attained near $a \approx 0.33$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 68$. \square

Thus we can assume from this point onwards that there is only one filament curve.


To summarize, we have reduced to the situation where we have

- a single filament curve α ,
- a curve β following α except for one edge e_1 , which has a double e'_1 , and
- a curve γ following α except for one edge e_2 , which has a double e'_2 ,

where β and γ are disjoint, thus in particular e_1, e'_1, e_2 , and e'_2 are distinct.

Lemma 7.10. *Either $\lambda^n \geq 17.9$ or there is an edge e_3 other from e_1, e'_1, e_2 , and e'_2 .*

Proof. Suppose there is no edge e_3 other from e_1, e'_1, e_2 , and e'_2 , then the curve complex G is exactly



Without loss of generality suppose $p \leq q$. The clique polynomial of G is

$$Q(t) = 1 - 2t^p - 2t^q + 4t^{p+q} - t^m,$$

where we have ordered the terms in increasing order of their exponents, except possibly $p = q$ or $p + q = m$.

By [Theorem 3.5](#) and [Theorem 4.12\(4\)](#), we know that $Q(t)$ is reciprocal. This can only happen if $p = q = \frac{m}{3}$, in which case we have

$$\begin{aligned} Q(t) &= 1 - 4t^{\frac{m}{3}} + 4t^{\frac{2m}{3}} - t^m \\ &= 1 - 4x^{\frac{1}{3}} + 4x^{\frac{2}{3}} - x \end{aligned}$$


where we changed variables $x = t^m$.

The reciprocal of the smallest positive root of this polynomial in x is $9 + 4\sqrt{5} \geq 17.9$. Thus $\lambda^n \geq \lambda^m \geq \lambda(G, w)^m \geq 17.9$. \square

Let μ be the curve obtained by traversing e_3 then following α until the path closes up, and let μ' be the curve defined similarly but using the double e'_3 of e_3 .

Lemma 7.11. *If μ does not intersect β and γ , then $\lambda^n \geq 27$.*

Proof. In this case the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$ and $p + q + u \leq n$ is attained when $m = n$ and $p = q = u = \frac{n}{3}$. In this case, the clique polynomial of G_1 is


$$\begin{aligned} Q_1(t) &= (1 - 2t^{\frac{n}{3}})^3 - t^n \\ &= (1 - 2x^{\frac{1}{3}})^3 - x \end{aligned}$$

where we changed variables $x = t^n$.

The reciprocal of the smallest positive root of this polynomial in x is 27. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 27$. \square

Lemma 7.12. *If μ intersects β but not γ , then $\lambda^n \geq 14.6$.*

Proof. In this case the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n$, $p + q \leq n$, and $q + u \leq n$ is attained when $m = n$ and $p = u = n - q$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= (1 - 2t^q)(1 - 4t^{n-q}) - t^n \\ &= (1 - 2x^a)(1 - 4x^{1-a}) - x \end{aligned}$$

where we changed variables $x = t^n$ and $a = \frac{q}{n}$.


Via a standard calculus exercise, one can compute that the minimum value of the reciprocal of the smallest positive root of this polynomial in x is $9 + 4\sqrt{2} \geq 14.6$, attained near $a \approx 0.37$. Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.6$. \square

Thus we can assume μ intersects both β and γ . In other words, e_3 must go from a vertex of β to a vertex of γ , or vice versa. Let us call an edge satisfying this property a **bridge**. We can thus assume that every edge other than e_1, e'_1, e_2 , and e'_2 is a bridge.

Lemma 7.13. *If there are at least two bridges that are not doubles of each other, then $\lambda^n \geq 17.8$.*

Proof. Let e_3 and e_4 be two bridges that are not doubles of each other. We can construct curves μ and ν passing through e_3 and e_4 as above.

Thus the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions $m \leq n, p + q \leq n, u \leq n$, and $v \leq n$ is attained when $m = n$ and $p = q = \frac{n}{2}$ and $u = v = n$. In this case, the clique polynomial of G_1 is

$$\begin{aligned} Q_1(t) &= (1 - 2t^{\frac{n}{2}})^2 - 5t^n + t^{2n} \\ &= (1 - 2x^{\frac{1}{2}})^2 - 5x + x^2 \end{aligned}$$


where we changed variables $x = t^n$.

The reciprocal of the smallest positive root of this polynomial in x is ≥ 17.8 . Thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq 17.8$. \square

Thus we can assume that e_3 is the only bridge. We obtain our lower bound in this specific scenario.

Lemma 7.14. *If there are only two bridges (which are necessarily doubles of each other), then $\lambda^n \geq \min\{14.5, \underline{\delta}_n^n\}$.*

Proof. In this case the growth rate of the curve complex G is bounded from below by that of the following graph G_1 :



By convexity and monotonicity, the minimum growth rate of G_1 subject to the conditions

- $p + q \leq m$,
- $u \leq m$,
- p, q, u are integers, and
- the characteristic polynomial $P(t) = Q(t^{-1})$ of g_*^{real} has a unique root of maximum modulus (which follows from the Perron-Frobenius theorem)

is attained when $u = m$ and

$$(p, q) = \begin{cases} (k, k + 1) & \text{if } m = 2k + 1 \\ (2k - 1, 2k + 1) & \text{if } m = 4k \\ (2k - 1, 2k + 3) & \text{if } m = 4k + 2 \end{cases}$$

In this case, the reciprocal of the smallest positive root of the clique polynomial is $\underline{\delta}_m$.

For $n \in \underline{N}$, we have $\underline{\delta}_m \geq \underline{\delta}_n$ for $m \leq n$, thus $\lambda \geq \lambda(G_1, w_1) \geq \underline{\delta}_m \geq \underline{\delta}_n$. For $n \notin \underline{N}$, we have $\underline{\delta}_m^n \geq 14.5$ for $m \leq n$, thus $\lambda^n \geq \lambda(G_1, w_1)^n \geq \underline{\delta}_m^n \geq 14.5$. \square

8. CASE II: ONE OF β AND γ PASSES THROUGH BOTH FILAMENTS AND PETALS WHILE THE OTHER PASSES THROUGH PETALS ONLY

The goal of this section is to prove the following proposition. As in case I, the assumptions in the proposition are more general than the stated hypothesis for this case. The added generality will come into play in the later cases.

Proposition 8.1. *Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β passes through both petals and filaments, and
- γ passes through petals only.


Then $\lambda^n \geq 14.5$.

Up to replacing β with a curve that it contains, we can assume that it is embedded. Similarly, up to replacing γ with a curve that it contains, we can assume that it is embedded. We denote the lengths of β and γ by p and q respectively. Throughout this section, we fix the canonical system of petal curves given by [Proposition 5.4\(2\)](#).

First, we rule out β and γ intersecting each other.

Lemma 8.2. *If β and γ intersect, then $\lambda^n \geq 14.5$*

Proof. In this case, the growth rate of G is bounded from below by that of the following graph G_1 :




Here, since γ is a curve that passes through petals only, we have $q \leq r_0$, thus $p + 2q \leq p + q + r_0 \leq n$.

Subject to the conditions $m \leq n$ and $p + 2q \leq n$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.5$, where the minimum growth rate is attained near $p \approx 0.70n$, $q \approx 0.15n$. \square

Thus we can assume from this point onwards that β and γ are disjoint.

Lemma 8.3. *If γ is not a petal curve, then $\lambda^n \geq 33$.*

Proof. In this case, the growth rate of G is bounded from below by that of the following graph G_1 :




Subject to the conditions $m \leq n$ and $p + q + r_0 \leq n$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 33$, where the minimum growth rate is attained near $p \approx 0.41n$, $q \approx 0.20n$, $r_0 \approx 0.39n$. \square

Thus we can assume from this point onwards that γ is a petal curve. Since we are assuming that β and γ are disjoint, β must intersect some petal curve other than γ . Since Γ is strongly connected, there is some curve μ intersecting γ and some filament curve or some petal curve other than γ .

Lemma 8.4. *If μ does not pass through filaments, then $\lambda^n \geq 32$.*

Proof. In this case, the growth rate of G is bounded from below by that of the following graph G_1 :




Here the bottom right vertex is the sum of the petal curves other than γ . In particular $q + s = r_0$, thus $p + 2q + s = p + q + r_0 \leq n$.

Subject to the conditions $p + 2q + s \leq n$ and $u \leq q + s$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 32$, where the minimum growth rate is attained near $p \approx 0.32n$, $q \approx 0.08n$, $s \approx 0.52n$. \square


Lemma 8.5. *If μ passes through filaments, then $\lambda^n \geq 14.5$.*

Proof. We first suppose that μ intersects β and petal curves other than γ . In this case, the growth rate of G is bounded from below by that of the following graph G_1 :




Subject to the conditions $m \leq n$, $p + 2q + s \leq n$, and $u \leq m + q + s$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 18.9$, where the minimum growth rate is attained near $p \approx 0.49n$, $q \approx 0.06n$, $s \approx 0.39n$.

Next, we suppose that μ intersects petal curves other than γ , but does not intersect β . In this case, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the conditions $m \leq n$, $p + 2q + s \leq n$, and $p + u \leq m + q + s$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 21$, where the minimum growth rate is attained near $p \approx 0.38n$, $q \approx 0.06n$, $s \approx 0.50n$.

The remaining case is if μ does not intersect petal curves other than γ . In this case, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the conditions $m \leq n$, $p + 2q + s \leq n$, and $s + u \leq m + q + s$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.5$, where the minimum growth rate is attained near $p \approx 0.59n$, $q \approx 0.07n$, $s \approx 0.27n$. \square

9. CASE III: ONE OF β AND γ PASSES THROUGH FILAMENTS ONLY WHILE THE OTHER PASSES THROUGH PETALS ONLY AND IS NOT PERSISTENT

The goal of this section is to prove the following proposition. As in cases I and II, the assumptions in the proposition are more general than the stated hypothesis in this case.

Proposition 9.1. *Let $g : \tau \rightarrow \tau$ be a floral train track that carries f . Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β passes through filaments only and is not a filament curve, and
- γ passes through petals only and is not persistent.


Then $\lambda^n \geq 14.8$.

Up to replacing β with a curve that it contains, we can assume that it is embedded. Similarly, up to replacing γ with a curve that it contains, we can assume that it is embedded. We denote the lengths of β and γ by p and q respectively.

We fix some system of petal curves that does not contain γ . Let r be the sum of lengths of the petal curves.

Lemma 9.2. *If γ intersects at least two petal curves, then $\lambda^n \geq 33$.*

Proof. In this case, the growth rate of G is bounded from below by that of the following graph G_1 :




where s and t are lengths of two petal curves which γ intersect.

Subject to the conditions $m \leq n$ and $p + q + s + t \leq n$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 33$ where the minimum growth rate is attained near $p \approx 0.21n$, $q \approx 0.40n$, $s = t \approx 0.20n$. \square

Thus we can assume from this point onwards that γ only intersects one petal curve δ . Since $\gamma \neq \delta$, there is an edge ε of γ at which γ exits δ . By Proposition 5.9, there is another edge ε' exiting δ at the same vertex as ε and which is not a double of ε . We let μ be a curve that passes through ε' . Note that μ intersects γ and δ .

Lemma 9.3. *If μ does not pass through filaments, then $\lambda^n \geq 17$.*


Proof. In this case, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the conditions $m \leq n$, $p + q + r \leq n$, and $u \leq r$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 17$ where the minimum growth rate is attained near $p \approx 0.27n$, $q \approx 0.24n$, $r \approx 0.49n$. \square


Lemma 9.4. *If μ passes through filaments, then $\lambda^n \geq 14.8$.*

Proof. We first suppose that μ intersects β . In this case, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the conditions $m \leq n$, $p + q + r \leq n$, and $u \leq m + r$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 17$, where the minimum growth rate is attained near $p \approx 0.37n$, $q \approx 0.24n$, $r \approx 0.39n$.

Thus we can assume that μ does not intersect β . In this case, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the conditions $m \leq n$, $p + q + r \leq n$, and $p + u \leq m + r$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 14.8$ where the minimum growth rate is attained near $p \approx 0.33n$, $q \approx 0.26n$, $r \approx 0.41n$. \square

10. CASE IV: ONE OF β AND γ PASSES THROUGH FILAMENTS ONLY WHILE THE OTHER PASSES THROUGH PETALS ONLY AND IS PERSISTENT

The goal of this section is to prove the following proposition.

Proposition 10.1. *Define β and γ as in Section 6.2. Suppose*

- *one of β and γ passes through filaments only, while*
- *the other passes through petals only and is persistent.*

Then $\lambda^n \geq 14.8$.

Note that unlike the previous 3 cases, Proposition 10.1 is only stated for the specific β and γ defined in Section 6.2. This is because we will need some specific properties of these curves towards the end of its proof.

For now, we first prove the following proposition, which restricts the possibilities for the petal curves.

Proposition 10.2. *Fix some system of petal curves. Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β passes through filaments only and is not a filament curve, and
- γ passes through petals only and is a petal curve.

Then either $\lambda^n \geq 14.8$, or

- γ is the only petal curve,

- every curve other than γ has to pass through filaments, and
- for every vertex e on γ , if μ_1 and μ_2 are two curves exiting γ through e , then μ_1 and μ_2 intersect a common filament curve.

Up to replacing β with a curve that it contains, we can assume that it is embedded. We denote the lengths of β and γ by p and q respectively. We will also denote the sum of the lengths of the petal curves other than γ by s .

By case III ([Proposition 9.1](#)), if there is a curve that passes through petals only and intersects γ , then we are done. Thus we can assume that such a curve does not exist.

On the other hand, since Γ is strongly connected, there must be an edge ε exiting γ . By [Proposition 5.9](#), there is another edge exiting γ at the same vertex as ε and which is not a double of ε . Let μ and ν be two curves passing through each of the two edges. By our assumption, μ and ν must pass through filaments. Thus they have doubles μ' and ν' . By construction, μ, μ', ν, ν' intersect each other.

Lemma 10.3. *If μ and ν meet some petal curve other than γ , then $\lambda^n \geq 15.8$.*

Proof. If μ and ν both meet β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, and $u, v \leq n + q + s$, we compute that $\lambda^n \geq 17.4$, where the minimum growth rate is attained near $p \approx 0.43n$, $q \approx 0.12n$, $s \approx 0.33n$.

If μ meets β but ν does not, then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq n + q + s$, and $p + v \leq n + q + s$, we compute that $\lambda^n \geq 17.5$, where the minimum growth rate is attained near $p \approx 0.36n$, $q \approx 0.14n$, $s \approx 0.36n$.

If μ and ν both do not meet β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, and $p + u, p + v \leq n + q + s$, we compute that $\lambda^n \geq 15.8$, where the minimum growth rate is attained near $p \approx 0.32n$, $q \approx 0.14n$, $s \approx 0.40n$. \square

Lemma 10.4. *If μ meets some petal curve other than γ , but ν does not, then $\lambda^n \geq 16.8$.*

Proof. If μ and ν both meet β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq n + q + s$, and $v \leq n + q$, we compute that $\lambda^n \geq 17.6$, where the minimum growth rate is attained near $p \approx 0.46n$, $q \approx 0.17n$, $s \approx 0.20n$.

If μ meets β but ν does not, then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq n + q + s$, and $p + v \leq n + q$, we compute that $\lambda^n \geq 18.6$, where the minimum growth rate is attained near $p \approx 0.37n$, $q \approx 0.21n$, $s \approx 0.21n$.

If ν meets β but μ does not, then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $p + u \leq n + q + s$, and $v \neq n + q$, we compute that $\lambda^n \geq 18.5$, where the minimum growth rate is attained near $p \approx 0.36n$, $q \approx 0.21n$, $s \approx 0.22n$.

If μ and ν both do not meet γ , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $p + u \leq n + q + s$, and $p + v \neq n + q$, we compute that $\lambda^n \geq 16.8$, where the minimum growth rate is attained near $p \approx 0.32n$, $q \approx 0.23n$, $s \approx 0.22n$. \square

Similarly, if ν intersects some petal curve other than γ , but μ does not, then $\lambda^n \geq 16.8$. Thus we can assume from this point that both μ and ν do not intersect petal curves other than γ .

Next, we rule out there being a petal curve other than γ . Suppose otherwise, then since Γ is strongly connected, there must be a curve ω intersecting such a petal curve δ and some filament curve or γ . Let ε be an edge where ω exits δ . There must be another edge exiting δ at the same vertex as ε , and which is not a double of ε . Let χ be a curve passing through such an edge.

We can assume that ω and χ do not intersect γ , since otherwise we could have chosen μ to be such a curve, in which case we are done by [Lemma 10.3](#) and [Lemma 10.4](#). In particular we can assume that ω passes through filaments.

Lemma 10.5. *If χ does not pass through filaments, then $\lambda^n \geq 15.7$.*

Proof. If ω intersects β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq m + q$, $w \neq n + s$, and $x \leq s$, we compute that $\lambda^n \geq 16$, where the minimum growth rate is attained near $p \approx 0.41n$, $q \approx 0.12n$, $s \approx 0.35n$.

If ω does not intersect β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq m + q$, $p + w \neq n + s$, and $x \leq s$, we compute that $\lambda^n \geq 15.7$, where the minimum growth rate is attained near $p \approx 0.36n$, $q \approx 0.13n$, $s \approx 0.38n$. \square

Lemma 10.6. *If χ passes through filaments, then $\lambda^n \geq 14.8$.*

Proof. If ω and χ both intersect β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq m + q$, and $w \leq m + s$, we compute that $\lambda^n \geq 14.8$, where the minimum growth rate is attained near $p \approx 0.48n$, $q \approx 0.13n$, $s \approx 0.26n$.

If μ and ν both intersect β , then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $u \leq m + q$, and $w \leq m + s$, we compute that $\lambda^n \geq 16.2$, where the minimum growth rate is attained near $p \approx 0.49n$, $q \approx 0.18n$, $s \approx 0.15n$.

If at least one of μ and ν , say μ , does not intersect β , and at least one of ω and χ , say ω , does not intersect β then using the inequalities $m \leq n$, $p + 2q + s \leq n$, $p + u \leq m + q$, $v \leq m + q$, $p + w \leq m + s$, and $x \leq m + s$, we compute that $\lambda^n \geq 16.6$, where the minimum growth rate is attained near $p \approx 0.35n$, $q \approx 0.20n$, $s \approx 0.25n$. \square

Thus we can assume from this point that γ is the only petal curve.

Next, we argue that we can assume μ and ν passes through a common filament curve.

Lemma 10.7. *If the collection of filament curves passed through by μ is disjoint from that of ν , then $\lambda^n \geq 19.5$.*

Proof. Without loss of generality, we can assume that β intersects a filament curve that is disjoint from ν . Let m_1 be the sum of lengths of filament curves that are disjoint from ν , and let m_2 be the sum of the remaining filament curves. Using the inequalities $m_1 + m_2 \leq n$, $p + 2q \leq n$, $u \leq m_1 + q$, and $v \leq m_2 + q$, we compute that $\lambda^n \geq 19.5$, where the minimum growth rate is attained near $m_1 \approx 0.61n$, $m_2 \approx 0.39n$, $p \approx 0.28n$, $q \approx 0.36n$. \square

This concludes the proof of [Proposition 10.2](#). For the rest of this section, we specialize to the canonical system of petal curves given by [Proposition 5.4\(2\)](#). In particular, the sum of the lengths of the petal curves is r_0 . To show [Proposition 10.1](#), we divide into two cases depending on whether β or γ is the curve that passes through filaments only.

10.1. Case IVa: γ is the curve that passes through filaments only. By [Proposition 4.15](#), there are at least two edges ε_1 and ε_2 entering γ in this case. We claim that we can extract curves μ, ν, ω, χ such that:

- each of these curves pass through both filaments and petals, in particular they each have doubles,
- these curves and their doubles are distinct, and
- μ intersects ν and ω intersects χ .

Indeed, we first pick a curve μ passing through ε_1 and a curve ω passing through ε_2 . If μ and ω exit γ from different vertices, then we can construct ν and χ by following the other edge exiting γ at the respective vertex. Otherwise by [Proposition 10.2](#), μ and ω pass through a common filament curve α . We can construct ν by traversing ε_2 , then exit γ by following μ until we reach α , then start following ω until we close up at ε_2 . We can construct χ symmetrically.

Proof of [Proposition 10.1](#) in case IVa. If at least one of μ and ν is disjoint from at least one of ω and χ , say μ and ω are disjoint, then using the inequalities $m \leq n$, $p + 2q \leq n$, $u + w \leq m + q$, and $v, x \leq m + q$, we compute that $\lambda^n \geq 20$, where the minimum growth rate is attained near $p \approx 0.33n$, $q \approx 0.34n$.

Thus we can assume that μ, ν, ω, χ intersect each other.

If all four curves intersect β , then using the inequalities $m \leq n$, $p + 2q \leq n$, and $u, v, w, x \leq n + q$, we compute that $\lambda^n \geq 17$, where the minimum growth rate is attained near $p \approx 0.24n$, $q \approx 0.38n$.

By similar computations:

- If exactly three of the curves intersect β , then $\lambda^n \geq 18.4$, where the minimum is attained near $p \approx 0.28n$, $q \approx 0.36n$.

- If exactly two of the curves intersect β , then $\lambda^n \geq 18.8$, where the minimum is attained near $p \approx 0.30n$, $q \approx 0.35n$.
- If exactly one of the curves intersects β , then $\lambda^n \geq 18.5$, where the minimum is attained near $p \approx 0.32n$, $q \approx 0.34n$.
- If all of the curves do not intersect β , then $\lambda^n \geq 17.2$, where the minimum is attained near $p \approx 0.34n$, $q \approx 0.33n$.

□

10.2. **Case IVb: β is the curve that passes through filaments only.** By [Equation \(6.2\)](#), we have $p = q = r_0$ in this case. By [Proposition 5.10](#), either

- (1) there are two other edges that exit γ , or
- (2) the edges from which μ and ν exit γ are the only edges that exit γ , in which case they have the same terminal vertex f , and there is a third edge ε entering f .

In case (i), we can construct four curves and reason as in [Section 10.1](#). Thus we assume that we are in case (ii). In this case, we can choose ν to pass through the same set of vertices as μ . We let ω be a curve passing through ε . In particular, ω intersects μ and ν .

Proof of [Proposition 10.1](#) in case IVb. If μ and ν intersect β , then subject to the conditions $m \leq n$, $3p \leq n$, and $u \leq n + p$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 16.6$.

Thus we can assume that μ and ν do not intersect β .

If ω intersects β and γ , then subject to the conditions $m \leq n$, $3p \leq n$, $p + u \leq n + p$, and $w \leq n + p$, we compute that $\lambda^n \geq 15.7$.

By similar computations,

- if ω intersects β but not γ , then $\lambda^n \geq 18.6$,
- if ω intersects γ but not β , then $\lambda^n \geq 15.2$,
- if ω does not intersect β and γ , then $\lambda^n \geq 16.1$.

□

11. CASE V: BOTH β AND γ PASS THROUGH PETALS ONLY, AND AT LEAST ONE OF THEM IS NOT PERSISTENT

The goal of this section is to prove the following proposition.

Proposition 11.1. *Define β and γ as in [Section 6.2](#). Suppose*

- *both β and γ pass through petals only, and*

- at least one of β and γ is not persistent.

Then $\lambda^n \geq 14.5$.

We first show that we are done if we can find a system of petal curves that does not contain both β and γ .

Proposition 11.2. *Fix some system of petal curves. Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$, and
- β and γ pass through petals only and are not petal curves.

Then $\lambda^n \geq 16$.

Proof. By [Lemma 7.3](#), we can assume that each of β and γ contains at most two curves. Under this assumption, there are three possibilities for β :


- (1) β contains one non-petal curve,
- (2) β contains two non-petal curves, or
- (3) β contains one non-petal curve and one petal curve,

and similarly for γ .

If both β and γ satisfy (1), then they are embedded. If at least one of β and γ satisfy (2), then up to replacing β and γ by non-petal curves that they contain, we can assume that β and γ are embedded.

If β satisfies (3), then the same statement holds unless the non-petal curve that β contains is also contained in γ . But in that case we have $\text{length}(\beta) \leq \text{length}(\gamma) + r_0$, which implies that $\text{length}(\beta) \leq \frac{n}{2}$. By [Lemma 7.2](#), we would then have $\lambda^n \geq 16$. Similarly, if γ satisfies (3), then we can assume that β and γ are embedded.

Under this assumption, the growth rate of G is bounded from below by that of the following graph G_1 :



Subject to the condition $p + q + r \leq n$, we compute that $\lambda^n \geq \lambda(G_1, w_1)^n \geq 16$ where the minimum growth rate is attained when $p = q = \frac{1}{4}$, $r = \frac{1}{2}$. □

Unlike case IV, β and γ will play a symmetric role in this case. Hence without loss of generality let us assume that γ is not persistent. Pick some system of petal curves that does not contain γ .

We first argue that we can assume γ only intersects one petal curve. Suppose otherwise that γ intersects petal curves δ_1 and δ_2 , then the cycle obtained by composing $\gamma, \delta_1, \delta_2$ has length $q + r \leq n$, so by [Lemma 7.3](#) we have $\lambda^n \geq 16$.

If the only petal curve that γ intersects is β , we argue that we can further assume γ passes through the vertices of β only. This is automatically true if every petal lies in one petal curve in our chosen system, for example if our system is the canonical one given by [Proposition 5.4\(2\)](#). So let us suppose that γ is contained in every system for which every petal lies in one petal curve.

On the other hand, by [Lemma 5.6](#), β is also non-persistent. The same reasoning as above shows that we can assume β is contained in every system for which every petal lies in one petal curve. But since β and γ intersect, we run into a contradiction.

Accordingly, we divide the rest of this section into two cases:

- (Va) γ passes through the vertices of β only.
- (Vb) γ intersects a petal curve $\delta \neq \beta$ and no other petal curve.

11.1. Case Va: γ passes through the vertices of β only. We first prove the following proposition, which restricts the possibilities for the remaining petal curves.

Proposition 11.3. *Fix some system of petal curves. Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β and γ pass through petals only,
- β is a petal curve, and
- γ is not a petal curve.

Then either $\lambda^n \geq 15.1$ or

- β is the only petal curve, and
- every curve other than β and γ passes through filaments.

Up to replacing γ with a curve that it contains, we can assume that it is embedded. We denote the lengths of β and γ by p and q respectively. Throughout this subsection, we will denote the sum of the length of the petal curves other than β by s .

If there is a curve $\mu \neq \gamma$ that passes through petals only and intersects β , then the curve obtained by composing β, γ, μ has length $p + q + r \leq n$, so by [Lemma 7.3](#) we have $\lambda^n \geq 16$. Thus we can assume from this point that any curve other than γ that intersects β must pass through filaments.

Since Γ is strongly connected, there must be a curve μ intersecting β and some other petal curve or filament curve. In fact, by our assumption, we know that μ must intersect some filament curve.

Lemma 11.4. *If μ intersects some petal curve other than β , then $\lambda^n \geq 17$.*

Proof. If μ intersects γ , then subject to the conditions $m \leq n$, $2p + q + s \leq n$, and $u \leq m + p + s$, we compute that $\lambda^n \geq 16$ where the minimum growth rate is attained near $p \approx 0.22n$, $q \approx 0.39n$, $s \approx 0.17n$.

If μ does not intersect γ , then we compute that $\lambda^n \geq 18$ where the minimum growth rate is attained near $p \approx 0.28n$, $q \approx 0.27n$, $s \approx 0.17n$. \square

Thus we can assume from this point that μ does not intersect any petal curves other than β .

Lemma 11.5. *If there is a petal curve other than β , then $\lambda^n \geq 15.1$.*

Proof. Since Γ is strongly connected, there is a curve ν intersecting a petal curve other than β , and β or some filament curve.

We can assume that ν does not pass through β , since otherwise we could have taken μ to be ν and be done by [Lemma 11.4](#). Thus ν passes through the filaments.

We separate into multiple cases depending on whether μ , ν , and γ intersect. By similar computations as before, we compute that $\lambda^n \geq 15.1$, with the minimum attained when μ intersects γ and ν , and near $p = q \approx 0.30n$. \square

This concludes the proof of [Proposition 11.3](#). With it, we can now prove [Proposition 11.1](#) in this subcase.

Proof of [Proposition 11.1](#) in case Va. By [Proposition 11.3](#), we can assume that β is the only petal curve and every curve other than β and γ passes through filaments.

Recall [Lemma 6.3](#). First suppose that b' is nonsingular. Let β' be a cycle determined by b' . The length of β' is $\leq p$, so β' cannot contain β . In particular β' must contain some curve other than β and γ . We replace β' with this curve.

By assumption, β' has to pass through filaments. Hence we can conclude in this setting by applying case II ([Proposition 8.1](#)) or case III ([Proposition 9.1](#)) to β' and γ .

A similar reasoning, now using the fact that $q \leq p$ to show that γ' cannot contain β , and using [Proposition 10.2](#), shows that if c' is nonsingular then we are done. Hence we can assume that both b' and c' are singular. Let β' and γ' be cycles determined by b' and c' which pass through petals. These exist by definition of petals. Let p' and q' be the lengths of β' and γ' respectively. [Lemma 6.4\(2a\)](#) states that $p + q + \frac{1}{3}p' + \frac{1}{3}q' \leq n$.

Note that β' and γ' must pass through filaments, since each of them either contains a curve other than β and γ , or is a concatenation of nontrivial powers of β and γ . But the latter case cannot occur by [Corollary 4.17](#), since either b or c lies in \mathcal{X}_O .

Thus β' and γ' must each contain a curve that passes through both filaments and petals. By replacing β' and γ' by these curves, we can assume that they are embedded.

Subject to the conditions $m \leq n$ and $p+q+\frac{1}{3}p'+\frac{1}{3}q' \leq n$, we compute that $\lambda^n \geq 29$ where the minimum growth rate is attained when $p \approx 0.37n$, $q \approx 0.21n$, $p' = q' \approx 0.63n$. \square

11.2. Case Vb: γ intersects a petal curve $\delta \neq \beta$ and no other petal curve. We first prove the following proposition, which restricts the possibilities for the remaining petal curves.

Proposition 11.6. *Fix some system of petal curves. Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- $\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,
- β and γ passes through petals only,
- β is a petal curve,
- γ is not a petal curve, and
- γ intersects a petal curve $\delta \neq \beta$ and no other petal curve.

Then either $\lambda^n \geq 14.5$ or

- β and δ are the only petal curves, and
- every curve other than β , γ , and δ passes through filaments.

Up to replacing γ with a curve that it contains, we can assume that it is embedded. We denote the lengths of β and γ by p and q respectively. Throughout this subsection, we will denote the length of δ by s .

We first rule out certain configurations of curves that pass through petals only.

If there is a curve μ that passes through petals only and which intersects β , then β is non-persistent. By applying the reasoning below [Proposition 11.2](#) (with the roles of β and γ switched), we reduce to case Va ([Proposition 11.3](#)). Thus we can assume from this point that every curve that passes through petals only does not intersect β .

Lemma 11.7. *If there is a curve μ that passes through petals only and which intersects γ , then $\lambda^n \geq 14.5$.*

Proof. Recall that by our assumption, μ does not intersect β . Since Γ is strongly connected, there must be a curve ν that intersects some petal curve other than β , and β or some filament curve. Again by our assumption, ν must pass through filaments.

We first tackle the case where ν intersects β . We separate into multiple cases depending on whether ν intersects γ and μ . We compute that $\lambda^n \geq 14.5$, with the minimum attained when ν intersects μ but not γ , and near $p \approx 0.05n$, $q \approx 0.18n$, $s \approx 0.72n$.

We then tackle the case where ν does not intersect β . Since Γ is strongly connected, there is a curve ω that intersects β and some petal curve other than β or some filament

curve. By assumption, ω passes through filaments. By the previous case, we can assume that ω does not intersect petal curves other than β . We can also assume that ν and ω intersect, since otherwise in our computations, we can sum up their vertices to reduce to the previous case as well.

We separate into multiple cases depending on whether ν intersects γ and μ . In all cases except the one where ν intersects μ but not γ , we have $\lambda^n \geq 14.5$, with the minimum attained when ν intersects both γ and μ , and near $p \approx 0.08n$, $q \approx 0.26$, $s \approx 0.58n$.

In the remaining case where ν intersects μ but not γ , we have to work harder. By [Proposition 5.9](#), there is a curve χ that exits β from the same vertex as ω . By assumption, χ passes through filaments, and by the reasoning above for ω , we can assume that χ does not intersect petal curves other than β . In this case, we compute that $\lambda^n \geq 17.4$, with the minimum growth rate is attained near $p \approx 0.10n$, $q \approx 0.24n$, $s \approx 0.66n$. \square

Thus we can assume from this point that every curve that passes through petals only does not intersect γ .

We now focus on showing that there are no petal curves other than β and δ .

Lemma 11.8. *If there is a curve μ that passes through petals only and which intersects δ and a petal curve other than β and δ , then $\lambda^n \geq 18.4$.*

Proof. Since Γ is strongly connected, there must be a curve ν that intersects some petal curve other than β , and β or some filament curve. By assumption, ν passes through filaments.

We first tackle the case where ν intersects β . We separate into multiple cases depending on whether ν intersects γ , δ , and other petal curves. We compute that $\lambda^n \geq 18.4$, with the minimum attained when ν intersects δ but not other petal curves, and near $p \approx 0.03n$, $q \approx 0.11n$, $s \approx 0.70n$.

We then tackle the case where ν does not intersect β . Since Γ is strongly connected, there is a curve ω that connects β to some other petal curve or the filaments. By assumption, ω passes through filaments. By the previous case, we can assume that ω does not intersect petal curves other than β . We can also assume that ν and ω intersect, since otherwise in our computations, we can sum up their vertices to reduce to the previous case as well.

We separate into multiple cases depending on whether ν intersects γ , δ , and other petal curves. We compute that $\lambda^n \geq 18.5$, with the minimum attained when ν intersects δ but not other petal curves and γ , and near $p \approx 0.05n$, $q \approx 0.12n$, $s \approx 0.64n$. \square

Thus we can assume from this point that every curve that passes through petals only and which intersects a petal curve other than β and δ does not intersect δ (nor, as previously assumed, β).

By [Proposition 5.9](#), there is a curve μ that exits δ from the same vertex as γ . By assumption, μ passes through filaments.

Lemma 11.9. *If μ passes through a petal curve other than β and δ , then $\lambda^n \geq 15$.*

Proof. If μ passes through β , then we compute that $\lambda^n \geq 17.8$, with the minimum growth rate attained near $p \approx 0.08n$, $q \approx 0.24n$, $s \approx 0.43n$.

Thus we can assume that μ does not pass through β . Since Γ is strongly connected, there is a curve ν that intersects β and some other petal curves or some filament curve. By assumption, ν passes through filaments. We may assume that μ and ν intersect, since otherwise in our computations, we can sum up their vertices to reduce to the previous case.

We separate into multiple cases depending on whether ν intersects γ , δ , and other petal curves. We compute that $\lambda^n \geq 15$, with the minimum attained when ν intersects δ but not other petal curves, and does not intersect γ , and near $p \approx 0.06n$, $q \approx 0.18$, $s \approx 0.60n$. \square

Lemma 11.10. *If μ passes through β and there is a petal curve other than β and δ , then $\lambda^n \geq 15$.*

Proof. By [Lemma 11.9](#), we can assume that μ does not pass through any petal curves other than β and δ . Since Γ is strongly connected, there is a curve ν intersecting a petal curve other than β and δ , and β or δ or some filament curve. By assumption, ν passes through filaments.

We separate into multiple cases depending on whether ν intersects β , γ , and δ . We compute that $\lambda^n \geq 15.4$, with the minimum attained when ν intersects δ but not β and γ , and near $p \approx 0.07n$, $q \approx 0.20$, $s \approx 0.59n$. \square

Lemma 11.11. *If μ does not pass through a petal curve other than δ and there is a petal curve other than β and δ , then $\lambda^n \geq 15$.*

Proof. Since Γ is strongly connected, there is a curve intersecting β and some other petal curve or some filament curve, and there is a curve intersecting a petal curve other than β and δ , and β or δ or some filament curve.

We first suppose that there is a single curve ν intersecting β , a petal curve other than β and δ , and δ or some filament curve.

By assumption, ν passes through the filaments. We separate into multiple cases depending on whether ν intersects γ and δ . We compute that $\lambda^n \geq 15.7$, with the minimum attained when ν does not intersect γ and δ , and near $p \approx 0.09n$, $q \approx 0.24$, $s \approx 0.37n$.

We then suppose that there a curve ν intersecting β , and δ or some filament curve, and there is a curve ω intersecting a petal curve other than β and δ , and δ or some filament curve.

By assumption, ν and ω pass through filaments. We separate into multiple cases depending on whether ν and ω intersects γ and δ . In all cases except the one where both ν and ω intersect δ but not γ , we have $\lambda^n \geq 15.4$, with the minimum attained when ν and ω do not intersect γ and δ , and near $p \approx 0.10n$, $q \approx 0.25$, $s \approx 0.40n$.

In the remaining case where both ν and ω intersect δ and do not intersect γ , we have to work harder. By [Proposition 5.9](#), there is a curve χ exiting β from the same vertex as ν and a curve v exiting a petal curve other than β and δ from the same vertex as ω . By previous computations, we can assume that both χ and v intersect δ and both do not intersect γ .

In this case, we have $\lambda^n \geq 17.4$, with the minimum attained near $p \approx 0.05n$, $q \approx 0.12n$, and $s \approx 0.70n$. \square

Thus we can assume from this point that there are no petal curves other than β and δ . To show [Proposition 11.6](#), it remains to show that any curve other than β , γ , and δ passes through filaments.

Lemma 11.12. *If there is a curve ν other than β , γ , and δ that passes through petals only, then $\lambda^n \geq 14.8$.*

Proof. By assumption, ν intersects δ but not β and γ .

Recall the curve μ . If μ intersects β but not ν , then $\lambda^n \geq 17.8$, with the minimum attained near $p \approx 0.08n$, $q \approx 0.24n$, $s \approx 0.60n$, $v \approx 0.31n$.

If μ intersects β and ν , let ω be a curve exiting β at the same vertex as μ . Then $\lambda^n \geq 16$, where the minimum is attained when ω intersects δ and ν but not γ , and near $p \approx 0.07n$, $q \approx 0.16n$, $s \approx 0.70n$, $v \approx 0.54n$.

If μ does not intersect β and does not intersect ν , let ω be a curve intersecting β and some filament curve. Then $\lambda^n \geq 15$, with the minimum attained when ω intersects δ and ν but not γ , and near $p \approx 0.05n$, $q \approx 0.17n$, $s \approx 0.54n$, $v \approx 0.37n$.

If μ does not intersect β but intersects ν , let ω and χ be curves intersecting β and some filaments curves, which exit β at the same vertex. Then $\lambda^n \geq 14.8$, with the minimum attained when ω and χ intersects δ and ν but not γ , and near $p \approx 0.06n$, $q \approx 0.12n$, $s \approx 0.76n$, $v \approx 0.64n$. \square

This concludes the proof of [Proposition 11.6](#). With it, we can now prove [Proposition 11.1](#) in this subcase.

Proof of Proposition 11.1 in case Vb. By Proposition 11.6, we can assume that β and δ are the only petal curve and every curve other than β , γ , and δ passes through filaments.

Recall Lemma 6.3. First suppose that b' is nonsingular. Let β' be a cycle determined by b' . The length of β' is $\leq p$, so β' cannot contain β . If β' contains δ , then by Proposition 11.3, we have $\lambda^n \geq 15.1$. Hence we can assume that β' contains some curve other than β , γ , and δ . We replace β' with this curve. By assumption, β' passes through filaments, so we can conclude by applying case II (Proposition 8.1) to β' and γ .

Hence we can assume that b' is singular. Let β' be a cycle determined by b' which passes through petals. Let p' be the length of β' .

We first suppose that β' passes through filaments. By picking $\kappa = 1$ in Lemma 6.4(1b), we get

$$(11.1) \quad p + q + s + \frac{1}{3}p' = q + (p + s) + \frac{1}{3}p' \leq n.$$

By picking $\kappa = 0$, we get

$$(11.2) \quad q + s + \frac{2}{3}p' = -p + q + (p + s) + \frac{2}{3}p' \leq n.$$

Up to replacing β' by a curve that it contains, we can assume that it is embedded. If β' intersects both β and δ , then using the constraint Equation (11.1), we compute that $\lambda^n \geq 22$ where the minimum is attained when μ does not intersect γ , and near $p \approx 0.09n$, $q \approx 0.23n$, $s \approx 0.30n$.

If β' intersects δ but not β , then using the constraint Equation (11.2), we compute that $\lambda^n \geq 20$ where the minimum is attained when μ does not intersect γ , and near $q \approx 0.22n$, $s \approx 0.41n$.

If β' intersects β but not δ , we let ν be a curve exiting δ at the same vertex as γ . If ν intersects β , then using the constraint Equation (11.1), we compute that $\lambda^n \geq 19.2$ where the minimum is attained near $p \approx 0.19n$, $q \approx 0.23n$, $s \approx 0.34n$.

If ν does not intersect β , then using the constraint Equation (11.1), we compute that $\lambda^n \geq 16.9$ where the minimum is attained near $p \approx 0.10n$, $q \approx 0.24n$, $s \approx 0.35n$.

Hence we can assume that β' does not pass through filaments. In this case, β' must be a concatenation of nontrivial powers of γ and δ . In fact, by Proposition 4.16 it could only be the concatenation of one copy of γ and one copy of δ . Thus $p' = q + s$.

We turn our attention to c' . First suppose that c' is nonsingular. Let γ' be a cycle determined by c' . If γ' passes through filaments, we can conclude by applying case II (Proposition 8.1) to β and γ' . Hence we can assume that γ' passes through petals only. Since $\text{length}(\gamma') \leq q$, we must have $\gamma' = \delta$.

By picking $\kappa = \frac{4}{5}$ in [Lemma 6.4\(1b\)](#), we get

$$(11.3) \quad \frac{4}{5}p + \frac{7}{5}q + \frac{7}{5}s = -\frac{1}{5}p + q + (p + s) + \frac{2}{5}(q + s) \leq n.$$

Let μ be a curve that exits δ at the same vertex as γ . If μ intersects β , let ν be a curve that exits β at the same vertex as μ . Using the constraint [Equation \(11.3\)](#), we compute that $\lambda^n \geq 14.7$, where the minimum is attained when ν does not meet γ or δ , and near $p \approx 0.21n$, $q \approx 0.34n$, $s \approx 0.26n$.

If μ does not intersect β , there exists curves ν and ω that intersect β , and other petal curves or filament curves. Using the constraint [Equation \(11.3\)](#), we compute that $\lambda^n \geq 14.5$, where the minimum is attained when ν and ω do not meet γ or δ , and near $p \approx 0.18n$, $q \approx 0.35n$, $s \approx 0.26n$.

Thus we can assume that c' is singular. Let γ' be a cycle determined by c' which passes through petals. Let q' be the length of γ' . If γ' does not pass through filaments, then by the same reasoning we used for β' , γ' must be the concatenation of one copy of γ and one copy of δ . But then this contradicts [Proposition 4.16](#). Hence we can assume that γ' passes through filaments.

Recall that $p' = q + s$. By picking $\kappa_1 = \frac{1}{2}$ and $\kappa_2 = \frac{5}{4}$ in [Lemma 6.4\(2b\)](#), we get

$$(11.4) \quad \frac{1}{2}p + \frac{3}{4}q + \frac{3}{2}s + \frac{1}{4}q' = -\frac{1}{2}p + \frac{1}{4}q + (p + s) + \frac{1}{2}(q + s) + \frac{1}{4}q' \leq n.$$

By picking $\kappa = 0$ in [Lemma 6.4\(1b\)](#), we get

$$(11.5) \quad q + s + \frac{2}{3}p' = -p + q + (p + s) + \frac{2}{3}p' \leq n.$$

Moreover, we claim that we can assume $q \leq s$. This is true if γ passes through the vertices in δ only, which is in turn automatically true if every petal lies in one petal curve in our chosen system. So let us suppose that γ is contained in every system for which every petal lies in one petal curve.

Meanwhile, if δ passes through the vertices in γ only, then we can swap the roles of γ and δ . So let us suppose that δ contains vertices that lie outside of γ as well. Now let us fix a system for which every petal lies in one petal curve. Let μ be a petal curve that contains a vertex in δ but not in γ . This curve μ will contradict our assumption that every curve other than β , γ , and δ passes through filaments.

Armed with these inequalities, we move on to the final computations in this proof.

Let μ be a curve in γ' that passes through both filaments and petals. If μ intersects both β and δ , then using the constraints [Equation \(11.4\)](#) and $q \leq s$, we compute that $\lambda^n \geq 17.9$ where the minimum is attained when μ does not intersect γ , and near $p \approx 0.14n$, $q = s \approx 0.27n$.

If μ intersects δ but not β , then using the constraints [Equation \(11.5\)](#) and $q \leq s$, we compute that $\lambda^n \geq 14.5$ where the minimum is attained when μ does not intersect γ , and near $q = s \approx 0.31n$.

If μ intersects β but not δ , we let ν be a curve exiting δ at the same vertex as γ . If ν intersects β , then using the constraints [Equation \(11.4\)](#) and $q \leq s$, we compute that $\lambda^n \geq 15$ where the minimum is attained near $p \approx 0.36n$, $q = s \approx 0.29n$.

If ν does not intersect β , then using the constraints [Equation \(11.4\)](#) and $q \leq s$, we compute that $\lambda^n \geq 16.9$ where the minimum is attained near $p \approx 0.15n$, $q = s \approx 0.30n$. \square

12. CASE VI: BOTH β AND γ PASS THROUGH PETALS ONLY, AND BOTH ARE PERSISTENT

The goal of this section is to prove the following proposition.

Proposition 12.1. *Define β and γ as in [Section 6.2](#). Suppose*

- *both β and γ pass through petals only, and*
- *both β and γ are persistent.*

Then $\lambda^n \geq 14.5$.

We first prove the following proposition, which restricts the possibilities for the remaining petal curves.

Proposition 12.2. *Fix some system of petal curves. Suppose there are two cycles β and γ in the directed graph Γ satisfying:*

- *$\text{length}(\beta) + \text{length}(\gamma) + r_0 \leq n$,*
- *β and γ passes through petals only, and*
- *β and γ are petal curves.*

Then either $\lambda^n \geq 14.5$, or:

- *There is at most one other petal curve δ .*
- *If there is exactly one other petal curve δ , then every curve other than β , γ , and δ either passes through filaments or only intersects δ and not β and γ .*
- *If there is exactly one other petal curve δ and there is a curve other than β , γ , and δ that passes through petals only, then*
 - *there is a curve intersecting β , δ , but not γ and*
 - *there is a curve intersecting γ , δ , but not β .*

If there are no petal curves other than β and γ , then we are done. So we assume that there is some petal curve δ other than β and γ . In this case, by [Proposition 11.2](#) and [Proposition 11.6](#), we can assume that every curve that intersects β or γ passes through filaments.

Lemma 12.3. *If there is some curve that passes through petals only and intersects multiple petal curves, then $\lambda^n \geq 14.9$.*

Proof. Since Γ is strongly connected, there is a curve μ that intersects β . Let ν be a curve that exits β at the same vertex as μ . By our assumption, μ and ν pass through filaments.

If μ and ν intersect γ , then we have $\lambda^n \geq 17.2$, with the minimum attained when

- μ and ν do not intersect other petal curves, in which case, since Γ is strongly connected, there is a curve ω intersecting a petal curve other than β and γ , and the filaments,
- ω does not meet β and γ , and
- μ , ν , and ω intersect each other,

and near $p = q \approx 0.17n$.

If μ intersects γ while ν does not intersect γ , then we have $\lambda^n \geq 17.5$, with the minimum attained when

- μ and ν do not intersect other petal curves, in which case, since Γ is strongly connected, there is a curve ω intersecting a petal curve other than β and γ , and the filaments,
- ω meets γ but not β , and
- μ , ν , and ω intersect each other,

and near $p \approx 0.14n$, $q \approx 0.18n$.

It remains to analyze the case when μ and ν do not intersect γ . In this case, symmetrically, we may assume that there are two curves ω and χ that intersect γ but not β . We have $\lambda^n \geq 14.9$, with the minimum attained when

- μ , ν , ω , χ all meet petal curves other than β and γ ,
- μ , ν , ω , χ intersect each other,

and near $p = q \approx 0.07n$. □

Thus we can assume from this point that any curve that passes through petals only but is not a petal curve only intersects one petal curve, which must be one other than β and γ .

The first item of [Proposition 12.2](#) follows from the lemma below.

Lemma 12.4. *If there are multiple petal curves other than β and γ , then $\lambda^n \geq 14.7$.*

Proof. Suppose there are two petal curves δ and η other than β and γ . We consider curves intersecting β and γ as in [Lemma 12.3](#).

If both δ and η intersect curves that pass through petals only, then $\lambda^n \geq 22$, with the minimum attained when

- there are two curves that intersect β, δ, η , but not γ ,
- there are two curves that intersect γ, δ, η , but not β , and
- all these curves intersect each other,

and near $p = q \approx 0.04n$, $\text{length}(\delta) = \text{length}(\eta) \approx 0.42n$.

If η intersects curves that pass through petals only while δ does not, then $\lambda^n \geq 15$, with the minimum attained when

- there are two curves that intersect β and η , but not γ and δ ,
- there are two curves that intersect γ and η , but not β and δ ,
- there are two curves that intersect δ and η , but not β and γ , and
- all these six curves intersect each other,

and near $p = q \approx 0.05n$, $\text{length}(\delta) \approx 0.07n$, $\text{length}(\eta) \approx 0.72n$.

If both δ and η do not intersect curves that pass through petals only, then $\lambda^n \geq 14.7$, with the minimum attained when there are two curves intersecting β, γ, δ , and η , and $p = q \approx 0.14n$, $\text{length}(\delta) = \text{length}(\eta) \approx 0.22n$. \square

The second item of [Proposition 12.2](#) follows from our assumption on curves that pass through petals only. We show the third item of [Proposition 12.2](#) below.

Lemma 12.5. *If there is exactly one other petal curve δ and there is a curve η which passes through petals only, then either $\lambda^n \geq 14.6$, or*

- *there is a curve intersecting β, δ , but not γ , and*
- *there is a curve intersecting γ, δ , but not β .*

Proof. By considering curves that intersect β and γ as above, we compute that $\lambda^n \geq 14.6$, with the minimum attained when there are two curves intersecting β, γ , and δ , and near $p = q \approx 0.11n$. \square

This concludes the proof of [Proposition 12.2](#).

For the rest of this section, we specialize to the canonical system of petal curves given by [Proposition 5.4\(2\)](#). To show [Proposition 12.1](#), we will divide into three subcases according to [Lemma 6.3](#):

(VIa) b' and c' are nonsingular.

(VIb) One of b' and c' is nonsingular and the other is singular.

(VIc) b' and c' are singular.

12.1. Case VIa: b' and c' are nonsingular. Let β' be a cycle determined by b' . If β' passes through filaments, then we can conclude by case II ([Proposition 8.1](#)) or [Proposition 10.2](#). If β' is not a petal curve, then we can conclude by [Proposition 11.2](#), [Proposition 11.3](#), or [Proposition 11.6](#). So we can assume that β' is a petal curve.

Similarly, let γ' be a cycle determined by c' . We can assume that γ' is a petal curve. In other words, we have 4 distinct petal curves β , γ , β' , and γ' , so we are done by [Proposition 12.2](#).

12.2. Case VIb: One of b' and c' is nonsingular and the other is singular. Here β and γ will play a symmetric role, so we suppose that b' is nonsingular and c' is singular.

Let β' be a cycle determined by b' . Arguing as in case VIa, we can assume that β' is a petal curve. By [Proposition 12.2](#), we can assume that β' is the only petal curve other than β and γ . Let p' be the length of β' .

Let γ' be a cycle determined by c' which passes through petals. Up to replacing γ' by a curve that it contains, we can assume that γ' is embedded. Let q' be the length of γ' .

Lemma 12.6. *If γ' passes through petals only, then $\lambda^n \geq 14.5$.*

Proof. By [Proposition 12.2](#), we can assume that γ' intersects β' but not β and γ , and that there is a curve μ intersecting β , β' , but not γ , and a curve ν intersecting γ , β' , but not β .

We consider curves that exit β at the same vertex as μ , or exit γ at the same vertex as ν . Using the fact that $p' \leq p$, we compute that $\lambda^n \geq 23$, with the minimum attained when

- there is a curve ω exiting β at the same vertex as μ , exiting γ at the same vertex as ν , and which intersects δ ,
- μ , ν , ω intersect each other,

and near $p = p' \approx 0.28n$, $q \approx 0.08n$. □

Lemma 12.7. *If γ' passes through filaments, then $\lambda^n \geq 15.7$.*

Proof. By choosing $\kappa = \frac{1}{2}$ in [Lemma 6.4\(1b\)](#) (with the roles of b and c reversed), we have

$$(12.1) \quad 2p + \frac{1}{2}q + p' + \frac{1}{2}q' \leq p - \frac{1}{2}q + (p + q + p') + \frac{1}{2}q' \leq n.$$

If γ' intersects at least two of the three petal curves β , γ , and β' , then, using [Equation \(12.1\)](#), we compute that $\lambda^n \geq 17.1$, where the minimum is attained when

- γ' intersects β and γ but not β' , in which case there must be a curve μ intersecting β' and some filament curve, and
- γ' and μ intersect,

and near $p \approx 0.08n$, $q \approx 0.24n$, $p' \approx 0.11n$.

If γ' only intersects one of the three petal curves, then we have to work harder. We consider curves that exit the one petal curve which γ' intersects at the same vertex as γ' , and consider curves that intersect the two remaining petal curves. Using [Equation \(12.1\)](#), we compute that $\lambda^n \geq 15.7$, where the minimum is attained when

- γ' intersects β' ,
- there are two curves μ and ν that intersect β and γ , and
- γ' intersects μ and ν ,

and near $p \approx 0.08n$, $q \approx 0.49n$, $p' \approx 0.21n$. □

12.3. Case VIc: b' and c' are singular. Let β' and γ' be cycles determined by b' and c' which pass through petals. By [Corollary 4.17](#), up to replacing these by curves that they contain, we can assume that they are embedded. Let p' and q' be the lengths of β' and γ' respectively.

Lemma 12.8. *If β' and γ' do not intersect filaments, then $\lambda^n \geq 14.5$.*

Proof. By [Proposition 12.2](#), we can assume that

- there is at most one other petal curve δ , and
- if there is one other petal curve δ , then every curve other than β , γ , and δ either passes through filaments or only intersects δ and not β and γ .

Thus, up to switching β' and γ' , there are two cases:

- (1) β' and γ' are both distinct from δ , or
- (2) β' equals δ while γ' is distinct from δ .

In case (1), we let s be the length of δ . We take $\kappa_1 = \kappa_2 = 1$ in [Lemma 6.4\(2b\)](#) to obtain

$$p + q + s + \frac{1}{3}p' + \frac{1}{3}q' \leq r + \frac{1}{3}p' + \frac{1}{3}q' \leq n.$$

Meanwhile, by the second statement in [Proposition 12.2](#), we can assume that

- there is a curve μ intersecting β , β' , but not γ , and
- there is a curve ν intersecting γ , β' , but not β .

In this case, we compute that $\lambda^n \geq 20$, where the minimum is attained when $\beta', \gamma', \mu, \nu$ intersect each other, and near $p = q \approx 0.09n$, $p' = q' \approx 0.59n$.

In case (2), we take $\kappa_1 = \kappa_2 = 1$ in [Lemma 6.4\(2b\)](#) to obtain

$$p + q + \frac{4}{3}p' + \frac{1}{3}q' \leq r + \frac{1}{3}p' + \frac{1}{3}q' \leq n.$$

We apply the second statement in [Proposition 12.2](#) as in case (1). By considering curves that exit β at the same vertex as μ , or exit γ at the same vertex as ν , using the fact that $q' \leq p'$, we compute that $\lambda^n \geq 15$ where the minimum is attained when

- there is a curve ω exiting β at the same vertex as μ and which intersects β', γ' but not γ , and
- there is a curve ω exiting γ at the same vertex as ν and which intersects β', γ' but not β ,

and $p = q \approx 0.14n$, $p' = q' \approx 0.43n$. □

Lemma 12.9. *If β' does not intersect filaments while γ' does, then $\lambda^n \geq 15.7$.*

Proof. The computations in this case are exactly the same as [Lemma 12.7](#), since in that lemma we did not appeal to any relation between p' and p . □

Lemma 12.10. *If β' and γ' intersect filaments, then $\lambda^n \geq 14.5$.*

Proof. By [Proposition 12.2](#), we can assume that there is at most one other petal curve δ . If δ exists, we let s be its length, otherwise we set $s = 0$. In [Lemma 6.4\(2b\)](#):

- If we take $\kappa_1 = \kappa_2 = \frac{1}{2}$, we have

$$(12.2) \quad \frac{1}{2}p + \frac{1}{2}q + s + \frac{1}{2}p' + \frac{1}{2}q' \leq -\frac{1}{2}p - \frac{1}{2}q + (p + q + s) + \frac{1}{2}p' + \frac{1}{2}q' \leq n.$$

- If we take $\kappa_1 = \frac{1}{2}$ and $\kappa_2 = 1$, we have

$$(12.3) \quad \frac{1}{2}p + q + s + \frac{1}{2}p' + \frac{1}{3}q' \leq -\frac{1}{2}p + (p + q + s) + \frac{1}{2}p' + \frac{1}{3}q' \leq n.$$

- If we take $\kappa_1 = \kappa_2 = 1$, we have

$$(12.4) \quad p + q + s + \frac{1}{3}p' + \frac{1}{3}q' \leq n.$$

We first suppose that β' intersects both β and γ . In this case, using [Equation \(12.2\)](#), we compute that $\lambda^n \geq 17.4$, where the minimum is attained when γ' intersects exactly one of β and γ , say, γ but not β , and does not intersect δ , and near $p \approx 0.23n$, $q \approx 0.47n$, $p' \approx 0.78n$, $q' \approx 0.52n$.

Then we suppose that δ exists and β' intersects δ . In this case, using [Equation \(12.2\)](#), we compute that $\lambda^n \geq 22$, where the minimum is attained when

- γ' intersects exactly one of β and γ , say, γ but not β , in which case there is a curve μ intersecting β and some filament curve,
- μ intersects β but not γ and δ , and
- μ intersects β' and γ' ,

and near $p \approx 0.19n$, $q \approx 0.22n$, $s \approx 0.13n$, $p' \approx 0.72n$, $q' \approx 0.60n$.

Thus we can assume from this point that β' intersects exactly one of β and γ , and does not intersect δ . Similarly, we can assume from this point that γ' intersects exactly one of β and γ , and does not intersect δ .

Suppose β' and γ' both intersect γ . By assumption, β' and γ' do not intersect β . Since Γ is strongly connected, there is some curve μ that intersects β .

If μ intersects γ , then using [Equation \(12.3\)](#), we compute that $\lambda^n \geq 16$, where the minimum is attained when μ does not intersect δ but intersects β' and γ' , and near $p \approx 0.14n$, $q \approx 0.64n$, $p' \approx 0.69n$, $q' \approx 0.78n$.

If μ does not intersect γ , then using [Equation \(12.4\)](#), we compute that $\lambda^n \geq 14.5$, where the minimum is attained when μ does not intersect δ but intersects β' and γ' , and near $p \approx 0.15n$, $q \approx 0.21n$, $p' = q' \approx 0.96n$.

Thus we can assume that at most one of β' and γ' intersect γ . Similarly, we can assume that at most one of β' and γ' intersect β . Up to switching β' and γ' , suppose that the only petal curve which β' intersects is β , and the only petal curve which γ' intersects is γ .

If β' and γ' intersect, then using [Equation \(12.2\)](#), we compute that $\lambda^n \geq 17$, where the minimum is attained near $p = q \approx 0.24n$, $p' = q' \approx 0.26n$. Thus we can assume that β' and γ' are disjoint.

We let μ be a curve that exits β at the same vertex as β' . If μ intersects γ , then using [Equation \(12.4\)](#), we compute that $\lambda^n \geq 14.6$, where the minimum is attained when μ intersects both β' and γ' , and near $p = q \approx 0.23n$, $p' = q' \approx 0.77n$.

If μ does not intersect γ , then we let ν be a curve that exits γ at the same vertex as γ' . Using [Equation \(12.4\)](#), we compute that $\lambda^n \geq 16.3$, where the minimum is attained when μ and ν intersect both β' and γ' and intersect each other, and near $p = q \approx 0.20n$, $p' = q' \approx 0.80n$. \square

13. DISCUSSION AND FURTHER QUESTIONS

13.1. Classification and persistence of petal curves. Let f be a pseudo-Anosov braid. Suppose $g : \tau \rightarrow \tau$ is a floral train track map that carries f . In [Section 5.1](#), we showed that by factorizing g into based folds, we can define a collection of filament curves and petal curves in the directed graph Γ associated to g_*^{real} .

In [Lemma 5.5](#), we showed that the filament curves are the cycles determined by the 1-pronged singularities. On the other hand, it is unclear whether the petal curves admit

such a natural description. In fact, it is not even clear if the petal curves depend on the factorization of g into based folds, or the choice of g itself.

If it is possible to obtain different petal curves by varying the factorization or choice of g , then it would seem that persistent petal curves should be rare. In particular, we ask the following question.

Question 13.1. Are the cycles determined by the primary singularities b and c persistent?

If the answer is ‘no’, then one can significantly reduce the amount of computations in showing [Theorem 1.1](#). For example, Case VI would be made completely redundant.

13.2. Reciprocity. One feature of the standardly embedded train track machinery which played a key role in [\[HT24\]](#) is the fact that the real transition matrix is reciprocal ([Theorem 4.4\(4\)](#)). This aspect of the machinery seems highly under-utilized in this paper. Indeed, the only place where we use reciprocity is in [Lemma 7.10](#).

In general, the minimal growth rate of a graph G is much larger if one only varies over the **reciprocal** weights, i.e. the weights $w : V(G) \rightarrow \mathbb{R}_+$ that make the clique polynomial $Q_{(G,w)}(t)$ reciprocal. However, a big problem here is that certain properties of the growth rate fail to make sense if one restricts to such reciprocal weights. More specifically, with respect to the items in [Proposition 3.2](#):

- (3) If H is an induced subgraph of G and w is a reciprocal weight for G , $w|_H$ may not be a reciprocal weight for H .
- (4) If H is a wide subgraph of G and w is a reciprocal weight for G , w may not be a reciprocal weight for H .
- (5) If w_1 and w_2 are two reciprocal weights, $tw_1 + (1 - t)w_2$ may not be a reciprocal weight.

This makes the minimization problems in our computations extremely difficult.

If one is able to formulate suitable generalizations of these monotonicity and convexity properties within the subset of reciprocal weights, then one should be able to obtain stronger lower bounds or bypass some of the computations,

13.3. The remaining values of δ_n . Recall that δ_n is the minimum dilatation among pseudo-Anosov braids with n strands. We now know the value of δ_n for all n except $n = 10, 12, 14, 18, 22, 26$. An obvious problem is to compute δ_n for these remaining values of n .

Venzke conjectured in [\[Ven08, Conjecture 5.4\]](#) that $\delta_n = \underline{\delta}_n$ for these remaining values of n . Note that this conjecture is false for $n = 10$. This is because one can construct a pseudo-Anosov braid with 10 strands and dilatation $\underline{\delta}_9 < \underline{\delta}_{10}$ by taking the pseudo-Anosov braid $\sigma_{3,5}$ with 9 strands in [\[HK06\]](#) (which is conjugate to the braid ψ_9 in [\[Ven08\]](#)) and puncturing out a fixed point.

We propose revising Venzke’s conjecture in the case $n = 10$ to:

Conjecture 13.2. *The minimum dilatation among pseudo-Anosov braids with 10 strands equals $\underline{\delta}_9 = |x^9 - 2x^5 - 2x^4 + 1|$.*

Theoretically, one should be able to repeat the computations in [LT11] to determine the values of these δ_n . However, the process might be rather tedious for these larger values of n .

One should also be able to improve on the results in this paper by more in-depth analysis of the directed graph Γ . Indeed, in all of our lemmas, except Lemma 7.10 and Lemma 7.14, the fact that

- (1) our polynomials have to be reciprocal, and
- (2) the exponents of our polynomials have to be integers

are not utilized at all. One could attempt to go through our computations, isolate out the lemmas that give the weakest lower bounds for the growth rates, redo the computations but imposing conditions (1) and (2), in order to get (slightly) stronger lower bounds. However, this seems equally, if not more tedious than the computations in [LT11].

13.4. The braids with minimum dilatation. Suppose that one can determine the value of δ_n for all n . The next natural question would then be the following:

Question 13.3. Which pseudo-Anosov braids with n strands have dilatation δ_n ?

We believe that for $n \in \mathbb{N}$, the techniques for answering Question 13.3 are already in place: Theorem 1.1 implies that such a pseudo-Anosov braid is carried by a floral train track, where the directed graph associated to the real transition matrix is of the very specific form in Lemma 7.14. In particular, the map g cannot consist of more than 3 based folds.

Theoretically, one should be able list out all floral train tracks with the correct amount of filaments and for each of them list out all maps with ≤ 3 based folds. For each map on the compiled list, one can then compute its transition matrix, and extract the maps whose transition matrix determines the desired directed graph. This would then be a complete list of pseudo-Anosov braids with dilatation δ_n , thereby answering Question 13.3 for these large values of n .

13.5. Other minimum dilatation problems. Let $\lambda_{g,n}$ be the minimum dilatation among pseudo-Anosov maps on the genus g surface with n punctures. The values of $\lambda_{g,n}$ have previously been only known for finitely many values of (g, n) , namely:

- $\lambda_{0,4} = \underline{\delta}_3$, $\lambda_{0,5} = \lambda_{0,6} = \underline{\delta}_5$, $\lambda_{0,7} = \underline{\delta}_7$, and $\lambda_{0,8} = \underline{\delta}_8$ by [SKL02], [HS07], and [LT11], as remarked in the introduction,
- $\lambda_{1,0} = \lambda_{1,1} = \underline{\delta}_3$ from classical arguments, and

- $\lambda_{2,0} = \delta_{\frac{1}{5}}$ by [CH08].

In this paper, we determine $\lambda_{0,n}$ for all large enough values of n . However, there is still a whole 2-parameter family of values of (g, n) that remain mysterious.

Of these, the 1-parameter family where $n = 0$ has received the most interest. In particular, we would be remiss not to mention the following well-known conjecture:

Conjecture 13.4 (Hironaka, [Hir10, Question 1.12]). *The minimum dilatations $\delta_{g,0}$ on the closed surfaces of genus g grow as*

$$\lim_{g \rightarrow \infty} \delta_{g,0}^g = \mu^2 \approx 2.618$$

where $\mu = \frac{1+\sqrt{5}}{2} \approx 1.618$ is the golden ratio.

A recurring phenomenon for $\lambda_{g,n}$ is the asymptotics

$$(13.1) \quad \log \lambda_{g,n} \asymp \frac{1}{|\chi(S_{g,n})|}$$

along certain 1-parameter families. For example, this is observed on the lines

- $g = 1$ by [Tsa09],
- $n = n_0$ for fixed $n_0 > 0$ by [Yaz20], and
- $n = m_0 g$ for fixed $m_0 > 0$ by [Val12].

Along each of these 1-parameter families, one can ask whether $\lambda_{g,n}^{|\chi(S_{g,n})|}$ converges to any number, and if yes, what the value of that number is.

We caution however, that Equation (13.1) does not hold along all 1-parameter families. For example, on the lines $g = g_0$ for fixed $g_0 \geq 2$, [Tsa09] shows that

$$\log \lambda_{g,n} \asymp \frac{\log |\chi(S_{g,n})|}{|\chi(S_{g,n})|}.$$

REFERENCES

- [BH95] M. Bestvina and M. Handel. Train-tracks for surface homeomorphisms. *Topology*, 34(1):109–140, 1995. doi:10.1016/0040-9383(94)E0009-9.
- [Bir74] Joan S. Birman. *Braids, links, and mapping class groups*. Annals of Mathematics Studies, No. 82. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1974.
- [CH08] Jin-Hwan Cho and Ji-Young Ham. The minimal dilatation of a genus-two surface. *Experiment. Math.*, 17(3):257–267, 2008. URL: <http://projecteuclid.org/euclid.em/1227121381>.
- [FLP12] Albert Fathi, François Laudenbach, and Valentin Poénaru. *Thurston’s work on surfaces*, volume 48 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 2012. Translated from the 1979 French original by Djun M. Kim and Dan Margalit.
- [FM12] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [FRW23] Ethan Farber, Braeden Reinoso, and Luya Wang. Fixed-point-free pseudo-Anosov homeomorphisms, knot floor homology and the cinquefoil. *Geometry and Topology (to appear)*, 2023. arXiv:2203.01402.

- [GS00] Massimiliano Goldwurm and Massimo Santini. Clique polynomials have a unique root of smallest modulus. *Inform. Process. Lett.*, 75(3):127–132, 2000. doi:[10.1016/S0020-0190\(00\)00086-7](https://doi.org/10.1016/S0020-0190(00)00086-7).
- [Hir10] Eriko Hironaka. Small dilatation mapping classes coming from the simplest hyperbolic braid. *Algebr. Geom. Topol.*, 10(4):2041–2060, 2010. doi:[10.2140/agt.2010.10.2041](https://doi.org/10.2140/agt.2010.10.2041).
- [HK06] Eriko Hironaka and Eiko Kin. A family of pseudo-Anosov braids with small dilatation. *Algebr. Geom. Topol.*, 6:699–738, 2006. doi:[10.2140/agt.2006.6.699](https://doi.org/10.2140/agt.2006.6.699).
- [HS07] Ji-Young Ham and Won Taek Song. The minimum dilatation of pseudo-Anosov 5-braids. *Experiment. Math.*, 16(2):167–179, 2007. URL: <http://projecteuclid.org/euclid.em/1204905873>.
- [HT24] Eriko Hironaka and Chi Cheuk Tsang. Standardly embedded train tracks and pseudo-Anosov maps with minimum expansion factor. *Groups, Geometry, and Dynamics (to appear)*, 2024. [arXiv:2210.13418](https://arxiv.org/abs/2210.13418).
- [Iva88] N. V. Ivanov. Coefficients of expansion of pseudo-Anosov homeomorphisms. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 167(Issled. Topol. 6):111–116, 191, 1988. doi:[10.1007/BF01099245](https://doi.org/10.1007/BF01099245).
- [LT11] Erwan Lanneau and Jean-Luc Thiffeault. On the minimum dilatation of braids on punctured discs. *Geom. Dedicata*, 152:165–182, 2011. Supplementary material available online. doi:[10.1007/s10711-010-9551-2](https://doi.org/10.1007/s10711-010-9551-2).
- [McM15] Curtis T. McMullen. Entropy and the clique polynomial. *J. Topol.*, 8(1):184–212, 2015. doi:[10.1112/jtopol/jtu022](https://doi.org/10.1112/jtopol/jtu022).
- [SKL02] Won Taek Song, Ki Hyoung Ko, and Jérôme E. Los. Entropies of braids. volume 11, pages 647–666. 2002. *Knots 2000 Korea*, Vol. 2 (Yongpyong). doi:[10.1142/S021821650200186X](https://doi.org/10.1142/S021821650200186X).
- [Tsa09] Chia-Yen Tsai. The asymptotic behavior of least pseudo-Anosov dilatations. *Geom. Topol.*, 13(4):2253–2278, 2009. doi:[10.2140/gt.2009.13.2253](https://doi.org/10.2140/gt.2009.13.2253).
- [Val12] Aaron D. Valdivia. Sequences of pseudo-Anosov mapping classes and their asymptotic behavior. *New York J. Math.*, 18:609–620, 2012. URL: http://nyjm.albany.edu:8000/j/2012/18_609.html.
- [Ven08] Rupert William Venzke. *Braid Forcing, Hyperbolic Geometry, and Pseudo-Anosov Sequences of Low Entropy*. PhD thesis, California Institute of Technology, 2008. doi:[10.7907/290Y-BY53](https://doi.org/10.7907/290Y-BY53).
- [Yaz20] Mehdi Yazdi. Pseudo-Anosov maps with small stretch factors on punctured surfaces. *Algebr. Geom. Topol.*, 20(4):2095–2128, 2020. doi:[10.2140/agt.2020.20.2095](https://doi.org/10.2140/agt.2020.20.2095).

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, 201 PRESIDENT KENNEDY AVENUE, MONTRÉAL, QC, CANADA H2X 3Y7

Email address: tsang.chi_cheuk@uqam.ca

FACULTÉ DES SCIENCES D’ORSAY, UNIVERSITÉ PARIS-SACLAY, F-91405 ORSAY CEDEX

Email address: xiangzhuo.zeng@etu-upsaclay.fr