

ON THE NUMERICAL RADIUS PARALLELISM AND THE NUMERICAL RADIUS BIRKHOFF ORTHOGONALITY

JIAYE BI, HUAYOU XIE, AND YONGJIN LI*

ABSTRACT. In this paper, we generalize the notions of numerical radius parallelism and numerical radius Birkhoff orthogonality, originally formulated for operators on Hilbert spaces, to operators on normed spaces. We then proceed to demonstrate their fundamental properties. Notably, our findings reveal that numerical radius parallelism lacks transitivity, and numerical radius Birkhoff orthogonality is neither left nor right additive. Additionally, we offer characterizations for both concepts. Furthermore, we establish a connection between numerical radius parallelism and numerical radius Birkhoff orthogonality.

1. INTRODUCTION

Throughout this paper, we assume that X is a normed space over the field $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}\}$ of dimension at least 2, B_X, S_X denote the closed unit ball and the unit sphere of X , respectively, $\mathbb{T} := \{\lambda \in \mathbb{F} : |\lambda| = 1\}$, and $\mathbb{B}(X)$ denote the normed space of all bounded linear operators on X .

The concept of *norm parallelism*, first introduced by Seddik [9], defines an element $x \in X$ as being *norm parallel* to $y \in X$, denoted as $x \parallel y$, if there exists $\lambda \in \mathbb{T}$ such that the equality $\|x + \lambda y\| = \|x\| + \|y\|$ is satisfied. This notion generalizes the concept of linear dependence, as elements that are linearly dependent are necessarily norm parallel. Conversely, the implication holds true only in the context of strictly convex spaces, as established in [5, Theorem 2.8]. Zamani and Moslehian [13, 14] contributed significantly to the characterizations of the norm parallelism in Hilbert C^* -modules. In particular, they established a relation between norm parallelism and Birkhoff orthogonality [1]. Recall that a vector x is said to be Birkhoff orthogonal to another vector y , denoted by $x \perp_B y$, if and only if $\|x + \alpha y\| \geq \|x\|$ holds for all $\alpha \in \mathbb{R}$.

Theorem 1. [14, Theorem 2.4] Let X be a normed space. For any $x, y \in X$, there exists $\lambda \in \mathbb{T}$ such that the following statements are equivalent:

- (i) $x \parallel y$.

2020 *Mathematics Subject Classification.* 46B20.

Key words and phrases. numerical radius parallelism; numerical radius Birkhoff orthogonality; norm parallelism; Birkhoff orthogonality.

* Corresponding author.

- (ii) $x \perp_B \|y\|x + \lambda\|x\|y$.
- (iii) $y \perp_B \|x\|y + \bar{\lambda}\|y\|x$.

Furthermore, Zamani [13] conducted an investigation into the characterization of norm parallelism within specific spaces of continuous functions. Wojcik [11] offered characterizations of norm parallelism for bounded linear operators between normed spaces and introduced a fascinating application pertinent to the invariant subspace problem. For those seeking a more profound understanding of norm parallelism, it is recommended to consult the references [7, 10, 12].

In Section 2, we introduce the concept of *numerical radius parallelism* in $\mathbb{B}(X)$ and establish some characterizations. Additionally, we show that numerical radius parallelism does not coincide with the linear dependence and lacks transitivity.

In Section 3, we introduce the concept of *numerical radius Birkhoff orthogonality* in $\mathbb{B}(X)$ and offer a characterization. Furthermore, we demonstrate that numerical radius Birkhoff orthogonality is neither additive on the left nor on the right. Finally, we provide a characterization of numerical radius parallelism in terms of numerical radius Birkhoff orthogonality.

2. NUMERICAL RADIUS PARALLELISM IN $\mathbb{B}(X)$

The *numerical radius* of $T \in \mathbb{B}(H)$, where $(H, \langle \cdot, \cdot \rangle)$ is a complex Hilbert space, is given by

$$\omega(T) := \sup\{|\langle Tx, x \rangle| : x \in S_H\}.$$

The *numerical radius* of bounded linear operators on a normed space X (cf. [3]) is the semi-norm defined on $\mathbb{B}(X)$ by

$$v(T) := \sup\{|x^*(Tx)| : x \in S_X, x^* \in J(x)\} \quad (T \in \mathbb{B}(X)),$$

where $J(x) := \{x^* \in S_{X^*} : x^*(x) = \|x\|\}$ is the *duality mapping* for x . Clearly, for any $x \in S_H$, there exist a unique $x^* \in S_{H^*}$ such that $x^*(x) = 1$, that is $x^* = \langle \cdot, x \rangle$. Consequently, for $T \in \mathbb{B}(H)$, we have $v(T) = \omega(T)$. This demonstrates that the concept of the numerical radius of bounded operators on a Hilbert space generalizes to the numerical radius of bounded operators on a normed space.

In the context of $\mathbb{B}(X)$, the alternative Daugavet equation [6]

$$\max_{\lambda \in \mathbb{T}} \|I + \lambda T\| = 1 + \|T\|$$

is a particular case of norm parallelism. In [3], it was shown that, the previous equation holds if and only if $v(T) = \|T\|$.

Mehrazin *et al.* [7] introduced the concept of numerical radius parallelism for Hilbert space operators and established a characterization, as follows.

Definition 2. [7, Definition 1.1] Let H be a complex Hilbert space. An element $T \in \mathbb{B}(H)$ is called *numerical radius parallel* to another element $S \in \mathbb{B}(H)$, denoted by $T \parallel_{\omega} S$, if $\omega(T + \lambda S) = \omega(T) + \omega(S)$ for some $\lambda \in \mathbb{T}$.

Theorem 3. [7, Theorem 2.2] Let $(H, \langle \cdot, \cdot \rangle)$ be a complex Hilbert space and $T, S \in \mathbb{B}(H)$. Then the following conditions are equivalent:

- (i) $T \parallel_{\omega} S$.
- (ii) There exists a sequence of unit vectors $\{x_n\}$ in H such that

$$\lim_{n \rightarrow \infty} |\langle T x_n, x_n \rangle \langle S x_n, x_n \rangle| = \omega(T)\omega(S).$$

In addition, if $\{x_n\}$ is a sequence of unit vectors in H satisfying (ii), then it also satisfies

$$\lim_{n \rightarrow \infty} |\langle T x_n, x_n \rangle| = \omega(T) \quad \text{and} \quad \lim_{n \rightarrow \infty} |\langle S x_n, x_n \rangle| = \omega(S).$$

We now extend the concept of numerical radius parallelism, which was originally defined on $\mathbb{B}(H)$, to $\mathbb{B}(X)$.

Definition 4. An element $T \in \mathbb{B}(X)$ is called *numerical radius parallel* to another element $S \in \mathbb{B}(X)$, denoted by $T \parallel_v S$, if $v(T + \lambda S) = v(T) + v(S)$ for some $\lambda \in \mathbb{T}$.

Remark 5. Since $A \mapsto v(A)$ is a semi-norm on $\mathbb{B}(X)$, the condition $T \parallel_v S$ holds whenever $v(T + \lambda S) \geq v(T) + v(S)$ for some $\lambda \in \mathbb{T}$. In particular, if $T, S \in \mathbb{B}(X)$ satisfy $v(T)v(S) = 0$, then $v(T + S) \geq |v(T) - v(S)| = v(T) + v(S)$ and hence $T \parallel_v S$.

If $T, S \in \mathbb{B}(X)$ are linearly dependent, say $S = \alpha T$ for some $\alpha \neq 0$. Then

$$v\left(T + \frac{\bar{\alpha}}{|\alpha|} S\right) = v(T + |\alpha|T) = (1 + |\alpha|)v(T) = v(T) + v(S).$$

Consequently, $T \parallel_v S$. However, the converse of this statement is invariably false, as established in the following theorem.

Theorem 6. Let X be a normed space of dimension at least 2. Then the numerical radius parallelism in $\mathbb{B}(X)$ dose not coincide with the linear dependence.

To establish this theorem, we need the following proposition.

Proposition 7. Let $T \in \mathbb{B}(X)$. Then $T \parallel_v \alpha I$ for all $\alpha \in \mathbb{F}$.

Proof. Without loss of generality, we assume that $\alpha \neq 0$. By the definition of $v(T)$, there exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |x_n^*(T x_n)| = v(T).$$

For every $n \in \mathbb{Z}_{\geq 1}$, there exists $\lambda_n \in \mathbb{T}$ such that

$$x_n^*(T x_n) = \lambda_n |x_n^*(T x_n)|.$$

By passing to a subsequence, we may assume that $\{\lambda_n\}$ converges to some $\lambda \in \mathbb{T}$. Thus

$$\lim_{n \rightarrow \infty} x_n^*(Tx_n) = \lambda v(T).$$

Observe that $v(\alpha I) = |\alpha|$, $\lambda \bar{\alpha}/|\alpha| \in \mathbb{T}$ and

$$\begin{aligned} & |x_n^*[(T + \lambda \bar{\alpha}/|\alpha| \cdot \alpha I)x_n]| \\ &= |\lambda_n x_n^*(Tx_n) + \lambda |\alpha| | \rightarrow |\lambda v(T) + \lambda v(\alpha I)| = v(T) + v(\alpha I) \end{aligned}$$

As a result,

$$v(T + \lambda \bar{\alpha}/|\alpha| \cdot \alpha I) \geq v(T) + v(\alpha I).$$

Consequently, $T \parallel_v \alpha I$. \square

Proof of Theorem 6. Let $x \in S_X$, $x^* \in J(x)$ and $T = x^*(\cdot)x$. We can apply the previous proposition to obtain $T \parallel_v I$. However, it is evident that T and I are linearly independent. \square

In the subsequent example, we shall demonstrate that the relations \parallel_v and \parallel are in general not comparable.

Example 8. Let $X = \mathbb{R}^2$ equipped with the norm defined by $\|(x, y)\| = (x^4 + y^4)^{1/4}$ for $x, y \in \mathbb{R}$. Consider

$$\begin{aligned} T : X &\rightarrow X, (x, y) \mapsto (0, x), \\ S : X &\rightarrow X, (x, y) \mapsto (y, x). \end{aligned}$$

Clearly,

$$\begin{aligned} \|T\| &= \|S\| = 1, \\ \|T + S\| &= 2, \end{aligned}$$

which implies that $T \parallel S$. We will show that $T \not\parallel_v S$. Actually, for any $z^* \in S_{X^*}$, there exist $\alpha, \beta \in \mathbb{R}$ such that

$$\alpha^{4/3} + \beta^{4/3} = 1$$

and

$$z^*(x, y) = \alpha x + \beta y$$

for all $(x, y) \in X$. Given that, in addition, $(x, y) \in S_X$ and $z^*(x, y) = 1$, we arrive at the following equation:

$$\alpha x + \beta y = 1 = (\alpha^{4/3} + \beta^{4/3})^{3/4} (x^4 + y^4)^{1/4}.$$

Pursuant to the equality conditions of the Hölder's inequality, we further deduce

$$|\alpha|^{4/3} |y|^4 = |\beta|^{4/3} |x|^4,$$

and hence

$$\alpha = x^3, \beta = y^3.$$

This leads to

$$\begin{aligned}
 v(T) &= \sup\{|z^*(Tz)| : z \in S_X, z^* \in J(z)\} \\
 &= \sup\{|\beta x| : x^4 + y^4 = 1, \alpha^{4/3} + \beta^{4/3} = 1, \alpha x + \beta y = 1\} \\
 &= \max\{xy^3 : x^4 + y^4 = 1\}, \\
 v(S) &= \sup\{|z^*(Sz)| : z \in S_X, z^* \in J(z)\} \\
 &= \sup\{|\alpha y + \beta x| : x^4 + y^4 = 1, \alpha^{4/3} + \beta^{4/3} = 1, \alpha x + \beta y = 1\} \\
 &= \max\{yx^3 + xy^3 : x^4 + y^4 = 1\}, \\
 v(T + S) &= \sup\{|z^*[(T + S)z]| : z \in S_X, z^* \in J(z)\} \\
 &= \sup\{|\alpha y + 2\beta x| : x^4 + y^4 = 1, \alpha^{4/3} + \beta^{4/3} = 1, \alpha x + \beta y = 1\} \\
 &= \max\{yx^3 + 2xy^3 : x^4 + y^4 = 1\}, \\
 v(T - S) &= \sup\{|z^*[(T - S)z]| : z \in S_X, z^* \in J(z)\} \\
 &= \sup\{|\beta x - \alpha y| : x^4 + y^4 = 1, \alpha^{4/3} + \beta^{4/3} = 1, \alpha x + \beta y = 1\} \\
 &= \max\{yx^3 : x^4 + y^4 = 1\} = \max\{xy^3 : x^4 + y^4 = 1\}.
 \end{aligned}$$

Under the condition $x^4 + y^4 = 1$, we have

$$\begin{aligned}
 (xy^3)^{1/4} &= \left[3^{3/4} x \left(\frac{y}{3^{1/4}} \right)^3 \right]^{1/4} \leq \frac{y}{3^{1/16}} \left[\frac{x^4 + 3 \left(\frac{y}{3^{1/4}} \right)^4}{4} \right]^{1/4} = \left(\frac{3^{3/4}}{4} \right)^{1/4}, \\
 (yx^3 + xy^3)^{1/2} &= \sqrt{2} \sqrt{xy} \left(\frac{x^2 + y^2}{2} \right)^{1/2} \leq \sqrt{2} \left(\frac{x^4 + y^4}{2} \right)^{1/4} \left(\frac{x^4 + y^4}{2} \right)^{1/4} = 1,
 \end{aligned}$$

with equality in the first inequality if and only if $3^{1/4}x = y$ and equality in the second inequality if and only if $x = y$. Consequently,

$$yx^3 + 2xy^3 = (yx^3 + xy^3) + xy^3 < 1 + \frac{3^{3/4}}{4},$$

since equality in both inequalities cannot be achieved simultaneously under the given condition. As a result

$$v(T \pm S) < v(S) + v(T),$$

leading to $T \not\|_v S$.

On the other hand,

$$\|T \pm I\| = \max \left\{ [x^4 + (x \pm y)^4]^{1/4} : x^4 + y^4 = 1 \right\} < 2 = \|T\| + \|I\|.$$

In fact, under the condition $x^4 + y^4 = 1$, we have

$$\begin{aligned}
 &x^4 + (x \pm y)^4 = 1 + (x \pm y)^4 - y^4 \\
 &= \begin{cases} 1 + x(x + 2y)[(x + y)^2 + y^2] < 16, & |x| < 1, \\ 1, & |x| = 1. \end{cases}
 \end{aligned}$$

Thus $T \not\parallel_v I$, despite $T \parallel_v I$ as established in Proposition 7.

One of our main results is the following characterization of numerical radius parallelism in $\mathbb{B}(X)$.

Theorem 9. *Let $T, S \in \mathbb{B}(X)$. Then the following conditions are equivalent:*

- (i) $T \parallel_v S$.
- (ii) *There exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that*

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| |x_n^*(Sx_n)| = v(T)v(S).$$

- (iii) *There exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that*

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| = v(T) \quad \text{and} \quad \lim_{n \rightarrow \infty} |x_n^*(Sx_n)| = v(S).$$

Proof. (i) \Rightarrow (ii): Suppose that there exists $\lambda \in \mathbb{T}$ such that

$$v(T + \lambda S) = v(T) + v(S).$$

By the definition of the numerical radius of $T + \lambda S$, there exist $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |x_n^*[(T + \lambda S)x_n]| = v(T + \lambda S).$$

Then the following sequence of inequalities hold:

$$\begin{aligned} v(T) &\geq |x_n^*(Tx_n)| \geq |x_n^*(Tx_n)| + |x_n^*(Sx_n)| - v(S) \\ &\geq |x_n^*[(T + \lambda S)x_n]| - v(S) \\ &\rightarrow v(T + \lambda S) - v(S) = v(T), \\ v(S) &\geq |x_n^*(Sx_n)| \geq |x_n^*(Tx_n)| + |x_n^*(Sx_n)| - v(T) \\ &\geq |x_n^*[(T + \lambda S)x_n]| - v(T) \\ &\rightarrow v(T + \lambda S) - v(T) = v(S). \end{aligned}$$

Thus

$$(2.1) \quad \lim_{n \rightarrow \infty} |x_n^*(Tx_n)| = v(T) \quad \text{and} \quad \lim_{n \rightarrow \infty} |x_n^*(Sx_n)| = v(S)$$

hold. Consequently,

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| |x_n^*(Sx_n)| = v(T)v(S).$$

(ii) \Rightarrow (iii): If $v(T) = 0$ or $v(S) = 0$, the proof reduces to a straightforward case. Therefore, we proceed under the assumption that $v(T)v(S) \neq 0$. Suppose that there exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| |x_n^*(Sx_n)| = v(T)v(S).$$

Then

$$\begin{aligned} v(T) &\geq |x_n^*(Tx_n)| \geq \frac{|x_n^*(Tx_n)| |x_n^*(Sx_n)|}{v(S)} \rightarrow \frac{v(T)v(S)}{v(S)} = v(T), \\ v(S) &\geq |x_n^*(Sx_n)| \geq \frac{|x_n^*(Tx_n)| |x_n^*(Sx_n)|}{v(T)} \rightarrow \frac{v(T)v(S)}{v(T)} = v(S), \end{aligned}$$

implying (2.1) hold.

(iii) \Rightarrow (i): Without loss of generality, we assume that $v(T)v(S) \neq 0$. Suppose that there exist $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that (2.1) hold. By passing to a subsequence, we may assume that there exist $\mu_1, \mu_2 \in \mathbb{T}$ such that

$$x_n^*(Tx_n) \rightarrow v(T)\mu_1, \quad x_n^*(Sx_n) \rightarrow v(S)\mu_2.$$

Let $\lambda = \mu_1 \overline{\mu_2}$, then $\lambda \in \mathbb{T}$ and

$$x_n^*[(T + \lambda S)x_n] = x_n^*(Tx_n) + \mu_1 \overline{\mu_2} x_n^*(Sx_n) \rightarrow v(T)\mu_1 + v(S)\mu_1.$$

It follows that

$$v(T + \lambda S) \geq \lim_{n \rightarrow \infty} |x_n^*[(T + \lambda S)x_n]| = v(T) + v(S).$$

This leads to $T \parallel_v S$. □

Remark 10. *If X is a Hilbert space equipped with the inner product $\langle \cdot, \cdot \rangle$, then $x \in S_X$ and $x^* \in J(x)$ imply that $x^* = \langle \cdot, x \rangle$. Consequently, by applying the current theorem, we are able to reestablish Theorem 3.*

Corollary 11. *Let X be a finite dimensional normed space and $T, S \in \mathbb{B}(X)$. Then the following conditions are equivalent:*

- (i) $T \parallel_v S$.
- (ii) *There exist $x \in S_X$ and $x^* \in J(x)$ such that*

$$|x^*(Tx)| |x^*(Sx)| = v(T)v(S).$$

- (iii) *There exist $x \in S_X$ and $x^* \in J(x)$ such that*

$$|x^*(Tx)| = v(T) \quad \text{and} \quad |x^*(Sx)| = v(S).$$

Proof. (iii) \Rightarrow (ii): Clearly.

- (ii) \Rightarrow (i): It follows immediately from Theorem 9.

(i) \Rightarrow (iii): Suppose that $T \parallel_v S$. By Theorem 9, there exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| = v(T) \quad \text{and} \quad \lim_{n \rightarrow \infty} |x_n^*(Sx_n)| = v(S).$$

Since X is finite dimensional, by passing to a subsequence, we can assume that $\{x_n\}$ and $\{x_n^*\}$ converge to some $x \in S_X$ and some $x^* \in S_{X^*}$, respectively. Then

$$\begin{aligned}
|x^*(x) - 1| &= |x^*(x) - x_n^*(x_n)| \\
&\leq |x^*(x - x_n)| + |(x^* - x_n^*)(x_n)| \\
&\leq \|x - x_n\| + \|x^* - x_n^*\| \rightarrow 0, \\
||x^*(Tx)| - v(T)| &\leq |x^*(Tx) - x_n^*(Tx_n)| + |x_n^*(Tx_n) - v(T)| \\
&\leq |x^*(Tx - Tx_n)| + |(x^* - x_n^*)(Tx_n)| + |x_n^*(Tx_n) - v(T)| \\
&\leq \|T\|\|x - x_n\| + \|T\|\|x^* - x_n^*\| + |x_n^*(Tx_n) - v(T)| \rightarrow 0, \\
||x^*(Sx)| - v(S)| &\leq |x^*(Sx) - x_n^*(Sx_n)| + |x_n^*(Sx_n) - v(S)| \\
&\leq |x^*(Sx - Sx_n)| + |(x^* - x_n^*)(Sx_n)| + |x_n^*(Sx_n) - v(S)| \\
&\leq \|S\|\|x - x_n\| + \|S\|\|x^* - x_n^*\| + |x_n^*(Sx_n) - v(S)| \rightarrow 0.
\end{aligned}$$

Consequently, $x^* \in J(x)$, $|x^*(Tx)| = v(T)$ and $|x^*(Sx)| = v(S)$ hold. \square

Concisely stated, based on the definition of numerical radius parallelism or Theorem 9, fundamental properties are derived immediately and listed below without proof.

Proposition 12. (i) *Numerical radius parallelism possesses symmetry, that is, if $T, S \in \mathbb{B}(X)$ satisfy $T \parallel_v S$, then $S \parallel_v T$.*

(ii) *Numerical radius parallelism possesses homogeneity, which means that if $T, S \in \mathbb{B}(X)$ satisfy $T \parallel_v S$, then $\alpha T \parallel_v \beta S$ for all $\alpha, \beta \in \mathbb{F}$.*

A natural question then arises regarding its transitivity, i.e., whether the following implication is true:

$$T, S, R \in \mathbb{B}(X) \setminus \{0\}, T \parallel_v S, S \parallel_v R \Rightarrow T \parallel_v R.$$

The answer is negative. Mehrazin *et al.* [7, Example] presented an example to show that the numerical radius parallelism of Hilbert space operators is in general not transitive. We will establish a more thorough result.

Theorem 13. *Let X be a normed space with dimension not less than 2. Then the numerical radius parallelism in $\mathbb{B}(X)$ fails to be transitive.*

To this end, we need the following proposition, which provides a sufficient condition for norm parallelism in terms of numerical radius parallelism.

Proposition 14. *Suppose that $x, y \in X$, $x^* \in J(x)$ and $y^* \in J(y)$ such that $x^*(\cdot)x \parallel_v y^*(\cdot)y$ holds. Then $x \parallel y$.*

Proof. Without losing generality, we assume that $\|x\|\|y\| \neq 0$. Let $T = x^*(\cdot)x$ and $S = y^*(\cdot)y$. Then

$$\begin{aligned} v(T) &= \sup\{|z^* [x^*(z)x]| : z \in S_X, z^* \in S_{X^*}, z^*(z) = 1\} \leq \|x\|, \\ v(T) &\geq x^* \left[x^* \left(\frac{x}{\|x\|} \right) x \right] = \|x\|. \end{aligned}$$

It follows that $v(T) = \|x\|$. Similarly, $v(S) = \|y\|$. By Theorem 9, there exist sequences $\{z_n\} \subset S_X$ and $\{z_n^*\} \subset S_{X^*}$ with $z_n^*(z_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |z_n^* [x^*(z_n)x]| = \|x\| \quad \text{and} \quad \lim_{n \rightarrow \infty} |z_n^* [y^*(z_n)y]| = \|y\|.$$

Observe that

$$\|x\| = \lim_{n \rightarrow \infty} |z_n^* [x^*(z_n)x]| \leq \liminf_{n \rightarrow \infty} |z_n^*(x)| \leq \limsup_{n \rightarrow \infty} |z_n^*(x)| \leq \|x\|.$$

Thus $|z_n^*(x)| \rightarrow \|x\|$. Similarly, $|z_n^*(y)| \rightarrow \|y\|$. By passing to a subsequence, we may assume that there exist $\mu_1, \mu_2 \in \mathbb{T}$ such that

$$z_n^*(x) \rightarrow \|x\|\mu_1, \quad z_n^*(y) \rightarrow \|y\|\mu_2.$$

Let $\lambda = \mu_1 \overline{\mu_2}$, then $\lambda \in \mathbb{T}$ and

$$\|x\| + \|y\| \geq \|x + \lambda y\| \geq |z_n^*(x) + \lambda z_n^*(y)| \rightarrow \|\|x\|\mu_1 + \|y\|\mu_1 \overline{\mu_2} \mu_2\| = \|x\| + \|y\|.$$

As a result, $\|x + \lambda y\| = \|x\| + \|y\|$, leading to $x \parallel y$. \square

Recall that a point $x \in X$ is said to be a smooth point if its duality mapping $J(x)$ is a singleton. It is worth noting that, as a particular instance of Rademacher's theorem [8], almost every point (in the sense of Lebesgue measure) in a finite dimensional normed space is a smooth point. For further reference, see [2, Proposition 2.3].

Proof of Theorem 13. Let Y be a 2 dimensional subspace of X , $x \in S_Y$ a smooth point of Y , $f \in S_{Y^*}$ with $f(x) = 1$ and $y \in S_Y$ satisfies $f(y) = 0$. We claim that $x \not\parallel y$. If not, then there exists $\lambda \in \mathbb{T}$ such that

$$\|x + \lambda y\| = \|x\| + \|y\| = 2.$$

It follows that there exists $g \in S_{Y^*}$ such that

$$2 = g(x + \lambda y) = g(x) + \lambda g(y).$$

This leads to $g(x) = 1$ and $|g(y)| = 1$. Since x is a smooth point of Y , $f = g$. But then $g(y) = 0$, a contradiction. The Hahn-Banach theorem yields $x^*, y^* \in S_{X^*}$ such that $x^*(x) = y^*(y) = 1$. Applying Proposition 14, we arrive at $x^*(\cdot)x$ is not numerical radius parallel to $y^*(\cdot)y$. Since $x^*(\cdot)x \parallel_v I$ and $I \parallel_v y^*(\cdot)y$ hold as established in Proposition 7, we conclude that the numerical radius parallelism in $\mathbb{B}(X)$ is not transitive. \square

It is noteworthy that the converse of Proposition 14 holds true provided that X is a complex Hilbert space, as demonstrated in [7, Corollary 2.6]. In this context, we proceed to present some weaker conditions that suffice for the converse of Proposition 14 to be valid.

The subsequent theorem focus on *rotund points*, defined as elements of S_X that do not lie within any line segment contained in S_X . It is worth recalling that X is strictly convex (or smooth, resp.) precisely when S_X consists entirely of rotund points (or smooth points, resp.).

Theorem 15. *Let X be a normed space with dimension not less than 2. Then the implication (i) \Rightarrow (ii) holds.*

(i) *Every point $x \in S_X$ is a rotund point or a smooth point in any 2-dimensional subspace including x .*

(ii) *If $x, y \in X$ satisfy $x \parallel y$. Then for any $x^* \in J(x)$ and $y^* \in J(y)$, the condition $x^*(\cdot)x \parallel_v y^*(\cdot)y$ holds.*

In particular, if X is strictly convex or smooth, then (ii) hold.

Proof. Suppose that (i) holds and that $x, y \in X$ satisfy $x \parallel y$. By the homogeneity of \parallel and \parallel_v , we may assume that $x, y \in S_X$. Then there exists $\lambda \in \mathbb{T}$ such that $\|x + \lambda y\| = 2$.

Case 1: If x and y are linearly independent. Then x can not be a rotund point of $\text{span}\{x, y\}$. Otherwise we have $x = \lambda y$, a contradiction. Thus x is a smooth point of $\text{span}\{x, y\}$. Let $f \in J(x + \lambda y)$. Then

$$2 = f(x + \lambda y) = f(x) + f(\lambda y) \leq |f(x)| + |f(\lambda y)| \leq 2.$$

This leads to $f(x) = f(\lambda y) = 1$. Since x is a smooth point of $\text{span}\{x, y\}$, we arrive at $x^* = f$ on $\text{span}\{x, y\}$. Hence

$$\begin{aligned} v(x^*(\cdot)x + y^*(\cdot)y) &\geq |x^*(\lambda y)f(x) + y^*(\lambda y)f(y)| \\ &= |f(\lambda y)f(x) + y^*(y)f(\lambda y)| \\ &= 2 = v(x^*(\cdot)x) + v(y^*(\cdot)y), \end{aligned}$$

which entails that $x^*(\cdot)x \parallel_v y^*(\cdot)y$.

Case 2: If x and y are linearly dependent, then there exists $\mu \in \mathbb{T}$ such that $\mu x = y$. It follows that

$$\begin{aligned} v(x^*(\cdot)x + y^*(\cdot)y) &\geq |x^*(x)x^*(x) + y^*(x)x^*(y)| \\ &= |1 + y^*(x)x^*(\mu x)| = |1 + y^*(\mu x)x^*(x)| \\ &= |1 + y^*(y)x^*(x)| = 2 = v(x^*(\cdot)x) + v(y^*(\cdot)y), \end{aligned}$$

which implies that $x^*(\cdot)x \parallel_v y^*(\cdot)y$. □

Remark 16. *The implication (ii) \Rightarrow (i) is invalid. To illustrate this, consider the space $X = \mathbb{R}^2$ equipped with the norm of element (x_1, x_2) defined by*

$$\|(x_1, x_2)\| := \begin{cases} x_1^2 + x_2^2 & x_1 x_2 \geq 0, \\ \max\{|x_1|, |x_2|\} & x_1 x_2 \leq 0, \end{cases}$$

Then $y = (-1, 1) \in S_X$ is neither a rotund point nor a smooth point. Observe that every point in S_X that is distinct from $\pm y$ is a smooth point. By analogy with the proof of the previous theorem, it can be shown that (ii) holds (note that in Case 1 of the proof, the sole requirement is that x be a smooth point).

Nevertheless, we present an example to show that (ii) in the preceding theorem may fail in general.

Example 17. Let X be the \mathbb{F} -vector space \mathbb{F}^2 , where $\mathbb{F} = \mathbb{R}$ or \mathbb{C} , equipped with the norm of element (x_1, x_2) defined by $\|(x_1, x_2)\| = |x_1| + |x_2|$. Let $x = (1, 0)$ and $y = (0, 1)$. Then $x \parallel y$ since

$$\|(1, 0) + (0, 1)\| = 2 = \|(1, 0)\| + \|(0, 1)\|.$$

Identify X^* with \mathbb{F}^2 equipped with the norm of element (x_1, x_2) defined by $\|(x_1, x_2)\| = \max\{|x_1|, |x_2|\}$. Let $x^* = (1, 0)$ and $y^* = (0, 1)$. Then $x^* \in J(x)$ and $y^* \in J(y)$. We claim that $x^*(\cdot)x \not\parallel_v y^*(\cdot)y$. If not, then there exist $z = (z_1, z_2) \in S_X$, $z^* = (a, b) \in J(z)$ and $\lambda \in \mathbb{T}$ such that

$$\begin{aligned} 2 &\geq |az_1| + |bz_2| = |az_1 + \lambda bz_2| = |x^*(z)z^*(x) + \lambda y^*(z)y^*(x)| \\ &= v(x^*(\cdot)x + \lambda y^*(\cdot)y) = v(x^*(\cdot)x) + v(y^*(\cdot)y) = 2, \end{aligned}$$

leading to $|az_1| = |bz_2| = 1$. Thus $|z_1| \geq |az_1| = 1$ and $|z_2| \geq |bz_2| = 1$, contradicts to $|z_1| + |z_2| = 1$.

3. NUMERICAL RADIUS BIRKHOFF ORTHOGONALITY

Mal *et al.* [4] introduced the concept of numerical radius Birkhoff orthogonality for Hilbert space operators, as follows.

Definition 18. [4, Definition 1.1 and 1.2] Let H be a Hilbert space. An element $T \in \mathbb{B}(H)$ is called *numerical radius Birkhoff orthogonal* to another element $S \in \mathbb{B}(H)$, denoted by $T \perp_{\omega_B} S$, if $\omega(T + \alpha S) \geq \omega(T)$ for all $\alpha \in \mathbb{F}$.

As an application, they derive numerical radius inequalities for bounded linear operators on a complex Hilbert space, yielding improved lower bounds for the numerical radius of a matrix compared to existing estimates.

We now extend the concept of numerical radius Birkhoff orthogonality, which was originally defined on $\mathbb{B}(H)$, to $\mathbb{B}(X)$.

Definition 19. An element $T \in \mathbb{B}(X)$ is called *numerical radius Birkhoff orthogonal* to another element $S \in \mathbb{B}(X)$, denoted by $T \perp_{vB} S$, if $v(T + \alpha S) \geq v(T)$ for all $\alpha \in \mathbb{F}$.

Remark 20. Since $A \mapsto v(A)$ is a semi-norm on $\mathbb{B}(X)$, the condition $T \perp_{vB} S$ holds whenever $v(T)v(S) = 0$.

The following proposition follows directly from the definition of the numerical radius Birkhoff orthogonality.

Proposition 21. *Numerical radius Birkhoff orthogonality possesses homogeneity, that is, if $T, S \in \mathbb{B}(X)$ satisfy $T \perp_{vB} S$, then $\alpha T \perp_{vB} \beta S$ for all $\alpha, \beta \in \mathbb{F}$.*

A natural question then arises regarding its left additivity and right additivity, i.e., whether the following implications are true:

$$T \perp_{vB} R, S \perp_{vB} R \Rightarrow T + S \perp_{vB} R,$$

$$T \perp_{vB} S, T \perp_{vB} R \Rightarrow T \perp_{vB} S + R.$$

The answer is negative. More precisely, we will establish the following theorem.

Theorem 22. *Let X be a normed space of dimension at least 2. Then the numerical radius Birkhoff orthogonality in $\mathbb{B}(X)$ is neither additive on the left nor on the right.*

To establish this theorem, we need the following proposition.

Proposition 23. *Let X be a normed space. Then $I \perp_{vB} T$ whenever $T \in \mathbb{B}(X)$ is not injective.*

Proof. There exists $x \in S_X$ such that $Tx = 0$. Let $x^* \in J(x)$. Then for any $\alpha \in \mathbb{F}$, we have

$$v(I + \alpha T) \geq x^* [(I + \alpha T)(x)] = 1 = v(I),$$

leading to the desired result. \square

Proof of Theorem 22. Let $x \in S_X$ and $x^* \in J(x)$. Clearly, $x^*(\cdot)x$ and $I - x^*(\cdot)x$ are not injective. We can apply the previous proposition to arrive at

$$I \perp_{vB} x^*(\cdot)x, \quad I \perp_{vB} I - x^*(\cdot)x.$$

Since $v(I - I) = 0 < 1 = v(I)$, i.e., $I \not\perp_{vB} I$, we conclude that the numerical radius Birkhoff orthogonality in $\mathbb{B}(X)$ is not additive on the right.

On the other hand, suppose for contrary that the numerical radius Birkhoff orthogonality in $\mathbb{B}(X)$ is additive on the left. Observe that

$$\begin{aligned} v\left(-x^*(\cdot)x + \alpha[I - x^*(\cdot)x]\right) &\geq \left| -x^*(x)x^*(x) + \alpha[x^*(x) - x^*(x)x^*(x)] \right| \\ &= 1 = v\left(-x^*(\cdot)x\right) \end{aligned}$$

for any $\alpha \in \mathbb{F}$. Thus $-x^*(\cdot)x \perp_{vB} I - x^*(\cdot)x$. Since $I \perp_{vB} I - x^*(\cdot)x$, we obtain

$$I - x^*(\cdot)x \perp_{vB} I - x^*(\cdot)x.$$

Notice that there exists $y \in S_X$ such that $x^*(y) = 0$. Let $y^* \in J(y)$. However,

$$v\left([I - x^*(\cdot)x] - [I - x^*(\cdot)x]\right) = 0 < 1 = |y^*(y) - x^*(y)y^*(x)| \leq v\left(I - x^*(\cdot)x\right),$$

a contradiction. \square

We now establish a characterization of numerical radius Birkhoff orthogonality in $\mathbb{B}(X)$, which generalizes the characterizations of numerical radius Birkhoff orthogonality for Hilbert space operators [4, Theorem 2.3 and 2.5].

Theorem 24. *The condition $T \perp_{vB} S$ holds if and only if for any $\lambda \in \mathbb{T}$ there exist $\{x_{\lambda,n}\}_{n=1}^\infty \subset S_X$ and $\{x_{\lambda,n}^*\}_{n=1}^\infty \subset S_{X^*}$ with $x_{\lambda,n}^*(x_{\lambda,n}) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$, such that the following conditions hold:*

$$\lim_{n \rightarrow \infty} |x_{\lambda,n}^*(Tx_{\lambda,n})| = v(T), \quad \lim_{n \rightarrow \infty} \operatorname{Re} \left[\overline{\lambda x_{\lambda,n}^*(Tx_{\lambda,n})} x_{\lambda,n}^*(Sx_{\lambda,n}) \right] \geq 0.$$

Proof. Suppose that $T \perp_{vB} S$. Then for any $\lambda \in \mathbb{T}$ and all $n \in \mathbb{Z}_{\geq 1}$, we have

$$v\left(T + \frac{\lambda}{\sqrt{n}}S\right) \geq v(T).$$

Observe that, for each $n \in \mathbb{Z}_{\geq 1}$, there exist $x_{\lambda,n} \in S_X$ and $x_{\lambda,n}^* \in S_{X^*}$ such that $x_{\lambda,n}^*(x_{\lambda,n}) = 1$ and

$$\left| x_{\lambda,n}^* \left[\left(T + \frac{\lambda}{\sqrt{n}}S \right) (x_{\lambda,n}) \right] \right| \geq \frac{n}{n+1} v\left(T + \frac{\lambda}{\sqrt{n}}S\right) \geq \frac{n}{n+1} v(T).$$

It follows that

$$\begin{aligned} & \left| x_{\lambda,n}^*(Tx_{\lambda,n}) \right| \geq \left| x_{\lambda,n}^* \left[\left(T + \frac{\lambda}{\sqrt{n}}S \right) (x_{\lambda,n}) \right] \right| - \frac{|x_{\lambda,n}^*(Sx_{\lambda,n})|}{\sqrt{n}} \\ & \geq \frac{n}{n+1} v(T) - \frac{v(S)}{\sqrt{n}}, \\ & \operatorname{Re} \left[\overline{\lambda x_{\lambda,n}^*(Tx_{\lambda,n})} x_{\lambda,n}^*(Sx_{\lambda,n}) \right] \\ & = \frac{\sqrt{n}}{2} \left\{ \left| x_{\lambda,n}^* \left[\left(T + \frac{\lambda}{\sqrt{n}}S \right) (x_{\lambda,n}) \right] \right|^2 - |x_{\lambda,n}^*(Tx_{\lambda,n})|^2 - \frac{1}{n} |x_{\lambda,n}^*(Sx_{\lambda,n})|^2 \right\} \\ & \geq \frac{\sqrt{n}}{2} \left\{ \left[\frac{n}{n+1} v(T) \right]^2 - v(T)^2 - \frac{v(S)^2}{n} \right\} = -\frac{(2n+1)\sqrt{n}}{2(n+1)^2} v(T)^2 - \frac{v(S)^2}{2\sqrt{n}}. \end{aligned}$$

Since the sequences $\{x_{\lambda,n}^*(Tx_{\lambda,n})\}$ and $\{x_{\lambda,n}^*(Sx_{\lambda,n})\}$ are bounded, by passing to a subsequence, we can assume that they converge. Consequently,

$$\lim_{n \rightarrow \infty} |x_{\lambda,n}^*(Tx_{\lambda,n})| \geq v(T), \quad \lim_{n \rightarrow \infty} \operatorname{Re} \left[\overline{\lambda x_{\lambda,n}^*(Tx_{\lambda,n})} x_{\lambda,n}^*(Sx_{\lambda,n}) \right] \geq 0.$$

Since $\lim_{n \rightarrow \infty} |x_{\lambda,n}^*(Tx_{\lambda,n})| \leq v(T)$ holds, we establish the necessity.

It remains to prove the sufficiency. For any $\alpha \in \mathbb{F}$, there exists $\lambda \in \mathbb{T}$ such that $\alpha = |\alpha|\lambda$. As a result,

$$\begin{aligned} v(T + \alpha S)^2 &\geq \limsup_{n \rightarrow \infty} |x_{\lambda,n}^* [(T + |\alpha|\lambda S)(x_{\lambda,n})]|^2 \\ &= \limsup_{n \rightarrow \infty} \left\{ |x_{\lambda,n}^*(Tx_{\lambda,n})|^2 + |\alpha|^2 |x_{\lambda,n}^*(Sx_{\lambda,n})|^2 \right. \\ &\quad \left. + 2|\alpha| \operatorname{Re} \left[\overline{\lambda x_{\lambda,n}^*(Tx_{\lambda,n})} x_{\lambda,n}^*(Sx_{\lambda,n}) \right] \right\} \\ &\geq \lim_{n \rightarrow \infty} |x_{\lambda,n}^*(Tx_{\lambda,n})|^2 = v(T)^2, \end{aligned}$$

i.e., $v(T + \alpha S) \geq v(T)$. Therefore, $T \perp_{vB} S$ holds. \square

The following theorem, analogies to Theorem 1, provides a characterization of numerical radius parallelism in terms of numerical radius Birkhoff orthogonality.

Theorem 25. *Let X be a normed space and $T, S \in \mathbb{B}(X)$. Then $T \parallel_v S$ if and only if there exists $\lambda \in \mathbb{T}$ such that $T \perp_{vB} v(S)T + \lambda v(T)S$.*

Proof. Without lose of generality, we assume that $v(T)v(S) \neq 0$.

Suppose that $T \parallel_v S$. Then by Theorem 9, there exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\lim_{n \rightarrow \infty} |x_n^*(Tx_n)| = v(T) \quad \text{and} \quad \lim_{n \rightarrow \infty} |x_n^*(Sx_n)| = v(S).$$

By passing to a subsequence, we may assume that $\{x_n^*(Tx_n)\}$ and $\{x_n^*(Sx_n)\}$ exist.

Let

$$\lambda = - \lim_{n \rightarrow \infty} \frac{v(S)x_n^*(Tx_n)}{v(T)x_n^*(Sx_n)}.$$

Then $\lambda \in \mathbb{T}$. For every $\mu \in \mathbb{T}$ and $n \in \mathbb{Z}_{\geq 1}$, let $x_{\mu,n} = x_n$ and $x_{\mu,n}^* = x_n^*$. It follows that

$$\lim_{n \rightarrow \infty} x_{\mu,n}^*(v(S)T + \lambda v(T)S)(x_{\mu,n}) = 0$$

for all $\mu \in \mathbb{T}$. Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} |x_{\mu,n}^*(Tx_{\mu,n})| &= v(T), \\ \lim_{n \rightarrow \infty} \operatorname{Re} \left[\overline{\mu x_{\mu,n}^*(Tx_{\mu,n})} x_{\mu,n}^*(v(S)T + \lambda v(T)S)(x_{\mu,n}) \right] &= 0, \end{aligned}$$

for all $\mu \in \mathbb{T}$. Applying Theorem 24, we obtain $T \perp_{vB} v(S)T + \lambda v(T)S$.

Conversely, suppose that there exists $\lambda \in \mathbb{T}$ such that $T \perp_{vB} v(S)T + \lambda v(T)S$. By Theorem 24, there exist sequences $\{x_n\} \subset S_X$ and $\{x_n^*\} \subset S_{X^*}$ with $x_n^*(x_n) = 1$ for all $n \in \mathbb{Z}_{\geq 1}$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} |x_n^*(Tx_n)| &= v(T), \\ \lim_{n \rightarrow \infty} \operatorname{Re} \left[-\overline{x_n^*(Tx_n)} x_n^*(v(S)T + \lambda v(T)S)(x_n) \right] &\geq 0. \end{aligned}$$

By passing to a subsequence, we may assume that $\{x_n^*(Tx_n)\}$ and $\{x_n^*(Sx_n)\}$ exist. Let

$$\mu = \lim_{n \rightarrow \infty} \frac{x_n^*(Tx_n)}{v(T)}.$$

Then $\mu \in \mathbb{T}$ and

$$\begin{aligned} & \operatorname{Re} \left[-\lambda \bar{\mu} v(T)^2 \lim_{n \rightarrow \infty} x_n^*(Sx_n) - v(T)^2 v(S) \right] \\ &= \lim_{n \rightarrow \infty} \operatorname{Re} \left[-\overline{x_n^*(Tx_n)} x_n^*(v(S)T + \lambda v(T)S)(x_n) \right] \geq 0. \end{aligned}$$

Thus

$$v(S) \leq \operatorname{Re} \left[-\lambda \bar{\mu} \lim_{n \rightarrow \infty} x_n^*(Sx_n) \right] \leq \lim_{n \rightarrow \infty} |x_n^*(Sx_n)| \leq v(S),$$

leading to

$$\lim_{n \rightarrow \infty} |x_n^*(Sx_n)| = v(S).$$

Consequently, we can invoke Theorem 9 to conclude that $T \parallel_v S$. □

Data Availability

No data were used to support this study.

Conflicts of Interest

The author(s) declare(s) that there is no conflict of interest regarding the publication of this paper.

Funding Statement

This work was supported by the National Natural Science Foundation of P. R. China (No.11971493 and 12071491).

REFERENCES

- [1] G. Birkhoff. Orthogonality in linear metric spaces. *Duke Math. J.*, 1(2):169–172, 1935.
- [2] A. Blanco and A. Turnšek. On maps that preserve orthogonality in normed spaces. *Proc. Roy. Soc. Edinburgh Sect. A*, 136(4):709–716, 2006.
- [3] J. Duncan, C. M. McGregor, J. D. Pryce, and A. J. White. The numerical index of a normed space. *J. London Math. Soc. (2)*, 2:481–488, 1970.
- [4] A. Mal, K. Paul, and J. Sen. Birkhoff-James orthogonality and numerical radius inequalities of operator matrices. *Monatsh. Math.*, 197(4):717–731, 2022.
- [5] A. Mal, D. Sain, and K. Paul. On some geometric properties of operator spaces. *Banach J. Math. Anal.*, 13(1):174–191, 2019.
- [6] M. Martín and T. Oikhberg. An alternative Daugavet property. *J. Math. Anal. Appl.*, 294(1):158–180, 2004.
- [7] M. Mehrazin, M. Amyari, and A. Zamani. Numerical radius parallelism of Hilbert space operators. *Bull. Iranian Math. Soc.*, 46(3):821–829, 2020.
- [8] H. Rademacher. Über partielle und totale differenzierbarkeit von Funktionen mehrerer Variablen und über die Transformation der Doppelintegrale. *Math. Ann.*, 79(4):340–359, 1919.
- [9] A. Seddik. Rank one operators and norm of elementary operators. *Linear Algebra Appl.*, 424(1):177–183, 2007.
- [10] P. Wójcik and A. Zamani. From norm derivatives to orthogonalities in Hilbert C^* -modules. *Linear Multilinear Algebra*, 71(6):875–888, 2023.
- [11] P. Wójcik. Norm-parallelism in classical M -ideals. *Indag. Math. (N.S.)*, 28(2):287–293, 2017.
- [12] A. Zamani. The operator-valued parallelism. *Linear Algebra Appl.*, 505:282–295, 2016.
- [13] A. Zamani. Characterizations of norm-parallelism in spaces of continuous functions. *Bull. Iranian Math. Soc.*, 45(2):557–567, 2019.
- [14] A. Zamani and M. S. Moslehian. Norm-parallelism in the geometry of Hilbert C^* -modules. *Indag. Math. (N.S.)*, 27(1):266–281, 2016.

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P. R. CHINA
 Email address: bjjiaye@mail2.sysu.edu.cn

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P. R. CHINA
 Email address: xiehy33@mail2.sysu.edu.cn

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P. R. CHINA
 Email address: stslj@mail.sysu.edu.cn