

LIOUVILLE THEOREM FOR V -HARMONIC HEAT FLOWS

HAN LUO, WEIKE YU, AND XI ZHANG

ABSTRACT. In this paper, we investigate V -harmonic heat flows from complete Riemannian manifolds with nonnegative Bakry-Emery Ricci curvature to complete Riemannian manifolds with sectional curvature bounded above. We give a gradient estimate of ancient solutions to this flow and establish a Liouville type theorem, which is a complementarity of [CQ23].

1. INTRODUCTION

In 1975, Yau [Yau75] established the celebrated Liouville theorem which shows that on a complete Riemannian manifold of nonnegative Ricci curvature, there does not exist any nontrivial harmonic function bounded from one side. Later, Souplet–Zhang [SZ06] proved the following parabolic version of Yau’s Liouville theorem.

Theorem 1.1. *Let (M, g) be a complete Riemannian manifold with nonnegative Ricci curvature. Then we have*

(1) *Let $u : M \times (-\infty, 0] \rightarrow (0, \infty)$ be a positive ancient solution to the heat equation. If*

$$u(x, t) = \exp[o(d(x) + \sqrt{|t|})]$$

near infinity, then u must be a constant. Here $d(x)$ denotes the Riemannian distance from a fixed point;

(2) *Let $u : M \times (-\infty, 0] \rightarrow \mathbb{R}$ be an ancient solution to the heat equation. If*

$$u(x, t) = o(d(x) + \sqrt{|t|})$$

near infinity, then u is a constant.

Recently, Kunikawa-Sakurai [KS21a, KS21b] have proved some analogue Liouville type theorems for an ancient super Ricci flow under a growth condition concerning reduced distance. One of their main results is stated as follows.

Theorem 1.2. *Let $(M, g(t))_{t \in [0, \infty)}$ be an admissible, complete backward super Ricci flow. We assume*

$$\mathcal{D}(U) \geq 0, \quad \mathcal{H}(U) \geq -\frac{\tilde{S}}{t}, \quad \tilde{S} \geq 0$$

for all vector fields U . Here $S = \frac{1}{2}\partial_t g$, $\tilde{S} = \text{tr} S$, $\mathcal{D}(U) = -\partial_t \tilde{S} - \Delta \tilde{S} - 2\|S\|^2 + 4\text{div} S(U) - 2g(\nabla \tilde{S}, U) + 2\text{Ric}(U, U) - 2S(U, U)$ and $\mathcal{H}(U) = -\partial_t \tilde{S} -$

$\frac{1}{t}\tilde{S} - 2g(\nabla\tilde{S}, U) + 2S(U, U)$. Let (N, h) be a complete Riemannian manifold with sectional curvature bounded above by a positive constant κ . Assume that an open geodesic ball $B_{\pi/2\sqrt{\kappa}}(y_0)$ of radius $\pi/2\sqrt{\kappa}$ centered at y_0 in N does not meet the cut locus $\text{Cut}(y_0)$ of y_0 . Suppose that $u : M \times [0, \infty) \rightarrow N$ is a solution to backward harmonic map heat flow. If the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$ and u satisfies a growth condition

$$\frac{1}{\cos \sqrt{\kappa}\rho(u(x, t))} = o(\sigma(x, t)^{1/2} + t^{1/4})$$

near infinity, then u is constant. Here $\rho : N \rightarrow \mathbb{R}$ is the Riemannian distance function from a fixed point $y_0 \in N$.

Furthermore, using the above theorem, Kunikawa-Sakurai improved the Liouville theorem of Choi [Cho82] as follows:

Theorem 1.3. *Let (M, g) be a complete Riemannian manifold with non-negative Ricci curvature, and let (N, h) be a complete Riemannian manifold with sectional curvature bounded above by a positive constant κ . Assume that $B_{\pi/2\sqrt{\kappa}}(y_0)$ does not meet $\text{Cut}(y_0)$. Let $u : M \rightarrow N$ be a harmonic map. If the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$ and u satisfies a growth condition*

$$\frac{1}{\cos \sqrt{\kappa}\rho(u(x))} = o(d(x)^{1/2})$$

near infinity, then u is constant.

From now on, we suppose that (M, g) and (N, h) are both complete Riemannian manifolds. Recall that a map $u : M \rightarrow N$ is called a V -harmonic map if it solves

$$(1.1) \quad \tau_V(u) := \tau(u) + du(V) = 0.$$

According to [CJW15], we have learnt that some interesting maps are exactly V -harmonic maps, such as Hermitian harmonic maps(c.f. [JY93, LZ07, Zha12]), Weyl harmonic maps(c.f. [Kok09]), affine harmonic maps(c.f. [JS09, JS11]) and harmonic maps from Finsler manifolds into Riemannian manifolds(c.f. [Cen00, HS08, HW09, Mo01, MY05]). Moreover, the results and methods of V -harmonic maps have been applied successfully in submanifolds geometry and mean curvature flows(c.f. [CQ16, Qiu22a, Qiu22b, QS22]), since Gauss maps of a self-shrinker or a translating soliton can be seen as a V -harmonic map.

The heat flow associated to (1.1) is

$$(1.2) \quad \frac{\partial}{\partial t}u = \tau_V(u) = \tau(u) + du(V).$$

Chen-Jost-Wang [CJW15] proved the global existence and convergence of the V -harmonic map heat flow from compact manifolds into regular balls

by using the maximum principle and continuous method. In [CJQ12], Chen–Jost–Qiu gave the existence theorems for V -harmonic maps between complete manifolds. When the source manifold is a complete noncompact Riemannian manifold with nonnegative Bakry–Emery Ricci curvature, Chen–Qiu [CQ23] obtained Souplet–Zhang type and Choi type Liouville theorems of ancient solutions to (1.2) under certain growth condition near infinity. Inspired by the above works, this paper aims to establish the analogue of the Liouville theorem for ancient solutions to (1.2).

Suppose the vector field V satisfies

$$(1.3) \quad g(V, \nabla r) \leq v(r)$$

for some nondecreasing function $v(\cdot)$ satisfying

$$(1.4) \quad \lim_{r \rightarrow \infty} \frac{|v(r)|}{r} = C_V < +\infty,$$

where r is the Riemannian distance of (M, g) from a fixed point. Our main results are as follows:

Theorem 1.4. *Let (M, g) be a complete Riemannian manifold with nonnegative Bakry–Emery Ricci curvature, and (N, h) be a complete Riemannian manifold with sectional curvature bounded above by a positive constant κ . Assume that $B_{\pi/2\sqrt{\kappa}}(y_0)$ does not meet $\text{Cut}(y_0)$. Suppose that the vector field V satisfies (1.3) for some nondecreasing function $v(\cdot)$ satisfying (1.4). Let $u : M \times (-\infty, 0] \rightarrow N$ be an ancient solution to (1.2). If the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$, and if u satisfies a growth condition*

$$\frac{1}{\cos \sqrt{\kappa} \rho(u(x))} = o(r(x)^{1/2} + |t|^{1/4})$$

near infinity, then u is constant.

A direct application of Theorem 1.4 is the following result.

Corollary 1.5. *Let (M, g) be a complete Riemannian manifold with nonnegative Bakry–Emery Ricci curvature, and (N, h) be a complete Riemannian manifold with sectional curvature bounded above by a positive constant κ . Assume that $B_{\pi/2\sqrt{\kappa}}(y_0)$ does not meet $\text{Cut}(y_0)$. Suppose that the vector field V satisfies (1.3) for some nondecreasing function $v(\cdot)$ satisfying (1.4). Let $u : M \rightarrow N$ be a V -harmonic map. If the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$, and if u satisfies a growth condition*

$$\frac{1}{\cos \sqrt{\kappa} \rho(u(x))} = o(d(x)^{1/2})$$

near infinity, then u is constant.

2. PROOF OF THE MAIN THEOREM

Before giving our proof of main results, we prove a lemma we need. For $R, T > 0$, we define

$$\begin{aligned} B_R(x_0) &= \{x \in M \mid r(x) < R\}, \\ Q_{R,T} &= \overline{B_R(x_0)} \times [-T, 0] \subset M \times (-\infty, +\infty). \end{aligned}$$

Here $r(x)$ is the Riemannian distance function from a fixed point x_0 on M . As in [KS21b, Theorem 5.1], we set

$$(2.1) \quad \varphi = 1 - \cos \sqrt{\kappa} \rho, \quad b = \frac{1}{2} \left(1 + \sup_{Q_{R,T}} \varphi \circ u \right)$$

and

$$\omega = \frac{\|du\|^2}{(b - \varphi \circ u)^2},$$

where $u : M \times (-\infty, 0] \rightarrow N$ is an ancient solution of (1.2), then we have the following:

Lemma 2.1. *Let (M, g) be a complete Riemannian manifold with Bakky-Emery Ricci curvature*

$$\text{Ric}_V = \text{Ric} - \frac{1}{2} L_V g \geq -A$$

where Ric is the Ricci curvature of M , L_V is the Lie derivative along the vector field V on M , and A is a nonnegative constant. Let (N, h) be a Riemannian manifold with sectional curvature bounded above by a constant $\kappa > 0$. Assume that $B_{\pi/2\sqrt{\kappa}}(y_0)$ does not meet the cut locus $\text{Cut}(y_0)$ of $y_0 \in N$. Let $u : M \times (-\infty, 0] \rightarrow N$ be an ancient solution to V -harmonic map heat flow (1.2). Suppose that the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$. Then we have

$$(\Delta_V - \partial_t)\omega - \frac{2g(\nabla\omega, \nabla(\varphi \circ u))}{b - \varphi \circ u} \geq 2\kappa(1-b)(b - \varphi \circ u)\omega^2 - 2A\omega,$$

where $\Delta_V = \Delta + g(V, \nabla \cdot)$ is the V -Laplace operator.

Proof. Straightforward computations give that

$$\begin{aligned}
 \nabla\omega &= \frac{\nabla\|du\|^2}{(b-\varphi\circ u)^2} + 2\frac{\|du\|^2\nabla(\varphi\circ u)}{(b-\varphi\circ u)^3}, \\
 \Delta_V\omega &= \frac{\Delta_V\|du\|^2}{(b-\varphi\circ u)^2} + \frac{4g(\nabla\|du\|^2, \nabla(\varphi\circ u))}{(b-\varphi\circ u)^3} + \frac{2\|du\|^2\Delta_V(\varphi\circ u)}{(b-\varphi\circ u)^3} \\
 &\quad + \frac{6\|\nabla(\varphi\circ u)\|^2\|du\|^2}{(b-\varphi\circ u)^4} \\
 &= \frac{2g(\nabla\omega, \nabla(\varphi\circ u))}{b-\varphi\circ u} + \frac{2\|du\|^2\Delta_V(\varphi\circ u)}{(b-\varphi\circ u)^3} + \frac{\Delta_V\|du\|^2}{(b-\varphi\circ u)^2} \\
 &\quad + \frac{2g(\nabla\|du\|^2, \nabla(\varphi\circ u))}{(b-\varphi\circ u)^3} + \frac{2\|\nabla(\varphi\circ u)\|^2\|du\|^2}{(b-\varphi\circ u)^4}, \\
 \partial_t\omega &= \frac{\partial_t\|du\|^2}{(b-\varphi\circ u)^2} + \frac{2\|du\|^2\partial_t(\varphi\circ u)}{(b-\varphi\circ u)^3}.
 \end{aligned}$$

Then we get

$$\begin{aligned}
 (\Delta_V - \partial_t)\omega - \frac{2g(\nabla\omega, \nabla(\varphi\circ u))}{b-\varphi\circ u} &= \frac{(\Delta_V - \partial_t)\|du\|^2}{(b-\varphi\circ u)^2} + \frac{2\|du\|^2(\Delta_V - \partial_t)(\varphi\circ u)}{(b-\varphi\circ u)^3} \\
 &\quad + \frac{2g(\nabla\|du\|^2, \nabla(\varphi\circ u))}{(b-\varphi\circ u)^3} + \frac{2\|\nabla(\varphi\circ u)\|^2\|du\|^2}{(b-\varphi\circ u)^4}.
 \end{aligned}$$

Due to Hessian comparison Theorem [KS21b],

$$\begin{aligned}
 (\Delta_V - \partial_t)(\varphi\circ u) &= \sum_{i=1}^m \nabla^2\varphi(du(e_i), du(e_i)) + \langle \nabla\varphi\circ u, \tau_V(u) - \partial_t u \rangle_h \\
 &= \sum_{i=1}^m \nabla^2\varphi(du(e_i), du(e_i)) \\
 &\geq \kappa \cos\sqrt{\kappa}\rho\circ u \|du\|^2.
 \end{aligned}$$

The parabolic Bochner formula in [Qiu17, Lemma 1] tells us that

$$\begin{aligned}
 (\Delta_V - \partial_t)\|du\|^2 &= 2\|\nabla du\|^2 + 2\sum_{i=1}^m \langle du(\text{Ric}_V(e_i)), du(e_i) \rangle \\
 &\quad - 2\sum_{i,j=1}^m R^N(du(e_i), du(e_j), du(e_i), du(e_j)) \\
 &\geq 2\|\nabla du\|^2 - 2A\|du\|^2 - 2\kappa\|du\|^4.
 \end{aligned}$$

Hence,

$$\begin{aligned}
(\Delta_V - \partial_t)\omega - \frac{2g(\nabla\omega, \nabla(\varphi \circ u))}{b - \varphi \circ u} &= \frac{(\Delta_V - \partial_t)\|du\|^2}{(b - \varphi \circ u)^2} + \frac{2\|du\|^2(\Delta_V - \partial_t)(\varphi \circ u)}{(b - \varphi \circ u)^3} \\
&\quad + \frac{2g(\nabla\|du\|^2, \nabla(\varphi \circ u))}{(b - \varphi \circ u)^3} + \frac{2\|\nabla(\varphi \circ u)\|^2\|du\|^2}{(b - \varphi \circ u)^4} \\
&\geq \frac{1}{(b - \varphi \circ u)^2}(\|\nabla du\|^2 - 2A\|du\|^2 - 2\kappa\|du\|^4) \\
&\quad + \frac{\kappa \cos \sqrt{\kappa}\rho \circ u\|du\|^4}{(b - \varphi \circ u)^3} + \frac{2g(\nabla\|du\|^2, \nabla(\varphi \circ u))}{(b - \varphi \circ u)^3} \\
&\quad + \frac{2\|\nabla(\varphi \circ u)\|^2\|du\|^2}{(b - \varphi \circ u)^4} \\
&\geq 2\kappa(b - \varphi \circ u)^2 \left(\frac{\cos \sqrt{\kappa}\rho}{b - \varphi \circ u} - 1 \right) \omega^2 \\
&\quad - \frac{2A\|du\|^2}{(b - \varphi \circ u)^2} + 2\mathcal{F} \\
&= 2\kappa(1 - b)(b - \varphi \circ u)\omega^2 - \frac{2A\|du\|^2}{(b - \varphi \circ u)^2} + 2\mathcal{F},
\end{aligned}$$

where

$$\mathcal{F} = \frac{\|\nabla du\|^2}{(b - \varphi \circ u)^2} + \frac{g(\nabla\|du\|^2, \nabla(\varphi \circ u))}{(b - \varphi \circ u)^3} + \frac{\|\nabla(\varphi \circ u)\|^2\|du\|^2}{(b - \varphi \circ u)^4}.$$

By the inequality of arithmetic-geometric means, Kato and Cauchy-Schwarz inequalities, we have

$$\begin{aligned}
\frac{\|\nabla du\|^2}{(b - \varphi \circ u)^2} + \frac{\|\nabla(\varphi \circ u)\|^2\|du\|^2}{(b - \varphi \circ u)^4} &\geq \frac{2\|\nabla du\|\|\nabla(\varphi \circ u)\|\|du\|}{(b - \varphi \circ u)^3} \\
&\geq \frac{\|\nabla\|du\|^2\|\nabla(\varphi \circ u)\|}{(b - \varphi \circ u)^3} \\
&\geq \frac{|g(\nabla\|du\|^2, \nabla(\varphi \circ u))|}{(b - \varphi \circ u)^3},
\end{aligned}$$

so $\mathcal{F} \geq 0$, which ends the proof. \square

Theorem 2.2. *Let (M^m, g) be a complete Riemannian manifold with Bakry-Emery Ricci curvature*

$$\text{Ric}_V \geq -A$$

where $A \geq 0$ is a constant. Let (N, h) be a complete Riemannian manifold with sectional curvature bounded above by a positive constant κ . Assume that $B_{\pi/2\sqrt{\kappa}}(y_0)$ does not meet $\text{Cut}(y_0)$. Suppose that the vector field V satisfies (1.3) for some nondecreasing function $v(\cdot)$ satisfying (1.4). Let $u : M \times (-\infty, 0] \rightarrow N$ be an ancient solution to the V -harmonic map heat flow (1.2). Suppose that the image of u is contained in $B_{\pi/2\sqrt{\kappa}}(y_0)$. Then

there is a positive constant $C_m > 0$ depending only on m such that for any $R, T > 0$,

$$\sup_{Q_{R/2, T/4}} \frac{\|du\|}{b - \varphi \circ u} \leq \frac{C_m}{\sqrt{\kappa}} \left(\frac{1}{R} + \frac{1}{\sqrt{T}} + \sqrt{A} \right) \sup_{Q_{R, T}} \left(\frac{1}{\cos \sqrt{\kappa} \rho \circ u} \right)^2.$$

Firstly, recall the following elementary lemma:

Lemma 2.3 (cf. [SZ06]). *Let $R, T > 0, \alpha \in (0, 1)$. Then there is a smooth function $\psi = \psi(r, t) : [0, \infty) \times (-\infty, 0] \rightarrow [0, 1]$, which is supported on $[0, R] \times [-T, 0]$, and a constant $C_\alpha > 0$ depending only on α such that the following hold:*

- (1) $\psi \equiv 1$ on $[0, R/2] \times [-T/4, 0]$;
- (2) $\partial_r \psi \leq 0$ on $[0, \infty) \times (-\infty, 0]$, and $\partial_r \psi \equiv 0$ on $[0, R/2] \times (-\infty, 0]$;
- (3) we have

$$\frac{|\partial_r \psi|}{\psi^\alpha} \leq \frac{C_\alpha}{R}, \quad \frac{|\partial_r^2 \psi|}{\psi^\alpha} \leq \frac{C_\alpha}{R^2}, \quad \frac{|\partial_t \psi|}{\psi^{1/2}} \leq \frac{C}{T},$$

where $C > 0$ is a universal constant.

Proof of Theorem 2.2. We take a function $\psi : [0, \infty) \times (-\infty, 0] \rightarrow [0, 1]$ in Lemma 2.3 with $\alpha = 3/4$, and set

$$\psi(x, t) := \psi(r(x), t).$$

Denote

$$\Phi = (\Delta_V - \partial_t)(\psi\omega) - \frac{2g(\nabla\psi, \nabla(\psi\omega))}{\psi} - \frac{2g(\nabla(\psi\omega), \nabla(\varphi \circ u))}{b - \varphi \circ u}.$$

By direct computations and using Lemma 2.1, we have

$$\begin{aligned} \Phi &= \psi(\Delta_V - \partial_t)\omega - \frac{2\psi g(\nabla\omega, \nabla(\varphi \circ u))}{b - \varphi \circ u} - \frac{2\omega \|\nabla\psi\|^2}{\psi} \\ &\quad + \omega(\Delta_V - \partial_t)\psi - \frac{2\omega g(\nabla\psi, \nabla(\varphi \circ u))}{b - \varphi \circ u} \\ (2.2) \quad &\geq 2\kappa(1-b)(b - \varphi \circ u)\psi\omega^2 - 2A\psi\omega \\ &\quad + \omega(\Delta_V - \partial_t)\psi - \frac{2\omega \|\nabla\psi\|^2}{\psi} - \frac{2\omega g(\nabla\psi, \nabla(\varphi \circ u))}{b - \varphi \circ u}. \end{aligned}$$

Rewrite (2.2) as

$$\begin{aligned} 2\kappa(1-b)(b - \varphi \circ u)\psi\omega^2 &\leq 2A\psi\omega - \omega(\Delta_V - \partial_t)\psi \\ &\quad + \frac{2\omega \|\nabla\psi\|^2}{\psi} + \frac{2\omega g(\nabla\psi, \nabla(\varphi \circ u))}{b - \varphi \circ u} + \Phi. \end{aligned}$$

By the Young inequality and $0 \leq \psi \leq 1$, for any $\epsilon > 0$, we have

$$2A\psi\omega \leq \epsilon\psi^2\omega^2 + \frac{A^2}{\epsilon} \leq \epsilon\psi\omega^2 + \frac{A^2}{\epsilon}.$$

Lemma 2.3 tells us that

$$(2.3) \quad \frac{|\partial_r \psi|^2}{\psi^{3/2}} \leq \frac{C_{3/4}^2}{R^2} \quad \text{and} \quad \frac{|\partial_r^2 \psi|^2}{\psi^{3/2}} \leq \frac{C_{3/4}^2}{R^4}.$$

According to [CJQ12], we know

$$(2.4) \quad \begin{aligned} \Delta_V r &\leq \sqrt{(m-1)A} \coth\left(\sqrt{\frac{A}{m-1}} r\right) + v(r) \\ &\leq \left(\sqrt{(m-1)A} + \frac{m-1}{r}\right) + v(r) \end{aligned}$$

where m is the dimension of M . Using (2.3) and (2.4), we get at any $x \notin \text{Cut}(x_0)$

$$\begin{aligned} -\omega(\Delta_V - \partial_t)\psi &\leq -\omega(\partial_r \psi(\Delta_V - \partial_t)r + \partial_r^2 \psi \|\nabla r\|^2 - \partial_t \psi) \\ &\leq \omega(|\partial_r \psi|(\sqrt{(m-1)A} + \frac{m-1}{R} + v(R)) + |\partial_r^2 \psi| + |\partial_t \psi|) \\ &\leq (\epsilon\psi\omega^2 + \frac{(\sqrt{(m-1)A} + \frac{m-1}{R} + v(R))^2 |\partial_r \psi|^2}{4\epsilon\psi}) \\ &\quad + (\epsilon\psi\omega^2 + \frac{1}{4} \frac{|\partial_r^2 \psi|^2}{\epsilon\psi}) + (\epsilon\psi\omega^2 + \frac{1}{4} \frac{|\partial_t \psi|^2}{\epsilon\psi}) \\ &\leq 3\epsilon\psi\omega^2 + \frac{C_{3/4}^2(1+3(m-1)^2)\psi^{1/2}}{4\epsilon R^4} + \frac{3C_{3/4}^2 v^2(R)\psi^{1/2}}{4\epsilon R^2} \\ &\quad + \frac{C^2}{4\epsilon T^2} + \frac{3C_{3/4}^2(m-1)A\psi^{1/2}}{4\epsilon R^2} \\ &\leq 3\epsilon\psi\omega^2 + \frac{C_{3/4}^2(1+3(m-1)^2)}{4\epsilon R^4} + \frac{3C_{3/4}^2 v^2(R)}{4\epsilon R^2} \\ &\quad + \frac{C^2}{4\epsilon T^2} + \frac{3C_{3/4}^2(m-1)A}{4\epsilon R^2}, \end{aligned}$$

Due to the Young inequality, the following two inequalities hold respectively

$$\begin{aligned} \frac{2\omega\|\nabla\psi\|^2}{\psi} &\leq \epsilon\psi\omega^2 + \frac{\|\nabla\psi\|^4}{\epsilon\psi^3} \\ &\leq \epsilon\psi\omega^2 + \frac{C_{3/4}^4}{\epsilon} \frac{1}{R^4}, \\ \frac{2\omega g(\nabla\psi, \nabla(\varphi \circ u))}{b - \varphi \circ u} &\leq \frac{2\omega\|\nabla\psi\|\|\nabla(\varphi \circ u)\|}{b - \varphi \circ u} \\ &\leq 2\sqrt{\kappa}\omega^{3/2}\|\nabla\psi\| \\ &\leq \frac{3\epsilon}{4}\psi\omega^2 + \frac{4\kappa^2}{\epsilon^3} \frac{\|\nabla\psi\|^4}{\psi^3} \\ &\leq \frac{3\epsilon}{4}\psi\omega^2 + \frac{4\kappa^2 C_{3/4}^4}{\epsilon^3} \frac{1}{R^4}. \end{aligned}$$

The above estimates lead to

$$(2.5) \quad \begin{aligned} 2\kappa(1-b)(b-\varphi \circ u)\psi\omega^2 &\leq \frac{23\epsilon}{4}\psi\omega^2 + \frac{C^2}{4\epsilon} \frac{1}{T^2} + \frac{3C_{3/4}^2 v^2(R)}{4\epsilon} \frac{1}{R^2} \\ &\quad + \frac{C_{3/4}^2}{\epsilon} \left(\frac{1+3(m-1)^2}{4} + C_{3/4}^2 + \frac{4\kappa^2 C_{3/4}^2}{\epsilon^2} \right) \frac{1}{R^4} \\ &\quad + \frac{A^2}{\epsilon} + \frac{3C_{3/4}^2 (m-1)A}{4\epsilon R^2} + \Phi. \end{aligned}$$

at every point in $Q_{R,T}$, where the universal constants $C_{3/4}, C > 0$ are given in Lemma 2.3.

Suppose that the maximum of $\psi\omega$ in $Q_{R,T}$ is reached at (\bar{x}, \bar{t}) . We may assume that $\psi\omega$ is positive at (\bar{x}, \bar{t}) , or else the theorem follows trivially. Then at point (\bar{x}, \bar{t}) , we have

$$\nabla(\psi\omega) = 0, \quad \Delta_V(\psi\omega) \leq 0, \quad \partial_t(\psi\omega) \geq 0.$$

Thus

$$(2.6) \quad \Phi(\bar{x}, \bar{t}) = (\Delta_V - \partial_t)(\psi\omega) - \frac{2g(\nabla\psi, \nabla(\psi\omega))}{\psi} - \frac{2g(\nabla(\psi\omega), \nabla(\varphi \circ u))}{b - \varphi \circ u} \leq 0$$

at point (\bar{x}, \bar{t}) . Set

$$\delta = (1-b)(b - \varphi(u(\bar{x}, \bar{\tau}))).$$

Note that $\delta \in (0, \frac{1}{4}]$. In fact, from (2.1), we obtain that $0 \leq \varphi < 1$ and $\frac{1}{2} \leq b < 1$, and thus $\delta > 0$. By (2.1) and the inequality $ab \leq (\frac{a+b}{2})^2$ for any $a, b > 0$, we get $\delta \leq \frac{(\cos \sqrt{\kappa\rho \circ u})^2}{4} \leq \frac{1}{4}$. Since

$$\frac{1}{1-b} = 2 \sup_{Q_{R,T}} \frac{1}{\cos \sqrt{\kappa\rho \circ u}}$$

and

$$\frac{1}{b - \varphi(u(\bar{x}, \bar{\tau}))} \leq \frac{2}{1 - \sup_{Q_{R,T}} \varphi \circ u} = 2 \sup_{Q_{R,T}} \frac{1}{\cos \sqrt{\kappa\rho \circ u}},$$

it follows that

$$\frac{1}{\delta} \leq 4 \sup_{Q_{R,T}} \left(\frac{1}{\cos \sqrt{\kappa\rho \circ u}} \right)^2.$$

Setting $\epsilon = 4\kappa\delta/23$ in (2.5), combining with (2.6), we get

$$\begin{aligned} \kappa\delta\psi\omega^2 &\leq \frac{23C_{3/4}^2}{4\kappa\delta} \left(\frac{1+3(m-1)^2}{4} + C_{3/4}^2 + \frac{529C_{3/4}^2}{4\delta^2} \right) \frac{1}{R^4} \\ &\quad + \frac{23C^2}{16\kappa\delta} \frac{1}{T^2} + \frac{69C_{3/4}^2 v^2(R)}{16\kappa\delta} \frac{1}{R^2} + \frac{23}{4\kappa\delta} A^2 + \frac{69C_{3/4}^2 (m-1)A}{16\kappa\delta R^2} \end{aligned}$$

Hence

$$\begin{aligned}
(\psi\omega)^2 &\leq \frac{23C_{3/4}^2}{4\kappa^2\delta^2} \left(\frac{1+3(m-1)^2}{4} + C_{3/4}^2 + \frac{529C_{3/4}^2}{4\delta^2} \right) \frac{1}{R^4} \\
&\quad + \frac{23C^2}{16\kappa^2\delta^2} \frac{1}{T^2} + \frac{69C_{3/4}^2 v^2(R)}{16\kappa^2\delta^2} \frac{1}{R^2} + \frac{23}{4\kappa^2\delta^2} A^2 + \frac{69C_{3/4}^2 (m-1)A}{16\kappa^2\delta^2} \frac{1}{R^2} \\
&\leq \frac{23C_{3/4}^2}{4\kappa^2\delta^2} \left(\frac{1+6(m-1)^2}{4} + C_{3/4}^2 + \frac{529C_{3/4}^2}{4\delta^2} \right) \frac{1}{R^4} \\
&\quad + \frac{23C^2}{16\kappa^2\delta^2} \frac{1}{T^2} + \frac{69C_{3/4}^2 C_V}{16\kappa^2\delta^2} \frac{1}{R^4} + \frac{92+69C_{3/4}^2}{16\kappa^2\delta^2} A^2
\end{aligned}$$

Owing to $\delta \in (0, 1)$, there is a positive constant \bar{c}_m depending on m such that

$$(\psi\omega)^2(\bar{x}, \bar{t}) \leq \frac{\bar{c}_m}{\kappa^2\delta^4} \left(\frac{1}{R^4} + \frac{1}{T^2} + A^2 \right).$$

It follows that, for all $(x, t) \in Q_{R,T}$,

$$\psi\omega(x, t) \leq \psi\omega(\bar{x}, \bar{t}) \leq \frac{\bar{c}_m^{1/2}}{\kappa\delta^2} \left(\frac{1}{R^2} + \frac{1}{T} + A \right).$$

Using $\psi \equiv 1$ on $Q_{R/2, T/4}$, we know

$$\begin{aligned}
\frac{\|du\|}{b - \varphi \circ u} &\leq \frac{\bar{c}_m^{1/4}}{\sqrt{\kappa}\delta} \left(\frac{1}{R} + \frac{1}{\sqrt{T}} + \sqrt{A} \right) \\
&\leq \frac{4\bar{c}_m^{1/4}}{\sqrt{\kappa}} \left(\frac{1}{R} + \frac{1}{\sqrt{T}} + \sqrt{A} \right) \sup_{Q_{R,T}} \left(\frac{1}{\cos \sqrt{\kappa}\rho \circ u} \right)^2
\end{aligned}$$

on $Q_{R/2, T/4}$. This finishes the proof. \square

We are now in a position to show Theorem 1.4.

Proof of Theorem 1.4. Let $u : M \rightarrow N$ be an ancient solution to V-harmonic heat flow. For $R > 0$ we put

$$\mathcal{A}_R := \sup_{Q_{R,R^2}} \left(\frac{1}{\cos \sqrt{\kappa}\rho \circ u} \right)^2.$$

From the growth condition, we know that $\mathcal{A}_R = o(R)$ as $R \rightarrow \infty$. For any fixed point $(x, t) \in M \times (-\infty, 0)$ and a sufficiently large $R > 0$, due to Theorem 2.2, we know when $A = 0$,

$$\frac{\|du\|}{b} \leq \frac{\|du\|}{b - \varphi \circ u} \leq \frac{2c_m \mathcal{A}_R}{R}$$

at (x, t) . Note that $b \leq 1$. We complete the proof by letting $R \rightarrow \infty$. \square

ACKNOWLEDGEMENTS

REFERENCES

- [Cen00] Centore, Paul. Finsler Laplacians and minimal-energy maps. *Internat. J. Math.*, 11(1):1–13, 2000.
- [CJQ12] Chen, Qun and Jost, Jürgen and Qiu, Hongbing. Existence and Liouville theorems for V -harmonic maps from complete manifolds. *Ann. Global Anal. Geom.*, 42(4):565–584, 2012.
- [CJW15] Chen, Qun and Jost, Jürgen and Wang, Guofang. A maximum principle for generalizations of harmonic maps in Hermitian, affine, Weyl, and Finsler geometry. *J. Geom. Anal.*, 25(4):91–94, 2015.
- [CQ16] Chen, Qun and Qiu, Hongbing. Rigidity of self-shrinkers and translating solitons of mean curvature flows. *Adv. Math.*, 294:517–531, 2016.
- [CQ23] Chen, Qun and Qiu, Hongbing. Liouville theorems for ancient solutions to the V -harmonic map heat flows. *Nonlinear Anal.*, 233:Paper No. 113294, 9, 2023.
- [Cho82] Choi, Hyeong In. On the Liouville theorem for harmonic maps. *Proc. Amer. Math. Soc.*, 85(1):91–94, 1982.
- [HS08] Han, Jingwei and Shen, Yibing. Harmonic maps from complex Finsler manifolds. *Pacific J. Math.*, 236(2):314–356, 2008.
- [HW09] von der Mosel, Heiko and Winklmann, Sven. On weakly harmonic maps from Finsler to Riemannian manifolds. *Ann. Inst. H. Poincaré C Anal. Non Linéaire.*, 26(1):39–57, 2009.
- [JS09] Jost, Jürgen and Şimsir, Fatma Muazzez. Affine harmonic maps. *Analysis (Munich)*, 29(2):185–197, 2009.
- [JS11] Jost, J. and Şimsir, F. M.. Non-divergence harmonic maps. *Contemp. Math.*, 542:231–238, 2011.
- [JY93] Jost, Jürgen and Yau, Shing-Tung. A nonlinear elliptic system for maps from Hermitian to Riemannian manifolds and rigidity theorems in Hermitian geometry. *Acta Math.*, 170(2):221–254, 1993.
- [Kok09] Kokarev, Gerasim. On pseudo-harmonic maps in conformal geometry. *Proc. Lond. Math. Soc. (3)*, 99(1):168–194, 2009.
- [KS21a] Kunikawa, Keita and Sakurai, Yohei. Liouville theorem for heat equation along ancient super Ricci flow via reduced geometry. *J. Geom. Anal.*, 31(12):11899–11930, 2021.
- [KS21b] Kunikawa, Keita and Sakurai, Yohei. Liouville theorems for harmonic map heat flow along ancient super Ricci flow via reduced geometry. *Calc. Var. Partial Differential Equations*, 60(5):Paper No. 199, 24, 2021.
- [LZ07] Li, Zhenyang and Zhang, Xi. Hermitian harmonic maps into convex balls. *Canad. Math. Bull.*, 50(1):113–122, 2007.
- [Mo01] Mo, Xiaohuan. Harmonic maps from Finsler manifolds. *Illinois J. Math.*, 45(4):1331–1345, 2001.
- [MY05] Mo, Xiaohuan and Yang, Yunyan. The existence of harmonic maps from Finsler manifolds to Riemannian manifolds. *Sci. China Ser. A.*, 48(1):115–130, 2005.
- [Qiu17] Qiu, Hongbing. The heat flow of V -harmonic maps from complete manifolds into regular balls. *Proc. Amer. Math. Soc.*, 145(5):2271–2280, 2017.
- [Qiu22a] Qiu, Hongbing. Rigidity of symplectic translating solitons. *J. Geom. Anal.*, 32(11):Paper No. 284, 19, 2022.
- [Qiu22b] Qiu, Hongbing. A Bernstein type result of translating solitons. *Calc. Var. Partial Differential Equations*, 61(6):Paper No. 228, 9, 2022.
- [QS22] Qiu, Hongbing and Sun, Linlin. Rigidity theorems of spacelike entire self-shrinking graphs in the pseudo-Euclidean space. *J. Funct. Anal.*, 281(9):Paper No. 109189, 24, 2021.

- [SZ06] Souplet, Philippe and Zhang, Qi S. Sharp gradient estimate and Yau's Liouville theorem for the heat equation on noncompact manifolds. *Bull. London Math. Soc.*, 38(6):1045–1053, 2006.
- [Yau75] Shing Tung Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.
- [Zha12] Zhang, Xi. Hermitian harmonic maps between almost Hermitian manifolds. *Recent developments in geometry and analysis*, 485–493, Adv. Lect. Math. (ALM), 23, Int. Press, Somerville, MA, 2012.

HAN LUO: SCHOOL OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF SCIENCE AND TECHNOLOGY, NANJING 210094, JIANGSU, PEOPLE'S REPUBLIC OF CHINA
Email address: 19110180023@fudan.edu.cn

WEIKE YU: SCHOOL OF MATHEMATICAL SCIENCES, MINISTRY OF EDUCATION KEY LABORATORY OF NSLSCS, NANJING NORMAL UNIVERSITY, NANJING, 210023, JIANGSU, PEOPLE'S REPUBLIC OF CHINA
Email address: wkyu2018@outlook.com

XI ZHANG: SCHOOL OF MATHEMATICS AND STATISTICS, NANJING UNIVERSITY OF SCIENCE AND TECHNOLOGY, NANJING 210094, JIANGSU, PEOPLE'S REPUBLIC OF CHINA
Email address: mathzx@ustc.edu.cn