# A ZERO-DENSITY ESTIMATE FOR *L*-FUNCTIONS ASSOCIATED WITH GL(3) HECKE–MAASS CUSP FORMS

QINGFENG SUN AND HUI WANG

ABSTRACT. In this paper, we establish an asymptotic formula for the twisted second moment of L-functions associated with Hecke–Maass cusp forms for  $SL(3, \mathbb{Z})$ , and further deduce a weighted zero-density estimate for these L-functions in the spectral aspect which may have important applications in other problems.

#### 1. INTRODUCTION

The zero-density estimates for automorphic L-functions play a significant role in many branches of analytic number theory. For example, it is very effective in some applications to remove the Riemann Hypothesis. Let L(s) be an automorphic L-functions which is absolutely convergent for  $\operatorname{Re}(s) > 1$ , has an analytic continuation to the whole complex plane, and satisfies a standard functional equation. For T > 0 and any fixed  $\sigma \ge 0$ , we define

$$N_L(\sigma;T) = \#\{\rho = 1/2 + \beta + i\gamma : L(\rho) = 0, \sigma \leq \beta, |\gamma| \leq T\}.$$

For the Riemann zeta function  $\zeta(s)$ , the most classical result can be tracked back to Ingham's famous work [22] on zero-density estimates, where it was proved that

$$N_{\zeta}(\sigma;T) \ll T^{3(1-\sigma)/(2-\sigma)} (\log T)^5.$$
 (1.1)

Later, number theorists such as Halász, Montgomery, Huxley, Jutila, Heath–Brown and Ivić improved this result (see, for example, [14, 15, 21, 23, 26, 41]). Recently, Guth and Maynard [13] proved that

$$N_{\mathcal{C}}(\sigma;T) \ll T^{15(1-\sigma)/(3+5\sigma)+\varepsilon}$$

Combining this with Ingham's estimate (1.1) when  $\sigma \leq 7/10$ , they further obtained (see [13, Eq. (1.4)])

$$N_{\zeta}(\sigma;T) \ll T^{30(1-\sigma)/13+\varepsilon}.$$

Here the exponent 30/13 improves on the previous exponent of 12/5 due to Huxley [21].

It is worth mentioning that Selberg [43, Theorem 1] made a major contribution to this problem by proving the uniform bound that for  $0 \le \sigma \le 1/2$ ,

$$N_{\zeta}(\sigma;T) \ll T^{1-\sigma/4} \log T. \tag{1.2}$$

Date: December 4, 2024.

<sup>2010</sup> Mathematics Subject Classification. 11F12, 11F66, 11F67, 11F72.

Key words and phrases. GL(3) L-functions, mollification, zero-density estimate, Kuznetsov trace formula.

Q. Sun was partially supported by the National Natural Science Foundation of China (Grant Nos. 12471005 and 12031008) and the Natural Science Foundation of Shandong Province (Grant No. ZR2023MA003).

The crucial feature of (1.2) is that the power of  $\log T$  matches the true order of the number of zeros of  $\zeta(s)$  up to height T, so that the estimate is still useful even  $\sigma$  is on the order of  $1/\log T$ . Later, Selberg [44, Theorem 4] proved an analogous zero-density estimate in the family of Dirichlet *L*-functions  $L(s, \chi)$  corresponding to the primitive character  $\chi$  modulo a large positive integer q.

For GL(2) *L*-functions, there have been fruitful results. In the case that f is a holomorphic cusp form of even integral weight k > 0 and level N with respect to the Hecke congruence subgroup  $\Gamma_0(N)$ , Luo [38] considered the case N = 1 and first established the analogous zerodensity estimate as Selberg [43] in *t*-aspect for L(s, f),

$$N_L(\sigma;T) \ll_f T^{1-\sigma/72} \log T,$$

uniformly for  $0 \le \sigma \le 1/2$ . Later, Kowalski–Michel [29] and Hough [19] extended Luo's result to the family of *L*-functions associated to holomorphic forms f of large prime level N and large weight k, respectively. In the case f is a Hecke–Maass cusp form Laplacian eigenvalue  $1/4 + t_f^2$ with  $t_f \ge 0$  for the full modular group, Liu–Streipel [36] recently established a weighted zerodensity result in the spectral aspect by means of the twisted second moment of *L*-functions L(s, f) at  $s = 1/2 + \sigma + i\tau$  with  $\sigma > 0$ .

In this paper, we aim to establish the analogous zero-density estimate for GL(3) *L*-functions in the spectral aspect. Let  $\phi_j$  run over an orthogonal basis of Hecke–Maass cusp forms for SL(3, Z). Each  $\phi_j$  has spectral parameter  $\nu_j = (\nu_{j,1}, \nu_{j,2}, \nu_{j,3})$ , the Langlands parameter  $\mu_j = (\mu_{j,1}, \mu_{j,2}, \mu_{j,3})$ , and the Hecke eigenvalues  $A_j(m, n)$  with  $A_j(1, 1) = 1$ . The *L*-function associated with  $\phi_j$  is defined by

$$L(s,\phi_j) = \sum_{n=1}^{\infty} \frac{A_j(1,n)}{n^s} = \prod_p \left( 1 - \frac{A_j(1,p)}{p^s} + \frac{A_j(p,1)}{p^{2s}} - \frac{1}{p^{3s}} \right)^{-1}, \quad \text{Re}(s) > 1.$$

As in Blomer–Buttcane [3], we consider the generic case, i.e., away from the Weyl chamber walls and away from self-dual forms. Let  $\mu_0 = (\mu_{0,1}, \mu_{0,2}, \mu_{0,3})$  and  $\nu_0 = (\nu_{0,1}, \nu_{0,2}, \nu_{0,3})$ , and satisfy the corresponding relations (2.2) and (2.1) below. We also assume

$$|\mu_{0,i}| \simeq |\nu_{0,i}| \simeq T := \|\mu_0\|, \quad 1 \le i \le 3,$$

where

$$\|\mu_0\| = \sqrt{\|\mu_{0,1}\|^2 + \|\mu_{0,2}\|^2 + \|\mu_{0,3}\|^2}$$

Throughout this paper, we let  $M = T^{\theta}$  for any fixed  $0 < \theta < 1$ . Define a test function  $h_{T,M}(\mu)$ (depending on  $\mu_0$ ) for  $\mu = (\mu_1, \mu_2, \mu_3) \in \mathbb{C}^3$  by

$$h_{T,M}(\mu) := P(\mu)^2 \left(\sum_{w \in \mathcal{W}} \psi\left(\frac{w(\mu) - \mu_0}{M}\right)\right)^2,$$

where

$$\psi(\mu) = \exp\left(\mu_1^2 + \mu_2^2 + \mu_3^2\right)$$

and

$$P(\mu) = \prod_{0 \le n \le A} \prod_{k=1}^{3} \frac{\left(\nu_k - \frac{1}{3}(1+2n)\right)\left(\nu_k + \frac{1}{3}(1+2n)\right)}{|\nu_{0,k}|^2}$$

for some fixed large A > 0. Here

$$\mathcal{W} := \left\{ I, \ w_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \ w_3 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \ w_4 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \ w_5 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \ w_6 = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}$$

is the Weyl group of  $SL(3,\mathbb{R})$ . The function  $h_{T,M}(\mu)$  has the localizing effect at a ball of radius M about  $w(\mu_0)$  for each  $w \in \mathcal{W}$ , and other nice properties stated in §3.4 below. Then with the normalizing factor

$$\mathcal{N}_j := \|\phi_j\|^2 \prod_{k=1}^3 \cos\left(\frac{3}{2}\pi\nu_{j,k}\right),$$

our main theorem is the zero-density estimate of weighted version in the spectral aspect for the L-functions associated with GL(3) Hecke–Maass forms.

**Theorem 1.1.** For  $\sigma > 0$  and H > 0, we let

$$N(\sigma, H; \phi_j) := \# \{ \rho = 1/2 + \beta + i\gamma : L(\rho, \phi_j) = 0, \beta > \sigma, |\gamma| < H \}$$

and

$$N(\sigma, H) := \frac{1}{\mathcal{H}} \sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} N(\sigma, H; \phi_j).$$
(1.3)

Let  $2/\log T < \sigma < 1/2$ . For some sufficiently small  $\delta$ ,  $\theta_1 > 0$ , we have

$$N(\sigma, H) \ll HT^{-\delta\sigma} \log T,$$

uniformly in  $3/\log T < H < T^{\theta_1}$ .

**Remark 1.** Theorem 1.1 shows that there are only very few *L*-functions with zeros of imaginary part less than  $T^{\theta_1}$  and real part greater than  $1/2 + C/\log T$ .

**Remark 2.** Note that it seems difficult to prove an analogous result without harmonic weights (see, e.g., [8]). However, this weighted zero-density estimate is useful for certain problems, such as removing the Generalized Riemann Hypothesis (GRH) in the work of Liu and Liu [35]. And, the detailed proof of this application will be presented in the thesis of the second author.

The main ingredient to prove Theorem 1.1 is the following asymptotic formula of the harmonic twisted second moment of  $L(s, \phi_j)$ .

**Theorem 1.2.** Let  $0 < \sigma \leq 1/2$ ,  $|\tau| < T^{1/4}$ , and  $\ell_1$ ,  $\ell_2 < T^{1/3}$  be square-free integers. Let the notations be as above and in § 3.4 below. Then for any  $\varepsilon > 0$  we have

$$\begin{split} &\sum_{j} \frac{h_{T,M}(\mu_{j})}{\mathcal{N}_{j}} A_{j}(\ell_{1},\ell_{2}) \left| L(1/2 + \sigma + i\tau,\phi_{j}) \right|^{2} \\ &= \zeta(1+2\sigma) \frac{1}{(\ell_{1}\ell_{2})^{1/2+\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \frac{1}{192\pi^{5}} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \mathrm{d}_{\operatorname{spec}}\mu \\ &+ \zeta(1-2\sigma) \frac{1}{(\ell_{1}\ell_{2})^{1/2-\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \frac{1}{192\pi^{5}} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^{3} \left(-\frac{\mu_{k}^{2}}{4\pi^{2}}\right)^{-\sigma} \mathrm{d}_{\operatorname{spec}}\mu \\ &+ O\left((\ell_{1}\ell_{2})^{-1/2+\sigma} T^{9/4-6\sigma} M^{2} + T^{2-3\sigma+\varepsilon} M^{2} + T^{83/42-41\sigma/21+\varepsilon} M^{2}(\ell_{1}\ell_{2})^{\vartheta}\right), \end{split}$$
(1.4)

where  $\vartheta$  is defined in (3.5) below.

Note that both sides of (1.4) are analytic in  $\sigma$  near 0. Letting  $\sigma \to 0$ , we have the following corollary.

**Corollary 1.3.** Let  $|\tau| < T^{1/4}$  and  $\ell_1$ ,  $\ell_2 < T^{1/3}$  be square-free integers. For any  $\varepsilon > 0$ , we have

$$\begin{split} &\sum_{j} \frac{h_{T,M}(\mu_{j})}{\mathcal{N}_{j}} A_{j}(\ell_{1},\ell_{2}) \left| L(1/2+i\tau,\phi_{j}) \right|^{2} \\ &= \frac{1}{192\pi^{5}} \frac{1}{(\ell_{1}\ell_{2})^{1/2}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \left( -\log(\ell_{1}\ell_{2})+2\gamma+\sum_{k=1}^{3}\log\left(-\frac{\mu_{k}^{2}}{4\pi^{2}}\right) \right) d_{\operatorname{spec}}\mu \\ &+ O\left((\ell_{1}\ell_{2})^{-1/2}T^{9/4}M^{2}+T^{2+\varepsilon}M^{2}+T^{83/42+\varepsilon}M^{2}(\ell_{1}\ell_{2})^{\vartheta}\right). \end{split}$$

The problem of moments of L-functions is another important topic in analytic number theory, with many applications, such as in the subconvexity problem for L-functions and for the non-vanishing of central L-values. And to obtain positive-proportional non-vanishing results, one typically employs the mollification method à la Selberg and studies the mollified first and second moments of L-values at the central point s = 1/2 (see, for example [1, 20, 27, 29-34, 37, 39, 45, 48], etc.).

In this paper, we also investigate the asymptotic formula of the mollified second moment of  $L(s, \phi_j)$ . Define the mollifier  $M(s, \phi_j)$  for  $L(s, \phi_j)$  by

$$M(s,\phi_j) := \sum_{\ell \leqslant L^2} \frac{A_j(1,\ell)}{\ell^s} x_\ell, \tag{1.5}$$

where

$$x_{\ell} := \mu(\ell) \frac{1}{2\pi i} \int_{(3)} \frac{(L^2/\ell)^u - (L/\ell)^u}{u^2} \frac{\mathrm{d}u}{\log L} = \begin{cases} \mu(\ell), & \ell \leq L, \\ \mu(\ell) \frac{\log(L^2/\ell)}{\log L}, & L < \ell \leq L^2, \\ 0, & \ell > L^2, \end{cases}$$
(1.6)

here  $\mu(\ell)$  is the Möbius function and  $L = T^{\delta}$  for some suitably small constant  $\delta > 0$ . Then we have the following important proposition.

**Proposition 1.4.** For sufficiently small  $\delta$ ,  $\theta_1 > 0$ , we have

$$\frac{1}{\mathcal{H}} \sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} \left| L(1/2 + \sigma + i\tau, \phi_j) M(1/2 + \sigma + i\tau, \phi_j) \right|^2 \le 1 + O(T^{-2\sigma\delta}), \tag{1.7}$$

uniformly for  $1/\log T \leq \sigma \leq 1$  and  $|\tau| < T^{\theta_1}$ , and here

$$\mathcal{H} := \frac{1}{192\pi^5} \int_{\text{Re}(\mu)=0} h_{T,M}(\mu) d_{\text{spec}}\mu.$$
 (1.8)

Moreover, for all  $\tau$ , we have

$$L(3/2 + i\tau, \phi_j)M(3/2 + i\tau, \phi_j) = 1 + O(T^{-2\delta}).$$

Structure of this paper. This paper is organized as follows. In §§ 2–4, we review some essential facts and useful tools, including GL(3) Hecke–Maass cusp forms and their *L*-functions, Kuznetsov trace formula and the argument principle. In § 5 and § 6, we provide the asymptotic formulas for the harmonic twisted and mollified second moment of  $L(s, \phi_j)$ , respectively, i.e., the detailed proofs of Theorem 1.2 and Proposition 1.4. In §7, we first give an application of Kuznetsov trace formula and then present the detailed proof of Theorem 1.1.

**Notation.** Through out the paper,  $\varepsilon$  is an arbitrarily small positive number and B is a sufficiently large positive number which may not be the same at each occurrence.

As usual,  $e(z) = \exp(2\pi i z) = e^{2\pi i z}$ . The symbol  $k \sim K$  means  $K < k \leq 2K$ . We write f = O(g) or  $f \ll g$  to mean  $|f| \leq Cg$  for some unspecified positive constant C. We denote  $f \approx g$  to mean that  $c_1g \leq |f| \leq c_2g$  for some positive constants  $c_1$  and  $c_2$ . The sign  $\overline{d}$  denotes the multiplicative inverse of  $d \pmod{c}$ .

### 2. GL(3) Hecke–Maass L-functions

In this section, we review the L-functions associated with GL(3) Hecke–Maass cusp forms and provide the corresponding approximate functional equations.

2.1. Hecke–Maass cusp forms and their *L*-functions. Let  $G = GL(3, \mathbb{R})$  with maximal compact subgroup  $K = O(3, \mathbb{R})$  and center  $Z \cong \mathbb{R}^{\times}$ . Let  $\mathbb{H}_3 = G/(K \cdot Z)$  be the generalized upper half-plane. For  $0 \leq c \leq \infty$ , let

$$\Lambda_c = \left\{ \mu = (\mu_1, \mu_2, \mu_3) \in \mathbb{C}^3 \ \middle| \ \mu_1 + \mu_2 + \mu_3 = 0, \ |\operatorname{Re}(\mu_k)| \le c, \ k = 1, 2, 3 \right\},\$$

and

$$\Lambda_c' = \left\{ \mu \in \Lambda_c \ \Big| \ \{-\mu_1, -\mu_2, -\mu_3\} = \{\overline{\mu_1}, \overline{\mu_2}, \overline{\mu_3}\} \right\}.$$

Here  $\mu$  is the Langlands parameter of a Hecke–Maass form  $\phi$  in  $L^2(SL(3,\mathbb{Z})\backslash\mathbb{H})$ . We will simultaneously use  $\mu$  and spectral parameters  $\nu = (\nu_1, \nu_2, \nu_3)$  coordinates, defined by

$$\nu_1 = \frac{1}{3}(\mu_1 - \mu_2), \quad \nu_2 = \frac{1}{3}(\mu_2 - \mu_3), \quad \nu_3 = \frac{1}{3}(\mu_3 - \mu_1).$$
(2.1)

And we have

$$\mu_1 = 2\nu_1 + \nu_2, \quad \mu_2 = \nu_2 - \nu_1, \quad \mu_3 = -\nu_1 - 2\nu_2.$$
 (2.2)

Let  $\phi$  be a Hecke–Maass cusp form for  $SL(3,\mathbb{Z})$  with Fourier coefficients  $A_{\phi}(m,n) \in \mathbb{C}$  for  $(m,n) \in \mathbb{N}^2$ . We have the following Hecke relation (see Goldfeld [11, Theorem 6.4.11])

$$A_{\phi}(m,1)A_{\phi}(1,n) = \sum_{d|(m,n)} A_{\phi}\left(\frac{m}{d},\frac{n}{d}\right).$$

$$(2.3)$$

The standard *L*-function of  $\phi$  is given by

$$L(s,\phi) := \sum_{n=1}^{\infty} \frac{A_{\phi}(1,n)}{n^s}, \text{ for } \operatorname{Re}(s) > 1.$$

The dual form of  $\phi$  is denoted by  $\phi$  with the Langlands parameter  $(-\mu_1, -\mu_2, -\mu_3)$  and the coefficients  $A_{\phi}(m, n) = \overline{A_{\phi}(m, n)} = A_{\phi}(n, m)$ . Then  $L(s, \phi)$  admits an analytic continuation to all  $s \in \mathbb{C}$  and satisfies the functional equation

$$\Lambda(s,\phi) = L(s,\phi) \prod_{j=1}^{3} \Gamma_{\mathbb{R}}(s-\mu_j) = L(1-s,\widetilde{\phi}) \prod_{j=1}^{3} \Gamma_{\mathbb{R}}(1-s+\mu_j) = \Lambda(1-s,\widetilde{\phi}),$$

where  $\Gamma_{\mathbb{R}}(s) = \pi^{-s/2} \Gamma(s/2)$ .

By [3], the Langlands parameter of  $\phi$  lies in  $\Lambda'_{5/14} \subset \Lambda_{5/14}$ , and the non-exceptional parameters lie in  $\Lambda'_0 = \Lambda_0$ .

2.2. Stirling's formula. By the Stirling asymptotic formula (see [42, §8.4, Eq. (4.03)]), for  $|\arg(z)| \leq \pi - \varepsilon$ ,  $|z| \gg 1$  and any  $\varepsilon > 0$ ,

$$\log \Gamma(z) = \left(z - \frac{1}{2}\right) \log z - z + \frac{1}{2} \log(2\pi) + \sum_{j=1}^{K_1} \frac{B_{2j}}{2j(2j-1)z^{2j-1}} + O_{K_1,\varepsilon}\left(\frac{1}{|z|^{2K_1+1}}\right),$$

where  $B_j$  is *j*-th Bernoulli number. Thus for  $z = \sigma_1 + it$ ,  $\sigma_1$  fixed and  $|t| \ge 2$ ,

$$\Gamma(\sigma_1 + it) = \sqrt{2\pi}(it)^{\sigma_1 - 1/2} e^{-\pi|t|/2} \left(\frac{|t|}{e}\right)^{it} \left(1 + \sum_{j=1}^{K_2} \frac{c_j}{t^j} + O_{\sigma_1, K_2, \varepsilon} \left(\frac{1}{|t|^{K_2 + 1}}\right)\right),$$

where the constant  $c_j$  depends on  $j, \sigma_1$  and  $\varepsilon$ .

2.3. The approximate functional equation. Let  $G(u) = (\cos \frac{\pi u}{A})^{-100A}$ , where A is a positive integer. Then G(u) is even and holomorphic inside the strip  $-\frac{A}{2} < \operatorname{Re}(u) < \frac{A}{2}$ . For  $\sigma > 0$  and  $\tau \in \mathbb{R}$ , we will use the following approximate functional equation of  $|L(1/2 + \sigma + i\tau, \phi_j)|^2$ .

**Lemma 2.1.** Let  $\phi_j$  be a GL(3) Maass form with Langlands parameters

$$\mu_j = (\mu_{j,1}, \mu_{j,2}, \mu_{j,3}) \in \Lambda'_{5/14}.$$

For  $s = 1/2 + \sigma + i\tau$  with  $0 < \sigma \leq 1/2$  and  $\tau \in \mathbb{R}$ , we have

$$|L(s,\phi_j)|^2 = \sum_r \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{A_j(m,n)}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} W_{\mu_j}(s,r^2mn) + \pi^{6\sigma} \sum_r \frac{1}{r^{1-2\sigma}} \sum_{m,n} \frac{A_j(m,n)}{(mn)^{1/2-\sigma}} \left(\frac{m}{n}\right)^{i\tau} \tilde{W}_{\mu_j}(s,r^2mn),$$

where

$$W_{\mu_j}(s,y) := \frac{1}{2\pi i} \int_{(3)} (\pi^3 y)^{-u} \prod_{k=1}^3 \frac{\Gamma\left(\frac{s+u-\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{s-\mu_{j,k}}{2}\right)} \frac{\Gamma\left(\frac{\overline{s}+u+\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_{j,k}}{2}\right)} G(u) \frac{\mathrm{d}u}{u},\tag{2.4}$$

and

$$\tilde{W}_{\mu_j}(s,y) := \frac{1}{2\pi i} \int_{(3)} (\pi^3 y)^{-u} \prod_{k=1}^3 \frac{\Gamma\left(\frac{1-s+u+\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_{j,k}}{2}\right)} \frac{\Gamma\left(\frac{1-\overline{s}+u-\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{s-\mu_{j,k}}{2}\right)} G(u) \frac{\mathrm{d}u}{u}.$$
(2.5)

Proof. See e.g., Iwaniec–Kowalski [24, §5.2].

Note that the sum in (1.3) is essentially supported on the generic forms which satisfy

$$|\mu_{j,k}| \approx |\nu_{j,k}| \approx T, \quad 1 \leqslant k \leqslant 3.$$

$$(2.6)$$

So we assume that  $\phi_j$  also satisfies the above relation. The functions  $W_{\mu_j}(s, y)$  and  $\tilde{W}_{\mu_j}(s, y)$  satisfy the following properties.

**Lemma 2.2.** Assume the condition (2.6), and let  $s = 1/2 + \sigma + i\tau$  with  $0 < \sigma \leq 1/2$  and  $|\tau| < T^{1/4}$ . For any  $\varepsilon > 0$ , we have

$$y^{i}W_{\mu_{j}}^{(i)}(s,y), \quad y^{i}\tilde{W}_{\mu_{j}}^{(i)}(s,y) \ll_{i,B} \left(1+\frac{y}{T^{3}}\right)^{-B}, \quad for \ any \ integer \ i \ge 0,$$
 (2.7)

$$W_{\mu_j}(s,y) = \frac{1}{2\pi i} \int_{\varepsilon-i\log T}^{\varepsilon+i\log T} (\pi^3 y)^{-u} \prod_{k=1}^3 \frac{\Gamma\left(\frac{s+u-\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{s-\mu_{j,k}}{2}\right)} \frac{\Gamma\left(\frac{\overline{s}+u+\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_{j,k}}{2}\right)} G(u) \frac{\mathrm{d}u}{u} + O_{\varepsilon,B}(y^{\varepsilon}T^{-B}),$$
(2.8)

$$\tilde{W}_{\mu_j}(s,y) = \frac{1}{2\pi i} \int_{\varepsilon-i\log T}^{\varepsilon+i\log T} (\pi^3 y)^{-u} \prod_{k=1}^3 \frac{\Gamma\left(\frac{1-s+u+\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_{j,k}}{2}\right)} \frac{\Gamma\left(\frac{1-\overline{s}+u-\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{s-\mu_{j,k}}{2}\right)} G(u) \frac{\mathrm{d}u}{u} + O_{\varepsilon,B}(y^{\varepsilon}T^{-B}),$$
(2.9)

and for  $y \ll T^3$ ,

$$W_{\mu_j}(s,y) = 1 + O_B\left(\left(\frac{y}{T^3}\right)^B\right),\tag{2.10}$$

$$\tilde{W}_{\mu_j}(s,y) = \prod_{k=1}^3 \frac{\Gamma\left(\frac{1-s+\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_{j,k}}{2}\right)} \frac{\Gamma\left(\frac{1-\overline{s}-\mu_{j,k}}{2}\right)}{\Gamma\left(\frac{s-\mu_{j,k}}{2}\right)} + O_B\left(\left(\frac{y}{T^3}\right)^B\right),\tag{2.11}$$

provided for any positive integer B < A/2.

*Proof.* Shifting the line of integration to  $\operatorname{Re}(u) = B$  and using Stirling's formula, we can get (2.7). For (2.8), we move the line of integration in (2.4) to  $\operatorname{Re}(u) = \varepsilon$  without passing any poles, and bound the integral for  $|\operatorname{Im}(u)| \leq \log T$  using Stirling's formula and the exponential decay of G(u). The main term in (2.10) is the residue from the pole at u = 0 when we shift the integral line to  $\operatorname{Re}(u) = -B$ , while the residues at the other poles are exponentially small in view of (2.6). Similarly, we can get (2.9) and (2.11), respectively.

By Stirling's formula, if  $|u| \ll |z|^{1/2}$ , then

$$\frac{\Gamma(z+u)}{\Gamma(z)} = z^u \left( 1 + \sum_{n=1}^{N_1} \frac{P_n(u)}{z^n} + O\left(\frac{(1+|u|)^{2N_1+2}}{|z|^{N_1+1}}\right) \right),$$
(2.12)

for certain polynomials  $P_n(u)$  of degree 2n. For  $\mu_j \in \Lambda'_{1/2}$  and  $\tau < T^{1/4}$ , we have

$$W_{\mu_j}(1/2 + \sigma + i\tau, y) = \frac{1}{2\pi i} \int_{(3)} (\pi^3 y)^{-u} \prod_{k=1}^3 \prod_{\pm} \left(\frac{1/2 + \sigma \mp i\tau \pm \mu_{j,k}}{2}\right)^{u/2} \cdot \left(1 + \sum_{n=1}^{N_1} \frac{2^n P_n(u)}{(1/2 + \sigma \mp i\tau \pm \mu_{j,k})^n} + O\left(\frac{(1 + |u|)^{2N_1 + 2}}{|\mu_{j,k}|^{N_1 + 1}}\right)\right) G(u) \frac{\mathrm{d}u}{u}, \quad (2.13)$$

and

$$\begin{split} \tilde{W}_{\mu_j}(1/2 + \sigma + i\tau, y) &= \frac{1}{2\pi i} \int_{(3)} (\pi^3 y)^{-u} \prod_{k=1}^3 \prod_{\pm} \left( \frac{1/2 + \sigma \mp i\tau \pm \mu_{j,k}}{2} \right)^{-\sigma + u/2} \\ &\cdot \left( 1 + \sum_{n=1}^{N_2} \frac{2^n P_n(u)}{(1/2 + \sigma \mp i\tau \pm \mu_{j,k})^n} + O\left(\frac{(1 + |u|)^{2N_2 + 2}}{|\mu_{j,k}|^{N_2 + 1}}\right) \right) G(u) \frac{\mathrm{d}u}{u}, \end{split}$$

for a different collection of polynomials  $P_n(u)$ .

### 3. The Kuznetsov trace formula

In this section, we state the Kuznetsov trace formula for the particular test function  $h_{T,M}$  described in §1 above.

### 3.1. The minimal Eisenstein series and its Fourier coefficients. Let

$$U_3 = \left\{ \begin{pmatrix} 1 & * & * \\ 0 & 1 & * \\ 0 & 0 & 1 \end{pmatrix} \right\} \cap \mathrm{SL}(3, \mathbb{Z}).$$

For  $z \in \mathbb{H}_3$  and  $\operatorname{Re}(\nu_1), \operatorname{Re}(\nu_2)$  sufficiently large, we define the minimal Eisenstein series

$$E(z;\mu) := \sum_{\gamma \in U_3 \setminus \mathrm{SL}(3,\mathbb{Z})} I_{\nu_1,\nu_2}(\gamma z),$$

where

$$I_{\nu_1,\nu_2}(z) := y_1^{1+\nu_1+2\nu_2} y_2^{1+2\nu_1+\nu_2},$$

and

$$z = \begin{pmatrix} y_1 y_2 & y_1 x_2 & x_3 \\ 0 & y_1 & x_1 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & x_2 & x_3 \\ 0 & 1 & x_1 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_1 y_2 & 0 & 0 \\ 0 & y_1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$
(3.1)

with  $y_1, y_2 > 0$  and  $x_1, x_2, x_3 \in \mathbb{R}$ .  $E(z; \mu)$  has Langlands parameter  $\mu$  and admits a meromorphic continuation in  $\nu_1$  and  $\nu_2$ . The Fourier coefficients  $A_{\mu}(m_1, m_2)$  are defined by (see Goldfeld [11, Theorem 10.8.6])

$$A_{\mu}(m,1) = \sum_{d_1d_2d_3 = m} d_1^{\mu_1} d_2^{\mu_2} d_3^{\mu_3},$$

and satisfy the symmetry and Hecke relation (see Goldfeld [11, Theorem 6.4.11])

$$A_{\mu}(m,1) = \overline{A_{\mu}(1,m)},$$
  

$$A_{\mu}(m_1,m_2) = \sum_{d \mid (m_1,m_2)} \mu(d) A_{\mu}(m_1/d,1) A_{\mu}(1,m_2/d).$$
(3.2)

Hence we have

$$|A_{\mu}(m_1, m_2)| \ll_{\varepsilon} (m_1 m_2)^{\varepsilon}, \quad \text{if } \mu \in (i\mathbb{R})^3.$$

The *L*-function associated with  $E(z; \mu)$  is given by

$$L(s, E(z; \mu)) = \sum_{m \ge 1} \frac{A_{\mu}(1, m)}{m^s} = \zeta(s + \mu_1)\zeta(s + \mu_2)\zeta(s + \mu_3),$$
(3.3)

for  $\operatorname{Re}(s) > 1$ . In order to state the Kuznetsov trace formula in the §3.4, we introduce the normalized factor for the minimal Eisenstein series is

$$\mathcal{N}_{\mu} = \frac{1}{16} \prod_{k=1}^{3} |\zeta(1+3\nu_k)|^2$$

corresponding to the minimal Eisenstein series  $E(z; \mu)$ , where

$$\mu = (\mu_1, \mu_2, \mu_3).$$

Recall that (see [46, Eq. (3.11.8)]) we have

$$\frac{1}{\zeta(1+it)} \ll \log(1+|t|),$$

which implies that

$$\frac{1}{\mathcal{N}_{\mu}} \ll \prod_{k=1}^{3} \log^2(1+|\nu_k|).$$

### 3.2. The maximal Eisenstein series and its Fourier coefficients. Let

$$P_{2,1} = \left\{ \begin{pmatrix} * & * & * \\ * & * & * \\ 0 & 0 & * \end{pmatrix} \right\} \cap \mathrm{SL}(3,\mathbb{Z}).$$

Let  $\mu \in \mathbb{C}$  have sufficiently large real part, and let  $g : \mathrm{SL}(2,\mathbb{Z}) \setminus \mathbb{H}_2 \to \mathbb{C}$  be a Hecke–Maass cusp form with ||g|| = 1, Langlands parameter  $\mu_g \in i\mathbb{R}$  and Hecke eigenvalue  $\lambda_g(m)$ . The maximal Eisenstein series twisted by a Maass form g is defined by

$$E(z;\mu;g) := \sum_{\gamma \in P_{2,1} \setminus \mathrm{SL}(3,\mathbb{Z})} \det(\gamma z)^{1/2+\mu} g(\mathfrak{m}_{P_{2,1}}(\gamma z)),$$

where z is defined as in (3.1), and

$$\mathfrak{m}_{P_{2,1}}: \mathbb{H}_3 \to \mathbb{H}_2, \quad \begin{pmatrix} y_1 y_2 & y_1 x_2 & x_3 \\ & y_1 & x_1 \\ & & 1 \end{pmatrix} \mapsto \begin{pmatrix} y_2 & x_2 \\ & 1 \end{pmatrix}$$

is the restriction to the upper left corner. It has a meromorphic continuation in  $\mu$ . The Fourier coefficients are determined by

$$B_{\mu,g}(m,1) = \sum_{d_1d_2=m} \lambda_g(d_1) d_1^{\mu} d_2^{-2\mu},$$

and satisfy the symmetry and Hecke relation as (3.2) above (see Goldfeld [11, Proposition 10.9.3 and Theorem 6.4.11])

$$B_{\mu,g}(m,1) = \overline{B_{\mu,g}(1,m)},$$
  

$$B_{\mu,g}(m_1,m_2) = \sum_{d \mid (m_1,m_2)} \mu(d) B_{\mu,g}(m_1/d,1) B_{\mu,g}(1,m_2/d).$$
(3.4)

Recall that the current best known estimate towards the Ramanujan–Petersson conjecture for Fourier coefficients of GL(2) Maass cusp forms is due to Kim–Sarnak [28, Appendix 2],

$$|\lambda_g(n)| \ll_{\varepsilon} n^{\vartheta + \varepsilon}, \quad n \in \mathbb{N}, \tag{3.5}$$

here  $\vartheta = 7/64$ . The Ramanujan–Petersson conjecture predicts  $\vartheta = 0$ . Hence we have

$$|B_{\mu,g}(m_1,m_2)| \ll_{\varepsilon} (m_1m_2)^{\vartheta+\varepsilon}, \quad \text{if } \mu \in i\mathbb{R}.$$

The *L*-function associated with  $E(z; \mu; g)$  is given by

$$L(s, E(z; \mu; g)) = \sum_{m \ge 1} \frac{B_{\mu,g}(1, m)}{m^s} = \zeta(s - 2\mu)L(s + \mu, g),$$
(3.6)

for sufficiently large  $\operatorname{Re}(s)$ . Note that L(s,g) is the Hecke–Maass *L*-function associated with g with real Fourier coefficients  $\lambda_q(n)$ . The complete *L*-function is defined by

$$\Lambda(s, E(z; \mu; g)) = \prod_{k=1}^{3} \Gamma_{\mathbb{R}}(s + \mu'_k) L(s, E(z; \mu; g)) = \Lambda(1 - s, E(z; -\mu; g)),$$
(3.7)

where

$$\mu'_1 = \mu + \mu_g, \quad \mu'_2 = \mu - \mu_g, \quad \mu'_3 = -2\mu.$$
 (3.8)

We also introduce the normalizing factor

$$\mathcal{N}_{\mu,g} = 8L(1, \mathrm{Ad}^2 g) |L(1+3\mu, g)|^2,$$

where  $L(s, \mathrm{Ad}^2 g)$  is the adjoint square L-function of g. And we have the lower bounds

$$L(1, \mathrm{Ad}^{2}g) \gg (1 + |\mu_{g}|)^{-\varepsilon}, \quad L(1 + it, g) \gg (1 + |t| + |\mu_{g}|)^{-\varepsilon}$$

which follow from [17, 18, 25] and [10]. Therefore, for  $\mu \in i\mathbb{R}$ , it follows that

$$\frac{1}{\mathcal{N}_{\mu,g}} \ll (1 + |\mu| + |\mu_g|)^{\varepsilon}.$$
(3.9)

3.3. The Kloosterman sums. For  $n_1$ ,  $n_2$ ,  $m_1$ ,  $m_2$ ,  $D_1$ ,  $D_2 \in \mathbb{N}$ , we need the relevant Kloosterman sums

$$\tilde{S}(n_1, n_2, m_1; D_1, D_2) := \sum_{\substack{C_1 \ ( \bmod D_1), C_2 \ ( \bmod D_2) \\ (C_1, D_1) = (C_2, D_2/D_1) = 1}} e\left(n_2 \frac{C_1 C_2}{D_1} + m_1 \frac{C_2}{D_2/D_1} + n_1 \frac{C_1}{D_1}\right)$$

for  $D_1|D_2$ , and

 $S(n_1, m_2, m_1, n_2; D_1, D_2)$ 

$$:= \sum_{\substack{B_1, C_1 \pmod{D_1} \\ B_2, C_2 \pmod{D_2} \\ D_1 C_2 + B_1 B_2 + D_2 C_1 \equiv 0 \pmod{D_1 D_2} \\ (B_j, C_j, D_j) = 1}} e\left(\frac{n_1 B_1 + m_1 (Y_1 D_2 - Z_1 B_2)}{D_1} + \frac{m_2 B_2 + n_2 (Y_2 D_1 - Z_2 B_1)}{D_2}\right),$$

where  $Y_j B_j + Z_j C_j \equiv 1 \pmod{D_j}$  for j = 1, 2. We have the standard (Weil-type) bounds

$$\tilde{S}(n_1, n_2, m_1; D_1, D_2) \ll ((m_1, D_2/D_1)D_1^2, (n_1, n_2, D_1)D_2)(D_1D_2)^{\varepsilon},$$
(3.10)

and

$$S(n_1, m_2, m_1, n_2; D_1, D_2) \ll (D_1 D_2)^{1/2 + \varepsilon} ((D_1, D_2)(m_1 n_1, [D_1, D_2])(m_2 n_2, [D_1, D_2]))^{1/2 + \varepsilon}.$$

The first bound is due to Larsen [5, Appendix], and the second is due to Stevens [6, Theorem 2].

3.4. The Kuznetsov trace formula. Define the spectral measure on the hyperplane  $\mu_1 + \mu_2 + \mu_3 = 0$  by

$$d_{\rm spec}\mu = {\rm spec}(\mu)d\mu,$$

where

$$\operatorname{spec}(\mu) := \prod_{k=1}^{3} \left( 3\nu_k \tan\left(\frac{3\pi}{2}\nu_k\right) \right) \quad \text{and} \quad \mathrm{d}\mu = \mathrm{d}\mu_1 \mathrm{d}\mu_2 = \mathrm{d}\mu_2 \mathrm{d}\mu_3 = \mathrm{d}\mu_1 \mathrm{d}\mu_3$$

Following [7, Theorems 2 & 3], we define the following integral kernels in terms of Mellin–Barnes representations. For  $s \in \mathbb{C}$ ,  $\mu \in \Lambda_{\infty}$ , define the meromorphic function

$$\tilde{G}^{\pm}(s,\mu) := \frac{\pi^{-3s}}{12288\pi^{7/2}} \left( \prod_{k=1}^{3} \frac{\Gamma(\frac{1}{2}(s-\mu_k))}{\Gamma(\frac{1}{2}(1-s+\mu_k))} \pm i \prod_{k=1}^{3} \frac{\Gamma(\frac{1}{2}(1+s-\mu_k))}{\Gamma(\frac{1}{2}(2-s+\mu_k))} \right),$$
(3.11)

and for  $s = (s_1, s_2) \in \mathbb{C}^2$ ,  $\mu \in \Lambda_{\infty}$ , define the meromorphic function

$$G(s,\mu) := \frac{1}{\Gamma(s_1 + s_2)} \prod_{k=1}^{3} \Gamma(s_1 - \mu_k) \Gamma(s_2 + \mu_k).$$

The latter is essentially the double Mellin transform of the GL(3) Whittaker function. We also define the following trigonometric functions

$$S^{++}(s,\mu) := \frac{1}{24\pi^2} \prod_{k=1}^3 \cos\left(\frac{3}{2}\pi\nu_k\right),$$

$$S^{+-}(s,\mu) := -\frac{1}{32\pi^2} \frac{\cos(\frac{3}{2}\pi\nu_2)\sin(\pi(s_1-\mu_1))\sin(\pi(s_2+\mu_2))\sin(\pi(s_2+\mu_3))}{\sin(\frac{3}{2}\pi\nu_1)\sin(\frac{3}{2}\pi\nu_3)\sin(\pi(s_1+s_2))},$$

$$S^{-+}(s,\mu) := -\frac{1}{32\pi^2} \frac{\cos(\frac{3}{2}\pi\nu_1)\sin(\pi(s_1-\mu_1))\sin(\pi(s_1-\mu_2))\sin(\pi(s_2+\mu_3))}{\sin(\frac{3}{2}\pi\nu_2)\sin(\frac{3}{2}\pi\nu_3)\sin(\pi(s_1+s_2))},$$

$$S^{--}(s,\mu) := \frac{1}{32\pi^2} \frac{\cos(\frac{3}{2}\pi\nu_3)\sin(\pi(s_1-\mu_2))\sin(\pi(s_2+\mu_2))}{\sin(\frac{3}{2}\pi\nu_2)\sin(\frac{3}{2}\pi\nu_1)}.$$

For  $y \in \mathbb{R} \setminus \{0\}$  with  $\operatorname{sgn}(y) = \epsilon$ , let

$$K_{w_4}(y;\mu) := \int_{-i\infty}^{i\infty} |y|^{-u} \tilde{G}^\epsilon(u,\mu) \frac{\mathrm{d}u}{2\pi i}.$$
(3.12)

Here we quote the result of [3, Lemma 4].

**Lemma 3.1.** For  $y \in \mathbb{R} \setminus \{0\}$  and  $\mu \in \Lambda_0$ , we have

$$K_{w_4}(y;\mu) = \frac{1}{3072\pi^5} \int_0^\infty J_{\mu_1-\mu_2}^-(2\sqrt{u}) \left(\frac{\pi^3|y|}{u^{3/2}}\right)^{-\mu_3} \exp\left(-\frac{2i\pi^3 y}{u}\right) \frac{\mathrm{d}u}{u} + \frac{1}{3072\pi^5} \int_0^\infty \tilde{K}_{\mu_1-\mu_2}(2\sqrt{u}) \left(\frac{\pi^3|y|}{u^{3/2}}\right)^{-\mu_3} \exp\left(\frac{2i\pi^3 y}{u}\right) \frac{\mathrm{d}u}{u},$$
(3.13)

where

$$\tilde{K}_{it}(x) = \int_{-\infty}^{\infty} \cos(2x \sinh v) \exp(itv) dv,$$
$$J_{it}^{-}(x) = \int_{-\infty}^{\infty} \cos(2x \cosh v) \exp(itv) dv,$$

for  $t \in \mathbb{R}$ , x > 0.

**Remark 3.** As in [3, §5], the integrals in (3.13) are not absolutely convergent at 0, but since  $\exp(\pm 2i\pi^3 y/u)$  is highly oscillating in a neighborhood of u = 0, they exist in a Riemann sense, and the portion 0 < u < 1 can be made absolutely convergent after partial integration.

**Remark 4.** For the Bessel functions, we have the uniform bounds (see [3, Eq. (4.13)])

$$\frac{\partial^k}{\partial x^k} \tilde{K}_{it}(x), \quad \frac{\partial^k}{\partial x^k} J_{it}^-(x) \ll_k \left(1 + \frac{|t|}{x}\right)^k, \tag{3.14}$$

for  $|t|, x \ge 1$  and non-negative integer k.

For 
$$y = (y_1, y_2) \in (\mathbb{R} \setminus \{0\})^2$$
 with  $\operatorname{sgn}(y_1) = \epsilon_1$ ,  $\operatorname{sgn}(y_2) = \epsilon_2$ , let  
 $K_{w_6}^{\epsilon_1, \epsilon_2}(y; \mu) := \int_{-i\infty}^{i\infty} \int_{-i\infty}^{i\infty} |4\pi^2 y_1|^{-s_1} |4\pi^2 y_2|^{-s_2} G(s, \mu) S^{\epsilon_1, \epsilon_2}(s, \mu) \frac{\mathrm{d}s_1 \mathrm{d}s_2}{(2\pi i)^2}.$ 

We generally follow the Barnes integral convention that the contour should pass to the right of all of the poles of the Gamma functions in the form  $\Gamma(s_j + a)$  and to the left of all of the poles of the Gamma functions \* in the form  $\Gamma(a - s_j)$ . Moreover, we choose the contour such that all integrals are absolutely convergent, which can always be arranged by shifting the unbounded part appropriately.

Now we can state the Kuznetsov trace formula in the version of Buttcane [7, Theorems 2–4].

**Lemma 3.2.** Let  $n_1, n_2, m_1, m_2 \in \mathbb{N}$  and let h be a function that is holomorphic on  $\Lambda_{1/2+\delta}$  for some  $\delta > 0$ , symmetric under the Weyl group  $\mathcal{W}$ , of rapid decay when  $|\operatorname{Im}(\mu_j)| \to \infty$ , and satisfies

$$h(3\nu_i \pm 1) = 0, \quad j = 1, 2, 3.$$

Then we have

$$\mathcal{C} + \mathcal{E}_{\min} + \mathcal{E}_{\max} = \Delta + \Sigma_4 + \Sigma_5 + \Sigma_6,$$

where

$$\begin{aligned} \mathcal{C} &:= \sum_{j} \frac{h(\mu_{j})}{\mathcal{N}_{j}} \overline{A_{j}(m_{1}, m_{2})} A_{j}(n_{1}, n_{2}), \\ \mathcal{E}_{\min} &:= \frac{1}{24(2\pi i)^{2}} \iint_{\operatorname{Re}(\mu)=0} \frac{h(\mu)}{\mathcal{N}_{\mu}} \overline{A_{\mu}(m_{1}, m_{2})} A_{\mu}(n_{1}, n_{2}) d\mu_{1} d\mu_{2}, \\ \mathcal{E}_{\max} &:= \sum_{g} \frac{1}{2\pi i} \int_{\operatorname{Re}(\mu)=0} \frac{h(\mu + \mu_{g}, \mu - \mu_{g}, -2\mu)}{\mathcal{N}_{\mu,g}} \overline{B_{\mu,g}(m_{1}, m_{2})} B_{\mu,g}(n_{1}, n_{2}) d\mu, \end{aligned}$$

<sup>\*</sup>Such Gamma functions may occur through the functional equation  $(\sin(\pi(s_1 + s_2))\Gamma(s_1 + s_2))^{-1} = \frac{1}{\pi}\Gamma(1 - s_1 - s_2).$ 

and

$$\begin{split} \Delta &:= \delta_{n_1,m_1} \delta_{n_2,m_2} \frac{1}{192\pi^5} \int_{\operatorname{Re}(\mu)=0} h(\mu) d_{\operatorname{spec}} \mu, \\ \Sigma_4 &:= \sum_{\epsilon = \pm 1} \sum_{\substack{D_2 \mid D_1 \\ m_2 D_1 = n_1 D_2^2}} \frac{\tilde{S}(-\epsilon n_2, m_2, m_1; D_2, D_1)}{D_1 D_2} \Phi_{w_4} \left(\frac{\epsilon m_1 m_2 n_2}{D_1 D_2}\right), \\ \Sigma_5 &:= \sum_{\epsilon = \pm 1} \sum_{\substack{D_1 \mid D_2 \\ m_1 D_2 = n_2 D_1^2}} \frac{\tilde{S}(\epsilon n_1, m_1, m_2; D_1, D_2)}{D_1 D_2} \Phi_{w_5} \left(\frac{\epsilon n_1 m_1 m_2}{D_1 D_2}\right), \\ \Sigma_6 &:= \sum_{\epsilon_1, \epsilon_2 = \pm 1} \sum_{D_1, D_2} \frac{S(\epsilon_2 n_2, \epsilon_1 n_1, m_1, m_2; D_1, D_2)}{D_1 D_2} \Phi_{w_6} \left(-\frac{\epsilon_2 m_1 n_2 D_2}{D_1^2}, -\frac{\epsilon_1 m_2 n_1 D_1}{D_2^2}\right), \end{split}$$

with

$$\Phi_{w_4}(y) := \int_{\operatorname{Re}(\mu)=0} h(\mu) K_{w_4}(y;\mu) \mathrm{d}_{\operatorname{spec}}\mu, 
\Phi_{w_5}(y) := \int_{\operatorname{Re}(\mu)=0} h(\mu) K_{w_4}(-y;-\mu) \mathrm{d}_{\operatorname{spec}}\mu, 
\Phi_{w_6}(y_1,y_2) := \int_{\operatorname{Re}(\mu)=0} h(\mu) K_{w_6}^{\operatorname{sgn}(y_1),\operatorname{sgn}(y_2)}((y_1,y_2);\mu) \mathrm{d}_{\operatorname{spec}}\mu.$$
(3.15)

The function  $h_{T,M}(\mu)$  localizes at a ball of radius M about  $w(\mu_0)$  for each  $w \in \mathcal{W}$ . We have

$$\mathscr{D}_k h_{T,M}(\mu) \ll_k M^{-k}, \tag{3.16}$$

for any differential operator  $\mathscr{D}_k$  of order k, which we use frequently when we integrate by parts, and sufficiently many differentiations can save arbitrarily many powers of T. Moreover, by trivial estimate, we have

$$\int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \mathrm{d}_{\operatorname{spec}} \mu \approx T^3 M^2.$$
(3.17)

3.5. The weight functions. For the weight functions, we will need the following results in Blomer–Buttcane [3, Lemma 1, Lemmas 8 & 9].

**Lemma 3.3.** For some large enough constant B > 0, we have

$$\Phi_{w_4}(y) \ll |y|^{1/10} T^B, \quad \Phi_{w_5}(y) \ll |y|^{1/10} T^B, \quad \Phi_{w_6}(y_1, y_2) \ll |y_1 y_2|^{3/5} T^B.$$

**Remark 5.** As in [3, §3.6], Lemma 3.3 shows more quantitatively that with our particular choice of h we can truncate the  $D_1$ ,  $D_2$ -sums at  $D_1$ ,  $D_2 \ll T^B$  for some sufficiently large B at the cost of an error term  $O(T^{-1000})$ , provided that  $n_1$ ,  $n_2$ ,  $m_1$ ,  $m_2 \ll T^{10}$ , say.

**Lemma 3.4.** (1) If  $0 < |y| \leq T^{3-\varepsilon}$ , then for any constant B > 0, we have

$$\Phi_{w_4}(y) \ll_{\varepsilon,B} T^{-B}.$$

(2) If  $T^{3-\varepsilon} < |y|$ , then

$$|y|^k \Phi_{w_4}^{(k)}(y) \ll_{k,\varepsilon} T^{3+2\varepsilon} M^2 (T+|y|^{1/3})^k.$$

**Lemma 3.5.** Let  $\Upsilon := \min\{|y_1|^{1/3}|y_2|^{1/6}, |y_1|^{1/6}|y_2|^{1/3}\}.$ (1) If  $\Upsilon \ll T^{1-\varepsilon}$ , then we have

$$\Phi_{w_6}(y_1, y_2) \ll T^{-B}.$$

(2) If  $\Upsilon \gg T^{1-\varepsilon}$ , then we have

$$\begin{aligned} |y_1|^{k_1} |y_2|^{k_2} \frac{\partial^{k_1}}{\partial y_1^{k_1}} \frac{\partial^{k_2}}{\partial y_2^{k_2}} \Phi_{w_6}(y_1, y_2) \\ \ll_{k_1, k_2, \varepsilon} T^{3+2\varepsilon} M^2 (T + |y_1|^{1/2} + |y_1|^{1/3} |y_2|^{1/6})^{k_1} (T + |y_2|^{1/2} + |y_2|^{1/3} |y_1|^{1/6})^{k_2}. \end{aligned}$$

**Remark 6.** In the proof of Lemmas 3.3–3.5, the only two properties of  $h_{T,M}(\mu)$  used are (3.16) and (3.17). Here, we remark that  $h_i(\mu)$  also satisfies these two inequalities by using properties of  $W_{\mu,N_1}$  or  $\tilde{W}_{\mu,N_2}$  (see §5 below). Thus,  $\Phi_{i,w_4}(y)$ ,  $\Phi_{i,w_5}(y)$  and  $\Phi_{i,w_6}(y)$  defined as in (3.15) by using the test function  $h_i(\mu)$ , also satisfy the corresponding bounds in Lemmas 3.3–3.5, since the difference that we use (3.16) and (3.17) instead of [3, (3.7) and (3.8)] has no influence.

Moreover, we also need the following lemma, which gives some strong bounds on the  $\Phi_w$ functions.

**Lemma 3.6.** (1) For 
$$y_1, y_2 \in (T^{-100}, T^{100})$$
,  $\epsilon_1, \epsilon_2 \in \{\pm 1\}$ , we have  $\Phi_{w_6}(\epsilon_1 y_1, \epsilon_2 y_2) \ll T^{3+\varepsilon}$ .  
(2) For  $y \in (T^{-100}, T^{100})$ ,  $\epsilon \in \{\pm 1\}$ , we have  $\Phi_{w_4}(\epsilon y) \ll T^{3+\varepsilon}$ .

*Proof.* The proof closely follows [8, Proposition 4.1], which we give here for completeness. For the completely supported test function  $h_{T,M}$  described above, we define

$$\check{h}_{T,M}(v_1, v_2) = \frac{3}{T^3} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) v_1^{-\nu_1} v_2^{-\mu_2} \mathrm{d}_{\operatorname{spec}} \mu.$$

The support of  $h_{T,M}(\mu)$  allows us to remove the tangents from spec( $\mu$ ) at a negligible cost, and integration by parts in  $\check{h}_{T,M}(v_1, v_2)$  constrains  $v_1, v_2$  to a region  $v_i = 1 + O(M^{-1}T^{\varepsilon})$ . Then we have the trivial bound  $\check{h}_{T,M}(v_1, v_2) \ll M^2$ . Now for  $y_1, y_2 \in (T^{-100}, T^{100}), \epsilon_1, \epsilon_2 \in \{\pm 1\}$ , [8, Lemma 4.2] (using the Weyl-invariance of

 $h_{T,M}(\mu)$  implies

$$\begin{split} \Phi_{w_6}(\epsilon_1 y_1, \epsilon_2 y_2) &:= \frac{T^3}{8192\pi^4} \sum_{\substack{d \in \{0,1\}^2\\\eta \in \{\pm 1\}^3}} (\eta_1 \eta_2 \eta_3 \epsilon_1)^{d_1} \epsilon_2^{d_2} \int_0^\infty \int_0^\infty \check{h}_{T,M} \left( \frac{y_2 z_1^{3/2}}{y_1}, \frac{y_1}{y_2 z_2^{3/2}} \right) \\ &\times \operatorname{sgn}(1 + \eta_1 z_1)^{d_1 + d_2} \operatorname{sgn}(1 + \eta_2 z_2)^{d_1 + d_2} \operatorname{sgn}(1 + \eta_3 z_3)^{d_1 + d_2} \\ &\times \left( \frac{2}{\pi} K_0 \left( 4\pi \sqrt{|z_4|} \right) - (-1)^{d_3} Y_0 \left( 4\pi \sqrt{|z_4|} \right) \right) \frac{\mathrm{d}z_1 \mathrm{d}z_2}{z_1 z_2}, \end{split}$$

where  $K_0$  and  $Y_0$  are the usual Bessel functions and

$$z_3 = \frac{y_1^2}{y_2^2 z_1 z_2}, \quad z_4 = y_2 (1 + \eta_1 \sqrt{z_1}) (1 + \eta_2 \sqrt{z_2}) (1 + \eta_3 \sqrt{z_3}),$$
  
$$d_3 = d_1 + d_2 - 2d_1 d_2.$$

Then using the trivial bound  $\ll \log(3 + \sqrt{|z_4|^{-1}})$  for  $K_0$ ,  $Y_0$ -Bessel functions and changing variables  $\frac{y_2 z_1^{3/2}}{y_1} \mapsto z_1'$  and  $\frac{y_1}{y_2 z_2^{3/2}} \mapsto z_2'$ , respectively, we imply directly

$$\Phi_{w_6}(\epsilon_1 y_1, \epsilon_2 y_2) \ll T^{3+\varepsilon}.$$

For  $y \in (T^{-100}, T^{100})$ ,  $\epsilon \in \{\pm 1\}$ , [8, Lemma 4.2] again implies that

$$\Phi_{w_4}(\epsilon y) := \frac{T^3}{64\pi^{13/2}} \sum_{d \in \{0,1\}} (\epsilon i)^{1-d} \int_0^\infty \int_0^\infty \check{h}_{T,M}\left(\frac{z_1^3}{y}, \frac{z_1^3 z_2^3}{y^2}\right) \\ \times \sin 2\pi \left(\frac{d}{2} + z_1\right) \sin 2\pi \left(\frac{d}{2} + z_2\right) \sin 2\pi \left(\frac{d}{2} + \frac{y}{z_1 z_2}\right) \frac{\mathrm{d}z_1 \mathrm{d}z_2}{z_1 z_2},$$

and trivially bounding the integral gives  $\Phi_{w_4}(\epsilon y) \ll T^{3+\epsilon}$ .

## 4. The argument principle

In this section, we use the following version of the argument principle, introduced by Selberg [44, Lemma 14] and Conrey–Soundararajan [9, Lemma 2.1].

**Lemma 4.1.** Let  $\omega$  be a holomorphic function, which is non-zero in some half-plane  $\operatorname{Re}(s) \geq W$ . Let  $\mathcal{B}$  be the rectangular box with vertices  $W_0 \pm iH$ ,  $W_1 \pm iH$  where H > 0 and  $W_0 < W < W_1$ . Then

$$4H \sum_{\substack{\beta+i\gamma\in\mathcal{B}\\\omega(\beta+i\gamma)=0}} \cos\left(\frac{\pi\gamma}{2H}\right) \sinh\left(\frac{\pi(\beta-W_0)}{2H}\right)$$
$$= \int_{-H}^{H} \cos\left(\frac{\pi t}{2H}\right) \log|\omega(W_0+it)| dt$$
$$+ \int_{W_0}^{W_1} \sinh\left(\frac{\pi(\alpha-W_0)}{2H}\right) \log|\omega(\alpha+iH)\omega(\alpha-iH)| d\alpha$$
$$- \operatorname{Re} \int_{-H}^{H} \cos\left(\frac{\pi(W_1-W_0+it)}{2iH}\right) \log\omega(W_1+it) dt.$$

**Remark 7.** The special feature of this lemma, which allows it to hold uniformly even for small  $H \approx 1/\log T$ , is that only the real part of the logarithm appears in the portion of the integral within the critical strip, so that this part can be bounded using the second moment estimate from Proposition 1.4.

#### QINGFENG SUN AND HUI WANG

### 5. The twisted second moment

In this section, we devote to giving the **proof of Theorem 1.2**. Using the approximate functional equation in Lemma 2.1, we can rewrite the twisted second moment as

$$\begin{split} &\sum_{j} \frac{h_{T,M}(\mu_{j})}{\mathcal{N}_{j}} A_{j}(\ell_{1},\ell_{2}) \left| L(1/2 + \sigma + i\tau,\phi_{j}) \right|^{2} \\ &= \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{j} \frac{h_{T,M}(\mu_{j})}{\mathcal{N}_{j}} A_{j}(\ell_{1},\ell_{2}) \overline{A_{j}(n,m)} W_{\mu_{j}} \left(\frac{1}{2} + \sigma + i\tau, r^{2}mn\right) \\ &+ \pi^{6\sigma} \sum_{r} \frac{1}{r^{1-2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2-\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{j} \frac{h_{T,M}(\mu_{j})}{\mathcal{N}_{j}} A_{j}(\ell_{1},\ell_{2}) \overline{A_{j}(n,m)} \tilde{W}_{\mu_{j}} \left(\frac{1}{2} + \sigma + i\tau, r^{2}mn\right) \\ &\coloneqq \mathcal{M}_{1}(\ell_{1},\ell_{2}) + \mathcal{M}_{2}(\ell_{1},\ell_{2}). \end{split}$$

In view of (2.7), we may truncate the summations over m, n and r to  $r^2mn \leq T^{3+\varepsilon}$  with a negligible error term.

Now we use the Kuznetsov trace formula (see Lemma 3.2) for  $\mathcal{M}_i(\ell_1, \ell_2)$ , i = 1, 2, respectively, with the test functions

$$h_1(\mu) = h_{T,M}(\mu) W_\mu \left(\frac{1}{2} + \sigma + i\tau, r^2 mn\right),$$
(5.1)

and

$$h_2(\mu) = h_{T,M}(\mu) \tilde{W}_{\mu} \left(\frac{1}{2} + \sigma + i\tau, r^2 mn\right).$$
(5.2)

It turns out we are led to estimate

$$\mathcal{M}_1(\ell_1, \ell_2) = \sum_r \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \left(\Delta^{(1)} + \Sigma_4^{(1)} + \Sigma_5^{(1)} + \Sigma_6^{(1)} - \mathcal{E}_{\max}^{(1)} - \mathcal{E}_{\min}^{(1)}\right), \quad (5.3)$$

and

$$\mathcal{M}_{2}(\ell_{1},\ell_{2}) = \pi^{6\sigma} \sum_{r} \frac{1}{r^{1-2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2-\sigma}} \left(\frac{m}{n}\right)^{i\tau} \left(\Delta^{(2)} + \Sigma_{4}^{(2)} + \Sigma_{5}^{(2)} + \Sigma_{6}^{(2)} - \mathcal{E}_{\max}^{(2)} - \mathcal{E}_{\min}^{(2)}\right),$$
(5.4)

where for i = 1, 2,

$$\begin{split} \Delta^{(i)} &:= \delta_{\ell_1,n} \delta_{\ell_2,m} \frac{1}{192\pi^5} \int_{\operatorname{Re}(\mu)=0} h_i(\mu) \mathrm{d}_{\operatorname{spec}} \mu, \\ \Sigma_4^{(i)} &:= \sum_{\epsilon = \pm 1} \sum_{\substack{D_2 \mid D_1 \\ mD_1 = \ell_1 D_2^2}} \frac{\tilde{S}(-\epsilon \ell_2, m, n; D_2, D_1)}{D_1 D_2} \Phi_{i,w_4} \left(\frac{\epsilon n m \ell_2}{D_1 D_2}\right), \\ \Sigma_5^{(i)} &:= \sum_{\epsilon = \pm 1} \sum_{\substack{D_1 \mid D_2 \\ nD_2 = \ell_2 D_1^2}} \frac{\tilde{S}(\epsilon \ell_1, n, m; D_1, D_2)}{D_1 D_2} \Phi_{i,w_5} \left(\frac{\epsilon \ell_1 n m}{D_1 D_2}\right), \\ \Sigma_6^{(i)} &:= \sum_{\epsilon_1, \epsilon_2 = \pm 1} \sum_{D_1, D_2} \frac{S(\epsilon_2 \ell_2, \epsilon_1 \ell_1, n, m; D_1, D_2)}{D_1 D_2} \Phi_{i,w_6} \left(-\frac{\epsilon_2 n \ell_2 D_2}{D_1^2}, -\frac{\epsilon_1 m \ell_1 D_1}{D_2^2}\right), \end{split}$$

with  $\Phi_{i,w_4}(y)$ ,  $\Phi_{i,w_5}(y)$  and  $\Phi_{i,w_6}(y)$  defined as in (3.15) by using the test functions  $h_i(\mu)$  given by (5.1) and (5.2), respectively; moreover, we have

$$\begin{aligned} \mathcal{E}_{\max}^{(i)} &:= \sum_{g} \frac{1}{2\pi i} \int_{\text{Re}(\mu)=0} \frac{h_i(\mu + \mu_g, \mu - \mu_g, -2\mu)}{\mathcal{N}_{\mu,g}} \overline{B_{\mu,g}(n,m)} B_{\mu,g}(\ell_1, \ell_2) d\mu, \\ \mathcal{E}_{\min}^{(i)} &:= \frac{1}{24(2\pi i)^2} \iint_{\text{Re}(\mu)=0} \frac{h_i(\mu)}{\mathcal{N}_{\mu}} \overline{A_{\mu}(n,m)} A_{\mu}(\ell_1, \ell_2) d\mu_1 d\mu_2. \end{aligned}$$

5.1. The diagonal terms  $\Delta^{(1)}$  and  $\Delta^{(2)}$ . Note that we have  $\ell_1 = n$  and  $\ell_2 = m$ . Thus we infer that the diagonal terms in (5.3) and (5.4) are

$$\frac{1}{192\pi^5} \sum_{r} \frac{1}{r^{1+2\sigma}} \frac{1}{(\ell_1 \ell_2)^{1/2+\sigma}} \left(\frac{\ell_2}{\ell_1}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) W_{\mu} \left(1/2 + \sigma + i\tau, r^2 \ell_1 \ell_2\right) \mathrm{d}_{\operatorname{spec}}\mu, \quad (5.5)$$

and

$$\frac{\pi^{6\sigma}}{192\pi^5} \sum_{r} \frac{1}{r^{1-2\sigma}} \frac{1}{(\ell_1\ell_2)^{1/2-\sigma}} \left(\frac{\ell_2}{\ell_1}\right)^{i\tau} \int_{\mathrm{Re}(\mu)=0} h_{T,M}(\mu) \tilde{W}_{\mu} \left(1/2 + \sigma + i\tau, r^2\ell_1\ell_2\right) \mathrm{d}_{\mathrm{spec}}\mu.$$
(5.6)

Using the definitions in (2.4) and (2.5) of  $W_{\mu}(s, y)$  and  $\tilde{W}_{\mu}(s, y)$ , respectively, we have (5.5) and (5.6) equal to

$$\frac{1}{192\pi^{5}} \frac{1}{(\ell_{1}\ell_{2})^{1/2+\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \frac{1}{2\pi i} \int_{(3)} \zeta(1+2\sigma+2u) (\pi^{3}\ell_{1}\ell_{2})^{-u} \\
\cdot \prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2+\sigma+i\tau+u-\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{1/2+\sigma-i\tau+u+\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma-i\tau+\mu_{k}}{2}\right)} G(u) \frac{\mathrm{d}u}{u} \mathrm{d}_{\operatorname{spec}}\mu,$$
(5.7)

and

$$\frac{\pi^{6\sigma}}{192\pi^{5}} \frac{1}{(\ell_{1}\ell_{2})^{1/2-\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \frac{1}{2\pi i} \int_{(3)} \zeta(1-2\sigma+2v) (\pi^{3}\ell_{1}\ell_{2})^{-v} \\
\cdot \prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2-\sigma+i\tau+v-\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{1/2-\sigma-i\tau+v+\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma-i\tau+\mu_{k}}{2}\right)} G(v) \frac{\mathrm{d}v}{v} \mathrm{d}_{\operatorname{spec}}\mu.$$
(5.8)

We shift the *u*-integral in (5.7) to  $\operatorname{Re}(u) = -B$  for a large 0 < B < A/2. The main contributions are coming from the residues at two simple poles at u = 0 and  $u = -\sigma$ . Hence, by the residue theorem, we infer that

$$\zeta(1+2\sigma)\frac{1}{(\ell_{1}\ell_{2})^{1/2+\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \frac{1}{192\pi^{5}} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \operatorname{d}_{\operatorname{spec}}\mu \\ -\frac{\pi^{3\sigma}}{384\pi^{5\sigma}} \frac{G(\sigma)}{(\ell_{1}\ell_{2})^{1/2}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2+i\tau-\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{1/2-i\tau+\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma-i\tau+\mu_{k}}{2}\right)} \operatorname{d}_{\operatorname{spec}}\mu.$$

Similarly, we shift the v-integral in (5.8) to  $\operatorname{Re}(v) = -B$  for a large 0 < B < A/2. The main contributions are coming from the residues at two simple poles at v = 0 and  $v = \sigma$ , which are

$$\frac{\pi^{6\sigma}}{192\pi^{5}} \frac{\zeta(1-2\sigma)}{(\ell_{1}\ell_{2})^{1/2-\sigma}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2-\sigma+i\tau-\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{1/2-\sigma-i\tau+\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma-i\tau+\mu_{k}}{2}\right)} d_{\operatorname{spec}}\mu + \frac{\pi^{3\sigma}}{384\pi^{5}\sigma} \frac{G(\sigma)}{(\ell_{1}\ell_{2})^{1/2}} \left(\frac{\ell_{2}}{\ell_{1}}\right)^{i\tau} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2+i\tau-\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{1/2-i\tau+\mu_{k}}{2}\right)}{\Gamma\left(\frac{1/2+\sigma-i\tau+\mu_{k}}{2}\right)} d_{\operatorname{spec}}\mu.$$

Note that the contribution from the residue at  $u = -\sigma$  from (5.7) and the residue at  $v = \sigma$  from (5.8) are canceled. By Stirling's formula and the exponentially decay of G, the contributions from the residues at the other places and the integral at  $\operatorname{Re}(u) = \operatorname{Re}(v) = -B$  are negligible. Moreover, using (2.12) and the Taylor expansion

$$(1+x)^{\alpha} = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!}x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}x^3 + o(x^3),$$

we have

$$\prod_{k=1}^{3} \frac{\Gamma\left(\frac{1/2 - \sigma + i\tau - \mu_k}{2}\right)}{\Gamma\left(\frac{1/2 + \sigma + i\tau - \mu_k}{2}\right)} \frac{\Gamma\left(\frac{1/2 - \sigma - i\tau + \mu_k}{2}\right)}{\Gamma\left(\frac{1/2 + \sigma - i\tau + \mu_k}{2}\right)} = \left(-\frac{\mu_1^2 \mu_2^2 \mu_3^2}{64}\right)^{-\sigma} + O\left(T^{-6\sigma - 3/4}\right),$$

provided for  $|\tau| \ll T^{1/4}$ . Thus, we obtain the main terms of Theorem 1.2 and the first error term  $O((\ell_1\ell_2)^{-1/2+\sigma}T^{9/4-6\sigma}M^2)$  by mean of (3.17).

5.2. The terms  $\Sigma_4^{(i)}$  and  $\Sigma_5^{(i)}$  for i = 1, 2. In this subsection, we will focus on the contribution from the  $w_4$ -term in  $\Sigma_4^{(1)}$ , that is,

$$\sum_{\epsilon=\pm 1} \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{\substack{D_2|D_1\\mD_1=\ell_1D_2^2}} \frac{\tilde{S}(-\epsilon\ell_2,m,n;D_2,D_1)}{D_1D_2} \Phi_{1,w_4}\left(\frac{\epsilon nm\ell_2}{D_1D_2}\right),$$

since the contributions from the  $w_4$ -term in  $\Sigma_4^{(2)}$  and the  $w_5$ -term in  $\Sigma_5^{(i)}$  (i = 1, 2) are very similar. Inserting a smooth partition of unity into m, n-sums, we are led to estimate

$$\sum_{\epsilon=\pm 1} \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{W_1(\frac{m}{\mathcal{M}}) W_2(\frac{n}{\mathcal{N}})}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{\substack{\delta,D\\m\delta=\ell_1D}} \frac{\tilde{S}(-\epsilon\ell_2,m,n;D,D\delta)}{D^2\delta} \Phi_{1,w_4}\left(\frac{\epsilon nm\ell_2}{D^2\delta}\right), \quad (5.9)$$

where  $\mathcal{M}, \mathcal{N} \gg 1, \mathcal{M}\mathcal{N} \ll T^{3+\varepsilon}$  (by (2.7)) and  $W_i(x)$  (i = 1, 2) are compactly supported in [1, 2] and satisfy  $x^j W_i^{(j)}(x) \ll_j 1$ , for any integer  $j \ge 0$ . Using the definitions (3.15), (5.1) and (2.4) of  $\Phi_{w_4}(y), h_1(\mu)$  and  $W_{\mu}(s, y)$ , respectively, we get that  $\Phi_{1,w_4}(y)$  is given by

$$\int_{\mathrm{Re}(\mu)=0} h_{T,M}(\mu) \frac{1}{2\pi i} \int_{(3)} (\pi^3 r^2 m n)^{-u} \prod_{k=1}^3 \prod_{\pm} \frac{\Gamma\left(\frac{\frac{1}{2} + \sigma \pm i\tau + u \mp \mu_k}{2}\right)}{\Gamma\left(\frac{\frac{1}{2} + \sigma \pm i\tau \mp \mu_k}{2}\right)} G(u) \frac{\mathrm{d}u}{u} K_{w_4}(y;\mu) \mathrm{d}_{\mathrm{spec}}\mu.$$

Now we consider the new resulting  $\mu$ -integral

$$\int_{\mathrm{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^{3} \frac{\Gamma\left(\frac{\frac{1}{2}+\sigma+i\tau+u-\mu_{k}}{2}\right)}{\Gamma\left(\frac{\frac{1}{2}+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{\frac{1}{2}+\sigma-i\tau+u+\mu_{k}}{2}\right)}{\Gamma\left(\frac{\frac{1}{2}+\sigma-i\tau+\mu_{k}}{2}\right)} K_{w_{4}}(y;\mu) \mathrm{d}_{\mathrm{spec}}\mu$$

with  $|\operatorname{Im}(u)| \ll T^{\varepsilon}$ , since the exponentially decay of G(u). With the help of (2.12), we have

$$\prod_{k=1}^{3} \frac{\Gamma\left(\frac{\frac{1}{2}+\sigma+i\tau+u-\mu_{k}}{2}\right)}{\Gamma\left(\frac{\frac{1}{2}+\sigma+i\tau-\mu_{k}}{2}\right)} \frac{\Gamma\left(\frac{\frac{1}{2}+\sigma-i\tau+u+\mu_{k}}{2}\right)}{\Gamma\left(\frac{\frac{1}{2}+\sigma-i\tau+\mu_{k}}{2}\right)} \ll T^{O(1)}$$

Then by applying the similar argument as Blomer–Buttcane [3, Lemma 1], we can truncate the D,  $\delta$ -sums at some  $T^{B_1}$  for some large  $B_1$  at the cost of a negligible error. In other words, we can use Lemma 3.4 with  $\Phi_{1,w_4}(y)$  replacing  $\Phi_{w_4}(y)$ . Let

$$W_{\mu,N_1}(1/2 + \sigma + i\tau, y) = \frac{1}{2\pi i} \int_{(3)} (\pi^3 y)^{-u} \prod_{k=1}^3 \prod_{\pm} \left(\frac{1/2 + \sigma \mp i\tau \pm \mu_k}{2}\right)^{u/2} \cdot \left(1 + \sum_{n=1}^{N_1} \frac{2^n P_n(u)}{(1/2 + \sigma \mp i\tau \pm \mu_k)^n}\right) G(u) \frac{\mathrm{d}u}{u}$$

Then with the help of (2.13), we can replace  $W_{\mu}$  by  $W_{\mu,N_1}$  in (5.9) with a negligible error if we choose  $N_1$  to be large enough.

Denote  $\Phi_{1,w_4}^{\star}(y)$  as in (3.15) with  $h_{T,M}(\mu)W_{\mu,N_1}\left(\frac{1}{2}+\sigma+i\tau,r^2mn\right)$  replacing  $h_1(\mu)$ . Then, in (5.9), we can replace  $\Phi_{1,w_4}(y)$  by  $\Phi_{1,w_4}^{\star}(y)$  with a negligible error term if we choose  $N_1$  to be large enough. Hence, together with the above truncation, we only need to consider the contribution from  $\Phi_{1,w_4}^{\star}(y)$ , where  $|y| > T^{-B_2}$  (by Lemma 3.3).

Note that in the original proof of Lemma 3.4 (see [3, Lemma 8]), the only two properties of  $h_{T,M}(\mu)$  which are used are (3.16) and (3.17). Obviously,  $h_{T,M}(\mu)W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, r^2mn\right)$  also satisfies these two properties. So we have

$$\Phi_{1,w_4}^{\star}(y) \ll_{\varepsilon,B_2} T^{-B_2},$$

provided for  $T^{-B_2} \leq |y| \leq T^{3-\varepsilon}$ . Then by Lemma 3.4, we can truncate the D,  $\delta$ -sums, again with a negligible error, at

$$\frac{mn\ell_2}{D^2\delta} \ge T^{3-\varepsilon},\tag{5.10}$$

or in other words,

$$D^2 \delta \leqslant \frac{\mathcal{MN}\ell_2}{T^{3-\varepsilon}} \ll \ell_2 T^{\varepsilon}$$

Note that we have  $m\delta = \ell_1 D$  now, which implies  $\mathcal{M}$  is small. That is,

$$\mathcal{M} \leqslant \ell_1 D / \delta \ll \ell_1 \ell_2^{1/2} T^{\varepsilon} / \delta^{3/2}.$$

And by (5.10) we have

$$\mathcal{N} \gg T^{3-\varepsilon} D^2 \delta / (m\ell_2) = T^{3-\varepsilon} D \delta^2 / (\ell_1 \ell_2) \gg T^{3-\varepsilon} / (\ell_1 \ell_2)$$

Therefore, by the above evaluation and the definition of  $\Phi_{1,w_4}^{\star}(y)$ , we only need to estimate

$$\sum_{\epsilon=\pm 1} \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{W_1(\frac{m}{\mathcal{M}}) W_2(\frac{n}{\mathcal{N}})}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{\substack{\delta,D\\m\delta=\ell_1D}} \frac{\tilde{S}(-\epsilon\ell_2,m,n;D,D\delta)}{D^2\delta} \times \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) W_{\mu,N_1}\left(\frac{1}{2}+\sigma+i\tau,r^2mn\right) K_{w_4}\left(\frac{\epsilon nm\ell_2}{D^2\delta};\mu\right) \mathrm{d}_{\operatorname{spec}}\mu.$$
(5.11)

We apply the Poisson summation formula for the n-sum, getting

$$\sum_{n=1}^{\infty} \frac{W_2(\frac{n}{N})}{n^{1/2+\sigma+i\tau}} \tilde{S}(-\epsilon\ell_2, m, n; D, D\delta) W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, r^2mn\right) K_{w_4}\left(\frac{\epsilon nm\ell_2}{D^2\delta}; \mu\right)$$

$$= \sum_{\substack{a(\text{mod }\delta)}} \tilde{S}(-\epsilon\ell_2, m, a; D, D\delta) \sum_{\substack{n\equiv a(\text{mod }\delta)}} \frac{W_2(\frac{n}{N})}{n^{1/2+\sigma+i\tau}} W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, r^2mn\right) K_{w_4}\left(\frac{\epsilon nm\ell_2}{D^2\delta}; \mu\right)$$

$$= \frac{\mathcal{N}^{1/2-\sigma-i\tau}}{\delta} \sum_{\substack{a(\text{mod }\delta)}} \tilde{S}(-\epsilon\ell_2, m, a; D, D\delta) \sum_{n\in\mathbb{Z}} e\left(\frac{an}{\delta}\right)$$

$$\cdot \int_{-\infty}^{\infty} \frac{W_2(x)}{x^{1/2+\sigma+i\tau}} W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, r^2mx\mathcal{N}\right) K_{w_4}\left(\frac{\epsilon x\mathcal{N}m\ell_2}{D^2\delta}; \mu\right) e\left(-\frac{n\mathcal{N}x}{\delta}\right) dx.$$
(5.12)

Inserting (5.12) into (5.11), using Lemma 3.1 and letting

$$\rho_1 = \operatorname{Im}(\mu_1 + \mu_2), \quad \rho_2 = \operatorname{Im}(\mu_1 - \mu_2),$$

we get the resulting  $\rho$ -integral involving  $\tilde{K}$ -function is

$$\int_{\mathbb{R}^2} h_{T,M}(\mu) W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, r^2 m x \mathcal{N}\right)$$
$$\cdot \int_0^\infty \tilde{K}_{i\rho_2}(2\sqrt{u}) \exp\left(i\rho_1 \log \frac{\pi^3 |y|}{u^{3/2}}\right) \exp\left(\frac{2i\pi^3 y}{u}\right) \frac{\mathrm{d}u}{u} \operatorname{spec}(\mu) \mathrm{d}\rho, \tag{5.13}$$

where

$$y = \frac{\epsilon x \mathcal{N} m \ell_2}{D^2 \delta},$$

and  $\rho = (\rho_1, \rho_2)$ .

We use the strategy in [3, Lemma 8] to deal with it. By using (3.16), the definition of  $W_{\mu,N_1}$ , and integration of by parts, we get that the  $\rho_1$ -integral is negligible unless

$$u \asymp |y|^{2/3}.$$

For the *u*-integral in (5.13), by differentiating k times with respect to y and by using (3.14), we see that each differentiation produces a factor

$$T|y|^{-1} + u^{-1} \approx (T + |y|^{1/3})|y|^{-1}.$$

We can truncate the integral involving  $J^{-}$ -function in the same way, and finally get

$$|y|^{k} \frac{\partial^{k}}{\partial y^{k}} K_{w_{4}}(y;\mu) \ll (T+|y|^{1/3})^{k}.$$
(5.14)

Now we go back to the x-integral in (5.12). With the help of (5.14), the definition of  $W_{\mu,N_1}$ , and the bounds for  $\mathcal{M}$  and  $\mathcal{N}$ , we integrate by parts and show that the x-integral is negligible unless n = 0. By opening  $\tilde{S}(-\epsilon \ell_2, m, a; D, D\delta)$  and computing the a-sum, we obtain

$$\begin{split} \sum_{a \pmod{\delta}} \tilde{S}(-\epsilon \ell_2, m, a; D, D\delta) \\ &= \sum_{a \pmod{\delta}} \sum_{\substack{C_1 \pmod{D}, C_2 \pmod{D\delta} \\ (C_1, D) = (C_2, \delta) = 1}} e\left(m\frac{\bar{C}_1 C_2}{D} + a\frac{\bar{C}_2}{\delta} - \epsilon \ell_2 \frac{C_1}{D}\right) = 0, \end{split}$$

unless  $\delta = 1$ . With the help of this and the fact  $m = \ell_1 D$ , we see that the contribution from n = 0 to (5.11) is

$$\mathcal{N}^{1/2-\sigma-i\tau}\ell_{1}^{-1/2-\sigma+i\tau}\sum_{\epsilon=\pm1}^{\infty}\sum_{r}\frac{1}{r^{1+2\sigma}}\sum_{D=1}^{\infty}W_{1}\left(\frac{\ell_{1}D}{\mathcal{M}}\right)\frac{1}{D^{3/2+\sigma-i\tau}}\sum_{\substack{C_{1}(\text{mod }D)\\(C_{1},D)=1}}e\left(-\frac{\epsilon\ell_{2}C_{1}}{D}\right)$$
$$\cdot\int_{\text{Re}(\mu)=0}h_{T,M}(\mu)\int_{-\infty}^{\infty}\frac{W_{2}(x)}{x^{1/2+\sigma+i\tau}}W_{\mu,N_{1}}\left(\frac{1}{2}+\sigma+i\tau,x\mathcal{N}r^{2}\ell_{1}D\right)K_{w_{4}}\left(\frac{\epsilon x\mathcal{N}\ell_{1}\ell_{2}}{D};\mu\right)dxd_{\text{spec}}\mu.$$

$$(5.15)$$

By inserting the definition (3.12) of  $K_{w_4}(y;\mu)$ , the above double integrals become

$$\int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \\ \times \int_{-i\infty}^{i\infty} \left( \int_{-\infty}^{\infty} x^{-u-1/2-\sigma-i\tau} W_2(x) W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, x \mathcal{N}r^2 \ell_1 D\right) \mathrm{d}x \right) \left| \frac{\ell_1 \ell_2 \mathcal{N}}{D} \right|^{-u} \\ \times \tilde{G}^{\epsilon}(u,\mu) \frac{\mathrm{d}u}{2\pi i} \mathrm{d}_{\operatorname{spec}} \mu.$$
(5.16)

By integration by parts with respect to the above *x*-integral, we can restrict the *u*-integral to  $|\tau + \text{Im}(u)| \leq T^{\varepsilon}$ . Inserting the definition (3.11) of  $\tilde{G}^{\epsilon}(u,\mu)$  into (5.16), we obtain the corresponding  $\mu$ -integral is

$$\int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, x \mathcal{N}r^2 \ell_1 D\right) \\ \times \left(\prod_{k=1}^3 \frac{\Gamma(\frac{1}{2}(u-\mu_k))}{\Gamma(\frac{1}{2}(1-u+\mu_k))} + i\epsilon \prod_{k=1}^3 \frac{\Gamma(\frac{1}{2}(1+u-\mu_k))}{\Gamma(\frac{1}{2}(2-u+\mu_k))}\right) d_{\operatorname{spec}}\mu.$$

We only need to consider the first part, since the second part is very similar. For fixed  $\sigma_1 \in \mathbb{R}$ ,  $t \in \mathbb{R}$  and  $|t| \ge 1$ , we have the Stirling formula

$$\Gamma(\sigma_1 + it) = \sqrt{2\pi} e^{\pi |t|/2} |t|^{\sigma_1 - 1/2} e^{it(\log |t| - 1)} e^{i(\sigma_1 - 1/2)\lambda\pi/2} \left(1 + O(|t|^{-1})\right),$$

where  $\lambda = 1$ , if  $t \ge 1$ , and  $\lambda = -1$ , if  $t \le -1$ . For convenience, we denote  $\mu_1 = it_1, \mu_2 = it_2, \mu_3 = -i(t_1 + t_2) = it_3$  and  $\operatorname{Im}(u) = t$ . Here, in the essential integrated range, we have  $|t_k| = |t_k - t'_k| = T$  for  $1 \le k, k' \le 3$  and  $k \ne k'$ , since we are considering the generic case. Without

loss of generality, we assume  $t_1 > 0$ ,  $t_2 > 0$  and  $t_3 < 0$ , and get

$$\prod_{k=1}^{3} \frac{\Gamma(\frac{1}{2}(u-\mu_{k}))}{\Gamma(\frac{1}{2}(1-u+\mu_{k}))}$$

$$= 2^{3/2} e^{i(-3t(\log 2+1)+\pi/4)} \frac{e^{i(t-t_{1})\log(t_{1}-t)+i(t-t_{2})\log(t_{2}-t)+i(t_{1}+t_{2}+t)\log(t_{1}+t_{2}+t)}}{(t_{1}-t)^{1/2}(t_{2}-t)^{1/2}(t_{1}+t_{2}+t)^{1/2}} + O(T^{-5/2}).$$
(5.17)

By the trivial estimate, the contribution from the error term in (5.17) to (5.15) is

$$\ll \mathcal{N}^{1/2-\sigma} \ell_1^{-1/2-\sigma} \sum_r \frac{1}{r^{1+2\sigma}} \sum_{D=1}^{\infty} W_1\left(\frac{\ell_1 D}{\mathcal{M}}\right) \frac{1}{D^{3/2+\sigma}} \bigg| \sum_{\substack{C_1 \pmod{D} \\ (C_1,D)=1}} e\left(-\frac{\epsilon \ell_2 C_1}{D}\right) \bigg| \cdot T^{-5/2} \\ \cdot \int_{|\tau + \operatorname{Im}(u)| \leqslant T^{\varepsilon}} \bigg| \int_{-\infty}^{\infty} x^{-u-1/2-\sigma-i\tau} W_2(x) \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) W_{\mu,N_1}\left(\frac{1}{2} + \sigma + i\tau, x \mathcal{N}r^2 \ell_1 D\right) \mathrm{d}_{\operatorname{spec}} \mu \mathrm{d}x \bigg| \mathrm{d}u \\ \ll \mathcal{N}^{1/2-\sigma} \ell_1^{-1/2-\sigma} T^{-5/2} T^{3+\varepsilon} M^2 \sum_{D \asymp \frac{\mathcal{M}}{\ell_1}} \frac{(D,\ell_2)}{D^{3/2+\sigma}} \ll T^{2-3\sigma+\varepsilon} M^2.$$

For the main term in (5.17), we denote the phase function for the  $\mu_1$ -integral is

$$\phi(t_1) := (t - t_1) \log(t_1 - t) + (t_1 + t_2 + t) \log(t_1 + t_2 + t).$$

Note that

$$\phi'(t_1) = \log \frac{t_1 + t_2 + t}{t_1 - t} \gg 1.$$

And hence, by partial integration many times, the  $\mu_1$ -integral is negligible. We finally prove that the contribution from the  $w_4$ -term in  $\Sigma_4^{(1)}$  to (5.3) is  $O(T^{2-3\sigma+\varepsilon}M^2)$ , which is the second error term of Theorem 1.2.

5.3. The terms  $\Sigma_6^{(1)}$  and  $\Sigma_6^{(2)}$ . Similarly, we only consider the contribution from  $w_6$ -term in  $\Sigma_6^{(1)}$ . By Lemma 3.5, we can truncate the sums at

$$\frac{(n\ell_2)^{1/3}(m\ell_1)^{1/6}}{D_1^{1/2}} \ge T^{1-\varepsilon}, \quad \text{and} \quad \frac{(n\ell_2)^{1/6}(m\ell_1)^{1/3}}{D_2^{1/2}} \ge T^{1-\varepsilon},$$

which means

$$\frac{(mn\ell_1\ell_2)^{1/2}}{(D_1D_2)^{1/2}} \ge T^{2-\varepsilon}.$$

This gives us

$$D_1 D_2 \ll \frac{mn\ell_1\ell_2}{T^{4-\varepsilon}},$$

which is impossible provided that  $\ell_1, \ell_2 \leq T^{1/3}$ , since the essential sums of n and m are truncated at  $mn \leq T^{3+\varepsilon}$ . Thus, the contribution of the  $w_6$ -term is  $O(T^{-B})$ .

5.4. The terms  $\mathcal{E}_{\max}^{(1)}$  and  $\mathcal{E}_{\max}^{(2)}$ . We only treat the contribution of the maximal Eisenstein series  $\mathcal{E}_{\max}^{(1)}$ , since the other contribution can be handled similarly. Firstly, we recall Weyl's law on GL<sub>2</sub> (see Venkov [47, Theorem 7.3] and Hejhal [16, pg. 119, Theorem 8.1]),

$$\sharp \{g : t_g \leq T\} = \frac{T^2}{12} - \frac{T\log T}{2\pi} + C_0 T + O\left(\frac{T}{\log T}\right),$$
(5.18)

where  $C_0$  is a constant. Next we use the subconvexity bounds for GL(1) and GL(2) *L*-functions to handle the contribution from this part. More precisely, we denote

$$\Theta := \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \sum_{g} \frac{1}{2\pi i} \int_{\operatorname{Re}(\mu)=0} \frac{h_1(\mu')}{\mathcal{N}_{\mu,g}} \overline{B_{\mu,g}(n,m)} B_{\mu,g}(\ell_1,\ell_2) \mathrm{d}\mu,$$

where  $\mu'$  is defined in (3.8). Inserting the definitions (5.1) and (2.4) into  $h_1(\mu)$  and  $W_{\mu}(s, y)$ , respectively, and using (3.4) and (3.6), we get

$$\Theta = \frac{1}{(2\pi i)^2} \sum_g \int_{\operatorname{Re}(\mu)=0} \frac{h_{T,M}(\mu')}{\mathcal{N}_{\mu,g}} B_{\mu,g}(\ell_1,\ell_2) \int_{(3)} \pi^{-3u} \zeta(\overline{s}+u+2\mu)\zeta(s+u-2\mu)$$
$$\times L(\overline{s}+u-\mu,g) L(s+u+\mu,g) \prod_{k=1}^3 \frac{\Gamma\left(\frac{s+u-\mu'_k}{2}\right)}{\Gamma\left(\frac{s-\mu'_k}{2}\right)} \frac{\Gamma\left(\frac{\overline{s}+u+\mu'_k}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu'_k}{2}\right)} G(u) \frac{\mathrm{d}u}{u} \mathrm{d}\mu,$$

where  $s = 1/2 + \sigma + i\tau$ . By the definition of  $h_{T,M}$ , we have  $h_{T,M}(\mu')$  is negligibly small unless

$$|\mu'_k - \mu_{0,k}| \le M, \quad for \ k = 1, 2, 3$$

Recall that  $|\mu_{0,k}| \approx T$  and (3.9). We now shift the line of integration to  $\operatorname{Re}(u) = \varepsilon$  and use Stirling's formula, the subconvexity bounds (see Bourgain [4, Theorem 5], and Meurman [40])

$$\zeta(1/2+it) \ll (1+|t|)^{13/84+\varepsilon}, \quad L(1/2+it,g) \ll (1+|t|)^{1/3+\varepsilon}$$
(5.19)

and [24, (5.20)] to get

$$\Theta \ll T^{41/42 - 41\sigma/21 + \varepsilon} M(\ell_1 \ell_2)^{\vartheta} \sum_{T - M \leqslant \mu_g \leqslant T + M} 1.$$

Finally, using Weyl's law (5.18), we conclude that

$$\Theta \ll T^{83/42 - 41\sigma/21 + \varepsilon} M^2 (\ell_1 \ell_2)^\vartheta, \tag{5.20}$$

which gives the third error term of Theorem 1.2.

5.5. The terms  $\mathcal{E}_{\min}^{(1)}$  and  $\mathcal{E}_{\min}^{(2)}$ . Similarly, to treat the contribution of the minimal Eisenstein series, we define

$$\Xi := \sum_{r} \frac{1}{r^{1+2\sigma}} \sum_{m,n} \frac{1}{(mn)^{1/2+\sigma}} \left(\frac{m}{n}\right)^{i\tau} \frac{1}{24(2\pi i)^2} \iint_{\operatorname{Re}(\mu)=0} \frac{h_1(\mu)}{\mathcal{N}_{\mu}} \overline{A_{\mu}(n,m)} A_{\mu}(\ell_1,\ell_2) \mathrm{d}\mu_1 \mathrm{d}\mu_2.$$

Inserting the definitions (5.1) and (2.4) into  $h_1(\mu)$  and  $W_{\mu}(s, y)$ , and using (3.2) and (3.3), we get

$$\begin{split} \Xi &= \frac{1}{24(2\pi i)^2} \iint_{\operatorname{Re}(\mu)=0} \frac{h_{T,M}(\mu)}{\mathcal{N}_{\mu}} A_{\mu}(\ell_1,\ell_2) \\ &\times \frac{1}{2\pi i} \int_{(3)} \pi^{-3u} \prod_{k=1}^3 \zeta(s+u+\mu_k) \zeta(\overline{s}+u-\mu_k) \frac{\Gamma\left(\frac{s+u-\mu_k}{2}\right)}{\Gamma\left(\frac{s-\mu_k}{2}\right)} \frac{\Gamma\left(\frac{\overline{s}+u+\mu_k}{2}\right)}{\Gamma\left(\frac{\overline{s}+\mu_k}{2}\right)} G(u) \frac{\mathrm{d}u}{u} \mathrm{d}\mu_1 \mathrm{d}\mu_2, \end{split}$$

where  $s = 1/2 + \sigma + i\tau$ . We shift the line of integration to  $\operatorname{Re}(u) = \varepsilon$  and use Stirling's formula, the subconvexity bound (5.19), and [24, (5.20)] to conclude that

$$\Xi \ll T^{13/14 - 13\sigma/7 + \varepsilon} M^2 (\ell_1 \ell_2)^\vartheta,$$

which is smaller than the contribution of the maximal Eisenstein series given in (5.20).

#### 6. The mollified second moment

In this section, we will give the **proof of Proposition 1.4**. Recall the definition of  $M(s, \phi_j)$  from (1.5). By applying the Hecke relation (2.3), we obtain

$$|M(1/2 + \sigma + i\tau, \phi_j)|^2 = \sum_{\ell_1} \sum_{\ell_2} \frac{A_j(1, \ell_2) A_j(1, \ell_1)}{\ell_1^{1/2 + \sigma - i\tau} \ell_2^{1/2 + \sigma + i\tau}} x_{\ell_1} x_{\ell_2}$$
$$= \sum_d \sum_{(\ell_1, d)=1} \sum_{(\ell_2, d)=1} \frac{A_j(\ell_1, \ell_2)}{d^{1+2\sigma} \ell_1^{1/2 + \sigma - i\tau} \ell_2^{1/2 + \sigma + i\tau}} x_{d\ell_1} x_{d\ell_2}$$

Using the above relation and Theorem 1.2, the left-hand side of (1.7) is transformed into

$$\begin{split} &\frac{1}{\mathcal{H}} \sum_{d} \sum_{(\ell_1,d)=1} \sum_{(\ell_2,d)=1} \frac{x_{d\ell_1} x_{d\ell_2}}{d^{1+2\sigma} \ell_1^{1/2+\sigma-i\tau} \ell_2^{1/2+\sigma+i\tau}} \sum_j \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} A_j(\ell_1,\ell_2) |L(1/2+\sigma+i\tau,\phi_j)|^2 \\ &= \zeta(1+2\sigma) \sum_{d} \sum_{(\ell_1,d)=1} \sum_{(\ell_2,d)=1} \frac{x_{d\ell_1} x_{d\ell_2}}{(d\ell_1\ell_2)^{1+2\sigma}} \\ &+ \zeta(1-2\sigma) \sum_{d} \sum_{(\ell_1,d)=1} \sum_{(\ell_2,d)=1} \frac{x_{d\ell_1} x_{d\ell_2}}{d^{1+2\sigma} \ell_1\ell_2} \frac{1}{\mathcal{H}} \frac{1}{192\pi^5} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^3 \left(-\frac{\mu_k^2}{4\pi^2}\right)^{-\sigma} \mathrm{d}_{\operatorname{spec}} \mu \\ &+ O\left(\sum_{d} \sum_{\ell_1} \sum_{\ell_2} \frac{|x_{d\ell_1} x_{d\ell_2}|}{(d^2\ell_1\ell_2)^{1/2+\sigma}} \left((\ell_1\ell_2)^{-1/2+\sigma} T^{-3/4-6\sigma} + T^{-1-3\sigma+\varepsilon} + T^{-43/42-41\sigma/21+\varepsilon} (\ell_1\ell_2)^\vartheta\right)\right) \\ &= \mathcal{S}_1 + \mathcal{S}_2 + \mathcal{E}_1 + \mathcal{E}_2 + \mathcal{E}_3. \end{split}$$

Next, we apply a strategy similar to that of Goldfeld–Huang [12, §4] to deal with the terms  $S_1$  and  $S_2$ .

6.1. The treatment of  $S_1$ . By the integral definition (1.6) of  $x_{d\ell_i}$ , (i = 1, 2), we have

$$S_1 = \zeta(1+2\sigma) \frac{1}{(2\pi i)^2} \int_{(3)} \int_{(3)} G_1(u,v;\sigma) \frac{L^{2u} - L^u}{u^2} \frac{L^{2v} - L^v}{v^2} \frac{\mathrm{d}u\mathrm{d}v}{(\log L)^2}$$
(6.1)

where

$$G_{1}(u,v;\sigma) = \sum_{d} \frac{\mu^{2}(d)}{d^{1+u+v+2\sigma}} \sum_{(\ell_{1},d)=1} \sum_{(\ell_{2},d)=1} \frac{\mu(\ell_{1})\mu(\ell_{2})}{\ell_{1}^{1+u+2\sigma}\ell_{2}^{1+v+2\sigma}}$$
  
= 
$$\prod_{p} \left(1 - p^{-1-u-2\sigma} - p^{-1-v-2\sigma} + p^{-1-u-v-2\sigma} + p^{-2-u-v-4\sigma}\right)$$
  
= 
$$\frac{\zeta(1+u+v+2\sigma)}{\zeta(1+u+2\sigma)\zeta(1+v+2\sigma)} H_{1}(u,v;\sigma).$$

The Euler product defining  $H_1(u, v; \sigma)$  converges absolutely in the region

$$\operatorname{Re}(u) + 2\sigma > -1/2$$
,  $\operatorname{Re}(v) + 2\sigma > -1/2$ ,  $\operatorname{Re}(u) + \operatorname{Re}(v) + 2\sigma > -1/2$ .

**Remark 8.** Recall that in the region  $\operatorname{Re}(s) \ge 1 - c/\log(|\operatorname{Im}(s)| + 3)$  (here c is some positive constant),  $\zeta(s)$  is analytic except for a single pole at s = 1, and has no zeros and satisfies (see e.g., [46, (3.11.7) and (3.11.8)])

 $\zeta(s)^{-1} \ll \log(|\operatorname{Im}(s)| + 3), \quad \text{and} \quad \zeta'(s)/\zeta(s) \ll \log(|\operatorname{Im}(s)| + 3).$ 

To evaluate the integral in (6.1), we shift both contours to the line  $\operatorname{Re}(u) = \operatorname{Re}(v) = 1/\log T$ without passing any poles and truncate the *v*-integral at  $|\operatorname{Im}(v)| \leq T$  with an error of  $O(T^{-1+\varepsilon})$ . Now, we have

$$S_{1} = \frac{\zeta(1+2\sigma)}{(2\pi i)^{2}} \int_{\left(\frac{1}{\log T}\right)} \int_{\frac{1}{\log T}-iT}^{\frac{1}{\log T}+iT} \frac{\zeta(1+u+v+2\sigma)}{\zeta(1+u+2\sigma)\zeta(1+v+2\sigma)} H_{1}(u,v;\sigma) \\ \cdot \frac{L^{2u}-L^{u}}{u^{2}} \frac{L^{2v}-L^{v}}{v^{2}} \frac{\mathrm{d}u\mathrm{d}v}{(\log L)^{2}} + O(T^{-1+\varepsilon}).$$
(6.2)

Shifting the *u*-integral to  $C_1$  given by

$$C_1 := \left\{ u : \operatorname{Re}(u) = -2\sigma - \frac{c}{\log^{3/4}(2 + |\operatorname{Im}(u)|)} \right\},\$$

where c > 0 is a suitable constant such that  $\zeta(s)$  has no zero in the region  $\operatorname{Re}(s) > 1 - c \log^{-3/4}(2 + |\operatorname{Im}(s)|)$  and  $c \log^{-3/4}(2) < 1/3$ , we encounter two simple poles at u = 0 and  $u = -2\sigma - v$ . The first pole yields a residue

$$\frac{1}{2\pi i} \int_{\frac{1}{\log T} - iT}^{\frac{1}{\log T} + iT} H_1(0, v; \sigma) \frac{L^{2v} - L^v}{v^2} \frac{\mathrm{d}v}{\log L} = 1 + O(T^{-\delta/2 - 2\delta\sigma + \delta\varepsilon}).$$
(6.3)

Here we extend the integration to the full line with the error term  $O(T^{-1+\varepsilon})$ , and shift the contour to  $\operatorname{Re}(v) = -1/2 - 2\sigma + \varepsilon$ . The second pole yields a residue

$$\frac{\zeta(1+2\sigma)}{(\log L)^2} \frac{1}{2\pi i} \int_{\frac{1}{\log T} - iT}^{\frac{1}{\log T} + iT} \frac{L^{-4\sigma} - L^{-v-4\sigma} + L^{-2\sigma} - L^{v-2\sigma}}{\zeta(1+v+2\sigma)\zeta(1-v)} H_1(-v-2\sigma,v;\sigma) \frac{\mathrm{d}v}{(v+2\sigma)^2 v^2}.$$
 (6.4)

Note that the contributions from the first three terms, that is,  $\frac{L^{-4\sigma}-L^{-v-4\sigma}+L^{-2\sigma}}{(v+2\sigma)^2v^2}$ , are small. Indeed, by shifting the integration to the contour  $C_{11}$  where

$$\mathcal{C}_{11} := \left[\frac{1}{\log T} - iT, \frac{1}{\log T} - i\right] \cup \left[\frac{1}{\log T} - i, \frac{1}{4} - i\right] \cup \left[\frac{1}{4} - i, \frac{1}{4} + i\right]$$
$$\cup \left[\frac{1}{4} + i, \frac{1}{\log T} + i\right] \cup \left[\frac{1}{\log T} + i, \frac{1}{\log T} + iT\right],$$

and using the standard bounds for the Riemann zeta function, one can show that the contributions from the the first three terms are bounded by

$$O\left(\frac{T^{-2\delta\sigma}}{\log^2 T}\right).\tag{6.5}$$

Now we consider the contribution from  $\frac{L^{\nu-2\sigma}}{(\nu+2\sigma)^2v^2}$  in (6.4). We move the line of integration to the contour  $C_{12}$  where

$$\mathcal{C}_{12} := \left[\frac{1}{\log T} - iT, -2\sigma - \frac{c}{\log^{3/4} T} - iT\right]$$
$$\cup \left[-2\sigma - \frac{c}{\log^{3/4} T} - iT, -2\sigma - \frac{c}{\log^{3/4} T} + iT\right]$$
$$\cup \left[-2\sigma - \frac{c}{\log^{3/4} T} + iT, \frac{1}{\log T} + iT\right],$$

for the same constant c in the definition of  $C_1$ , and pick up contributions from two simple poles at v = 0 and  $v = -2\sigma$ . The pole at v = 0 contributes

$$\frac{L^{-2\sigma}}{(2\sigma\log L)^2}H_1(-2\sigma,0;\sigma).$$
(6.6)

The pole at  $v = -2\sigma$  contributes

$$-\frac{L^{-4\sigma}}{(2\sigma\log L)^2}H_1(0, -2\sigma; \sigma).$$
(6.7)

Moreover, we find the *v*-integral on  $C_{12}$  can be bounded by

$$O\left(T^{-4\delta\sigma}e^{-c'\log^{1/4}T}\right), \quad \text{for some } c' > 0.$$
(6.8)

The remaining integral, for u of (6.2) on  $C_1$ , is bounded by

$$O\left(T^{-2\delta\sigma}e^{-c'\log^{1/4}T}\right), \quad \text{for some } c' > 0, \tag{6.9}$$

in view of standard bounds for  $\zeta(s)$  in the zero-free region. Combining (6.3) and (6.5)–(6.9), we conclude that

$$S_1 = 1 + O(T^{-2\delta\sigma}).$$
 (6.10)

6.2. The treatment of  $S_2$ . Recall that

$$S_2 = \zeta(1-2\sigma) \sum_d \sum_{(\ell_1,d)=1} \sum_{(\ell_2,d)=1} \frac{x_{d\ell_1} x_{d\ell_2}}{d^{1+2\sigma} \ell_1 \ell_2} \frac{1}{\mathcal{H}} \frac{1}{192\pi^5} \int_{\operatorname{Re}(\mu)=0} h_{T,M}(\mu) \prod_{k=1}^3 \left(-\frac{\mu_k^2}{4\pi^2}\right)^{-\sigma} \mathrm{d}_{\operatorname{spec}}\mu.$$

Using the trivial estimates, we obtain

$$S_2 \ll T^{-6\sigma} \sum_{d \leqslant L^2} \sum_{\ell_1 \leqslant L^2/d} \sum_{\ell_2 \leqslant L^2/d} \frac{1}{d^{1+2\sigma} \ell_1 \ell_2} \ll T^{-6\sigma} \log^2 T.$$
(6.11)

Next we focus on getting a better bound for  $S_2$ , and define

$$\Delta := \sum_{d} \sum_{(\ell_1, d)=1} \sum_{(\ell_2, d)=1} \frac{x_{d\ell_1} x_{d\ell_2}}{d^{1+2\sigma} \ell_1 \ell_2}.$$

By the definition (1.6) of  $x_{d\ell_i}$ , (i = 1, 2), we have

$$\Delta = \frac{1}{(2\pi i)^2} \int_{(3)} \int_{(3)} G_2(u, v; \sigma) \frac{L^{2u} - L^u}{u^2} \frac{L^{2v} - L^v}{v^2} \frac{\mathrm{d}u \mathrm{d}v}{(\log L)^2}$$
(6.12)

where

$$G_{2}(u,v;\sigma) = \sum_{d} \frac{\mu^{2}(d)}{d^{1+u+v+2\sigma}} \sum_{(\ell_{1},d)=1} \sum_{(\ell_{2},d)=1} \frac{\mu(\ell_{1})\mu(\ell_{2})}{\ell_{1}^{1+u}\ell_{2}^{1+v}}$$
$$= \prod_{p} \left(1 - p^{-1-u} - p^{-1-v} + p^{-1-u-v-2\sigma} + p^{-2-u-v}\right)$$
$$= \frac{\zeta(1+u+v+2\sigma)}{\zeta(1+u)\zeta(1+v)} H_{2}(u,v;\sigma).$$

The Euler product defining  $H_2(u, v; \sigma)$  converges absolutely in the region

 $\min\big\{\operatorname{Re}(u)+\operatorname{Re}(v)+2\sigma,\,\operatorname{Re}(u),\,\operatorname{Re}(v)\big\}>-1/2.$ 

We may assume that  $\sigma < \frac{100 \log \log T}{\log T}$  since otherwise  $T^{-2\sigma} < T^{-\sigma}/(\log T)^{100}$  and the integral may be bounded directly by using standard bounds for  $\zeta(s)$  and  $\zeta^{-1}(s)$  to the right of the 1-line (see also the trivial bound in (6.11)). To study (6.12), we shift the u, v-integrals to the lines  $\operatorname{Re}(u) = \operatorname{Re}(v) = 1/\log T$  without passing any poles and truncate the v-integral at  $|\operatorname{Im}(v)| \leq T$  with the error term  $O(T^{-1+\varepsilon})$ . Now we have

$$\Delta = \frac{1}{(2\pi i)^2} \int_{\left(\frac{1}{\log T}\right)} \int_{\frac{1}{\log T} - iT}^{\frac{1}{\log T} + iT} \frac{\zeta(1 + u + v + 2\sigma)}{\zeta(1 + u)\zeta(1 + v)} H_2(u, v; \sigma) \\ \cdot \frac{L^{2u} - L^u}{u^2} \frac{L^{2v} - L^v}{v^2} \frac{\mathrm{d}u\mathrm{d}v}{(\log L)^2} + O(T^{-1+\varepsilon}).$$
(6.13)

Shifting the *u*-integral to  $C_2$  given by

$$C_2 := \left\{ u : \operatorname{Re}(u) = -\frac{c}{\log^{3/4}(2 + |\operatorname{Im}(u)|)} \right\},\$$

for the same constant c in the definition of  $C_1$ , we encounter a pole at  $u = -v - 2\sigma$ , which yields a residue

$$\frac{1}{(\log L)^2} \frac{1}{2\pi i} \int_{\frac{1}{\log T} - iT}^{\frac{1}{\log T} + iT} \frac{L^{-4\sigma} - L^{-v-4\sigma} + L^{-2\sigma} - L^{v-2\sigma}}{\zeta(1 - v - 2\sigma)\zeta(1 + v)} H_2(-v - 2\sigma, v; \sigma) \frac{\mathrm{d}v}{(v + 2\sigma)^2 v^2}.$$
 (6.14)

Note that the contributions from the first three terms, that is,  $\frac{L^{-4\sigma}-L^{-v-4\sigma}+L^{-2\sigma}}{(v+2\sigma)^2v^2}$ , are small. Indeed, by shifting the integration to the contour  $C_{21}$  where

$$\mathcal{C}_{21} := \left[\frac{1}{\log T} - iT, \frac{1}{\log T} - i\right] \cup \left[\frac{1}{\log T} - i, \frac{1}{4} - i\right] \cup \left[\frac{1}{4} - i, \frac{1}{4} + i\right] \cup \left[\frac{1}{4} + i, \frac{1}{\log T} + i\right] \cup \left[\frac{1}{\log T} + i, \frac{1}{\log T} + iT\right],$$

and using the standard bounds for the Riemann zeta function, one can show that the contributions from the the first three terms are bounded by

$$O\left(\frac{T^{-2\delta\sigma}}{\log^2 T}\right).\tag{6.15}$$

Now we consider the contribution from  $\frac{L^{v-2\sigma}}{(v+2\sigma)^2v^2}$  in (6.14). We move the line of integration in (6.14) to the contour  $C_{22}$  where

$$\begin{aligned} \mathcal{C}_{22} &:= \left[ \frac{1}{\log T} - iT, -\frac{c}{\log^{3/4} T} - iT \right] \cup \left[ -\frac{c}{\log^{3/4} T} - iT, -\frac{c}{\log^{3/4} T} + iT \right] \\ & \cup \left[ -\frac{c}{\log^{3/4} T} + iT, \frac{1}{\log T} + iT \right], \end{aligned}$$

for the same constant c in the definition of  $C_1$ , and pick up contributions from the poles at v = 0and  $v = -2\sigma$ . The pole at v = 0 contributes

$$-\frac{L^{-2\sigma}}{\zeta(1-2\sigma)(2\sigma\log L)^2}H_2(-2\sigma,0;\sigma).$$
(6.16)

The pole at  $v = -2\sigma$  contributes

$$\frac{L^{-4\sigma}}{\zeta(1-2\sigma)(2\sigma\log L)^2}H_2(0,-2\sigma;\sigma).$$
(6.17)

Similarly, we find the integration in (6.14) on  $\mathcal{C}_{22}$  can be bounded by

$$O\left(T^{-2\delta\sigma}e^{-c'\log^{1/4}T}\right), \quad \text{for some } c' > 0.$$
(6.18)

The remaining integral, for u of (6.13) on  $\mathcal{C}_2$ , is bounded by

$$O\left(e^{-c'\log^{1/4}T}\right), \quad \text{for some } c' > 0.$$
(6.19)

Combining (6.13) and (6.15)–(6.19), and noting that  $H_2(-2\sigma, 0; \sigma) = H_2(0, -2\sigma; \sigma) = 1$ , we obtain

$$\Delta = \frac{L^{-2\sigma}(-1+L^{-2\sigma})}{(2\sigma\log L)^2} + O\left(\frac{T^{-2\delta\sigma}}{\log^2 T}\right).$$

Further, we conclude that

$$S_2 \ll T^{-2\sigma\delta}.\tag{6.20}$$

### 6.3. The contribution of the error terms. By the trivial estimates, we have

$$\mathcal{E}_1 \ll T^{-3/4 - 6\sigma} \sum_{d \leqslant L^2} \sum_{\ell_1 \leqslant L^2/d} \sum_{\ell_2 \leqslant L^2/d} \frac{1}{d^{1 + 2\sigma} \ell_1 \ell_2} \ll T^{-3/4 - 6\sigma + \varepsilon}, \tag{6.21}$$

$$\mathcal{E}_{2} \ll T^{-1-3\sigma+\varepsilon} \sum_{d \leq L^{2}} \sum_{\ell_{1} \leq L^{2}/d} \sum_{\ell_{2} \leq L^{2}/d} \frac{1}{d^{1+2\sigma} (\ell_{1}\ell_{2})^{1/2+\sigma}} \ll T^{-1-3\sigma+(2-4\sigma)\delta+\varepsilon},$$
(6.22)

and

$$\mathcal{E}_{3} \ll T^{-43/42 - 41\sigma/21 + \varepsilon} \sum_{d \leqslant L^{2}} \sum_{\ell_{1} \leqslant L^{2}/d} \sum_{\ell_{2} \leqslant L^{2}/d} \frac{(\ell_{1}\ell_{2})^{7/64}}{d^{1 + 2\sigma}(\ell_{1}\ell_{2})^{1/2 + \sigma}}$$

$$\ll T^{-43/42 - 41\sigma/21 + 39\delta/16 - 4\sigma\delta + \varepsilon} \ll T^{-2\sigma\delta},$$
(6.23)

provided for  $\delta \leq 1/3$ . Thus, combining (6.10) and (6.20)–(6.23), we complete the proof of Proposition 1.4.

## 7. Proof of Theorem 1.1

In the proof of Theorem 1.1, we shall need the following summation formula, which is an application of the GL(3) Kuznetsov trace formula.

**Theorem 7.1.** Let  $P = m_1 n_1 m_2 n_2 \neq 0$ . We have for any  $\varepsilon > 0$  that

$$\sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} \overline{A_j(m_1, m_2)} A_j(n_1, n_2)$$
$$= \delta_{n_1, m_1} \delta_{n_2, m_2} \mathcal{H} + O\left( (TP)^{\varepsilon} (TP^{1/2} + T^3 + TM^2 P^{\vartheta}) \right),$$

where  $\mathcal{H}$  is defined in (1.8) and  $\vartheta = 7/64$  is defined in (3.5).

*Proof.* The proof follows closely the strategy of Buttcane–Zhou [8, Theorem 3.3] or Blomer [2, Theorem 5], which is given here for completeness. Firstly, Lemmas 3.4 and 3.5 imply that for any fixed B > 0,

$$\Phi_{w_6}\left(-\frac{\epsilon_2 m_1 n_2 D_2}{D_1^2}, -\frac{\epsilon_1 m_2 n_1 D_1}{D_2^2}\right) \ll T^{-B},$$

unless  $D_1 D_2 \ll P T^{-4} (TP)^{\varepsilon}$ , and

$$\Phi_{w_4}\left(\frac{\epsilon m_1 m_2 n_2}{D_1 D_2}\right) \ll T^{-B},$$

unless  $D_1 D_2 \ll PT^{-3}(TP)^{\varepsilon}$ . As mentioned as before, in order to use these lemmas, it is necessary to know the  $D_1$ ,  $D_2$ -sums may be terminated at some high power of T. At that point we may

replace  $h_{T,M}$  with a smooth, compactly supported test function satisfying (3.16). In the current context, [2, Lemma 3] becomes

$$\sum_{\epsilon_1,\epsilon_2=\pm 1} \sum_{D_1 \leqslant X_1} \sum_{D_2 \leqslant X_2} \frac{|S(\epsilon_2 n_2,\epsilon_1 n_1, m_1, m_2; D_1, D_2)|}{D_1 D_2} \ll (X_1 X_2)^{1/2+\varepsilon} (m_1 n_2, m_2 n_1)^{\varepsilon},$$

as the proof goes through essentially unchanged. So we use the upper bounds  $\Phi(y) \ll T^{3+\varepsilon}$  from Lemma 3.6, and obtain  $\Sigma_6 \ll P^{1/2}T(TP)^{\varepsilon}$ . Note that the different sharp of the summation region may be handled by dyadic partitions. Similarly, Larsen's bound (3.10) implies

$$\sum_{\epsilon=\pm 1} \sum_{\substack{D_2|D_1\\m_2D_1=n_1D_2\\D_1D_2\ll X}} \frac{\left|S(-\epsilon n_2, m_2, m_1; D_2, D_1)\right|}{D_1D_2} \ll X^{\varepsilon} \sum_{\substack{n_1|m_2D_1\\m_2D_1^2\ll n_1X}} \frac{(m_1, D_1)}{D_1} \ll X^{\varepsilon} (m_2n_1)^{\varepsilon},$$

So we have  $\Sigma_4 \ll T^3(TP)^{\varepsilon}$ . The treatment of the  $w_5$ -term is identical.

As discussed in § 5.4 and § 5.5, here we only treat the contribution of the maximal Eisenstein series, since the minimal Eisenstein series can be handled similarly and the contribution will be smaller. Combining the known bounds on the *L*-functions and the GL(2) Weyl law together with definitions of  $h_{T,M}$ , we see that the essential region of the integration on  $\mu$  is of length  $\approx M$ , and the essential number of the sum of  $\mu_g$  is of size  $\approx TM$ . Thus we imply the trivial bound  $\mathcal{E}_{\min} + \mathcal{E}_{\max} \ll TM^2 P^{\vartheta} (TP)^{\varepsilon}$  and complete the proof of this theorem.

We record here frequently used estimate

$$\mathcal{H} = \frac{1}{192\pi^5} \int_{\mathrm{Re}(\mu)=0} h_{T,M}(\mu) \mathrm{d}_{\mathrm{spec}}\mu = CT^3 M^2 + O(T^2 M^3),$$

for some constant C > 0 (see Buttcane–Zhou [8, pg. 15]). Taking  $m_1 = n_1 = m_2 = n_2 = 1$  in Theorem 7.1 immediately gives

**Corollary 7.2.** With the assumptions in Theorem 7.1. We have for small  $\varepsilon > 0$  that

$$\sum_{j} \frac{h_{T,M}(\mu_j)}{N_j} = CT^3 M^2 + O(T^2 M^3 + T^{3+\varepsilon}).$$

Now we give the **proof of Theorem 1.1**. We will apply Lemma 4.1 to  $L(s, \phi_j)M(s, \phi_j)$  with box bounded by  $1/2 + 1/\log T \pm 2iH$  and  $3/2 \pm 2iH$ . For  $2/\log T < \sigma < 1/2$ , we have

$$\begin{split} N(\sigma,H;\phi_j) &:= \sum_{\substack{L(1/2+\beta+i\gamma,\phi_j)=0\\\beta>\sigma,\,|\gamma|2/\log T,\,|\gamma|1/\log T,\,|\gamma|1/\log T,\,|\gamma|$$

Since the zeros of 
$$L(s, \phi_j)$$
 are also zeros of the function  $L(s, \phi_j)M(s, \phi_j)$ , we use Lemma 4.1 with  $\omega(s) = L(s, \phi_j)M(s, \phi_j) := LM(s, \phi_j)$ ,  $W_0 = 1/\log T$  and  $W_1 = 1$ , and get that  $N(\sigma, H; \phi_j)$  is  $\leq \log T \left( \int_{-H}^{H} \cos\left(\frac{\pi t}{2H}\right) \log |LM(1/2 + 1/\log T + it, \phi_j)| dt + \int_{\frac{1}{\log T}}^{1} \sinh\left(\frac{\pi(\alpha - 1/\log T)}{2H}\right) \log |LM(1/2 + \alpha + iH, \phi_j)LM(1/2 + \alpha - iH, \phi_j)| d\alpha - \operatorname{Re} \int_{-H}^{H} \cos\left(\frac{\pi(1 - 1/\log T + it)}{2iH}\right) \log LM(3/2 + it, \phi_j) dt \right).$ 

Since always

$$\log|x| \leqslant \frac{|x|^2 - 1}{2},$$

and the definition (1.3),  $N(\sigma, H)$  is

$$\leq \frac{\log T}{2} \bigg\{ \int_{-H}^{H} \cos\left(\frac{\pi t}{2H}\right) \frac{1}{\mathcal{H}} \sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} \left( |LM(1/2 + 1/\log T + it, \phi_j)|^2 - 1 \right) \mathrm{d}t \\ + \int_{\frac{1}{\log T}}^{1} \sinh\left(\frac{\pi(\alpha - 1/\log T)}{2H}\right) \frac{1}{\mathcal{H}} \sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} \\ \times \left( |LM(1/2 + \alpha + iH, \phi_j)|^2 + |LM(1/2 + \alpha - iH, \phi_j)|^2 - 2 \right) \mathrm{d}\alpha \\ - 2\operatorname{Re} \int_{-H}^{H} \cos\left(\frac{\pi(1 - 1/\log T + it)}{2iH}\right) \frac{1}{\mathcal{H}} \sum_{j} \frac{h_{T,M}(\mu_j)}{\mathcal{N}_j} \log LM(3/2 + it, \phi_j) \mathrm{d}t \bigg\}.$$

Using Proposition 1.4 and Corollary 7.2, for some sufficiently small  $\delta$ ,  $\theta_1 > 0$ , we conclude that

$$N(\sigma, H) \ll HT^{-\sigma\delta} \log T,$$

uniformly in  $3/\log T < H < T^{\theta_1}$ .

Acknowledgements The authors would like to thank Professor Sheng-Chi Liu for his valuable suggestions on this topic. H. Wang would like to thank the Alfréd Rényi Institute of Mathematics for providing a great working environment and the China Scholarship Council for its financial support.

### References

- O. Balkanova, B. Huang and A. Södergren, Non-vanishing of Maass form L-functions at the central point. Proc. Amer. Math. Soc. 149 (2021), no. 2, 509–523.
- [2] V. Blomer, Applications of the Kuznetsov formula on GL(3). Invent. Math. 194 (2013), no. 3, 673–729.
- [3] V. Blomer and J. Buttcane, On the subconvexity problem for L-functions on GL(3). Ann. Sci. Éc. Norm. Supér. (4) 53 (2020), no. 6, 1441–1500.
- [4] J. Bourgain, Decoupling, exponential sums and the Riemann zeta function. J. Amer. Math. Soc. 30 (2017), no. 1, 205–224.
- [5] D. Bump, S. Friedberg and D. Goldfeld, Poincaré series and Kloosterman sums for SL(3,ℤ). Acta Arith. 50 (1988), no. 1, 31–89.
- [6] J. Buttcane, On sums of SL(3, Z) Kloosterman sums. Ramanujan J. 32 (2013), no. 3, 371-419.

#### QINGFENG SUN AND HUI WANG

- [7] J. Buttcane, The spectral Kuznetsov formula on SL(3). Trans. Amer. Math. Soc. 368 (2016), no. 9, 6683–6714.
- [8] J. Buttcane and F. Zhou, Plancherel distribution of Satake parameters of Maass cusp forms on GL<sub>3</sub>. Int. Math. Res. Not. IMRN 2020, no. 5, 1417–1444.
- J.B. Conrey and K. Soundararajan, Real zeros of quadratic Dirichlet L-functions. Invent. Math. 150 (2002), no. 1, 1–44.
- [10] S.S. Gelbart, E.M. Lapid and P. Sarnak, A new method for lower bounds of L-functions. C. R. Math. Acad. Sci. Paris 339 (2004), no. 2, 91–94.
- [11] D. Goldfeld, Automorphic forms and L-functions for the group GL(n, ℝ). With an appendix by Kevin A. Broughan. Cambridge Studies in Advanced Mathematics, 99. Cambridge University Press, Cambridge, 2006. xiv+493 pp.
- [12] D. Goldfeld and B. Huang, Super-positivity of a family of L-functions in the level aspect. Res. Math. Sci. 5 (2018), no. 2, Paper No. 16, 26 pp.
- [13] L. Guth and J. Maynard, New large value estimates for Dirichlet polynomials. arXiv: 2405.20552
- [14] G. Halász, Über die Mittelwerte multiplikativer zahlentheoretischer Funktionen. (German) Acta Math. Acad. Sci. Hungar. 19 (1968), 365–403.
- [15] D.R. Heath-Brown, Zero density estimates for the Riemann zeta-function and Dirichlet L-functions. J. London Math. Soc. (2) 19 (1979), no. 2, 221–232.
- [16] D.A. Hejhal, The Selberg trace formula for PSL(2, ℝ). Vol. 2. Lecture Notes in Mathematics, 1001. Springer-Verlag, Berlin, 1983. viii+806 pp.
- [17] J. Hoffstein and P. Lockhart, Coefficients of Maass forms and the Siegel zero. With an appendix by Dorian Goldfeld, Hoffstein and Daniel Lieman. Ann. of Math. (2) 140 (1994), no. 1, 161–181.
- [18] J. Hoffstein and D. Ramakrishnan, Siegel zeros and cusp forms. Internat. Math. Res. Notices 1995, no. 6, 279–308.
- [19] B. Hough, Zero-density estimate for modular form L-functions in weight aspect. Acta Arith. 154 (2012), no. 2, 187–216.
- [20] B. Huang, S. Liu and Z. Xu, Mollification and non-vanishing of automorphic L-functions on GL(3). Israel J. Math. 227 (2018), no. 2, 597–622.
- [21] M.N. Huxley, On the difference between consecutive primes. Invent. Math. 15 (1972), 164–170.
- [22] A.E. Ingham, On the estimation of  $N(\sigma, T)$ . Quart. J. Math. Oxford Ser. 11 (1940), 291–292.
- [23] A. Ivić, The Riemann zeta-function. Theory and applications. Reprint of the 1985 original [Wiley, New York; MR0792089]. Dover Publications, Inc., Mineola, NY, 2003. xxii+517 pp.
- [24] H. Iwaniec and E. Kowalski, Analytic number theory. American Mathematical Society Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004. xii+615 pp.
- [25] H. Jacquet and J.A. Shalika, A non-vanishing theorem for zeta functions of GL<sub>n</sub>. Invent. Math. 38 (1976/77), no. 1, 1–16.
- [26] M. Jutila, Zero-density estimates for L-functions. Acta Arith. 32 (1977), no. 1, 55–62.
- [27] R. Khan, Non-vanishing of the symmetric square L-function at the central point. Proc. Lond. Math. Soc. (3) 100 (2010), no. 3, 736–762.
- [28] H.H. Kim, Functoriality for the exterior square of GL<sub>4</sub> and the symmetric fourth of GL<sub>2</sub>. With appendix 1 by Dinakar Ramakrishnan and appendix 2 by Kim and Peter Sarnak. J. Amer. Math. Soc. 16 (2003), no. 1, 139–183.
- [29] E. Kowalski and P. Michel, The analytic rank of  $J_0(q)$  and zeros of automorphic L-functions. Duke Math. J. **100** (1999), no. 3, 503–542.
- [30] E. Kowalski and P. Michel, A lower bound for the rank of  $J_0(q)$ . Acta Arith. 94 (2000), no. 4, 303–343.
- [31] E. Kowalski, P. Michel and J. VanderKam, Mollification of the fourth moment of automorphic L-functions and arithmetic applications. Invent. Math. 142 (2000), no. 1, 95–151.
- [32] E. Kowalski, P. Michel and J. VanderKam, Non-vanishing of high derivatives of automorphic L-functions at the center of the critical strip. J. Reine Angew. Math. 526 (2000), 1–34.
- [33] E. Kowalski, P. Michel and J. VanderKam, Rankin-Selberg L-functions in the level aspect. Duke Math. J. 114 (2002), no. 1, 123–191.
- [34] S. Liu, Nonvanishing of central L-values of Maass forms. Adv. Math. 332 (2018), 403–437.
- [35] S. Liu and S. Liu, A GL<sub>3</sub> analog of Selberg's result on S(t). Ramanujan J. 56 (2021), no. 1, 163–181.
- [36] S. Liu and J. Streipel, The twisted second moment of L-functions associated to Hecke-Maass forms. Int. J. Number Theory 20 (2024), no. 3, 849–866.

- [37] W. Luo, Nonvanishing of L-values and the Weyl law. Ann. of Math. (2) 154 (2001), no. 2, 477–502.
- [38] W. Luo, Zeros of Hecke L-functions associated with cusp forms. Acta Arith. 71 (1995), no. 2, 139–158.
- [39] W. Luo, Nonvanishing of the central L-values with large weight. Adv. Math. 285 (2015), 220–234.
- [40] T. Meurman, On the order of the Maass L-function on the critical line. Number theory, Vol. I (Budapest, 1987), 325–354, Colloq. Math. Soc. János Bolyai, 51, North–Holland, Amsterdam, 1990.
- [41] H.L. Montgomery, Zeros of L-functions. Invent. Math. 8 (1969), 346-354.
- [42] F. Olver, Asymptotics and special functions. Reprint of the 1974 original [Academic Press, New York; MR0435697]. AKP Classics. A K Peters, Ltd., Wellesley, MA, 1997. xviii+572 pp.
- [43] A. Selberg, Contributions to the theory of the Riemann zeta-function. Arch. Math. Naturvid. 48 (1946), no. 5, 89–155.
- [44] A. Selberg, Contributions to the theory of Dirichlet's L-functions. Skr. Norske Vid.-Akad. Oslo I 1946 (1946), no. 3, 62 pp.
- [45] K. Soundararajan, Nonvanishing of quadratic Dirichlet L-functions at  $s = \frac{1}{2}$ . Ann. of Math. (2) **152** (2000), no. 2, 447–488.
- [46] E.C. Titchmarsh, The theory of the Riemann zeta-function. Second edition. Edited and with a preface by D.
   R. Heath-Brown. The Clarendon Press, Oxford University Press, New York, 1986. x+412 pp.
- [47] A. B. Venkov, Spectral theory of automorphic functions and its applications. Translated from the Russian by N. B. Lebedinskaya. Mathematics and its Applications (Soviet Series), 51. Kluwer Academic Publishers Group, Dordrecht, 1990. xiv+176 pp.
- [48] Z. Xu, Nonvanishing of automorphic L-functions at special points. Acta Arith. 162 (2014), no. 4, 309–335.

School of Mathematics and Statistics, Shandong University, Weihai, Weihai, Shandong 264209, China

Email address: qfsun@sdu.edu.cn

SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN, SHANDONG 250100, CHINA *Email address*: wh0315@mail.sdu.edu.cn