

FACET-HAMILTONIAN CYCLES IN THE B -PERMUTAHEDRON

NASTARAN BEHROOZANIA, SOFIA BRENNER, ARTURO MERINO, TORSTEN MÜTZE,
CHRISTIAN RIECK, AND FRANCESCO VERCIANI

ABSTRACT. We construct facet-Hamiltonian cycles in the B -permutahedron, resolving a conjecture raised in a recent paper by Akitaya, Cardinal, Felsner, Kleist and Lauff [<https://arxiv.org/abs/2411.02172>].

1. INTRODUCTION

In a recent paper, Akitaya, Cardinal, Felsner, Kleist and Lauff [ACF⁺24, Conjecture 1] conjectured that the B -permutahedron admits a so-called facet-Hamiltonian cycle; see Section 2 for definitions. In this short note, we prove this conjecture (Theorem 4 below), by constructing rhombic strips in the face lattice of hypercubes. This preprint serves as a teaser for many more upcoming results on various notions of Hamiltonian cycles that visit faces of polytopes.

2. PRELIMINARIES

Let $(P, <)$ be a poset. We say that two distinct elements $x, y \in P$ with $x < y$ are in a *cover relation*, denoted $x < y$, if there is no $z \in P$ with $x < z < y$. The *cover graph* of P , denoted $G(P)$, has as vertices all elements of P , and an edge (x, y) for every cover relation $x < y$. A poset is *graded* if there is a function $\rho : P \rightarrow \mathbb{Z}$ such that $\rho(y) = \rho(x) + 1$ if (x, y) is a cover relation in P . The function ρ is called *rank function*, and $\rho(x)$ is called the *rank* of $x \in P$. The *downset* of some $x \in P$ is the set of all $y \in P$ for which $y \leq x$. A *chain* in P is a sequence of elements (x_1, \dots, x_k) from P such that $x_1 < x_2 < \dots < x_k$. It corresponds to a path in the cover graph in which the ranks are strictly increasing.

A *polytope* P is the convex hull of a finite set of points in \mathbb{R}^n . Equivalently, it is a bounded intersection of half-spaces. If P is n -dimensional, we sometimes refer to it as an *n -polytope*. We write $G(P)$ for the graph of the skeleton of P . A *face* of P is the intersection of P with a hyperplane such that all of P lies on the same side or on the hyperplane. The combinatorial structure of P is captured by its *face lattice*, denoted $L(P)$, which is the inclusion order of all faces of P . Figure 1 (a) shows a 3-dimensional polytope, and part (c) of the figure shows its face lattice. The face lattice has the empty set \emptyset as its unique minimum, and the full polytope P as its unique maximum. We take as the rank function the dimensions of the faces, where the unique minimum \emptyset of $L(P)$ is assigned rank (dimension) -1 . Furthermore, the 0-dimensional

(Nastaran Behrooznia) DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF WARWICK, UNITED KINGDOM

(Sofia Brenner) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, GERMANY

(Arturo Merino) UNIVERSIDAD DE O'HIGGINS, RANCAGUA, CHILE

(Torsten Mütze) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, GERMANY

(Christian Rieck) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, GERMANY

(Francesco Verciani) INSTITUT FÜR MATHEMATIK, UNIVERSITÄT KASSEL, GERMANY

E-mail addresses: nastaran.behrooznia@warwick.ac.uk, sofia.brenner@mathematik.uni-kassel.de, arturo.merino@uoh.cl, tmuetze@mathematik.uni-kassel.de, christian.rieck@mathematik.uni-kassel.de, francesco.verciani@mathematik.uni-kassel.de .

This project was supported by German Science Foundation grant 522790373.




FIGURE 1. (a) A 3-polytope P ; (b) a Schlegel diagram of P , a 3-connected plane graph; (c) the face lattice of P ; (d1)+(d2) two distinct facet-Hamiltonian cycles of P (of different lengths); (e) a rhombic strip of $G(L(P))$. The rhombi are colored according to ranks for clarity. In this and all of the following pictures of rhombic strips in our paper, we display a grayed-out copy of the leftmost vertices at the right-hand side, in order to depict the ‘wrap-around’ edges on the cylinder.

faces are called *vertices* and the 1-dimensional faces are called *edges*. If P is n -dimensional, then the faces of dimension $n - 1$ are called *facets*. We denote these three sets by $V(P)$, $E(P)$, and $F(P)$, respectively. Given two distinct faces $F, G \in L(P)$, we write $F \subset G$ if F is a subspace of G , i.e., a subset of G that is a face of P . Furthermore, we write $F \subseteq G$ if F and G form a cover relation in $L(P)$, which means that $F \subset G$ and the dimension/rank of G is exactly one higher than that of F . A *flag* of P is a chain in $L(P)$ that starts at \emptyset and ends at P , i.e., this chain contains exactly one face from each rank.

A *facet-Hamiltonian cycle* of a polytope P is a cycle C in the skeleton graph $G(P)$ with the property that every facet of P has a nonempty and connected intersection with C ; see Figure 1 (d1)+(d2). In terms of the face lattice $L(P)$, such a cycle enters and leaves the downset of every facet exactly once.

The *cylinder* is the quotient of the set \mathbb{R}^2 under the equivalence relation $(x, y) \sim (x + 1, y)$. Given a graded poset P , a *rhombic strip* is a spanning subgraph of the cover graph $G(P)$ embedded on the cylinder subject to the following conditions; see Figure 1 (e):

- the height of vertices in the embedding is given by their rank;

- edges are drawn as straight lines and there are no crossings between them;
- every face is a *rhombus*, i.e., a 4-cycle (a, b, c, d) , where a and c have the same rank r , and b and d have ranks $r - 1$ and $r + 1$, respectively.

Given strings x and y , we write xy for their concatenation. This operation extends to sets of strings Y in the natural way, specifically $xY := \{xy \mid y \in Y\}$. Similarly, for a sequence of strings $Y = (y_1, \dots, y_n)$ we define $xY := (xy_1, \dots, xy_n)$. For a string x and integer n , we write x^n for the n -fold concatenation of x with itself. Given a sequence $x = (x_1, \dots, x_n)$, we write $\text{rev}(x) = (x_n, x_{n-1}, \dots, x_1)$ for the reversed sequence.

3. RHOMBIC STRIPS IN FACE LATTICES OF SIMPLICES

The *Boolean lattice* Q_n is the inclusion order of all 2^n subsets of $[n] := \{1, \dots, n\}$. We identify subsets of $[n]$ by their characteristic vectors of length n , i.e., we think of $G(Q_n)$ as the graph that has $\{0, 1\}^n$ as its vertex set, and an edge between any two bitstrings that differ in a single bit. The *n -simplex* Δ_n is the convex hull of the n unit vectors in \mathbb{R}^n ; see Figure 2. It has n vertices, and any k -subset of vertices forms a $(k - 1)$ -face, for $k = 1 \dots, n$, which is itself a simplex. Specifically, the 1-simplex is a point, the 2-simplex is a line segment, the 3-simplex is a triangle, the 4-simplex is a tetrahedron, etc. Furthermore, the face lattice of the n -simplex Δ_n is the Boolean lattice Q_n .




FIGURE 2. The 2- and 3-simplex and their face lattices, the Boolean lattices of dimension 3 and 4, respectively.

Theorem 1 (Folklore). *For any $n \geq 2$, the graph $G(L(\Delta_n)) = G(Q_n)$ has a rhombic strip.*

We revisit two proofs for Theorem 1, and later combine the techniques presented here for constructing rhombic strips for the face lattice of the hypercube. The following proofs are based on two classical constructions of the binary reflected Gray code, and in fact yield the same rhombic strip, under an isomorphism that renames elements of the ground set.

Proof 1 of Theorem 1. A rhombic strip is a spanning subgraph of $G(Q_n)$ with certain additional properties, and hence it has $\{0, 1\}^n$ as its vertex set. We specify such a subgraph by the set of edges of $G(Q_n)$ that it contains. Specifically, we construct a rhombic strip $R_n \subseteq G(Q_n)$ by induction on n ; see Figures 3 and 4. The construction also maintains two chains $C_n = (x_0, \dots, x_n) \subseteq R_n$ and $D_n = (y_0, \dots, y_n) \subseteq R_n$ with $x_0 = y_0 = 0^n$ and $x_n = y_n = 1^n$, where C_n contains the leftmost vertices across all ranks and D_n contains the rightmost vertices across all ranks, and both chains are connected by edges $Z_n := \{(x_i, y_{i+1}) \mid i = 1, \dots, n - 2\} \subseteq R_n$. We refer to the edges in Z_n as *zipper edges*, as they connect the left and right side of the plane drawing to a rhombic strip on the cylinder, like a zipper that connects two pieces of fabric.





FIGURE 3. Schematic illustration of proof 1 of Theorem 1.

FIGURE 4. The rhombic strips obtained from proof 1 of Theorem 1 for $n = 2, 3, 4, 5$.

For the base case $n = 2$ of the construction we take $R_2 := G(Q_2)$, $C_2 := (00, 01, 11)$ and $D_2 := (00, 10, 11)$, and in this case we have $Z_2 = \emptyset$.

For the induction step, let $R_n, C_n = (x_0, \dots, x_n), D_n = (y_0, \dots, y_n)$ be given for some $n \geq 2$. We construct the rhombic strip $R_{n+1} \subseteq G(Q_{n+1})$ as follows: We remove the zipper edges from R_n , and consider the plane graph $R' := R_n \setminus Z_n$. We take a copy of $0R'$, i.e., the graph R' in which all vertices are prefixed with a 0-bit, maintaining the same embedding. In addition, we take a mirrored copy of $1R'$, i.e., the graph R' in which all vertices are prefixed with a 1-bit and the embedding is mirrored along a vertical line. The second (mirrored) graph is embedded to the right of the first one and one unit higher than the first one (note that prefixing with a 1-bit increases the ranks). We connect the two graphs by the edges

$$E := \{(0y_i, 1y_i) \mid i = 0, \dots, n\}$$

and the zipper edges

$$Z_{n+1} := \{(0x_i, 1x_i) \mid i = 1, \dots, n-1\},$$

which yields the rhombic strip

$$R_{n+1} := 0R' \cup 1R' \cup E \cup Z_{n+1}.$$

Furthermore, the new leftmost and rightmost chain are

$$C_{n+1} := (0C_n, 1^{n+1}) \text{ and } D_{n+1} := (0^{n+1}, 1C_n),$$

respectively.

One can check that the vertices on the chains C_n and D_n , respectively, have the explicit form $x_i = 0^{n-i}1^i$ for $i = 0, \dots, n$ and $y_i = 10^{n-i}1^{i-1}$ for $i = 1, \dots, n$. \square

Proof 2 of Theorem 1. An x -monotone path in a drawing of a graph is a path with the property that its vertices are embedded with strictly increasing abscissa values. We construct a rhombic strip $R_n \subseteq G(Q_n)$ by induction on n ; see Figures 5 and 6. The construction maintains a partition $R_n = P_n \cup A_n \cup B_n \cup Z_n$ into four sets with the following properties:

- P_n is an x -monotone Hamiltonian path of $G(Q_n)$ that starts at the vertex 0^n and ends at a rank 0 vertex.
- The sequence $C_n = (x_0, \dots, x_n)$ of leftmost vertices across all ranks is a chain $C_n \subseteq R_n$, and the sequence $(0^n, D_n) = (0^n, y_1, \dots, y_n)$ of rightmost vertices across all ranks is a chain $(0^n, D_n) \subseteq R_n$, and the two chains are connected by the zipper edges $Z_n := \{(x_i, y_{i+1}) \mid i = 1, \dots, n-2\} \subseteq R_n$.
- The edges A_n and B_n are the edges of R_n strictly above and below P_n , respectively.

For the base case $n = 2$ of the construction we take $R_2 := G(Q_2)$ with $P_2 := (00, 10, 11, 01)$, which gives $C_2 = (00, 10, 11)$, $D_2 = (01, 11)$, $Z_2 = \emptyset$, $A_2 = \emptyset$ and $B_2 = \{(00, 01)\}$.

For the induction step, let P_n, C_n, D_n, A_n, B_n be given for some $n \geq 2$. We construct the rhombic strip R_{n+1} as follows: We consider the plane graphs $R^\times := R_n \setminus (A_n \cup Z_n) = P_n \cup B_n$




FIGURE 5. Schematic illustration of proof 2 of Theorem 1.




FIGURE 6. The rhombic strips obtained from proof 2 of Theorem 1 for $n = 2, 3, 4, 5$.

and $R_\times := R_n \setminus (B_n \cup Z_n) = P_n \cup A_n$. We take a copy of $0R_\times$ and a copy of $1R_\times$, maintaining the same embedding. Both graphs are embedded at the same positions horizontally, but the second graph is placed one unit higher than the first one, so that the corresponding vertices on the two copies of P_n line up vertically (note that prefixing with 1 increases the ranks). Let $P_n =: (p_1, \dots, p_N)$, $N = 2^n$, be the sequence of vertices on the Hamiltonian path P_n . Then we connect the two graphs by the edges

$$E := \{(0p_i, 1p_i) \mid i = 1, \dots, N\}$$

and the zipper edges

$$Z_{n+1} := 1Z_n \cup \{(1p_N, 10^n)\},$$

which yields the rhombic strip

$$R_{n+1} := 0R_\times \cup 1R_\times \cup E \cup Z_{n+1}.$$

The new Hamiltonian path is

$$P_{n+1} := (0p_1, 1p_1, 1p_2, 0p_2, 0p_3, 1p_3, \dots, 0p_{N-1}, 1p_{N-1}, 1p_N, 0p_N), \quad (1)$$

which becomes x -monotone by slight perturbations of the vertical edges. Furthermore, the new leftmost and rightmost chain are

$$C_{n+1} := (0^{n+1}, 1C_n) \text{ and } D_{n+1} := (0p_N, 1D_n),$$

respectively (with the exception of 0^{n+1} , which is not part of D_{n+1}), and the sets of edges above and below P_{n+1} are

$$\begin{aligned} A_{n+1} &:= 1A_n \cup \{(1p_i, 1p_{i+1}) \mid i = 2, 4, 6, \dots, N-2\} \text{ and} \\ B_{n+1} &:= 0B_n \cup \{(0p_i, 0p_{i+1}) \mid i = 1, 3, 5, \dots, N-1\}, \end{aligned} \quad (2)$$

respectively. Note that in (1) and (2) we have used that $N = 2^n$ is even.




FIGURE 7. The dual graph of a rhombic strip of $G(Q_n)$ is an n -Venn diagram. Vertices and regions are encoded by their characteristic vectors of length n .

One can check that the vertices x_i and y_i on the chains C_n and D_n , respectively, have the explicit form $x_i = 1^i 0^{n-i}$ for $i = 0, \dots, n$ and $y_i = 1^{i-1} 0^{n-i} 1$ for $i = 1, \dots, n$. \square

Maybe surprisingly, both aforementioned proofs actually yield the same rhombic strip, up to reversal of the vertex labels, which can be checked for $n = 2, 3, 4, 5$ by carefully comparing Figures 4 and 6, and which can easily be shown in general using induction.

There is an interesting connection between rhombic strips in $G(Q_n)$ and Venn diagrams. Specifically, the dual graph of a rhombic strip of $G(Q_n)$ is an n -Venn diagram, i.e., a diagram of n simple closed curves in the plane, such that every of the 2^n possible intersection patterns is represented by exactly one of the regions in the diagram; see Figure 7. Consequently, from any of the known constructions of Venn diagrams (see [Rus97, RSW06]), we obtain a rhombic strip of $G(L(\Delta_n)) = G(Q_n)$ by considering the dual graph. The two aforementioned proofs can of course also be dualized into the language of Venn diagrams.

4. RHOMBIC STRIPS IN FACE LATTICES OF HYPERCUBES

The n -(*hyper*)*cube* is the convex hull of the point set $\{0, 1\}^n$; see Figure 8. Equivalently, it is the n -fold Cartesian product of the unit interval $[0, 1]$. The skeleton of this polytope is isomorphic to the cover graph of the Boolean lattice Q_n , and so we use Q_n to also denote the polytope. Each face F of Q_n can be encoded by a ternary string $x_F \in \{0, 1, -\}^n$, where the set of vertices contained in F is given by replacing in x_F all occurrences of $-$ by either 0 or 1; see Figure 8. The dimension of the face F is given by the number of $-$ s in the string x_F . For example, $x_F = 01-- = \{0100, 0101, 0110, 0111\}$ represents a 2-dimensional face F (a quadrilateral), and $-^n = Q_n$ is the entire polytope.

The cover graph of the face lattice $G(L(Q_n))$ has as vertex set $\{0, 1, -\}^n \cup \{\emptyset\}$, and as edges all pairs of ternary strings that differ in replacing a single 0 or 1 by $-$, or vice versa, plus the edges (\emptyset, x) for all $x \in \{0, 1\}^n$.

Theorem 2. *For any $n \geq 1$, the graph $G(L(Q_n))$ has a rhombic strip.*

For proving this result, we combine the techniques developed in the two proofs of Theorem 1 presented in Section 3. We use induction and embed three copies of the previous structures, one prefixed with 0, one prefixed with $-$, and one prefixed with 1. The two structures prefixed with 0 and $-$ are connected as in the second proof via ‘stacking’, and the two structures prefixed with $-$ and 1 are connected as in the first proof via ‘mirroring’.




FIGURE 8. The 2- and 3-cube and their face lattices.

Proof. We construct a rhombic strip $R_n \subseteq G(L(Q_n))$ by induction on n ; see Figures 9 and 10. We define the abbreviation $B'_n := \{(\emptyset, x) \mid x \in \{0, 1\}^n\}$ for the set of edges between the minimum \emptyset of $L(Q_n)$ and all rank 0 faces. The construction maintains a partition $R_n = P_n \cup A_n \cup B_n \cup B'_n \cup Z_n$ into five sets with the following properties:

- P_n is an x -monotone Hamiltonian path of $G(L(Q_n)) \setminus \{\emptyset\}$ that starts and ends at a rank 0 vertex.
- The sequence $(\emptyset, C_n) = (\emptyset, x_0, \dots, x_n)$ of leftmost vertices across all ranks is a chain $(\emptyset, C_n) \subseteq R_n$, and the sequence $(\emptyset, D_n) = (\emptyset, y_0, \dots, y_n)$ of rightmost vertices across all ranks is a chain $(\emptyset, D_n) \subseteq R_n$, and the two chains are connected by the zipper edges $Z_n := \{(x_i, y_{i+1}) \mid i = 0, \dots, n-2\} \subseteq R_n$.
- The edges A_n and $B_n \cup B'_n$ are the edges of R_n strictly above and below P_n , respectively.

For the base case $n = 1$ of the construction we take $R_1 := G(L(Q_1))$ with $P_1 := (0, -, 1)$, which gives $C_1 = (0, -)$, $D_1 = (1, -)$, $Z_1 = \emptyset$, $A_1 = \emptyset$ and $B_1 = \emptyset$.

For the induction step, let $P_n, C_n = (x_0, \dots, x_n), D_n = (y_0, \dots, y_n), A_n, B_n$ be given for some $n \geq 1$. We construct the rhombic strip R_{n+1} as follows: We consider the plane graphs $R^\times := R_n \setminus (A_n \cup B'_n \cup Z_n) = P_n \cup B_n$, $R_\times := R_n \setminus (B_n \cup B'_n \cup Z_n) = P_n \cup A_n$ and $R' := R_n \setminus (B'_n \cup Z_n) = P_n \cup A_n \cup B_n$. We take a copy of $0R^\times$ and a copy of $-R_\times$, maintaining the same embedding, and a vertically mirrored copy of $1R'$. The first two graphs are embedded at the same positions horizontally, but the second graph is placed one unit higher than the first one, so that the corresponding vertices on the two copies of P_n line up vertically (note that prefixing with $-$ increases the ranks). The third (mirrored) graph is embedded to the right of the first two and at the same height as the first one. Let $P_n =: (p_1, \dots, p_N)$, $N = 3^n$, be the sequence of vertices on the Hamiltonian path P_n . Then we connect the first and second graph by the edges

$$E := \{(0p_i, -p_i) \mid i = 1, \dots, N\},$$




FIGURE 9. Schematic illustration of the proof of Theorem 2.

the second and third graph by the edges

$$F := \{(1y_i, -y_i) \mid i = 0, \dots, n\},$$

and the third and first graph by the zipper edges

$$Z_{n+1} := \{(1x_i, -x_i) \mid i = 0, \dots, n-1\},$$

which yields the rhombic strip

$$R_{n+1} := 0R^\times \cup -R^\times \cup 1R' \cup E \cup F \cup Z_{n+1} \cup B'_{n+1}.$$

The new Hamiltonian path (for $G(L(Q_{n+1})) \setminus \{\emptyset\}$) is

$$P_{n+1} := (0p_1, -p_1, -p_2, 0p_2, 0p_3, -p_3, \dots, -p_{N-1}, 0p_{N-1}, 0p_N, -p_N, 1 \operatorname{rev}(P_n)), \quad (3)$$

which becomes x -monotone by slight perturbations of the vertical edges in the first part. Furthermore, the new leftmost and rightmost chain are

$$C_{n+1} := (0p_1, -C_n) \text{ and } D_{n+1} := (1C_n, -^{n+1}),$$

respectively (with the exception of \emptyset , which is neither part of C_{n+1} nor D_{n+1}), and the sets of edges above and below P_{n+1} (in addition to B'_{n+1}) are

$$\begin{aligned} A_{n+1} &:= -A_n \cup 1A_n \cup \{(-p_i, -p_{i+1}) \mid i = 2, 4, 6, \dots, N-1\} \cup (F \setminus \{(1p_N, -p_N)\}) \text{ and} \\ B_{n+1} &:= 0B_n \cup 1B_n \cup \{(0p_i, 0p_{i+1}) \mid i = 1, 3, 5, \dots, N-2\}, \end{aligned} \quad (4)$$




FIGURE 10. The rhombic strips obtained from the proof of Theorem 2 for $n = 1, 2, 3, 4$.

respectively. Note that in (3) and (4) we have used that $N = 3^n$ is odd.

One can check that the vertices x_i and y_i on the chains C_n and D_n , respectively, have the explicit form $x_i = -^i 0^{n-i}$ for $i = 0, \dots, n$ and $y_i = 1^{-i} 0^{n-1-i}$ for $i = 0, \dots, n-1$. \square

5. TRUNCATION

The *permutahedron* Π_n is the convex hull of $\{(\pi(1), \dots, \pi(n)) \mid \pi \in S_n\}$, where S_n denotes the set of all permutations on $[n]$; see the top right of Figure 11. The permutahedron is $(n-1)$ -dimensional, and its edges connect pairs of permutations that differ in an adjacent transposition.¹

The *B -permutahedron* $\overline{\Pi}_n$ is the convex hull of $\{(\pm\pi(1), \dots, \pm\pi(n)) \mid \pi \in S_n\}$, i.e., it is the convex hull of all signed permutations. Its edges connect pairs of signed permutations that either differ in an adjacent transposition, preserving all signs, or in a complementation of the sign of the first entry; see the bottom right of Figure 11.

Given a polytope P , the *omnitruncation* of P , denoted $\text{tr}(P)$, is the simple polytope of the same dimension as P that has one vertex for each flag of P . Geometrically, it is obtained by truncating all the proper faces of P . It is well-known that truncating the simplex yields the permutahedron, i.e., we have $\text{tr}(\Delta_n) = \Pi_n$, and truncating the hypercube yields the B -permutahedron, i.e., we have $\text{tr}(Q_n) = \overline{\Pi}_n$; see Figure 11.

In their paper [ACF⁺24], the authors established the following connection between rhombic strips in the face lattice $L(P)$ of a polytope P and facet-Hamiltonian cycle in the omnitruncated polytope $\text{tr}(P)$.

Lemma 3. *If $G(L(P))$ has a rhombic strip, then $G(\text{tr}(P))$ has a facet-Hamiltonian cycle.*

¹By this we mean a transposition of adjacent positions, i.e., $\pi(i) \leftrightarrow \pi(i+1)$, which is true if vertices are labeled by the inverse of the permutation that defines the vertex coordinates in our definition via the convex hull.




FIGURE 11. (a) Truncating the simplex yields the permutahedron. (b) Truncating the hypercube yields the B -permutahedron. The facet-Hamiltonian cycles in the permutahedron and B -permutahedron are obtained via Lemma 3 from the rhombic strips shown in Figures 4 and 10, respectively. The highlighted vertex corresponds to the leftmost chain in the rhombic strip. Each of the highlighted four-cycles highlights two possibilities for the facet-Hamiltonian cycle to walk around this face, corresponding to two options of sweeping the chain through the rhombic strip.

The facet-Hamiltonian cycle in $G(\text{tr}(P))$ is obtained by sweeping a chain from left-to-right through the rhombic strip of $G(L(P))$, changing one element of the chain at a time, moving it across a rhombus. If several elements in the chain can be changed, then we have freedom in constructing several different facet-Hamiltonian cycles; see Figure 11.

Applying Lemma 3 for $P := \Delta_n$ and using Theorem 1 thus proves that the permutahedron Π_n has a facet-Hamiltonian cycle; see Figure 11 (a). In fact, we obtain many different such cycles, corresponding to the different constructions discussed in Section 3 that establish Theorem 1.

Similarly, applying Lemma 3 for $P := Q_n$ and using Theorem 2 yields the following result, which affirmatively resolves Conjecture 1 raised by Akitaya, Cardinal, Felsner, Kleist and Lauff [ACF⁺24]; see Figure 11 (b).

Theorem 4. *For any $n \geq 2$, the B -permutahedron $G(\overline{\Pi}_n)$ has a facet-Hamiltonian cycle.*

REFERENCES

- [ACF⁺24] H. Akitaya, J. Cardinal, S. Felsner, L. Kleist, and R. Lauff. Facet-Hamiltonicity, 2024. <https://arxiv.org/abs/2411.02172>. An extended abstract was accepted for presentation at SODA 2025.
- [RSW06] F. Ruskey, C. D. Savage, and S. Wagon. The search for simple symmetric Venn diagrams. *Notices Amer. Math. Soc.*, 53(11):1304–1312, 2006.
- [Rus97] F. Ruskey. A survey of Venn diagrams. *Electron. J. Combin.*, 4(1):Dynamic Survey 5, 10 HTML documents, 1997.