

SOME CHARACTERIZATIONS OF RIEMANNIAN MANIFOLDS ENDOWED WITH A CONFORMAL VECTOR FIELDS

A. BARROS¹, I. EVANGELISTA² AND E. VIANA³

ABSTRACT. The aim of this article is to investigate the presence of a conformal vector ξ with conformal factor ρ on a compact Riemannian manifold M with or without boundary ∂M . We firstly prove that a compact Riemannian manifold (M^n, g) , $n \geq 3$, with constant scalar curvature, with boundary ∂M totally geodesic, in such way that the traceless Ricci curvature is zero in the direction of $\nabla\rho$, is isometric to a standard hemisphere. In the 4-dimensional case, under the condition $\int_M |\mathring{Ric}|^2 \langle \xi, \nabla\rho \rangle dM \leq 0$, we show that, either M is isometric to a standard sphere, or M is isometric to a standard hemisphere. Finally, we give a partial answer for the cosmic no-hair conjecture.

1. INTRODUCTION

Conformal mappings and conformal vector fields were intensively studied during the last 150 years, in particular in Riemannian geometry of dimension $n \geq 3$. Namely, there have been notable findings concerning the characterization of an n -sphere $\mathbb{S}^n(c)$ as well as an hemisphere $\mathbb{S}_+^n(c)$, for some positive constant c , utilizing a non-Killing conformal vector field, as documented in [5, 6, 7, 8, 9, 10, 11, 12] and [13].

Before we proceed, given an n -dimensional Riemannian manifold (M^n, g) with Levi-Civita connection ∇ , we recall that a conformal vector field ξ on M is said to be closed if the 1-form ξ^\flat is closed. This is easily seen to be equivalent to the existence of a smooth function $\rho : M \rightarrow \mathbb{R}$, called the conformal factor of ξ , such that

$$\nabla_X \xi = \rho X,$$

for all $X \in \mathfrak{X}(M)$. Whence we have

$$\rho = \frac{1}{n} \operatorname{div} \xi,$$

where div is the divergence operators on M . Henceforth, we also use the notation $\langle \cdot, \cdot \rangle$ for the metric g or inner product induced by g on tensor spaces.

It's important to note that closed vector fields are also referred to as concircular vector fields. These fields have surfaced in the examination of conformal mappings that preserve geodesic circles and boast intriguing applications in general relativity (cf. [20]). The conformal nature of a vector field remains unchanged under a conformal alteration of the metric. However, a closed conformal vector field remains closed only if the conformal alteration maintains a constant conformal factor; in such instances, the factor of the field remains unaltered. Moreover, a closed and conformal vector field X is labelled parallel if ρ vanishes entirely. We highlight that numerous manifolds host non parallel closed; certain existence outcomes and explicit examples are detailed in [3] and [16].

After these preliminary remarks we may state our main results as follows.

2010 *Mathematics Subject Classification*. Primary: 53C20, 53A30.

Key words and phrases. Conformal gradient vector fields; cosmic no-hair conjecture; manifolds with boundary; hemisphere of the Euclidean sphere.

Theorem 1. *Let ξ be a non trivial conformal vector field with conformal factor ρ on a compact Riemannian manifold (M^n, g) with smooth totally geodesic boundary ∂M of dimension $n \geq 3$ and $\rho|_{\partial M} = 0$. If the scalar curvature R of (M^n, g) is constant and $\mathring{Ric}(\nabla\rho) = 0$, then M is isometric to a hemisphere $\mathbb{S}_+^n(c)$ for some positive constant c .*

Next, we present two rigidity results for the 4-dimensional case.

Theorem 2. *Let (M^4, g) be a compact Riemannian manifold with constant scalar curvature and ξ be a non trivial conformal vector field on M^4 with conformal factor ρ . If $\int_M |\mathring{Ric}|^2 \langle \xi, \nabla\rho \rangle dM \leq 0$, then M^4 is isometric to a standard sphere $\mathbb{S}^4(c)$ for some positive constant c .*

Theorem 3. *Let (M^4, g) be a compact Riemannian manifold with smooth totally geodesic boundary ∂M and constant scalar curvature. Let ξ be a non trivial conformal vector field on M^4 with conformal factor ρ , which satisfies $\rho = 0$ on ∂M . If $\int_M |\mathring{Ric}|^2 \langle \xi, \nabla\rho \rangle dM \leq 0$, then M^4 is isometric to a hemisphere $\mathbb{S}_+^4(c)$ for some positive constant c .*

In order to state our last result, we define the static triple as follows:

Definition 1. *A complete and connected Riemannian manifold (M^n, g) with a (possibly non empty) boundary ∂M is said to be static if there exists a non-negative function f on M satisfying*

$$(1.1) \quad (\Delta f)g + fRic = \nabla^2 f,$$

in $M \setminus \partial M$, and $\partial M = f^{-1}(0)$, where Ric , Δ and ∇^2 stand, respectively, for the Ricci tensor, the Laplacian operator and the Hessian form on M^n .

In this case, (M^n, g, f) is called a static triple or simply a static metric.

We highlight that, in General Relativity, Equation (1.1) appears as static solutions of Einstein field equations. Furthermore, Corvino et al. [4], Proposition 2.1] showed that a static metric also has constant scalar curvature R . Boucher et al. [2] formulated a classic conjecture, called *cosmic no-hair conjecture*, as follows:

The only n -dimensional compact static triple (M^n, g, f) with positive scalar curvature and connected boundary Σ is given by a round hemisphere \mathbb{S}_+^n , where the function f is taken as the height function.

Now we give a partial answer for the cosmic no-hair conjecture. We point out that this result is similar to that one derived in [15] for critical metrics.

Theorem 4. *Let (M^n, g, ρ) be a simply connected, compact static metric with smooth boundary ∂M totally geodesic and endowed with a smooth conformal vector field ξ which conformal factor is ρ . If $\psi := n\rho^2 - \langle \nabla\rho, \xi \rangle \neq 0$, then (M^n, g) is isometric to a hemisphere $\mathbb{S}_+^n(c)$ for some positive constant c .*

This paper is organized as follows. In Section 2 we review the necessary knowledge about conformal vector fields in Riemannian geometry, as well as properties of static metrics. In Section 3 we give the proofs of Theorems 1, 2, 3 and 4.

2. PRELIMINARIES AND KEY LEMMAS

In this section, we will present some basic results of static metrics, as well as we will present the basic concepts about conformal vector fields and basic results that will be used in the proofs of the main theorems of this work.

Note that tracing (1.1) we arrive at

$$(2.1) \quad \Delta f = -\frac{Rf}{n-1},$$

and, by using again (1.1), it is not difficult to check that

$$(2.2) \quad f\overset{\circ}{Ric} = \overset{\circ}{\nabla}^2 f,$$

where $\overset{\circ}{T}$ stands for the traceless of the tensor T .

We recall that a smooth vector field $\xi \in \mathfrak{X}(M)$ is said to be *conformal* if

$$(2.3) \quad \mathcal{L}_\xi g = 2\rho g$$

for a smooth function ρ on M , where \mathcal{L}_ξ is the Lie derivative in the direction of ξ . The function ρ is the conformal factor of ξ (cf. [1]). If ξ is the gradient of a smooth function on M , then ξ is said to be a *conformal gradient vector field*. In this case, ξ is also closed. We say that ξ is a non trivial conformal vector field if it is a non-Killing conformal vector field.

Remember from Equation (2.3) that a vector field ξ on Riemannian manifold (M^n, g) is called conformal if $\mathcal{L}_\xi g$ is a multiple of g . As a straightforward consequence of Koszul's formula, we have the following identity for ξ on M ,

$$2g(\nabla_X \xi, Y) = \mathcal{L}_\xi g(X, Y) + d\eta(X, Y), \quad X, Y \in \mathfrak{X}(M),$$

where η stands for the dual 1-form associated to Z , that is, $\eta(Y) = g(Z, Y)$. We note that we can define φ the following skew symmetric (1,1)-tensor:

$$d\eta(X, Y) = 2g(\varphi(X), Y), \quad X, Y \in \mathfrak{X}(M).$$

Therefore, by using the above equations, we arrive at

$$(2.4) \quad \nabla_X \xi = \rho X + \varphi(X), \quad X \in \mathfrak{X}(M).$$

Note that φ gives us an idea of how much of the field ξ is not closed vector field. There are several papers involving closed conformal vector fields (see, e.g., [3, 14, 19]).

Observe that we can identify φ with a skew symmetric (0, 2)-tensor and ξ with the tensor $\xi(Y) = g(\xi, Y)$, $Y \in \mathfrak{X}(M)$, to rewrite (2.4) as follows

$$(2.5) \quad \nabla \xi = \rho g + \varphi,$$

where $\varphi(X, Y) := g(\varphi(X), Y)$. Moreover, we adopt the following expression for the curvature tensor

$$R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z.$$

Furthermore, on a Riemannian manifold (M^n, g) , the Ricci operator S is defined using the Ricci tensor Ric (see [1]) by

$$(2.6) \quad Ric(X, Y) = g(SX, Y), \quad X, Y \in \mathfrak{X}(M).$$

Whence, one can use Equation (2.4) to get

$$(2.7) \quad R(X, Y)\xi = X(\rho)Y - Y(\rho)X + (\nabla\varphi)(X, Y) - (\nabla\varphi)(Y, X),$$

where $(\nabla\varphi)(X, Y) = \nabla_X(\varphi Y) - \varphi(\nabla_X Y)$.

Using the above equation and the expression for the Ricci tensor

$$Ric(X, Y) = \sum_{i=1}^n g(R(e_i, X)Y, e_i),$$

where $\{e_1, \dots, e_n\}$ is a local orthonormal frame, we obtain

$$(2.8) \quad Ric(\xi, Y) = -(n-1)Y(\rho) - \sum_{i=1}^n g(Y, (\nabla\varphi)(e_i, e_i)),$$

where we used the skew symmetry of the operator φ . The above relation gives

$$(2.9) \quad S(\xi) = -(n-1)\nabla\rho - \sum_{i=1}^n (\nabla\varphi)(e_i, e_i).$$

Now, using again, Equation (2.4), we compute the action of the rough Laplace operator Δ on the vector field ξ and find

$$(2.10) \quad \Delta\xi = \nabla\rho + \sum_{i=1}^n (\nabla\varphi)(e_i, e_i).$$

The following lemma, obtained previously in [[12], Lemma 1], will be useful.

Lemma 1. *Let (M^n, g) be a smooth compact Riemannian manifold with smooth totally geodesic boundary ∂M and ξ a smooth conformal vector field on M with conformal factor ρ satisfying $\rho|_{\partial M} = 0$. Denote by div and $\text{div}_{\partial M}$ the divergence operators on M and ∂M , respectively, and by ξ^T the tangential part of ξ on ∂M . Then,*

$$(2.11) \quad \text{div}(\xi) = n\rho, \quad \text{div}_{\partial M}(\xi^T) = (n-1)\rho.$$

Furthermore,

$$(2.12) \quad \int_M g(\nabla\rho, \xi) dM = -n \int_M \rho^2 dM.$$

We present below properties that are valid, in general, for conformal vector fields.

Lemma 2. *Let ξ be a non trivial conformal vector field on a Riemannian manifold (M^n, g) with conformal factor ρ . Then*

$$(2.13) \quad \begin{aligned} & \text{(i)} \quad \mathcal{L}_\xi \mathring{Ric} + (n-2)\mathring{\nabla}^2\rho = 0. \\ & \text{(ii)} \quad \text{div}(\mathcal{L}_\xi \mathring{Ric}) = -(n-2)\mathring{Ric}(\nabla\rho) - \frac{(n-1)(n-2)}{n}\nabla\Delta\rho - \frac{(n-2)R}{n}\nabla\rho. \text{ When the} \\ & \quad \text{sacalar curvature } R \text{ is constant, we have} \\ & \text{(iii)} \quad \text{div}(\mathcal{L}_\xi \mathring{Ric}) = -(n-2)\mathring{Ric}(\nabla\rho). \end{aligned}$$

$$\begin{aligned} \text{div}(\rho^m \mathcal{L}_\xi \mathring{Ric}(\nabla\rho)) &= -\rho^{m-1} [m(n-2)\mathring{\nabla}^2\rho(\nabla\rho, \nabla\rho) + \rho g(\text{div}(\mathcal{L}_\xi \mathring{Ric}), \nabla\rho) \\ & \quad + (n-2)\rho|\mathring{\nabla}^2\rho|^2]. \end{aligned}$$

Moreover,

$$\int_M \rho^{m-1} \left(m(n-2)\mathring{\nabla}^2\rho(\nabla\rho, \nabla\rho) + \rho g(\text{div}(\mathcal{L}_\xi \mathring{Ric}), \nabla\rho) + (n-2)\rho|\mathring{\nabla}^2\rho|^2 \right) dM = 0.$$

Proof. For the first item we will use Lemma 3.2 due to Hwang and Yun [15] (see also, e.g., [21, 22]), which gives

$$\mathcal{L}_\xi \mathring{Ric} = -(n-2)\nabla^2\rho + \frac{n-2}{n}\Delta\rho g = -(n-2)\mathring{\nabla}^2\rho.$$

The first part of item (ii) follows immediately from Bochner formula. Furthermore, if the scalar curvature R is constant, then, by Lemma 2.2 in [15], ρ satisfies

$$(2.14) \quad \Delta\rho = -\frac{R}{n-1}\rho,$$

which give us

$$\text{div}(\mathcal{L}_\xi \mathring{Ric}) = -(n-2)\mathring{Ric}(\nabla\rho).$$

For the last item, we observe that

$$\begin{aligned} \operatorname{div}(\rho^m \mathcal{L}_\xi \mathring{Ric}(\nabla \rho)) &= m\rho^{m-1} \mathcal{L}_\xi \mathring{Ric}(\nabla \rho, \nabla \rho) + \rho^m \operatorname{div}(\mathcal{L}_\xi \mathring{Ric}(\nabla \rho)) \\ &= m\rho^{m-1} \mathcal{L}_\xi \mathring{Ric}(\nabla \rho, \nabla \rho) + \rho^m g(\operatorname{div}(\mathcal{L}_\xi \mathring{Ric}), \nabla \rho) + \rho^m g(\mathcal{L}_\xi \mathring{Ric}, \nabla^2 \rho). \end{aligned}$$

Next we use the first item to deduce

$$\begin{aligned} \operatorname{div}(\rho^m \mathcal{L}_\xi \mathring{Ric}(\nabla \rho)) &= -m(n-2)\rho^{m-1} \mathring{\nabla}^2 \rho(\nabla \rho, \nabla \rho) + \rho^m g(\operatorname{div}(\mathcal{L}_\xi \mathring{Ric}), \nabla \rho) \\ &\quad - (n-2)\rho^m |\mathring{\nabla}^2 \rho|^2 \\ &= -\rho^{m-1} [m(n-2) \mathring{\nabla}^2 \rho(\nabla \rho, \nabla \rho) + \rho g(\operatorname{div}(\mathcal{L}_\xi \mathring{Ric}), \nabla \rho) \\ &\quad + (n-2)\rho |\mathring{\nabla}^2 \rho|^2]. \end{aligned}$$

So, the integral equation follows immediately by integrating equation contained in the third item. \square

The following proposition will be used in the proof of the Theorems 1 and 3.

Proposition 1. *Let ξ be a non trivial conformal vector field with conformal factor ρ on a compact Riemannian manifold (M^n, g) with smooth totally geodesic boundary ∂M of dimension $n \geq 3$ and $\rho|_{\partial M} = 0$. If the scalar curvature R of (M, g) is constant, then*

$$(2.15) \quad \int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\mathring{\nabla}^2 \rho|^2 \right) dM = 0,$$

or, equivalently,

$$(2.16) \quad \int_M \left(Ric(\nabla \rho, \nabla \rho) + |\nabla^2 \rho|^2 - (\Delta \rho)^2 \right) dM = 0.$$

Proof. By relation (2.14), we have $R > 0$, since $\rho = 0$ on ∂M . Taking the divergence of item (i) in Lemma 2, using Bochner formula, and (2.13), we get

$$\begin{aligned} \operatorname{div}(\mathcal{L}_\xi \mathring{Ric}(\nabla \rho)) &= \langle \mathcal{L}_\xi \mathring{Ric}, \nabla^2 \rho \rangle + \operatorname{div}(\mathcal{L}_\xi \mathring{Ric}, \nabla \rho) \\ &= -(n-2) |\mathring{\nabla}^2 \rho|^2 - (n-2) \mathring{Ric}(\nabla \rho, \nabla \rho). \end{aligned}$$

Since $n \geq 3$, using once more (2.13), integrating over M , and applying the Divergence Theorem, one gets

$$\int_{\partial M} \mathring{\nabla}^2 \rho(\nabla \rho, \nu) d\sigma = \int_M \left(|\mathring{\nabla}^2 \rho|^2 + \mathring{Ric}(\nabla \rho, \nabla \rho) \right) dM,$$

where ν is a unit outward normal vector field along ∂M . Nevertheless, choosing a local orthonormal frame $\{e_1, \dots, e_n\}$, such that $e_n = \nu$, we have

$$\begin{aligned} \int_{\partial M} \mathring{\nabla}^2 \rho(\nabla \rho, \nu) d\sigma &= \int_{\partial M} \left(\nabla^2 \rho(\nabla \rho, \nu) - \frac{\Delta \rho}{n} g(\nabla \rho, \nu) \right) d\sigma \\ &= \int_{\partial M} \left(\sum_{i=1}^n \rho_{i\nu} \rho_i - \frac{1}{n} \sum_{i=1}^n \rho_{ii} \rho_\nu \right) d\sigma \\ &= \int_{\partial M} \left(\sum_{i=1}^{n-1} \rho_{i\nu} \rho_i - \frac{1}{n} \sum_{i=1}^{n-1} \rho_{ii} \rho_\nu + \left(1 - \frac{1}{n}\right) \rho_{\nu\nu} \rho_\nu \right) d\sigma. \end{aligned}$$

Using that ∂M is totally geodesic one gets

$$\sum_{i=1}^{n-1} \rho_{ii} = \Delta^{\partial M} \rho \quad \text{and} \quad \sum_{i=1}^{n-1} \rho_{i\nu} \rho_i = g(\nabla^{\partial M} \rho_\nu, \nabla^{\partial M} \rho).$$

Applying the Divergence Theorem on ∂M we obtain

$$\begin{aligned} \int_{\partial M} \mathring{\nabla}^2 \rho (\nabla \rho, \nu) d\sigma &= \int_{\partial M} \left(g(\nabla^{\partial M} \rho_\nu, \nabla^{\partial M} \rho) - \frac{1}{n} \rho_\nu \Delta^{\partial M} \rho + \left(1 - \frac{1}{n}\right) \rho_{\nu\nu} \rho_\nu \right) d\sigma \\ &= - \int_{\partial M} \left(\rho \Delta^{\partial M} \rho_\nu - \frac{1}{n} \rho \Delta^{\partial M} \rho_\nu + \left(1 - \frac{1}{n}\right) \rho_{\nu\nu} \rho_\nu \right) d\sigma \\ &= \frac{(n-1)}{n} \int_{\partial M} \rho_{\nu\nu} \rho_\nu d\sigma, \end{aligned}$$

and therefore,

$$(2.17) \quad \int_M \left(|\mathring{\nabla}^2 \rho|^2 + \mathring{Ric}(\nabla \rho, \nabla \rho) \right) dM = \frac{(n-1)}{n} \int_{\partial M} \rho_{\nu\nu} \rho_\nu d\sigma.$$

Using relation (2.14) in Bochner formula and integrating over M , we have

$$\frac{1}{2} \int_M \operatorname{div}(\nabla |\nabla \rho|^2) dM = \int_M \left(|\nabla^2 \rho|^2 + Ric(\nabla \rho, \nabla \rho) - \frac{R}{n-1} |\nabla \rho|^2 \right) dM.$$

Since $\operatorname{div}(\rho \nabla \rho) = -\frac{R}{n-1} \rho^2 + |\nabla \rho|^2$ and $\rho = 0$ on ∂M , we have

$$\frac{1}{2} \int_{\partial M} \langle \nabla |\nabla \rho|^2, \nu \rangle d\sigma = \int_M \left(Ric(\nabla \rho, \nabla \rho) + |\nabla^2 \rho|^2 - (\Delta \rho)^2 \right) dM.$$

Noting that $\mathring{Ric}(\nabla \rho, \nabla \rho) = Ric(\nabla \rho, \nabla \rho) - \frac{R}{n} |\nabla \rho|^2$, and $\int_M |\nabla \rho|^2 dM = \frac{R}{n-1} \int_M \rho^2 dM$, we get

$$\int_{\partial M} \langle \nabla_{\nabla \rho} \nabla \rho, \nu \rangle d\sigma = \int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\nabla^2 \rho|^2 - \frac{1}{n} (\Delta \rho)^2 \right) dM.$$

We already showed that

$$\int_{\partial M} \langle \nabla_{\nabla \rho} \nabla \rho, \nu \rangle d\sigma = \int_{\partial M} \left(g(\nabla^{\partial M} \rho_\nu, \nabla^{\partial M} \rho) + \rho_{\nu\nu} \rho_\nu \right) d\sigma = \int_{\partial M} \rho_{\nu\nu} \rho_\nu d\sigma,$$

hence,

$$(2.18) \quad \int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\mathring{\nabla}^2 \rho|^2 \right) dM = \int_{\partial M} \rho_{\nu\nu} \rho_\nu d\sigma.$$

By Equations (2.17) and (2.18), we achieve $\int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\mathring{\nabla}^2 \rho|^2 \right) dM = 0$, that finishes the proof of the proposition. \square

Next, we present a useful lemma which is an important tool to prove our main results.

Lemma 3. *Let ξ be a nontrivial conformal vector field on a Riemannian manifold (M^n, g) with conformal factor ρ . Then*

- (i) $\operatorname{div}(|\mathring{Ric}|^2 \xi) = -2(n-2) \langle \nabla^2 \rho, \mathring{Ric} \rangle + (n-4) \rho |\mathring{Ric}|^2$.
- (ii) $\operatorname{div}(\rho |\mathring{Ric}|^2 \xi) = |\mathring{Ric}|^2 \langle \nabla \rho, \xi \rangle - 2(n-2) \rho \langle \nabla^2 \rho, \mathring{Ric} \rangle + (n-4) \rho^2 |\mathring{Ric}|^2$.
- (iii) $n \operatorname{div}(|\mathring{Ric}|^2 \xi + 2(n-2) \mathring{Ric}(\nabla \rho)) = (n-2)^2 \langle \nabla R, \nabla \rho \rangle + n(n-4) \rho |\mathring{Ric}|^2$.

Proof. In fact, firstly we note that

$$\operatorname{div}(|\mathring{Ric}|^2 \xi) = \langle \nabla |\mathring{Ric}|^2, \xi \rangle + n \rho |\mathring{Ric}|^2.$$

By Yano [22], we have

$$\mathcal{L}_\xi |\mathring{Ric}|^2 = -2(n-2) \langle \nabla^2 \rho, \mathring{Ric} \rangle - 4 \rho |\mathring{Ric}|^2,$$

and using that $\mathcal{L}_\xi |\mathring{Ric}|^2 = \langle \nabla |\mathring{Ric}|^2, \xi \rangle$, we obtain

$$\operatorname{div}(|\mathring{Ric}|^2 \xi) = -2(n-2)\langle \nabla^2 \rho, \mathring{Ric} \rangle + (n-4)\rho |\mathring{Ric}|,$$

which establishes (i).

On the other hand, from the first item we obtain

$$\operatorname{div}(\rho |\mathring{Ric}|^2 \xi) = |\mathring{Ric}|^2 \langle \nabla \rho, \xi \rangle - 2(n-2)\rho \langle \nabla^2 \rho, \mathring{Ric} \rangle + (n-4)\rho^2 |\mathring{Ric}|^2.$$

Now using that $\operatorname{div}(\mathring{Ric}(\nabla \rho)) = (\operatorname{div} \mathring{Ric})(\nabla \rho) + \langle \mathring{Ric}, \nabla^2 \rho \rangle$, and $\operatorname{div} \mathring{Ric} = \frac{n-2}{2n} \nabla R$, we have

$$\langle \mathring{Ric}, \nabla^2 \rho \rangle = \operatorname{div}(\mathring{Ric}(\nabla \rho)) - \frac{n-2}{2n} \langle \nabla R, \nabla \rho \rangle,$$

and, by the first item, again, we infer

$$\operatorname{div}(|\mathring{Ric}|^2 \xi) = -2(n-2) \operatorname{div}(\mathring{Ric}(\nabla \rho)) + \frac{(n-2)^2}{n} \langle \nabla R, \nabla \rho \rangle + (n-4)\rho |\mathring{Ric}|^2,$$

which implies that

$$(2.19) \quad n \operatorname{div}(|\mathring{Ric}|^2 \xi + 2(n-2)\mathring{Ric}(\nabla \rho)) = (n-2)^2 \langle \nabla R, \nabla \rho \rangle + n(n-4)\rho |\mathring{Ric}|^2.$$

Thus we finish the proof of the lemma. \square

Now we present some properties when we have a 4-dimensional manifold.

Proposition 2. *Let (M^4, g) be a Riemannian manifold with constant scalar curvature and ξ be a nontrivial conformal vector field on M^4 with conformal factor ρ . If $f : M^4 \rightarrow \mathbb{R}$ is a smooth function, then*

$$(2.20) \quad \operatorname{div}\left(f^m(|\mathring{Ric}|^2 \xi + 4\mathring{Ric}(\nabla \rho))\right) = |\mathring{Ric}|^2 \langle \xi, \nabla f^m \rangle + 4\mathring{Ric}(\nabla \rho, \nabla f^m).$$

Proof. In fact, by item (iii) of Lemma 3, since the scalar curvature is constant, and $n = 4$, we have $\operatorname{div}(|\mathring{Ric}|^2 \xi + 4\mathring{Ric}(\nabla \rho)) = 0$. Hence,

$$\begin{aligned} \operatorname{div}\left(f^m(|\mathring{Ric}|^2 \xi + 4\mathring{Ric}(\nabla \rho))\right) &= \langle |\mathring{Ric}|^2 \xi + 4\mathring{Ric}(\nabla \rho), \nabla f^m \rangle \\ &= |\mathring{Ric}|^2 \langle \xi, \nabla f^m \rangle + 4\mathring{Ric}(\nabla \rho, \nabla f^m). \end{aligned}$$

\square

Consequently, we have the following corollary.

Corollary 1. *Let (M^4, g) be a Riemannian manifold with constant scalar curvature R , and let ξ be a non trivial conformal vector field on M^4 with conformal factor ρ . If M^4 is either closed, or, compact with smooth boundary ∂M , where $\rho = 0$ on ∂M , then the following holds:*

$$\int_M \left(|\mathring{Ric}|^2 \langle \xi, \nabla \rho \rangle + 4\mathring{Ric}(\nabla \rho, \nabla \rho) \right) dM = 0.$$

3. PROOF OF THEOREMS

In this section, we will present the proofs of the theorems.

3.1. Proof of Theorem 1. Since $\mathring{Ric}(\nabla \rho) = 0$, by hypothesis, using Proposition 1, we deduce

$$\int_M |\mathring{\nabla}^2 \rho|^2 dM = 0.$$

Therefore, by the hypothesis on ρ , we can apply [[18], Theorem B] to ensure that M is isometric to a geodesic ball on $\mathbb{S}^n(c)$. Since ∂M is totally geodesic, we conclude that M is isometric to a hemisphere $\mathbb{S}_+^n(c)$.

3.2. Proof of Theorem 2. Using Bochner formula one gets

$$\int_M \left(|\nabla^2 \rho|^2 + Ric(\nabla \rho, \nabla \rho) + \langle \nabla \Delta \rho, \nabla \rho \rangle \right) dM = 0.$$

Since R is constant, we have $\Delta \rho = -\frac{R}{n-1}\rho$. Then,

$$\int_M \left(|\nabla^2 \rho|^2 + Ric(\nabla \rho, \nabla \rho) - \frac{R}{n-1} |\nabla \rho|^2 \right) dM = 0.$$

Using once more that ρ is an eigenfunction, we establish that $\int_M |\nabla \rho|^2 dM = \frac{R}{n-1} \int_M \rho^2 dM$.

On the other hand, $\mathring{Ric}(\nabla \rho, \nabla \rho) = Ric(\nabla \rho, \nabla \rho) - \frac{R}{n} |\nabla \rho|^2$ and $|\mathring{\nabla}^2 \rho|^2 = |\nabla^2 \rho|^2 - \frac{1}{n} (\Delta \rho)^2$, give us

$$\int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\mathring{\nabla}^2 \rho|^2 \right) dM = 0.$$

Using Corollary 1, we finally get

$$\int_M |\mathring{Ric}|^2 \langle \xi, \nabla \rho \rangle dM = 4 \int_M |\mathring{\nabla}^2 \rho|^2 dM.$$

Therefore, by the hypothesis we assume that $\int_M |\mathring{Ric}|^2 \langle \xi, \nabla \rho \rangle dM \leq 0$. Whence we can use Obata's result [17] to conclude that (M^4, g) is isometric to a standard sphere $\mathbb{S}^4(c)$.

3.3. Proof of Theorem 3. In fact, by Proposition 1, we have

$$\int_M \left(\mathring{Ric}(\nabla \rho, \nabla \rho) + |\mathring{\nabla}^2 \rho|^2 \right) dM = 0.$$

Combining with Corollary 1, we get

$$\int_M |\mathring{Ric}|^2 \langle \xi, \nabla \rho \rangle dM = 4 \int_M |\mathring{\nabla}^2 \rho|^2 dM.$$

Therefore, by the hypothesis on ρ , we can apply [[18], Theorem B] to conclude that M is isometric to a geodesic ball on $\mathbb{S}^4(c)$. Since ∂M is totally geodesic, we conclude that M is isometric to a hemisphere $\mathbb{S}_+^4(c)$.

3.4. Proof of Theorem 4. Using the relation (2.2) and item (ii) in Lemma 3, we obtain

$$\operatorname{div}(\rho |\mathring{Ric}|^2 \xi) = |\mathring{Ric}|^2 \langle \nabla \rho, \xi \rangle - n |\mathring{\nabla}^2 \rho|^2.$$

Integrating over M and using the Divergence Theorem, one gets

$$(3.1) \quad \int_M |\mathring{Ric}|^2 \langle \nabla \rho, \xi \rangle dM = n \int_M |\mathring{\nabla}^2 \rho|^2 dM.$$

Hence, by Equations (2.2) and (3.1), we have

$$\int_M |\mathring{Ric}|^2 \psi dM = 0,$$

where $\psi = n\rho^2 - \langle \nabla \rho, \xi \rangle$ as defined in the theorem. Since $\psi \neq 0$, we obtain $|\mathring{Ric}|^2 = 0$.

Again, by relation (2.2) and the definition of $\mathring{\nabla}^2 \rho$, yield

$$(3.2) \quad \nabla^2 \rho = \frac{\Delta \rho}{n} g.$$

Since $\Delta \rho = -\frac{R}{n-1}\rho$ and $\rho = 0$ on ∂M , it is not difficult to prove that $R > 0$. Then,

$$\nabla^2 \rho = -\frac{R\rho}{n(n-1)} g.$$

Therefore, by the hypothesis on ρ , we can apply [[18], Theorem B] to guarantee that M is isometric to a geodesic ball on $\mathbb{S}^n(c)$. Since ∂M is totally geodesic, we conclude that M is isometric to a hemisphere $\mathbb{S}_+^n(c)$.

REFERENCES

- [1] Besse, A.: Einstein manifolds. Springer-Verlag, Berlin (1987)
- [2] Boucher, W., Gibbons, G., Horowitz, G.: Uniqueness theorem for anti-de Sitter spacetime. *Physical Review D*. **30** (3), 2447–2451 (1984)
- [3] Caminha, A.: The geometry of closed conformal vector fields on Riemannian spaces. *Bulletin of the Brazilian Mathematical Society. New Series*. **42** (2), 277–300 (2011)
- [4] Corvino, J., Eichmair, M., Miao, P.: Deformation of scalar curvature and volume. *Math. Ann.* **357**, 551–584 (2013)
- [5] Deshmukh, S.: Characterizing spheres by conformal vector fields. *Ann. Univ. Ferrara*. **56**, 231–236 (2010)
- [6] Deshmukh, S., Al-Solamy, F.: Conformal gradient vector fields on a compact Riemannian manifold. *Colloquium Math.* **112** (1), 157–161 (2008)
- [7] Deshmukh, S., Alsolamy, F.: Conformal vector fields and conformal transformations on a Riemannian manifold. *Balkan J. Geom. Appl.* **17** (1), 9–16 (2012)
- [8] Deshmukh, S., Alsolamy, F.: Conformal vector fields on a Riemannian manifold. *Balkan J. Geom. Appl.* **19** (2), 86–93 (2014)
- [9] Deshmukh, S., Ishan, A., Vilcu, G.: Conformal Vector Fields and the De-Rham Laplacian on a Riemannian Manifold. *Mathematics*. **9** (8), 863–871 (2021)
- [10] Deshmukh, S.: Geometry of conformal vector fields. *Arab J. Math. Sci.* **23**, 44–73 (2017)
- [11] Ejiri, N.: A negative answer to a conjecture of conformal transformations of Riemannian manifolds. *J. Math. Soc. Jpn.* **33** (2), 261–266 (1981)
- [12] Evangelista, I., Viana, E.: Conformal gradient vector fields on Riemannian manifolds with boundary. *Colloq. Math.* **159**, 231–241 (2020)
- [13] Evangelista, I., Freitas, A., Viana, E.: Conformal vector fields and the de-Rham Laplacian on a Riemannian manifold with boundary. *Results Math.* **77**, 217–231 (2022)
- [14] Hicks, N.: Closed vector fields. *Pacific Journal of Mathematics*. **15** (1), 141–151 (1965)
- [15] Hwang, S., Yun, G.: Conformal Vector Fields and Their Applications to Einstein-Type Manifolds. *Results Math.* **79**, 45–64 (2024)
- [16] Kühne, W., Rademacher, H.: Conformal vector fields on pseudo-Riemannian spaces. *Differential Geom. Appl.* **7**, 237–250 (1997)
- [17] Obata, M.: Certain conditions for a Riemannian manifold to be isometric with a sphere. *J. Math. Soc. Jpn.* **14** (3), 333–340 (1962)
- [18] Reilly, R. C.: Geometric applications of the solvability of Neumann problems on a Riemannian manifold. *Arch. Ration. Mech. Anal.* **75**, 23–29 (1980)
- [19] Tanno, S., Weber, W. C.: Closed conformal vector fields. *Journal of Differential Geometry*. **3**, 361–366 (1969)
- [20] Takeno, H.: Conircular scalar field in spherically symmetric space-times I, Hiroshima Univ. (1969)
- [21] Yano, K.: *Integral Formulas in Riemannian Geometry*. Marcel Dekker Inc, New York (1970)
- [22] Yano, K., Obata, M.: Conformal changes of Riemannian metrics. *J. Differ. Geom.* **4**, 53–72 (1970)

(A. Barros) UNIVERSIDADE FEDERAL DO CEARÁ - UFC, DEPARTAMENTO DE MATEMÁTICA, CAMPUS DO PICI, AV. HUMBERTO MONTE, BLOCO 914, 60455-760, FORTALEZA / CE, BRAZIL.

Email address: abbarros@mat.ufc.br

(I. Evangelista) UNIVERSIDADE FEDERAL DO DELTA DO PARNAÍBA- UFDPAR, CURSO DE MATEMÁTICA, AV. SÃO SEBASTIÃO, 2819, 64202-020, PARNAÍBA /PI, BRAZIL

Email address: evangelistaisrael@ufdpar.edu.br

(E. Viana) INSTITUTO FEDERAL DE EDUCAÇÃO, CIÊNCIA E TECNOLOGIA DO CEARÁ - CAMPUS CAUCAIA, 61609-090 CAUCAIA/CE, BRAZIL.

Email address: emanuel.mendonca@ifce.edu.br