# RIGIDITY RESULTS FOR A TRIPLE JUNCTION SOLUTION OF ALLEN-CAHN SYSTEM

### ZHIYUAN GENG

ABSTRACT. For the two dimensional Allen-Cahn system with a triple-well potential, previous results established the existence of a minimizing solution  $u : \mathbb{R}^2 \to \mathbb{R}^2$  with a triple junction structure at infinity. We show that along each of three sharp interfaces, u is asymptotically invariant in the direction of the interface and can be well-approximated by the 1D heteroclinic connections between two phases. Consequently, the diffuse interface is located in an O(1) neighborhood of the sharp interface, and becomes nearly flat at infinity. This generalizes all the results for the triple junction solution with symmetry hypotheses to the non-symmetric case. The proof relies on refined sharp energy lower and upper bounds, alongside a precise estimate of the diffuse interface location.

#### 1. INTRODUCTION

In this paper, we investigate the bounded entire solution of the system

(1.1) 
$$\Delta u - W_u(u) = 0, \quad u : \mathbb{R}^2 \to \mathbb{R}^2,$$

which is minimizing on compact sets the associated Allen-Cahn energy:

$$E(u,\Omega) := \int_{\Omega} \left( \frac{1}{2} |\nabla u|^2 + W(u) \right) \, dx.$$

Specifically for W we assume

(H1).  $W \in C^2(\mathbb{R}^2; [0, +\infty)), \{z : W(z) = 0\} = \{a_1, a_2, a_3\}, W_u(u) \cdot u > 0 \text{ if } |u| > M \text{ and}$  $c_2|\xi^2| \ge \xi^T W_{uu}(a_i)\xi \ge c_1|\xi|^2, \ i = 1, 2, 3.$ 

for some positive constants  $c_1 < c_2$  depending on W.

(H2). For  $i \neq j$ ,  $i, j \in \{1, 2, 3\}$ , there exists a unique (up to translation) minimizing heteroclinic connection  $U_{ij} \in W^{1,2}(\mathbb{R}, \mathbb{R}^2)$  that minimizes the one dimensional energy functional

$$J(U) := \int_{\mathbb{R}} \left( \frac{1}{2} |U'|^2 + W(U) \right) d\eta, \quad \lim_{\eta \to -\infty} U(\eta) = a_i, \ \lim_{\eta \to +\infty} U(\eta) = a_j.$$

The connection  $U_{ij}$  is non-degenerate in the sense that 0 is a simple eigenvalue of the operator  $\mathcal{T}: W^{2,2}(\mathbb{R},\mathbb{R}^2) \to L^2(\mathbb{R},\mathbb{R}^2)$ :

(1.2) 
$$\mathcal{T}\varphi := -\varphi'' + W_{uu}(U_{ij})\varphi.$$

Let  $\sigma_{ij}$  denote the minimal energy  $J(U_{ij})$ . Assume

(1.3) 
$$\sigma_{ij} \equiv \sigma, \quad \forall i \neq j \in \{1, 2, 3\}$$

Department of Mathematics, Purdue University, 150 N. University Street, West Lafayette, IN  $47907\mathchar`-2067$ 

*E-mail address*: geng42@purdue.edu.

Key words and phrases. Allen-Cahn system, diffuse interface, triple junction solution, heteroclinic connection, De Giorgi conjecture.

The minimizing solution we seek for is a diffuse analogue of the planar minimizing partition. Let  $\Omega$  (could be the whole  $\mathbb{R}^2$ ) denote a two dimensional domain. A partition  $\mathcal{P} = \{P_1, P_2, P_3\}$  of  $\Omega$  is called a locally minimizing partition of  $\Omega$ , if for any compact set  $K \subset \Omega$ ,

$$E_0(\{P_1, P_2, P_3\}, K) \le E_0(\{A_1, A_2, A_3\}, K),$$

where  $E_0(\cdot)$  denotes the functional

(1.4) 
$$E_0(\{D_1, D_2, D_3\}, K) := \sum_{1 \le i < j \le 3} \sigma \mathcal{H}^1(\partial^* D_i \cap \partial^* D_j \cap K),$$

and  $\mathcal{A} = \{A_i\}_{i=1}^3$  is any 3-partition of  $\Omega$  such that  $\mathcal{P} \sqcup (\Omega \setminus K) = \mathcal{A} \sqcup (\Omega \setminus K)$ . Here  $\partial^*$  refers to the reduced boundary of a set of finite perimeter. Analogously, for any function  $v \in BV_{loc}(\mathbb{R}^2, \{a_1, a_2, a_3\})$ , we can define the energy functional

(1.5) 
$$E_0^*(v,K) := E_0(\{v^{-1}(a_i)\}_{i=1}^3, K).$$

v is called a minimizing partition map on  $\Omega$  if v minimizes  $E_0^*(\cdot, K)$  for any  $K \in \Omega$ .

In particular, when  $\Omega = \mathbb{R}^2$ , we consider the following minimizing 3-partition of  $\mathbb{R}^2$ ,

(1.6)  
$$\mathcal{P} := \{\mathcal{D}_1, \mathcal{D}_2, \mathcal{D}_3\},\$$
$$\mathcal{D}_i := \{(r\cos\theta, r\sin\theta) : r \in (0, \infty), \theta \in (\frac{2(i-1)\pi}{3}, \frac{2i\pi}{3})\}, i = 1, 2, 3,$$

which is a partition of the plane into three sectors centered at the origin with opening angles of  $\frac{2\pi}{3}$ . The sharp interface that separates these sub-domains is denoted by

$$\partial \mathcal{P} := \partial \mathcal{D}_1 \cup \partial \mathcal{D}_2 \cup \partial \mathcal{D}_3.$$

The *triple junction map* is defined by

(1.7) 
$$u_{\mathcal{P}} := a_1 \chi_{\mathcal{D}_1} + a_2 \chi_{\mathcal{D}_2} + a_3 \chi_{\mathcal{D}_3}$$

where  $\chi_{\Omega}$  represents the characteristic function of domain  $\Omega$ . The minimality of  $\mathcal{P}$  and  $u_{\mathcal{P}}$  defined above is related to the Steiner point of triangle in classical Euclidean geometry.

Here is our main result.

**Theorem 1.1.** Let  $u : \mathbb{R}^2 \to \mathbb{R}^2$  be a minimizing entire solution of (1.1) with a triple junction structure at infinity, i.e.

(1.8) 
$$u(Rz) \xrightarrow{L^1_{loc}} u_{\mathcal{P}} \quad as \ R \to \infty.$$

For any  $i, j \in \{1, 2, 3\}$ , let  $\mathbf{e}_{ij}$  denote the unit vector representing the direction of  $\partial \mathcal{D}_i \cap \partial \mathcal{D}_j$  and  $\mathbf{e}_{ij}^{\perp}$  as its orthogonal unit vector. Then there exists a constant  $h_{ij}$ , such that

(1.9) 
$$\lim_{x \to +\infty} \|u(x\mathbf{e}_{ij} + y\mathbf{e}_{ij}^{\perp}) - U_{ij}(y - h_{ij})\|_{C^{2,\alpha}(\mathbb{R};\mathbb{R}^2)} = 0, \quad \forall \alpha \in (0,1).$$

Furthermore, the following pointwise estimate holds:

(1.10) 
$$|u(z) - a_i| \le C e^{-k \operatorname{dist}(z, \partial \mathcal{P})}, \quad \text{for } z \in \mathcal{D}_i,$$

where the positive constants C, k depend only on W and u.

*Remark* 1.1. We consider the case where all  $\sigma_{ij}$  are equal for simplicity. The result can be extended to the general  $\sigma_{ij}$  case without significant technical difficulty.

3

*Remark* 1.2. By [29, Proposition 4.3] and [21, Theorem 1.2], the condition (1.8) can be relaxed to the following: there exists a point  $z \in \mathbb{R}^2$  such that

$$\operatorname{dist}(u(z), \Lambda) > 0,$$

where  $\Lambda = \overline{U_{12}(\mathbb{R}) \cup U_{23}(\mathbb{R}) \cup U_{31}(\mathbb{R})}$  denotes the closure of three heteroclinic connections. This remarkably weak condition is sufficient to establish (1.8).

In the scalar case of (1.1), where  $u : \mathbb{R}^N \to \mathbb{R}$  and W(u) has only two energy wells  $u = \pm 1$ , the solutions are closely connected to the minimal surface theory, leading to the famous conjecture of De Giorgi [11]. The conjecture proposes that in dimension  $N \leq 8$ , any entire solution (not necessarily minimizing) that is monotonic in one variable only depends on that variable, meaning that all level sets are parallel hyperplanes orthogonal to the direction of monotonicity. There are many deep results contributing to the understanding of De Giorgi's conjecture and the relationship between Allen-Cahn equation and minimal surfaces, see for example [6, 14, 22, 30, 36, 12, 13, 23, 26, 28] and the expository papers [31, 10] for a detailed account.

In the vector-valued case, minimizing solutions are related to minimal partitions. The convergence of the vector-valued Allen-Cahn system to a minimal partition problem can be established by the  $\Gamma$ -convergence technique, see [7, 34, 16, 20] for further details. For the two dimensional case, recent developments have provided significant insights into the geometric and analytic description of fine structures of the minimizing Allen-Cahn solutions. Bethuel [8] discovered a new monotonicity formula and successfully generalized many results on the regularity of interfaces from the scalar case to the vectorial case. Fusco [19] studied the connectivity of the diffuse interface and showed the existence of a connected network with a well-defined structure that separates all phases. Alikakos and Fusco [3] investigated two examples with carefully designed Dirichlet boundary data, where a sharp lower bound for the energy minimizer could be derived, which in turn yielded precise pointwise estimates.

As for the triple junction solution on  $\mathbb{R}^2$ , the first existence result was due to Bronsard, Gui and Schatzman [9], where an entire solution to (1.1) was constructed in the equivariant class of the reflection group  $\mathcal{G}$  corresponding to the symmetries of the equilateral triangle. The triplewell potential is also assumed invariant under  $\mathcal{G}$ . The solution is obtained as a minimizer in the equivariant class  $u(gx) = gu(x), g \in \mathcal{G}$ , hence is not necessarily stable under general perturbations. The result was later extended to the three dimensional case by Gui and Schatzman [25]. More recently, Fusco [18] established the result of [9] in the equivariant class of the rotation subgroup of  $\mathcal{G}$  only, thus eliminating the two reflections. For a more detailed discussion, we refer to the book [4] and the references therein.

On bounded domains, several constructions of triple junction solutions without symmetry assumptions have been established. For instance, Sternberg and Ziemer [35] constructed solutions on clover-shaped domains in  $\mathbb{R}^2$  using  $\Gamma$ -convergence; while Flores, Padilla, and Tonegawa [15] extended the construction to more general domains via a mountain pass argument. Regarding entire solutions without symmetry assumptions, Schatzman [32] demonstrated the existence of a 2D solution connecting two heteroclinic connections for a double-well potential. Her result was later revisited in [17, 27, 33].

Recently, Alikakos and the author [5], and Sandier and Sternberg [29], independently established the existence of an entire minimizing solution, characterized by a triple junction structure at infinity without imposing symmetry assumptions. Using distinct methods, both studies obtained comparable results saying that along a subsequence  $R_k \to \infty$ , the rescaled function  $u_{R_k}(z) := u(R_k z)$ converges in  $L^1_{loc}(\mathbb{R}^2)$  to a triple junction map  $u_{\mathcal{P}}$  of the form (1.7). Following these two works, the author [21] showed the uniqueness of the blow-down limit  $u_{\mathcal{P}}$ , thus establishing that  $u(Rz) \to u_{\mathcal{P}}$ in  $L^1_{loc}$  as  $R \to \infty$ . These results will be discussed in more detail later in Section 2.

### RIGIDITY OF TRIPLE JUNCTION

A noteworthy byproduct of [29] is the first vectorial analogue of the De Giorgi's conjecture for 2D minimizing Allen-Cahn solutions of two phases. It states that if the blow-down limit of u consists of only two phases  $a_1$  and  $a_2$ , separated by a straight line, then u must be invariant along the direction of this line. This result follows from sharp energy lower and upper bounds, which force the directional derivative of u along the interface to be negligible, thereby ensuring the flatness of the interface. However, the same argument does not apply to the triple junction solution due to the lack of a sharp lower bound. The current paper addresses this challenge. Theorem 1.1 establishes that the diffuse interface defined in (3.18) forms a strip of width O(1), which is asymptotically flat at infinity, and that u is nearly an one-dimensional solution along the interface. This result generalizes the complete results of [9] to the non-symmetric setting, as well as confirms a De Giorgi-type conjecture for Allen Cahn solutions with triple phases.

We now outline some key steps in the proof of Theorem 1.1. We start with the minimizing entire solution constructed in [29] and [5], which satisfies condition (1.8) for the partition  $\mathcal{P}$  defined in (1.6). Moreover, as shown in [21], the intersection of the diffuse interface  $\Gamma_{\delta} := \{z : \min_{i=1,2,3} |u(z) - z_i| > \delta\}$  with  $\mathcal{P}_{\delta}$  is contained in an  $\mathcal{O}(\mathcal{P}^{\beta})$  pairs have a diffuse interface  $\mathcal{P}_{\delta}$  for a second start of the short interface  $\mathcal{P}_{\delta}$  for a second start of the short interface  $\mathcal{P}_{\delta}$  for a second start of the short interface  $\mathcal{P}_{\delta}$  for a second start of the second start of the short interface  $\mathcal{P}_{\delta}$  for a second start of the second st

 $a_i \geq \delta$  with  $B_R$  is contained in an  $O(R^{\beta})$  neighborhood of the sharp interface  $\partial \mathcal{P}$ , for some  $\beta \in (\frac{1}{2}, 1)$ . Outside this  $O(R^{\beta})$  neighborhood, the distance of u(z) to  $a_i$  is controlled by the estimate

$$|u(z) - a_i| \le C e^{-k(\operatorname{dist}(z,\partial \mathcal{P}) - CR^\beta)}, \quad z \in B_R \cap D_i \cap \{\operatorname{dist}(z,\partial \mathcal{P}) > CR^\beta\}.$$

This exponential decay implies that on any large circle  $\partial B_R$ , the restriction  $u|_{\partial B_R}$  approaches three phases respectively. Thus there must be at least three phase transitions along  $\partial B_R$ , which contribute a minimal energy of  $3\sigma$  in the tangential direction. Integrating with the radius R yields the sharp energy bound

(1.11) 
$$3\sigma R - C \le \int_{B_R} \left(\frac{1}{2}|\nabla u|^2 + W(u)\right) dz \le 3\sigma R + C,$$

as established in Lemma 3.3. From this energy bound, one can update the power  $\beta$ , which measures the closeness of  $\Gamma_{\delta}$  to  $\partial \mathcal{P}$ , to  $\frac{1}{2}$ .

Since the tangential deformation and the potential take most of the energy, the radial deformation is relatively small:

$$\int_{\mathbb{R}^2} |\partial_r u|^2 \, dx dy < \infty.$$

Assume the  $a_1$ - $a_3$  interface lies along the positive x-axis. Through technical arguments, it follows that

(1.12) 
$$\int_{\{x>0\}} |\partial_x u|^2 \, dx \, dy < \infty.$$

For most x > 0, the vertical slice  $l_x := \{(x, y) : y \in \mathbb{R}\}$  connects  $a_3$  to  $a_1$  from  $(x, -\infty)$  to  $(x, \infty)$ , with an energy close to  $\sigma$ . By [32], this implies that  $u|_{l_x}$  is close to the heteroclinic connection  $U_{31}$ in the  $H^1$  norm. Let h(x) denotes the optimal translation of  $U_{31}$  such that the  $L^2$  distance between  $u(x, \cdot)$  and  $U_{31}(\cdot - h(x))$  is minimized. As shown in [32], h(x) is a  $C^2$  function of x provided  $u(x, \cdot)$ stays sufficiently close to  $U_{31}$ . A direct calculation yields

$$|h'(x)| \sim C \int_{-\infty}^{\infty} |\partial_x u(x,y)|^2 \, dy,$$

which together with (1.12) implies that h(x) converges to some finite value  $h_{31}$  as  $x \to \infty$ . Consequently,  $u(x, \cdot)$  converges to  $U_{31}(\cdot - h_{31})$  in  $L^2$ . This  $L^2$  convergence can be upgraded to  $C^{2,\alpha}$ convergence by standard Schauder estimates, thus proving (1.9). Furthermore, this result indicates that the diffuse interface locates within an O(1) neighborhood of  $\partial \mathcal{P}$ . From this, the pointwise estimate (1.10) follows from the comparison principle in elliptic theory.

The article is organized as follows. In Section 2, we review fundamental estimates for the minimizing solution of (1.1) from [3, 4] and summarize results in [29, 5, 21] on the existence of an entire triple junction solution. The proof of Theorem 1.1 is presented in Section 3. We begin by proving (1.11) and refining the localization of the diffuse interface, then derive the horizontal deformation estimate (1.12), and finally conclude the proof by analyzing the optimal translation function h(x).

### 2. Preliminaries

Throughout the paper we denote by z = (x, y) a two dimensional point and by  $B_r(z)$  the two dimensional disk centered at the point z with radius r. In addition, we let  $B_r$  denote the disk centered at the origin. Moreover, without specific explanation, C denotes a constant that depends on the potential W and on the solution u. C may have distinct values in various estimates. We first recall the following basic results (without proofs) which play a crucial part in our analysis.

**Lemma 2.1** (Lemma 2.1 in [3]). The hypotheses on W imply the existence of  $\delta_W > 0$ , and constants c, C > 0 such that

$$\begin{aligned} |u - a_i| &= \delta \\ \Rightarrow \frac{1}{2}c\delta^2 \leq W(u) \leq \frac{1}{2}C\delta^2, \quad \forall \delta < \delta_W, \ i = 1, 2, 3. \end{aligned}$$

Moreover if  $\min_{1 \le i \le N} |u - a_i| \ge \delta$  for some  $\delta < \delta_W$ , then  $W(u) \ge \frac{1}{2}c\delta^2$ .

**Lemma 2.2** (Lemma 2.3 in [3]). Take  $i \neq j \in \{1, ..., N\}$ ,  $\delta < \delta_W$  and  $s_+ > s_-$  be two real numbers. Let  $v : (s_-, s_+) \to \mathbb{R}^2$  be a smooth map that minimizes the energy functional

$$J_{(s_{-},s_{+})}(v) := \int_{s_{-}}^{s_{+}} \left(\frac{1}{2}|\nabla v|^{2} + W(v)\right) dx$$

subject to the boundary condition

$$|v(s_{-}) - a_i| = |v(s_{+}) - a_j| = \delta.$$

Then

$$J_{(s_-,s_+)}(v) \ge \sigma_{ij} - C\delta^2,$$

where C is the constant in Lemma 2.1.

**Lemma 2.3** (Variational maximum principle, [2]). There exists a positive constant  $r_0 = r_0(W)$  such that for any  $u \in W^{1,2}(\Omega, \mathbb{R}^2) \cap L^{\infty}(\Omega, \mathbb{R}^2)$  being a minimizer of  $E(\cdot, \Omega)$ , if u satisfies

 $|u(x) - a_i| \le r \text{ on } \partial\Omega, \text{ for some } r < r_0, i \in \{1, 2, 3\},\$ 

then

$$|u(x) - a_i| \le r \quad \forall x \in \Omega.$$

Suppose u is a bounded minimizing solution of (1.1). Schauder estimates imply that there exists a constant M such that

(2.13) 
$$||u||_{C^{2,\alpha}(\mathbb{R}^2,\mathbb{R}^2)} \le M, \quad \forall \alpha \in (0,1).$$

To characterize the blowdown limits of u, we invoke the following compactness result.

**Proposition 2.4** (Proposition 3.1 in [29]). Let  $u : \mathbb{R}^2 \to \mathbb{R}^2$  be a minimizing solution of (1.1) and  $\{r_j\} \to \infty$  be any sequence. Then there exists a subsequence  $\{r_{j_k}\}$  and a function  $u_0 \in BV_{loc}(\mathbb{R}^2, \{a_i\}_{i=1}^N)$  such that the blowdowns  $\{u_{r_{j_k}}(z) = u(r_{j_k}z)\}$  satisfy

$$u_{r_{j_k}} \to u_0 \quad in \ L^1_{loc}(\mathbb{R}^2, \mathbb{R}^2)$$

Here  $u_0$  is a minimizing partition map on  $\mathbb{R}^2$  in the sense that

$$E_0^*(u_0, K) \le E_0^*(v, K)$$

for every compact set K and every  $v \in BV_{loc}(\mathbb{R}^2, \{a_i\}_{i=1}^3)$  such that  $v = u_0$  on  $\mathbb{R}^2 \setminus K$ . Along the same subsequence, the following energy estimate holds

(2.14) 
$$E_{r_{i_k}}(u_{r_{j_k}}, K) \to E_0^*(u_0, K), \quad \forall K \in \mathbb{R}^2,$$

where  $E_R$  denotes the rescaled energy

$$E_R(v,\Omega) := \int_{\Omega} \left( \frac{1}{2R} |\nabla u|^2 + RW(u) \right) \, dz.$$

Note that the energy convergence result (2.14) is not explicitly stated in [29, Proposition 3.1], but can be derived from the  $\Gamma$ -convergence result in Baldo [7] that holds also without the mass constraint (see Gazoulis [20]).

By a "clearing-out" argument, the  $L_{loc}^1$  convergence can be improved to uniform convergence outside the support of  $\nabla u_0$ .

**Proposition 2.5** (Proposition 4.2 in [29]). Let  $\{u_{r_j}\}$  be a sequence of blowdowns of the minimizing solution u that converges in  $L^1_{loc}$  to  $u_0 \in BV_{loc}(\mathbb{R}^2, \{a_i\}_1^N)$ . Then  $\{u_{r_j}\}$  converges to  $u_0$  uniformly outside the support of  $\nabla u_0$ .

The uniform convergence result above can also be derived from the the vector version of the Caffarelli–Córdoba density estimate [1], see for example [4, Proposition 5.3].

Moreover, utilizing a Pohozaev identity and estimates of energy lower and upper bounds, [29] establishes an asymptotic monotonicity formula and an energy equipartition result, which further implies the homogeneity of the blowdown limit  $u_0$ . Related results are summarized below.

**Proposition 2.6** (Lemma 3,4, Lemma 3.5 & Theorem 3.6 in [29]). Let  $u : \mathbb{R}^2 \to \mathbb{R}^2$  be a minimizing solution of (1.1) and  $u_0$  be a blowdown limit guaranteed by Proposition 2.4. Then  $u_0$  is a homogeneous map, i.e.

$$u_0(z) = u_0(\frac{z}{|z|}), \quad \forall z \in \mathbb{R}^2$$

The following energy limits hold:

(2.15) 
$$\lim_{R \to \infty} \frac{1}{R} \int_{B_R} W(u) \, dz = \lim_{R \to \infty} \frac{1}{R} \int_{B_R} \frac{1}{2} |\nabla u|^2 \, dz = \lim_{R \to \infty} \frac{1}{R} \int_{B_R} \frac{1}{2} |\partial_T u|^2 \, dz = \frac{1}{2} E_0^*(u_0, B_1),$$

where  $\partial_T$  means the tangential derivative on  $\partial B_R$ . Moreover, for every positive  $\lambda_1 < \lambda_2$  it holds that

$$\lim_{R \to \infty} \frac{1}{R} \int_{B_{\lambda_2 R} \setminus B_{\lambda_1 R}} |\partial_r u|^2 \, dz = 0,$$
$$\lim_{R \to \infty} \frac{1}{R} \int_{B_{\lambda_2 R} \setminus B_{\lambda_1 R}} \left| W(u) - \frac{1}{2} |\nabla u|^2 \right| \, dz = 0$$

where  $\partial_r u = \nabla u \cdot \frac{x}{|x|}$  represents the radial derivative of u.

Proposition 2.4 and 2.6 together imply the following theorem.

**Theorem 2.7** (Theorem 1.1 in [29]). There exists a minimizing entire solution  $u : \mathbb{R}^2 \to \mathbb{R}^2$  to (1.1) such that for any compact set  $K \subset \mathbb{R}^2$ ,

(2.16) 
$$\lim_{R \to \infty} \left( \inf_{\theta} \| u_R(z) - u_{\mathcal{P}}(G_{\theta}z) \|_{L^1(K;\mathbb{R}^2)} \right)$$

where  $u_R = u(Rz)$  is the blowdown of u,  $u_P$  is the triple junction map defined in (1.7) and  $G_{\theta}$  denotes the rotation through an angle  $\theta$  about the origin.

Equivalently, this theorem tells that along any sequence  $R_i \to \infty$ , there is a subsequence, still denoted by  $\{R_i\}$ , such that  $u_{R_i}$  converges in  $L^1_{loc}$  to a triple junction map which may depend on the sequence. A comparable result has been obtained in [5] by N. Alikakos and the author around the same time with a different method.

Finally, we mention the following uniqueness result of the blow-down limit which is proved by the author in [21], thereby eliminating the possible rotation  $G_{\theta}$  in (2.16).

**Theorem 2.8.** For the entire solution u in Theorem 2.7, there exists a unique triple junction map  $u_{\mathcal{P}}$ , such that

(2.17) 
$$\lim_{R \to \infty} \|u_R - u_\mathcal{P}\|_{L^1_{loc}(\mathbb{R}^2)} = 0.$$

## 3. Proof of Theorem 1.1

Throughout this section,  $u : \mathbb{R}^2 \to \mathbb{R}^2$  denotes a minimizing solution of (1.1) satisfying Theorem 2.7 and 2.8. The minimizing partition  $\mathcal{P}$  and its associated triple junction map  $u_{\mathcal{P}}$  are given by (1.6) and (1.7) respectively. To prove Theorem 1.1, we focus on the interface  $\partial \mathcal{D}_1 \cap \partial \mathcal{D}_3 = \{(x,0) : x > 0\}$ . The proof is identical for the other two interfaces. In this setting, we pick

 $\mathbf{e}_{13} := (1,0)$  is the directional unit vector of the interface,  $\mathbf{e}_{13}^{\perp} := (0,1)$ .

By the hypothesis (H2), there exists a unique (up to a translation) minimizing heteroclinic connection  $U_{31}$ , such that  $U_{31}(-\infty) = a_3$ ,  $U_{31}(+\infty) = a_1$ . For convenience, in the rest of the paper we drop the subscript and simply write

$$U = U_{31}.$$

Define the diffuse interface of u as

(3.18) 
$$\Gamma_{\delta} := \{ z \in \mathbb{R}^2 : \min_{i=1,2,3} |u(z) - a_i| \ge \delta \}, \quad \delta > 0$$

3.1. Localization of the diffuse interface. From Theorem 2.8, we know u can be approximated by  $u_{\mathcal{P}}$  at large scales. Next we invoke a more quantified version of this approximation from [21], which states that the diffuse interface is located in a small neighborhood of  $\partial \mathcal{P}$ .

**Proposition 3.1.** There exist  $R_0$ , C,  $\alpha \in (0,1)$ , K, k depending on W and u, such that for every  $R > R_0$ , the following hold:

(1) There are points  $O_R \in B_R$ ,  $D_R^1, D_R^2, D_R^3 \in \partial B_R$  such that the line segments  $O_R D_R^1, O_R D_R^2, O_R D_R^3$  form a  $\frac{2\pi}{3}$  angle pairwisely, and the following energy estimate holds:

(3.19) 
$$\left| E(u, B_R) - \sigma \mathcal{H}^1(T_R) \right| \le CR^{\alpha}$$

where  $T_R$  denotes  $O_R D_R^1 \cup O_R D_R^2 \cup O_R D_R^3$ .

(2) The diffuse interface locates in an  $O(R^{\frac{\alpha+1}{2}})$  neighborhood of  $T_R$ , i.e.

(3.20) 
$$\Gamma_{\delta} \cap B_R \subset \{ z \in B_R : \operatorname{dist}(z, T_R) \le CR^{\frac{\alpha+1}{2}} \}.$$

Moreover,

(3.21) 
$$\min_{i=1,2,3} |u(z) - a_i| \le K e^{-k(\operatorname{dist}(z,T_R) - CR^{\frac{\alpha+1}{2}})^+}, \ \forall z \in B_{\frac{3}{4}R},$$

where  $(a)^+ = \max\{a, 0\}.$ 

(3) Let  $O_R D_R^1$  be the approximated  $a_1$ - $a_3$  interface at the scale R, and let  $\theta_R$  denote the angle between  $O_R D_R^1$  and the direction  $\mathbf{e}_{13}$ . Then

$$(3.22) \qquad \qquad |\theta_R - \theta_{2R}| \le CR^{\frac{\alpha-1}{2}}.$$

*Proof.* All the proofs can be found in [21], and we therefore omit the details here. Specifically, (1) follows directly from Section 4, Proposition 5.1 and Proposition 5.3 in [21], while (2) and (3) correspond to Proposition 6.1 and Lemma 8.1, respectively.  $\Box$ 

Now we are in the position to derive the pointwise estimate of the distance of u(z) to energy wells away from the  $a_1$ - $a_3$  sharp interface  $\{(x, 0) : x > 0\}$ .

**Lemma 3.2.** There exist  $\beta \in (\frac{1}{2}, 1)$  and positive constants  $R_0, C, K, k$  only depending on W and u, such that for any  $x \ge R_0$ ,

(3.23)  $|u(x,y) - a_1| \le K e^{-k(y - Cx^\beta)}, \quad y \ge Cx^\beta,$ 

(3.24) 
$$|u(x,y) - a_3| \le K e^{-k(|y| - Cx^\beta)}, \quad y \le -Cx^\beta,$$

(3.25) 
$$|\nabla u(x,y)| \le K e^{-k(y-Cx^{\beta})}, \quad |y| \ge Cx^{\beta}.$$

*Proof.* Let  $\beta = \frac{1+\alpha}{2}$ , where  $\alpha$  is the parameter in Proposition 3.1. We first show that for sufficiently large R,

(3.26) 
$$D_R^1 \in \{(x,y) : |y| \le C_0 R^\beta\}.$$

for  $C_0 = C_0(W, u)$  independent of R.

From (3.22), we have

(3.27)  
$$\begin{aligned} |\theta_R| &= |\theta_R - \theta_\infty| \\ &\leq \sum_{n=0}^{\infty} |\theta_{2^n R} - \theta_{2^{n+1} R}| \\ &\leq C(\sum_{n=0}^{\infty} 2^{n(\beta-1)}) R^{\beta-1} = C_1 R^{\beta-1} \end{aligned}$$

Furthermore, let  $C_2$  denote the constant C in (3.20). Set

$$C_0 := 100 \max\{C_1, C_2\}.$$

We show that this choice of  $C_0$  is sufficient to validate (3.26). Assume by contradiction there exists a sequence  $R_i \to \infty$ , such that  $D_{R_i}^1 = (x_i, y_i)$  satisfies  $|y_i| > C_0 R_i^{\beta}$ . Since  $|\theta_{R_i}| \le C_1 R_i^{\beta-1}$ , by an elementary geometry argument we obtain

$$\operatorname{dist}((0,0), l_i) > \frac{C_0}{2} R_i^\beta,$$

where  $l_i$  represents the straight line passing by  $D_{R_i}^1$  and  $O_{R_i}$ . Therefore,

$$r_i := \operatorname{dist}(O_{R_i}, (0, 0)) > \frac{C_0}{2} R_i^{\beta} \ge 50C_2 R_i^{\beta}.$$

We focus on the ball  $B_{r_i}$ . From (3.20) we have

(3.28) 
$$\Gamma_{\delta} \cap B_{r_i} \subset \Gamma_{\delta} \cap B_{R_i} \subset \{z : \operatorname{dist}(z, T_{R_i}) \le C_2 R_i^{\beta}\},\$$

which implies the  $\Gamma_{\delta} \cap B_{r_i}$  is contained within an  $\frac{r_i}{50}$ -neighborhood of a triod (i.e.  $T_{R_i}$ ) centered on  $\partial B_{r_i}$ . Outside this neighborhood, u(z) remains close to one of the energy wells  $a_i$ . It is clear that there exists a positive constant c > 0 such that

$$||u_{r_i} - u_{\mathcal{P}}||_{L^1(B_1)} > c, \quad \forall i.$$

As  $r_i \to \infty$ , this yields a contradiction with (2.17), thereby proving (3.26). Define

$$C := 100(C_0 + C_2).$$

We fix a sufficiently large x such that  $Cx^{\beta} < \frac{1}{2}x$ . If  $Cx^{\beta} < y < x$ , then  $z = (x, y) \in B_{2x}$ . By (3.26) and (3.27) we have

$$\operatorname{dist}(z, T_{4x}) \ge C_2(4x)^{\beta},$$

which with (3.21) shows that (3.23) holds for  $Cx^{\beta} < y < x$ .

For the case y > x, we proceed similarly, only replacing  $B_{4x}$  with  $B_{4y}$ , to obtain

$$|u(x,y) - a_1| \le K e^{-k(\operatorname{dist}((x,y),T_{4y}) - Cy^{\beta})}$$

When x, and thus y, is chosen sufficiently large, we have  $dist((x, y), T_{4y}) - Cy^{\beta} > \frac{1}{3}(y - Cx^{\beta})$ . Therefore, by updating k to  $\frac{k}{3}$ , we can verify (3.23) for the case y > x.

(3.24) can be proven in exactly the same manner as (3.23), while (3.25) follows directly from the standard elliptic regularity theory. This completes the proof.

For  $r_1 < r_2$ ,  $\theta_1 < \theta$ , we define the angular section

$$\mathcal{A}(r_1, r_2; \theta_1, \theta_2) := \{ z = (r \cos \theta, r \sin \theta) : r \in (r_1, r_2), \theta \in (\theta_1, \theta_2) \}.$$

Next we prove the following sharp energy lower and upper bounds on  $B_R$ .

**Lemma 3.3.** There are constants  $R_1$ , C, depending on W, u only, such that for  $R > R_1$ ,

$$(3.29) 3\sigma R - C \le E(u, B_R) \le 3\sigma R + C$$

*Proof.* Let  $R_1 := 10R_0$ , where  $R_0$  is the constant in Lemma 3.2. For any  $R > R_1$ , by the exponential decay results stated in Lemma (3.2), which also hold for the interfaces  $\partial \mathcal{D}_1 \cap \partial \mathcal{D}_2$  and  $\partial \mathcal{D}_2 \cap \partial \mathcal{D}_3$ , we have

(3.30) 
$$\max\left\{|u(r,\sqrt{3}r) - a_1|, |u(r,-\sqrt{3}r) - a_3|, |u(-2r,0) - a_2|\right\} \le Ke^{-k(\sqrt{3}r - Cr^{\beta})}, \ r \in [R_0, R].$$

This, together with Lemma 2.2, implies

$$E(u, B_R \setminus B_{R_0})$$

$$\geq \int_{R_0}^R \int_{\partial B_r} \left(\frac{1}{2} |\partial_T u|^2 + W(u)\right) d\mathcal{H}^1 dr$$

$$\geq 3\sigma(R - R_0) - \int_{R_0}^R 3C e^{-2k(\sqrt{3}r - Cr^\beta)} dr$$

$$\geq 3\sigma R - C(W, u),$$

which proves the lower bound in (3.29).

Now we fix two small constants  $\delta$  and  $\varepsilon$  depending on W and u. The energy limit (2.15) and Fubini's Theorem imply that for suitably large R, one can always find a  $\overline{R} \in (R, 2R)$  such that

$$3\sigma - \varepsilon \leq \int_{\partial B_{\bar{R}}} \left(\frac{1}{2} |\partial_T u|^2 + W(u)\right) d\mathcal{H}^1 \leq 3\sigma + \varepsilon.$$

Then we invoke the arguments from [21, Section 4] to obtain the following nice behavior of u on  $\partial B_{\bar{R}}$ , when  $\delta$  and  $\varepsilon$  are chosen suitably small.

(1) There are three arcs  $I_1, I_2, I_3 \subset \partial B_{\bar{R}}$  such that

$$z \in I_i \Rightarrow |u(z) - a_i| \le \delta, \quad i = 1, 2, 3.$$

(2) There are three arcs  $I_{12}, I_{23}, I_{13}$  denoting transitional arcs between  $I'_i s$ , such that

$$C_1 \leq \mathcal{H}^1(I_{ij}) \leq \frac{C_2}{\delta^2}, \ \forall i \neq j.$$
$$\bigcup_{i \neq j} I_{ij} = \partial B_{\bar{R}} \setminus (\bigcup_i I_i).$$

Here  $C_1, C_2$  are constants that only depends on W.

Now we construct an energy competitor V(z) on  $\partial B_{\bar{R}}$ . For  $z \in \partial B_{\bar{R}-1}$ , we define

$$v_1(z) = \begin{cases} a_i, & z \in \frac{\bar{R}-1}{R}I_i, \ i = 1, 2, 3\\ \text{smooth connection of } a_i \text{ to } a_j, & z \in \frac{\bar{R}-1}{\bar{R}}I_{ij}. \end{cases}$$

On the closed annulus  $\{\bar{R} - 1 \le |z| \le \bar{R}\}$ , we set

$$v(z) = \begin{cases} u(z), & |z| = \bar{R}, \\ v_1(z), & |z| = \bar{R} - 1, \\ \text{linear interpolation}, & |z| \in (\bar{R} - 1, \bar{R}). \end{cases}$$

We split the energy of v on the annulus into four parts:

$$\begin{split} E(v, B_{\bar{R}} \setminus B_{\bar{R}-1}) \\ &= \sum_{j=1}^{3} \int_{\{\theta: \bar{R}e^{i\theta} \in I_{j}\}} \int_{\bar{R}-1}^{\bar{R}} \frac{1}{2} |\partial_{r}v|^{2} r dr d\theta \\ &+ \sum_{j=1}^{3} \int_{\{\theta: \bar{R}e^{i\theta} \in I_{j}\}} \int_{\bar{R}-1}^{\bar{R}} \frac{1}{2} |\partial_{T}v|^{2} r dr d\theta \\ &+ \sum_{j=1}^{3} \int_{\{\theta: \bar{R}e^{i\theta} \in I_{j}\}} \int_{\bar{R}-1}^{\bar{R}} W(v) r dr d\theta \\ &+ \int_{\{\theta: \bar{R}e^{i\theta} \in I_{12} \cup I_{23} \cup I_{13}\}} \int_{\bar{R}-1}^{\bar{R}} \left(\frac{1}{2} |\nabla v|^{2} + W(v)\right) r dr d\theta \\ &=: \mathcal{E}_{1} + \mathcal{E}_{2} + \mathcal{E}_{3} + \mathcal{E}_{4}. \end{split}$$

For  $\mathcal{E}_4$ , since the length of  $I_{ij}$  is O(1) and  $|\nabla u|$  and W(u) are uniformly bounded, immediately one gets

$$\mathcal{E}_4 \le C(W, \delta).$$

For quantities  $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ , we estimate using Lemma 2.1:

$$\mathcal{E}_3 \leq \sum_{j=1}^3 \int_{\{\theta: \bar{R}e^{i\theta} \in I_j\}} \int_{\bar{R}-1}^{\bar{R}} C|u-a_j|^2 \, dr d\theta \leq C.$$

Combining all the estimates above implies

$$\int_{B_{\bar{R}} \setminus B_{\bar{R}-1}} \left( \frac{1}{2} |\nabla v|^2 + W(v) \right) \le C, \quad \text{ for some } C = C(W).$$

Here we can choose  $\delta$  such that Lemma 2.1 is applicable.

Then we follow the same construction of the energy competitor in [5, Appendix A] to complete the construction of v(z) inside  $B_{\bar{R}-1}$ , which satisfies

$$\int_{B_{\bar{R}-1}} \left( \frac{1}{2} |\nabla v|^2 + W(v) \right) \, dz \le 3\sigma(\bar{R}-1) + C(W).$$

We explain the rough idea: First we pick three midpoints of three transition layers,  $I_{12}$ ,  $I_{23}$ ,  $I_{13}$ , denoted by  $A_{12}$ ,  $A_{23}$ ,  $A_{13}$  respectively. For the approximated interface  $OA_{ij}$ , we define the function  $v(z) = U_{ij}(\text{dist}(z, OA_{ij}))$  within the annulus sector  $\mathcal{A}(R_0, \bar{R}-2; \theta(A_{ij}) - \frac{\pi}{6}, \theta_{ij} + \frac{\pi}{6})$ , where  $\text{dist}(\cdot, \cdot)$ represents the signed distance. Most of the energy will be concentrated in these three annulus sectors, which adds up to  $3\sigma(\bar{R} - R_0) - C$ . For the remaining regions of  $B_{\bar{R}-1}$ , we interpolate linearly, and the residual energy can be shown to be bound by a constant C. For the detailed proof, we refer interested readers to [5, Appendix A].

Therefore, we combine the estimates and the minimality of u to get

$$E(u, B_{\bar{R}}) \le E(v, B_{\bar{R}}) \le 3\sigma \bar{R} + C.$$

Recall that  $\overline{R} \in (R, 2R)$ . We utilize (3.30) again to bound the energy from below on the annulus  $B_{\overline{R}} \setminus B_R$ :

$$\int_{B_{\bar{R}}\setminus B_R} \left(\frac{1}{2}|\nabla u|^2 + W(u)\right) dz \ge 3(\bar{R} - R) - C.$$

Combining the two inequalities above we obtain the upper bound in (3.29), which completes the proof.

*Remark* 3.1. A direct consequence of Lemma 3.3 is

(3.32) 
$$\int_{\mathbb{R}^2} |\partial_r u|^2 \, dz \le C.$$

This follows from the fact that only tangential deformation is considered when estimating the lower bound in (3.31).

**Lemma 3.4.** For suitably small  $\delta$ , there exist  $R_0$ ,  $C_1$  and  $C_2$ , depending on W, u and  $\delta$ , such that for  $R > R_0$ , there exists an angle  $\theta^R \in (-C_1 R^{\beta-1}, C_1 R^{\beta-1})$  that satisfies

(3.33) 
$$\left(\Gamma_{\delta} \cap \mathcal{A}\left(R,9R;-\frac{\pi}{3},\frac{\pi}{3}\right)\right) \subset \mathcal{A}\left(R,9R;\theta^{R}-C_{2}R^{-\frac{1}{2}},\theta^{R}+C_{2}R^{-\frac{1}{2}}\right)$$

Moreover, there exist K, k depending on W and u, such that

$$(3.34) \quad |u(z) - a_1| \le K e^{-kr(\theta - \theta^R - C_2 R^{-\frac{1}{2}})}, \ z = (r\cos\theta, r\sin\theta) \in \mathcal{A}\left(2R, 8R; \theta^R + C_2 R^{-\frac{1}{2}}, \frac{\pi}{2}\right),$$

$$(3.35) \quad |u(z) - a_3| \le K e^{-kr(\theta^R - C_2 R^{-\frac{1}{2}} - \theta)}, \ z = (r\cos\theta, r\sin\theta) \in \mathcal{A}\left(2R, 8R; -\frac{\pi}{2}, \theta^R - C_2 R^{-\frac{1}{2}}\right).$$

*Proof.* For sufficiently large R, using Fubini Theorem, (2.15) and Proposition 3.1, we follow the same argument as in the proof of Lemma 3.3 to derive the existence of  $R_1 \in (\frac{1}{2}R, R), R_2 \in (9R, 10R)$ that satisfies the following properties:

- (1)  $3\sigma \varepsilon < \frac{1}{R_i} \int_{\partial B_{R_i}} \left( \frac{1}{2} |\partial_T u|^2 + W(u) \right) < 3\sigma + \varepsilon$ , for  $i = 1, 2, \varepsilon \ll 1$ . (2) There are  $\theta_1, \theta_2 \in (-CR^{\beta-1}, CR^{\beta-1})$  such that

(3.36) 
$$|u(z) - a_1| \le \frac{1}{2}\delta$$
, for  $z = (R_i \cos \theta, R_i \sin \theta), \ \theta \in (\theta_i + \frac{C}{R_i}, \frac{\pi}{3}), \ i = 1, 2,$ 

(3.37) 
$$|u(z) - a_3| \le \frac{1}{2}\delta$$
, for  $z = (R_i \cos \theta, R_i \sin \theta)$ ,  $\theta \in (-\frac{\pi}{3}, \theta_i - \frac{C}{R_i})$ ,  $i = 1, 2$ .

The inequalities above basically indicate that on  $\partial B_{R_i} \cap \{\theta \in (-\frac{\pi}{3}, \frac{\pi}{3})\}, u(z)$  will be uniformly close to  $a_1$  or  $a_3$  outside a transition layer of size O(1).

Next we claim that there exists a constant  $C_3 = C_3(W, u)$  such that

$$|\theta_1 - \theta_2| \le C_3 R^{-\frac{1}{2}}.$$

**Proof of the Claim:** We argue by contradiction. Assume that  $\theta_1 - \theta_2 > C_3 R^{-\frac{1}{2}}$  for some  $C_3 = C_3(W, u)$  to be determined. We can take R large enough to guarantee that

$$(\theta_1 - \frac{C}{R_1}) - (\theta_2 + \frac{C}{R_2}) > \frac{C_3}{2}R^{-\frac{1}{2}},$$

where the constant C appears in (3.36) and (3.37). Then for  $\theta \in (\theta_2 + \frac{C}{R_2}, \theta_1 - \frac{C}{R_1})$ , we have

(3.38) 
$$|u(R_1, \theta) - a_3| \le \delta, \quad |u(R_2, \theta) - a_1| \le \delta.$$

Now we calculate the energy on the annulus  $\mathcal{A}(R_1, R_2; 0, 2\pi)$ . Let  $\varphi \in (0, \frac{\pi}{2})$  be an angle whose value will be specified later, we compute

$$(3.39) E(u, \mathcal{A}(R_1, R_2; 0, 2\pi)) \\ \geq \int_{R_1}^{R_2} \int_{\partial B_r} \left(\frac{1}{2} |\partial_T u|^2 + \sin^2 \varphi W(u)\right) d\mathcal{H}^1 dr + \int_{\theta_2 + \frac{C}{R_2}}^{\theta_1 - \frac{C}{R_1}} \int_{R_1}^{R_2} \left(\frac{1}{2} |\partial_r u|^2 + \cos^2 \varphi W(u)\right) r dr d\theta \\ \geq \int_{R_1}^{R_2} \sin \varphi (3\sigma - Ce^{-kr}) dr + \frac{C_3}{2} R^{-\frac{1}{2}} \cdot (\cos \varphi (\sigma - C\delta^2)) \cdot (\frac{1}{2}R) \\ \geq \left[3\sin \varphi (R_2 - R_1) + \cos \varphi \frac{C_3 R^{\frac{1}{2}}}{8}\right] \sigma - C_4(W).$$

Since the estimate above applies to any  $\varphi$ , we take  $\varphi = \arctan \frac{3(R_2 - R_1)}{\frac{C_3}{8}R^{\frac{1}{2}}}$  to obtain

$$(3.40) \qquad E(u, \mathcal{A}(R_1, R_2; 0, 2\pi)) \\ \ge \left(9(R_2 - R_1)^2 + \frac{C_3^2 R}{64}\right)^{\frac{1}{2}} \sigma - C_4 \\ \ge \left(3(R_2 - R_1) + \frac{1}{2} \frac{C_3^2 R}{64 \cdot 3(R_2 - R_1)}\right) \sigma - C_4 \\ \ge 3(R_2 - R_1) \sigma + \frac{C_3^2}{3840} \sigma - C_4.$$

 $\mathbf{n}$ 

Applying (3.29) to  $R_1$  and  $R_2$  implies that

(3.41) 
$$E(u, \mathcal{A}(R_1, R_2; 0, 2\pi)) \le 3(R_2 - R_1)\sigma + C_5$$

Now we can take  $C_3(W, u)$  large enough such that

$$\frac{C_3^2\sigma}{3840} - C_4 > 2C_5$$

which yields a contradiction. This completes the proof of the claim.

We set

$$\theta^R := \frac{\theta_1 + \theta_2}{2}.$$

We further show that there exists a constant  $C_2(W, u)$  such that one can find  $\tilde{\theta}_1 \in (\theta^R, \theta^R + C_2 R^{-\frac{1}{2}})$ and  $\tilde{\theta}_3 \in (\theta^R - C_2 R^{-\frac{1}{2}}, \theta^R)$  that satisfy

(3.42) 
$$\begin{aligned} |u(r,\tilde{\theta}_1) - a_1| &\leq \delta, \quad \forall r \in (R_1, R_2), \\ |u(r,\tilde{\theta}_3) - a_3| &\leq \delta, \quad \forall r \in (R_1, R_2). \end{aligned}$$

The proof is based on the same idea as the proof of the claim. If for all  $\theta \in (\theta^R, \theta^R + C_2 R^{-\frac{1}{2}})$ , there exists at least one point  $r(\theta) \in (R_1, R_2)$  such that  $|u(r(\theta), \theta) - a_1| > \delta$ , then for such  $\theta$ , there will be some non trivial energy generated along the radial segment  $\{re^{i\theta}: r \in [R_1, R_2]\}$ . This is due to the boundary constraint (3.36). Using similar estimates as (3.39) and (3.40), which split the energy into tangential and radial parts, we can obtain

$$\int_{\mathcal{A}(R_1,R_2;0,2\pi)} \left(\frac{1}{2} |\nabla u|^2 + W(u)\right) dz \ge 3\sigma(R_2 - R_1) + K_1 C_2^2 \delta^2 - K_2,$$

where  $K_1, K_2$  depend on W, u. Taking  $C_2$  sufficiently large yields a contradiction with (3.41). The existence of  $\tilde{\theta}_3$  follows by the same argument.

Using (3.23), (3.36) and (3.42), we find

$$|u(z) - a_1| \le \delta, \quad \forall z \in \partial \mathcal{A}(R_1, R_2; \tilde{\theta}_1, \frac{\pi}{3}).$$

Invoking the variational maximum principle in [2], we obtain

$$|u(z) - a_1| \le \delta, \quad \forall z \in \mathcal{A}(R_1, R_2; \tilde{\theta}_1, \frac{\pi}{3})$$

Similarly,

$$|u(z) - a_3| \le \delta, \quad \forall z \in \mathcal{A}(R_1, R_2; -\frac{\pi}{3}, \tilde{\theta}_3).$$

Since  $\tilde{\theta}_1, \tilde{\theta}_3$  are in the  $C_2 R^{-\frac{1}{2}}$  neighborhood of  $\theta^R$ , (3.33) follows immediately. The exponential decay in (3.34) and (3.35) follows from standard estimates using the comparison principle in elliptic theory, see [4, Lemma 4.5 & Proposition 5.2] for detailed arguments. The proof of Lemma 3.4 is complete.

Given R large, applying (3.33) to radius R and 2R implies

$$\left( \Gamma_{\delta} \cap \mathcal{A}(2R, 9R; -\frac{\pi}{3}, \frac{\pi}{3}) \right)$$
  
 
$$\subset \left[ \mathcal{A} \left( 2R, 9R; \theta^{R} - C_{2}R^{-\frac{1}{2}}, \theta^{R} + C_{2}R^{-\frac{1}{2}} \right) \cap \mathcal{A} \left( 2R, 9R; \theta^{2R} - C_{2}(2R)^{-\frac{1}{2}}, \theta^{2R} + C_{2}(2R)^{-\frac{1}{2}} \right) \right].$$

Consequently, we get the similar estimate of the closeness for  $\theta_{13}^R$  at comparable scalings as in (3.22), but with the improved power,

$$|\theta^R - \theta^{2R}| \le CR^{-\frac{1}{2}}.$$

We argue as in (3.27) to get

$$|\theta^R| \le CR^{-\frac{1}{2}}.$$

Furthermore, following the proof of Lemma 3.2, we can refine the power  $\beta$  in (3.23)–(3.25) to  $\frac{1}{2}$ . Specifically, we have

(3.43) 
$$|u(x,y) - a_1| \le K e^{-k(y - Cx^{\frac{1}{2}})}, \quad y \ge Cx^{\frac{1}{2}},$$

(3.44) 
$$|u(x,y) - a_3| \le K e^{-k(|y| - Cx^{\frac{1}{2}})}, \quad y \le -Cx^{\frac{1}{2}}.$$

(3.45) 
$$|\nabla u(x,y)| \le K e^{-k(y-Cx^{\frac{1}{2}})}, \quad |y| \ge Cx^{\frac{1}{2}},$$

when  $x > R_0(W, u)$ . In short, Lemma 3.4 together with the analysis above indicate that the parameters  $(\alpha, \beta)$  in Proposition 3.1 and Lemma 3.2 can be strengthened to  $(0, \frac{1}{2})$ .

Considering a ray from the origin  $\{re^{i\theta} : r \in (0,\infty), \theta = \theta_0\}$ , we obtain the following direct consequence of the exponential decay estimates (3.43) and (3.44), which apply to all three sharp interfaces.

**Lemma 3.5.** If  $\theta \notin \{0, \frac{2\pi}{3}, \frac{4\pi}{3}\}$ , then there exist K, k and  $R_0$  such that

$$|u(r,\theta) - a_i| \le K e^{-k(r-R_0)}, \ \forall r \ge R_0,$$

where  $a_i$  is the phase associated with  $\theta$ , that is,  $\{re^{i\theta} : r \in (0,\infty), \theta \text{ fixed}\} \in \mathcal{D}_i$ .

Moreover, if  $\theta$  belongs to a compact set  $\mathcal{K} \subset ([0, 2\pi) \setminus \{0, \frac{2\pi}{3}, \frac{4\pi}{3}\})$ , then the constants  $K, k, R_0$  can be chosen uniformly with respect to  $\mathcal{K}$ .

The proof is omitted here, as it follows immediately from (3.43)–(3.45).

3.2. Small horizontal deformation. Let  $\mathbb{R}^2_{x+}$  denote the half plane  $\{(x, y) : x > 0\}$ . We will derive the key estimate that  $\int_{\mathbb{R}^2_{x+}} |\partial_x u|^2 dz$  is finite. For R sufficiently large, define an equilateral triangle  $S_R$  by

(3.46)  
$$\partial S_R = \{(x, y) : x = R, y \in [-\sqrt{3R}, \sqrt{3R}]\} \bigcup$$
$$\{(x, y) : y = \frac{\sqrt{3}}{3}(x + 2R), x \in [-2R, R]\} \bigcup$$
$$\{(x, y) : y = \frac{-\sqrt{3}}{3}(x + 2R), x \in [-2R, R]\}.$$

 $S_R$  is centered at the origin with a side length of  $2\sqrt{3}R$ . We further define

$$S_R^{13} := \{ z = (r\cos\theta, r\sin\theta) \in S_R : \theta \in (-\frac{\pi}{3}, \frac{\pi}{3}) \},$$
  

$$S_R^{12} := \{ z = (r\cos\theta, r\sin\theta) \in S_R : \theta \in (\frac{\pi}{3}, \pi) \},$$
  

$$S_R^{23} := \{ z = (r\cos\theta, r\sin\theta) \in S_R : \theta \in (\pi, \frac{5\pi}{3}) \},$$

which form a 3-partition of  $S_R$  such that  $S_R^{ij}$  is symmetric with respect to the  $a_i$ - $a_j$  interface. We now carry out some energy estimates on  $S_R$ .



FIGURE 1. Definition of  $S_R$  and  $S_R^{ij}$ . Red :  $a_1$ , green:  $a_2$ , blue:  $a_3$ .

Lemma 3.6. The following estimate holds:

(3.47) 
$$\lim_{R \to \infty} \frac{1}{R} E(u, S_R) = 3\sigma$$

*Proof.* Firstly, since  $B_R \subset S_R$ ,

(3.48) 
$$\lim_{R \to \infty} \frac{1}{R} E(u, S_R) \ge \lim_{R \to \infty} \frac{1}{R} E(u, B_R) = 3\sigma$$

For the inequality in the other direction, we consider a slightly larger ball  $B_{R_1}$  where

$$R_1(R) := \sqrt{R^2 + C^2 R} \sim R + \frac{C^2}{2}.$$

Here C is the constant in (3.43)–(3.45). By (2.15), we have

$$\lim_{R \to \infty} \frac{1}{R} E(u, B_{R_1}) = \lim_{R \to \infty} \frac{R_1}{R} \cdot 3\sigma = 3\sigma.$$

From definition,

$$S_R \setminus B_{R_1} = \bigcup_{i < j} (S_R^{ij} \setminus B_{R_1}),$$
  
$$S_R^{13} \setminus B_{R_1} = \{(x, y) : x \in [\frac{R_1}{2}, R), |y| \in [\sqrt{R_1^2 - x^2}, \sqrt{3}x)\}.$$

In particular, a point  $(x, y) \in S_R^{13} \setminus B_{R_1}$  satisfies

$$|y| \ge CR^{\frac{1}{2}} \ge Cx^{\frac{1}{2}}.$$

Utilizing estimates (3.43)-(3.45), we compute

$$\begin{aligned} &\frac{1}{R}E(u, S_R^{13} \setminus B_{R_1}) \\ \leq &\frac{2}{R} \left[ \int_{R_1/2}^{R} \left( \int_{\sqrt{R_1^2 - x^2}}^{\sqrt{3}x} Ce^{-2k(y - Cx^{\frac{1}{2}})} \, dy \right) dx \right] \\ \leq &\frac{C}{R} \int_{R_1/2}^{R} e^{-2k(\sqrt{R_1^2 - x^2} - Cx^{\frac{1}{2}})} \, dx \\ \leq &\frac{C}{R}. \end{aligned}$$

The same estimates also hold for  $S^{23}_R$  and  $S^{12}_R$ . Therefore,

(3.49)  
$$\lim_{R \to \infty} \frac{1}{R} E(u, S_R)$$
$$\leq \lim_{R \to \infty} \left( \int_{S_R \setminus B_{R_1}} + \int_{B_{R_1}} \right) \left( \frac{1}{2} |\nabla u|^2 + W(u) \right) dz$$
$$\leq \lim_{R \to \infty} \left( \frac{3C}{R} + 3\sigma \right) = 3\sigma.$$

(3.48) and (3.49) together imply (3.47), which completes the proof.

**Lemma 3.7.** There exists a constant C = C(W, u) such that for R sufficiently large, there is a  $\tilde{R} \in (R, 2R)$  satisfying

(3.50) 
$$3\sigma\tilde{R} - C \le \int_{S_{\tilde{R}}} \left(\frac{1}{2}|\nabla u|^2 + W(u)\right) dz \le 3\sigma\tilde{R} + C.$$

*Proof.* The lower bound follows immediately by the same arguments as in (3.30) and (3.31). We are only left to prove the upper bound.

Fix a small constant  $\varepsilon \ll 1$ , whose value will depend on the constant  $\delta$  introduced later. By Lemma (3.6) and Fubini's Theorem, for sufficiently large R, there exists a  $\tilde{R} \in (R, 2R)$  satisfying

(3.51) 
$$\int_{\partial S_{\tilde{R}}} \left( \frac{1}{2} |\partial_T u|^2 + W(u) \right) d\mathcal{H}^1 \le 3\sigma + \varepsilon.$$

We first observe that substituting  $r = \tilde{R}$  in (3.30) and Lemma 2.2 together yield

$$\int_{\{\partial S_{\tilde{R}} \setminus \partial S_{\tilde{R}}^{13}\}} \left(\frac{1}{2} |\partial_T u|^2 + W(u)\right) d\mathcal{H}^1 \ge 2\sigma - Ce^{-k\tilde{R}} \ge 2\sigma - \varepsilon,$$

when  $\tilde{R}$  is large enough. Combining this with (3.51), we obtain

(3.52) 
$$\int_{-\sqrt{3}\tilde{R}}^{\sqrt{3}\tilde{R}} \left(\frac{1}{2} |\partial_y u(\tilde{R}, y)|^2 + W(u(\tilde{R}, y))\right) dy \le \sigma + 2\varepsilon.$$

This further implies that there is a  $y_{\tilde{R}}$  satisfying  $|y_{\tilde{R}}| \leq C \tilde{R}^{\frac{1}{2}}$  and

(3.53) 
$$\begin{aligned} |u(\bar{R}, y) - a_1| &\leq \delta, \quad y \in (y_{\tilde{R}} + C(W, \delta), \sqrt{3R}), \\ |u(\tilde{R}, y) - a_3| &\leq \delta, \quad y \in (-\sqrt{3}\tilde{R}, y_{\tilde{R}} - C(W, \delta)), \end{aligned}$$

where  $\delta \ll 1$  is a fixed small constant depending only on W. Once  $\delta$  is fixed, the value of  $\varepsilon$  can be adjusted to ensure the existence of  $y_{\tilde{R}}$ .

Next we construct an energy competitor v on  $S_{\tilde{R}}$  such that v = u on  $\partial S_{\tilde{R}}$  and v satisfies the energy upper bound in (3.50). Then the upper bound for u follows from minimality.

Fix a large enough  $r_0$  such that  $x > 2Cx^{\frac{1}{2}}$  when  $x > r_0$ , where C is the constant in estimates (3.43) and (3.44). Additionally, assume  $\tilde{R}$  is much larger than  $r_0$ . For the region  $S_{\tilde{R}}^{13} \setminus S_{r_0}^{13}$ , we first define v(x, y) on the boundary:

$$(3.54) \quad v(x,y) = \begin{cases} a_1, & \text{on } \{x \in [r_0, \tilde{R} - 1], y = \sqrt{3}x\}, \\ a_3, & \text{on } \{x \in [r_0, \tilde{R} - 1], y = -\sqrt{3}x\}, \\ (\tilde{R} - x)a_1 + (x - (\tilde{R} - 1))u(\tilde{R}, \sqrt{3}\tilde{R}), & \text{on } \{x \in (\tilde{R} - 1, \tilde{R}), y = \sqrt{3}x\}, \\ (\tilde{R} - x)a_3 + (x - (\tilde{R} - 1))u(\tilde{R}, -\sqrt{3}\tilde{R}), & \text{on } \{x \in (\tilde{R} - 1, \tilde{R}), y = -\sqrt{3}x\}, \\ u(x, y), & \text{on } \{x \in \tilde{R}, y \in [-\sqrt{3}\tilde{R}, \sqrt{3}\tilde{R}]\}. \end{cases}$$

Let  $l_{13}$  denote the line segment connecting  $(r_0, 0)$  and  $(\tilde{R} - 1, y_{\tilde{R}})$ . The slope of  $l_{13}$  is  $O(\tilde{R}^{-\frac{1}{2}})$ . We define

$$v(x,y) = \begin{cases} U_{13}(\operatorname{dist}((x,y),l_{13})), & \text{on } S_1 := \{x \in [r_0, \tilde{R}-1], y \in [-x,x]\}, \\ \text{vertical linear interpolation between} \\ \{y = x\} \text{ and } \{y = \sqrt{3}x\}, & \text{on } S_2 := \{x \in [r_0, \tilde{R}-1], y \in (x,\sqrt{3}x)\}, \\ \text{vertical linear interpolation between} \\ \{y = -\sqrt{3}x\} \text{ and } \{y = -x\}, & \text{on } S_3 := \{x \in [r_0, \tilde{R}-1], y \in (-\sqrt{3}x, -x)\}, \\ \text{horizontal linear interpolation} \\ \text{between boundary data}, & \text{on } S_4 := \{x \in (\tilde{R}-1, \tilde{R}), y \in (-\sqrt{3}x, \sqrt{3}x)\} \end{cases}$$

We compute the energy in  $S_1$ – $S_4$ .

$$\begin{split} E(v,S_1) &\leq \sigma \sqrt{(\tilde{R} - 1 - r_0)^2 + C^2 \tilde{R}} \leq \sigma (\tilde{R} - r_0 + C). \\ E(v,S_2 \cup S_3) &\leq 2 \int_{r_0}^{\tilde{R} - 1} \int_x^{\sqrt{3}x} C e^{-ky} \, dy dx \leq C(W,u). \\ E(v,S_4) &\leq \int_{S_4} \frac{1}{2} |\partial_x u|^2 \, dz + \int_{S_4} \frac{1}{2} |\partial_y u|^2 \, dz + \int_{S_4} W(u) \, dz \end{split}$$

For fixed y, by definition we have

$$\int_{\tilde{R}-1}^{\tilde{R}} |\partial_x v(x,y)|^2 \, dx = |v(\tilde{R},y) - v(\tilde{R}-1,y)|^2.$$

Therefore,

$$\begin{split} &\int_{S_4} \frac{1}{2} |\partial_x v|^2 \, dz \\ \leq & \frac{1}{2} \int_{-\sqrt{3}(\tilde{R}-1)}^{\sqrt{3}(\tilde{R}-1)} |v(\tilde{R},y) - v(\tilde{R}-1,y)|^2 \, dy + C \\ \leq & C + \int_{y_{\tilde{R}}+C}^{\sqrt{3}(\tilde{R}-1)} \left( |v(\tilde{R}-1,y) - a_1|^2 + |u(\tilde{R},y) - a_1|^2 \right) \, dy \\ &\quad + \int_{-\sqrt{3}(\tilde{R}-1)}^{y_{\tilde{R}}-C} \left( |v(\tilde{R}-1,y) - a_3|^2 + |u(\tilde{R},y) - a_3|^2 \right) \, dy \\ \leq & C + C \int_{-\sqrt{3}(\tilde{R}-1)}^{\sqrt{3}(\tilde{R}-1)} \left( W(v(\tilde{R}-1,y)) + W(u(\tilde{R},y)) \right) \, dy \leq C \end{split}$$

where we have used (3.53) to get the second inequality. Similarly,

$$\begin{split} &\int_{S_4} W(v) \, dz \\ \leq C + C \int_{\tilde{R}-1}^{\tilde{R}} \left( \int_{y_{\tilde{R}}+C}^{\sqrt{3}(\tilde{R}-1)} |v(x,y) - a_1|^2 dy + \int_{-\sqrt{3}(\tilde{R}-1)}^{y_{\tilde{R}}-C} |v(x,y) - a_3|^2 \, dy \right) \, dx \\ \leq C + C \int_{-\sqrt{3}(\tilde{R}-1)}^{\sqrt{3}(\tilde{R}-1)} \left( W(v(\tilde{R}-1,y)) + W(v(\tilde{R},y)) \right) \, dy \leq C \end{split}$$

Moreover, we have

$$\int_{S_4} \frac{1}{2} |\partial_y v|^2 dz$$
  
$$\leq C + C \int_{-\sqrt{3}(\tilde{R}-1)}^{\sqrt{3}(\tilde{R}-1)} \left( |\partial_y v(\tilde{R}-1,y)|^2 + |\partial_y v(\tilde{R},y)|^2 \right) dy \leq C$$

Adding up all estimates above implies

$$(3.56) E(v, S^{13}_{\tilde{R}} \setminus S^{13}_{r_0}) \le \sigma \tilde{R} + C$$

The same estimate also holds for  $S^{12}_{\tilde{R}} \setminus S^{12}_{r_0}$  and  $S^{23}_{\tilde{R}} \setminus S^{23}_{r_0}$ .

Within the triangle  $S_{r_0}$ , the previous construction ensures that  $v|_{\partial S_{r_0}}$  is Lipschitz. Hence one can extend v to the interior of  $S_{r_0}$  such that

$$(3.57) E(v, S_{r_0}) \le C.$$

Combining (3.56) and (3.57) we get

$$E(v, S_{\tilde{R}}) \le 3\sigma \tilde{R} + C.$$

This completes the proof of Lemma 3.7.

Our next lemma asserts that the horizontal deformation of u on  $\mathbb{R}^2_{x+}$  is bounded and small.

**Lemma 3.8.** There is a constant C = C(W, u) such that

(3.58) 
$$\int_{\mathbb{R}^2_{x+}} |\partial_x u|^2 \, dx dy \le C.$$

*Proof.* In the following, we will use the universal constants K, k, and  $R_0$ , which depend only on W and u. When the constants differ across various estimates, they can be adjusted as needed without compromising the validity of the estimates.

For each x, define the vertical line

$$l_x := \{(x, y) : y \in (-\infty, \infty)\}.$$

By the localization of the diffuse interface, outside the finite disk  $B_{R_0}$ , the diffuse interface  $\Gamma_{\delta}$  is located in a small neighborhood of the limit interface  $\partial \mathcal{P}$ , where

$$\partial \mathcal{P} = \{(x,0) : x > 0\} \cup \{(x, -\sqrt{3}x) : x < 0\} \cup \{(x, \sqrt{3}x) : x < 0\}.$$

The size of the neighborhood is measured by  $O(R^{\frac{1}{2}})$ , i.e.

dist
$$(\Gamma_{\delta} \cap \partial B_R, \ \partial \mathcal{P} \cap \partial B_R) \leq C(W, \delta, u) R^{\frac{1}{2}}, \ \forall R \geq R_0.$$

Thanks to the exponential decay of the distance of u(x, y) to  $a_i$  away from the diffuse interface, it follows that for any x > 0, u(x, y) converges to  $a_1(a_3)$  exponentially as y goes to  $+\infty(-\infty)$ . And  $|\nabla u|$  converges exponentially to 0 as |y| goes to  $+\infty$ . Therefore,

$$\int_{-\infty}^{\infty} |\partial_x u(x,y)|^2 \, dy < \infty, \quad \forall x > 0.$$

Consequently, to prove (3.58), it suffices to show

(3.59) 
$$\int_{\{x>R_0\}} |\partial_x u|^2 \, dx \, dy < \infty,$$

for a constant  $R_0$  such that (3.43)–(3.45) hold for  $x > R_0$  and Lemma (3.7) applies for  $R > R_0$ . Applying (3.43) and (3.44) to  $x = R_0$  implies that

$$|u(R_0, y) - a_1| \le K e^{-k(y - CR_0^{\frac{1}{2}})}, \quad y \ge CR_0^{\frac{1}{2}},$$
$$|u(R_0, y) - a_3| \le K e^{-k(|y| - CR_0^{\frac{1}{2}})}, \quad y \le -CR_0^{\frac{1}{2}}$$

Furthermore, since  $\{(x, 0) : x < 0\}$  is the bisector of  $\mathcal{D}_2$ , by Lemma 3.5 we have

$$|u(x,0) - a_2| \le K e^{-k(|x| - R_0)}, \quad x < -R_0.$$

For any  $r < -2R_0$ , we consider the line segment  $\{(x, y) : x \in [r, R_0], y = \frac{\sqrt{3}}{3}(x - r)\}$ , which will be exponentially close to  $a_2$  and  $a_1$  at two endpoints, due to the estimates above. Therefore,

19

Lemma 2.2 implies that the 1D energy on this line segment is bounded from below by  $\sigma - Ce^{-k|r|}$ . We have for any  $R > R_0$ ,

(3.60) 
$$\int_{-2R}^{R_0} \int_0^{\frac{\sqrt{3}}{3}(x+2R)} \left(\frac{1}{2} |\nabla u|^2 + W(u)\right) dy dx$$
$$\geq \frac{1}{2} \int_{-2R}^{-2R_0} (\sigma - Ce^{-k|r|}) dr \geq \sigma R - C.$$

Similarly,

(3.61) 
$$\int_{-2R}^{R_0} \int_{-\frac{\sqrt{3}}{3}(x+2R)}^0 \left(\frac{1}{2}|\nabla u|^2 + W(u)\right) \, dy dx \ge \sigma R - C.$$

For any  $R > R_0$ , by Lemma 3.7 there exists a  $\tilde{R} \in (R, 2R)$  such that

$$(3.62) E(u, S_{\tilde{R}}) \le 3\sigma \tilde{R} + C$$

This, together with (3.60) and (3.61) applying to  $\tilde{R}$ , yields that

$$(3.63) E(u, S_{\tilde{R}} \cap \{x > R_0\}) \le \sigma \tilde{R} + C.$$

Note that for any  $x \in (R_0, \tilde{R})$ , the vertical line  $l_x$  intersects  $\partial S_{\tilde{R}}$  at two points,  $(x, \frac{\sqrt{3}}{3}(x+2\tilde{R}))$  and  $(x, -\frac{\sqrt{3}}{3}(x+2\tilde{R}))$ . (3.43) and (3.44) imply that

$$|u(x, \frac{\sqrt{3}}{3}(x+2\tilde{R})) - a_1| \le Ke^{-k\tilde{R}}, \quad |u(x, \frac{-\sqrt{3}}{3}(x+2\tilde{R})) - a_3| \le Ke^{-k\tilde{R}},$$

which leads to

$$\int_{-\frac{\sqrt{3}}{3}(x+2\tilde{R})}^{\frac{\sqrt{3}}{3}(x+2\tilde{R})} \left(\frac{1}{2} |\partial_y u(x,y)|^2 + W(u(x,y))\right) dy \ge \sigma - Ce^{-2k\tilde{R}}$$

thanks to Lemma 2.2. From (3.63) we have

(3.64)  

$$\int_{S_{\tilde{R}} \cap \{x > R_0\}} \frac{1}{2} |\partial_x u|^2 dz$$

$$\leq (\sigma \tilde{R} + C) - \int_{S_{\tilde{R}} \cap \{x > R_0\}} \left(\frac{1}{2} |\partial_y u|^2 + W(u)\right) dz$$

$$\leq (\sigma \tilde{R} + C) - (\tilde{R} - R_0)(\sigma - Ce^{-2k\tilde{R}}) \leq C,$$

where the last constant C depends only on W and u. The final step is to use (3.45) to get

$$\lim_{\tilde{R}\to\infty}\int_{R_0}^R\int_{|y|>\frac{\sqrt{3}}{3}(x+2\tilde{R})}|\partial_x u|^2\,dydx=0.$$

Combining this and (3.64) gives

$$\int_{R_0 < |x| < \tilde{R}} |\partial_x u|^2 \, dz \le C,$$

where C is independent of  $\tilde{R}$ . Now (3.59) follows immediately by letting  $\tilde{R} \to \infty$ . The proof is complete.

**Corollary 3.9.** There exists a constant C = C(W, u) such that

(3.65) 
$$\int_{\{0 < x < R\}} \left(\frac{1}{2} |\nabla u|^2 + W(u)\right) dz \le \sigma R + C, \quad \forall R > 0.$$

*Proof.* Let  $R_0 = R_0(W, u)$  be the same constant as in (3.59). By Lemma 3.5 we know that u(x, y) will exponentially converge to  $a_1(a_3)$  as y goes to  $\infty(-\infty)$  uniformly for  $x \in [0, R_0]$ . Therefore,

(3.66) 
$$\int_{\{0 < x < R_0\}} \left(\frac{1}{2} |\nabla u|^2 + W(u)\right) dz \le C.$$

Now (3.65) follows directly from (3.66), (3.63) and the pointwise estimates (3.43)–(3.45).  $\Box$ 

3.3. Approximation by the heteroclinic connection. Lemma 3.4 implies that the diffuse interface  $\Gamma_{\delta}$  is confined within an  $O(R^{\frac{1}{2}})$  neighborhood of the limiting interface. We now argue that  $O(R^{\frac{1}{2}})$  can be improved to O(1) and conclude the proof of Theorem 1.1.

Define the following notations:

- Denote by  $||v||_s$  the  $H^s(\mathbb{R})$  norm of a function  $v : \mathbb{R} \to \mathbb{R}^2$  for  $s \ge 0$ , In particular,  $||v||_0$  denotes the  $L^2$  norm.
- $\mathcal{U} := \{ U(\cdot m) : m \in \mathbb{R} \}$  is the set of translations of the connection  $U = U_{31}$ .

• 
$$S := \{ v \in H^1_{loc}(\mathbb{R}, \mathbb{R}^2) : \int_{\mathbb{R}} \left( \frac{1}{2} |v'|^2 + W(v) \right) \, dx < \infty, \lim_{x \to -\infty} v(x) = a_3, \lim_{x \to \infty} v(x) = a_1 \}$$

- $J(v) := \int_{\mathbb{R}} \left( \frac{1}{2} |v'|^2 + W(v) \right) dx, \forall v \in \mathcal{S}.$
- $d_s(v,\mathcal{U}) := \inf\{\|v-w\|_s, w \in \mathcal{U}\}.$

We rely on the following results of [32] saying that if the one dimensional energy of v(x) is close to  $\sigma$ , then v(x) can be well approximated by a translation of U.

**Proposition 3.10.** Suppose (H1), (H2) hold. There exist positive constants  $\varepsilon$  and  $\alpha$  such that the following hold:

(1) ([32, Lemma 2.1]) For s = 0, 1, if  $d_s(v, \mathcal{U}) \leq \varepsilon$ , then there is a unique  $h_s(v)$  such that

$$d_s(v,\mathcal{U}) = \|v - U(\cdot - h_s(v))\|_s$$

Moreover,  $h_s$  is a function of class  $C^{3-s}$  of v for s = 0, 1.

(2) ([32, Lemma 4.5]) If  $d_1(v, \mathcal{U}) \leq \varepsilon$ , then

(3.67) 
$$J(v) - \sigma \ge \alpha \|v - U(\cdot - h_0(v))\|_1^2 \ge \alpha d_1(v, \mathcal{U})^2.$$

(3) ([32, Corollary 4.6]) If  $d_1(v, \mathcal{U}) > \varepsilon$ , then  $J(v) > \alpha \varepsilon^2$ .

In the rest of this paper we fix the constants  $\varepsilon$  and  $\alpha$ , as defined in Proposition 3.10. Moreover, we may assume  $\varepsilon \ll 1$ , since  $\alpha$  in (3.67) is independent of  $d_1(v, \mathcal{U})$ . Define the set

(3.68) 
$$\mathcal{B} := \{x > 0 : d_1(u(x, \cdot)) \ge \varepsilon\}$$

From Corollary 3.9 and (3.67), it follows that

(3.69) 
$$|\mathcal{B}| \le \frac{C}{\alpha \varepsilon^2} = C(W, u).$$

Proposition 3.10 implies that for any  $x \in \mathbb{R}^+ \setminus \mathcal{B}$ , i.e., outside a set of finite measure, the slicing function  $u(x, \cdot)$  can be approximated closely in  $\|\cdot\|_s$  norm by a translated connection  $U(\cdot - h_s(x))$ , where  $h_s(x)$  is a  $C^2$  function of x on  $\mathbb{R}^+ \setminus \mathcal{B}$ , for s = 0 or 1.

We further recall the following identities for the 2D Allen-Cahn system, which will play a crucial role in our analysis.

**Proposition 3.11** (Lemma 8.2 in [32]). The following identities hold for u:

(3.70) 
$$\int_{-\infty}^{\infty} \left[ \frac{1}{2} (|\partial_y u|^2 - |\partial_x u|^2) + W(u(x,y)) \right] dy \equiv \sigma, \quad \forall x \in \mathbb{R}^+$$

(3.71) 
$$\int_{-\infty}^{\infty} (\partial_x u \partial_y u) \, dy \equiv 0, \quad \forall x \in \mathbb{R}^+$$

*Proof.* The above identities were first derived in [32], and later extended to a more general setting in [24]. See also [4, Section 3.4] for a more detailed discussion. Here we present a proof for completeness.

Define

$$G(x) := \int_{-\infty}^{\infty} \left[ \frac{1}{2} (|\partial_y u|^2 - |\partial_x u|^2) + W(u(x,y)) \right] \, dy, \quad x > 0$$

Thanks to the exponential decay estimates (3.43)–(3.45) as  $|y| \to \infty$ , we have

$$\frac{dG(x)}{dx} = \int_{-\infty}^{\infty} \left( \partial_y u \partial_{xy}^2 u - \partial_x u \partial_{xx}^2 u + W_u(u) \partial_x u \right) \, dy$$
$$= \int_{-\infty}^{\infty} \left( \partial_y u \partial_{xy}^2 u + \partial_x u \partial_{yy}^2 u \right) \, dy$$
$$= \int_{-\infty}^{\infty} \partial_y (\partial_x u \partial_y u) \, dy = 0.$$

Here we have utilized the equation  $\Delta u = W_u(u)$  and (3.45). Thus  $G(x) \equiv C_1$  for all x > 0. From (3.65), we deduce

$$\sigma R + C \ge C_1 R.$$

Letting  $R \to \infty$  yields  $C_1 \leq \sigma$ . On the other hand, since  $u(x, \cdot)$  connects  $a_3$  to  $a_1$  for every x, and  $\sigma$  is the minimal energy for such a connection, we also have

$$C_1 R \ge \sigma R - \int_{\{x>0\}} |\partial_x u|^2 \, dz \ge \sigma R - C,$$

Sending  $R \to \infty$  implies  $C_1 \ge \sigma$ . Therefore,  $C_1 = \sigma$  and identity (3.70) is proved.

For identity (3.71), we define

$$H(x) := \int_{-\infty}^{\infty} (\partial_x u \partial_y u) \, dy, \quad x > 0.$$

We compute

$$\begin{aligned} \frac{dH(x)}{dx} &= \int_{-\infty}^{\infty} (\partial_{xx}^2 u \partial_y u + \partial_x u \partial_{xy}^2 u) \, dy \\ &= \int_{-\infty}^{\infty} (-\partial_{yy}^2 u \partial_y u - W_u(u) \partial_y u + \partial_x(u) \partial_{xy}^2 u) \, dy \\ &= \int_{-\infty}^{\infty} -\partial_y (\frac{1}{2} |\partial_y u|^2 - \frac{1}{2} |\partial_x u|^2 + W(u)) \, dy = 0, \end{aligned}$$

where we use  $|\nabla u(x,y)| \to 0$  and  $W(u(x,y)) \to 0$  as  $|y| \to \infty$  in the last step. Thus  $H(x) \equiv C_2$  for all x > 0.

For any  $\tau > 0$ , (3.58) and (3.65) implies that there is a  $x_{\tau} > 0$  such that

$$\int_{-\infty}^{\infty} |\partial_x u(x_\tau, y)|^2 \, dy < \tau^2, \quad \int_{-\infty}^{\infty} |\partial_y u(x_\tau, y)|^2 \, dy < 2\sigma.$$

which further gives

$$|C_2| = |H(x_\tau)| \le \left(\int_{-\infty}^{\infty} |\partial_x u|^2 \, dy\right)^{\frac{1}{2}} \left(\int_{-\infty}^{\infty} |\partial_y u|^2 \, dy\right)^{\frac{1}{2}} \le \sqrt{2\sigma\tau}$$

Since  $\tau$  can be arbitrarily small,  $C_2 = 0$ . This proves (3.71).

From Proposition 3.10, there exists a function  $h_0(x) \in C^2(\mathbb{R}^+ \setminus \mathcal{B}, \mathbb{R})$  such that

$$d_0(u(x,\cdot),\mathcal{U}) = \|u(x,\cdot) - U(\cdot - h_0(x))\|_0, \quad x \in \mathbb{R}^+ \setminus \mathcal{B}$$

For simplicity in the following analysis, we will omit the subscript and denote it as h(x). We have the following identities.

(3.72) 
$$\int_{-\infty}^{\infty} (u(x,y) - U(y - h(x))) \cdot U'(y - h(x)) \, dy = 0, \quad x \in \mathbb{R}^+ \setminus \mathcal{B},$$

(3.73) 
$$h'(x) = \frac{\int_{-\infty}^{\infty} \partial_x u(x,y) \cdot U'(y-h(x)) \, dy}{\int_{-\infty}^{\infty} \left[ |U'(y-h(x))|^2 + U''(y-h(x))(u(x,y) - U(y-h(x))) \right] \, dy}.$$

Identity (3.72) follows directly from

$$\frac{d}{d\delta} \|u(x,\cdot) - U(\cdot - h(x) + \delta)\|_0^2 = 0, \quad \text{at } \delta = 0.$$

since U(y - h(x)) minimizes the  $L^2$  distance to u(x, y). And (3.73) follows from differentiating (3.72) with respect to x.

For the numerator of (3.73), we use (3.71) to deduce

(3.74)  

$$\begin{aligned} \left| \int_{-\infty}^{\infty} \partial_x u(x,y) \cdot U'(y-h(x)) \, dy \right| \\ &= \left| \int_{-\infty}^{\infty} \partial_x u(x,y) \cdot (U'(y-h(x)) - \partial_y u(x,y)) \, dy \right| \\ &\leq \|\partial_x u(x,\cdot)\|_0 \cdot \|u(x,\cdot) - U(\cdot - h(x))\|_1 \\ &\leq \frac{1}{2} \|\partial_x u(x,\cdot)\|_0^2 + \frac{1}{2} \|u(x,\cdot) - U(\cdot - h(x))\|_1^2. \end{aligned}$$

In particular, for  $x \in \mathbb{R} \setminus \mathcal{B}$ , we deduce from (3.67) and (3.70) that

$$(3.75) \qquad \left| \int_{-\infty}^{\infty} \partial_x u(x,y) \cdot U'(y-h(x)) \, dy \right|$$
$$\leq \frac{1}{2} \|\partial_x u(x,\cdot)\|_0^2 + \frac{1}{2\alpha} (J(u(x,\cdot)) - \sigma)$$
$$= \frac{\alpha+1}{2\alpha} \|\partial_x u(x,\cdot)\|_0^2$$

We establish the following lemma.

Lemma 3.12.  $\lim_{x\to\infty} d_0(u(x,\cdot),\mathcal{U}) = 0.$ 

*Proof.* We argue by contradiction. Suppose there exist  $\varepsilon_0 > 0$  and a sequence  $x_n \to \infty$  such that  $d_0(u(x_n, \cdot), \mathcal{U}) \ge \varepsilon_0$  for each n. Without loss of generality, we may assume  $\varepsilon_0 < \varepsilon$ .

When  $d_0(u(x, \cdot), \mathcal{U}) > \frac{\varepsilon_0}{10}$ , Proposition 3.10 implies that  $J(u(x, \cdot)) > \sigma + \alpha(\frac{\varepsilon_0}{\alpha})^2$ , and therefore the set  $\{x : d_0(u(x, \cdot), \mathcal{U}) > \frac{\varepsilon_0}{10}\}$  has finite measure. For a sufficiently large R, we can always find  $x_R > R$  such that

$$d_0(u(x_R,\cdot),\mathcal{U}) \leq \frac{\varepsilon_0}{10}.$$

We define

$$\bar{x}_R := \inf\{x : x > x_R, \, d_0(u(x_R, \cdot) = \varepsilon_0)\},\$$

where the existence of  $\bar{x}_R > x_R$  is guaranteed by the fact that  $x_n \to \infty$ . For any  $x \in (x_R, \bar{x}_R)$ , we have

$$(3.76) \qquad \qquad \frac{d}{dx} |d_0(u(x,\cdot),\mathcal{U})|^2 \\ = \frac{d}{dx} \int_{-\infty}^{\infty} |u(x,y) - U(y - h(x))|^2 dy \\ = \int_{-\infty}^{\infty} (\partial_x u(x,y) - U'(y - h(x))\partial_x h(x)) \cdot (u(x,y) - U(y - h(x))) dy \\ = \int_{-\infty}^{\infty} \partial_x u(x,y) \cdot (u(x,y) - U(y - h(x))) dy \\ \leq \frac{1}{2} \left[ ||\partial_x u(x,\cdot)||_0^2 + |d_0(u(x,\cdot),\mathcal{U})|^2 \right]$$

where we utilize (3.72) from the third line to the fourth line. Moreover, if  $x \in \mathbb{R} \setminus \mathcal{B}$ ,

$$d_0(u(x,\cdot),\mathcal{U})^2 \le \frac{1}{\alpha} \|\partial_x u(x,\cdot)\|_0^2$$

thanks to (3.67). Next we compute

$$(3.77) \qquad \frac{99\varepsilon_0^2}{100} = \int_{x_R}^{\bar{x}_R} \left( \frac{d}{dx} |d_0(u(x,\cdot),\mathcal{U})|^2 \right) dx \\ \leq \int_{(x_R,\bar{x}_R)\setminus\mathcal{B}} \frac{\alpha+1}{2\alpha} ||\partial_x u(x,\cdot)||_0^2 dx + \int_{(x_R,\bar{x}_R)\cap\mathcal{B}} \frac{1}{2} \left( ||\partial_x u(x,\cdot)||_0^2 + d_0(u(x,\cdot),\mathcal{U})^2 \right) dx \\ \leq C \int_{x_R}^{\bar{x}_R} ||\partial_x u||_0^2 dx + \varepsilon_0^2 |(x_R,\bar{x}_R)\cap\mathcal{B}|$$

Since both  $\int_{x>0} |\partial_x u|^2 dz$  and  $|\mathcal{B}|$  are bounded, the last line in (3.77) tends to 0 as  $R \to \infty$ , which yields a contradiction. The proof is complete.

From Lemma 3.12 and Proposition 3.10, there exists  $R_0 > 0$  such that h(x) is a well-defined  $C^3$  function on  $(R_0, \infty)$  and satisfies (3.72), (3.73) and  $d_0(u(x, \cdot), \mathcal{U}) < \varepsilon$ .

When  $x > R_0$ , we can estimate the denominator in (3.73) by

(3.78)  

$$\int_{-\infty}^{\infty} \left[ |U'(y - h(x))|^2 + U''(y - h(x))(u(x, y) - U(y - h(x))) \right] dy$$

$$\geq ||U'||_0^2 - \varepsilon ||U''||_0$$

$$\geq \frac{1}{2} ||U'||_0^2 \geq C,$$

where we use the fact that  $||U''||_0$  is uniformly bounded. By taking  $\varepsilon \ll 1$ , we can ensure that  $\varepsilon ||U''||_0 \le \frac{1}{2} ||U'||_0^2$ .

Combining (3.73), (3.74), (3.75) and (3.78), we obtain

(3.79) 
$$|h'(x)| \leq \begin{cases} C \|\partial_x u(x, \cdot)\|_0^2, & x \in \mathbb{R} \setminus \mathcal{B}, \\ C \left[ \|\partial_x u(x, \cdot)\|_0^2 + \|u(x, \cdot) - U(\cdot - h(x))\|_1^2 \right], & x \in \mathcal{B}. \end{cases}$$

where the constant C only depends on W.

We have the following result on the convergence of h(x).

**Lemma 3.13.** There exists a constant  $h_0$  such that  $\lim_{x \to +\infty} h(x) = h_0$ .

*Proof.* It suffices to show that

 $f^{\infty}$ 

$$\int_{R_0}^{\infty} |h'(x)| \, dx < \infty.$$

Using (3.79), (3.70), (3.58) and (3.69), we deduce that

$$\begin{split} &\int_{R_0} |h'(x)| \, dx \\ \leq C \int_{R_0}^{\infty} \|\partial_x u(x, \cdot)\|_0^2 \, dx + \int_{\mathcal{B}} \|u(x, \cdot) - U(\cdot - h(x))\|_1^2 \, dx \\ \leq C \int_{\{x > R_0\}} |\partial_x u|^2 \, dx + C \int_{\mathcal{B}} \left( \|\partial_y u(x, \cdot)\|_0^2 + \|U'\|_0^2 + \|u(x, \cdot) - U(\cdot - h(x))\|_0^2 \right) \, dx \\ \leq C \left( \int_{\{x > R_0\}} |\partial_x u|^2 \, dx + \int_{\mathcal{B}} J(u(x, \cdot)) \, dx + \|U'\|_0^2 + \varepsilon |\mathcal{B}| \right) \\ \leq C \left( \int_{\{x > R_0\}} |\partial_x u|^2 \, dx + \int_{\mathcal{B}} |\partial_x u|^2 \, dx + \sigma |\mathcal{B}| + \|U'\|_0^2 + \varepsilon |\mathcal{B}| \right) \\ \leq C(W, u). \end{split}$$

The proof is complete.

Lemma 3.12 and Lemma 3.13 together imply that

$$\lim_{x \to +\infty} \|u(x, \cdot) - U(y - h_0)\|_0^2 = 0.$$

Since  $u(x, \cdot) - U(y - h_0)$  are uniformly bounded in  $C^{2,\alpha}(\mathbb{R}, \mathbb{R}^2)$  for any  $\alpha \in (0, 1)$  due to (2.13), the  $L^2$  convergence above implies convergence in  $C^{2,\alpha}(\mathbb{R}, \mathbb{R}^2)$ , which establishes (1.9) for the  $a_1$ - $a_3$  interface. The same arguments also hold for the other two sharp interfaces.

To complete the proof of Theorem 1.1, it remains to prove that the solution u satisfies the pointwise estimate (1.10). To establish this we first observe that (1.9) implies the diffused interface defined in (3.18) must be contained in an  $R_0$ -neighborhood of the sharp interface  $\partial \mathcal{P}$ . Particularly, we have

$$\max_{x \in B(y, \frac{1}{2} \operatorname{dist}(x, \partial \mathcal{P}))} \left( \min_{i} |u(x) - a_{i}| \right) < \delta,$$

for all y satisfying dist $(y, \partial \mathcal{P}) > 2R_0$ . Equation (1.10) then follows from the comparison principle for elliptic equations. This concludes the proof of Theorem 1.1.

### Acknowledgment

The author would like to thank Nicholas D. Alikakos and Changyou Wang for their stimulating discussions and valuable suggestions. The research of Z. Geng was partially supported by AMS-Simons Travel Grant 25014588.

### RIGIDITY OF TRIPLE JUNCTION

### References

- ALIKAKOS, N. D., AND FUSCO, G. Density estimates for vector minimizers and applications. Discrete & Continuous Dynamical Systems 35, 12 (2015), 5631–5663.
- [2] ALIKAKOS, N. D., AND FUSCO, G. A maximum principle for systems with variational structure and an application to standing waves. Journal of the European Mathematical Society 17, 7 (2015), 1547–1567.
- [3] ALIKAKOS, N. D., AND FUSCO, G. Sharp lower bounds for vector Allen-Cahn energy and qualitative properties of minimizes under no symmetry hypotheses. Bulletin of the Hellenic Mathematical Society 67 (2023), 12–58.
- [4] ALIKAKOS, N. D., FUSCO, G., AND SMYRNELIS, P. Elliptic systems of phase transition type. Springer, 2018.
- [5] ALIKAKOS, N. D., AND GENG, Z. On the triple junction problem without symmetry hypotheses. Archive for Rational Mechanics and Analysis 248, 2 (2024), 24.
- [6] AMBROSIO, L., AND CABRÉ, X. Entire solutions of semilinear elliptic equations in ℝ<sup>3</sup> and a conjecture of De Giorgi. Journal of the American Mathematical Society 13, 4 (2000), 725–739.
- [7] BALDO, S. Minimal interface criterion for phase transitions in mixtures of Cahn-Hilliard fluids. Annales de l'Institut Henri Poincaré C, Analyse non linéaire 7, 2 (1990), 67–90.
- [8] BETHUEL, F. Asymptotics for two-dimensional vectorial Allen-Cahn systems, to appear, Acta Mathematica (2021).
- [9] BRONSARD, L., GUI, C., AND SCHATZMAN, M. A three-layered minimizer in  $\mathbb{R}^2$  for a variational problem with a symmetric three-well potential. *Communications on pure and applied mathematics* 49, 7 (1996), 677–715.
- [10] CHAN, H., AND WEI, J. On De Giorgi's conjecture: recent progress and open problems. Science China Mathematics 61 (2018), 1925–1946.
- [11] DE GIORGI, E. Convergence problems for functionals and operators. In Proc. Int. Meeting on Recent Methods in Nonlinear Analysis (1979), Pitagora, Bologna, pp. 131–188.
- [12] DEL PINO, M., KOWALCZYK, M., AND WEI, J. On De Giorgi's conjecture in dimension  $n \ge 9$ . Annals of Mathematics (2011), 1485–1569.
- [13] DEL PINO, M., KOWALCZYK, M., AND WEI, J. Entire solutions of the Allen-Cahn equation and complete embedded minimal surfaces of finite total curvature in ℝ<sup>3</sup>. Journal of Differential Geometry 93, 1 (2013), 67– 131.
- [14] FARINA, A., AND VALDINOCI, E. 1D symmetry for solutions of semilinear and quasilinear elliptic equations. Transactions of the American Mathematical Society 363, 2 (2011), 579–609.
- [15] FLORES, G., PADILLA, P., AND TONEGAWA, Y. Higher energy solutions in the theory of phase transitions: a variational approach. Journal of Differential Equations 169, 1 (2001), 190–207.
- [16] FONSECA, I., AND TARTAR, L. The gradient theory of phase transitions for systems with two potential wells. Proceedings of the Royal Society of Edinburgh Section A: Mathematics 111, 1-2 (1989), 89–102.
- [17] FUSCO, G. Layered solutions to the vector Allen-Cahn equation in ℝ<sup>2</sup>. minimizers and heteroclinic connections. Communications on Pure and Applied Analysis 16, 5 (2017), 1807.
- [18] FUSCO, G. Minimizing under relaxed symmetry constraints: Triple and N-junctions. Annali Scuola Normale Superiore - classe di Scienze (2022).
- [19] FUSCO, G. Connectivity of the diffuse interface and fine structure of minimizers in the Allen-Cahn theory of phase transitions. arXiv preprint arXiv:2401.11575 (2024).
- [20] GAZOULIS, D. On the Γ-convergence of the Allen–Cahn functional with boundary conditions. Proceedings of the Royal Society of Edinburgh Section A: Mathematics (2024), 1–23.
- [21] GENG, Z. Uniqueness of the blow-down limit for triple junction problem. arXiv preprint arXiv:2404.02859 (2024).
- [22] GHOUSSOUB, N., AND GUI, C. On a conjecture of De Giorgi and some related problems. Mathematische Annalen 311, 3 (1998), 481–491.
- [23] GUARACO, M. A. Min-max for phase transitions and the existence of embedded minimal hypersurfaces. Journal of Differential Geometry 108, 1 (2018), 91–133.
- [24] GUI, C. Hamiltonian identities for elliptic partial differential equations. Journal of Functional Analysis 254, 4 (2008), 904–933.
- [25] GUI, C., AND SCHATZMAN, M. Symmetric quadruple phase transitions. Indiana University mathematics journal (2008), 781–836.
- [26] LIU, Y., WANG, K., AND WEI, J. Global minimizers of the Allen–Cahn equation in dimension  $n \ge 8$ . Journal de Mathematiques Pures et Appliquees 108, 6 (2017), 818–840.
- [27] MONTEIL, A., AND SANTAMBROGIO, F. Metric methods for heteroclinic connections in infinite dimensional spaces. arXiv preprint arXiv:1709.02117 (2017).
- [28] PACARD, F., AND WEI, J. Stable solutions of the Allen-Cahn equation in dimension 8 and minimal cones. Journal of Functional Analysis 264, 5 (2013), 1131-1167.

- [29] SANDIER, E., AND STERNBERG, P. Allen-Cahn solutions with triple junction structure at infinity. Communications on Pure and Applied Mathematics 77, 11 (2024), 4163–4211.
- [30] SAVIN, O. Regularity of flat level sets in phase transitions. Annals of Mathematics (2009), 41-78.
- [31] SAVIN, O. Minimal surfaces and minimizers of the Ginzburg Landau energy. Cont. Math. Mech. Analysis AMS 526 (2010), 43–58.
- [32] SCHATZMAN, M. Asymmetric heteroclinic double layers. ESAIM: Control, Optimisation and Calculus of Variations 8 (2002), 965–1005.
- [33] SMYRNELIS, P. Connecting orbits in Hilbert spaces and applications to PDE. Communications on Pure & Applied Analysis 19, 5 (2020), 2797.
- [34] STERNBERG, P. The effect of a singular perturbation on nonconvex variational problems. Archive for Rational Mechanics and Analysis 101, 3 (1988), 209–260.
- [35] STERNBERG, P., AND ZEIMER, W. P. Local minimisers of a three-phase partition problem with triple junctions. Proceedings of the Royal Society of Edinburgh Section A: Mathematics 124, 6 (1994), 1059–1073.
- [36] WANG, K. A new proof of Savin's theorem on Allen-Cahn equations. Journal of the European Mathematical Society 19, 10 (2017), 2997–3051.