KÄHLER-RICCI SOLITONS ON BOUNDED PSEUDOCONVEX DOMAINS

ZEHAO SHA

ABSTRACT. In this paper, we study the Kähler-Ricci soliton on bounded pseudoconvex domains in \mathbb{C}^n . We proved that under certain assumptions, this soliton is a Kähler-Einstein metric. Building on the work of Huang and Xiao, we investigate an analogue of Cheng's conjecture for the Bergman Kähler-Ricci soliton. Additionally, we introduce some model domains as examples.

1. Introduction

In Kähler geometry, there is a classical question to find the "best" metric in a given Kähler class. The Kähler-Einstein metric is usually regarded as a strong contender for this question. Despite this, not every Kähler manifold admits a Kähler-Einstein metric. Extending the notion of Kähler-Einstein metrics, a Kähler-Ricci soliton is a Kähler metric g associated with a real holomorphic vector field X and a constant $\lambda \in \mathbb{R}$, satisfying the equation

$$\operatorname{Ric}(g) + \mathcal{L}_X g = \lambda g.$$

Moreover, if there exist potential functions $f \in C^{\infty}(M)$, such that $X = \frac{1}{2}\nabla f$, we call g a gradient Kähler-Ricci soliton. A Kähler-Ricci soliton g is said to be trivial if X is a Killing vector field, in which case g reduces to a Kähler-Einstein metric with Einstein constant λ . In [Ham88], Hamilton, first established that all compact gradient steady and expanding solitons are Einstein metrics. On compact Fano manifolds (Kähler manifolds with positive first Chern class), if there exists a non-trivial Kähler-Ricci soliton g associated with a non-trivial real holomorphic vector field X, then the Futaki invariant for $X^{1,0}$,

$$F(X^{1,0}) = \int_{M} |X^{1,0}|^{2} \omega_{g}^{n}$$

is strictly positive. Consequently, the existence of a non-trivial Kähler-Ricci soliton obstructs the existence of a Kähler-Einstein metric [Fut83].

In [Per02], Perelman presented that all compact solitons are gradient solitons, implying that steady and expanding solitons must be non-compact. The compactness assumption for Kähler manifolds is crucial for using several powerful tools such as the $\partial \overline{\partial}$ -lemma, the maximum principle, etc. This challenge makes the study of canonical metrics on complete Kähler manifolds significantly less developed compared to their compact counterparts. Bounded pseudoconvex domains in \mathbb{C}^n provide an ideal starting point for addressing this gap. These domains retain essential geometric features of non-compact Kähler manifolds while being simpler to analyze.

Notably, Cheng, Mok, and Yau [CY80, MY83] proved that a bounded domain carries a Kähler-Einstein metric if and only if it is pseudoconvex. Additionally, some significant examples of non-compact gradient Kähler-Ricci soliton were studied, such as in [FIK03, Cao96, Cao97].

In this paper, we study the case of bounded domains (Ω, g) equipped with a complete Kähler metric of C^1 -bounded geometry. It is known that a bounded domain in a complex manifold with C^1 -boundary admitting a Kähler metric is pseudoconvex [Ohs80]. Our main result is the following:

Theorem 1.1. Let $\Omega \subset \mathbb{C}^n$ be a bounded pseudoconvex domain, and let g be a complete Kähler metric on Ω of C^1 -bounded geometry. Suppose there exists a compact subset $K \subset \subset \Omega$ such that $0 > -C_1g \geq \mathrm{Ric}(g) \geq -C_2g$ in $\Omega \setminus K$ for some constants $C_2 \geq C_1 > 0$. If g is a Kähler-Ricci soliton associated with a real holomorphic vector field X whose dual 1-form with respect to g is closed, then g is Kähler-Einstein.

As an important application, based on Huang-Xiao's proof [HX21] for Cheng's conjecture, we have the following analogue statement:

Corollary 1.2. Let Ω be a bounded strictly pseudoconvex domain in \mathbb{C}^n with C^{∞} -boundary and let g_B be the Bergman metric. If g_B is a Kähler-Ricci soliton, then Ω is biholomorphic to the ball.

In general, not all complete Kähler metrics admit strictly negative Ricci curvature near the boundary on bounded pseudoconvex domains. For example, the boundary behavior of the Bergman metric on pseudoconvex domains has been studied in [KY96]. This observation raises an interesting question: what can be said about domains that do not satisfy our assumptions. Specifically, does there exist any bounded pseudoconvex domain in \mathbb{C}^n that admits a non-trivial Kähler-Ricci soliton?

It is worth noting that many domains equipped with complete Kähler metrics satisfy our assumption. In Section 4-6, We will provide a detailed explanation and several examples as applications.

2. PRELIMINARIES

2.1. **Notations.** Let (M^n, g) be an n-dimentional complete Kähler manifold. In local coordinates $(z^1, ..., z^n)$, the Kähler metric can be expressed as $g = g_{i\bar{j}}dz^id\bar{z}^j$, and we will use the Einstein summation convention.

Denote the matrix of metric components by $(g_{i\bar{j}})$ with the inverse $(g^{i\bar{j}})$ for $1 \leq i, j \leq n$, satisfying $g^{k\bar{l}}g_{k\bar{j}}=\delta^l_j$. The Ricci curvature $\mathrm{Ric}(g)$ is defined by

$$\operatorname{Ric}(g) = R_{i\bar{j}} dz^i d\bar{z}^j$$
 where $R_{i\bar{j}} = -\partial_i \partial_{\bar{j}} \log \left(\det(g_{k\bar{l}}) \right)$.

The scalar curvature $R(g)=\operatorname{tr}_g(\operatorname{Ric})=g^{i\bar{j}}R_{i\bar{j}}$ is defined by the trace of the Ricci curvature with respect to g, where ∂_i and $\partial_{\bar{j}}$ are short notations for $\frac{\partial}{\partial z^i}$ and $\frac{\partial}{\partial \bar{z}^j}$ respectively. For any two tensor A,B of type (m,n), the Hermitian inner product of A,B with respect to g is defined by

$$g(A, \bar{B}) := g^{i_1\bar{j}_1} \cdots g^{i_n\bar{j}_n} g_{k_1\bar{l}_1} \cdots g_{k_m\bar{l}_m} A^{k_1 \cdots k_m}_{i_1 \cdots i_m} \overline{B^{l_1 \cdots l_m}_{j_1 \cdots j_n}},$$

and the norm of A is defined by $|A|^2 := g(A, \bar{A})$.

Let ∇ be the associated Levi-Civita connection of the Kähler metric g. Covariant derivatives are denoted by

$$\nabla_i = \nabla_{\partial_i}$$
 and $\nabla_{\bar{j}} = \nabla_{\partial_{\bar{j}}}$.

It is remarkable that the covariant derivative $\nabla_i \nabla_{\bar{j}}$ coincides with the partial derivative $\partial_i \partial_{\bar{j}}$ when applied to C^2 functions, since mixed type Christoffel symbols $\Gamma_{i\bar{j}}$ vanish. For any C^{∞} function $f: M \to \mathbb{R}$, the gradient vector field of f is defined by

$$\nabla f := g^{i\bar{j}} \partial_i f \cdot \partial_{\bar{j}} + g^{i\bar{j}} \partial_{\bar{j}} f \cdot \partial_i.$$

Let $\nabla^2 f := (\partial_i \partial_{\bar{j}} f)$ be the complex Hessian of f, then the Laplacian is defined by the trace of $\nabla^2 f$ with respect to g, that is

$$\Delta f := \operatorname{tr}_g \left(\nabla^2 f \right) = g^{i\bar{j}} \partial_i \partial_{\bar{j}} f.$$

2.2. **Kähler-Einstein metric and Kähler-Ricci soliton.** We say a Kähler metric is Kähler-Einstein if the Ricci curvature is proportional to the metric, that is

$$Ric(g) = \lambda g,$$
 (2.1)

for some $\lambda \in \mathbb{R}$. By rescaling the metric, we can assume that $\lambda = 1, 0$ or -1.

A vector field X is said to be real if $\overline{X} = X$, where \overline{X} is the complex conjugate of X. Recall that a real holomorphic vector field X is a real vector field such that its (1,0)-part, that is $X^{1,0} := X - \sqrt{-1}JX$ is holomorphic for the fixed complex structure J. We say a Kähler metric g is a Kähler-Ricci soliton if there is an associated real holomorphic vector field X and a constant $X \in \mathbb{R}$ such that

$$Ric(q) + \mathcal{L}_X q = \lambda q, \tag{2.2}$$

where $\mathcal{L}_X g$ is the Lie derivative of the Kähler metric g along X. Suppose that there exists a C^{∞} real-valued function f such that $X = \frac{1}{2}\nabla f$, then we say g is a gradient Kähler-Ricci soliton and f is the potential function. The assumption of ∇f being a real holomorphic vector field is equivalent to $\nabla_i \nabla_j f = 0$ for all $1 \le i, j \le n$. The Kähler-Ricci soliton is said to be shrinking if $\lambda > 0$, steady if $\lambda = 0$, and expanding if $\lambda < 0$. Note that if the Lie derivative term vanishes, the soliton equation (2.2) reduces to the Kähler-Einstein equation (2.1).

2.3. Bounded geometry and Omori-Yau's maximum principle. To establish our main result, we first introduce the concept of (quasi-)bounded geometry. Recall that the injectivity radius at a point $x \in M$ is the maximum radius r of the ball B_r in the tangent space T_xM for which the exponential map $\exp_x : B_r \to \exp_x(B_r) \subset M$ is a diffeomorphism. The injectivity radius of M is the infimum of the injectivity radius at all points in M.

Definition 2.1. Let (M, g) be a complete Kähler manifold and let $k \ge 0$ be an integer. We say (M, g) has C^k -quasi-bounded geometry if for each non-negative integer $l \le k$, there exists a constant $C_l > 0$ such that

$$\sup_{M} |\nabla^{l} \operatorname{Rm}| \le C_{l}, \tag{2.3}$$

where $\mathrm{Rm}=\{R_{i\bar{j}k\bar{l}}\}$ is the Riemann curvature tensor of g and ∇^l is the convariant derivative of order l. Moreover, if (M,g) has a positive injectivity radius, then we say (M,g) has C^k -bounded geometry.

Next, let us introduce Omori-Yau's generalized maximum principle on non-compact manifolds which serves as a crucial tool in our approach.

Proposition 2.2 ([Omo67, Yau75]). Let (M, g) be a complete Kähler manifold. Assume that g has bounded sectional curvature, then for any function $u \in C^2(M)$ with $\sup_M u < +\infty$, there exists a sequence of points $\{z_k\}_{k\in\mathbb{N}} \subset M$ satisfying

$$(1) \lim_{k \to \infty} u(z_k) = \sup_{M} u, \ (2) \lim_{k \to \infty} |\nabla u(z_k)| = 0, \ (3) \lim_{k \to \infty} \nabla^2 u(z_k) \le 0,$$

where the last inequality holds in the sense of matrices.

Remark 2.3. If we replace the assumption of bounded sectional curvature in the generalized maximum principle with bounded Ricci curvature, then (3) is replaced by

$$(3')\lim_{k\to\infty}\Delta u(z_k)\leq 0.$$

3. Proof of Theorem 1.1

Let $\Omega\subset\mathbb{C}^n$ be a bounded pseudoconvex domain with a complete Kähler metric g. Suppose that g is a Kähler-Ricci soliton satisfying

$$\operatorname{Ric}(g) + \mathcal{L}_X g = \lambda g,$$

for a real holomorphic vector field X and a constant $\lambda \in \mathbb{R}$.

Proposition 3.1. If the Ricci curvature is bounded from below, that is $Ric(g) \ge Cg$ for some constant C, then $\lambda < 0$.

Proof. We first claim that C < 0. If not, suppose C > 0, then by Myers's theorem, (Ω, g) is compact which is a contradiction. By the general fact that there is no positive non-constant harmonic function on complete manifolds with non-negative Ricci curvature [SY10, Corollary

3.1], the case C=0 can also be ruled out since we are on bounded domains in \mathbb{C}^n . Thus we have C<0, and there exists a point $z\in\Omega$ such that $\mathrm{Ric}(g)=-\varepsilon g$ at z for some $\varepsilon>0$ small enough. This implies that the scalar curvature R(z)<0 which contradicts the sharp scalar curvature lower bound estimates for Ricci solitons [Cho23, Theorem 2.14] if we have $\lambda\geq0$.

Since our assumption ensures that the Ricci curvature is negatively pinched near the boundary, it is clear that the Ricci curvature is bounded from below globally. Without loss of generality, we may assume $\lambda=-1$. Let us recall the following equation for the scalar curvature R in this case.

Lemma 3.2 ([AMR16], Proposition 8.3). Let S := R + n, then

$$\Delta S - \langle X, \nabla S \rangle - S + |\operatorname{Ric}(g) + g|^2 = 0. \tag{3.1}$$

Notably, the scalar curvature can only achieve its minimum on the boundary thanks to the sharp scalar curvature lower bound estimates for Ricci solitons. By applying the generalized maximal principle to (3.1), we obtain the asymptotic behavior for the Kähler-Ricci soliton at boundary points where the scalar curvature achieves its minimum.

Proposition 3.3. Let $\Omega \subset \mathbb{C}^n$ be a bounded pseudoconvex domain, and let g be a complete Kähler-Ricci soliton associated with a real holomorphic vector field X such that $|\operatorname{Ric}(g)| \leq K$ for some constant K > 0. If R(p) = -n for some $p \in \partial \Omega$, that is for any sequence $\{p_m\}$ which converges to p, we have $R(p_m) \to -n$ as $p_m \to p$, and $|X|(z) \leq |\nabla R|^{-1}(z) \cdot d(z, \partial \Omega)$ for all $z \in \Omega$ where d is the Euclidean distance, then there exists a sequence $p_k \to p$ such that

$$\lim_{k \to \infty} |\operatorname{Ric}(g) + g|(p_k) = 0.$$

Proof. For any $p \in \partial \Omega$ such that R(p) = -n, we may choose a sequence $\{p_k\}$ satisfying the generalized maximum principle and it follows from the equation (3.1),

$$0 \le \lim_{k \to \infty} \Delta S(p_k) = \lim_{k \to \infty} \left(\langle \nabla S, X \rangle (p_k) + S(p_k) - |\operatorname{Ric}(g) + g|^2(p_k) \right).$$

Since we have $|X|(p_k) \leq |\nabla R|^{-1}(p_k) \cdot d(p_k, \partial \Omega)$, this yields

$$0 \leq \lim_{k \to \infty} \Delta S(p_k)$$

$$\leq \lim_{k \to \infty} d(p_k, \partial \Omega) + \lim_{k \to \infty} S(p_k) - \lim_{k \to \infty} |\operatorname{Ric}(g) + g|^2(p_k)$$

$$\leq -\lim_{k \to \infty} |\operatorname{Ric}(g) + g|^2(p_k)$$

$$\leq 0.$$

Thus, we have

$$\lim_{k \to \infty} |\operatorname{Ric}(g) + g|(p_k) = 0,$$

which is the conclusion.

It follows from the computation of the Bochner formula, we can derive the following equation for real holomorphic vector fields.

Lemma 3.4. Assume that g is a complete Kähler metric and X is a real holomorphic vector field, then we have

$$\Delta |X|^2 = |\nabla X|^2 - \text{Ric}(X, X). \tag{3.2}$$

Proof. We compute this formula in local coordinates. The assumption of X being real holomorphic is equivalent to $\nabla_i X_j = 0$ for any i, j, and we will use cancellation notation $\nabla_i X_j$ in the computation if we use this condition. Let $\nabla^i = g^{i\bar{j}} \nabla_{\bar{i}}$, then we have

$$\begin{split} \Delta |X|^2 &= \nabla^i \nabla_i \left(X^k \cdot X_k \right) \\ &= \nabla^i \left(\nabla_i X^k \cdot X_k + \underbrace{X^k}_i \nabla_i \widehat{X_k} \right) \\ &= \nabla^i X_k \cdot \nabla_i X^k + X_k \cdot \nabla^i \nabla_i X^k, \end{split}$$

where the cancellation notation means we use the real holomorphicity of X. Denote the first term by $|\nabla X|^2$ and commute ∇^i with ∇_i in the second term, we obtain

$$\Delta |X|^2 = |\nabla X|^2 + X_k \cdot \left(-R_l^k \cdot X^l + \nabla_i \nabla^i X^k \right)$$

$$= |\nabla X|^2 - R_{l\bar{k}} X^l X^{\bar{k}}$$

$$= |\nabla X|^2 - \text{Ric}(X, X),$$

which completes the proof.

Remark 3.5. On compact Kähler manifolds with non-positive first Chern class, as a direct consequence of the equation (3.2), there is no non-trivial holomorphic vector field.

Assume that g is a complete Kähler-Ricci soliton with C^1 -bounded geometry and ω_g is the corresponding Kähler form. Denote $X^{\mathfrak{b}} = (X^{\mathfrak{b}})^{1,0} + (X^{\mathfrak{b}})^{0,1}$ the dual 1-form of the real holomorphic vector field X where $(X^{\mathfrak{b}})^{0,1} := \omega_g(X^{1,0},\cdot)$. If $X^{\mathfrak{b}}$ is closed, then for any complex vector field Y, Z, we have

$$0 = dX^{\mathfrak{b}}(Y, Z) = (\nabla_Y X^{\mathfrak{b}})(Z) - (\nabla_Z X^{\mathfrak{b}})(Y).$$

We first claim that the real holomorphic vector field X is unique up to constant in our setting. Otherwise, suppose that X and \tilde{X} are two real holomorphic vector fields for g with the same soliton constant such that their dual 1-forms are closed. Taking $Y = X - \tilde{X}$, then for any $1 \le i, j \le n$, the following equations hold:

$$\begin{cases} \nabla_j Y_i = 0; \\ \nabla_{\bar{j}} Y_i = 0, \end{cases}$$

where the first equation follows from Y is real holomorphic, and the second equation follows from the soliton equation of X and \tilde{X} , respectively. Then applying $g^{k\bar{j}}\nabla_k$ to the second equation, we have

$$0 = g^{k\bar{j}} \nabla_k \nabla_{\bar{j}} Y_i = g^{k\bar{j}} \left(\nabla_{\bar{j}} \nabla_k Y_i - R_{k\bar{j}i\bar{l}} Y^{\bar{l}} \right) = -R_{i\bar{l}} Y^{\bar{l}},$$

which implies $\mathrm{Ric}(Y,Y)=0$. Since $\mathrm{Ric}(g)$ is negatively pinched outside a compact subset $K\subset\subset\Omega$, it follows that in $\Omega\setminus K$,

$$0 = \operatorname{Ric}(Y, Y) \le -C|Y|^2,$$

for some constant C>0. Thus, Y=0 in $\Omega\setminus K$. We then conclude Y=0 in Ω thanks to the real analyticity of Y.

We can now proceed to the proof of the main theorem.

Proof of Theorem 1.1. Consider the soliton equation in local coordinates,

$$R_{i\bar{j}} + \nabla_i X_{\bar{j}} + \nabla_{\bar{j}} X_i = -g_{i\bar{j}}. \tag{3.3}$$

Since $X^{\mathfrak{b}}$ is closed, we have

$$\nabla_i X_{\bar{i}} = \nabla_{\bar{i}} X_i.$$

Then applying $g^{k\bar{j}}\nabla_k$ to both sides of (3.3),

$$g^{k\bar{j}}\nabla_k R_{i\bar{j}} + 2g^{k\bar{j}}\nabla_k \nabla_{\bar{j}} X_i = 0.$$

By using contracted Bianchi identity and commuting ∇_k with $\nabla_{\bar{i}}$, we obtain

$$0 = g^{k\bar{j}} \nabla_k R_{i\bar{j}} + 2g^{k\bar{j}} \nabla_k \nabla_{\bar{j}} X_i$$

$$= \nabla_i R + 2g^{k\bar{j}} \left(\nabla_{\bar{j}} \nabla_k X_i - R_{k\bar{j}i\bar{l}} X^{\bar{l}} \right),$$

where $\nabla_{\bar{i}}\nabla_k X_i = 0$ since X is real holomorphic. Then, we obtain,

$$\nabla_i R = 2R_{i\bar{l}} X^{\bar{l}},$$

or in a coordinate-free version

$$g(\nabla R, \cdot) = 2\operatorname{Ric}(X, \cdot).$$
 (3.4)

It is clear that |X| is bounded on the compact subset $K \subset\subset \Omega$. On $\Omega\setminus K$, we have $-C_1g\leq \mathrm{Ric}\leq -C_2g$ for some constant $C_1,C_2>0$. Then it follows from (3.4),

$$g(\nabla R, X) = 2\operatorname{Ric}(X, X) \le -2C_2|X|^2,$$

which implies

$$|X|^2 \le -\frac{1}{2C_2}g(\nabla R, X) \le \frac{1}{2C_2}|X| \cdot |\nabla R|.$$

Thus, we have

$$\sup_{\Omega} |X| \leq \max \left\{ \max_{K} |X|, \frac{\sup_{\Omega} |\nabla R|}{2C_2} \right\} < +\infty,$$

thanks to g has C^1 -bounded geometry.

Plugging X into the soliton equation, we obtain

$$Ric(X, X) + \mathcal{L}_X g(X, X) = -|X|^2.$$

Observe that $\mathcal{L}_X g(X,X) = \nabla_X |X|^2$, we then have

$$Ric(X, X) = -\nabla_X |X|^2 - |X|^2.$$

We replace the Ricci curvature in (3.2) by this formula, then

$$\Delta |X|^2 = |\nabla X|^2 + |\nabla X|X|^2 + |X|^2 \ge |\nabla X|X|^2 + |X|^2.$$

Since $\sup_{\Omega} |X| < +\infty$, by the generalized maximum principle, there exists a sequence $\{z_k\}$ such that

$$0 \ge \lim_{k \to \infty} \Delta_g |X|^2(z_k) \ge \lim_{k \to \infty} \nabla_X |X|^2(z_k) + \lim_{k \to \infty} |X|^2(z_k) = \sup_{\Omega} |X|^2 \ge 0,$$

which implies |X| = 0. This concludes that g is Kähler-Einstein.

Remark 3.6. It is important to note that the assumption that $X^{\mathfrak{b}}$ is closed does not mean that X is a gradient vector field of some potential functions. In the special case of the gradient Kähler-Ricci soliton, we can apply the generalized maximum principle to the PDE

$$\Delta f - |\nabla f|^2 - f = 0,$$

of the potential function f, to show that f = 0. The fact that $\sup_{\Omega} f < +\infty$ can be obtained by using a similar argument.

4. STRICTLY PSEUDOCNVEX DOMAINS

Let Ω be a bounded strictly pseudoconvex domain in \mathbb{C}^n and let $A^2(\Omega)$ be the space of holomorphic functions in $L^2(\Omega)$. It is clear that $A^2(\Omega)$ is a Hilbert space. The Bergman kernel K(z) on Ω is a real analytic function given by

$$K(z) = \sum_{j=1}^{\infty} |\varphi_j(z)|^2, \quad \forall z \in \Omega,$$

where $\{\varphi_j\}_{j=1}^\infty$ is an orthonormal basis of $A^2(\Omega)$ with respect to the L^2 inner product. Since the Bergman kernel is positive and independent of the choice of any orthonormal basis [Kra01] on bounded domains, we then can define an invariant metric on Ω called the Bergman metric by

$$g_B := g_{i\bar{j}} dz^i d\overline{z}^j$$
 where $g_{i\bar{j}} = \partial_i \partial_{\bar{j}} \log K$.

The Bergman metric is a complete real analytic Kähler metric, with the real analyticity derived from the Bergman kernel. On strictly pseudoconvex domains with C^{∞} -boundary, the sectional curvature of the Bergman metric is asymptotic to a negative constant, and, the Bergman metric has C^{∞} -bounded geometry [GKK11]. Because of the boundary behavior of the Ricci curvature of g_B [KY96, Corollary 2], there always exists a compact subset $K \subset \Omega$ such that $\mathrm{Ric}(g_B)$ is bounded and strictly negative outside K.

Let $G := \det \left(g_{i\bar{j}}\right)$ denote the determinant of the Bergman metric. The Bergman invariant function B(z) := G(z)/K(z), introduced by Bergman in [BS51], is invariant under any biholomorphic map between two domains. A significant result regarding the Bergman invariant function, established by Diederich in [Die70], states that as we approach the boundary of strictly pseudoconvex domains, B tends to $(n+1)^n \pi^n/n!$. It was proved in [FW97] that if B(z) is constant in Ω then the Bergman metric is Kähler-Einstein.

Proposition 4.1. Let $\Omega \subset \mathbb{C}^n$ be a bounded strictly pseudoconvex domain with C^∞ -boundary. Assume that the Bergman invariant function satisfies $B(z) = e^{f(z)}$ for some $f \in C^\infty(\Omega)$ such that ∇f is a real holomorphic vector field, then the Bergman metric is a gradient Kähler-Ricci soliton.

It was conjectured by S.-Y. Cheng [Che79] that if the Bergman metric of a bounded strictly pseudoconvex domain with C^{∞} -boundary is Kähler–Einstein, then the domain is biholomorphic to the ball. In [HX21], Huang and Xiao provided an affirmative answer to Cheng's conjecture.

Theorem 4.2 ([HX21], Theorem 1.1). The Bergman metric of a bounded strictly pseudoconvex domain Ω with C^{∞} -boundary is Kähler-Einstein if and only if the domain is biholomorphic to the ball.

By combining Theorem 4.2 and 1.1, we immediately see the following:

Corollary 4.3. Let $\Omega \subset \mathbb{C}^n$ be a bounded strictly pseudoconvex domain with C^{∞} -boundary and let g_B be the Bergman metric. Suppose that g_B is a Kähler-Ricci soliton, then Ω is biholomorphic to the ball.

For the remaining part of this section, we assume that Ω admits a C^{k+2} , $k \geq 5$, defining function ρ such that $\rho = 0$, $d\rho \neq 0$ on $\partial\Omega$, $\left(\rho_{i\bar{j}}\right) > 0$ on $\overline{\Omega}$ and $\Omega = \{\rho < 0\}$, where $\rho_{i\bar{j}} := \partial_i \partial_{\bar{j}} \rho$. Let $\phi = -\log(-\rho)$, we can define another complete Kähler metric on Ω of C^{k-2} -bounded geometry introduced in [CY80], that is

$$g_{\rho}:=g_{i\overline{\jmath}}dz^id\overline{z}^j\quad\text{where}\quad g_{i\overline{\jmath}}:=\partial_i\partial_{\overline{\jmath}}\phi.$$

A direct computation gives

$$g_{i\bar{j}} = \frac{\rho_{i\bar{j}}}{-\rho} + \frac{\rho_{i}\rho_{\bar{j}}}{\rho^{2}}, \qquad g^{i\bar{j}} = (-\rho)\left(\rho^{i\bar{j}} + \frac{\rho^{i}\rho^{\bar{j}}}{\rho - |d\rho|^{2}}\right)$$

and

$$\det (g_{i\bar{j}}) = \left(-\frac{1}{\rho}\right)^n \det \left(\rho_{i\bar{j}} - \frac{\rho_i \rho_{\bar{j}}}{\rho}\right)$$
$$= \left(\frac{1}{\rho}\right)^{n+1} \det \left(\rho_{i\bar{j}}\right) \left(-\rho + |d\rho|^2\right),$$

where
$$\rho_i = \partial_i \rho$$
, $\left(\rho^{i\bar{j}}\right) = \left(\rho_{i\bar{j}}\right)^{-1}$, $\rho^i = \rho^{i\bar{j}}\rho_{\bar{j}}$ and $|d\rho|^2 = \rho^{i\bar{j}}\rho_i\rho_{\bar{j}}$.

Since we have

$$g^{i\bar{j}}g_ig_{\bar{j}} = \frac{|d\rho|^2}{|d\rho|^2 - \rho} \le 1,$$

it follows from [Dem12, p366 (2.4) Lemma] that, g_{ρ} is a complete Kähler metric. The Ricci curvature $Ric(g_{\rho})$ is given by

$$R_{i\bar{j}} = -\partial_i \partial_{\bar{j}} \left(\log \left(\det \left(g_{k\bar{l}} \right) \right) \right)$$

= $-(n+1)g_{i\bar{j}} - \partial_i \partial_{\bar{j}} \log \left[\det \left(\rho_{k\bar{l}} \right) \left(-\rho + |d\rho|^2 \right) \right].$

Note that $\det\left(\rho_{k\bar{l}}\right)\left(-\rho+|d\rho|^2\right)$ is a positive C^{k-2} function defined on $\overline{\Omega}$. It can be computed that

$$\partial_i \partial_{\bar{i}} \log \left[\det \left(\rho_{k\bar{l}} \right) \left(-\rho + |d\rho|^2 \right) \right]$$

is a tensor whose norm tends to 0 near the boundary, so the Ricci curvature asymptotically tends to -(n+1). Therefore, we have the following:

Corollary 4.4. Let Ω be a bounded strictly pseudoconvex domain in \mathbb{C}^n with C^k -boundary for $k \geq 5$ and let g_ρ be the complete Kähler metric defined above. Suppose that g_ρ is a Kähler-Ricci soliton, then g_ρ is the unique Kähler-Einstein metric on strictly pseudoconvex domains constructed by Cheng-Yau.

5. Uniformly squeezing domains

The uniform squeezing property was introduced by Liu et al. [LSY05] and by Yeung [Yeu09] independently in order to study canonical invariant metrics on complex manifolds.

Definition 5.1. A complex manifold M of dimension n is called uniformly squeezing if there exists 0 < r < R such that for any $p \in M$, there is a holomorphic map $f_p : M \to \mathbb{C}^n$ satisfying

- (1) $f_p(p) = 0$;
- (2) $f_p: M \to f_p(M)$ is biholomorphic;
- (3) $B_r \subset f_p(M) \subset B_R$, where B_r and B_R are Euclidean balls in \mathbb{C}^n with radius r and R centred at 0.

It is known that all bounded homogeneous domains, bounded domains that are covers of compact Kähler manifold, and strongly convex domains with C^2 -boundary satisfy the uniformly squeezing property. It was proved in [Yeu09] that uniformly squeezing domains equipped with the Bergman metric or Kähler-Einstein metric have C^{∞} -bounded geometry and thus, the Ricci curvature is bounded.

In [DGZ12], Deng-Guan-Zhang introduced the concept of the squeezing function to study the geometric and analytic properties of uniformly squeezing domains.

Definition 5.2. Let Ω be a bounded domain in \mathbb{C}^n . For any $z \in \Omega$ and any holomorphic embedding $f: \Omega \to B_1$ with f(z) = 0, we set

$$s_{\Omega}(z, f) := \sup\{r; B_r \subset f(\Omega)\}.$$

The squeezing function of Ω is defined by

$$s_{\Omega}(z) := \sup_{f} \{ s_{\Omega}(z, f) \}.$$

By definition, we see that $0 \le s_{\Omega}(z) \le 1$ and $s_{\Omega}(z)$ admits a positive lower bound if and only if Ω is uniformly squeezing. In [Zha15], Zhang established the curvature estimate of the Bergman metric in terms of the squeezing function.

Theorem 5.3 ([Zha15], Theorem 1.1). Let Ω be a bounded domain in \mathbb{C}^n and let s_{Ω} be the squeezing function. Denoting H(z,W), $\mathrm{Ric}(z,W)$ and R(z) by the holomorphic sectional curvature, the Ricci curvature and the scalar curvature at z in the direction W, respectively. Then we have

$$2 - 2\frac{n+2}{n+1}s_{\Omega}^{-4n}(z) \le H(z,W) \le 2 - 2\frac{n+2}{n+1}s_{\Omega}^{4n}(z)$$
$$(n+1) - (n+2)s_{\Omega}^{-2n}(z) \le \text{Ric}(z,W) \le (n+1) - (n+2)s_{\Omega}^{2n}(z),$$
$$n(n+1) - n(n+2)s_{\Omega}^{-2n}(z) \le R(z) \le n(n+1) - n(n+2)s_{\Omega}^{2n}(z).$$

Thanks to the curvature estimate, we see that if the squeezing function is close to 1 outside a compact subset $K \subset\subset \Omega$, then the Ricci curvature is strictly negative on $\Omega\setminus K$. We can conclude the following:

Corollary 5.4. Let Ω be a bounded domain in \mathbb{C}^n and let s_{Ω} be the squeezing function. Suppose there exists $0 < \varepsilon < 1$ such that

$$\left(\frac{n+1+\varepsilon}{n+2}\right)^{\frac{1}{2n}} \le s_{\Omega} \le 1,$$

in $\Omega \setminus K$ for some compact subset $K \subset\subset \Omega$. If g_B is a Kähler-Ricci soliton, then g_B is Kähler-Einstein.

It was proven in [DGZ16] that if Ω is a bounded strictly pseudoconvex domain with C^2 -boundary, then $\lim_{z\to\partial\Omega}s_\Omega(z)=1$. Conversely, this statement is not always true, and we refer the interested reader to see [FW18] for a counterexample. In [Zim18], Zimmer proved that for a bounded convex domain Ω with C^∞ -boundary, if $\lim_{z\to\partial\Omega}s_\Omega(z)=1$, then Ω is strictly pseudoconvex.

6. Examples

6.1. Thullen domain in \mathbb{C}^2 . The Thullen domain $\Omega_m := \{(z, w) \in \mathbb{C}^2; |z|^2 + |w|^{2m} < 1\}$ is a bounded pseudoconvex domain in \mathbb{C}^2 for any $m \geq 1$. All boundary points are strictly pseudoconvex away from the boundary part $\{|z| = 1\}$. If we set m = 1, the Thullen domain

becomes the ball. If m>1, then $-\log(1-|z|^2-|w|^{2m})$ is not strictly plurisubharmonic and therefore $g_{i\bar{j}}=-\partial_i\partial_{\bar{j}}\log(1-|z|^2-|w|^{2m})$ is not a Kähler metric. Instead, we consider another defining function $\rho(z,w)=(1-|z|^2)^{1/m}-|w|^2$, and the corresponding complete Kähler metric given by ρ introduced in [Seo12] is

$$g_m(z, w) = -\partial_i \partial_{\bar{i}} \log \rho(z, w).$$

The specific expression of g_m is

$$\frac{(1-|z|^2)^{\frac{1}{m}-2}}{m\rho^2} \begin{pmatrix} \rho + \frac{1}{m}|z|^2|w|^2 & w\overline{z}(1-|z|^2) \\ \overline{w}z(1-|z|^2) & m(1-|z|^2)^2 \end{pmatrix},$$

and the determinant of g_m is

$$G := \det g_m = \frac{\frac{1}{m} (1 - |z|^2)^{\frac{2}{m} - 2}}{\rho^3}.$$

Then the Ricci curvature $\operatorname{Ric}(g_m) = -\partial_i \partial_{\bar{j}} \log G$ is computed by

$$\operatorname{Ric}(g_m) = -\frac{3(1-|z|^2)^{\frac{1}{m}-2}}{m\rho^2} \begin{pmatrix} \frac{2}{3}(m-1)\rho^2(1-|z|^2)^{-\frac{1}{m}} + \rho + \frac{1}{m}|z|^2|w|^2 & w\overline{z}(1-|z|^2) \\ \overline{w}z(1-|z|^2) & m(1-|z|^2)^2 \end{pmatrix}.$$

It is clear that g_m is Kähler-Einstein if and only if m = 1. By a straightforward computation, we have the following.

Proposition 6.1. Suppose g_m is a Kähler-Ricci soliton, then m=1.

Proof. Assume g_m is a Kähler-Ricci soliton associated with a real holomorphic vector field $X = X^i \partial_i + \overline{X}^i \partial_{\overline{i}}$ where X^i is a holomorphic function, that is

$$R_{i\bar{j}} + \nabla_i X_{\bar{j}} + \nabla_{\bar{j}} X_i = \lambda g_{i\bar{j}}, \tag{6.1}$$

for some constant λ and i, j = 1 or 2. We then compute

$$\nabla_i X_{\bar{i}} + \nabla_{\bar{i}} X_i = X^k \partial_k g_{i\bar{i}} + \overline{X}^l \partial_{\bar{i}} g_{i\bar{i}} + \partial_i X^k g_{k\bar{i}} + \partial_{\bar{i}} \overline{X}^l g_{i\bar{l}}.$$

First, fixing i = j = 1. Denote $\partial_1 = \partial_z$, $\partial_2 = \partial_w$, we have

$$X^{k}\partial_{k}g_{1\bar{1}} = X^{1}\partial_{1}g_{1\bar{1}} + X^{2}\partial_{2}g_{1\bar{1}}$$

$$= \frac{X^{1}\overline{z}}{m^{3}\rho^{3}(1-|z|^{2})^{3-\frac{1}{m}}} \left(2m^{2}\left(1-|z|^{2}\right)^{\frac{2}{m}} + \left((1-m)|z|^{2} + 2m^{2} - 2m\right)|w|^{4} + \left(1-|z|^{2}\right)^{\frac{1}{m}}\left((m+1)|z|^{2} - 4m^{2} + 2m\right)|w|^{2}\right)$$

$$+ \frac{X^{2}\overline{w}}{m^{2}\rho^{3}(1-|z|^{2})^{2-\frac{1}{m}}} \left(\left(|z|^{2} - m\right)|w|^{2} + \left(1-|z|^{2}\right)^{\frac{1}{m}}\left(|z|^{2} + m\right)\right),$$

and

$$\begin{split} \partial_i X^k g_{k\bar{j}} = & \partial_1 X^1 g_{1\bar{1}} + \partial_1 X^2 g_{2\bar{1}} \\ = & \frac{\rho + \frac{1}{m} |z|^2 |w|^2}{m \rho^2 (1 - |z|^2)^{2 - \frac{1}{m}}} \cdot \partial_1 X^1 + \frac{z \overline{w}}{m \rho^2 (1 - |z|^2)^{1 - \frac{1}{m}}} \cdot \partial_1 X^2. \end{split}$$

By combining the above computations and rearranging all terms in (6.1), we obtain

$$2(m-1)\rho^{2}(1-|z|^{2})^{-\frac{1}{m}} + (3+\lambda)\left(\rho + \frac{1}{m}|z|^{2}|w|^{2}\right)$$

$$= \frac{X^{1}\overline{z} + \overline{X}^{1}z}{m^{2}(1-|z|^{2})\rho}\left(2m^{2}\left(1-|z|^{2}\right)^{\frac{2}{m}} + ((1-m)|z|^{2} + 2m^{2} - 2m)|w|^{4} + \left(1-|z|^{2}\right)^{\frac{1}{m}}\left((m+1)|z|^{2} - 4m^{2} + 2m\right)|w|^{2}\right)$$

$$+ \frac{X^{2}\overline{w} + \overline{X}^{2}w}{m\rho}\left(\left(|z|^{2} - m\right)|w|^{2} + \left(1 - |z|^{2}\right)^{\frac{1}{m}}\left(|z|^{2} + m\right)\right)$$

$$+ \left(\rho + \frac{1}{m}|z|^{2}|w|^{2}\right)\left(\partial_{1}X^{1} + \partial_{\overline{1}}\overline{X}^{1}\right) + \left(1 - |z|^{2}\right)\left(\partial_{1}X^{2}\overline{w}z + \partial_{\overline{1}}\overline{X}^{2}w\overline{z}\right).$$

$$(6.2)$$

Then, we take partial derivatives with respect to w and \overline{w} twice (that is, applying $\partial_2^2 \partial_{\overline{2}}^2$) to both sides of (6.2). By evaluating at (z,0), we get

$$\frac{4|z|^2}{m(1-|z|^2)} \left(X^1 \overline{z} + \overline{X}^1 z \right) + |z|^2 \left(\partial_2 X^2 + \partial_{\overline{2}} \overline{X}^2 \right) = 4m(m-1). \tag{6.3}$$

Next, fixing i = j = 2. By taking the same approach as in the above computations, we have

$$4\left(X^{1}\overline{z} + \overline{X}^{1}z\right) + m(1 - |z|^{2})\left(\partial_{2}X^{2} + \partial_{\overline{2}}\overline{X}^{2}\right) = 0, \tag{6.4}$$

which implies

$$\partial_2 X^2 + \partial_{\overline{2}} \overline{X}^2 = -\frac{4}{m(1-|z|^2)} \left(X^1 \overline{z} + \overline{X}^1 z \right). \tag{6.5}$$

If $\partial_2 X^2 + \partial_{\overline{2}} \overline{X}^2 \neq 0$, then plugging (6.5) into (6.3), we obtain

$$4m(m-1) = 0,$$

which yields m = 1, since m > 1.

If $\partial_2 X^2 + \partial_{\overline{2}} \overline{X}^2 = 0$, from (6.4), we have $X^1 \overline{z} + \overline{X}^1 z = 0$. Then, plugging into (6.3), we still have m = 1.

Remark 6.2. By the expression of $Ric(g_m)$, we can see that $Ric(g_m)$ is negatively pinched in Ω_m . Thus, Proposition 6.1 can be deduced from Theorem 1.1 directly.

We now consider the Bergman metric on the Thullen domain. The Bergman kernel of the Thullen domain Ω_m is given by

$$K(z,w) = \frac{m\pi^2}{m+1} \cdot \frac{(m+1)\left(1-|z|^2\right)^{\frac{1}{m}} - (m-1)|w|^2}{(m+1)\rho^3\left(1-|z|^2\right)^{2-\frac{1}{m}}}.$$

The Bergman metric g_B can then be computed. An interesting fact between the two metrics g_m and g_B on the Thullen domain is that they only coincide when Ω_m is the ball, that is m=1.

Proposition 6.3. Let $g_B = \lambda g_m$ for some $\lambda > 0$, then m = 1.

Proof. Assume $g_B = \lambda g_m$ for some $\lambda > 0$, then for some pluriharmonic functions h, we have

$$\log K + \lambda \log \rho = h. \tag{6.6}$$

Since the left-hand side of (6.6) only depends on |z| and |w|, so does the right-hand side. Then by the maximum principle, we obtain that h=C for some constant C. Evaluating at the origin, we get $C=\log\left(\frac{m\pi^2}{m+1}\right)$.

Thus, we have

$$\frac{(m+1)\left(1-|z|^2\right)^{\frac{1}{m}}-(m-1)|w|^2}{(m+1)\rho^{3-\lambda}\left(1-|z|^2\right)^{2-\frac{1}{m}}}=1,\tag{6.7}$$

which implies that the left-hand side of (6.7) is independent of the value of z and w. When we approach the boundary, namely $|z|^2 + |w|^{2m} \to 1$, we note that the numerator of (6.7) does not always vanish, whereas the denominator always vanishes if $\lambda \neq 3$. That forces $\lambda = 3$. Let $(z,w) \to (0,1) \in \partial \Omega_m$, we can verify that m=1.

As a significant example, the following computation verifies the recent work of Savale-Xiao [SX23] regarding Yau's question on bounded pseudoconvex domains with C^{∞} -boundary of finite type in dimension two.

Proposition 6.4. The Bergman metric g_B is Kähler-Einstein if and only if m=1.

Proof. The "if" part follows from the case of the ball. For the "only if" part, we use the same notations as in [AS83]. It is enough to show this at (0, w) with |w| < 1. Let

$$t = \frac{1 - |w|^2}{1 - r|w|^2}, \qquad |w| < 1,$$

where r = (m-1)/(m+1). We note that $0 \le r < 1$ and $0 < t \le 1$. Then the Bergman metric is given by

$$g(0,w) = \begin{pmatrix} \alpha/(1+r)t & 0\\ 0 & \beta(1-rt)^2/(1-r)^2t^2 \end{pmatrix}$$

and the Ricci curvature of the Bergman metric is given by

$$\operatorname{Ric}(0,w) = \begin{pmatrix} \frac{3\alpha^2\beta - 4A\beta - 2B\alpha}{\alpha\beta(1+r)t} & 0\\ 0 & \frac{(1-rt)^2(3\alpha\beta^2 - 4C\alpha - 2B\beta)}{\alpha\beta(1-r)^2t^2} \end{pmatrix},$$

where

$$\begin{cases}
\alpha = 3 + rt^{2}, & \beta = 3 - rt^{2}, \\
A = 6 + 4rt^{2} + (1 + r)rt^{3}, \\
B = 2(9 + 3rt^{2} - 3(1 + r)rt^{3} + 2r^{2}t^{4}) / (3 + rt^{2}), \\
C = 3(6 - 6rt^{2} + (1 + r)rt^{3}) / (3 - rt^{2}).
\end{cases}$$
(6.8)

Assume that the Bergman metric is Kähler-Einstein, that is

$$Ric(0, w) = -q(0, w).$$

Consider the $(1, \bar{1})$ -component of the Ricci curvature, after rearrangement we obtain

$$4\alpha^2\beta - 4A\beta - 2B\alpha = 0. ag{6.9}$$

Plugging (6.8) into (6.9), we get

$$r^{2}[rt^{2} - (1+r)t + 1] = 0.$$

Since $0 \le r < 1$, we have $rt^2 - (1+r)t + 1 > 0$. This implies r = 0.

It was proved in [Gon19] that the holomorphic bisectional curvature of g_B is negatively pinched in Ω_m . This implies that $\text{Ric}(g_B)$ is also negatively pinched. Thus, we can characterize the Thullen domain if g_m or g_B is a Kähler-Ricci soliton.

Theorem 6.5. Let Ω_m be the bounded Thullen domain in \mathbb{C}^2 . Then the following statements are equivalent.

- (1) g_m (or g_B) is Kähler-Einstein;
- (2) g_m (or g_B) is a Kähler-Ricci soliton;
- (3) there is a constant $\lambda > 0$ such that $g_m = \lambda g_B$;
- (4) m = 1 and Ω_m is the unit ball.
- 6.2. Hartogs domains. Let $C \in (0, +\infty)$ and let $F : [0, B) \to (0, +\infty)$ be a continuous decreasing function that is smooth on (0, B) satisfying

$$\frac{d}{dx}\left(\frac{xF'(x)}{F(x)}\right) < 0.$$

The bounded Hartogs domain $D_F \subset \mathbb{C}^n$ associated with the function F is a strictly pseudoconvex domain given by

$$D_F = \left\{ \left(z_0, z_1, \dots, z_{n-1} \right) \in \mathbb{C}^n; \left| z_0 \right|^2 < x_0, \left| z_1 \right|^2 + \dots + \left| z_{n-1} \right|^2 < F\left(\left| z_0 \right|^2 \right) \right\}.$$

There is a corresponding complete Kähler metric defined by

$$g_F = -\partial_i \partial_{\bar{j}} \log \left(F \left(|z_0|^2 \right) - |z_1|^2 - \dots - |z_{n-1}|^2 \right).$$

It was computed in [LZ10] that the Ricci curvature of g_F is negatively pinched near the boundary. Consequently, we have the following:

Corollary 6.6 (Theorem 1.2, [LZ10]). Let $D_F \subset \mathbb{C}^n$ be a bounded strictly pseudoconvex Hartogs domain equipped with a complete Kähler metric g_f defined above. Suppose g_F is a Kähler-Ricci soliton. Then g_F is Kähler-Einstein. Moreover, $F(x) = C_1 - C_2 x$ for some $C_1, C_2 > 0$, which implies that D_F is holomorphically isomorphic to an open subset of the complex hyperbolic space \mathbb{H}^n via the map

$$\varphi: D_F \to \mathbb{H}^n, \quad (z_0, z_1, \dots, z_{n-1}) \mapsto \left(\frac{z_0}{\sqrt{c_1/c_2}}, \frac{z_1}{\sqrt{c_1}}, \dots, \frac{z_{n-1}}{\sqrt{c_1}}\right).$$

REFERENCES

- [AMR16] L. J. Alías, P. Mastrolia, and M. Rigoli, *Maximum principles and geometric applications*, vol. 700, Springer, 2016.
- [AS83] K. Azukawa and M. Suzuki, The Bergman metric on a Thullen domain, Nagoya Math. J. 89 (1983), 1–11.
- [BS51] S. Bergman and M. Schiffer, *Kernel functions and conformal mapping*, Compositio Mathematica **8** (1951), 205–249.
- [Cao96] H.-D. Cao, *Existence of gradient Kähler-Ricci solitons*, Elliptic and parabolic methods in geometry (Minneapolis, MN, 1994) **1** (1996), 16.
- [Cao97] _____, Limits of solutions to the Kähler-Ricci flow, J. Diff. Geom. 45 (1997), no. 2, 257–272.
- [Che79] S.-Y. Cheng, Open problems, Conference on nonlinear problems in geometry held in Katata, September 3–8.
- [Cho23] B. Chow, Ricci solitons in low dimensions, vol. 235, American Mathematical Society, 2023.
- [CY80] S.-Y. Cheng and S.-T. Yau, On the existence of a complete Kähler metric on non-compact complex manifolds and the regularity of Fefferman's equation, Comm. Pure Appl. Math 33 (1980), 507–544.
- [Dem12] J.-P. Demailly, *Complex analytic and differential geometry*, https://www-fourier.ujf-grenoble.fr/~dema 2012.
- [DGZ12] F. Deng, Q. Guan, and L. Zhang, *Some properties of squeezing functions on bounded domains*, Pac. J. Math. **257** (2012), no. 2, 319–341.
- [DGZ16] ______, Properties of squeezing functions and global transformations of bounded domains, Trans. Amer. Math. Soc. **368** (2016), no. 4, 2679–2696.
- [Die 70] K. Diederich, Das randverhalten der Bergmanschen kernfunktion und metrik in streng pseudo-konvexen gebieten, Math. Ann. **187** (1970), no. 1, 9–36.
- [FIK03] M. Feldman, T. Ilmanen, and D. Knopf, *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*, J. Diff. Geom. **65** (2003), no. 2, 169–209.
- [Fut83] A. Futaki, An obstruction to the existence of Einstein Kähler metrics, Invent. Math. 73 (1983), no. 3, 437–443.
- [FW97] S. Fu and B. Wong, On strictly pseudoconvex domains with Kähler-Einstein Bergman metrics, Math. Res. Lett. 4 (1997), no. 5, 697–703.
- [FW18] J. Fornæss and E. Wold, A non-strictly pseudoconvex domain for which the squeezing function tends to 1 towards the boundary, Pacific J. Math. 297 (2018), no. 1, 79–86.
- [Gon19] S. Gontard, Curvatures of metrics in non-compact complex manifolds, Université Grenoble Alpes, 2019.
- [GKK11] R. E. Greene, K.-T. Kim, and S. G. Krantz, *The geometry of complex domains*, vol. 291, Springer Science & Business Media, 2011.
- [Ham88] R. Hamilton, *The Ricci flow on surfaces, Mathematics and general relativity*, Contemp. Math. **71** (1988), 237–261.
- [HX21] X. Huang and M. Xiao, *Bergman-Einstein metrics, a generalization of Kerner's theorem and Stein spaces with spherical boundaries*, J. Reine Angew. Math. **2021** (2021), no. 770, 183–203.
- [Kra01] S. Krantz, Function theory of several complex variables, vol. 340, American Mathematical Soc., 2001.
- [KY96] S. G. Krantz and J. Yu, *On the Bergman invariant and curvatures of the Bergman metric*, Illinois J. Math. **40** (1996), no. 2, 226–244.
- [LSY05] K. Liu, X. Sun, and S.T. Yau, Canonical metrics on the moduli space of Riemann surfaces ii, J. Differential Geom. **69** (2005), no. 1, 163–216.
- [LZ10] A. Loi and F. Zuddas, Canonical metrics on Hartogs domains, Osaka J. Math. 47 (2010), no. 2, 507–521.
- [MY83] N. Mok and S.-T. Yau, Completeness of the Kähler-Einstein metric on bound domains and the characterization of domain of holomorphy by curvature condition, Proceedings of Symposia in Pure mathematics, V.39 Part 1 (1983), 41—59.
- [Ohs80] T. Ohsawa, On complete Kähler domains with C^1 -boundary, Publ. Res. Inst. Math. Sci. **16** (1980), no. 3, 929–940.

- [Omo67] H. Omori, *Isometric immersions of Riemannian manifolds*, J. Math. Soc. Japan **19** (1967), no. 2, 205–214.
- [Per02] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv preprint math/0211159 (2002).
- [Seo12] A. Seo, *On a theorem of Paul Yang on negatively pinched bisectional curvature*, Pac. J. Math. **256** (2012), no. 1, 201–209.
- [SX23] N. Savale and M. Xiao, Kähler-Einstein Bergman metrics on pseudoconvex domains of dimension two, arXiv preprint arXiv:2309.10595 (2023).
- [SY10] R.-M. Schoen and S.-T. Yau, Lectures on differential geometry, International Press, 2010.
- [Yau75] S.-T. Yau, *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math. **28** (1975), no. 2, 201–228.
- [Yeu09] S.-K. Yeung, *Geometry of domains with the uniform squeezing property*, Adv. Math. **221** (2009), no. 2, 547–569.
- [Zha15] L. Zhang, *On curvature estimates of bounded domains*, Complex Analysis and Geometry, Springer Proceedings in Mathematics & statistics,, vol. 144, 2015, pp. 353–367.
- [Zim18] A. Zimmer, A gap theorem for the complex geometry of convex domains, Trans. Amer. Math. Soc. **370** (2018), no. 10, 7489–7509.

ZEHAO SHA, INSTITUT FOURIER, UNIVERSITÉ GRENOBLE ALPES, 100 RUE DES MATHÉMATIQUES, 38610 GIÈRES, FRANCE.

Email address: zehao.sha@univ-grenoble-alpes.fr