

On intermediate levels of nested occupancy scheme in random environment generated by stick-breaking: the case of heavy tails

Oksana Braganets* and Alexander Iksanov†

December 5, 2024

Abstract

We investigate a nested balls-in-boxes scheme in a random environment. The boxes follow a nested hierarchy, with infinitely many boxes in each level, and the hitting probabilities of boxes are random and obtained by iterated fragmentation of a unit mass. The hitting probabilities of the first-level boxes are given by a stick-breaking model $P_k = W_1 W_2 \cdot \dots \cdot W_{k-1} (1 - W_k)$ for $k \in \mathbb{N}$, where W_1, W_2, \dots are independent copies of a random variable W taking values in $(0, 1)$. The infinite balls-in-boxes scheme in the first level is known as a Bernoulli sieve. We assume that the mean of $|\log W|$ is infinite and the distribution tail of $|\log W|$ is regularly varying at ∞ . Denote by $K_n(j)$ the number of occupied boxes in the j th level provided that there are n balls and call the level j intermediate, if $j = j_n \rightarrow \infty$ and $j_n = o((\log n)^a)$ as $n \rightarrow \infty$ for appropriate $a > 0$. We prove that, for some intermediate levels j , finite-dimensional distributions of the process $(K_n(\lfloor j_n u \rfloor))_{u>0}$, properly normalized, converge weakly as $n \rightarrow \infty$ to those of a pathwise Lebesgue-Stieltjes integral, with the integrand being an exponential function and the integrator being an inverse stable subordinator. The present paper continues the line of investigation initiated in the articles Buraczewski, Dvogay and Iksanov (2020) and Iksanov, Marynych and Samoilenko (2022) in which the random variable $|\log W|$ has a finite second moment, and Iksanov, Marynych and Rashytov (2022) in which $|\log W|$ has a finite mean and an infinite second moment.

Key words: Bernoulli sieve; infinite occupancy scheme; perturbed random walk; random environment; weak convergence of finite-dimensional distributions; weighted branching process.

2020 Mathematics Subject Classification: 60F05, 60J80.

*Faculty of Computer Science and Cybernetics, Taras Shevchenko National University of Kyiv, Ukraine; e-mail address: oksanabraganets@knu.ua

†Faculty of Computer Science and Cybernetics, Taras Shevchenko National University of Kyiv, Ukraine; e-mail address: iksan@univ.kiev.ua

1 Introduction

1.1 Definition of the model

Let P_1, P_2, \dots be nonnegative random variables with an arbitrary joint distribution satisfying $\sum_{k \geq 1} P_k = 1$ almost surely (a.s.). The sequence $(P_k)_{k \geq 1}$ is interpreted as a *random environment*. An *occupancy scheme in random environment* is defined as follows. Conditionally on $(P_k)_{k \geq 1}$, balls are allocated independently over an infinite array of boxes $1, 2, \dots$ with probability P_k of hitting box k . The occupancy scheme in random environment is called *infinite*, if the number of positive probabilities is infinite a.s. One of the most popular infinite occupancy schemes in random environment is called *Bernoulli sieve*. It corresponds to $(P_k)_{k \geq 1}$ which follow a stick-breaking model

$$P_k = W_1 W_2 \cdots W_{k-1} (1 - W_k), \quad k \in \mathbb{N} := \{1, 2, \dots\}, \quad (1)$$

where W_1, W_2, \dots are independent copies of a random variable W taking values in $(0, 1)$. The Bernoulli sieve was introduced in [14] and further investigated in many articles. Surveys of earlier works can be found in [13, 17]. An incomplete list of more recent contributions includes [1, 11, 18, 25, 26, 27].

A *nested infinite occupancy scheme in random environment* is a hierarchical generalization of the infinite occupancy scheme in random environment which is defined by settling, in a consistent way, the sequence of the occupancy schemes in random environment on the tree of a deterministic weighted branching process with positive weights (a multiplicative counterpart of a branching random walk). The scheme was introduced in [3] and further investigated in [8, 9, 12, 20, 22]. Following [3, 12], we now provide the definition of the nested occupancy scheme in random environment. To this end, some preparations are needed. Let $\mathbb{V} := \bigcup_{n \geq 0} \mathbb{N}^n$ be the set of all possible individuals of some population. The initial ancestor is identified with the empty word \emptyset and its weight is $P(\emptyset) = 1$. On some probability space, let $((P_k(v))_{k \geq 1})_{v \in \mathbb{V}}$ denote a family of independent copies of $(P_k)_{k \geq 1}$. An individual $v = v_1 \dots v_j$ in the j th level (generation) whose weight is denoted by $P(v)$ produces an infinite number of offspring residing in the $(j + 1)$ st level. The offspring of the individual v are enumerated by $vk = v_1 \dots v_j k$, where $k \in \mathbb{N}$, and the weights of the offspring are denoted by $P(vk)$. It is postulated that $P(vk) = P(v)P_k(v)$. Note that, for each $j \in \mathbb{N}$, $\sum_{|v|=j} P(v) = 1$ a.s., where, by convention, $\sum_{|v|=j}$ means that the sum is taken over all individuals in the j th level.

We are ready to explain the construction of the nested occupancy scheme in random environment. We identify individuals with boxes, so that the random weights (probabilities) of boxes in the subsequent levels are formed by the vectors $(P(v))_{|v|=1} = (P_k)_{k \geq 1}$, $(P(v))_{|v|=2}$ and so on. The collection of balls is the same for all levels. The balls are allocated, conditionally on $(P(v))_{v \in \mathbb{V}}$, according to the following rule. At time 0, infinitely many balls are collected in the box \emptyset . At the time $n \in \mathbb{N}$, one of the balls leaves the root \emptyset and falls, independently of the $(n - 1)$ balls that have left the root earlier, into the box v in the first level with probability $P(v)$. Simultaneously, it falls into the box vi_1 in the second level with probability $P(vi_1)/P(v)$, into the box vi_1i_2 in the third level with probability $P(vi_1i_2)/P(vi_1)$ and so on, indefinitely. At time n , that is, when n balls have already spread over the tree, a box is deemed occupied provided it was hit by some ball (out of n) on its way over the levels.

In what follows, we assume that the probabilities $(P_k)_{k \geq 1}$ are given by (1) and that these and the outcome of throwing balls are defined on a common probability space. Let $K_n(j)$ be the number of occupied boxes at time n in the j th level. Assuming that the mean of $|\log W|$ is infinite and the distribution tail of $|\log W|$ is regularly varying at ∞ , we investigate the distributional behavior of $K_n(j)$ when $j = j_n \rightarrow \infty$ and $j_n = o((\log n)^a)$ as $n \rightarrow \infty$ for appropriate $a > 0$. We call *intermediate* the levels j satisfying the latter two properties. The analogous problem was studied in [9, 20] in the situation in which the variable $|\log W|$ has a finite second moment, and in [19] in the situation in which $|\log W|$ has a finite mean and an infinite second moment.

1.2 Our assumptions on the distribution of W

Our main result, Theorem 1, will be stated under certain assumptions on the distribution of $(|\log W|, |\log(1 - W)|)$. We think it is instructive to introduce the assumptions for an arbitrary random vector (ξ, η) and then specialize for $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$.

Let $(\xi_k, \eta_k)_{k \geq 1}$ be independent copies of a random vector (ξ, η) with positive arbitrary dependent components. Let F and G be the distribution functions of ξ and η , respectively. Denote by $(S_k)_{k \geq 0}$ the zero-delayed standard random walk with increments ξ_k , that is, $S_0 := 0$ and $S_k := \xi_1 + \dots + \xi_k$ for $k \in \mathbb{N}$. Put

$$T_k := S_{k-1} + \eta_k, \quad k \in \mathbb{N}.$$

The random sequence $T = (T_k)_{k \geq 1}$ is known in the literature as a (globally) perturbed random walk, see [17] for a survey and [2, 6, 7, 8, 21] for more recent contributions. A connection with the occupancy scheme treated in the paper is justified by the fact that if $(P_k)_{k \geq 1}$ is given by (1), then $(-\log P_k)_{k \geq 1}$ is a perturbed random walk with $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$.

Put $N(t) := \sum_{k \geq 1} \mathbb{1}_{\{T_k \leq t\}}$ and $V(t) := \mathbb{E}N(t)$ for $t \geq 0$. Plainly,

$$V(t) = \int_{[0, t]} G(t - y) dU(y), \quad t \geq 0, \quad (2)$$

where, for $t \geq 0$, $U(t) = \sum_{i \geq 0} \mathbb{P}\{S_i \leq t\}$ is the renewal function.

Assume that

$$\mathbb{P}\{\xi > t\} \sim ct^{-\alpha}, \quad t \rightarrow \infty \quad (3)$$

for some $c > 0$ and $\alpha \in (0, 1)$. A standard result of renewal theory states that the latter limit relation is equivalent to

$$U(t) \sim Ct^\alpha, \quad t \rightarrow \infty,$$

where $C = C_\alpha := (c\Gamma(1 + \alpha)\Gamma(1 - \alpha))^{-1}$ and Γ is the Euler gamma-function. To prove this, one uses the equality $\int_{[0, \infty)} e^{-st} dU(t) = (1 - \varphi(s))^{-1}$ for $s > 0$, where $\varphi(s) := \mathbb{E}e^{-s\xi}$ for $s \geq 0$, and Karamata's Tauberian theorem (Theorem 1.7.1 on p. 37 in [5]). Since $\int_{[0, \infty)} e^{-st} dV(t) = \mathbb{E}e^{-s\eta}(1 - \mathbb{E}e^{-s\xi})^{-1}$ for $s > 0$, the same reasoning enables us to conclude that (3) is equivalent to $V(t) \sim Ct^\alpha$ as $t \rightarrow \infty$. It will become clear soon that the first-order behavior of V is not sufficient for the purposes of the present work. A two-term expansion

of V is needed. It will be shown in Section 3 that the following two assumptions ensure the required expansion of V .

ASSUMPTION *A*: $U(t) = Ct^\alpha + O(t^\theta)$ as $t \rightarrow \infty$ for some $\rho \in [0, \alpha)$.

ASSUMPTION *B*: Either $\limsup_{t \rightarrow \infty} (1 - G(t))/(1 - F(t)) = m_0 \in [0, \infty)$ or $m_0 = \infty$ and that $\limsup_{t \rightarrow \infty} t^\theta (1 - G(t)) = m_1 \in [0, \infty)$ necessarily for some $\theta \in (0, \alpha)$.

It seems to be a non-trivial problem to provide necessary and sufficient conditions for Assumption *A* formulated in terms of the distribution of ξ . In the case where the distribution of ξ is absolutely continuous (with respect to Lebesgue measure) sufficient conditions in terms of the densities considered as functions of a complex argument can be found in [28]. In Section 3 we give sufficient conditions of a different flavor.

Observe that when applied to the vector $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$ Assumptions *A* and *B* regulate the behavior of W near 0 and near 1, respectively. If (a) $\mathbb{E}e^{-s|\log W|} = \exp(-c\Gamma(1-\alpha)s^\alpha)$ for $s \geq 0$, that is, the distribution of $|\log W|$ is α -stable or (b) $\mathbb{E}e^{-s|\log W|} = (1 + (c/\kappa)\Gamma(1-\alpha)s^\alpha)^{-\kappa}$ for $s \geq 0$ and some $\kappa > 0$, then Assumption *A* holds with $\rho = 0$ and Assumption *B* holds with $m_0 = 0$. This statement will be proved in Section 3.

2 Main result

To formulate our main result, we recall the definition of an *inverse stable subordinator*. Let $(\mathcal{Z}_\alpha(u))_{u \geq 0}$ be an α -stable subordinator with $-\log \mathbb{E}e^{-s\mathcal{Z}_\alpha(1)} = \Gamma(1-\alpha)s^\alpha$ for $s \geq 0$. The process $\mathcal{Z}_\alpha^\leftarrow = (\mathcal{Z}_\alpha^\leftarrow(u))_{u \geq 0}$ defined by

$$\mathcal{Z}_\alpha^\leftarrow(u) := \inf\{v \geq 0 : \mathcal{Z}_\alpha(v) > u\}, \quad u \geq 0$$

is called *inverse α -stable subordinator*.

We write $\xrightarrow{\text{f.d.d.}}$ to denote weak convergence of finite-dimensional distributions. Also, $[x]$ denotes the integer part of $x \in \mathbb{R}$.

Theorem 1. *Under Assumptions *A* and *B* imposed on $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$, let $(j_n)_{n \geq 1}$ be a sequence of positive numbers satisfying $\lim_{n \rightarrow \infty} j_n = \infty$ and*

$$j_n = o\left((\log n)^{\min(1/3, (\alpha-\beta)/(\alpha-\beta+1))}\right), \quad n \rightarrow \infty,$$

where $\beta = \rho$ if $m_0 < \infty$ and $\beta = \max(\rho, \alpha - \theta)$ if $m_0 = \infty$ and $m_1 < \infty$. Then

$$\left(\frac{c j_n^\alpha}{\rho_{[j_n u]-1} (\log n)^{\alpha [j_n u]}} K_n([j_n u]) \right)_{u>0} \xrightarrow{\text{f.d.d.}} \left(\int_0^\infty e^{-\alpha u y} d\mathcal{Z}_\alpha^\leftarrow(y) \right)_{u>0}, \quad n \rightarrow \infty,$$

where

$$\rho_i := \frac{(C\Gamma(\alpha+1))^i}{\Gamma(\alpha i+1)}, \quad i \in \mathbb{N}_0 := \{0, 1, \dots\}, \quad (4)$$

and Γ is the Euler gamma-function.

Now we discuss the structure of the limit process in Theorem 1. It can be seen from the proof that the integrator $\mathcal{Z}_\alpha^\leftarrow$ describes the fluctuations of the number of occupied boxes in the first level, properly normalized. The process $\mathcal{Z}_\alpha^\leftarrow$ is a.s. continuous singular

with respect to Lebesgue measure. The integrand $y \mapsto e^{-\alpha y}$ which is formed by the renewal structure of the underlying tree makes the process $\left(\int_0^\infty e^{-\alpha y} d\mathcal{Z}_\alpha^-(y)\right)_{u>0}$ a.s. infinitely differentiable. Thus, when passing from the first level to intermediate levels the fluctuations of the number of occupied boxes smooth out.

Next, we intend to set a non-rigorous link between Theorem 1 and a limit theorem for $K_n(j)$ with fixed j . Assume that $\mathbb{P}\{|\log W| > x\} \sim x^{-\alpha} \ell(x)$ as $x \rightarrow \infty$ for some $\alpha \in (0, 1)$ and some ℓ slowly varying at ∞ . A specialization of a functional limit theorem given in Theorem 7 of [8] reads

$$\left(\frac{(\ell(\log n))^j K_n(j)}{(\log n)^{\alpha j}}\right)_{j \geq 1} \xrightarrow{d} \left(\int_{[0,1]} (1-y)^{\alpha(j-1)} d\mathcal{Z}_\alpha^-(y)\right)_{j \geq 1}, \quad n \rightarrow \infty,$$

where \xrightarrow{d} denotes convergence in distribution. Observe that $(\mathcal{Z}_\alpha^-(y/j))_{y \geq 0}$ has the same distribution as $(j^{-\alpha} \mathcal{Z}_\alpha^-(y))_{y \geq 0}$ and that, with $y > 0$ fixed, $\lim_{j \rightarrow \infty} (1-y/j)^{\alpha(j-1)} = e^{-\alpha y}$. Hence,

$$j^\alpha \int_{[0,1]} (1-y)^{\alpha(j-1)} d\mathcal{Z}_\alpha^-(y) = j^\alpha \int_{[0,j]} (1-y/j)^{\alpha(j-1)} d\mathcal{Z}_\alpha^-(y/j) \stackrel{d}{=} \int_{[0,j]} (1-y/j)^{\alpha(j-1)} d\mathcal{Z}_\alpha^-(y),$$

where $\stackrel{d}{=}$ denotes equality of distributions. It is reasonable to expect that the distributional limit of the last integral as $j \rightarrow \infty$ is $\int_{[0,\infty)} e^{-\alpha y} d\mathcal{Z}_\alpha^-(y)$. This is in line with Theorem 1.

The remainder of the article is structured as follows. As has already been announced, we discuss in Section 3 Assumptions *A* and *B* and their consequences. Section 4 contains auxiliary results on a general branching process generated by a perturbed random walk. The proof of Theorem 1 is given in Section 5.

3 Discussion of Assumptions *A* and *B*

We start by showing that Assumptions *A* and *B* secure a two-term expansion of V .

Lemma 1. *Assume that Assumptions *A* and *B* hold. Then there exists $D > 0$ such that*

$$|V(t) - Ct^\alpha| \leq Dt^\beta, \quad t \geq 0, \tag{5}$$

where $\beta = \rho$ if $m_0 < \infty$, $\beta = \max(\rho, \alpha - \theta)$ if $m_0 = \infty$ and $m_1 < \infty$; $C > 0$, $\alpha \in (0, 1)$ and $\rho \in [0, \alpha)$ are as defined in Assumption *A*; m_0 , m_1 and $\theta \in (0, \alpha)$ are as defined in Assumption *B*.

Proof. Assuming that $\limsup_{t \rightarrow \infty} (1 - G(t))/(1 - F(t)) = m_0 \in [0, \infty)$ we shall show that

$$\int_{[0,t]} (1 - G(t-y)) dU(y) = O(1), \quad t \rightarrow \infty. \tag{6}$$

Indeed, given $m_2 > m_0$ there exists $t_0 > 0$ such that $1 - G(t) \leq m_2(1 - F(t))$ whenever $t \geq t_0$. With this at hand,

$$\int_{[0,t-t_0]} (1 - G(t-y)) dU(y) \leq m_2 \int_{[0,t]} (1 - F(t-y)) dU(y) = m_2.$$

The last equality is another form of writing the renewal equation for U . Alternatively, it is equivalent to $\mathbb{P}\{\inf\{k \geq 0 : S_k > t\} < \infty\} = 1$. By subadditivity of U (see, for instance, Theorem 1.7 on p. 10 in [23]),

$$\int_{(t-t_0, t]} (1 - G(t-y)) dU(y) \leq U(t) - U(t-t_0) \leq U(t_0),$$

and (6) follows. Equality (2) is equivalent to

$$V(t) = U(t) - \int_{[0, t]} (1 - G(t-y)) dU(y), \quad t \geq 0.$$

Hence, Assumption A entails $V(t) = Ct^\alpha + O(t^\rho)$ as $t \rightarrow \infty$. Since $V(0) = 0$, we can find $D > 0$ such that $|V(t) - Ct^\alpha| \leq Dt^\rho$ for all $t \geq 0$.

Assume now that $m_0 = \infty$ and that $\limsup_{t \rightarrow \infty} t^\theta(1 - G(t)) = m_1 \in [0, \infty)$ for some $\theta \in (0, \alpha)$. We shall prove that

$$\int_{[0, t]} (1 - G(t-y)) dU(y) = O(t^{\alpha-\theta}), \quad t \rightarrow \infty. \quad (7)$$

Given $m_3 > m_1$ and given $C^* > C$ there exists t_1 such that $U(t) \leq C^*t^\alpha$ and $1 - G(t) \leq m_3t^{-\theta}$ whenever $t \geq t_1$. We already know from the previous paragraph that subadditivity of U ensures $\int_{(t-t_1, t]} (1 - G(t-y)) dU(y) = O(1)$ as $t \rightarrow \infty$. Further, we obtain with the help of integration by parts,

$$\begin{aligned} \int_{[0, t-t_1]} (1 - G(t-y)) dU(y) &\leq m_3 \int_{[0, t-t_1]} (t-y)^{-\theta} dU(y) = \int_{[t_1, t]} y^{-\theta} d(U(t) - U(t-y)) \\ &= t^{-\theta}(U(t) - U(0)) - t_1^{-\theta}(U(t) - U(t-t_1)) + \theta \int_{t_1}^t (U(t) - U(t-y)) y^{-\theta-1} dy. \end{aligned}$$

The first term is $O(t^{\alpha-\theta})$ and the second term is $O(1)$. Another appeal to subadditivity of U yields

$$\int_{t_1}^t (U(t) - U(t-y)) y^{-\theta-1} dy \leq \int_{t_1}^t U(y) y^{-\theta-1} dy \leq C^* \int_{t_1}^t y^{\alpha-\theta-1} dy = O(t^{\alpha-\theta}),$$

and (7) follows. Mimicking the argument from the first part of the proof we arrive at (5) with $\beta = \max(\rho, \alpha - \theta)$. \square

Let $(\mathcal{Z}_\alpha(u))_{u \geq 0}$ be an α -stable subordinator as defined at the beginning of Section 2. Now we give a sufficient condition for Assumption A which works for a special class of the distributions of ξ . Namely, assume that the distribution of ξ coincides with that of $\mathcal{Z}_\alpha(c)X^{1/\alpha}$, where X is a positive random variable of mean one which is independent of $\mathcal{Z}_\alpha(c)$ for some $c > 0$. Equivalently, this assumption means that the standard random walk $(S_n)_{n \geq 0}$ has the same distribution as $(\mathcal{Z}_\alpha(c\hat{S}_n))_{n \geq 0}$. Here, $\hat{S}_0 := 0$, $\hat{S}_n := X_1 + \dots + X_n$ for $n \in \mathbb{N}$, X_1, X_2, \dots are independent copies of X , and $(\hat{S}_n)_{n \geq 0}$ is independent of $(\mathcal{Z}_\alpha(u))_{u \geq 0}$.

Put $\hat{U}(t) := \sum_{n \geq 0} \mathbb{P}\{\hat{S}_n \leq t\}$ for $t \geq 0$. Now we obtain a formula which connects U and \hat{U} . To this end, we need the following property: for fixed $t > 0$, $\mathcal{Z}_\alpha(ct)$ has the same

distribution as $t^{1/\alpha} \mathcal{Z}_\alpha(c)$. As a consequence,

$$\begin{aligned} U(t) &= \sum_{n \geq 0} \mathbb{P}\{S_n \leq t\} = \sum_{n \geq 0} \mathbb{P}\{\mathcal{Z}_\alpha(c \hat{S}_n) \leq t\} = \sum_{n \geq 0} \mathbb{P}\{\hat{S}_n^{1/\alpha} \mathcal{Z}_\alpha(c) \leq t\} \\ &= \sum_{n \geq 0} \mathbb{P}\{\hat{S}_n \leq (\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha\} = \mathbb{E} \hat{U}((\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha), \quad t \geq 0. \end{aligned} \quad (8)$$

Let $d > 0$. Recall that the distribution of a positive random variable is called *d-arithmetic* if it is concentrated on the lattice $(nd)_{n \geq 1}$ and not concentrated on $(nd_1)_{n \geq 1}$ for any $d_1 > d$. The distribution of Y is called *nonarithmetic* if it is not *d-arithmetic* for all $d > 0$.

Invoking (8) enables us to provide sufficient conditions which ensure Assumption A.

Lemma 2. *Assume that $\mathbb{E}X^\vartheta < \infty$ for some $\vartheta \in (1, 2]$. Then*

$$U(t) = Ct^\alpha + O(t^{\alpha(2-\vartheta)}), \quad t \rightarrow \infty,$$

that is, Assumption A holds with $\rho = \alpha(2-\vartheta)$. Here, as before, $C = (c\Gamma(1+\alpha)\Gamma(1-\alpha))^{-1}$.

Proof. We intend to use Lorden's inequality for \hat{U} . Although, in many sources it is only stated for nonarithmetic distributions, it holds and takes the same form for both nonarithmetic and arithmetic distributions, see [10] for an elegant proof.

If $\mathbb{E}X^2 < \infty$, then, by Lorden's inequality, $\hat{U}(t) \leq t + \mathbb{E}X^2$ for all $t \geq 0$ (recall that $\mathbb{E}X = 1$). Hence,

$$U(t) = \mathbb{E} \hat{U}((\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha) \leq \mathbb{E}(\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha + \mathbb{E}X^2.$$

It remains to note that, by Lemma 9,

$$\mathbb{E}(\mathcal{Z}_\alpha(c))^{-\alpha} = \frac{\alpha}{\Gamma(1+\alpha)} \int_0^\infty s^{\alpha-1} e^{-c\Gamma(1-\alpha)s^\alpha} ds = C.$$

If $\mathbb{E}X^\vartheta < \infty$ for some $\vartheta \in (1, 2)$, then $\hat{U}(t) = t + O(t^{2-\vartheta})$ as $t \rightarrow \infty$, see, for instance, formula (3.13) on p. 433 in [7]. In particular, for some $t_0 > 0$ and a constant $C^* > 0$ $\hat{U}(t) \leq t + C^*t^{2-\vartheta}$ whenever $t \geq t_0$. Using (8) yields

$$\begin{aligned} U(t) &\leq \mathbb{E}((\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha + C^*(\mathcal{Z}_\alpha(c))^{-\alpha(2-\vartheta)} t^{\alpha(2-\vartheta)}) \mathbb{1}_{\{(\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha \geq t_0\}} \\ &\quad + \mathbb{E} \hat{U}((\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha) \mathbb{1}_{\{(\mathcal{Z}_\alpha(c))^{-\alpha} t^\alpha < t_0\}} \leq Ct^\alpha + C^* \mathbb{E}(\mathcal{Z}_\alpha(c))^{-\alpha(2-\vartheta)} t^{\alpha(2-\vartheta)} + \hat{U}(t_0) \\ &= Ct^\alpha + O(t^{\alpha(2-\vartheta)}), \quad t \rightarrow \infty. \end{aligned}$$

□

Next, we show that Assumptions A and B hold true, with $(\xi, \eta) = (|\log W|, |\log(1-W)|)$, if, for instance, (a) $|\log W| = \mathcal{Z}_\alpha(c)$ for some $c > 0$, that is, the distribution of $|\log W|$ is α -stable; (b) $\varphi(s) := \mathbb{E}e^{-s\xi} = (1 + (c/\kappa)\Gamma(1-\alpha)s^\alpha)^{-\kappa}$ for $s \geq 0$, some $\alpha \in (0, 1)$ and some $\kappa > 0$. In both cases, $|\log W|$ has the same distribution as $\mathcal{Z}_\alpha(c)X^{1/\alpha}$, where $\mathbb{P}\{X = 1\} = 1$ in the case (a) and X is independent of $\mathcal{Z}_\alpha(c)$ and has a gamma distribution with parameters κ and κ in the case (b). To justify the latter claim, observe

that $s \mapsto \varphi(((c\Gamma(1-\alpha))^{-1}s)^{1/\alpha}) = (1+s/\kappa)^{-\kappa}$ is the Laplace-Stieltjes transform of the aforementioned gamma distribution. Thus, in both cases $\mathbb{E}X^2 < \infty$ and, according to Lemma 2, Assumption *A* holds with $\rho = 0$.

Now, temporarily ignoring the cases (a) and (b), we show that Assumption *B* holds true, with $m_0 = 0$, whenever $1-F(x) = \mathbb{P}\{|\log W| > x\} \sim cx^{-\alpha}$ as $x \rightarrow \infty$ and $\mathbb{E}|\log W|^{-\gamma} < \infty$ for some $\gamma > 0$. Indeed, by Markov's inequality,

$$\begin{aligned} 1 - G(x) &= \mathbb{P}\{|\log(1 - W)| > x\} = \mathbb{P}\{|\log W| < |\log(1 - e^{-x})|\} \\ &= \mathbb{P}\{|\log W|^{-\gamma} > |\log(1 - e^{-x})|^{-\gamma}\} \leq \mathbb{E}|\log W|^{-\gamma} (-\log(1 - e^{-x}))^\gamma \sim \mathbb{E}|\log W|^{-\gamma} e^{-\gamma x} \end{aligned}$$

as $x \rightarrow \infty$, and the claim follows.

We already know that $\mathbb{E}(\mathcal{Z}_\alpha(c))^{-\alpha} < \infty$. Hence, Assumption *B* holds, with $m_0 = 0$, in the case (a). Although we do not need such a precision, we note in passing that in the case (a), with c and α satisfying $c\Gamma(1-\alpha) = 1$, by Lemma 1 in [15], the following asymptotic relation holds

$$\mathbb{P}\{|\log(1 - W)| > x\} \sim c_1 e^{-\frac{\alpha x}{2(1-\alpha)}} \exp(-c_2 |\log(1 - e^{-x})|^{-\frac{\alpha}{1-\alpha}}), \quad x \rightarrow \infty$$

for explicitly known positive constants c_1 and c_2 .

In the case (b), using Lemma 9 we infer $\mathbb{E}X^{-\gamma} < \infty$ for all $\gamma \in (0, \min(\kappa, 1))$. Thus, again, Assumption *B* holds, with $m_0 = 0$. To make the presentation symmetric, we note, without going into details, that Tauberian and Abelian theorems for Laplace transforms can be used to show that

$$\mathbb{P}\{|\log(1 - W)| > x\} \sim \frac{\kappa^\kappa}{(c\Gamma(1-\alpha))^\kappa \Gamma(1+\alpha\kappa)} e^{-\alpha\kappa x}, \quad x \rightarrow \infty.$$

4 An auxiliary general branching process

To prove Theorem 1, we need some auxiliary results on a general branching process generated by T (a perturbed random walk). Here is its definition in terms of a population of individuals initiated at time 0 by one individual, the ancestor. An individual born at time $s \geq 0$ produces an offspring whose birth times have the same distribution as $(s + T_k)_{k \geq 1}$. All individuals act independently of each other. An individual resides in the j th generation if it has exactly j ancestors.

For $t \geq 0$ and $j \in \mathbb{N}$, denote by $N_j(t)$ the number of the j th generation individuals with birth times smaller than or equal to t and put $V_j(t) := \mathbb{E}N_j(t)$. Then $N_1(t) = N(t)$, and, for $j \geq 2$, N_j admits the following decomposition

$$N_j(t) = \sum_{r \geq 1} N_{j-1}^{(r)}(t - T_r) \mathbb{1}_{\{T_r \leq t\}}, \quad t \geq 0,$$

where, for $r \in \mathbb{N}$, $N_{j-1}^{(r)}(t)$ is the number of the j th generation individuals who are descendants of the first-generation individual with birth time T_r , and whose birth times fall in $[T_r, T_r + t]$. By the branching property, $(N_{j-1}^{(1)}(t))_{t \geq 0}$, $(N_{j-1}^{(2)}(t))_{t \geq 0}, \dots$ are independent copies of $(N_{j-1}(t))_{t \geq 0}$ which are also independent of T . As a consequence, $V_1(t) = V(t)$, and

$$V_j(t) = (V_{j-1} * V)(t) = \int_{[0, t]} V_{j-1}(t - y) dV(y), \quad j \geq 2, \quad t \geq 0, \quad (9)$$

that is, $V_j = V^{*(j)}$ is the j -fold Lebesgue-Stieltjes convolution of V with itself. The remainder of this section is concerned with obtaining precise and asymptotic estimates for V_j .

According to Lemma 1, Assumptions A and B ensure that inequality (5) holds true. Now we prove that V_n admits a similar estimate.

Lemma 3. *Suppose (5). Then*

$$|V_n(t) - \rho_n t^{\alpha n}| \leq \sum_{i=0}^{n-1} \binom{n}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{n-i}}{\Gamma(\alpha i + \beta(n-i) + 1)} (Ct^\alpha)^i (Dt^\beta)^{n-i}, \quad n \in \mathbb{N}, t \geq 0, \quad (10)$$

where ρ_n is as given in (4).

Proof. We use mathematical induction. For $n = 1$, (10) reduces to (5). Assume that (10) holds with $n - 1$ replacing n . Using (9) for the first equality we conclude that

$$\begin{aligned} |V_n(t) - \rho_n t^{\alpha n}| &= \left| \int_{[0,t]} V_{n-1}(t-y) dV(y) - \rho_n t^{\alpha n} \right| \\ &\leq \int_{[0,t]} |V_{n-1}(t-y) - \rho_{n-1}(t-y)^{\alpha(n-1)}| dV(y) + \left| \rho_{n-1} \int_{[0,t]} (t-y)^{\alpha(n-1)} dV(y) - \rho_n t^{\alpha n} \right|. \end{aligned}$$

Invoking the induction assumption and then integration by parts we obtain

$$\begin{aligned} A_n(t) &:= \int_{[0,t]} |V_{n-1}(t-y) - \rho_{n-1}(t-y)^{\alpha(n-1)}| dV(y) \\ &\leq \sum_{i=0}^{n-2} \binom{n-1}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{n-1-i}}{\Gamma(\alpha i + \beta(n-1-i) + 1)} D^{n-1-i} C^i \int_{[0,t]} (t-y)^{\alpha i + \beta(n-1-i)} dV(y) \\ &= \sum_{i=0}^{n-2} \binom{n-1}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{n-1-i}}{\Gamma(\alpha i + \beta(n-1-i) + 1)} D^{n-1-i} C^i \int_0^t V(t-y) d(y^{\alpha i + \beta(n-1-i)}). \end{aligned}$$

In view of (5),

$$\begin{aligned} \int_0^t V(t-y) d(y^{\alpha i + \beta(n-1-i)}) &\leq \int_0^t (C(t-y)^\alpha + D(t-y)^\beta) d(y^{\alpha i + \beta(n-1-i)}) \\ &= C \frac{\Gamma(\alpha+1)\Gamma(\alpha i + \beta(n-1-i) + 1)}{\Gamma(\alpha(i+1) + \beta(n-1-i) + 1)} t^{\alpha(i+1) + \beta(n-1-i)} \\ &\quad + D \frac{\Gamma(\beta+1)\Gamma(\alpha i + \beta(n-1-i) + 1)}{\Gamma(\alpha i + \beta(n-i) + 1)} t^{\alpha i + \beta(n-i)}, \end{aligned}$$

whence

$$\begin{aligned}
A_n(t) &\leq \sum_{i=0}^{n-2} \binom{n-1}{i} \frac{(\Gamma(\alpha+1))^{i+1}(\Gamma(\beta+1))^{n-1-i}}{\Gamma(\alpha(i+1)+\beta(n-1-i)+1)} D^{n-1-i} C^{i+1} t^{\alpha(i+1)+\beta(n-1-i)} \\
&\quad + \sum_{i=0}^{n-2} \binom{n-1}{i} \frac{(\Gamma(\alpha+1))^i(\Gamma(\beta+1))^{n-i}}{\Gamma(\alpha i+\beta(n-i)+1)} D^{n-i} C^i t^{\alpha i+\beta(n-i)} \\
&\quad = (n-1) \frac{(\Gamma(\alpha+1))^{n-1} \Gamma(\beta+1)}{\Gamma(\alpha(n-1)+\beta+1)} D C^{n-1} t^{\alpha(n-1)+\beta} \\
&\quad + \sum_{i=1}^{n-2} \left(\binom{n-1}{i-1} + \binom{n-1}{i} \right) \frac{(\Gamma(\alpha+1))^i(\Gamma(\beta+1))^{n-i}}{\Gamma(\alpha i+\beta(n-i)+1)} D^{n-i} C^i t^{\alpha i+\beta(n-i)} + \frac{(\Gamma(\beta+1))^n}{\Gamma(\beta n+1)} D^n t^{\beta n} \\
&\quad = (n-1) \frac{(\Gamma(\alpha+1))^{n-1} \Gamma(\beta+1)}{\Gamma(\alpha(n-1)+\beta+1)} D C^{n-1} t^{\alpha(n-1)+\beta} \\
&\quad \quad + \sum_{i=0}^{n-2} \binom{n}{i} \frac{(\Gamma(\alpha+1))^i(\Gamma(\beta+1))^{n-i}}{\Gamma(\alpha i+\beta(n-i)+1)} D^{n-i} C^i t^{\alpha i+\beta(n-i)}. \quad (11)
\end{aligned}$$

To obtain the first equality we have changed the variable $i \rightarrow i-1$ under the first sum appearing in the first inequality, singled out the term which corresponds to $i = n-1$ and also singled out the term of the second sum which corresponds to $i = 0$.

Further, we infer

$$\begin{aligned}
&\left| \rho_{n-1} \int_{[0,t]} (t-y)^{\alpha(n-1)} dV(y) - \rho_n t^{\alpha n} \right| \\
&\quad = \rho_{n-1} \left| \int_{[0,t]} (t-y)^{\alpha(n-1)} dV(y) - C \int_{[0,t]} (t-y)^\alpha d(y^{\alpha(n-1)}) \right| \\
&\quad = \left| \rho_{n-1} \int_0^t (V(t-y) - C(t-y)^\alpha) d(y^{\alpha(n-1)}) \right| \leq D \rho_{n-1} \int_0^t (t-y)^\beta d(y^{\alpha(n-1)}) \\
&\quad = D \frac{(C\Gamma(\alpha+1))^{n-1} \Gamma(\alpha(n-1)+1) \Gamma(\beta+1)}{\Gamma(\alpha(n-1)+1) \Gamma(\alpha(n-1)+\beta+1)} t^{\alpha(n-1)+\beta} \\
&\quad \quad = \frac{(\Gamma(\alpha+1))^{n-1} \Gamma(\beta+1)}{\Gamma(\alpha(n-1)+\beta+1)} D C^{n-1} t^{\alpha(n-1)+\beta}
\end{aligned}$$

having utilized integration by parts for the second equality and (5) for the first inequality. The sum of the last expression and the first term on the right-hand side of (11) is equal to the term of the sum in (10) which corresponds to $i = n-1$. The proof of Lemma 3 is complete. \square

We proceed with a technical estimate of an analytic flavor.

Lemma 4. For $t \geq 0$ and positive integer j satisfying

$$2D\Gamma(\beta+1)j(\alpha(j-1)+\beta+1)^{\alpha-\beta} \leq C\Gamma(\alpha+1)t^{\alpha-\beta}, \quad (12)$$

the following inequality holds

$$\sum_{i=0}^{j-1} \binom{j}{i} \frac{(\Gamma(\alpha+1))^i(\Gamma(\beta+1))^{j-i}}{\Gamma(\alpha i+\beta(j-i)+1)} (Ct^\alpha)^i (Dt^\beta)^{j-i} \leq \frac{21}{10} D C^{j-1} j \frac{(\Gamma(\alpha+1))^{j-1} \Gamma(\beta+1)}{\Gamma(\alpha(j-1)+\beta+1)} t^{\alpha(j-1)+\beta}.$$

Proof. Write

$$\begin{aligned} & \sum_{i=0}^{j-1} \binom{j}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{j-i}}{\Gamma(\alpha i + \beta(j-i) + 1)} (Ct^\alpha)^i (Dt^\beta)^{j-i} \\ &= DC^{j-1} j \frac{(\Gamma(\alpha+1))^{j-1} \Gamma(\beta+1)}{\Gamma(\alpha(j-1) + \beta + 1)} t^{\alpha(j-1) + \beta} \\ & \quad + \sum_{i=0}^{j-2} \binom{j}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{j-i}}{\Gamma(\alpha i + \beta(j-i) + 1)} (Ct^\alpha)^i (Dt^\beta)^{j-i}. \end{aligned}$$

Assuming that t and j satisfy (12) we intend to show that the second term on the right-hand side does not exceed the first term times 11/10. Indeed, using the inequality

$$\binom{j}{i} \leq j^{j-i} \quad (13)$$

and (39) we infer

$$\begin{aligned} & \frac{\Gamma(\alpha(j-1) + \beta + 1)}{DC^{j-1} j (\Gamma(\alpha+1))^{j-1} \Gamma(\beta+1)} t^{\alpha(j-1) + \beta} \sum_{i=0}^{j-2} \binom{j}{i} \frac{(\Gamma(\alpha+1))^i (\Gamma(\beta+1))^{j-i}}{\Gamma(\alpha i + \beta(j-i) + 1)} (Ct^\alpha)^i (Dt^\beta)^{j-i} \\ & \leq \frac{C\Gamma(\alpha+1) t^{\alpha-\beta}}{D\Gamma(\beta+1) j} \sum_{i=0}^{j-2} \left(\frac{D\Gamma(\beta+1) j}{C\Gamma(\alpha+1) t^{\alpha-\beta}} \right)^{j-i} \frac{\Gamma(\alpha(j-1) + \beta + 1)}{\Gamma(\alpha i + \beta(j-i) + 1)} \\ & \leq \frac{11}{10} \frac{C\Gamma(\alpha+1) t^{\alpha-\beta}}{D\Gamma(\beta+1) j (\alpha(j-1) + \beta + 1)^{\alpha-\beta}} \sum_{i=0}^{j-2} \left(\frac{D\Gamma(\beta+1) j (\alpha(j-1) + \beta + 1)^{\alpha-\beta}}{C\Gamma(\alpha+1) t^{\alpha-\beta}} \right)^{j-i} \\ & \leq \frac{11}{10} \frac{D\Gamma(\beta+1) j (\alpha(j-1) + \beta + 1)^{\alpha-\beta}}{C\Gamma(\alpha+1) t^{\alpha-\beta}} \left(1 - \frac{D\Gamma(\beta+1) j (\alpha(j-1) + \beta + 1)^{\alpha-\beta}}{C\Gamma(\alpha+1) t^{\alpha-\beta}} \right)^{-1} \leq \frac{11}{10}. \end{aligned}$$

The penultimate inequality is secured by $\sum_{i=0}^{j-2} (\dots)^{j-i} \leq \sum_{i \geq 2} (\dots)^i$. The function $x \mapsto x(1-x)^{-1}$ is increasing on $[0, 1)$ and equal to 1 at $x = 1/2$. This fact in combination with (12) justifies the last inequality. \square

Now we are ready to provide a simplification of formula (10) for convolution powers n and arguments t satisfying (12), with $j = n$.

Corollary 1. *Suppose (5). Then, for $t \geq 0$ and positive integer j satisfying (12),*

$$|V_j(t) - \rho_j t^{\alpha j}| \leq \frac{21}{10} DC^{j-1} j \frac{(\Gamma(\alpha+1))^{j-1} \Gamma(\beta+1)}{\Gamma(\alpha(j-1) + \beta + 1)} t^{\alpha(j-1) + \beta} \quad (14)$$

and

$$V_j(t) \leq \frac{331}{100} \rho_j t^{\alpha j}, \quad t \geq 0. \quad (15)$$

Proof. Inequality (14) is an immediate consequence of Lemmas 3 and 4.

A part of (14) reads

$$V_j(t) \leq \rho_j t^{\alpha j} + \frac{21}{10} DC^{j-1} j \frac{(\Gamma(\alpha+1))^{j-1} \Gamma(\beta+1)}{\Gamma(\alpha(j-1) + \beta + 1)} t^{\alpha(j-1) + \beta}.$$

To prove (15) we bound from above the ratio of the terms on the right-hand side of the last inequality

$$\begin{aligned} \frac{21DC^{j-1}j(\Gamma(\alpha+1))^{j-1}\Gamma(\beta+1)t^{\alpha(j-1)+\beta}}{10\Gamma(\alpha(j-1)+\beta+1)\rho_j t^{\alpha j}} &= \frac{21Dj\Gamma(\beta+1)\Gamma(\alpha j+1)}{10C\Gamma(\alpha+1)t^{\alpha-\beta}\Gamma(\alpha(j-1)+\beta+1)} \\ &\leq \frac{21Dj\Gamma(\beta+1)}{10C\Gamma(\alpha+1)t^{\alpha-\beta}} \cdot \frac{11}{10}(\alpha j+1)^{\alpha-\beta} \leq \frac{231(\alpha j+1)^{\alpha-\beta}}{200(\alpha(j-1)+\beta+1)^{\alpha-\beta}} \\ &\leq \frac{231}{200} \left(1 + \frac{\alpha-\beta}{\alpha(j-1)+\beta+1}\right)^\alpha \leq \frac{231}{100}. \end{aligned}$$

Here, the first inequality follows from Lemma 10, the second literally repeats (12) and the last is a consequence of $\frac{\alpha-\beta}{\alpha(j-1)+\beta+1} < 1$, which in turn is secured by $\alpha \in (0, 1)$ and $\beta \geq 0$. \square

Corollary 2. *Suppose (5). Then, for all $\gamma > 0$,*

$$\limsup_{t \rightarrow \infty} \left| \frac{V_j(y)}{\rho_j y^{\alpha j}} - 1 \right| = 0 \quad (16)$$

whenever $j = j(t)$ satisfies $j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$, and particularly

$$V_j(t) \sim \rho_j t^{\alpha j}, \quad t \rightarrow \infty. \quad (17)$$

Proof. If $j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$, then (12) holds for large j and t and

$$\begin{aligned} \sup_{y \geq \gamma t} \frac{DC^{j-1}j \frac{(\Gamma(\alpha+1))^{j-1}\Gamma(\beta+1)}{\Gamma(\alpha(j-1)+\beta+1)} y^{\alpha(j-1)+\beta}}{\rho_j y^{\alpha j}} &\leq \frac{DC^{j-1}j \frac{(\Gamma(\alpha+1))^{j-1}\Gamma(\beta+1)}{\Gamma(\alpha(j-1)+\beta+1)}}{\rho_j t^{\alpha-\beta}} \\ &\sim \frac{D\Gamma(\beta+1)\alpha^{\alpha-\beta} j^{\alpha-\beta+1}}{C\Gamma(\alpha+1)t^{\alpha-\beta}} \rightarrow 0 \end{aligned}$$

as $t \rightarrow \infty$. Here, we have used a standard asymptotic relation

$$\Gamma(x+a)/\Gamma(x) \sim x^a, \quad x \rightarrow \infty \quad (18)$$

for fixed $a > 0$, with $x = 1 + \alpha j - (\alpha - \beta)$ and $a = \alpha - \beta$. Thus, according to (14), the claim follows. \square

5 Proof of Theorem 1

We start by collecting several (additional) auxiliary results to be used in the proof of Theorem 1.

Recall that $f : [0, \infty) \rightarrow [0, \infty)$ is called *directly Riemann integrable* (dRi) function on $[0, \infty)$, if $\bar{\sigma}(h) < \infty$ for each $h > 0$ and $\lim_{h \rightarrow 0^+} (\bar{\sigma}(h) - \underline{\sigma}(h)) = 0$, where

$$\bar{\sigma}(h) := h \sum_{n \geq 1} \sup_{(n-1)h \leq y < nh} f(y) \quad \text{and} \quad \underline{\sigma}(h) := h \sum_{n \geq 1} \inf_{(n-1)h \leq y < nh} f(y).$$

A function $f : \mathbb{R} \rightarrow [0, \infty)$ is called dRi on \mathbb{R} , if the same two conditions hold, with $n \in \mathbb{Z}$ replacing $n \geq 1$ in the definition of the integral sums.

The next result follows from the proof of Lemma 4.5 in [20].

Lemma 5. Let $f : [0, \infty) \rightarrow [0, \infty)$ be dRi on $[0, \infty)$ and j a positive integer, possibly dependent on t and possibly divergent to ∞ together with t . Then

$$\int_{[0,t]} f(t-y)dV_j(y) = O(V_{j-1}(t)), \quad t \rightarrow \infty.$$

Lemma 6. Suppose (5) and let $j = j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$. Then

$$\int_{(t,\infty)} e^{t-y}dV_j(y) = O(V_{j-1}(t)), \quad t \rightarrow \infty.$$

Proof. Let $h : \mathbb{R} \rightarrow [0, \infty)$ be a dRi function on \mathbb{R} satisfying $h(t) = 0$ for $t > 0$. We start as in the proof of Lemma 4.6 in [20]: for $t \geq 0$,

$$\int_{(t,\infty)} h(t-y)dV_j(y) = \int_{[0,t]} h_1(t-y)dV_{j-1}(y) + \int_{(t,\infty)} h_2(t-y)dV_{j-1}(y),$$

where $h_1(t) = \int_{(t,\infty)} h(t-y)dV(y)$ and $h_2(t) = \int_{[0,\infty)} h(t-y)dV(y)$ for $t \in \mathbb{R}$. By Lemma A.1 in [20], $h_1(t) \leq b$ for some $b > 0$ and all $t \geq 0$, whence

$$\int_{[0,t]} h_1(t-y)dV_{j-1}(y) = O(V_{j-1}(t)), \quad t \rightarrow \infty.$$

Now we put $h(t) = e^t \mathbb{1}_{(-\infty,0]}(t)$ and note that all the formulae given in the preceding part of the present proof hold for this h . Plainly, $h_2(t) = e^t \int_{[0,\infty)} e^{-y}dV(y) = \kappa e^t$ for $t \leq 0$, where $\kappa := \mathbb{E}e^{-\eta}(1 - \mathbb{E}e^{-\xi})^{-1}$. Integrating by parts yields

$$\begin{aligned} & \int_{(t,\infty)} h_2(t-y)dV_{j-1}(y) \\ &= \kappa \int_{(t,\infty)} e^{t-y}dV_{j-1}(y) = -\kappa V_{j-1}(t) + \kappa \int_t^\infty e^{t-y}V_{j-1}(y)dy =: \kappa C_j(t) \end{aligned}$$

for $t \geq 0$. By Corollary 2, (5) together with our choice of $j = j(t)$ ensures (16). In view of (16), given $\varepsilon > 0$ we obtain, for large enough t ,

$$\begin{aligned} 0 \leq C_j(t) &\leq -V_{j-1}(t) + (1 + \varepsilon)\rho_{j-1} \int_t^\infty e^{t-y}y^{\alpha(j-1)}dy = -V_{j-1}(t) \\ &+ (1+\varepsilon)(C\Gamma(\alpha+1))^{j-1} \left(\sum_{k=0}^{[\alpha(j-1)]} \frac{t^{\alpha(j-1)-k}}{\Gamma(\alpha(j-1)+1-k)} + \frac{1}{\Gamma(\{\alpha(j-1)\})} \int_t^\infty e^{t-y}y^{\{\alpha(j-1)\}-1}dy \right), \end{aligned}$$

where $\{x\}$ denotes the fractional part of $x \in \mathbb{R}$. The expression in the parantheses is asymptotically equivalent to $t^{\alpha(j-1)}/\Gamma(\alpha(j-1)+1)$ whenever $j(t) = o(t)$ as $t \rightarrow \infty$. Hence, recalling that, according to (17) (which holds according to Corollary 2), $V_{j-1}(t) \sim \rho_{j-1}t^{\alpha(j-1)}$ as $t \rightarrow \infty$ we conclude that $\limsup_{t \rightarrow \infty} (C_j(t)/V_{j-1}(t)) \leq \varepsilon$ and thereupon $C_j(t) = o(V_{j-1}(t))$ as $t \rightarrow \infty$ upon sending $\varepsilon \rightarrow 0+$. \square

Lemma 7. Suppose (5) and let $j = j(t) \rightarrow \infty$ and $j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$. Then

$$\lim_{s \rightarrow \infty} \limsup_{t \rightarrow \infty} \frac{j^\alpha}{\rho_{j-1}t^{\alpha j}} \int_{(st/j,t]} y^\alpha d_y(-V_{j-1}(t-y)) = 0.$$

Proof. Integrating by parts we conclude that

$$\begin{aligned} \frac{j^\alpha}{\rho_{j-1}t^{\alpha j}} \int_{(st/j, t]} y^\alpha d_y(-V_{j-1}(t-y)) &= \frac{s^\alpha V_{j-1}(t(1-s/j))}{\rho_{j-1}t^{\alpha(j-1)}} \\ &+ \frac{\alpha j^\alpha}{\rho_{j-1}t^{\alpha j}} \int_{st/j}^t V_{j-1}(t-y)y^{\alpha-1}dy. \end{aligned}$$

In view of (16) which holds true according to Corollary 2,

$$\frac{s^\alpha V_{j-1}(t(1-s/j))}{\rho_{j-1}t^{\alpha(j-1)}} \sim s^\alpha(1-s/j)^{\alpha(j-1)} \rightarrow s^\alpha e^{-\alpha s}, \quad t \rightarrow \infty.$$

The right-hand side converges to 0 as $s \rightarrow \infty$. Further, we infer with the help of (10) that

$$\begin{aligned} \frac{j^\alpha}{\rho_{j-1}t^{\alpha j}} \int_{st/j}^t V_{j-1}(t-y)y^{\alpha-1}dy &\leq \frac{j^\alpha}{\rho_{j-1}t^{\alpha j}} \left(\int_{st/j}^t \rho_{j-1}(t-y)^{\alpha(j-1)}y^{\alpha-1}dy \right. \\ &\left. + \int_{st/j}^t \sum_{i=0}^{j-2} \binom{j-1}{i} \frac{(\Gamma(\alpha+1))^i(\Gamma(\beta+1))^{j-i-1}}{\Gamma(\alpha i + \beta(j-i-1)+1)} (C(t-y)^\alpha)^i (D(t-y)^\beta)^{j-i-1} y^{\alpha-1} dy \right). \quad (19) \end{aligned}$$

Changing the variable we obtain for the first term

$$\frac{j^\alpha}{t^{\alpha j}} \int_{st/j}^t (t-y)^{\alpha(j-1)}y^{\alpha-1}dy = \int_s^j (1-y/j)^{\alpha(j-1)}y^{\alpha-1}dy \rightarrow \int_s^\infty e^{-\alpha y}y^{\alpha-1}dy, \quad t \rightarrow \infty$$

having utilized the Lebesgue dominated convergence theorem for the limit relation. The right-hand side converges to 0 as $s \rightarrow \infty$. Invoking $y^{\alpha-1} \leq (st/j)^{\alpha-1}$ for $y \in [st/j, t]$ and then evaluating the remaining integral we conclude that the second term in (19) does not exceed

$$\begin{aligned} &\frac{j^\alpha}{\rho_{j-1}t^{\alpha j}} \left(\frac{st}{j}\right)^{\alpha-1} \sum_{i=0}^{j-2} \binom{j-1}{i} \frac{(C\Gamma(\alpha+1))^i (D\Gamma(\beta+1))^{j-i-1}}{\Gamma(\alpha i + \beta(j-i-1)+1)} \frac{(t-st/j)^{\alpha i + \beta(j-i-1)+1}}{\alpha i + \beta(j-i-1)+1} \\ &= s^{\alpha-1} j \sum_{i=0}^{j-2} \binom{j-1}{i} \frac{\Gamma(\alpha(j-1)+1)}{\Gamma(\alpha i + \beta(j-i-1)+1)} \left(\frac{D\Gamma(\beta+1)t^\beta}{C\Gamma(\alpha+1)t^\alpha}\right)^{j-i-1} \frac{(1-s/j)^{\alpha i + \beta(j-i-1)+1}}{\alpha i + \beta(j-i-1)+1} \\ &\leq \frac{s^{\alpha-1}}{\beta} \sum_{i=0}^{j-2} \binom{j-1}{i} \frac{\Gamma(\alpha(j-1)+1)}{\Gamma(\alpha i + \beta(j-i-1)+1)} \left(\frac{D\Gamma(\beta+1)t^\beta}{C\Gamma(\alpha+1)t^\alpha}\right)^{j-i-1} \\ &\leq \frac{11s^{\alpha-1}}{10\beta} \sum_{i=0}^{j-2} \left((j-1)(\alpha(j-1)+1)^{\alpha-\beta} \frac{D\Gamma(\beta+1)}{C\Gamma(\alpha+1)t^{\alpha-\beta}} \right)^{j-i-1} \\ &\leq \frac{11s^{\alpha-1}}{10\beta} \frac{D\Gamma(\beta+1)}{C\Gamma(\alpha+1)} (j-1) \left(\frac{\alpha(j-1)+1}{t} \right)^{\alpha-\beta} \left(1 - \frac{D\Gamma(\beta+1)}{C\Gamma(\alpha+1)} (j-1) \left(\frac{\alpha(j-1)+1}{t} \right)^{\alpha-\beta} \right)^{-1} \\ &\rightarrow 0, \quad t \rightarrow \infty. \end{aligned}$$

We have used $0 < (1-s/j)^{\alpha i + \beta(j-i-1)+1} \leq 1$ and $\frac{1}{\alpha i + \beta(j-i-1)+1} \leq \frac{1}{\beta j}$ for the first inequality, formula (13) and Lemma 10 for the second. Our assumption $j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$ entails $\lim_{t \rightarrow \infty} (j(t)-1)((\alpha(j(t)-1)+1)/t)^{\alpha-\beta} = 0$, thereby justifying the limit relation. \square

A major part of the proof of Theorem 1 is covered by Theorems 2 and 3 given next.

Theorem 2. *Suppose (5) and let $j = j(t) \rightarrow \infty$ and $j(t) = o(t^{\frac{\alpha-\beta}{\alpha-\beta+1}})$ as $t \rightarrow \infty$. Then*

$$\left(\frac{c(j(t))^\alpha}{\rho_{[j(t)u]_{-1}} t^{\alpha[j(t)u]}} \sum_{r \geq 1} V_{[j(t)u]_{-1}}(t - T_r) \mathbb{1}_{\{T_r \leq t\}} \right)_{u>0} \xrightarrow{\text{f.d.d.}} \left(\int_{[0, \infty)} e^{-\alpha u y} dZ_\alpha^-(y) \right)_{u>0}, \quad t \rightarrow \infty,$$

where ρ_n is as given in (4), and Z_α^- is an inverse α -stable subordinator.

Our proof of Theorem 2 uses an auxiliary technical result, which is a slight reformulation of Lemma A.5 in [16]. Let \mathcal{D} denote the Skorokhod space of càdlàg functions defined on $[0, \infty)$. We assume that the space \mathcal{D} is endowed with the J_1 -topology. Comprehensive information about the J_1 -topology on \mathcal{D} can be found in [4].

Lemma 8. *Let $0 \leq a < b < \infty$ and, for each $k \in \mathbb{N}$, $y_k : [0, \infty) \rightarrow \mathbb{R}$ be a right-continuous bounded and nondecreasing function. Assume that $\lim_{k \rightarrow \infty} x_k = x$ on \mathcal{D} and that, for each $t \geq 0$, $\lim_{k \rightarrow \infty} y_k(t) = y(t)$, where $y : [0, \infty) \rightarrow \mathbb{R}$ is a bounded continuous function. Then*

$$\lim_{k \rightarrow \infty} \int_{[a, b]} x_k(t) dy_k(t) = \int_{[a, b]} x(t) dy(t).$$

Proof of Theorem 2. For notational simplicity we write j for $j(t)$. In view of the Cramér-Wold device and the equality $\int_{[0, \infty)} e^{-\alpha u y} dZ_\alpha^-(y) = \alpha u \int_0^\infty Z_\alpha^-(y) e^{-\alpha u y} dy$ for $u > 0$, obtained with the help of integration by parts, it suffices to show that, for any $\ell \in \mathbb{N}$, any positive $\lambda_1, \dots, \lambda_\ell$ and any $0 < u_1 < \dots < u_\ell$,

$$\sum_{i=1}^{\ell} \lambda_i \frac{c j^\alpha \sum_{r \geq 1} V_{[ju_i]_{-1}}(t - T_r) \mathbb{1}_{\{T_r \leq t\}}}{\rho_{[ju_i]_{-1}} t^{\alpha[ju_i]}} \xrightarrow{d} \sum_{i=1}^{\ell} \lambda_i \alpha u_i \int_0^\infty Z_\alpha^-(y) e^{-\alpha u_i y} dy, \quad t \rightarrow \infty,$$

where, as before, \xrightarrow{d} denotes convergence in distribution. Write, for any $s > 0$ and sufficiently large t ,

$$\begin{aligned} \frac{c j^\alpha \sum_{r \geq 1} V_{[ju_i]_{-1}}(t - T_r) \mathbb{1}_{\{T_r \leq t\}}}{\rho_{[ju_i]_{-1}} t^{\alpha[ju_i]}} &= \frac{c j^\alpha \int_{[0, t]} V_{[ju_i]_{-1}}(t - y) dN(y)}{\rho_{[ju_i]_{-1}} t^{\alpha[ju_i]}} \\ &= \frac{c j^\alpha \int_{[0, t]} N(y) d(-V_{[ju_i]_{-1}}(t - y))}{\rho_{[ju_i]_{-1}} t^{\alpha[ju_i]}} \\ &= \frac{1}{\rho_{[ju_i]_{-1}} t^{\alpha([ju_i]_{-1})}} \int_{[0, s]} \frac{c N(yt/j)}{(t/j)^\alpha} d_y(-V_{[ju_i]_{-1}}(t(1 - y/j))) \\ &\quad + \frac{c j^\alpha}{\rho_{[ju_i]_{-1}} t^{\alpha[ju_i]}} \int_{(st/j, t]} N(y) d_y(-V_{[ju_i]_{-1}}(t - y)). \end{aligned}$$

Using (16) we obtain, for each fixed $y > 0$, as $t \rightarrow \infty$,

$$V_{[ju_i]_{-1}} \left(t \left(1 - \frac{y}{j} \right) \right) \sim \rho_{[ju_i]_{-1}} t^{\alpha([ju_i]_{-1})} \left(1 - \frac{y}{j} \right)^{\alpha([ju_i]_{-1})} \sim \rho_{[ju_i]_{-1}} t^{\alpha([ju_i]_{-1})} e^{-\alpha u_i y}. \quad (20)$$

Under the sole assumption $\mathbb{P}\{\xi > t\} \sim ct^{-\alpha}$ as $t \rightarrow \infty$, an application of part (B.4) of Theorem 3.2 in [1] yields

$$(c(t/j)^{-\alpha}N(yt/j))_{y \geq 0} \Longrightarrow (\mathcal{Z}_\alpha^-(y))_{y \geq 0}, \quad t \rightarrow \infty$$

on \mathcal{D} , where \Longrightarrow denotes weak convergence on a function space. Here, we have used the fact that the assumptions imposed on j ensure that $\lim_{t \rightarrow \infty} (t/j(t)) = \infty$. Let $(t_k)_{k \geq 1}$ be any sequence of positive numbers satisfying $\lim_{k \rightarrow \infty} t_k = \infty$. According to the Skorokhod representation theorem, there exist $\hat{\mathcal{Z}}_\alpha^-$ a version of \mathcal{Z}_α^- and $((\hat{N}_{t_k}(y))_{y \geq 0})_{k \geq 1}$ a version of $((c(t_k/j(t_k))^{-\alpha}N(yt_k/j(t_k)))_{y \geq 0})_{k \geq 1}$ such that, for all $T > 0$,

$$\lim_{k \rightarrow \infty} \sup_{y \in [0, T]} |\hat{N}_{t_k}(y) - \hat{\mathcal{Z}}_\alpha^-(y)| = 0 \quad \text{a.s.}$$

In view of (20),

$$\lim_{k \rightarrow \infty} \frac{V_{\lfloor j(t_k)u_i \rfloor - 1}(t_k(1 - y/j(t_k)))}{\rho_{\lfloor j(t_k)u_i \rfloor - 1} t_k^{\alpha(\lfloor j(t_k)u_i \rfloor - 1)}} = e^{-\alpha u_i y}.$$

Using Lemma 8, with $x_k = \hat{N}_{t_k}$, $x = \hat{\mathcal{Z}}_\alpha^-$, and y_k and y given by the functions on the left-hand and the right-hand side of the last centered formula, respectively, we obtain

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{1}{\rho_{\lfloor j(t_k)u_i \rfloor - 1} t_k^{\alpha(\lfloor j(t_k)u_i \rfloor - 1)}} \int_{[0, s]} \hat{N}_{t_k}(y) d_y(-V_{\lfloor j(t_k)u_i \rfloor - 1}(t_k(1 - y/j(t_k)))) \\ = \alpha u_i \int_0^s \hat{\mathcal{Z}}_\alpha^-(y) e^{-\alpha u_i y} dy \quad \text{a.s.} \end{aligned}$$

Since the diverging sequence $(t_k)_{k \geq 1}$ is arbitrary, this entails

$$\begin{aligned} \sum_{i=1}^{\ell} \lambda_i \frac{1}{\rho_{\lfloor j u_i \rfloor - 1} t^{\alpha(\lfloor j u_i \rfloor - 1)}} \int_{[0, s]} c(t/j)^{-\alpha} N(yt/j) d_y(-V_{\lfloor j u_i \rfloor - 1}(t(1 - y/j))) \\ \xrightarrow{d} \sum_{i=1}^{\ell} \lambda_i \alpha u_i \int_0^s \mathcal{Z}_\alpha^-(y) e^{-\alpha u_i y} dy, \quad t \rightarrow \infty. \end{aligned}$$

In view of

$$\lim_{s \rightarrow \infty} \sum_{i=1}^{\ell} \lambda_i \alpha u_i \int_0^s \mathcal{Z}_\alpha^-(y) e^{-\alpha u_i y} dy = \sum_{i=1}^{\ell} \lambda_i \alpha u_i \int_0^\infty \mathcal{Z}_\alpha^-(y) e^{-\alpha u_i y} dy \quad \text{a.s.},$$

we are left with showing that, for all $\varepsilon > 0$,

$$\lim_{s \rightarrow \infty} \limsup_{t \rightarrow \infty} \mathbb{P} \left\{ \frac{j^\alpha}{\rho_{\lfloor j u_i \rfloor - 1} t^{\alpha \lfloor j u_i \rfloor}} \int_{(st/j, t]} N(y) d_y(-V_{\lfloor j u_i \rfloor - 1}(t - y)) > \varepsilon \right\} = 0.$$

Noting that $\mathbb{E}N(y) = V(y) \leq 2Cy^\alpha$ for large y , this limit relation is ensured by Markov's inequality and Lemma 7. \square

Theorem 3. *Suppose (5) and let $t \mapsto j(t)$ be an integer-valued function satisfying $j(t) = o(t^{1/3})$ as $t \rightarrow \infty$ and $\lim_{t \rightarrow \infty} j(t) = \infty$. Then*

$$\frac{(j(t))^\alpha}{\rho_{j(t)-1} t^{\alpha j(t)}} \left(N_{j(t)}(t) - \sum_{r \geq 1} V_{j(t)-1}(t - T_r) \mathbb{1}_{\{T_r \leq t\}} \right) \xrightarrow{\mathbb{P}} 0, \quad t \rightarrow \infty.$$

Proof. Although the scheme of the proof is analogous to that of Theorem 3.1 in [20], technical details are essentially different at places.

For $j \in \mathbb{N}$ and $t \geq 0$, put $D_j(t) := \text{Var } N_j(t)$ and

$$I_j(t) := \mathbb{E} \left(\sum_{r \geq 1} V_{j-1}(t - T_r) \mathbb{1}_{\{T_r \leq t\}} - V_j(t) \right)^2$$

with the convention that $V_0(t) = 1$ for $t \geq 0$, so that $I_1(t) = D_1(t)$. We shall use a formula: for $j \geq 2$ and $t \geq 0$,

$$\begin{aligned} D_j(t) &= \mathbb{E} \left(\sum_{r \geq 1} (N_{j-1}^{(r)}(t - T_r) - V_{j-1}(t - T_r)) \mathbb{1}_{\{T_r \leq t\}} \right)^2 \\ &\quad + \mathbb{E} \left(\sum_{r \geq 1} V_{j-1}(t - T_r) \mathbb{1}_{\{T_r \leq t\}} - V_j(t) \right)^2 = \int_{[0, t]} D_{j-1}(t - y) dV(y) + I_j(t). \end{aligned} \quad (21)$$

Iterating (21) yields

$$\int_{[0, t]} D_{j-1}(t - y) dV(y) = \sum_{k=1}^{j-1} \int_{[0, t]} I_k(t - y) dV_{j-k}(y), \quad j \geq 2, \quad t \geq 0. \quad (22)$$

First we show that I_j is bounded from above by a nonnegative and nondecreasing function. We need an inequality obtained in the proof of Theorem 3.1 in [20]

$$\begin{aligned} I_j(t) &\leq V_{j-1}(t) V_j(t) + 2 \int_{[0, t]} V_{j-1}(t - y) V_j(t - y) dU(y) - (V_j(t))^2 \\ &\leq V_{j-1}(t) V_j(t) + 2 \int_{[0, t]} V_{j-1}(t - y) V_j(t - y) dU(y). \end{aligned} \quad (23)$$

Put $\tilde{U}(x) := \sum_{i \geq 1} \mathbb{P}\{S_i \leq x\}$ for $x \in \mathbb{R}$ and note that $U(x) = \tilde{U}(x) + 1$ for $x \geq 0$. Assumption A together with $\tilde{U}(0) = 0$ entails

$$\tilde{U}(x) \leq Cx^\alpha + C_1x^\rho, \quad x \geq 0$$

for some constant $C_1 > 0$. Integrating by parts and then using the last inequality yields

$$\begin{aligned} \int_{[0, t]} V_{j-1}(t - y) V_j(t - y) dU(y) &= V_{j-1}(t) V_j(t) + \int_{[0, t]} V_{j-1}(t - y) V_j(t - y) d\tilde{U}(y) \\ &= V_{j-1}(t) V_j(t) + \int_{[0, t]} \tilde{U}(t - y) d(V_{j-1}(y) V_j(y)) \\ &\leq V_{j-1}(t) V_j(t) + C \int_{[0, t]} (t - y)^\alpha d(V_{j-1}(y) V_j(y)) + C_1 \int_{[0, t]} (t - y)^\rho d(V_{j-1}(y) V_j(y)). \end{aligned}$$

With this at hand, integrating by parts once again we arrive at

$$\begin{aligned} I_j(t) &\leq V_{j-1}(t) V_j(t) + 2 \int_{[0, t]} V_{j-1}(t - y) V_j(t - y) dU(y) \leq 3V_{j-1}(t) V_j(t) \\ &\quad + 2C\alpha \int_0^t V_{j-1}(y) V_j(y) (t - y)^{\alpha-1} dy + 2C_1\rho \int_0^t V_{j-1}(y) V_j(y) (t - y)^{\rho-1} dy =: h_j(t). \end{aligned} \quad (24)$$

The function h_j is a nonnegative and nondecreasing function that we were looking for. Combining (24) with (21) and (22) we infer

$$D_{j-1}(t) = \sum_{k=1}^{j-2} \int_0^t I_k(t-y) dV_{j-k-1}(y) + I_{j-1}(t) \\ \leq \sum_{k=1}^{j-2} h_k(t) V_{j-k-1}(t) + h_{j-1}(t), \quad j \geq 2, t \geq 0. \quad (25)$$

Next, we obtain an upper bound for h_j (and for D_j) which is valid for large arguments. Fix $j \in \mathbb{N}$ and $s \geq 0$ satisfying

$$2D\Gamma(\beta+1)j(\alpha(j-1) + \beta + 1)^{\alpha-\beta} \leq C\Gamma(\alpha+1)s^{\alpha-\beta} \quad (26)$$

and also

$$2(\alpha(j-2) + 1)^{2\alpha} \leq C\Gamma(\alpha+1)s^\alpha, \quad (27)$$

$$33(2\alpha j + 1)^\alpha \leq 20C\Gamma(\alpha+1)s^\alpha, \quad (28)$$

$$11C_1\Gamma(\rho)(2\alpha j + 1)^{\alpha-\rho} \leq 10\Gamma(\alpha+1)s^{\alpha-\rho}, \quad (29)$$

and

$$j \geq 1/\alpha + 2. \quad (30)$$

Plainly, these inequalities trivially hold for large enough s whenever $j = j(s) = o(s^{1/2})$ as $s \rightarrow \infty$.

Put $r := (331/10)^2$. In view of (26), we conclude with the help of formula (15) that, for $1 \leq k \leq j$,

$$V_{k-1}(s)V_k(s) \leq r\rho_{k-1}\rho_k s^{\alpha(2k-1)}.$$

Substituting this into (24) we obtain

$$h_k(s) \leq 3r\rho_{k-1}\rho_k s^{\alpha(2k-1)} \\ + 2C\alpha r\rho_{k-1}\rho_k \int_0^s s^{\alpha(2k-1)}(s-y)^{\alpha-1} dy + 2C_1\rho r\rho_{k-1}\rho_k \int_0^s s^{\alpha(2k-1)}(s-y)^{\rho-1} dy \\ = 3r\rho_{k-1}\rho_k s^{\alpha(2k-1)} + 2rC\alpha\rho_{k-1}\rho_k B(\alpha(2k-1) + 1, \alpha) s^{2\alpha k} \\ + 2rC_1\rho\rho_{k-1}\rho_k B(\alpha(2k-1) + 1, \rho) B(\alpha(2k-1) + 1, \rho) s^{2\alpha k - \alpha + \rho}, \quad (31)$$

where B is the Euler beta-function. The second term on the right-hand side is of the highest order. Consider the ratio of the first and the second terms in (31):

$$\frac{3r\rho_{k-1}\rho_k s^{\alpha(2k-1)}}{2rC\alpha\rho_{k-1}\rho_k B(\alpha(2k-1) + 1, \alpha) s^{2\alpha k}} = \frac{3\Gamma(2\alpha k + 1)}{2C\Gamma(\alpha+1)\Gamma(\alpha(2k-1) + 1) s^\alpha} \\ \leq \frac{33(2\alpha k + 1)^\alpha}{20C\Gamma(\alpha+1) s^\alpha} \leq \frac{33(2\alpha j + 1)^\alpha}{20C\Gamma(\alpha+1) s^\alpha} \leq 1,$$

where the first inequality is justified by Lemma 10 and the last inequality is secured by (28). The ratio of the third and the second terms in (31) can be estimated as follows:

$$\begin{aligned} \frac{2rC_1\rho\rho_{k-1}\rho_k B(\alpha(2k-1)+1, \rho)s^{2\alpha k-\alpha+\rho}}{2rC\alpha\rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha)s^{2\alpha k}} &= \frac{C_1\Gamma(\rho)\Gamma(2\alpha k+1)}{\Gamma(\alpha+1)s^{\alpha-\rho}\Gamma(2\alpha k-\alpha+\rho+1)} \\ &\leq \frac{11C_1\Gamma(\rho)(2\alpha k+1)^{\alpha-\rho}}{10\Gamma(\alpha+1)s^{\alpha-\rho}} \leq \frac{11C_1\Gamma(\rho)(2\alpha j+1)^{\alpha-\rho}}{10\Gamma(\alpha+1)s^{\alpha-\rho}} \leq 1, \end{aligned}$$

where the first inequality follows from Lemma 10 and the last inequality is ensured by (29). Inequality (31) in combination with the two estimates enables us to conclude that, for $1 \leq k \leq j$,

$$h_k(s) \leq 6rC\alpha\rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha) =: C_2\rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha)s^{2\alpha k}.$$

Invoking (25) we further obtain

$$\begin{aligned} D_{j-1}(s) &\leq C_2\rho_{j-2}\rho_{j-1} B(\alpha(2j-3)+1, \alpha)s^{2\alpha(j-1)} \\ &\quad + \sum_{k=1}^{j-2} C_2\rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha)s^{2\alpha k} V_{j-k-1}(s). \end{aligned} \quad (32)$$

Write with the help of (4) and (15)

$$\begin{aligned} &\sum_{k=1}^{j-2} \rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha)s^{2\alpha k} V_{j-k-1}(s) \\ &\leq \frac{331}{100} \sum_{k=1}^{j-2} \rho_{k-1}\rho_k B(\alpha(2k-1)+1, \alpha)s^{2\alpha k} \rho_{j-k-1} s^{\alpha(j-k-1)} \\ &= \frac{331\Gamma(\alpha)}{100} \sum_{k=1}^{j-2} \frac{(C\Gamma(\alpha+1))^{j+k-2} \Gamma(\alpha(2k-1)+1)}{\Gamma(\alpha(k-1)+1)\Gamma(\alpha k+1)\Gamma(\alpha(j-k-1)+1)\Gamma(2\alpha k+1)} s^{\alpha(j+k-1)} \\ &\leq \frac{331(2\alpha+1)\Gamma(\alpha)}{100} \sum_{k=1}^{j-2} \frac{(C\Gamma(\alpha+1))^{j+k-2}}{(\Gamma(\alpha(k-1)+1))^2 \Gamma(\alpha(j-k-1)+1)} s^{\alpha(j+k-1)}. \end{aligned} \quad (33)$$

To justify the last inequality, we use the fact that the gamma-function Γ is increasing on $[2, \infty)$, whence, for $k \geq 1$,

$$\begin{aligned} \frac{1}{\Gamma(\alpha k+1)} &= \frac{\alpha k+1}{\Gamma(\alpha k+2)} \leq \frac{\alpha k+1}{\Gamma(\alpha(k-1)+2)} = \frac{\alpha k+1}{(\alpha(k-1)+1)\Gamma(\alpha(k-1)+1)} \\ &\leq \frac{\alpha+1}{\Gamma(\alpha(k-1)+1)} \end{aligned}$$

and

$$\frac{\Gamma(\alpha(2k-1)+1)}{\Gamma(2\alpha k+1)} = \frac{\Gamma(\alpha(2k-1)+2)}{\Gamma(2\alpha k+2)} \frac{2\alpha k+1}{\alpha(2k-1)+1} \leq \frac{2\alpha k+1}{\alpha(2k-1)+1} \leq \frac{2\alpha+1}{\alpha+1}.$$

The right-hand side of (33), with the multiplicative constant omitted, can be bounded from above as follows:

$$\begin{aligned} & \frac{1}{(\Gamma(\alpha(j-2)+1))^2} \sum_{k=1}^{j-2} \frac{(C\Gamma(\alpha+1))^{j+k-2} (\Gamma(\alpha(j-2)+1))^2}{(\Gamma(\alpha(k-1)+1))^2 \Gamma(\alpha(j-k-1)+1)} s^{\alpha(j+k-1)} \\ & \leq \frac{\text{const}}{(\Gamma(\alpha(j-2)+1))^2} \sum_{k=1}^{j-2} (C\Gamma(\alpha+1))^{j+k-2} (\alpha(j-2)+1)^{2\alpha(j-k-1)} s^{\alpha(j+k-1)}. \end{aligned} \quad (34)$$

Here and hereafter, const denotes a constant whose value is of no importance and may change from one appearance to another. The last inequality is a consequence of

$$\frac{\Gamma(\alpha(j-2)+1)}{\Gamma(\alpha(k-1)+1)} \leq \frac{11}{10} (\alpha(j-2)+1)^{\alpha(j-k-1)},$$

which follows from Lemma 10, and $\frac{1}{\Gamma(\alpha(j-k-1)+1)} \leq \frac{1}{\min_{z \in [1,2]} \Gamma(z)} < \infty$ for $1 \leq k \leq j-2$ (the minimum cannot be equal to 0 because $\Gamma(z)$ is the moment of order $z-1$ of an exponentially distributed random variable with mean one). We proceed by bounding from above the right-hand side of (34), up to the multiplicative constant:

$$\begin{aligned} & \frac{(C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2} \sum_{k=1}^{j-2} \left(\frac{(\alpha(j-2)+1)^{2\alpha}}{C\Gamma(\alpha+1)s^\alpha} \right)^{j-k-1} \\ & = \frac{(C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2} \sum_{k=1}^{j-2} \left(\frac{(\alpha(j-2)+1)^{2\alpha}}{C\Gamma(\alpha+1)s^\alpha} \right)^k \\ & \leq \frac{(C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2} \sum_{k \geq 1} \left(\frac{(\alpha(j-2)+1)^{2\alpha}}{C\Gamma(\alpha+1)s^\alpha} \right)^k \leq \frac{(C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2}. \end{aligned}$$

Here, the last inequality is secured by (27). Combining this with (32) we conclude that

$$D_{j-1}(s) \leq C_2 \rho_{j-2} \rho_{j-1} B(\alpha(2j-3)+1, \alpha) s^{2\alpha(j-1)} + \frac{\text{const} (C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2}.$$

Condition (30) entails $2j \geq 1/\alpha + 3$. This together with monotonicity of the gamma-function on $[2, \infty)$ proves

$$\frac{\Gamma(\alpha)(C\Gamma(\alpha+1))^{2j-3}}{\Gamma^2(\alpha(j-2)+1) \rho_{j-2} \rho_{j-1} B(\alpha(2j-3)+1, \alpha)} = \frac{\Gamma(\alpha(j-1)+1) \Gamma(\alpha(2j-2)+1)}{\Gamma(\alpha(j-2)+1) \Gamma(\alpha(2j-3)+1)} \geq 1$$

and thereupon

$$D_{j-1}(s) \leq \text{const} \frac{(C\Gamma(\alpha+1))^{2j-3} s^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2}. \quad (35)$$

Using (35) we infer

$$\begin{aligned} \int_{[0,t]} D_{j-1}(t-y) dV(y) &= \int_{[0,t-j]} D_{j-1}(t-y) dV(y) + \int_{(t-j,t]} D_{j-1}(t-y) dV(y) \\ &\leq \text{const} \frac{(C\Gamma(\alpha+1))^{2j-3}}{(\Gamma(\alpha(j-2)+1))^2} \int_{[0,t]} (t-y)^{2\alpha(j-1)} dV(y) + \max_{s \in [0,j]} D_{j-1}(s) U(j) \end{aligned} \quad (36)$$

having utilized $V(t) - V(t - j) \leq U(j)$, see formula (40) in [7] and its proof. Integrating by parts and then invoking (5) we estimate the last integral as follows:

$$\begin{aligned} \int_{[0,t]} (t-y)^{2\alpha(j-1)} dV(y) &\leq 2\alpha(j-1) \int_0^t (Cy^\alpha + Dy^\beta)(t-y)^{2\alpha(j-1)-1} dy \\ &= 2\alpha(j-1)(CB(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)} + DB(2\alpha(j-1), \beta+1)t^{2\alpha(j-1)+\beta}) \\ &= O(jB(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)}), \quad t \rightarrow \infty. \end{aligned}$$

To justify the last equality, note that, as $j, t \rightarrow \infty$,

$$B(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)} \sim \text{const} \frac{t^{\alpha(j-1)}}{j^{\alpha+1}}, \quad B(2\alpha(j-1), \beta+1)t^{2\alpha(j-1)+\beta} \sim \text{const} \frac{t^{2\alpha(j-1)+\beta}}{j^{\beta+1}}.$$

Hence, $B(2\alpha(j-1), \beta+1)t^{2\alpha(j-1)+\beta} = o(B(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)})$ as a consequence of $j = j(t) = o(t^{1/3})$. Thus, the first term on the right-hand side of (36) is of the order

$$\frac{(C\Gamma(\alpha+1))^{2j-3}}{(\Gamma(\alpha(j-2)+1))^2} jB(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)}.$$

The product of this and the squared normalization (appearing in the theorem) vanishes under the assumption $j = j(t) = o(t^{1/3})$ as $t \rightarrow \infty$:

$$\begin{aligned} &\frac{(C\Gamma(\alpha+1))^{2j-3} jB(2\alpha(j-1), \alpha+1)t^{\alpha(2j-1)}}{(\Gamma(\alpha(j-2)+1))^2} \frac{j^{2\alpha}}{\rho_{j-1}^2 t^{2\alpha j}} \\ &= \frac{\Gamma(2\alpha(j-1))}{\Gamma(\alpha(2j-1)+1)} \frac{(\Gamma(\alpha(j-1)+1))^2 j^{2\alpha+1}}{(\Gamma(\alpha(j-2)+1))^2 C t^\alpha} \sim \frac{\alpha^{2\alpha}}{C 2^{\alpha+1}} \frac{j^{3\alpha}}{t^\alpha} \rightarrow 0, \quad t \rightarrow \infty. \end{aligned}$$

Further, using (35) and (15),

$$\begin{aligned} \max_{s \in [0, j]} D_{j-1}(s)U(j) &\leq (D_{j-1}(j) + (V_{j-1}(j))^2)U(j) \\ &\leq \left(\text{const} \frac{(C\Gamma(\alpha+1))^{2j-3} j^{2\alpha(j-1)}}{(\Gamma(\alpha(j-2)+1))^2} + r\rho_{j-1}^2 j^{2\alpha(j-1)} \right) U(j). \quad (37) \end{aligned}$$

Assumption A entails $U(j) \sim Cj^\alpha$ as $j \rightarrow \infty$. Now we show that the right-hand side of (37) times the squared normalization vanishes whenever $j = j(t) = o(t)$ and $j(t) \rightarrow \infty$ as $t \rightarrow \infty$. We start with the first term:

$$\begin{aligned} \frac{j^{2\alpha j} (C\Gamma(\alpha+1))^{2j-3}}{\rho_{j-1}^2 t^{2\alpha j} (\Gamma(\alpha(j-2)+1))^2} U(j) &= \frac{1}{C\Gamma(\alpha+1)} \left(\frac{j}{t} \right)^{2\alpha j} \frac{(\Gamma(\alpha(j-1)+1))^2}{(\Gamma(\alpha(j-2)+1))^2} U(j) \\ &\sim \frac{\alpha^{2\alpha}}{C\Gamma(\alpha+1)} \left(\frac{j}{t} \right)^{2\alpha j} j^{3\alpha} \rightarrow 0, \quad t \rightarrow \infty, \end{aligned}$$

because $(j/t)^{2\alpha j}$ converges to zero faster than any negative power of j . As for the second term in (37), arguing similarly we infer

$$\frac{j^{2\alpha j}}{t^{2\alpha j}} U(a_j) \sim \left(\frac{j}{t} \right)^{2\alpha j} Cj^\alpha \rightarrow 0, \quad t \rightarrow \infty.$$

An application of Markov's inequality completes the proof of Theorem 3. \square

We are ready to prove Theorem 1.

Proof of Theorem 1. Equality (1) is equivalent to

$$|\log P_k| = |\log W_1| + \dots + |\log W_{k-1}| + |\log(1 - W_k)|, \quad k \in \mathbb{N}.$$

Thus, $(|\log P_k|)_{k \geq 1}$ is a globally perturbed random walk generated by $(|\log W|, |\log(1 - W)|)$, that is, in the notation of Section 1.2, $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$.

For $j \in \mathbb{N}$ and $t > 0$, put $\psi_j(t) := \sum_{|v|=j} \mathbb{1}_{\{P(v) \geq 1/t\}}$ and observe that $\psi_j(n) = N_j(\log n)$ for $j, n \in \mathbb{N}$. We shall use a decomposition

$$\begin{aligned} K_n(\lfloor j_n u \rfloor) &= \left(K_n(\lfloor j_n u \rfloor) - \psi_{\lfloor j_n u \rfloor}(n) \right) + \left(\psi_{\lfloor j_n u \rfloor}(n) - \sum_{r \geq 1} V_{\lfloor j_n u \rfloor - 1}(\log n - T_r) \mathbb{1}_{\{T_r \leq \log n\}} \right) \\ &\quad + \sum_{r \geq 1} V_{\lfloor j_n u \rfloor - 1}(\log n - T_r) \mathbb{1}_{\{T_r \leq \log n\}} =: Y_1(n, u) + Y_2(n, u) + Y_3(n, u). \end{aligned}$$

It suffices to show that, as $n \rightarrow \infty$,

$$\frac{j_n^\alpha Y_i(n, u)}{\rho_{\lfloor j_n u \rfloor - 1}(\log n)^{\alpha \lfloor j_n u \rfloor}} \xrightarrow{\mathbb{P}} 0, \quad i = 1, 2, \quad (38)$$

where $\xrightarrow{\mathbb{P}}$ denotes convergence in probability, and

$$\left(\frac{c j_n^\alpha Y_3(n, u)}{\rho_{\lfloor j_n u \rfloor - 1}(\log n)^{\alpha \lfloor j_n u \rfloor}} \right)_{u > 0} \xrightarrow{\text{f.d.d.}} \left(\int_0^\infty e^{-\alpha y} d\mathcal{Z}_\alpha^-(y) \right)_{u > 0}.$$

We are going to apply Theorems 2 and 3 and Lemmas 5 and 6. While doing so, we replace t with $\log n$ and choose any diverging positive function $t \mapsto j(t)$ satisfying $j(\log n) = j_n$ and $j(t) = o(t^{\min(\frac{1}{3}, \frac{\alpha - \beta}{\alpha - \beta + 1})})$ as $t \rightarrow \infty$. By Lemma 1, Assumptions A and B entail (5). With this at hand, relation (38) with $i = 2$ follows from Theorem 3. The limit relation involving Y_3 is ensured by Theorem 2. In view of Markov's inequality, relation (38) with $i = 1$ follows if we can prove that, with $u > 0$ fixed,

$$\lim_{n \rightarrow \infty} \frac{j_n^\alpha \mathbb{E}|K_n(\lfloor j_n u \rfloor) - \psi_{\lfloor j_n u \rfloor}(n)|}{\rho_{\lfloor j_n u \rfloor - 1}(\log n)^{\alpha \lfloor j_n u \rfloor}} = 0.$$

In Section 6 of [9], see the top of p. 21, it was shown that

$$\mathbb{E}|K_n(\lfloor j_n u \rfloor) - \psi_{\lfloor j_n u \rfloor}(n)| \leq n \int_{(n, \infty)} x^{-1} d\mathbb{E}\psi_{\lfloor j_n u \rfloor}(x) + \int_{[1, n]} e^{-n/x} d\mathbb{E}\psi_{\lfloor j_n u \rfloor}(x).$$

By Lemma 6 applied in the particular setting $(\xi, \eta) = (|\log W|, |\log(1 - W)|)$, so that $V_{\lfloor j_n u \rfloor}(\log x) = \mathbb{E}\psi_{\lfloor j_n u \rfloor}(x)$ for $x \geq 1$, we obtain, as $n \rightarrow \infty$,

$$n \int_{(n, \infty)} x^{-1} d\mathbb{E}\psi_{\lfloor j_n u \rfloor}(x) = \int_{(\log n, \infty)} e^{\log n - x} dV_{\lfloor j_n u \rfloor}(x) = O(V_{\lfloor j_n u \rfloor - 1}(\log n))$$

for each fixed $u > 0$. The function f defined by $f(x) = \exp(-e^x)$ is decreasing and Lebesgue integrable on $[0, \infty)$. Hence, it is dRi on $[0, \infty)$, see, for instance, Lemma 6.2.1 (a) in [17]. By Lemma 5, with the so defined f , as $n \rightarrow \infty$,

$$\int_{[1, n]} e^{-n/x} d\mathbb{E}\psi_{\lfloor j_n u \rfloor}(x) = \int_{[0, \log n]} \exp(-e^{\log n - x}) dV_{\lfloor j_n u \rfloor}(x) = O(V_{\lfloor j_n u \rfloor - 1}(\log n))$$

for each fixed $u > 0$. Invoking (17) and the assumption on the growth rate of j_n which particularly entails $j_n = o(\log n)$ as $n \rightarrow \infty$ we infer

$$\frac{j_n^\alpha V_{[j_n u]-1}(\log n)}{\rho_{[j_n u]-1}(\log n)^{\alpha[j_n u]}} \sim \frac{j_n^\alpha}{(\log n)^\alpha} \rightarrow 0, \quad n \rightarrow \infty$$

thereby arriving at (38) with $i = 1$. The proof of Theorem 1 is complete. \square

6 Appendix

The following formula is rather standard. We give a proof for completeness.

Lemma 9. *Let $\gamma > 0$ and η be a positive random variable with the Laplace transform ℓ . Then*

$$\mathbb{E}\eta^{-\gamma} = \frac{\gamma}{\Gamma(1 + \gamma)} \int_0^\infty s^{\gamma-1} \ell(s) ds,$$

where Γ is the Euler gamma-function. Here, both sides of the equality may be infinite.

Proof. Let ζ be an exponentially distributed random variable of unit mean which is independent of η . Assuming that $\mathbb{E}\eta^{-\gamma} < \infty$, write

$$\mathbb{E}(\eta/\zeta)^{-\gamma} = \mathbb{E}\zeta^\gamma \mathbb{E}\eta^{-\gamma} = \Gamma(1 + \gamma) \mathbb{E}\eta^{-\gamma}.$$

On the other hand, using $\mathbb{P}\{\zeta > \eta/s\} = \ell(1/s)$ for $s > 0$ we infer

$$\mathbb{E}(\eta/\zeta)^{-\gamma} = \gamma \int_0^\infty s^{-\gamma-1} \mathbb{P}\{\eta/\zeta \leq s\} ds = \gamma \int_0^\infty s^{-\gamma-1} \ell(1/s) ds = \gamma \int_0^\infty s^{\gamma-1} \ell(s) ds.$$

\square

Finally, we give an estimate for the gamma-function.

Lemma 10. *For $x, y \geq 0$,*

$$\frac{\Gamma(x+1+y)}{\Gamma(x+1)} \leq \frac{11}{10} (x+1+y)^y. \quad (39)$$

Proof. Using the inequality given on p. 573 in [24]

$$1 < \frac{\Gamma(z)}{\sqrt{2\pi} z^{z-\frac{1}{2}} e^{-z}} < 1 + \frac{1}{12z} + \frac{1}{288z^2}, \quad z > 0,$$

we obtain

$$\begin{aligned} \frac{\Gamma(x+1+y)}{\Gamma(x+1)} &\leq \frac{(x+1+y)^{x+1+y-\frac{1}{2}} e^{-x-1-y} \left(1 + \frac{1}{12(x+1+y)} + \frac{1}{288(x+1+y)^2}\right)}{(x+1)^{x+1-\frac{1}{2}} e^{-x-1}} \\ &\leq \left(1 + \frac{y}{x+1}\right)^{x+1-\frac{1}{2}} (x+1+y)^y e^{-y} \left(1 + \frac{1}{12} + \frac{1}{288}\right) \leq \frac{11}{10} (x+1+y)^y. \end{aligned}$$

\square

References

- [1] G. Alsmeyer, A. Iksanov, and A. Marynych. Functional limit theorems for the number of occupied boxes in the Bernoulli sieve. *Stochastic Process. Appl.*, 127(3):995–1017, 2017.
- [2] B. Basrak, M. Conroy, M. Olvera-Cravioto, and Z. Palmowski. Importance sampling for maxima on trees. *Stoch. Proc. Appl.*, 148:139–179, 2022.
- [3] J. Bertoin. Asymptotic regimes for the occupancy scheme of multiplicative cascades. *Stochastic Process. Appl.*, 118(9):1586–1605, 2008.
- [4] P. Billingsley. *Convergence of probability measures*. Wiley, second edition, 1999.
- [5] N. Bingham, C. Goldie, and J. Teugels. *Regular variation*. Cambridge University Press, 1987.
- [6] V. Bohdanskyyi, V. Bohun, A. Marynych, and I. Samoilenko. Arithmetic properties of multiplicative integer-valued perturbed random walks. *Mod. Stoch. Appl.*, 11:133–148, 2024.
- [7] V. Bohun, A. Iksanov, A. Marynych, and B. Rashytov. Renewal theory for iterated perturbed random walks on a general branching process tree: intermediate generations. *J. Appl. Probab.*, 59(2):421–446, 2022.
- [8] O. Braganets and A. Iksanov. A limit theorem for a nested infinite occupancy scheme in random environment. *Austr. J. Statist.*, 52:1–12, 2023.
- [9] D. Buraczewski, B. Dvovgay, and A. Iksanov. On intermediate levels of nested occupancy scheme in random environment generated by stick-breaking I. *Electron. J. Probab.*, 25:1–24, 2020.
- [10] H. Carlsson and O. Nerman. An alternative proof of Lorden’s renewal inequality. *Adv. in Appl. Probab.*, 18(4):1015–1016, 1986.
- [11] J.-J. Duchamps, J. Pitman, and W. Tang. Renewal sequences and record chains related to multiple zeta sums. *Trans. Amer. Math. Soc.*, 371(8):5731–5755, 2019.
- [12] A. Gnedin and A. Iksanov. On nested infinite occupancy scheme in random environment. *Probab. Theory Related Fields*, 177:855–890, 2020.
- [13] A. Gnedin, A. Iksanov, and A. Marynych. The Bernoulli sieve: an overview. In *21st International Meeting on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms (AofA’10)*, Discrete Math. Theor. Comput. Sci. Proc., AM, pages 329–341. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2010.
- [14] A. V. Gnedin. The Bernoulli sieve. *Bernoulli*, 10(1):79–96, 2004.
- [15] J. Hawkes. A lower Lipschitz condition for the stable subordinator. *Z. Wahrscheinlichkeitstheorie verw. Geb.*, 17:23–32, 1971.

- [16] A. Iksanov. Functional limit theorems for renewal shot noise processes with increasing response functions. *Stoch. Proc. Appl.*, 123(6):1987–2010, 2013.
- [17] A. Iksanov. *Renewal theory for perturbed random walks and similar processes*. Birkhäuser/Springer, Cham, 2016.
- [18] A. Iksanov, W. Jedidi, and F. Bouzeffour. A law of the iterated logarithm for the number of occupied boxes in the Bernoulli sieve. *Statist. Probab. Lett.*, 126:244–252, 2017.
- [19] A. Iksanov, A. Marynych, and B. Rashytov. Stable fluctuations of iterated perturbed random walks in intermediate generations of a general branching process tree. *Lith. Math. J.*, 62(4):447–466, 2022.
- [20] A. Iksanov, A. Marynych, and I. Samoilenko. On intermediate levels of a nested occupancy scheme in a random environment generated by stick-breaking II. *Stochastics*, 94:1077–1101, 2022.
- [21] A. Iksanov, B. Rashytov, and I. Samoilenko. Renewal theory for iterated perturbed random walks on a general branching process tree: early generations. *J. Appl. Probab.*, 60(1):45–67, 2023.
- [22] A. Joseph. A phase transition for the heights of a fragmentation tree. *Random Structures Algorithms*, 39(2):247–274, 2011.
- [23] K. V. Mitov and E. Omev. *Renewal processes*. Springer, Cham, 2014.
- [24] G. Nemes. Error bounds and exponential improvements for the asymptotic expansions of the gamma function and its reciprocal. *Proc. Roy. Soc. Edinburgh Sect. A*, 145(3):571–596, 2015.
- [25] J. Pitman and W. Tang. Regenerative random permutations of integers. *Ann. Probab.*, 47(3):1378–1416, 2019.
- [26] J. Pitman and Y. Yakubovich. Extremes and gaps in sampling from a GEM random discrete distribution. *Electron. J. Probab.*, 22:Paper No. 44, 26, 2017.
- [27] J. Pitman and Y. Yakubovich. Gaps and interleaving of point processes in sampling from a residual allocation model. *Bernoulli*, 25(4B):3623–3651, 2019.
- [28] J. S. W. Wong and R. Wong. On asymptotic solutions of the renewal equation. *J. Math. Anal. Appl.*, 53:243–250, 1976.