(σ, τ) -Derivations of Number Rings with Coding Theory Applications

Praveen Manju and Rajendra Kumar Sharma

Department of Mathematics, Indian Institute of Technology Delhi, Hauz Khas, New Delhi-110016, India¹

Abstract

In this article, we study (σ, τ) -derivations of number rings by considering them as commutative unital Z-algebras. We begin by characterizing all (σ, τ) -derivations and inner (σ, τ) -derivations of the ring of algebraic integers of a quadratic number field. Then we characterize all (σ, τ) -derivations of the ring of algebraic integers $\mathbb{Z}[\zeta]$ of a p^{th} -cyclotomic number field $\mathbb{Q}(\zeta)$ (p odd rational prime and ζ a primitive p^{th} -root of unity). We also conjecture (using SAGE and MATLAB) an "if and only if" condition for a (σ, τ) -derivation D on $\mathbb{Z}[\zeta]$ to be inner. We further characterize all (σ, τ) -derivations and inner (σ, τ) -derivations of the bi-quadratic number ring $\mathbb{Z}[\sqrt{m}, \sqrt{n}]$ (m, n distinct square-free rational integers). In each of the above cases, we also determine the rank and an explicit basis of the derivation algebra consisting of all (σ, τ) -derivations of the number ring. Finally, we give the applications of the above work in coding theory by giving the notion of a Hom-IDD code.

Keywords: (σ, τ) -derivation, inner (σ, τ) -derivation, number ring, ring of algebraic integers, quadratic, cyclotomic, bi-quadratic

Mathematics Subject Classification (2010): Primary: 13N15, 11R04; Secondary: 11C20, 11R11, 11R16, 11R18.

1 Introduction

Let us denote a commutative ring with unity by R, an algebra over R by \mathcal{A} , and two different non-zero R-algebra endomorphisms of \mathcal{A} by σ , τ . An R-linear map $d : \mathcal{A} \to \mathcal{A}$ that satisfies $d(\alpha\beta) = d(\alpha)\beta + \alpha d(\beta)$ for every $\alpha, \beta \in \mathcal{A}$, is called a derivation on \mathcal{A} . It is called inner if $d(\alpha) = \beta\alpha - \alpha\beta$ for all $\alpha \in \mathcal{A}$, for some $\beta \in \mathcal{A}$. An R-linear map $D : \mathcal{A} \to \mathcal{A}$ that satisfies $D(\alpha\beta) = D(\alpha)\tau(\beta) + \sigma(\alpha)D(\beta)$ for all $\alpha, \beta \in \mathcal{A}$, is called a (σ, τ) -derivation on \mathcal{A} . Again, it is called inner if $D(\alpha) = \beta\tau(\alpha) - \sigma(\alpha)\beta$ for every $\alpha \in \mathcal{A}$, for some $\beta \in \mathcal{A}$.

¹*E-mail addresses:* praveenmanjuiitd@gmail.com(Corresponding Author: Praveen Manju), rksharmaiitd@gmail.com(Rajendra Kumar Sharma).

 $\mathcal{D}_{(\sigma,\tau)}(\mathcal{A})$ denotes the set of all (σ,τ) -derivations on \mathcal{A} . Defining componentwise sum and module action, $\mathcal{D}_{(\sigma,\tau)}(\mathcal{A})$ becomes an R- as well as \mathcal{A} -module. If 1 is the unity in \mathcal{A} , σ , τ are unital, that is, $\sigma(1) = \tau(1) = 1$ and D is a (σ,τ) -derivation of \mathcal{A} , then D(1) = 0.

We shall need some definitions and results from algebraic number theory for which we refer the reader to [13], [18]. Still, we summarize a few, as follows: A complex number is an algebraic number if it is a zero of some non-constant polynomial over \mathbb{Q} . A number field is a finite field extension of \mathbb{Q} that contains algebraic numbers. A number ring is any subring of a number field. An algebraic integer is a complex number that is a zero of some monic polynomial over \mathbb{Z} . The ring of algebraic integers of a number field K, denoted by O_K , is defined as $O_K = K \cap \mathbb{B}$, where \mathbb{B} is the ring of all algebraic integers. Thus O_K contains precisely all those algebraic integers which belong to K. A subring of O_K which is also a \mathbb{Z} -module of rank $[K:\mathbb{Q}]$ is called an order of K. A set of algebraic numbers $\{\alpha_1, \alpha_2, ..., \alpha_r\}$ is called an integral basis of K (or O_K) if each element in O_K is uniquely expressible as a \mathbb{Z} -linear combination of $\alpha_1, \alpha_2, ..., \alpha_r$. Every integral basis is a \mathbb{Q} -basis. An integral basis having form $\{1, \theta, ..., \theta^{n-1}\}$ for some $\theta \in O_K$, is called a power basis of K. A number field K that has a power basis is called monogenic. A number field K that has degree 2 over \mathbb{Q} is called quadratic. A number field that has the form $K = \mathbb{Q}(\zeta)$, where $\zeta = e^{2\pi i/m}$ $(m \ge 1)$ is a primitive complex m^{th} root of unity, is called an m^{th} cyclotomic field. We state below some results that we may need later.

Theorem 1.1 ([18, Theorem 2.2]). Let K be a number field. Then $K = \mathbb{Q}(\theta)$ for some algebraic number θ , or in other words, number fields are finite simple extensions of \mathbb{Q} . In fact, $K = \mathbb{Q}(\theta)$ for some algebraic integer θ .

Theorem 1.2 ([18, Theorem 2.16]). A number field K always has an integral basis. Further, the additive group of O_K is free abelian having rank equal to the degree of K.

Theorem 1.3 ([18, Proposition 3.1]). Every quadratic field is of the form $\mathbb{Q}(\sqrt{d})$ for some square-free rational integer d.

Theorem 1.4 ([18, Theorem 3.2]). Let $K = \mathbb{Q}(\sqrt{d})$ be a quadratic field. Then

- (i) $O_K = \mathbb{Z}[\sqrt{d}]$ when $d \not\equiv 1 \pmod{4}$,
- (ii) $O_K = \mathbb{Z}[\frac{1+\sqrt{d}}{2}]$ when $d \equiv 1 \pmod{4}$.

Theorem 1.5 ([18, Theorem 3.3]). Let $K = \mathbb{Q}(\sqrt{d})$ be a quadratic field. Then \mathcal{B} is an integral basis of K, where

- (a) $\mathcal{B} = \{1, \sqrt{d}\}$ when $d \not\equiv 1 \pmod{4}$.
- (b) $\mathcal{B} = \{1, \frac{1+\sqrt{d}}{2}\}$ when $d \equiv 1 \pmod{4}$.

Theorem 1.6 ([18, Lemma 3.4]). If p is an odd prime and ζ is a primitive p^{th} root of unity, then the minimal polynomial of ζ over \mathbb{Q} is $\phi_p(x) = x^{p-1} + x^{p-2} + \ldots + 1$. Hence $[\mathbb{Q}(\zeta) : Q] = p - 1$.

Theorem 1.7 ([18, Theorem 3.5]). Let p be an odd prime and $K = \mathbb{Q}(\zeta)$ be a p^{th} cyclotomic field. Then $O_K = \mathbb{Z}[\zeta]$. Hence, $\{1, \zeta, \zeta^2, ..., \zeta^{p-2}\}$ is an integral basis of $K = \mathbb{Q}(\zeta)$.

Theorem 1.8 ([13, Chapter 2]). Let $m \in \mathbb{N}$ and $K = \mathbb{Q}(\zeta)$ be an m^{th} cyclotomic field, then $O_K = \mathbb{Z}[\zeta]$.

Derivations have been studied in various fields like algebra, functional analysis, Fourier analysis, measure theory, operator theory, harmonic analysis, differential equations, differential geometry, etc. They play an essential role in many important branches of mathematics and physics. The notion of derivation in rings has been applied to and studied in various algebras to produce their theory of derivations. For example, BCI-algebras [15], von Neumann algebras [3], incline algebras which have many applications [11], MV-algebras [10], [16], Banach algebras [17], lattices that have a very crucial role in various fields like information theory: information recovery, management of information access and cryptanalysis [4]. Differentiable manifolds, operator algebras, \mathbb{C}^* -algebras, and representation theory of Lie groups are continued to be studied using derivations [12]. Derivations on rings and algebras help in the study of their structure. For a historical account and more applications of derivations, we refer the reader to [14], [2], [8]. The idea of a (s_1, s_2) -derivation was introduced by Jacobson [9]. These derivations were later on commonly called as (σ, τ) or (θ, ϕ) -derivation. These derivations have been highly studied in prime and semiprime rings and have been principally used in solving functional equations [3]. Derivations, and especially (σ, τ) -derivations of rings have various applications in coding theory [7], [6]. For more applications of (σ, τ) -derivations, we refer the reader to [1].

There is no literature available on twisted derivations of number rings except in $\begin{bmatrix} 5 \end{bmatrix}$ where the author studies (σ, τ) -derivations of the ring of algebraic integers of a quadratic number field. In this article, we mainly study (σ, τ) -derivations of number rings. We partition the article into three sections. In Section 2, we first obtain results on (σ, τ) -derivations and inner (σ, τ) -derivations of a commutative unital algebra \mathcal{A} over a commutative ring R with unity 1. We find a necessary condition for an R-linear map D on A to be a (σ, τ) -derivation and give counterexamples showing that the condition is not sufficient. This motivates us to study these derivations in certain number rings where this condition is both necessary as well as sufficient. We also determine a necessary and sufficient condition for a (σ, τ) -derivation on \mathcal{A} to be inner. In Section 3, we apply our results obtained in Section 2 to study (σ, τ) -derivations and inner (σ, τ) -derivations of number rings by considering them as commutative unital Zalgebras. Section 3 is further subdivided into three subsections. In Subsection 3.1, we study (σ, τ) -derivations and inner (σ, τ) -derivations of the ring of algebraic integers of a quadratic number field. We obtain necessary and sufficient conditions for a \mathbb{Z} -linear map D on the rings of algebraic integers of a quadratic field to be a (σ, τ) -derivation. We also classify the inner (σ, τ) -derivations of the ring of algebraic integers of a quadratic field. In Subsection 3.2, we study (σ, τ) -derivations and inner (σ, τ) -derivations of the ring of algebraic integers of a cyclotomic number field. We obtain necessary and sufficient conditions for a \mathbb{Z} -linear map D on the rings of algebraic integers $\mathbb{Z}[\zeta]$ of a p^{th} cyclotomic field $\mathbb{Q}(\zeta)$ to be a (σ, τ) -derivation. We prove that $\mathcal{D}_{(\sigma,\tau)}(\mathbb{Z}[\zeta])$ is a \mathbb{Z} -module having rank p-1. Further, in this process, we also propose two conjectures giving a necessary and sufficient condition on a (σ, τ) -derivation of $\mathbb{Z}[\zeta]$ to be inner. This is done with the help of SAGE and MATLAB. In Subsection 3.3, a beautiful application of Lemma 2.1 is given in determining all (σ, τ) -derivations of the number ring $\mathbb{Z}[\sqrt{m},\sqrt{n}]$ for every pair (σ,τ) of distinct ring endomorphisms σ and τ of $\mathbb{Z}[\sqrt{m}, \sqrt{n}]$. As a result, we establish that $D_{(\sigma,\tau)}(\mathbb{Z}[\sqrt{m}, \sqrt{n}])$ is a \mathbb{Z} -module having rank 4. Further, we obtain a necessary and sufficient condition for a (σ, τ) -derivation of $\mathbb{Z}[\sqrt{m}, \sqrt{n}]$ to be inner. We discuss the applications of our work in coding theory in Section 4 by giving the notion of a Hom-IDD code. In Section 5, we conclude our findings.

2 Some useful results on (σ, τ) -derivations of commutative algebras

 \mathcal{A} , in this section, denotes a commutative algebra with unity 1 over a commutative ring R having unity 1 and σ , τ are two different non-zero unital R-algebra endomorphisms of \mathcal{A} . First, we have an important lemma below.

Lemma 2.1. Let \mathcal{A} be of finite rank n as an R-module and let $\{\alpha_1, \alpha_2, ..., \alpha_n\}$ be an R-basis of \mathcal{A} . Then an R-linear map $D : \mathcal{A} \to \mathcal{A}$ is a (σ, τ) -derivation if and only if

$$D(\alpha_i \alpha_j) = D(\alpha_i)\tau(\alpha_j) + \sigma(\alpha_i)D(\alpha_j)$$

for every $i, j \in \{1, 2, ..., n\}$.

Proof. Let $D : \mathcal{A} \to \mathcal{A}$ be an *R*-linear map, and $x, y \in \mathcal{A}$. Then $x = \sum_{i=1}^{n} a_i \alpha_i$ and $y = \sum_{j=1}^{n} b_j \alpha_j$ for some $a_i, b_j \in R$ $(i, j \in \{1, 2, ..., n\})$. Observe that

$$xy = \left(\sum_{i=1}^{n} a_i \alpha_i\right) \left(\sum_{j=1}^{n} b_j \alpha_j\right) = \sum_{i,j} (a_i b_j)(\alpha_i \alpha_j),$$

and

$$D(xy) = \sum_{i,j=1}^{n} (a_i b_j) D(\alpha_i \alpha_j) = \sum_{i,j=1}^{n} (a_i b_j) (D(\alpha_i) \tau(\alpha_j) + \sigma(\alpha_i) D(\alpha_j))$$
$$= D\left(\sum_{i=1}^{n} a_i \alpha_i\right) \tau\left(\sum_{j=1}^{n} b_j \alpha_j\right) + \sigma\left(\sum_{i=1}^{n} a_i \alpha_i\right) D\left(\sum_{j=1}^{n} b_j \alpha_j\right)$$
$$= D(x)\tau(y) + \sigma(x)D(y).$$

Hence D is a (σ, τ) -derivation. The converse is straightforward.

For a non-negative integer k, define S_k as the set containing all ordered pairs (i, j) for which i + j = k. Note that the sets S_k 's are pairwise disjoint. Therefore, $|\bigcup_{i=0}^m S_i| = \sum_{i=0}^m |S_i| = 1 + 2 + 3 + \dots + m + (m+1) = \frac{(m+1)(m+2)}{2}$ for every non-negative integer m.

Lemma 2.2. For a (σ, τ) -derivation D of \mathcal{A} ,

$$D(\alpha^k) = \left(\sum_{(i,j)\in S_{k-1}} \sigma(\alpha^i)\tau(\alpha^j)\right) D(\alpha),$$

for every $\alpha \in \mathcal{A}$ and for every $k \in \mathbb{N}$.

Proof. We use induction. For k = 1, the equality trivially holds. Let the result hold for k = n, that is, $D(\alpha^n) = \left(\sum_{(i,j)\in S_{n-1}} \sigma(\alpha^i)\tau(\alpha^j)\right) D(\alpha)$. Then

$$D(\alpha^{n+1}) = D(\alpha^n \alpha) = D(\alpha^n)\tau(\alpha) + \sigma(\alpha^n)D(\alpha)$$

= $\left(\sum_{i+j=n-1} \sigma(\alpha^i)\tau(\alpha^{j+1}) + \sigma(\alpha^n)\tau(\alpha^0)\right)D(\alpha)$
= $\left(\sum_{i+j=n} \sigma(\alpha^i)\tau(\alpha^j)\right)D(\alpha) = \left(\sum_{(i,j)\in S_n} \sigma(\alpha^i)\tau(\alpha^j)\right)D(\alpha).$

Induction is complete and so is proof.

The theorem is now straightforward.

Theorem 2.3. Let \mathcal{A} be of finite rank n and suppose that \mathcal{A} has an R-basis of the form $\{1, \alpha, \alpha^2, ..., \alpha^{n-1}\}$ for some $\alpha \in \mathcal{A}$. If an R-linear map $D : \mathcal{A} \to \mathcal{A}$ is a (σ, τ) -derivation, then

$$D(\alpha^k) = \left(\sum_{(i,j)\in S_{k-1}} \sigma(\alpha^i)\tau(\alpha^j)\right) D(\alpha)$$
(2.1)

for all $k \in \{1, 2, ..., n-1\}$.

The converse of the Theorem 2.3 is not true. Some examples are given below.

Example 1. Let R[X] be the polynomial ring over a commutative unital ring R. Let $f(X) \in R[X]$ be monic with degree n and $\mathcal{A} = \frac{R[X]}{\langle f(X) \rangle}$. Then \mathcal{A} is a commutative R-algebra with unity $\overline{1} = 1 + f(X)$. Suppose $\alpha = X + f(X)$. Then $\{1, \alpha, \alpha^2, ..., \alpha^{n-1}\}$ is an R-basis of the R-module \mathcal{A} . We denote the zero element $0 + \langle f(X) \rangle$ of \mathcal{A} by $\overline{0}$.

In particular, take $R = \mathbb{Z}$ and $f(X) = X^4 - 1$ so that $\{\overline{1}, \alpha, \alpha^2, \alpha^3\}$ is a \mathbb{Z} -basis of $\mathcal{A} = \frac{R[X]}{\langle f(X) \rangle}$. Let σ and τ be \mathbb{Z} -algebra endomorphisms of \mathcal{A} given by $\sigma(\alpha) = \alpha$ and $\tau(\alpha) = \alpha^2$. Then $\sigma(\overline{1}) = \overline{1}$ and $\tau(\overline{1}) = \overline{1}$. Define $D : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with $D(\overline{1}) = \overline{0}$ and

$$D(\alpha^{r}) = \left(\sum_{(i,j)\in S_{r-1}} \sigma(\alpha^{i})\tau(\alpha^{j})\right) D(\alpha), \,\forall r \in \{1,2,3\}$$

 $\alpha^4 = \overline{1}$ so that $D(\alpha^4) = \overline{0}$. Also,

$$\left(\sum_{(i,j)\in S_3}\sigma(\alpha^i)\tau(\alpha^j)\right)D(\alpha) = \left(\sigma(\alpha^3) + \sigma(\alpha^2)\tau(\alpha) + \sigma(\alpha)\tau(\alpha^2) + \tau(\alpha^3)\right)D(\alpha)$$
$$= \left(\alpha^3 + \alpha^4 + \alpha^5 + \alpha^6\right)D(\alpha) = \left(\bar{1} + \alpha + \alpha^2 + \alpha^3\right)D(\alpha).$$

Now $\overline{1} + \alpha + \alpha^2 + \alpha^3 \neq \overline{0}$ since $\{\overline{1}, \alpha, \alpha^2, \alpha^3\}$ is a basis of \mathcal{A} . Also, \mathcal{A} is an integral domain, so if $D(\alpha) \neq 0$, then the above expression cannot be zero. Further, observe that $D(\alpha^4) \neq \left(\sum_{(i,j)\in S_3} \sigma(\alpha^i)\tau(\alpha^j)\right) D(\alpha)$. Hence by Lemma 2.2, D cannot be a (σ, τ) -derivation of \mathcal{A} . Example 2. Consider the set

$$\mathcal{A} = \{ aI + bA + cA^2 \mid a, b, c \in \mathbb{Z} \},\$$

where $A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$. Then \mathcal{A} is a commutative subalgebra of the \mathbb{Z} -algebra $M_3(\mathbb{Z})$ with

unity I and $\{I, A, A^2\}$ is a \mathbb{Z} -basis of \mathcal{A} . Let σ and τ be \mathbb{Z} -algebra endomorphisms of \mathcal{A} given by $\sigma(aI + bA + cA^2) = aI + bA + cA^2$ and $\tau(aI + bA + cA^2) = aI + bA^2 + cA$ for all $a, b, c \in \mathbb{Z}$. Then $\sigma(I) = \tau(I) = I$. Define $D : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(I) = 0 and such that

$$D(A^r) = \left(\sum_{(i,j)\in S_{r-1}} \sigma(A^i)\tau(A^j)\right) D(A), \quad \text{for } r = 1, 2.$$

Then $D(A^3) = D(I) = 0$. Also,

$$\left(\sum_{(i,j)\in S_2}\sigma(A^i)\tau(A^j)\right)D(A) = \left(\sigma(A^2) + \sigma(A)\tau(A) + \tau(A^2)\right)D(A) = \left(I + A + A^2\right)D(A).$$

Again, $I + A + A^2 \neq 0$, so if $D(A) \in \mathcal{A}$ is such that $(I + A + A^2) D(A) \neq 0$, for example, D(A) = A or A^2 , then $D(A^3) \neq \left(\sum_{(i,j)\in S_2} \sigma(A^i)\tau(A^j)\right) D(A)$. Hence, D is not a (σ, τ) -derivation.

Example 3. Let $n \in \mathbb{N}$ and $A \in M_{n \times n}(\mathbb{Z})$ be an idempotent matrix. Then the subset

 $\mathcal{A} = \{ aI + bA \mid a, b \in \mathbb{Z} \}$

of $M_{n \times n}(\mathbb{Z})$ is a commutative \mathbb{Z} -algebra with unity I and $\{I, A\}$ is a basis of \mathcal{A} . Let σ and τ be \mathbb{Z} -algebra endomorphisms of \mathcal{A} given by $\sigma(aI + bA) = (a + b)I$ and $\tau(aI + bA) = aI + bA$ for all $a, b \in \mathbb{Z}$. Define $D : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(I) = 0 and such that

$$D(A^r) = \left(\sum_{(i,j)\in S_{r-1}} \sigma(A^i)\tau(A^j)\right) D(A), \quad \text{for } r = 1.$$
$$\left(\sum_{(i,j)\in S_1} \sigma(A^i)\tau(A^j)\right) D(A) = (\sigma(A) + \tau(A)) D(A) = (I+A) D(A).$$

 $D(A) \neq (I + A) D(A)$ provided $AD(A) \neq 0$. Hence, D is not a (σ, τ) -derivation of \mathcal{A} provided D(A) is chosen in $M_{n \times n}(\mathbb{Z})$ such that $AD(A) \neq 0$.

Example 4. Let $n \in \mathbb{N}$ and $A \in M_{n \times n}(\mathbb{Z})$ be a nilpotent matrix, that is, $A^k = 0$ for some least positive integer k. Then the subset

$$\mathcal{A} = \{ \sum_{i=0}^{k-1} a_i A^i \mid a_i \in \mathbb{Z}, \, \forall \, i \in \{0, 1, ..., k-1\} \}$$

of $M_{n \times n}(\mathbb{Z})$ is a commutative \mathbb{Z} -algebra with unity I and $\{A^0 = I, A, ..., A^{k-1}\}$ forms a basis of the \mathbb{Z} -module \mathcal{A} . Let σ and τ be \mathbb{Z} -algebra endomorphisms of \mathcal{A} given by $\sigma(\sum_{i=0}^{k-1} a_i A^i) = \sum_{i=0}^{k-1} a_i A^i$ and $\tau(\sum_{i=0}^{k-1} a_i A^i) = \sum_{i=0}^{k-1} a_i B^i$ for all $a_i \in \mathbb{Z}$ $(0 \le i \le k-1)$, where B = mA for some $m \in \mathbb{N}$ and $m \ne 1$. Define $D : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(I) = 0 and such that

$$D(A^{r}) = \left(\sum_{(i,j)\in S_{r-1}} \sigma(A^{i})\tau(A^{j})\right) D(A), \quad \text{for } r \in \{1, ..., k-1\}.$$

Then $D(A^k) = 0$. Also,

$$\left(\sum_{(i,j)\in S_{k-1}}\sigma(A^i)\tau(A^j)\right)D(A) = \left(\sum_{i+j=k-1}A^iB^j\right)D(A) = \left(\sum_{j=0}^{k-1}m^jA^{k-1}\right)D(A)$$
$$= \left(\frac{1-m^k}{1-m}\right)A^{k-1}D(A).$$

 $0 \neq \left(\frac{1-m^k}{1-m}\right) A^{k-1}D(A)$ provided $A^{k-1}D(A) \neq 0$. Hence, D cannot be a (σ, τ) -derivation of \mathcal{A} provided D(A) is chosen in $M_{n \times n}(\mathbb{Z})$ such that $A^{k-1}D(A) \neq 0$.

If R is a ring and G is a group, then the group ring of G over R is defined as the set

$$RG = \{ \sum_{g \in G} a_g g \mid a_g \in R, \forall g \in G \text{ and } |\operatorname{supp}(\alpha)| < \infty \},\$$

where for $\alpha = \sum_{g \in G} a_g g$, $\operatorname{supp}(\alpha)$ is the support of α that consists of elements from G that appear in the expression of α . RG is a ring with respect to the componentwise addition and multiplication defined respectively by: For $\alpha = \sum_{g \in G} a_g g$, $\beta = \sum_{g \in G} b_g g$ in RG,

$$(\sum_{g\in G} a_g g) + (\sum_{g\in G} b_g g) = \sum_{g\in G} (a_g + b_g)g \text{ and } \alpha\beta = \sum_{g,h\in G} a_g b_h gh.$$

If the ring R is commutative having unity 1 and the group G is abelian having identity e, then RG becomes a commutative unital algebra over R with unity 1 = 1e. If σ and τ are algebra endomorphisms of RG which are R-linear extensions of group homomorphisms of Gand $D: RG \to RG$ is a (σ, τ) -derivation, then

$$D(e) = D(ee) = D(e)\tau(e) + D(e)\sigma(e) = D(e) + D(e)$$

so that D(e) = 0.

Example 5. Let $C_n = \langle g \mid g^n = 1 \rangle$ be a cyclic group having order $n \ge 2$ and consider the group ring $\mathbb{Z}C_n$. Let σ and τ be \mathbb{Z} -algebra endomorphisms of $\mathbb{Z}G$ which are \mathbb{Z} -linear extensions of group homomorphisms of G, say, $\sigma(g) = 1$ and $\tau(g) = g$. Let $D : \mathbb{Z}C_n \to \mathbb{Z}C_n$ be a \mathbb{Z} -linear map with D(1) = 0 and

$$D(g^{r}) = \left(\sum_{(i,j)\in S_{r-1}} \sigma(g^{i})\tau(g^{j})\right) D(g), \,\forall r \in \{1, ..., n-1\}.$$

So $D(g^n) = D(1) = 0$. Also,

$$\left(\sum_{(i,j)\in S_{n-1}}\sigma(g^i)\tau(g^j)\right)D(g) = (\sigma(g^{n-1}) + \sigma(g^{n-2})\tau(g) + \sigma(g^{n-3})\tau(g^2) + \dots + \sigma(g^2)\tau(g^{n-3}) + \sigma(g)\tau(g^{n-2}) + \tau(g^{n-1}))D(g)$$
$$= \left(e + g + \dots + g^{n-1}\right)D(g)$$

Note that $e + g + ... + g^{n-1} \neq 0$ since $e, g, ..., g^{n-1}$ are linearly independent being basis elements of $\mathbb{Z}C_n$. So if $D(g) \in \mathbb{Z}C_n$ is such that $(e + g + ... + g^{n-1}) D(g) \neq 0$, then $D(g^n) \neq (\sum_{(i,j)\in S_{n-1}} \sigma(g^i)\tau(g^j)) D(g)$. Hence, D cannot be a (σ, τ) -derivation of $\mathbb{Z}C_n$.

Lemma 2.4. Let \mathcal{A} be of finite rank n and $\{\alpha_1, \alpha_2, ..., \alpha_n\}$ be an R-basis of \mathcal{A} . Let $D : \mathcal{A} \to \mathcal{A}$ be a (σ, τ) -derivation. Suppose that there exists some $\beta \in \mathcal{A}$ such that $D(\alpha_i) = \beta(\tau - \sigma)(\alpha_i)$ for all $i \in \{1, 2, ..., n\}$. Then $D(\alpha) = \beta(\tau - \sigma)(\alpha)$ for all $\alpha \in \mathcal{A}$.

Proof. For $\alpha = \sum_{i=1}^{n} a_i \alpha_i \in \mathcal{A}$,

$$D(\alpha) = \sum_{i=1}^{n} a_i D(\alpha_i) = \sum_{i=1}^{n} a_i \beta(\tau - \sigma)(\alpha_i) = \beta \left(\sum_{i=1}^{n} a_i \tau(\alpha_i) - \sum_{i=1}^{n} a_i \sigma(\alpha_i) \right)$$
$$= \beta \left(\tau \left(\sum_{i=1}^{n} a_i \alpha_i \right) - \sigma \left(\sum_{i=1}^{n} a_i \alpha_i \right) \right)$$
$$= \beta(\tau - \sigma)(\alpha)$$

Since $\alpha \in \mathcal{A}$ is arbitrary, therefore, D is inner.

The theorem now follows immediately.

Theorem 2.5. Let \mathcal{A} be of finite rank n and $\{\alpha_1, \alpha_2, ..., \alpha_n\}$ be an R-basis of \mathcal{A} . Then $a \ (\sigma, \tau)$ -derivation $D : \mathcal{A} \to \mathcal{A}$ is inner if and only if there exists some $\beta \in \mathcal{A}$ such that $D(\alpha_i) = \beta(\tau - \sigma)(\alpha_i)$ for all $i \in \{1, 2, ..., n\}$.

Lemma 2.6. Let $D : \mathcal{A} \to \mathcal{A}$ be a (σ, τ) -derivation and $\alpha \in \mathcal{A}$. If there exists some $\beta \in \mathcal{A}$ such that $D(\alpha) = \beta(\tau - \sigma)(\alpha)$, then

$$D(\alpha^n) = \beta(\tau - \sigma)(\alpha^n)$$

for all $n \in \mathbb{N}$.

Proof. We again use induction on n. If n = 1, the result holds trivially. Now suppose that $D(\alpha^k) = \beta(\tau - \sigma)(\alpha^k)$. Then

$$D(\alpha^{k+1}) = D(\alpha^k)\tau(\alpha) + \sigma(\alpha^k)D(\alpha) = \beta(\tau - \sigma)(\alpha^k)\tau(\alpha) + \sigma(\alpha^k)\beta(\tau - \sigma)(\alpha)$$

= $\beta \left(\tau(\alpha^{k+1}) - \sigma(\alpha^k)\tau(\alpha) + \sigma(\alpha^k)\tau(\alpha) - \sigma(\alpha^{k+1})\right)$
= $\beta(\tau - \sigma)(\alpha^{k+1})$

Induction is complete and so is proof.

Lemma 2.7. Let \mathcal{A} be of finite rank n and suppose that \mathcal{A} has an R-basis of the form $\{1, \alpha, \alpha^2, ..., \alpha^{n-1}\}$ for some $\alpha \in \mathcal{A}$. Let $D : \mathcal{A} \to \mathcal{A}$ be a (σ, τ) -derivation. If there exists some $\beta \in \mathcal{A}$ such that $D(\alpha) = \beta(\tau - \sigma)(\alpha)$, then D is inner.

Proof. Since D(1) = 0 and $\sigma(1) = \tau(1) = 1$, so $D(1) = \beta(\tau - \sigma)(1)$. Further since $D(\alpha) = \beta(\tau - \sigma)(\alpha)$, therefore, by Lemma 2.6,

$$D(\alpha^k) = \beta(\tau - \sigma)(\alpha^k)$$

for all $k \in \mathbb{N}$. So $D(\alpha^k) = \beta(\tau - \sigma)(\alpha^k)$ for every $k \in \{0, 1, ..., n - 1\}$. Now we use Lemma 2.4.

The theorem below follows immediately.

Theorem 2.8. Let \mathcal{A} be of finite rank n and suppose that \mathcal{A} has an R-basis of the form $\{1, \alpha, \alpha^2, ..., \alpha^{n-1}\}$ for some $\alpha \in \mathcal{A}$. Then a (σ, τ) -derivation $D : \mathcal{A} \to \mathcal{A}$ is inner if and only if there exists some $\beta \in \mathcal{A}$ such that $D(\alpha) = \beta(\tau - \sigma)(\alpha)$.

3 (σ, τ) -derivations of number rings

We know by Theorem 1.1 that if K is a number field, then $K = \mathbb{Q}(\theta)$ for some algebraic integer θ . Theorem 2.3 gives the lemma below:

Lemma 3.1. Let $K = \mathbb{Q}(\theta)$ (θ an algebraic integer) be a number field of degree n such that $\{1, \theta, ..., \theta^{n-1}\}$ is a power basis of K. Let σ and τ be two different non-zero \mathbb{Z} -algebra endomorphisms of O_K . If a \mathbb{Z} -linear map $D: O_K \to O_K$ is a (σ, τ) -derivation, then

$$D(\theta^k) = \left(\sum_{(i,j)\in S_{k-1}} \sigma(\theta^i) \tau(\theta^j)\right) D(\theta)$$

for all $k \in \{1, 2, ..., n-1\}$.

3.1 (σ, τ) -derivations of quadratic number rings

By Theorem 1.3, if K is a quadratic field, then $K = \mathbb{Q}(\sqrt{d})$ for some square-free rational integer d. In this section, we take $K = \mathbb{Q}(\sqrt{d})$ and σ , τ as two different non-zero ring endomorphisms of O_K . Note that any ring endomorphism of the ring O_K is a unital \mathbb{Z} algebra endomorphism of the unital \mathbb{Z} -algebra O_K . As an application of Lemma 2.1, we present a different proof of a result of [5] stated as follows.

Theorem 3.2 ([5, Theorem 4.2]). Every \mathbb{Z} -linear map $D: O_K \to O_K$ with D(1) = 0 is a (σ, τ) -derivation of O_K .

Proof. Only two possibilities arise: $d \not\equiv 1 \pmod{4}$ and $d \equiv 1 \pmod{4}$. Since $(\sqrt{d})^2 = d$, so if θ is a non-zero ring endomorphism of O_K , then $(\theta(\sqrt{d}))^2 = d$ or that $\theta(\sqrt{d}) = \pm \sqrt{d}$.

When $d \not\equiv 1 \pmod{4}$, then from Theorem 1.4, $O_K = \mathbb{Z}[\sqrt{d}]$ and from Theorem 1.5, $\{1, \sqrt{d}\}$ is a \mathbb{Z} -basis of O_K . In this case, exactly two possibilities arise. For every $a, b \in \mathbb{Z}$,

- (a) $\sigma(a + b\sqrt{d}) = a + b\sqrt{d}; \tau(a + b\sqrt{d}) = a b\sqrt{d},$
- (b) $\sigma(a+b\sqrt{d}) = a b\sqrt{d}; \tau(a+b\sqrt{d}) = a + b\sqrt{d}.$

When $d \equiv 1 \pmod{4}$, then by Theorem 1.4, $O_K = \mathbb{Z}[\frac{1+\sqrt{d}}{2}]$ and by Theorem 1.5, $\{1, \frac{1+\sqrt{d}}{2}\}$ is a \mathbb{Z} -basis of O_K . Thus in this case, too, exactly two possibilities arise. For all $a, b \in \mathbb{Z}$,

(a)
$$\sigma(a + b(\frac{1+\sqrt{d}}{2})) = a + b(\frac{1+\sqrt{d}}{2}); \tau(a + b(\frac{1+\sqrt{d}}{2})) = a + b(\frac{1-\sqrt{d}}{2}),$$

(b) $\sigma(a + b(\frac{1+\sqrt{d}}{2})) = a + b(\frac{1-\sqrt{d}}{2}); \tau(a + b(\frac{1+\sqrt{d}}{2})) = a + b(\frac{1+\sqrt{d}}{2}).$

Now the proof trivially follows from Lemma 2.1.

Further, in the same paper [5], the author has proved the theorem stated below.

Theorem 3.3 ([5, Theorem 4.2]). Let $d \not\equiv 1 \pmod{4}$. Let $D : O_K \to O_K$ be a (σ, τ) -derivation and $D(\sqrt{d}) = c_0 + c_1\sqrt{d}$ for some $c_0, c_1 \in \mathbb{Z}$. If 2d divides c_0 and c_1 is even, then D is inner.

We present a different proof of this theorem using Theorem 2.8 and also prove that the conditions in the above theorem are necessary.

Theorem 3.4. Let $d \not\equiv 1 \pmod{4}$. Let $D : O_K \to O_K$ be a (σ, τ) -derivation and $D(\sqrt{d}) = c_0 + c_1\sqrt{d}$ for some $c_0, c_1 \in \mathbb{Z}$. Then D is inner if and only if c_0 is divisible by 2d and c_1 is even. In particular, O_K has non-trivial outer (σ, τ) -derivations.

Proof. Since $d \not\equiv 1 \pmod{4}$, so $\{1, \sqrt{d}\}$ is a basis of the \mathbb{Z} -module $O_K = \mathbb{Z}[\sqrt{d}]$.

 $O_K = \mathbb{Z}[\sqrt{d}]$ has precisely two different non-zero ring endomorphisms: $\phi_1(a + b\sqrt{d})$ = $a + b\sqrt{d}$ and $\phi_2(a + b\sqrt{d}) = a - b\sqrt{d}$ $(a, b \in \mathbb{Z})$. Therefore, $(\sigma, \tau) = (\phi_1, \phi_2)$ or (ϕ_2, ϕ_1) . We prove the result for $(\sigma, \tau) = (\phi_2, \phi_1)$ since the other follows similarly.

First, let D be inner. Then there exists some $\beta = b_0 + b_1 \sqrt{d}$ for some $b_0, b_1 \in \mathbb{Z}$ such that $D(\sqrt{d}) = \beta(\tau - \sigma)(\sqrt{d})$. Therefore,

$$c_0 + c_1\sqrt{d} = D(\sqrt{d}) = \beta(\tau - \sigma)(\sqrt{d})$$
$$= (b_0 + b_1\sqrt{d})(\tau(\sqrt{d}) - \sigma(\sqrt{d}))$$
$$= 2db_1 + 2b_0\sqrt{d}$$

Since $\{1, \sqrt{d}\}$ is a basis of the \mathbb{Z} -module $O_K = \mathbb{Z}[\sqrt{d}]$, therefore,

$$2db_1 = c_0$$
 and $2b_0 = c_1$.

Obviously, $b_0 = \frac{c_1}{2}$ and $b_1 = \frac{c_0}{2d}$. Since $b_0, b_1 \in \mathbb{Z}$, therefore, c_1 is divisible by 2 and c_0 is divisible by 2d.

Conversely, let c_0 be divisible by 2d and c_1 be even. Define $\beta = \left(\frac{c_0}{2d}\right) + \left(\frac{c_1}{2}\right)\sqrt{d}$. Then it can be verified that $\beta(\tau - \sigma)(\sqrt{d}) = D(\sqrt{d})$ as $D(\sqrt{d}) = c_0 + c_1\sqrt{d}$. Now the result follows immediately by Theorem 2.8.

Theorem 3.5. Let $d \equiv 1 \pmod{4}$. Let $D: O_K \to O_K$ be a (σ, τ) -derivation and $D(\sqrt{d}) = c_0 + c_1(\frac{1+\sqrt{d}}{2})$ for some $c_0, c_1 \in \mathbb{Z}$. Then D is inner if and only if d divides $-c_0 + c_1(\frac{d-1}{2})$ and $2c_0 + c_1$. In particular, the following conditions hold:

- (i) O_K has non-trivial outer (σ, τ) -derivations.
- (ii) D is inner if d divides both c_0 and c_1 .

Proof. Since $d \equiv 1 \pmod{4}$, so $\{1, \frac{1+\sqrt{d}}{2}\}$ is a basis of the \mathbb{Z} -module $O_K = \mathbb{Z}[\frac{1+\sqrt{d}}{2}]$. O_K has precisely two different non-zero ring endomorphisms, namely, $\phi_1(a+b(\frac{1+\sqrt{d}}{2})) = a+b(\frac{1+\sqrt{d}}{2})$ and $\phi_2(a+b(\frac{1+\sqrt{d}}{2})) = a+b(\frac{1-\sqrt{d}}{2})$ $(a,b\in\mathbb{Z})$. Therefore, $(\sigma,\tau) = (\phi_1,\phi_2)$ or (ϕ_2,ϕ_1) . We prove the result for $(\sigma,\tau) = (\phi_2,\phi_1)$ since the other follows similarly.

First, let *D* be inner. Then there exists some $\beta = b_0 + b_1(\frac{1+\sqrt{d}}{2})$ for some $b_0, b_1 \in \mathbb{Z}$ such that $D(\frac{1+\sqrt{d}}{2}) = \beta(\tau - \sigma)(\frac{1+\sqrt{d}}{2})$. Therefore,

$$c_{0} + c_{1}\left(\frac{1+\sqrt{d}}{2}\right) = D\left(\frac{1+\sqrt{d}}{2}\right) = \beta\left(\tau-\sigma\right)\left(\frac{1+\sqrt{d}}{2}\right)$$
$$= \left(b_{0} + b_{1}\left(\frac{1+\sqrt{d}}{2}\right)\right)\left(\tau\left(\frac{1+\sqrt{d}}{2}\right) - \sigma\left(\frac{1+\sqrt{d}}{2}\right)\right)$$
$$= \left(b_{0} + b_{1}\left(\frac{1+\sqrt{d}}{2}\right)\right)\left(\left(\frac{1+\sqrt{d}}{2}\right) - \left(\frac{1-\sqrt{d}}{2}\right)\right)$$
$$= \left(-b_{0} + b_{1}\left(\frac{d-1}{2}\right)\right) + (2b_{0} + b_{1})\left(\frac{1+\sqrt{d}}{2}\right)$$

Since $\{1, \frac{1+\sqrt{d}}{2}\}$ is a basis of $O_K = \mathbb{Z}[\frac{1+\sqrt{d}}{2}]$, therefore,

$$-b_0 + b_1\left(\frac{d-1}{2}\right) = c_0$$
 and $2b_0 + b_1 = c_1$.

Note that since $d \equiv 1 \pmod{4}$, so $\frac{d-1}{2} \in \mathbb{Z}$. Solving, we get,

$$b_0 = \frac{1}{d} \left(-c_0 + c_1 \left(\frac{d-1}{2} \right) \right)$$
 and $b_1 = \frac{1}{d} \left(2c_0 + c_1 \right)$.

Since $b_0, b_1 \in \mathbb{Z}$, therefore, d divides $-c_0 + c_1\left(\frac{d-1}{2}\right)$ and $2c_0 + c_1$. Conversely, let d divide $-c_0 + c_1\left(\frac{d-1}{2}\right)$ and $2c_0 + c_1$. Define

$$\beta = \left(\frac{1}{d}\left(-c_0 + c_1\left(\frac{d-1}{2}\right)\right)\right) + \left(\frac{1}{d}\left(2c_0 + c_1\right)\right)\left(\frac{1+\sqrt{d}}{2}\right).$$

Then it can be verified that $\beta(\tau - \sigma)\left(\frac{1+\sqrt{d}}{2}\right) = D\left(\frac{1+\sqrt{d}}{2}\right)$ as $D\left(\frac{1+\sqrt{d}}{2}\right) = c_0 + c_1\left(\frac{1+\sqrt{d}}{2}\right)$. The result now follows immediately by Theorem 2.8.

3.2 (σ, τ) -derivations of cyclotomic number rings

In this section, $K = \mathbb{Q}(\zeta)$ denotes a p^{th} cyclotomic field, where p is an odd rational prime. By Theorem 1.7, the ring of algebraic integers of K is $O_K = \mathbb{Z}[\zeta]$ and $\{1, \zeta, \zeta^2, ..., \zeta^{p-2}\}$ is an integral basis of K. Further, σ and τ denote any two different non-zero ring endomorphisms of $O_K = \mathbb{Z}[\zeta]$.

Lemma 3.6. Let for each $k \in \{0, 1, ..., p-2\}$, S_k 's be sets as defined in Section 2. Then

$$\sum_{(i,j)\in \bigcup_{k=0}^{p-2}S_k}\sigma(\zeta^i)\tau(\zeta^j)=0.$$

Proof. All the non-zero ring endomorphisms ϕ_i of O_K are given by $\phi_i(\zeta) = \zeta^i$, where $i \in \{1, 2, ..., p-1\}$. Since σ and τ are non-zero and different, therefore,

$$\sigma(\zeta) = \zeta^u$$
 and $\tau(\zeta) = \zeta^{u+v}$

for some $u, v \in \{1, 2, ..., p-1\}$ such that $u + v \neq p$. Now,

(

$$\sum_{i,j)\in \cup_{k=0}^{p-2}S_k}\sigma(\zeta^i)\tau(\zeta^j) = \sum_{k=0}^{p-2}\sum_{(i,j)\in S_k}\sigma(\zeta^i)\tau(\zeta^j).$$

For any $k \in \{0, 1, ..., p - 2\},\$

$$\sum_{(i,j)\in S_k} \sigma(\zeta^i)\tau(\zeta^j) = \sum_{(i,j)\in S_k} \zeta^{ku}\zeta^{jv}.$$

There are three observations for each $k \in \{0, 1, ..., p-2\}$:

- (a) The sum $\sum_{(i,j)\in S_k} \sigma(\zeta^i)\tau(\zeta^j)$ contains k+1 terms and all these k+1 terms are different from each other since $u+v\neq p$.
- (b) The sum $\sum_{(i,j)\in \bigcup_{k=0}^{p-2}S_k} \sigma(\zeta^i)\tau(\zeta^j)$ contains exactly $\frac{p(p-1)}{2}$ terms.
- (c) Since the prime p is odd, therefore, for each $k \in \{1, 2, ..., p-1\}, \zeta^k$ is a primitive p^{th} root of unity.

These together imply that each term from the set $\{1, \zeta, \zeta^2, ..., \zeta^{p-1}\}$ is being repeated $\frac{p-1}{2}$ times in the sum $\sum_{(i,j)\in \cup_{k=0}^{p-2}S_k} \sigma(\zeta^i)\tau(\zeta^j)$. Hence,

$$\sum_{(i,j)\in \bigcup_{m=0}^{p-2}S_m} \sigma(\zeta^i)\tau(\zeta^j) = \frac{p-1}{2} \left(1+\zeta+\zeta^2+\ldots+\zeta^{p-1}\right) = 0,$$

since $1 + \zeta + \zeta^2 + \ldots + \zeta^{p-1} = 0$ by Theorem 1.6.

This proves the required result.

Lemma 3.7. Let $D: O_K \to O_K$ be a \mathbb{Z} -linear map with D(1) = 0 and

$$D(\zeta^k) = \left(\sum_{(i,j)\in S_{k-1}} \sigma(\zeta^i)\tau(\zeta^j)\right) D(\zeta)$$
(3.1)

for all $k \in \{1, 2, ..., p-2\}$. Then D is a (σ, τ) -derivation.

Proof. According to Lemma 2.1, the result can be concluded by establishing that

$$D(\zeta^{i+j}) = D(\zeta^i)\tau(\zeta^j) + \sigma(\zeta^i)D(\zeta^j)$$
(3.2)

for all $i, j \in \{0, 1, ..., p - 2\}$.

The relations (3.2) hold trivially when at least one of i or j is 0, using the fact that $\sigma(1) = \tau(1) = 1$ and D(1) = 0. So now let $i, j \in \{1, 2, ..., p-2\}$. Using (3.1), we get:

$$D(\zeta^{i})\tau(\zeta^{j}) = \left(\sum_{(s,t)\in S_{i-1}}\sigma(\zeta^{s})\tau(\zeta^{t})\right)D(\zeta)\tau(\zeta^{j}) = \left(\sum_{(s,t)\in S_{i-1}}\sigma(\zeta^{s})\tau(\zeta^{t+j})\right)D(\zeta)$$
$$= \left(\sum_{s=0}^{i-1}\sigma(\zeta^{s})\tau(\zeta^{i-1-s+j})\right)D(\zeta)$$

and

$$\begin{aligned} \sigma(\zeta^{i})D(\zeta^{j}) &= \sigma(\zeta^{i}) \left(\sum_{(s,t)\in S_{j-1}} \sigma(\zeta^{s})\tau(\zeta^{t}) \right) D(\zeta) &= \left(\sum_{(s,t)\in S_{j-1}} \sigma(\zeta^{i+s})\tau(\zeta^{t}) \right) D(\zeta) \\ &= \left(\sum_{s=0}^{j-1} \sigma(\zeta^{i+s})\tau(\zeta^{j-1-s}) \right) D(\zeta) \\ &= \left(\sum_{s=i}^{i+j-1} \sigma(\zeta^{s})\tau(\zeta^{j-1+i-s}) \right) D(\zeta). \end{aligned}$$

Therefore,

$$D(\zeta^{i})\tau(\zeta^{j}) + \sigma(\zeta^{i})D(\zeta^{j}) = \left(\sum_{(s,t)\in S_{i+j-1}}\sigma(\zeta^{s})\tau(\zeta^{t})\right)D(\zeta).$$
(3.3)

Since $i, j \in \{1, 2, ..., p-2\}$, so $i + j \in \{2, 3..., 2(p-2)\}$. We partition the proof into the following four cases.

Case 1: $i + j \leq p - 2$. Since $i + j \leq p - 2$, so by (3.1), $D(\zeta^{i+j}) = \left(\sum_{(s,t)\in S_{i+j-1}} \sigma(\zeta^s)\tau(\zeta^t)\right) D(\zeta)$. Then by (3.3), $D(\zeta^i)\tau(\zeta^j) + \sigma(\zeta^i)D(\zeta^j) = D(\zeta^{i+j})$. Therefore, in this case, relations (3.2) hold.

Case 2: i + j = p - 1. By Lemma 3.6, $\sum_{(i,j) \in \cup_{k=0}^{p-2} S_k} \sigma(\zeta^i) \tau(\zeta^j) = 0$.

$$\Rightarrow \sum_{k=0}^{p-2} \sum_{(i,j)\in S_k} \sigma(\zeta^i)\tau(\zeta^j) = 0.$$

$$\Rightarrow \sum_{(i,j)\in S_{p-2}} \sigma(\zeta^i)\tau(\zeta^j) = -\left(\sum_{k=0}^{p-3} \sum_{(i,j)\in S_k} \sigma(\zeta^i)\tau(\zeta^j)\right).$$

$$\Rightarrow \left(\sum_{(i,j)\in S_{p-2}} \sigma(\zeta^i)\tau(\zeta^j)\right) D(\zeta) = -\left(\sum_{k=0}^{p-3} \left(\sum_{(i,j)\in S_k} \sigma(\zeta^i)\tau(\zeta^j)\right) D(\zeta)\right).$$
Therefore, using (3.1) and (3.3),

$$D(\zeta^{i})\tau(\zeta^{j}) + \sigma(\zeta^{i})D(\zeta^{j}) = -\left(\sum_{k=0}^{p-3} D(\zeta^{k+1})\right) = -D(1+\zeta+\zeta^{2}+\ldots+\zeta^{p-2}) = D(\zeta^{p-1}).$$

So (3.2) holds in this case too.

Case 3: i + j = p.

Since the ring endomorphisms σ and τ are non-zero and different, therefore, $\sigma(\zeta) = \zeta^u$ and $\tau(\zeta) = \zeta^{u+v}$ for some $u, v \in \{1, 2, ..., p-1\}$ such that $u + v \neq p$. Now by (3.3),

$$D(\zeta^{i})\tau(\zeta^{j}) + \sigma(\zeta^{i})D(\zeta^{j}) = \left(\sum_{(s,t)\in S_{i+j-1}}\sigma(\zeta^{s})\tau(\zeta^{t})\right)D(\zeta) = \left(\sum_{(s,t)\in S_{p-1}}\sigma(\zeta^{s})\tau(\zeta^{t})\right)D(\zeta)$$
$$= \left(\sum_{s=0}^{p-1}\sigma(\zeta^{s})\tau(\zeta^{p-1-s})\right)D(\zeta) = \zeta^{(p-1)u}\left(\sum_{s=0}^{p-1}\zeta^{(p-1-s)v}\right)D(\zeta)$$
$$= \zeta^{(p-1)u}\left(1 + \zeta^{v} + \zeta^{2v} + \dots + \zeta^{(p-1)v}\right)D(\zeta) = 0,$$

as ζ^v satisfies the cyclotomic polynomial. Therefore, $D(\zeta^i)\tau(\zeta^j) + \sigma(\zeta^i)D(\zeta^j) = D(\zeta^p)$ as $\zeta^p = 1$ and D(1) = 0. So (3.2) holds in this case as well.

Case 4: $p < i + j \le 2(p - 2)$. Then i + j = m + p for some $m \in \{1, 2, ..., p - 2\}$. By (3.1),

$$D(\zeta^{i+j}) = D(\zeta^m) = \left(\sum_{(s,t)\in S_{m-1}} \sigma(\zeta^s)\tau(\zeta^t)\right) D(\zeta) = \left(\sum_{t=0}^{m-1} \sigma(\zeta^{m-1-t})\tau(\zeta^t)\right) D(\zeta)$$
$$= \zeta^{(m-1)u} \left(\sum_{t=0}^{m-1} \zeta^{tv}\right) D(\zeta).$$

Further, by (3.3), $D(\zeta^i)\tau(\zeta^j) + \sigma(\zeta^i)D(\zeta^j) = \left(\sum_{(s,t)\in S_{i+j-1}} \sigma(\zeta^s)\tau(\zeta^t)\right)D(\zeta)$

$$= \left(\sum_{(s,t)\in S_{m+p-1}} \sigma(\zeta^{s})\tau(\zeta^{t})\right) D(\zeta) = \left(\sum_{t=0}^{m+p-1} \sigma(\zeta^{m+p-1-t})\tau(\zeta^{t})\right) D(\zeta)$$
$$= \zeta^{(m+p-1)u} \left(\sum_{t=0}^{m+p-1} \zeta^{tv}\right) D(\zeta) = \zeta^{(m+p-1)u} \left(\sum_{t=0}^{m-1} \zeta^{tv} + \sum_{t=m}^{m+p-1} \zeta^{tv}\right) D(\zeta)$$
$$= \zeta^{(m-1)u} \left(\sum_{t=0}^{m-1} \zeta^{tv} + \zeta^{mv} \left(\sum_{t=0}^{p-1} \zeta^{tv}\right)\right) D(\zeta) = \zeta^{(m-1)u} \left(\sum_{t=0}^{m-1} \zeta^{tv}\right) D(\zeta)$$

since $\sum_{t=0}^{p-1} \zeta^{tv} = 0$ as $v \in \{1, 2, ..., p-1\}$. Therefore, $D(\zeta^{i+j}) = D(\zeta^i)\tau(\zeta^j) + \sigma(\zeta^i)D(\zeta^j)$. This gives again the relations (3.2). We conclude that the relations (3.2) hold for all $i, j \in \{0, 1, ..., p-2\}$. The result now can be concluded from Lemma 2.1.

As a consequence of Lemma 3.1 and Lemma 3.7, we have the main theorem.

Theorem 3.8. Let $D: O_K \to O_K$ be a \mathbb{Z} -linear map with D(1) = 0. Then D is a (σ, τ) -derivation if and only if the following relations hold for all $k \in \{1, 2, ..., p-2\}$:

$$D(\zeta^k) = \left(\sum_{(i,j)\in S_{k-1}} \sigma(\zeta^i)\tau(\zeta^j)\right) D(\zeta).$$

Corollary 3.9. The \mathbb{Z} -module $\mathcal{D}_{(\sigma,\tau)}(O_K)$ is finitely generated of rank p-1.

Proof. For every $i \in \{0, 1, ..., p-2\}$, define $D_i : O_K \to O_K$ as a \mathbb{Z} -linear map with D(1) = 0 and

$$D_i(\zeta) = \zeta^i$$

More precisely, for each $i \in \{0, 1, ..., p-2\}$, let $D_i : O_K \to O_K$ be a map defined by

$$D_i\left(\sum_{j=0}^{p-2} a_j \zeta^j\right) = \sum_{j=1}^{p-2} a_j D_i(\zeta^j),$$

where for each $j \in \{1, 2, ..., p-2\}, D_i(\zeta^j)$ is defined as

$$D_i(\zeta^j) = \left(\sum_{(s,t)\in S_{j-1}} \sigma(\zeta^s)\tau(\zeta^t)\right) D_i(\zeta).$$

Then by Theorem 3.8, for each $i \in \{0, 1, ..., p-2\}$, $D_i : O_K \to O_K$ is a (σ, τ) -derivation. Further, it can be easily verified that $\{D_0, D_1, ..., D_{p-2}\}$ forms a linearly independent subset of the \mathbb{Z} -module $\mathcal{D}_{(\sigma,\tau)}(O_K)$ that generates it. Hence the result is proved.

We observe from the preceding corollary that the rank of the \mathbb{Z} -module $\mathcal{D}_{(\sigma,\tau)}(O_K)$ is equal to p-1, the degree of $K = \mathbb{Q}(\zeta)$ and which, by Theorem 1.2, is also equal to the rank of the free abelian (additive) group $O_K = \mathbb{Z}[\zeta]$.

Some immediate corollaries can be stated below.

Corollary 3.10. Let $K = Q(\zeta)$ be a 3th cyclotomic field. Then any \mathbb{Z} -linear map $D : O_K \to O_K$ with D(1) = 0 is a (σ, τ) -derivation.

Corollary 3.11. There always exists a non-zero (σ, τ) -derivation of $O_K = \mathbb{Z}[\zeta]$.

We propose the following conjecture.

Conjecture 3.12. Suppose $\beta = \sum_{i=0}^{p-2} b_i \zeta^i \in O_K$ and $\beta(\tau - \sigma)(\zeta) = \sum_{i=0}^{p-2} \left(\sum_{j=0}^{p-2} a_{ij} b_j \right) \zeta^i$. Then $A = [a_{ij}]$ is a $(p-1) \times (p-1)$ matrix with determinant p. We have verified the conjecture for all odd primes less than 100 using SAGE and MAT-LAB. Calculations for primes 3, 5, 7, 11, 13 were done on SAGE (explicit matrix A and det(A)were found for all possible values of the pair $(\sigma(\zeta), \tau(\zeta))$ (see the appendix)). We then developed a code on MATLAB and calculations for the remaining primes were done by running that MATLAB code. For larger primes, MATLAB took a long time to provide the output after running the code. Nevertheless, we believe the conjecture to be true for all odd primes. As a consequence, we propose another conjecture which too will hold once the above conjecture is proved.

Let $\beta = \sum_{i=0}^{p-2} b_i \zeta^i \in K$, D be a (σ, τ) -derivation of O_K and $D(\zeta) = \sum_{i=0}^{p-2} c_i \zeta^i \in O_K$. Put $X^T = (b_0 \ b_1 \ \dots \ b_{p-2})$ and $C^T = (c_0 \ c_1 \ \dots \ c_{p-2})$. We denote by \mathbb{Z}^{p-1} the set of all $(p-1) \times 1$ column matrices with entries from \mathbb{Z} . Also, Adj(A) denotes the adjoint of the matrix A.

Note that since $\sigma \neq \tau$, therefore, $D(\zeta) = \beta(\tau - \sigma)(\zeta)$ always has a solution β in K. Furthermore, $D(\zeta) = \beta(\tau - \sigma)(\zeta)$ has a solution β in O_K if and only if AX = C has a solution in \mathbb{Z}^{p-1} .

Conjecture 3.13. With the above notations, D is inner if and only if $\frac{1}{p}(Adj(A)C) \in \mathbb{Z}^{p-1}$. In particular, the following conditions hold:

- (i) O_K has non-trivial outer (σ, τ) -derivations.
- (ii) If p divides c_i for each $i \in \{0, 1, ..., p-2\}$, then D is inner.

Given below are some interesting examples.

Example 3.14. Let $p \geq 5$. Then not every \mathbb{Z} -linear map $D : \mathbb{Z}[\zeta] \longrightarrow \mathbb{Z}[\zeta]$ with D(1) = 0is a (σ, τ) -derivation. In fact, there exists a non-zero \mathbb{Z} -linear map $D : \mathbb{Z}[\zeta] \longrightarrow \mathbb{Z}[\zeta]$ with D(1) = 0 which is not a (σ, τ) -derivation for every pair (σ, τ) ($\sigma \neq 0, \tau \neq 0, \sigma \neq \tau$). As an example, define $D : \mathbb{Z}[\zeta] \longrightarrow \mathbb{Z}[\zeta]$ by $D(\sum_{i=0}^{p-2} a_i \zeta^i) = a_1 D(\zeta)$, where $D(\zeta) \in O_K \setminus \{0\}$, say, $D(\zeta) = \zeta$. Then D is a non-zero well-defined \mathbb{Z} -linear map with D(1) = 0. But in view of Theorem 3.8, D is not a (σ, τ) -derivation, since

$$D(\zeta^2) \neq \left(\sum_{(i,j)\in S_1} \sigma(\zeta^i)\tau(\zeta^j)\right) D(\zeta).$$

Example 3.15. Let p = 5. Let σ and τ be given by $\sigma(\zeta) = \zeta$ and $\tau(\zeta) = \zeta^2$.

(i) Define $D: O_K \to O_K$ by

$$D(\sum_{i=0}^{3} a_i \zeta^i) = \sum_{i=1}^{3} a_i D(\zeta^i), \ \forall \ a_i \in \mathbb{Z} \ (i \in \{0, 1, 2, 3\}),$$

where for each $i \in \{1, 2, 3\}, D(\zeta^i)$ is defined as

$$D(\zeta^i) = \left(\sum_{(s,t)\in S_{i-1}} \sigma(\zeta^s)\tau(\zeta^t)\right) D(\zeta).$$

Note that D is a \mathbb{Z} -linear map with D(1) = 0. Take $D(\zeta) \in O_K \setminus \{0\}$. Then by Theorem 3.8, D is a non-zero (σ, τ) -derivation. But as shown below, D is not inner if in particular, we take $D(\zeta) = \zeta$. Thus in a cyclotomic field K, it is not necessary that every (σ, τ) -derivation of its ring of algebraic integers O_K is inner.

(ii) If possible, suppose that the (σ, τ) -derivation $D : O_K \to O_K$ is inner. Then there exists some $\beta \in O_K$ such that $D(\alpha) = \beta(\sigma - \tau)(\alpha), \forall \alpha \in O_K$. Since $\beta \in O_K$, so $\beta = \sum_{i=0}^{3} b_i \zeta^i$ for some unique $b_i \in \mathbb{Z}$ for all $i \in \{0, 1, 2, 3\}$.

$$\beta(\sigma - \tau)(\zeta) = \beta(\sigma(\zeta) - \tau(\zeta))$$

= $(b_0 + b_1\zeta + b_2\zeta^2 + b_3\zeta^3)(\zeta - \zeta^2)$
= $(b_2 - 2b_3) + (b_0 + b_2 - b_3)\zeta + (-b_0 + b_1 + b_2 - b_3)\zeta^2 + (-b_1 + 2b_2 - b_3)\zeta^3$

Now $\beta(\sigma - \tau)(\zeta) = D(\zeta)$ with $D(\zeta) = \zeta$ gives

$$(b_2 - 2b_3) + (b_0 + b_2 - b_3)\zeta + (-b_0 + b_1 + b_2 - b_3)\zeta^2 + (-b_1 + 2b_2 - b_3)\zeta^3 = 0 + 1\zeta + 0\zeta^2 + 0\zeta^3.$$

 $\Rightarrow b_2 - 2b_3 = 0; b_0 + b_2 - b_3 = 1; -b_0 + b_1 + b_2 - b_3 = 0; -b_1 + 2b_2 - b_3 = 0.$ This system of linear equations can be written as AX = C, where

$$A = \begin{pmatrix} 0 & 0 & 1 & -2 \\ 1 & 0 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 0 & -1 & 2 & -1 \end{pmatrix}, \ X^{T} = \begin{pmatrix} b_{0} & b_{1} & b_{2} & b_{3} \end{pmatrix} \text{ and } C^{T} = \begin{pmatrix} 0 & 1 & 0 & 0 \end{pmatrix}.$$

This can have a (unique) integral solution if and only if A is a unimodular matrix, that is, the determinant of A is either 1 or -1. But the determinant of A is 5. Therefore, there is no integer solution. Hence a contradiction. Therefore, the given (σ, τ) -derivation $D: O_K \to O_K$ is not inner for $D(\zeta) = \zeta$.

3.3 (σ, τ) -derivations of bi-quadratic number rings

Here we present an elegant application of Lemma 2.1 in classifying all (σ, τ) -derivations and inner (σ, τ) -derivations of the bi-quadratic number ring $\mathbb{Z}[\sqrt{m}, \sqrt{n}]$. Throughout this subsection, K denotes the quartic number field $K = \mathbb{Q}(\sqrt{m}, \sqrt{n})$, where m and n are distinct square-free rational integers, \mathcal{A} the number ring $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ with \mathbb{Z} -basis $\{1, \sqrt{m}, \sqrt{n}, \sqrt{mn}\}$, and σ and τ any two non-zero ring endomorphisms of \mathcal{A} .

Lemma 3.16. If $D : \mathcal{A} \to \mathcal{A}$ is a non-zero (σ, τ) -derivation of \mathcal{A} , then precisely one case amongst (i), (ii) or (iii) holds.

(i) $D(\sqrt{m}) = 0$, $D(\sqrt{n}) \neq 0$, $D(\sqrt{mn}) \neq 0$ and $D(\sqrt{mn}) = \pm\sqrt{m}D(\sqrt{n})$. Further, in this case, the possible values of the pair (σ, τ) are $(\phi_1, \phi_2), (\phi_2, \phi_1), (\phi_3, \phi_4), (\phi_4, \phi_3)$. In fact, $D(\sqrt{mn}) = \sqrt{m}D(\sqrt{n})$ if $(\sigma, \tau) = (\phi_1, \phi_2)$ or (ϕ_2, ϕ_1) and $D(\sqrt{mn}) = -\sqrt{m}D(\sqrt{n})$ if $(\sigma, \tau) = (\phi_3, \phi_4)$ or (ϕ_4, ϕ_3) .

- (ii) $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) = 0$, $D(\sqrt{mn}) \neq 0$ and $D(\sqrt{mn}) = \pm \sqrt{n}D(\sqrt{m})$. Further, in this case, the possible values of the pair (σ, τ) are $(\phi_1, \phi_3), (\phi_2, \phi_4), (\phi_3, \phi_1), (\phi_4, \phi_2)$. In fact, $D(\sqrt{mn}) = \sqrt{n}D(\sqrt{m})$ if $(\sigma, \tau) = (\phi_1, \phi_3)$ or (ϕ_3, ϕ_1) and $D(\sqrt{mn}) = -\sqrt{n}D(\sqrt{m})$ if $(\sigma, \tau) = (\phi_2, \phi_4)$ or (ϕ_4, ϕ_2) .
- (iii) $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) \neq 0$, $D(\sqrt{mn}) = 0$ and $\sqrt{m}D(\sqrt{n}) = \pm\sqrt{n}D(\sqrt{m})$. Further, in this case, the possible values of the pair (σ, τ) are $(\phi_1, \phi_4), (\phi_2, \phi_3), (\phi_4, \phi_1), (\phi_3, \phi_2)$.

In fact,
$$\sqrt{m}D(\sqrt{n}) = \sqrt{n}D(\sqrt{m})$$
 if $(\sigma, \tau) = (\phi_1, \phi_4)$ or (ϕ_4, ϕ_1) and $\sqrt{m}D(\sqrt{n}) = -\sqrt{n}D(\sqrt{m})$ if $(\sigma, \tau) = (\phi_2, \phi_3)$ or (ϕ_3, ϕ_2) .

Here $\phi_1, \phi_2, \phi_3, \phi_4$ are the non-zero ring endomorphisms of $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ given by

- (a) $\phi_1(\sqrt{m}) = \sqrt{m}, \ \phi_1(\sqrt{n}) = \sqrt{n}.$
- (b) $\phi_2(\sqrt{m}) = \sqrt{m}, \ \phi_2(\sqrt{n}) = -\sqrt{n}.$
- (c) $\phi_3(\sqrt{m}) = -\sqrt{m}, \ \phi_3(\sqrt{n}) = \sqrt{n}.$
- (d) $\phi_4(\sqrt{m}) = -\sqrt{m}, \ \phi_4(\sqrt{n}) = -\sqrt{n}.$

Proof. If ϕ is a non-zero ring endomorphisms of \mathcal{A} , then $(\phi(\sqrt{m}))^2 = m$ and $(\phi(\sqrt{n}))^2 = n$. So $\phi(\sqrt{m}) = \pm \sqrt{m}$ and $\phi(\sqrt{n}) = \pm \sqrt{n}$. Therefore, \mathcal{A} has precisely the four non-zero ring endomorphisms $\phi_1, \phi_2, \phi_3, \phi_4$ given in the table below.

Table 1

	1	\sqrt{m}	\sqrt{n}	\sqrt{mn}
ϕ_1	1	\sqrt{m}	\sqrt{n}	\sqrt{mn}
ϕ_2	1	\sqrt{m}	$-\sqrt{n}$	$-\sqrt{mn}$
ϕ_3	1	$-\sqrt{m}$	\sqrt{n}	$-\sqrt{mn}$
ϕ_4	1	$-\sqrt{m}$	$-\sqrt{n}$	\sqrt{mn}

Put $\alpha_1 = 1$, $\alpha_2 = \sqrt{m}$, $\alpha_3 = \sqrt{n}$ and $\alpha_4 = \sqrt{mn}$. Then by Lemma 2.1, $D(\alpha_i \alpha_j) = D(\alpha_i)\tau(\alpha_j) + \sigma(\alpha_i)D(\alpha_j)$, $\forall i, j \in \{1, 2, 3, 4\}$. Also, D is \mathbb{Z} -linear. So we get the following table 2.

(i, j)	α_i	α_j	$D(\alpha_i \alpha_j)$	$D(\alpha_i)\tau(\alpha_j) + \sigma(\alpha_i)D(\alpha_j)$
(1, 1)	1	1	0	0
(1, 2)	1	\sqrt{m}	$D(\sqrt{m})$	$D(\sqrt{m})$
(1,3)	1	\sqrt{n}	$D(\sqrt{n})$	$D(\sqrt{n})$
(1, 4)	1	\sqrt{mn}	$D(\sqrt{mn})$	$D(\sqrt{mn})$
(2,1)	\sqrt{m}	1	$D(\sqrt{m})$	$D(\sqrt{m})$
(2, 2)	\sqrt{m}	\sqrt{m}	0	$(\sigma(\sqrt{m}) + \tau(\sqrt{m}))D(\sqrt{m})$
(2,3)	\sqrt{m}	\sqrt{n}	$D(\sqrt{mn})$	$D(\sqrt{m})\tau(\sqrt{n}) + \sigma(\sqrt{m})D(\sqrt{n})$
(2,4)	\sqrt{m}	\sqrt{mn}	$mD(\sqrt{n})$	$D(\sqrt{m})\tau(\sqrt{mn}) + \sigma(\sqrt{m})D(\sqrt{mn})$
(3, 1)	\sqrt{n}	1	$D(\sqrt{n})$	$D(\sqrt{n})$
(3, 2)	\sqrt{n}	\sqrt{m}	$D(\sqrt{mn})$	$D(\sqrt{n})\tau(\sqrt{m}) + \sigma(\sqrt{n})D(\sqrt{m})$
(3, 3)	\sqrt{n}	\sqrt{n}	0	$(\sigma(\sqrt{n}) + \tau(\sqrt{n}))D(\sqrt{n})$
(3, 4)	\sqrt{n}	\sqrt{mn}	$nD(\sqrt{m})$	$D(\sqrt{n})\tau(\sqrt{mn}) + \sigma(\sqrt{n})D(\sqrt{mn})$
(4, 1)	\sqrt{mn}	1	$D(\sqrt{mn})$	$D(\sqrt{mn})$
(4, 2)	\sqrt{mn}	\sqrt{m}	$mD(\sqrt{n})$	$D(\sqrt{mn})\tau(\sqrt{m}) + \sigma(\sqrt{mn})D(\sqrt{m})$
(4, 3)	\sqrt{mn}	\sqrt{n}	$nD(\sqrt{m})$	$D(\sqrt{mn})\tau(\sqrt{n}) + \sigma(\sqrt{mn})D(\sqrt{n})$
(4, 4)	\sqrt{mn}	\sqrt{mn}	0	$(\sigma(\sqrt{mn}) + \tau(\sqrt{mn}))D(\sqrt{mn})$

Table 2

From Table 2 containing the values of $D(\alpha_i \alpha_j)$ and $D(\alpha_i)\tau(\alpha_j) + \sigma(\alpha_i)D(\alpha_j)$ $(i, j \in \{1, 2, 3, 4\})$, we see that the relation $D(\alpha_i \alpha_j) = D(\alpha_i)\tau(\alpha_j) + \sigma(\alpha_i)D(\alpha_j)$ is being satisfied by the pairs

$$(i, j) = (1, 1), (1, 2), (1, 3), (1, 4), (2, 1), (3, 1), (4, 1).$$

Then using the fact that the Z-algebra $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ is commutative, we get that D will be a (σ, τ) -derivation if and only if it satisfies the following nine equations simultaneously.

$$(\sigma(\sqrt{m}) + \tau(\sqrt{m}))D(\sqrt{m}) = 0 \tag{3.4}$$

$$D(\sqrt{m})\tau(\sqrt{n}) + \sigma(\sqrt{m})D(\sqrt{n}) = D(\sqrt{mn})$$
(3.5)

$$D(\sqrt{m})\tau(\sqrt{mn}) + \sigma(\sqrt{m})D(\sqrt{mn}) = mD(\sqrt{n})$$
(3.6)

$$(\sigma(\sqrt{n}) + \tau(\sqrt{n}))D(\sqrt{n}) = 0 \tag{3.7}$$

$$D(\sqrt{n})\tau(\sqrt{mn}) + \sigma(\sqrt{n})D(\sqrt{mn}) = nD(\sqrt{m})$$
(3.8)

$$(\sigma(\sqrt{mn}) + \tau(\sqrt{mn}))D(\sqrt{mn}) = 0$$
(3.9)

$$D(\sqrt{n})\tau(\sqrt{m}) + \sigma(\sqrt{n})D(\sqrt{m}) = D(\sqrt{mn})$$
(3.10)

$$D(\sqrt{mn})\tau(\sqrt{m}) + \sigma(\sqrt{mn})D(\sqrt{m}) = mD(\sqrt{n})$$
(3.11)

$$D(\sqrt{mn})\tau(\sqrt{n}) + \sigma(\sqrt{mn})D(\sqrt{n}) = nD(\sqrt{m})$$
(3.12)

Since $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ is an integral domain, so in view of equations (3.4), (3.7) and (3.9), there are four possible cases.

Case 1: All of $D(\sqrt{m})$, $D(\sqrt{n})$ and $D(\sqrt{mn})$ are non-zero. Since $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ is an integral domain, therefore, from (3.4), (3.7) and (3.9),

 $\tau(\sqrt{m}) = -\sigma(\sqrt{m}), \ \tau(\sqrt{n}) = -\sigma(\sqrt{n}) \text{ and } \tau(\sqrt{mn}) = -\sigma(\sqrt{mn}).$

But from Table 1, we observe that there is no such pair (σ, τ) which satisfies the above three equations simultaneously. Therefore, this case is not possible.

Case 2: Exactly one of $D(\sqrt{m})$, $D(\sqrt{n})$ and $D(\sqrt{mn})$ is non-zero.

Then there are three possible subcases.

Subcase 2.1: $D(\sqrt{m}) = 0$, $D(\sqrt{n}) \neq 0$, $D(\sqrt{mn}) \neq 0$.

By equations (3.7) and (3.9) respectively, $\tau(\sqrt{n}) = -\sigma(\sqrt{n})$ and $\tau(\sqrt{mn}) = -\sigma(\sqrt{mn})$. Also, by (3.5) and (3.10), $\tau(\sqrt{m}) = \sigma(\sqrt{m})$. Therefore, in view of Table 1, the possible values of the pair (σ, τ) are

$$(\phi_1, \phi_2), (\phi_2, \phi_1), (\phi_3, \phi_4), (\phi_4, \phi_3).$$

Further, from (3.5), $D(\sqrt{mn}) = \sigma(\sqrt{m})D(\sqrt{n})$ and $\sigma(\sqrt{m}) = \pm\sqrt{m}$, therefore, D must satisfy $D(\sqrt{mn}) = \pm\sqrt{m}D(\sqrt{n})$. More precisely, we have the following table 3:

Table 3

(σ, τ)	$D(\sqrt{mn}) = \pm \sqrt{m} D(\sqrt{n})?$
(ϕ_1,ϕ_2)	$D(\sqrt{mn}) = \sqrt{m}D(\sqrt{n})$
(ϕ_2,ϕ_1)	$D(\sqrt{mn}) = \sqrt{m}D(\sqrt{n})$
(ϕ_3,ϕ_4)	$D(\sqrt{mn}) = -\sqrt{m}D(\sqrt{n})$
(ϕ_4,ϕ_3)	$D(\sqrt{mn}) = -\sqrt{m}D(\sqrt{n})$

Subcase 2.2: $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) = 0$, $D(\sqrt{mn}) \neq 0$.

By equations (3.4) and (3.9) respectively, $\tau(\sqrt{m}) = -\sigma(\sqrt{m})$ and $\tau(\sqrt{mn}) = -\sigma(\sqrt{mn})$. Also, by (3.5) and (3.10), $\tau(\sqrt{n}) = \sigma(\sqrt{n})$. Therefore, in view of Table 1, the possible values of the pair (σ, τ) are

$$(\phi_1, \phi_3), (\phi_2, \phi_4), (\phi_3, \phi_1), (\phi_4, \phi_2).$$

Further from (3.10), $D(\sqrt{mn}) = \sigma(\sqrt{n})D(\sqrt{m})$ and $\sigma(\sqrt{n}) = \pm\sqrt{n}$, therefore, D must satisfy $D(\sqrt{mn}) = \pm\sqrt{n}D(\sqrt{m})$. More precisely, we have the following table 4:

Table 4	Ta	bl	e	4
---------	----	----	---	---

(σ, τ)	$D(\sqrt{mn}) = \pm \sqrt{n} D(\sqrt{m})?$
(ϕ_1,ϕ_3)	$D(\sqrt{mn}) = \sqrt{n}D(\sqrt{m})$
(ϕ_2,ϕ_4)	$D(\sqrt{mn}) = -\sqrt{n}D(\sqrt{m})$
(ϕ_3,ϕ_1)	$D(\sqrt{mn}) = \sqrt{n}D(\sqrt{m})$
(ϕ_4,ϕ_2)	$D(\sqrt{mn}) = -\sqrt{n}D(\sqrt{m})$

Subcase 2.3: $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) \neq 0$, $D(\sqrt{mn}) = 0$.

By equations (3.4) and (3.7) respectively, $\tau(\sqrt{m}) = -\sigma(\sqrt{m})$ and $\tau(\sqrt{n}) = -\sigma(\sqrt{n})$. Also, by equations (3.6) and (3.11), $\tau(\sqrt{mn}) = \sigma(\sqrt{mn})$. Therefore, in view of Table 1, the possible values of the pair (σ, τ) are

$$(\phi_1, \phi_4), \ (\phi_2, \phi_3), \ (\phi_4, \phi_1), \ (\phi_3, \phi_2).$$

Further from equation (3.11), $mD(\sqrt{n}) = \sigma(\sqrt{mn})D(\sqrt{m})$ and $\sigma(\sqrt{mn}) = \pm\sqrt{mn}$, therefore, D must satisfy $mD(\sqrt{n}) = \pm\sqrt{mn}D(\sqrt{m})$ so that $\sqrt{m}D(\sqrt{n}) = \pm\sqrt{n}D(\sqrt{m})$. More precisely, we have the following table 5:

Table	5
TODIO	\mathbf{U}

(σ, τ)	$\sqrt{m}D(\sqrt{n}) = \sqrt{n}D(\sqrt{m})?$
(ϕ_1,ϕ_4)	$\sqrt{m}D(\sqrt{n}) = \sqrt{n}D(\sqrt{m})$
(ϕ_2,ϕ_3)	$\sqrt{m}D(\sqrt{n}) = -\sqrt{n}D(\sqrt{m})$
(ϕ_3,ϕ_2)	$\sqrt{m}D(\sqrt{n}) = -\sqrt{n}D(\sqrt{m})$
(ϕ_4,ϕ_1)	$\sqrt{m}D(\sqrt{n}) = \sqrt{n}D(\sqrt{m})$

Case 3: Exactly two of $D(\sqrt{m})$, $D(\sqrt{n})$ or $D(\sqrt{mn})$ are 0.

Then there are three possible subcases.

Subcase 3.1: $D(\sqrt{m}) = 0, \ D(\sqrt{n}) = 0, \ D(\sqrt{mn}) \neq 0.$

- Subcase 3.2: $D(\sqrt{m}) = 0$, $D(\sqrt{n}) \neq 0$, $D(\sqrt{mn}) = 0$.
- Subcase 3.3: $D(\sqrt{m}) \neq 0, D(\sqrt{n}) = 0, D(\sqrt{mn}) = 0.$

Then in each of the three possible subcases, D will become a zero derivation using equations (3.4) to (3.12) and the facts that $\{1, \sqrt{m}, \sqrt{n}, \sqrt{mn}\}$ is a \mathbb{Z} -basis of the \mathbb{Z} -module $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ and $D : \mathcal{A} \to \mathcal{A}$ is \mathbb{Z} -linear with D(1) = 0. But this gives a contradiction because D is non-zero. Therefore, this case is not possible.

Case 4: All $D(\sqrt{p})$, $D(\sqrt{q})$ and $D(\sqrt{pq})$ are zero. The proof is similar to Case 3.

The proof of the following lemma is trivial using Lemma 2.1.

Lemma 3.17. Let $D : \mathcal{A} \to \mathcal{A}$ be any \mathbb{Z} -linear map with D(1) = 0. Then D is a (σ, τ) -derivation in the following cases.

(i)
$$D(\sqrt{m}) = 0, \ D(\sqrt{n}) \neq 0 \ and \ D(\sqrt{mn}) \neq 0.$$

(a) If $D(\sqrt{mn}) = \sqrt{m}D(\sqrt{n}) \ and \ (\sigma, \tau) = (\phi_1, \phi_2) \ or \ (\phi_2, \phi_1).$
(b) If $D(\sqrt{mn}) = -\sqrt{m}D(\sqrt{n}) \ and \ (\sigma, \tau) = (\phi_3, \phi_4) \ or \ (\phi_4, \phi_3).$
(ii) $D(\sqrt{m}) \neq 0, \ D(\sqrt{n}) = 0 \ and \ D(\sqrt{mn}) \neq 0.$

(a) If
$$D(\sqrt{mn}) = \sqrt{n}D(\sqrt{m})$$
 and $(\sigma, \tau) = (\phi_1, \phi_3)$ or (ϕ_3, ϕ_1) .
(b) If $D(\sqrt{mn}) = -\sqrt{n}D(\sqrt{m})$ and $(\sigma, \tau) = (\phi_2, \phi_4)$ or (ϕ_4, ϕ_2) .

(iii) $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) \neq 0$ and $D(\sqrt{mn}) = 0$.

(a) If
$$\sqrt{m}D(\sqrt{n}) = \sqrt{n}D(\sqrt{m})$$
 and $(\sigma, \tau) = (\phi_1, \phi_4)$ or (ϕ_4, ϕ_1) .
(b) If $\sqrt{m}D(\sqrt{n}) = -\sqrt{n}D(\sqrt{m})$ and $(\sigma, \tau) = (\phi_2, \phi_3)$ or (ϕ_3, ϕ_2) .

Theorem 3.18. A non-zero \mathbb{Z} -linear map $D : \mathcal{A} \to \mathcal{A}$ with D(1) = 0 is a (σ, τ) -derivation if and only if exactly one of the cases (i), (ii) or (iii) of Lemma 3.16 holds.

Corollary 3.19. The \mathbb{Z} -module $\mathcal{D}_{(\sigma,\tau)}(\mathcal{A})$ is finitely generated of rank 4.

Proof. $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ has exactly four distinct ring endomorphisms as given in Table 1. In view of Theorem 3.18, there are three possible cases.

Case 1: $(\sigma, \tau) \in \{(\phi_1, \phi_2), (\phi_2, \phi_1), (\phi_3, \phi_4), (\phi_4, \phi_3)\}.$

For each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(1) = 0 such that

$$D_1(\sqrt{n}) = 1$$
, $D_2(\sqrt{n}) = \sqrt{m}$, $D_3(\sqrt{n}) = \sqrt{n}$, $D_4(\sqrt{n}) = \sqrt{mn}$.

More precisely, for each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ by

$$D_i(a_1 + a_2\sqrt{m} + a_3\sqrt{n} + a_4\sqrt{mn}) = \begin{cases} (a_3 + a_4\sqrt{m})D_i(\sqrt{n}) & \text{if } (\sigma,\tau) = (\phi_1,\phi_2) \text{ or } (\phi_2,\phi_1) \\ (a_3 - a_4\sqrt{m})D_i(\sqrt{n}) & \text{if } (\sigma,\tau) = (\phi_3,\phi_4) \text{ or } (\phi_4,\phi_3) \end{cases}$$

where for each $i \in \{1, 2, 3, 4\}$, $D_i(\sqrt{n})$ is defined as above.

Case 2: $(\sigma, \tau) \in \{(\phi_1, \phi_3), (\phi_3, \phi_1), (\phi_2, \phi_4), (\phi_4, \phi_2)\}.$ For each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(1) = 0 such that

$$D_1(\sqrt{m}) = 1, \ D_2(\sqrt{m}) = \sqrt{m}, \ D_3(\sqrt{m}) = \sqrt{n}, \ D_4(\sqrt{m}) = \sqrt{mn}$$

More precisely, for each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ by

$$D_i(a_1 + a_2\sqrt{m} + a_3\sqrt{n} + a_4\sqrt{mn}) = \begin{cases} (a_2 + a_4\sqrt{n})D_i(\sqrt{m}) & \text{if } (\sigma,\tau) = (\phi_1,\phi_3) \text{ or } (\phi_3,\phi_1) \\ (a_2 - a_4\sqrt{n})D_i(\sqrt{m}) & \text{if } (\sigma,\tau) = (\phi_2,\phi_4) \text{ or } (\phi_4,\phi_2) \end{cases}$$

where for each $i \in \{1, 2, 3, 4\}$, $D_i(\sqrt{m})$ is defined as above.

Case 3: $(\sigma, \tau) \in \{(\phi_1, \phi_4), (\phi_4, \phi_1), (\phi_2, \phi_3), (\phi_3, \phi_2)\}.$

Suppose gcd(m, n) = k. Then m = kr and n = ks for some $r, s \in \mathbb{Z}$.

For each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ as a \mathbb{Z} -linear map with D(1) = 0 such that

$$D_1(\sqrt{m}) = m, \ D_2(\sqrt{m}) = \sqrt{m}, \ D_3(\sqrt{m}) = r\sqrt{n}, \ D_4(\sqrt{m}) = \sqrt{mn}$$

so that

$$D_1(\sqrt{n}) = \sqrt{mn}, \ D_2(\sqrt{n}) = \sqrt{n}, \ D_3(\sqrt{n}) = s\sqrt{m}, \ D_4(\sqrt{n}) = n.$$

More precisely, for each $i \in \{1, 2, 3, 4\}$, define $D_i : \mathcal{A} \to \mathcal{A}$ by

$$D_i(a_1 + a_2\sqrt{m} + a_3\sqrt{n} + a_4\sqrt{mn}) = a_2D_i(\sqrt{m}) + a_3D_i(\sqrt{n})$$

where for each $i \in \{1, 2, 3, 4\}$, $D_i(\sqrt{m})$ and $D_i(\sqrt{n})$ are defined as above.

In all three cases, it can be easily verified that $\{D_1, D_2, D_3, D_4\}$ is a linearly independent subset of the \mathbb{Z} -module $\mathcal{D}_{(\sigma,\tau)}(\mathcal{A})$ that generates it.

We note from the preceding corollary that the rank of the Z-module $\mathcal{D}_{(\sigma,\tau)}(\mathcal{A})$ is equal to 4, the degree of $K = \mathbb{Q}(\sqrt{m}, \sqrt{n})$ and which, by Theorem 1.2, is also equal to the rank of the free abelian group $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ as a Z-module.

Corollary 3.20. There always exists a non-zero (σ, τ) -derivation of \mathcal{A} .

Corollary 3.21. There always exists a non-zero \mathbb{Z} -linear map $D : \mathcal{A} \to \mathcal{A}$ with D(1) = 0 that is not a (σ, τ) -derivation. In fact, there exists a non-zero \mathbb{Z} -linear map $D : \mathcal{A} \to \mathcal{A}$ with D(1) = 0 which is not a (σ, τ) -derivation for every pair (σ, τ) of any two different non-zero ring endomorphisms of \mathcal{A} .

Theorem 3.22. Let $D : \mathcal{A} \to \mathcal{A}$ be a non-zero (σ, τ) -derivation.

- (i) If $D(\sqrt{m}) = 0$, $D(\sqrt{n}) \neq 0$ and $D(\sqrt{mn}) \neq 0$, then D is inner if and only if $\frac{D(\sqrt{n})}{2\sqrt{n}} \in \mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$. In particular, \mathcal{A} has non-trivial outer (σ, τ) -derivations, where $(\sigma, \tau) \in \{(\phi_1, \phi_2), (\phi_2, \phi_1), (\phi_3, \phi_4), (\phi_4, \phi_3)\}$.
- (ii) If $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) = 0$ and $D(\sqrt{mn}) \neq 0$, then D is inner if and only if $\frac{D(\sqrt{m})}{2\sqrt{m}} \in \mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$. In particular, \mathcal{A} has non-trivial outer (σ, τ) -derivations, where $(\sigma, \tau) \in \{(\phi_1, \phi_3), (\phi_2, \phi_4), (\phi_3, \phi_1), (\phi_4, \phi_2)\}$.
- (iii) If $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) \neq 0$ and $D(\sqrt{mn}) = 0$, then D is inner if and only if either $\frac{D(\sqrt{m})}{2\sqrt{m}} \in \mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ or $\frac{D(\sqrt{n})}{2\sqrt{n}} \in \mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$. In particular, \mathcal{A} has non-trivial outer (σ, τ) -derivations, where $(\sigma, \tau) \in \{(\phi_1, \phi_4), (\phi_2, \phi_3), (\phi_4, \phi_1), (\phi_3, \phi_2)\}$.

Proof. $\mathcal{A} = \mathbb{Z}[\sqrt{m}, \sqrt{n}]$ has exactly four distinct ring endomorphisms as given in Table 1. In view of Theorem 3.18, there are three possible cases.

Case 1: $D(\sqrt{m}) = 0$, $D(\sqrt{n}) \neq 0$ and $D(\sqrt{mn}) \neq 0$.

In this case, $(\sigma, \tau) \in \{(\phi_1, \phi_2), (\phi_2, \phi_1), (\phi_3, \phi_4), (\phi_4, \phi_3)\}$. In view of Subcase 2.1 in the proof of Lemma 3.16, in this case,

$$\tau(\sqrt{m}) = \sigma(\sqrt{m}), \ \tau(\sqrt{n}) = -\sigma(\sqrt{n}) \ \text{and} \ \tau(\sqrt{mn}) = -\sigma(\sqrt{mn}).$$

Note that $(\tau - \sigma)(1) = 0$, $(\tau - \sigma)(\sqrt{m}) = 0$, $(\tau - \sigma)(\sqrt{n}) = 2\tau(\sqrt{n})$ and $(\tau - \sigma)(\sqrt{mn}) = 2\tau(\sqrt{mn}) = 2\tau(\sqrt{m})\tau(\sqrt{n})$. By Theorem 2.5, D is inner if and only if there is a $\beta \in \mathcal{A}$ with

$$D(1) = \beta(\tau - \sigma)(1), \quad D(\sqrt{m}) = \beta(\tau - \sigma)(\sqrt{m}),$$
$$D(\sqrt{n}) = \beta(\tau - \sigma)(\sqrt{n}), \quad D(\sqrt{mn}) = \beta(\tau - \sigma)(\sqrt{mn})$$

Note that $D(1) = \alpha(\tau - \sigma)(1)$ and $D(\sqrt{m}) = \alpha(\tau - \sigma)(\sqrt{m})$ hold for all $\alpha \in \mathcal{A}$. So D will be inner if and only if

$$D(\sqrt{n}) = \beta(\tau - \sigma)(\sqrt{n})$$
 and $D(\sqrt{mn}) = \beta(\tau - \sigma)(\sqrt{mn}),$

that is, if and only if

$$D(\sqrt{n}) = 2\beta\tau(\sqrt{n})$$
 and $D(\sqrt{mn}) = 2\beta\tau(\sqrt{m})\tau(\sqrt{n}).$

Further since in this case, $D(\sqrt{mn}) = \begin{cases} \sqrt{m}D(\sqrt{n}) & \text{if } \tau = \phi_1 \text{ or } \phi_2 \\ -\sqrt{m}D(\sqrt{n}) & \text{if } \tau = \phi_3 \text{ or } \phi_4 \end{cases}$ and

$$2\beta\tau(\sqrt{m})\tau(\sqrt{n}) = \begin{cases} 2\beta\sqrt{mn} \text{ if } \tau = \phi_1 \text{ or } \tau = \phi_4 \\ -2\beta\sqrt{mn} \text{ if } \tau = \phi_2 \text{ or } \tau = \phi_3 \end{cases} \text{ therefore,}$$

 $D(\sqrt{mn}) = 2\beta\tau(\sqrt{m})\tau(\sqrt{n}) \text{ if and only if } D(\sqrt{n}) = \begin{cases} 2\beta\sqrt{n} & \text{if } \tau = \phi_1 \text{ or } \phi_3 \\ -2\beta\sqrt{n} & \text{if } \tau = \phi_2 \text{ or } \phi_4 \end{cases}$

From the above discussion, it can be easily concluded that D is inner if and only if

$$D(\sqrt{n}) = \begin{cases} 2\beta\sqrt{n} & \text{if } \tau = \phi_1 \text{ or } \phi_3 \\ -2\beta\sqrt{n} & \text{if } \tau = \phi_2 \text{ or } \phi_4 \end{cases}$$

Therefore, D is inner if and only if there is a $\beta \in \mathcal{A}$ with

$$\beta = \begin{cases} \frac{D(\sqrt{n})}{2\sqrt{n}} & \text{if } \tau = \phi_1 \text{ or } \phi_3 \\ -\frac{D(\sqrt{n})}{2\sqrt{n}} & \text{if } \tau = \phi_2 \text{ or } \phi_4 \end{cases}$$

Therefore, in this case, D is inner if and only if $\frac{D(\sqrt{n})}{2\sqrt{n}} \in \mathcal{A}$.

Case 2: $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) = 0$ and $D(\sqrt{mn}) \neq 0$. As in Case 1, it can be shown that D is inner if and only if there is a $\beta \in \mathcal{A}$ with

$$\beta = \begin{cases} \frac{D(\sqrt{m})}{2\sqrt{m}} & \text{if } \tau = \phi_1 \text{ or } \phi_2 \\ -\frac{D(\sqrt{m})}{2\sqrt{m}} & \text{if } \tau = \phi_3 \text{ or } \phi_4 \end{cases}$$

Therefore, in this case, D is inner if and only if $\frac{D(\sqrt{m})}{2\sqrt{m}} \in \mathcal{A}$.

Case 3: $D(\sqrt{m}) \neq 0$, $D(\sqrt{n}) \neq 0$ and $D(\sqrt{mn}) = 0$. On similar lines, it can be shown that in this case, D is inner if and only if either $\frac{D(\sqrt{m})}{2\sqrt{m}} \in \mathcal{A}$ or $\frac{D(\sqrt{n})}{2\sqrt{n}} \in \mathcal{A}$ (as $\frac{D(\sqrt{m})}{2\sqrt{m}} = \frac{D(\sqrt{n})}{2\sqrt{n}}$). Hence the result.

The corollary is immediate.

Corollary 3.23. There always exists a non-zero (σ, τ) -inner derivation of \mathcal{A} .

4 Coding Theory Applications

In this section, we give the applications of the work of previous sections in coding theory. We give the notion of a Hom-IDD code over finite fields.

4.1 IDD Code in O_K

Let $K = \mathbb{Q}(\theta)$ (θ an algebraic integer) be a number field of degree n over \mathbb{Q} , which is a normal extension of \mathbb{Q} and $\mathcal{B} = \{\theta_1 = \theta, \theta_2, ..., \theta_n\}$ be an ordered integral basis of O_K . Let $\sigma, \tau : O_K \to O_K$ be two different \mathbb{Z} -algebra endomorphisms of O_K . Let D be a (σ, τ) derivation of O_K .

The range Im(D) of D is a \mathbb{Z} -submodule of the \mathbb{Z} -module O_K with the generating set $D(\mathcal{B}) = \{D(\theta_i) | 1 \leq i \leq n\}$. It generates an (n, r)-code in O_K , where r = rank(Im(D)), the rank of the \mathbb{Z} -module Im(D).

Suppose $S = \{D(\theta_{i_1}), D(\theta_{i_2}), ..., D(\theta_{i_s})\}$ is a Z-linearly independent subset of Im(D). Then S generates a submodule of Im(D) with rank s. Thus, S generates an (n, s)-code.

Definition 4.1. Let $T = \{\theta_{k_1}, \theta_{k_2}, ..., \theta_{k_t}\}$ $(\{k_1, k_2, ..., k_t\} \subseteq \{1, 2, ..., n\})$ be a subset of the integral basis $\{\theta_1, \theta_2, ..., \theta_n\}$ such that D(T) is \mathbb{Z} -linearly independent subset of O_K (or Im(D)). Then the submodule $W = \langle D(T) \rangle$ generates an (n, t)-code, and is called as Image of Derivation-Derived Code, or simply an IDD Code.

4.2 Equivalent IDD Code in \mathbb{Z}^n

In this subsection, we construct a code in \mathbb{Z}^n equivalent to an IDD code.

Suppose $\Delta : \mathbb{Z}^n \to O_K$ be the mapping given by

$$\Delta(a_1, a_2, \dots, a_n) = \sum_{i=1}^n a_i \theta_i$$

For each $i \in \{1, 2, ..., n\}$, suppose $D(\theta_i) = \sum_{j=1}^n a_{ij}\theta_j$. Then $\Delta^{-1}(D(\theta_i)) = (a_{i1}, a_{i2}, ..., a_{in})$.

Construct an $n \times n$ matrix B with the rows $\Delta^{-1}(D(\theta_1)), \Delta^{-1}(D(\theta_2)), ..., \Delta^{-1}(D(\theta_n))$. More precisely,

$$B = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}.$$

Theorem 4.2. Any rows $i_1, i_2, ..., i_t$ of B are \mathbb{Z} -linearly independent if and only if the set $\{D(\theta_{i_1}), D(\theta_{i_2}), ..., D(\theta_{i_t})\}$ is \mathbb{Z} -linearly independent.

Below, we give the steps of construction:

- (i) Suppose W is an Image of Derivations-Derived (IDD) code. Then $W = \langle S \rangle$, for some Z-linearly independent subset S of $D(\mathcal{B})$, say, $S = \{D(\theta_{k_1}), D(\theta_{k_2}), ..., D(\theta_{k_s})\}$.
- (ii) Pick $\underline{w} \in \mathbb{Z}^s$ and $\underline{w} = (\lambda_1, \lambda_2, ..., \lambda_s)$.
- (iii) Write \underline{w} as \underline{x} in \mathbb{Z}^n with λ_j in position k_j and zero elsewhere.

(iv) <u>x</u> can be mapped to an element O_K by $x = \Delta(\underline{x}) = \sum_{j=1}^s \lambda_j \theta_{k_j}$ and a codeword

$$D(x) = D(\Delta(\underline{x})) = D(\sum_{j=1}^{s} \lambda_j \theta_{k_j}) = \sum_{j=1}^{s} \lambda_j D(\theta_{k_j})$$

equated with a codeword in \mathcal{E} given by

$$\Delta^{-1}(D(x)) = \underline{x}B.$$

(v) Thus, we obtain an (n, s) code

 $\mathcal{E} = \{ \underline{x}B \mid \underline{x} \text{ extends } \underline{w} \in \mathbb{Z}^s \text{ to } \mathbb{Z}^n \}.$

4.3 Hom-IDD Code in $(\mathbb{Z}_q)^n$

For a positive integer q and for an equivalent IDD code C in \mathbb{Z}^n , we construct a Hom-IDD code $\Omega(C)$ in $(\mathbb{Z}_q)^n$.

Lemma 4.3. Let q be a positive integer. Then the map $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$ defined by

$$\Omega(\lambda_1, \lambda_2, ..., \lambda_n) = (\overline{\lambda_1}, \overline{\lambda_2}, ..., \overline{\lambda_n}),$$

where $\overline{\lambda_i} = \lambda_i \pmod{q}$ for each $i \in \{1, 2, ..., n\}$ is a \mathbb{Z} -module homomorphism.

Proof. Both \mathbb{Z}^n and $(\mathbb{Z}_q)^n$ are abelian groups with respect to the usual operations of componentwise addition. And since every abelian group is a \mathbb{Z} -module with respect to the usual operation of scalar multiplication. More precisely, if M is an additive abelian group, then M is a \mathbb{Z} -module with respect to the scalar multiplication operation defined by: $\mathbb{Z} \times M \to M$, where

$$(\lambda, m) \mapsto \lambda m,$$

where

$$\lambda m = \begin{cases} \underbrace{m + m + \dots + m}_{\lambda \text{ times}} & \text{if } \lambda > 0\\ \underbrace{(-m) + (-m) + \dots + (-m)}_{-\lambda \text{ times}} & \text{if } \lambda < 0 \\ 0 & \text{if } \lambda = 0 \end{cases}$$

First note that Ω is indeed a well-defined map from \mathbb{Z}^n to $(\mathbb{Z}_q)^n$. Let $x = (\lambda_1, \lambda_2, ..., \lambda_n), y = (\mu_1, \mu_2, ..., \mu_n) \in \mathbb{Z}^n$ and $\lambda \in \mathbb{Z}$. Then

$$\Omega(x+_1 y) = \Omega(\lambda_1 + \mu_1, \lambda_2 + \mu_2, ..., \lambda_n + \mu_n)$$

= $(\overline{\lambda_1 + \mu_1}, \overline{\lambda_2 + \mu_2}, ..., \overline{\lambda_n + \mu_n}),$

where for each $i \in \{1, 2, ..., n\}$, $\overline{\lambda_i + \mu_i} = (\lambda_i + \mu_i) \pmod{q}$.

Now, for $i \in \{1, 2, ..., n\}$, we have that

$$\overline{\lambda_i + \mu_i} = (\lambda_i + \mu_i) \pmod{q}$$
$$= (\lambda_i \pmod{q} + \mu_i \pmod{n}) \pmod{q}$$
$$= (\overline{\lambda_i} + \overline{\mu_i}) \pmod{q}$$
$$= \overline{\lambda_i} \oplus_q \overline{\mu_i}$$

So from above, we get that

$$\Omega(x+_1 y) = (\overline{\lambda_1 + \mu_1}, \overline{\lambda_2 + \mu_2}, ..., \overline{\lambda_n + \mu_n})$$

= $(\overline{\lambda_1} \oplus_q \overline{\mu_1}, \overline{\lambda_2} \oplus_q \overline{\mu_2}, ..., \overline{\lambda_n} \oplus_q \overline{\mu_n})$
= $(\overline{\lambda_1}, \overline{\lambda_2}, ..., \overline{\lambda_n}) +_2 (\overline{\mu_1}, \overline{\mu_2}, ..., \overline{\mu_n})$
= $\Omega(x) +_2 \Omega(y)$

Now, first suppose that $\lambda > 0$.

$$\lambda \odot_1 x = \underbrace{x + 1 x + 1 \dots + 1 x}_{\lambda \text{ times}}$$

$$= \underbrace{(\lambda_1 + \lambda_1 + \dots + \lambda_1)}_{\lambda \text{ times}}, \underbrace{\lambda_2 + \lambda_2 + \dots + \lambda_2}_{\lambda \text{ times}}, \dots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \lambda_n + \dots + \lambda_n}_{\lambda \text{ times}}, \ldots, \underbrace{\lambda_n + \dots +$$

Now,

$$\Omega(\lambda \odot_1 x) = \Omega(\lambda \lambda_1, \lambda \lambda_2, ..., \lambda \lambda_n)$$
$$= (\overline{\lambda \lambda_1}, \overline{\lambda \lambda_2}, ..., \overline{\lambda \lambda_n})$$

Now, for each $i \in \{1, 2, ..., n\}$,

$$\begin{aligned} \overline{\lambda\lambda_i} &= (\lambda\lambda_i) (\text{mod } q) \\ &= (((\lambda(\text{mod } q))(\lambda_i(\text{mod } q))))(\text{mod } q) \\ &= (\overline{\lambda} \ \overline{\lambda_i})(\text{mod } q) \\ &= (\lambda\lambda_i)(\text{mod } q) \\ &= (\lambda\lambda_i)(\text{mod } q). \end{aligned}$$
$$\lambda \odot_2 \Omega(x) &= \underbrace{\Omega(x) +_2 \Omega(x) +_2 \dots +_2 \Omega(x)}_{\lambda \text{ times}} \\ &= (\underbrace{\overline{\lambda_1} \oplus_q \ \overline{\lambda_1} \oplus_q \dots \oplus_q \overline{\lambda_1}}_{\lambda \text{ times}}, \underbrace{\overline{\lambda_2} \oplus_q \dots \oplus_q \overline{\lambda_2}}_{\lambda \text{ times}}, \dots, \underbrace{\overline{\lambda_n} \oplus_q \ \overline{\lambda_n} \oplus_q \dots \oplus_q \overline{\lambda_n}}_{\lambda \text{ times}}) \end{aligned}$$

Now, let
$$i \in \{1, 2, ..., n\}$$
. Then we get that

$$\underbrace{\overline{\lambda_i} \oplus_q \overline{\lambda_i} \oplus_q ... \oplus_q \overline{\lambda_i}}_{\lambda \text{ times}} = \underbrace{(\overline{\lambda_i} + \overline{\lambda_i} + ... + \overline{\lambda_i})}_{\lambda \text{ times}} (\text{mod } q)$$

$$= \underbrace{(\lambda_i + \lambda_i + ... + \lambda_i)}_{\lambda \text{ times}} (\text{mod } q)$$

$$= (\lambda \lambda_i) (\text{mod } q)$$

So we have proved that $\Omega(\lambda \odot_1 x) = \lambda \odot_2 \Omega(x)$ for $\lambda > 0$. The above equation holds trivially for $\lambda = 0$. Now, suppose $\lambda < 0$. Then

$$\begin{split} \lambda \odot_1 x &= \underbrace{(-x) +_1 (-x) +_1 \dots +_1 (-x)}_{-\lambda \text{ times}} \\ &= \underbrace{((-\lambda_1) + (-\lambda_1) + \dots + (-\lambda_1)}_{-\lambda \text{ times}}, \underbrace{(-\lambda_2) + (-\lambda_2) + \dots + (-\lambda_2)}_{-\lambda \text{ times}}, \dots, \\ \underbrace{(-\lambda_n) + (-\lambda_n) + \dots + (-\lambda_n))}_{-\lambda \text{ times}} \\ &= ((-\lambda)(-\lambda_1), (-\lambda)(-\lambda_2), \dots, (-\lambda)(-\lambda_n)) \\ &= (\lambda\lambda_1, \lambda\lambda_2, \dots, \lambda\lambda_n) \end{split}$$

Then as shown above, we will have

$$\Omega(\lambda \odot_1 x) = (\overline{\lambda \lambda_1}, \overline{\lambda \lambda_2}, ..., \overline{\lambda \lambda_n}),$$

where for each $i \in \{1, 2, ..., n\}, \overline{\lambda \lambda_i} = (\lambda \lambda_i) \pmod{q}$.

Using the fact that Ω is a group homomorphisms from the additive group \mathbb{Z}^n to the additive group $(\mathbb{Z}_q)^n$, we get that

$$\begin{split} \lambda \odot_2 \Omega(x) &= \underbrace{(-\Omega(x)) +_2 (-\Omega(x)) +_2 \dots +_2 (-\Omega(x))}_{-\lambda \text{ times}} \\ &= \underbrace{\Omega(-x) +_2 \Omega(-x) +_2 \dots +_2 \Omega(-x)}_{-\lambda \text{ times}} \\ &= \underbrace{(\overline{-\lambda_1} \oplus_q \overline{-\lambda_1} \oplus_q \dots \oplus_q \overline{-\lambda_1}, \overline{-\lambda_2} \oplus_q \overline{-\lambda_2} \oplus_q \dots \oplus_q \overline{-\lambda_2}}_{-\lambda \text{ times}}, \dots, \\ &\underbrace{\overline{-\lambda_n} \oplus_q \overline{-\lambda_n} \oplus_q \dots \oplus_q \overline{-\lambda_n}}_{-\lambda \text{ times}} \end{split}$$

Now, for
$$i \in \{1, 2, ..., n\}$$
,

$$\underbrace{-\lambda_i \oplus_q -\lambda_i \oplus_q \dots \oplus_q -\lambda_i}_{-\lambda \text{ times}} = \underbrace{(-\lambda_i) + (-\lambda_i) + \dots + (-\lambda_i)}_{-\lambda \text{ times}}) \pmod{q}$$

$$= \underbrace{((-\lambda_i) + (-\lambda_i) + \dots + (-\lambda_i))}_{-\lambda \text{ times}} \pmod{q}$$

$$= ((-\lambda)(-\lambda_i)) \pmod{q}$$

$$= (\lambda\lambda_i) \pmod{q}$$

So we have proved that $\Omega(\lambda \odot_1 x) = \lambda \odot_2 \Omega(x)$ for $\lambda < 0$. In all possible cases, we have proved that $\Omega(\lambda \odot_1 x) = \lambda \odot_2 \Omega(x)$ for all $\lambda \in \mathbb{Z}$. Therefore, Ω is a \mathbb{Z} -module homomorphism from the \mathbb{Z} -module \mathbb{Z}^n to the \mathbb{Z} -module $(\mathbb{Z}_q)^n$. Hence proved. **Lemma 4.4.** (i) If W is a \mathbb{Z} -submodule of \mathbb{Z}^n , then $\Omega(W)$ is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$.

(ii) If V is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$, then $\Omega^{-1}(V) = \{ w \in \mathbb{Z}^n \mid \Omega(w) \in V \}$ is a \mathbb{Z} -submodule of \mathbb{Z}^n .

Proof. (i) Let W be a \mathbb{Z} -submodule of the \mathbb{Z} -module \mathbb{Z}^n .

Since $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$, therefore, $\Omega(W) \subseteq (\mathbb{Z}_q)^n$. Also, $\Omega(W)$ is non-empty as W being a \mathbb{Z} -submodule of \mathbb{Z}^n is non-empty.

Let $x, y \in \Omega(W)$ and $\lambda \in \mathbb{Z}$.

Then $x = \Omega(w)$ and $y = \Omega(w')$ for some $w, w' \in W$.

 $x+y = \Omega(w) + \Omega(w') = \Omega(w+w')$ as $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$ is a \mathbb{Z} -module homomorphism and $W \subseteq \mathbb{Z}^n$.

As $w, w' \in W$ and W is a Z-submodule of \mathbb{Z}^n , therefore, $w + w' \in W$. This implies that $x + y = \Omega(w + w') \in \Omega(W)$.

Further, since $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$ is a \mathbb{Z} -module homomorphism, therefore, $\lambda x = \lambda \Omega(w) = \Omega(\lambda w)$.

Since $\lambda \in \mathbb{Z}$, $w \in W$ and W is a Z-submodule of \mathbb{Z}^n , therefore, $\lambda w \in W$.

This implies that $\lambda x = \Omega(\lambda w) \in \Omega(W)$.

Therefore, $\Omega(W)$ is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$.

Hence proved.

(ii) Let V be a Z-submodule of the Z-module $(\mathbb{Z}_q)^n$.

Clearly, $\Omega^{-1}(V)$ is a subset of \mathbb{Z}^n . Also, $\Omega^{-1}(V)$ is non-empty as V being a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$ is non-empty (note that $0 \in V$ and $\Omega(0) = 0$).

Let $w, w' \in \Omega^{-1}(V)$ and $\lambda \in \mathbb{Z}$.

Then $w, w' \in \mathbb{Z}^n$ such that $\Omega(w), \Omega(w') \in V$.

But then the fact that V is a Z-submodule of $(\mathbb{Z}_q)^n$ implies that $\Omega(w) + \Omega(w') \in V$. Since by Lemma 4.3, $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$ is a Z-module homomorphism and $w, w' \in \mathbb{Z}^n$, so

 $\Omega(w+w') = \Omega(w) + \Omega(w')$. Therefore, from above, we get that $\Omega(w+w') \in V$.

 $\Rightarrow w + w' \in \Omega^{-1}(V).$

Also, since $\lambda \in \mathbb{Z}$, $\Omega(w) \in V$ and V is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$, therefore, $\lambda \Omega(w) \in V$. Since by Lemma 4.3, $\Omega : \mathbb{Z}^n \to (\mathbb{Z}_q)^n$ is a \mathbb{Z} -module homomorphism, $w \in \mathbb{Z}^n$ and $\lambda \in \mathbb{Z}$,

therefore, $\Omega(\lambda w) = \lambda \Omega(w)$. Therefore, from above, we get that $\Omega(\lambda w) \in V$. $\Rightarrow \lambda w \in \Omega^{-1}(V)$. Therefore, $\Omega^{-1}(V)$ is a \mathbb{Z} -submodule of \mathbb{Z}^n .

Lemma 4.5. A subset W is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$ if and only if W is a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$.

Proof. Since $(\mathbb{Z}_q)^n$ is a commutative unital ring with respect to the usual operations of componentwise addition module q and multiplication module q, therefore, $(\mathbb{Z}_q)^n$ is also a \mathbb{Z}_q -module with respect to the usual operation of componentwise scalar multiplication module q.

Let \odot_2 denote the scalar multiplication of the \mathbb{Z} -module $(\mathbb{Z}_q)^n$ and \odot denote the scalar multiplication of the \mathbb{Z}_q -module $(\mathbb{Z}_q)^n$.

Hence proved.

First, let W be a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$. We show that W is a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$. W being a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$ is a non-empty subset of $(\mathbb{Z}_q)^n$. Let $w, w' \in W$ and $\lambda \in \mathbb{Z}_q$. Suppose $w = (\lambda_1, \lambda_2, ..., \lambda_n)$ for some $\lambda_i \in \mathbb{Z}_q$ $(1 \le i \le n)$. Since $w, w' \in W$ and W is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$, therefore, $w + w' \in W$. $\lambda \in \mathbb{Z}_q$ implies that $\lambda \in \mathbb{Z}$. Now,

$$\lambda \odot_2 w = \underbrace{w +_2 w +_2 \dots +_2 w}_{\lambda \text{ times}} \\ = \underbrace{\left(\underbrace{\lambda_1 \oplus_q \lambda_1 \oplus_q \dots \oplus_q \lambda_1}_{\lambda \text{ times}}, \underbrace{\lambda_2 \oplus_q \lambda_2 \oplus_q \dots \oplus_q \lambda_2}_{\lambda \text{ times}}, \dots, \underbrace{\lambda_n \oplus_q \lambda_n \oplus_q \dots \oplus_q \lambda_n}_{\lambda \text{ times}}\right)}_{\lambda \text{ times}}$$

Also,

$$\lambda \odot w = (\lambda \otimes_q \lambda_1, \lambda \otimes_q \lambda_2, ..., \lambda \otimes_q \lambda_n)$$

Claim: For each $i \in \{1, 2, ..., n\}$, $\underbrace{\lambda_i \oplus_q \lambda_i \oplus_q ... \oplus_q \lambda_i}_{\lambda \text{ times}} = \lambda \otimes_q \lambda_1$.

So let $i \in \{1, 2, ..., n\}$. Then we get

$$\underbrace{\lambda_i \oplus_q \lambda_i \oplus_q \dots \oplus_q \lambda_i}_{\lambda \text{ times}} = \underbrace{(\lambda_i + \lambda_i + \dots + \lambda_i)}_{\lambda \text{ times}} (\text{mod } q)$$
$$= (\lambda \lambda_i) (\text{mod } q)$$
$$= \lambda \otimes_q \lambda_i$$

So we have proved that $\lambda \odot_2 w = \lambda \odot w$.

Since W is a \mathbb{Z} -submodule, therefore, $\lambda \odot_2 w \in W$.

Further, since $\lambda \odot_2 w = \lambda \odot w$, therefore, $\lambda \odot w \in W$.

Since $w, w' \in W$ and $\lambda \in \mathbb{Z}_q$ are arbitrary, therefore, $w + w', \lambda \odot w \in W$ for all $w, w' \in W$ and $\lambda \in \mathbb{Z}_q$.

Therefore, W is also a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$. Conversely, let W be a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$. We show that W is also a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$. Being a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$, W is a non-empty subset of $(\mathbb{Z}_q)^n$. Let $w, w' \in W$ and $\lambda \in \mathbb{Z}$. Since W is a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$, therefore, $w + w' \in W$. Since $w \in W \subseteq (\mathbb{Z}_q)^n$, so $w = (\lambda_1, \lambda_2, ..., \lambda_n)$ for some $\lambda_i \in \mathbb{Z}_q$ $(1 \le i \le n)$. Now, if $\lambda > 0$, then

$$\lambda \odot_2 w = \underbrace{w +_2 w +_2 \dots +_2 w}_{\lambda \text{ times}} = \underbrace{\left(\underbrace{\lambda_1 \oplus_q \lambda_1 \oplus_q \dots \oplus_q \lambda_1}_{\lambda \text{ times}}, \underbrace{\lambda_2 \oplus_q \lambda_2 \oplus_q \dots \oplus_q \lambda_2}_{\lambda \text{ times}}, \dots, \underbrace{\lambda_n \oplus_q \lambda_n \oplus_q \dots \oplus_q \lambda_n}_{\lambda \text{ times}}\right)}_{\lambda \text{ times}}.$$

 $\overline{\lambda} = \lambda \pmod{q}$. Then

$$\overline{\lambda} \odot w = (\overline{\lambda} \otimes_q \lambda_1, \overline{\lambda} \otimes_q \lambda_2, ..., \overline{\lambda} \otimes_q \lambda_n).$$

Claim: For each $i \in \{1, 2, ..., n\}$, $\underbrace{\lambda_i \oplus_q \lambda_i \oplus_q ... \oplus_q \lambda_i}_{\lambda \text{ times}} = \overline{\lambda} \otimes_q \lambda_i$.

We get

$$\underbrace{\lambda_i \oplus_q \lambda_i \oplus_q \dots \oplus_q \lambda_i}_{\lambda \text{ times}} = \underbrace{(\lambda_i + \lambda_i + \dots + \lambda_i)}_{\lambda \text{ times}} (\text{mod } q)$$
$$= (\lambda \lambda_i) (\text{mod } q)$$
$$= (\overline{\lambda \lambda_i}) (\text{mod } q)$$
$$= \overline{\lambda} \otimes_q \lambda_i$$

because $\overline{\lambda_i} = \lambda_i$ as $\lambda_i \in \mathbb{Z}_q$.

So we have proved that for $\lambda > 0$, $\lambda \odot_2 w = \overline{\lambda} \odot w$. For $\lambda = 0$, the result holds trivially. If $\lambda < 0$, then

$$\lambda \odot_2 w = \underbrace{(-w) +_2 (-w) +_2 \dots +_2 (-w)}_{-\lambda \text{ times}} = \underbrace{(\underbrace{(-\lambda_1) \oplus_q (-\lambda_1) \oplus_q \dots \oplus_q (-\lambda_1)}_{-\lambda \text{ times}}, \underbrace{(-\lambda_n) \oplus_q (-\lambda_n) \oplus_q \dots \oplus_q (-\lambda_n)}_{-\lambda \text{ times}}, \underbrace{(-\lambda_n) \oplus_q (-\lambda_n) \oplus_q \dots \oplus_q (-\lambda_n)}_{-\lambda \text{ times}})$$

Claim: For each $i \in \{1, 2, ..., n\}$, $\underbrace{(-\lambda_i) \oplus_q (-\lambda_i) \oplus_q ... \oplus_q (-\lambda_i)}_{-\lambda \text{ times}} = \overline{\lambda} \otimes_q \lambda_i$.

We get

$$\underbrace{(-\lambda_i) \oplus_q (-\lambda_i) \oplus_q \dots \oplus_q (-\lambda_i)}_{-\lambda \text{ times}} = \underbrace{((-\lambda_i) + (-\lambda_i) + \dots + (-\lambda_i))}_{-\lambda \text{ times}} (\text{mod } q)$$
$$= ((-\lambda)(-\lambda_i))(\text{mod } q)$$
$$= (\lambda\lambda_i)(\text{mod } q)$$
$$= (\overline{\lambda\lambda_i})(\text{mod } q)$$
$$= \overline{\lambda} \otimes_q \lambda_i$$

because $\overline{\lambda_i} = \lambda_i$ as $\lambda_i \in \mathbb{Z}_q$.

So we have proved that for $\lambda < 0$, $\lambda \odot_2 w = \overline{\lambda} \odot w$. Therefore, for any λ , $\lambda \odot_2 w = \overline{\lambda} \odot w$. But since $\overline{\lambda} \in \mathbb{Z}_q$, $w \in W$ and W is a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$, therefore, $\overline{\lambda} \odot w \in W$. And then the fact that $\lambda \odot_2 w = \overline{\lambda} \odot w$ implies that $\lambda \odot_2 w \in W$. Therefore, W is also a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$.

Suppose W is an IDD code and \mathcal{E} is its corresponding equivalent IDD code in \mathbb{Z}^n . Then \mathcal{E} is a \mathbb{Z} -submodule of the \mathbb{Z} -module $(\mathbb{Z})^n$. By Lemma 4.4 (i), $\Omega(\mathcal{E})$ is a \mathbb{Z} -submodule of $(\mathbb{Z}_q)^n$. Then by Lemma 4.5, $\Omega(\mathcal{E})$ is a \mathbb{Z}_q -submodule of $(\mathbb{Z}_q)^n$. Hence, $\Omega(\mathcal{E})$ is a code over \mathbb{Z}_q and in $(\mathbb{Z}_q)^n$. So we have the following definition.

Definition 4.6. Let q be a positive integer. With the notations of Subsection 4.1, let W be an IDD code and \mathcal{E} be its equivalent IDD code in \mathbb{Z}^n . Then the \mathbb{Z}_q -submodule $\Omega(\mathcal{E})$ of $(\mathbb{Z}_q)^n$ is called Hom-IDD code in $(\mathbb{Z}_q)^n$.

4.4 Example

Example 4.7. Let $K = \mathbb{Q}(\zeta)$, ζ primitive p^{th} -root of unity, where p = 17. Let $\sigma(\zeta) = \zeta$ and $\tau(\zeta) = \zeta^3$. Then by Theorem 3.8, the map $D: O_K \to O_K$ given by

$$D(\zeta) = 1 + \zeta + \zeta^{2} + \zeta^{3} + \zeta^{5} + \zeta^{7} + \zeta^{8} + \zeta^{11}$$

is a (σ, τ) -derivation of $O_K = \mathbb{Z}[\zeta]$. Below, we have obtained (binary) Hom-IDD codes of length n = 16 over the finite field \mathbb{Z}_2 with various parameters, using the construction discussed above.

Basis S_i	Code Description: $[n, k, d]$	Code Properties	Dual Code Description: [n, k, d]
S_1	[16, 8, 4]	LCD	[16, 8, 4]
S_2	[16, 8, 4]	non-LCD	[16, 8, 4]
S_3	[16, 7, 4]	LCD	[16, 9, 3]
S_4	[16, 9, 3]	non-LCD	[16, 7, 4]
S_5	[16, 7, 4]	non-LCD	[16, 9, 3]
S_6	[16, 6, 4]	non-LCD	[16, 10, 3]
S_7	[16, 5, 5]	non-LCD	[16, 11, 2]
S_8	[16, 4, 7]	non-LCD	[16, 12, 2]
S_9	[16, 5, 5]	LCD	[16, 11, 2]
S_{10}	[16, 8, 3]	non-LCD	[16, 8, 4]
S_{11}	[16, 9, 3]	LCD	[16, 7, 4]
S_{12}	[16, 8, 4]	LCD	[16, 8, 3]
S_{13}	[16, 6, 5]	LCD	[16, 10, 3]

Basis S_i
$S_1 = \{ D(\zeta), D(\zeta^2), D(\zeta^4), D(\zeta^5), D(\zeta^6), D(\zeta^7), D(\zeta^{10}), D(\zeta^{13}) \}$
$S_2 = \{ D(\zeta), D(\zeta^2), D(\zeta^4), D(\zeta^5), D(\zeta^6), D(\zeta^7), D(\zeta^9), D(\zeta^{12}) \}$
$S_3 = \{ D(\zeta), D(\zeta^2), D(\zeta^5), D(\zeta^6), D(\zeta^7), D(\zeta^9), D(\zeta^{12}) \}$
$S_4 = \{ D(\zeta), D(\zeta^2), D(\zeta^5), D(\zeta^6), D(\zeta^7), D(\zeta^9), D(\zeta^{10}), D(\zeta^{12}), D(\zeta^{14}) \}$
$S_5 = \{ D(\zeta), D(\zeta^2), D(\zeta^6), D(\zeta^7), D(\zeta^9), D(\zeta^{12}), D(\zeta^{14}) \}$
$S_6 = \{ D(\zeta), D(\zeta^2), D(\zeta^6), D(\zeta^7), D(\zeta^9), D(\zeta^{12}) \}$
$S_7 = \{ D(\zeta), D(\zeta^2), D(\zeta^5), D(\zeta^6), D(\zeta^7) \}$
$S_8 = \{ D(\zeta), D(\zeta^2), D(\zeta^6), D(\zeta^7) \}$
$S_9 = \{ D(\zeta), D(\zeta^2), D(\zeta^6), D(\zeta^7), D(\zeta^{15}) \}$
$S_{10} = \{ D(\zeta), D(\zeta^2), D(\zeta^6), D(\zeta^7), D(\zeta^{10}), D(\zeta^{13}), D(\zeta^{14}), D(\zeta^{15}) \}$
$S_{11} = \{ D(\zeta), D(\zeta^2), D(\zeta^4), D(\zeta^5), D(\zeta^6), D(\zeta^{10}), D(\zeta^{13}), D(\zeta^{14}), D(\zeta^{15}) \}$
$S_{12} = \{ D(\zeta), D(\zeta^4), D(\zeta^5), D(\zeta^6), D(\zeta^{10}), D(\zeta^{13}), D(\zeta^{14}), D(\zeta^{15}) \}$
$S_{13} = \{ D(\zeta), D(\zeta^4), D(\zeta^5), D(\zeta^6), D(\zeta^{10}), D(\zeta^{13}) \}$

5 Conclusion

In this article, we studied the (σ, τ) -derivations of number rings and their applications to coding theory. First, in Section 2, we obtained some useful results for the forthcoming sections. In Section 3, we studied (σ, τ) -derivations of the rings of algebraic integers of quadratic and cyclotomic number fields, and that of the bi-quadratic number ring. In quadratic case, we reformulated and reinvented the results formed in [5]. We obtained a characterization in Subsection 3.2 for a Z-linear map vanishing on unity to be a (σ, τ) -derivation of the ring of algebraic integers of a p^{th} -cyclotomic number field. We also conjectured a criterion under which a (σ, τ) -derivation of the ring of algebraic integers of a p^{th} cyclotomic number field is inner. In Subsection 3.3, we explicitly obtained all (σ, τ) -derivations of a bi-quadratic number ring. We also obtained a criterion under which a (σ, τ) -derivation of a bi-quadratic number ring is inner. Finally, in Section 4, we discussed the applications of our work in coding theory by giving the notion of Hom-IDD codes.

Declaration of Interest Statement

The authors report there are no competing interests to declare.

References

- [1] Aleksandr Alekseev, Andronick Arutyunov, and Sergei Silvestrov. On (σ, τ) -derivations of group algebra as category characters. arXiv preprint arXiv:2008.00390, 2020.
- [2] Mehsin Jabel Atteya. New types of permuting n-derivations with their applications on associative rings. *Symmetry*, 12(1):46, 2019.
- [3] M. Bresar. On the composition of (α, β) -derivations of rings and applications to von Neumann algebras. Acta Sct. Math, 56:369–375, 1992.

- [4] Muhammad Anwar Chaudhry and Zafar Ullah. On generalized (α, β) -derivations on lattices. *Quaestiones Mathematicae*, 34(4):417–424, 2011.
- [5] Dishari Chaudhuri. A note on (σ, τ) -derivations on commutative algebras. arXiv preprint arXiv:1912.12812, 2019.
- [6] Leo Creedon and Kieran Hughes. Derivations on group algebras with coding theory applications. *Finite Fields and Their Applications*, 56:247–265, 2019.
- [7] Boucher Delphine and Felix Ulmer. Linear codes using skew polynomials with automorphisms and derivations. *Designs, codes and cryptography*, 70:405–431, 2014.
- [8] M. Haetinger, C. Ashraf and S. Ali. On higher derivations: a survey. Int. J. Math., Game Theory and Algebra, 18:359–379, 2011.
- [9] N. Jacobson. Structure of Rings. CP37. American Mathematical Society, 1964.
- [10] L. Kamali Ardakani and Bijan Davvaz. f-derivations and (f,g)-derivations of MValgebras. Journal of Algebraic Systems, 1(1):11–31, 2013.
- [11] Kyung Ho Kim. On (f,g)-derivations of incline algebras. Journal of the Chungcheong Mathematical Society, 27(4):643–649, 2014.
- [12] Slawomir Klimek and Matt McBride. Unbounded derivations in algebras associated with monothetic groups. Journal of the Australian Mathematical Society, 111(3):345– 371, 2021.
- [13] Daniel A. Marcus. *Number fields*. Universitext. Springer, 2nd edition, 2018.
- [14] Ashraf Mohammad, Ali Shakir, and Haetinger Claus. On derivations in rings and their applications. The Aligarh Bull of Math, 25(2):79–107, 2006.
- [15] G Muhiuddin and Abdullah M Al-Roqi. On (α, β) -derivations in BCI-algebras. *Discrete Dynamics in Nature and Society*, 2012:11, 2012.
- [16] AŞCI Mustafa and Şahin Ceran. Generalized (f, g)-derivations of lattices. Mathematical Sciences and Applications E-Notes, 1(2):56–62, 2013.
- [17] Mohd Arif Raza and Nadeem Ur Rehman. On prime and semiprime rings with generalized derivations and non-commutative Banach algebras. *Proceedings-Mathematical Sciences*, 126(3):389–398, 2016.
- [18] Ian Stewart and David Tall. Algebraic Number Theory and Fermat's Last Theorem. A. K. Peters, 3rd edition, 2002.