

Pin Classes I: Growth Rates and Bounds

Ben Jarvis*

School of Mathematics and Statistics
The Open University, UK

December 6, 2024

Abstract

Pin sequences play an important role in the structural study of permutation classes. In this paper, we study the permutation classes that comprise all the finite subpermutations contained in an infinite pin sequence. We prove that these permutation classes have proper growth rates and establish a procedure for calculating these growth rates.

1 Introduction

Since their introduction by Brignall, Huczynska and Vatter [8], pin sequences have proven to be objects of considerable interest in the study of permutation classes, especially in connection with simple permutations and infinite antichains. Bassino, Bouvel and Rossin [2] showed that the class of all pin sequences has a rational generating function and established its growth rate, and recently, Brignall and Vatter [7] used pin sequences to construct uncountably many well-quasi-ordered permutation classes with distinct enumeration sequences.

Given an (infinite) pin sequence, the *pin class* is formed from all finite permutations that are contained in the sequence. In this paper we take up the study of pin classes in a more systematic manner than has previously been attempted. Our main result is that pin classes have *proper* growth rates, not merely upper growth rates as guaranteed by the Marcus-Tardos Theorem [12]. We also describe a procedure for finding the growth rate of a given pin class in terms of the recurrent subfactors of the defining pin sequence.

Pin classes are most naturally situated in the context of **centred permutations**, which we define in Section 2. We shall see in Section 3 that this change of perspective allows us to state a particularly nice structure theorem for *recurrent* pin classes in terms of the \boxplus -sum (an operation specific to centred permutations). This gives us an extraordinarily large family of permutation classes whose generating functions we can find by solving a simple combinatorial problem concerning subfactors of the defining pin sequence. In Section 4 we extend this theory to pin

*Ben.Jarvis@open.ac.uk

classes defined by non-recurrent pin sequences: in this case we can no longer find generating functions but we can still prove the existence of growth rates and find these by considering a centred class called the \boxplus -interior of a pin class.

2 Centred Permutation Classes

We refer the reader to Bevan [4] for background information and basic definitions concerning permutations and permutation classes.

2.1 Centred Permutations

Our main focus in this paper will be a particular method for converting a binary sequence into a permutation class; the resulting permutation classes will be known as *pin classes*. It shall transpire that pin classes are in fact most naturally understood not as classes of permutations in the ordinary sense, but as *centred permutations*. A centred permutation is essentially a permutation with an extra point, designated as the *origin*, which does not contribute to the length of the permutation. A formal definition follows:

Definition 2.1 (Centred Permutations). A centred permutation of length n is a permutation of length $n + 1$ in which one point is designated as the origin, which does not contribute to the length of the centred permutation.



Figure 1: Two centred permutations of length 3. Note that a centred permutation of length 3 consists of 4 points (no two of which share either an x - or y -coordinate) in the plane, of which one is designated as the origin, which does not contribute to the length. Here we denote the origin with an empty circle, whereas the ‘true’ points are represented by solid circles.

We usually distinguish between centred and uncentred permutations by placing a circle in the superscript: so π is a regular permutation and π° is a centred permutation. (Similarly \mathcal{C}° is a class of centred permutations, whereas \mathcal{C} is a class of uncentred permutations.)

Comment 2.2. By drawing horizontal and vertical axes through the origin, we can identify a centred permutation with a 2-by-2-gridded permutation; we shall usually draw these axes on in addition to the origin for clarity. This splits the plane into four *quadrants*, numbered 1 to 4 anticlockwise - see Fig. 2. Centred permutations containing points in only one of these four quadrants will play a special role in the theory - we call these *one-quadrant permutations*.

Every centred permutation σ° of length n is naturally associated with an *underlying* permutation σ of length n , obtained by removing the origin point from σ° . We also have the *filled-in*




Figure 2: The standard quadrant numbering: the origin point of a centred permutation splits the plane into four quadrants, numbered anticlockwise.

permutation σ^\bullet , the permutation of length $n + 1$ obtained by replacing the origin of σ° with a true point. Though we usually study centred permutation classes in order to understand their associated underlying permutation classes, it is the latter of these notions which provides us with the most convenient notation for a centred permutation:

Definition 2.3 (One-Line Notation for Centred Permutations). Suppose that σ° is a centred permutation of length n , with corresponding filled-in permutation σ^\bullet of length $n + 1$. Suppose that the k th entry of σ^\bullet is the origin of σ° . Then we can write σ° in one-line notation as follows:

$$\sigma^\circ = \sigma^\bullet(1)\sigma^\bullet(2)\dots\underline{\sigma^\bullet(k)}\dots\sigma^\bullet(n)\sigma^\bullet(n + 1)$$

In other words, this is the one-line notation for σ^\bullet with the centre point underlined. See Fig. 3 for an example.




Figure 3: The centred permutation $426\underline{3}51$.

Comment 2.4. When dealing with centred permutations, the origin does not count as part of the length of the permutation, but *does* need to be in the same place for two permutations to be the same (see Fig. 4 for an example).




Figure 4: The centred permutations $14\underline{2}3$ and $1\underline{2}43$ are both of length 3 (the origin point doesn't count) and both have the same underlying (uncentred) permutation 132, but they are distinct as centred permutations as the origins are in a different place.

Analogously with the uncentred case we define the notion of centred permutation containment:

Definition 2.5 (Centred Permutation Containment). Suppose that σ° is a centred permutation of length n and π° is a centred permutation of length $m \leq n$. We say that π° is *contained* in

σ° , denoted $\pi^\circ \leq \sigma^\circ$, if π^\bullet can be embedded in σ^\bullet in such a way that the origins match up. Explicitly, if

$$\pi^\circ = \pi^\bullet(1)\pi^\bullet(2)\dots\underline{\pi^\bullet(k)}\dots\pi^\bullet(m)\pi^\bullet(m+1)$$

and

$$\sigma^\circ = \sigma^\bullet(1)\sigma^\bullet(2)\dots\underline{\sigma^\bullet(l)}\dots\sigma^\bullet(n)\sigma^\bullet(n+1)$$

then $\pi^\circ \leq \sigma^\circ$ if and only if there is an ascending sequence of $m+1$ indices

$$1 \leq i_1 < i_2 < \dots < i_{m+1} \leq n+1$$

such that the sequence $\sigma^\bullet(i_1)\sigma^\bullet(i_2)\dots\sigma^\bullet(i_{m+1})$ is order-isomorphic to π^\bullet and $i_k = l$.

As with regular permutation containment, centred permutation containment is much easier to understand graphically - see Fig. 5 for an example. This is one of the reasons that we generally prefer to draw centered permutations out rather than work with one-line notation.




Figure 5: The centred permutation $14\underline{2}3$ is contained in $71834\underline{5}26$; the relevant embedding is shown by the points marked in red in the diagram of $71834\underline{5}26$. Note that the embedding must make the origins match up.

Centred permutation containment, of course, immediately allows us to define the centred analogue of a permutation class:

Definition 2.6 (Centred Permutation Class). A **centred permutation class** is a non-empty set \mathcal{C}° of centred permutations which is downward-closed under centred permutation containment. We call a centred permutation class **proper** if the set of all underlying (uncentred) permutations of centred permutations in \mathcal{C}° does not contain every (uncentred) permutation.

Note that $\sigma^\circ \leq \pi^\circ$ implies that $\sigma \leq \pi$. This means that, for every centred permutation class \mathcal{C}° , the set \mathcal{C} consisting of the underlying permutations of \mathcal{C}° is itself an uncentred permutation class. We call \mathcal{C} the **underlying** permutation class of \mathcal{C}° . We wish to use centred permutation classes as a means to study regular permutation classes: there are many interesting permutation classes that are most succinctly described as the underlying class of some given centred permutation class. In particular, we are interested in the growth rates of (centred and uncentred) permutation classes, so we recall the following:

Definition 2.7 (Growth Rates). Let (C_n) be a sequence of non-negative integers. The **upper** and **lower growth rates** of (C_n) are defined, respectively, to be

$$\overline{\text{gr}}((C_n)) = \limsup_{n \rightarrow \infty} \sqrt[n]{C_n}, \quad \text{and} \quad \underline{\text{gr}}((C_n)) = \liminf_{n \rightarrow \infty} \sqrt[n]{C_n},$$

when these quantities exist. If both quantities exist and $\overline{\text{gr}}((C_n)) = \underline{\text{gr}}((C_n))$, then the sequence has a **(proper) growth rate**, typically denoted $\text{gr}(C_n)$.

If \mathcal{C} is a (centred or uncentred) permutation class, we define \mathcal{C}_n to be the set of permutations of length n in \mathcal{C} and write $C_n = |\mathcal{C}_n|$. We call C_n the **enumeration sequence** of \mathcal{C}_n and say that \mathcal{C}_n has (upper, lower, proper) growth rate ρ if ρ is the (upper, lower, proper) growth rate of the enumeration sequence. We write $\text{gr}(\mathcal{C})$ for $\text{gr}(C_n)$, and analogously for upper and lower growth rates.

We plan to use centred permutation classes to study the growth rates of their underlying (uncentred) permutation classes, so the following is crucial to note:

Proposition 2.8. *Suppose that \mathcal{C}° is a proper centred permutation class whose underlying permutation class is \mathcal{C} . Then:*

1. *The upper growth rate of \mathcal{C}° exists and is equal to the upper growth rate of \mathcal{C} .*
2. *The lower growth rate of \mathcal{C}° exists and is equal to the lower growth rate of \mathcal{C} .*
3. *The proper growth rate $\text{gr}(\mathcal{C}^\circ)$ exists if and only if $\text{gr}(\mathcal{C})$ exists; if so then $\text{gr}(\mathcal{C}^\circ) = \text{gr}(\mathcal{C})$.*

Proof. As \mathcal{C}° is a proper centred permutation class, \mathcal{C} is, by definition, a proper permutation class. Hence by the Marcus-Tardos Theorem [12], \mathcal{C} has an upper growth rate. On noting that a centred permutation can be identified with a 2-by-2 gridded permutation, the fact that \mathcal{C}° has the same upper and lower growth rates follows immediately from Vatter [14, Proposition 2.1]. (Informally, there are $(n+1)^2$ ways of placing a centre in a permutation of length n ; given that this factor is polynomial it cannot affect the (exponential) asymptotics of the class.) \square

We also note the following immediate consequence of Vatter [14, Proposition 2.1]:

Lemma 2.9. *Let \mathcal{C}° be a proper centred permutation class and let $\mathcal{C}^{\circ+n}$ denote the class consisting of centred permutations in \mathcal{C}° with at most n points added anywhere. Then $\overline{\text{gr}}(\mathcal{C}^{\circ+n})$ exists and is equal to $\overline{\text{gr}}(\mathcal{C}^\circ)$. Similarly, $\underline{\text{gr}}(\mathcal{C}^{\circ+n}) = \underline{\text{gr}}(\mathcal{C}^\circ)$.*

We shall also require a centred analogue of the notion of an interval of a permutation:

Definition 2.10 (\circ -intervals of a Centred Permutation). A **centred-interval** (also written \circ -**interval**) of a centred permutation π° is an interval containing the origin. We call a centred interval non-trivial if it contains at least one point in addition to the origin. A non-trivial centred interval is **minimal** if it contains no smaller non-trivial centred interval.

Lemma 2.11 (Minimal non-trivial \circ -intervals in a centred permutation). *Let π° be a centred permutation. Then π° has either:*

- one minimal non-trivial \circ -interval \mathcal{J} ; or
- two minimal non-trivial \circ -intervals $\mathcal{J}_1, \mathcal{J}_2$. In this case, each \mathcal{J}_i will contain points in one quadrant only, and from opposite quadrants to each other.

Proof. Suppose that π° has two distinct minimal non-trivial \circ -intervals $\mathcal{J}_1, \mathcal{J}_2$. Then:

- as the intersection of two \circ -intervals is a \circ -interval, $\mathcal{J}_1 \cap \mathcal{J}_2$ is a \circ -interval.
- Hence $\mathcal{J} = \mathcal{J}_1 \cap \mathcal{J}_2$ is a \circ -interval contained in both \mathcal{J}_1 and \mathcal{J}_2 ; as these were assumed to be distinct we can deduce that \mathcal{J} is *strictly* contained in at least one of these. Hence, by the assumed minimality of \mathcal{J}_1 and \mathcal{J}_2 , $\mathcal{J} = \mathcal{J}_1 \cap \mathcal{J}_2 = \{\circ\}$
- Hence $\mathcal{J}_1, \mathcal{J}_2$ are non-trivial \circ -intervals with trivial overlap. Suppose that $\mathcal{J}_1, \mathcal{J}_2$ both have at least one point in the upper half-plane: say $p_1 \in \mathcal{J}_1, p_2 \in \mathcal{J}_2$. Note that $p_1 \notin \mathcal{J}_2, p_2 \notin \mathcal{J}_1$ (by $\mathcal{J}_1 \cap \mathcal{J}_2 = \{\circ\}$) and that one of p_1, p_2 must be lower than the other - without loss of generality, say that p_1 is lower than p_2 . But then p_1 is vertically between the origin and p_2 and hence slices the rectangle \mathcal{J}_2 , contradicting the assumption that \mathcal{J}_2 is a \circ -interval. Hence our supposition is impossible, and $\mathcal{J}_1, \mathcal{J}_2$ cannot both occupy the upper half-plane. See Fig. 6 for an illustration of this argument.
- By precisely the same logic $\mathcal{J}_1, \mathcal{J}_2$ cannot both occupy any half-plane, and thus the only possibility remaining is that \mathcal{J}_1 and \mathcal{J}_2 are both one-quadrant intervals occupying opposite quadrants, as required. Note that this argument immediately precludes the existence of a third minimal non-trivial \circ -interval \mathcal{J}_3 as this would have to be a one-quadrant interval opposite to both \mathcal{J}_1 and \mathcal{J}_2 .

□




Figure 6: The ‘interval’ π_2° is intersected by the point p_1 - and hence is not an interval at all.

2.2 The \boxplus -decomposition

The following operation on centred permutations will be fundamental in describing the structure of the permutation classes we work with in the next section:

Definition 2.12 (The Box Sum). Given two centred permutations π° and σ° their \boxplus -**sum** (read: **box-sum**), written $\pi^\circ \boxplus \sigma^\circ$, is obtained by inflating the origin of σ° with a copy of π° - see Fig. 7 for an illustration.




Figure 7: The \boxplus -sum of the centred permutations $\pi^\circ = 24135$ and $\sigma^\circ = 41352$ is 413685792 ; the origin of σ° is simply replaced ('inflated') by a copy of π° .

Comment 2.13. As the visual representation makes clear, the \boxplus -sum is associative. On the other hand, the \boxplus -sum is certainly *not* commutative, as discussed in more detail below.

Comment 2.14. The \boxplus -sum can be thought of as a generalisations of the direct and skew sums of (uncentred) permutations, two of the most fundamental operations in the study of permutation patterns - see Bevan [4] for the definitions of these operations. When π° and σ° are centred permutations whose points are in the first quadrant only, $\pi^\circ \boxplus \sigma^\circ$ is equivalent to the usual direct sum $\pi \oplus \sigma$ of the corresponding uncentred permutations. If, on the other hand, π° and σ° are centred permutations whose points are in the *second* quadrant only, then $\pi^\circ \boxplus \sigma^\circ$ is the usual skew sum $\sigma \ominus \pi$ of the uncentred permutations (note the reversed order).

Fig. 7 suggests a close relationship between the \boxplus -sum and \circ -intervals. Note that if \mathcal{J} is a \circ -interval of a centred permutation π° then \mathcal{J} contains the origin, and so the set of points contained in \mathcal{J} forms a centred permutation σ° . We say that \mathcal{J} **encloses** the centred permutation σ° , and we immediately observe the following:

Observation 2.15 (The \boxplus -sum and centred intervals). *Let π°, σ° be (centred) permutations, with σ° having strictly smaller length than π° . Then:*

1. $\pi^\circ = \sigma^\circ \boxplus \tau^\circ$ for some centred permutation τ° if and only if π° contains a (necessarily unique) \circ -interval \mathcal{J}_σ enclosing σ° .
2. If $\pi^\circ = \sigma^\circ \boxplus \tau^\circ$ then τ° is the permutation obtained from π° by deleting all non-origin points contained in the (unique) \circ -interval \mathcal{J}_σ .

We note the following immediate consequence of this result:

Corollary 2.16 (Left-cancellation). *Suppose $\sigma^\circ, \tau_1^\circ, \tau_2^\circ$ are centred permutations such that*

$$\sigma^\circ \boxplus \tau_1^\circ = \sigma^\circ \boxplus \tau_2^\circ.$$

Then $\tau_1^\circ = \tau_2^\circ$.

Proof. Let $\pi^\circ = \sigma^\circ \boxplus \tau_1^\circ = \sigma^\circ \boxplus \tau_2^\circ$. Then π° contains a unique \circ -interval \mathcal{J}_σ enclosing σ° by Observation 2.15. Deleting all (non-origin) points in \mathcal{J}_σ from π° will result in τ_1° , but by the same token will also result in τ_2° ; hence $\tau_1^\circ = \tau_2^\circ$. \square

Right-cancellation is also easy to prove, but will not be required here.

Definition 2.17 (\boxplus -decomposables and -indecomposables). We say that a centred permutation σ° is \boxplus -**decomposable** if it can be written as $\pi_1^\circ \boxplus \pi_2^\circ$ for two smaller centred permutations π_1° and π_2° . Conversely, σ° is \boxplus -**indecomposable** if it cannot be split up like this.

It will be important to note for later that σ° is \boxplus -decomposable if and only if there is a proper, non-trivial \circ -interval containing the origin; this is an immediate consequence of Observation 2.15. For example, the inner rectangle in Fig. 7 is an interval because there are no points in the ‘shadow’ of the rectangle in either the x - or y -direction.

Any \boxplus -decomposable centred permutation can be decomposed as a \boxplus -sum of \boxplus -indecomposables. This decomposition is not necessarily unique, as certain elements commute, as in Fig. 8.




Figure 8: The two \boxplus -indecomposable centred permutations $\underline{132}$ and $\underline{12}$ commute under the \boxplus -sum, due to being one-quadrant permutations from opposite quadrants.

More generally, we have:

Lemma 2.18 (Commuting elements under the \boxplus -sum).

Suppose π° and σ° are \boxplus -indecomposable centred permutations. Then

$$\pi^\circ \boxplus \sigma^\circ = \sigma^\circ \boxplus \pi^\circ$$

if and only if either $\pi^\circ = \sigma^\circ$ or π° and σ° are one-quadrant permutations from opposite quadrants.

Proof. One direction here is obvious: if either $\pi^\circ = \sigma^\circ$ or π° and σ° are one-quadrant permutations from opposite quadrants then clearly π° and σ° commute under the \boxplus -sum. For the converse, we assume that the centred permutations π° and σ° commute and aim to deduce that they must be of one of these two forms:

Consider the centred permutation $\tau^\circ = \pi^\circ \boxplus \sigma^\circ = \sigma^\circ \boxplus \pi^\circ$. This can be generated in two ways: either by inflating the origin of σ° by π° or vice versa. Thus there is a \circ -interval \mathcal{J}_1 of τ° enclosing the permutation π° and another \circ -interval \mathcal{J}_2 of τ° enclosing the permutation σ° . As π°, σ° are both \boxplus -indecomposable, \mathcal{J}_1 and \mathcal{J}_2 are both minimal non-trivial \circ -intervals. Hence, by Lemma 2.11, either $\mathcal{J}_1 = \mathcal{J}_2$, in which case $\pi^\circ = \sigma^\circ$, or \mathcal{J}_1 and \mathcal{J}_2 each contain points in only one quadrant, and from opposite quadrants to each other, in which case π°, σ° are one-quadrant permutations from opposite quadrants, as required. \square

We shall henceforth call a pair $\{\pi^\circ, \sigma^\circ\}$ of one-quadrant centred permutations from opposite quadrants a **commuting pair**.

In fact, commutativity of these commuting pairs is the *only* way that uniqueness of the \boxplus -decomposition can fail, as is demonstrated by the following result:

Theorem 2.19. *Let π° be a centred permutation. Then there exist \boxplus -indecomposables $\sigma_1^\circ, \sigma_2^\circ, \dots, \sigma_n^\circ$ such that*

$$\pi^\circ = \sigma_1^\circ \boxplus \sigma_2^\circ \boxplus \dots \boxplus \sigma_n^\circ.$$

Furthermore, this decomposition is unique up to commutativity of adjacent commuting pairs. In other words, if

$$\pi^\circ = \tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_k^\circ.$$

where $\tau_1^\circ, \tau_2^\circ, \dots, \tau_k^\circ$ are \boxplus -indecomposables, then:

1. $k = n$;
2. $[\sigma_1^\circ, \sigma_2^\circ, \dots, \sigma_n^\circ] = [\tau_1^\circ, \tau_2^\circ, \dots, \tau_k^\circ]$ as multisets;
3. the n -tuple $(\sigma_1^\circ, \sigma_2^\circ, \dots, \sigma_n^\circ)$ can be transformed into $(\tau_1^\circ, \tau_2^\circ, \dots, \tau_k^\circ)$ by a sequence of transpositions of adjacent commuting pairs.

Proof. On noting that centred permutations are equivalent to 2-by-2-gridded permutations, this follows from Bevan, Brignall and Ruškuc [3, Lemma 3.4]. We outline a proof here:

Existence is clear: just take minimal non-trivial \circ -intervals inductively. For uniqueness the key is to prove that if $\sigma_1^\circ \neq \tau_1^\circ$ then there is some $l \in \{2, 3, \dots, k\}$ such that $\sigma_1^\circ = \tau_1^\circ$ and $\sigma_1^\circ = \tau_1^\circ$ commutes with τ_j° for $j = 1, 2, \dots, l-1$. All three claims follow inductively from this.

To prove this, note that, as

$$\pi^\circ = \tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_k^\circ$$

there is, by Lemma 2.15 an increasing sequence of \circ -intervals $\mathcal{J}_1, \mathcal{J}_2, \dots, \mathcal{J}_k$ where \mathcal{J}_j encloses the centred permutation $\tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_j^\circ$. Also by Lemma 2.15 there is a unique \circ -interval \mathcal{J} enclosing σ_1° . Note that $\mathcal{J}_1 \cap \mathcal{J} = \{\circ\}$ (as these are distinct and \mathcal{J} is minimal due to \boxplus -indecomposability of σ_1°), and so by Lemma 2.11 these are both one-quadrant intervals from opposite quadrants - wlog say that σ_1° is entirely contained in quadrant 1 and τ_1° is in quadrant 3. Now, as \mathcal{J} is minimal $\mathcal{J}_j \cap \mathcal{J}$ is either $\{\circ\}$ or \mathcal{J} for all j . Further $\mathcal{J}_k \cap \mathcal{J} = \mathcal{J}$. As \mathcal{J}_j is an increasing sequence of intervals, this implies that there is some l such that $\mathcal{J}_l \cap \mathcal{J} = \mathcal{J}$ but $\mathcal{J}_j \cap \mathcal{J} = \{\circ\}$ for all $j \leq l-1$. As $\mathcal{J}_{l-1} \cap \mathcal{J} = \{\circ\}$, Lemma 2.11 implies that $\tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_{l-1}^\circ$ is entirely contained in quadrant 3, so σ_1° commutes with all of τ_j° for $j = 1, 2, \dots, l-1$. And as $\mathcal{J}_l \cap \mathcal{J} = \mathcal{J}$, $\tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_l^\circ$ contains a \circ -interval enclosing σ_1° .

But $\tau_1^\circ \boxplus \tau_2^\circ \boxplus \dots \boxplus \tau_l^\circ$ can be thought of as ' τ_1° plus some points in the third quadrant', whereas σ_1° is entirely contained in the first quadrant. Hence the interval enclosing σ_1° is in fact entirely contained in τ_1° , hence $\sigma_1^\circ \leq \tau_1^\circ$. But by \boxplus -indecomposability of τ_1° we conclude that $\sigma_1^\circ = \tau_1^\circ$. \square

2.3 The Generating Function Specification

We say that a centred permutation class \mathcal{C}° is **\boxplus -closed** if, for any $\pi^\circ, \sigma^\circ \in \mathcal{C}^\circ$, $\pi^\circ \boxplus \sigma^\circ \in \mathcal{C}^\circ$. For any centred permutation class \mathcal{C}° we define its **\boxplus -closure**, denoted $\boxplus\mathcal{C}^\circ$, to be the smallest \boxplus -closed class containing \mathcal{C}° . Equivalently, $\boxplus\mathcal{C}^\circ$ is the set of all centred permutations of the form

$$\sigma_1^\circ \boxplus \sigma_2^\circ \boxplus \cdots \boxplus \sigma_n^\circ$$

where all σ_i° are \boxplus -indecomposables in \mathcal{C}° . Of course, a centred permutation class is \boxplus -closed if and only if it is equal to its own \boxplus -closure. We aim in this section to describe a general method for determining the generating function of a \boxplus -closed centred permutation class from the enumeration sequence of its \boxplus -indecomposables.

Suppose we have a \boxplus -closed centred permutation class \mathcal{C}° . Then every element of \mathcal{C}° is of the form

$$\pi^\circ = \sigma_1^\circ \boxplus \sigma_2^\circ \boxplus \cdots \boxplus \sigma_n^\circ$$

where each σ_i° is a \boxplus -indecomposable in \mathcal{C}° . Suppose we have the generating function $g(z)$ of the (non-empty) \boxplus -indecomposables in \mathcal{C}° (crucially, for the class of permutation classes that will be the main focus of this paper, this generating function will be relatively easy to find). *If this decomposition were unique* we could then immediately obtain the generating function of the entire class by applying the **sequent operator**:

$$\begin{aligned} f(z) = \text{Seq}(f(z)) &= 1 + g(z) + g(z)^2 + g(z)^3 + \dots \\ &= \frac{1}{1 - g(z)} \end{aligned}$$

Of course, as was demonstrated in the previous section, this decomposition is *not* in general unique: the issue is that pairs of \boxplus -indecomposables from opposite quadrants commute, and so elements of \mathcal{C}° cannot be uniquely identified with n -tuples of \boxplus -indecomposables from the class. As it happens, however, we can easily amend the sequent operator to provide a specification of the generating function of \mathcal{C}° in terms of $g(z)$, though, unsurprisingly, we shall also have to use the generating functions $g_i(z)$ of the \boxplus -indecomposables of \mathcal{C}° contained entirely in the i th quadrant, for each $i \in \{1, 2, 3, 4\}$, as these are the elements that introduce commutativity into the \boxplus -decomposition:

Theorem 2.20 (Generating Function Specification for a \boxplus -closed Class). *Let \mathcal{C}° be a centred permutation class, with \boxplus -closure $\boxplus\mathcal{C}^\circ$. Let*

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z)$$

where $g(z)$ is the generating function of the (non-empty) \boxplus -indecomposables of \mathcal{C}° ; and $g_i(z)$ is the generating function of the (non-empty) one-quadrant \boxplus -indecomposables of \mathcal{C}° contained entirely in the i th quadrant. Then

$$f(z) = \text{Seq}(G(z)) = \frac{1}{1 - G(z)}$$

is the generating function of $\boxplus\mathcal{C}^\circ$.

If \mathcal{C}° is itself \boxplus -closed then $f(z)$ is in fact the generating function of \mathcal{C}° .

Proof. This follows almost immediately from much more general result of Cartier and Foata [10] on trace monoids; see also [11, Note V.10]. \square

Comment 2.21. We call the power series

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z)$$

the **amended G-sequence** of \mathcal{C}° . The generating function of $\boxplus\mathcal{C}^\circ$ is then given by the sequent operator not of $g(z)$ but of $G(z)$. We note something curious however: our usual intuition for the sequent operator is that it takes as its input a sequence of positive integers describing the number of distinguishable types of indecomposable ‘block’ of each length n in some collection; the sequent operator then gives us the enumeration sequence of the set of compound structures we can form by gluing any number of these blocks together. But this intuition breaks down when allowing amended G-sequences as inputs for the sequent operator as $G(z)$ can have negative coefficients. Of course, if $G(z)$ is the amended G-sequence of a centred permutation class then $\text{Seq}(G(z)) = (1 - G(z))^{-1}$ has non-negative coefficients by dint of the fact that this is the generating function of a combinatorial class; we know of no ‘internal’ condition for deciding whether a general sequence of integers is mapped to a sequence of non-negative integers by the sequent operator. There are in fact \boxplus -closed centred permutation classes whose amended G-sequences become unbounded and negative and yet (of course) their sequent operators give entirely positive sequences.

2.4 The Exponential Growth Theorem

We now have a method for determining the generating function of a \boxplus -closed centred permutation class. This may seem initially to be of limited interest, as we are mostly interested in the underlying (uncentred) classes, and these will of course have different generating functions to their centred counterparts. We recall, however, that if \mathcal{C}° is a centred permutation class with underlying class \mathcal{C} then \mathcal{C}° and \mathcal{C} have the same (upper) growth rate by Proposition 2.8, and by the following consequence of Pringsheim’s Theorem we can calculate this from the generating function of \mathcal{C}° :

Theorem 2.22 (Exponential Growth Theorem, Flajolet and Sedgewick [11]). *Suppose that $f(z)$ is the generating function of a proper centred permutation class \mathcal{C}° . Then the upper growth rate of \mathcal{C}° is equal to the reciprocal of the radius of convergence of $f(z)$.*

Proof. This is an immediate consequence of [11, Theorem IV.7], on noting that $f(z)$ certainly has non-negative coefficients. \square

Suppose now that \mathcal{C}° is a proper centred permutation class with amended G-sequence $G(z)$. Then the generating function of $\boxplus(\mathcal{C}^\circ)$ is given by

$$f(z) = \frac{1}{1 - G(z)}$$

By Theorem 2.22 the growth rate of $\boxplus\mathcal{C}^\circ$ is then equal to the reciprocal of the radius of convergence of $f(z)$. At first glance it looks like this radius of convergence should be equal to the

modulus of the smallest solution of $G(z) = 1$ (and we will see that this is true for the classes that we are interested in in this paper) but we need to be careful about $G(z)$ itself possibly having singularities before any solutions of $G(z) = 1$. For the time-being we will deal with this issue on a case-by-case basis.

2.5 Examples

Theorems 2.20 and 2.22 are remarkably useful in tandem, suggesting as they do a powerful method for constructing a large number of permutation classes whose growth rates we can easily calculate. For example, the \boxplus -closure of a given finite set of \boxplus -indecomposables is now easy to calculate, as demonstrated by the following examples:

Example 2.23 (\boxplus -closure of 41352). $\pi^\circ = 41352$ is a \boxplus -indecomposable centred permutation. We shall determine the growth rate of its \boxplus -closure $\mathcal{C}^\circ = \boxplus(\pi^\circ)$, that is, the smallest \boxplus -closed class containing π° . Equivalently, this is the downward closure of the set of centred permutations of the form $\pi^\circ \boxplus \pi^\circ \boxplus \dots \boxplus \pi^\circ$. As \mathcal{C}° is (by definition) \boxplus -closed we need only enumerate the \boxplus -indecomposables in \mathcal{C}° in each quadrant.

First, the only \boxplus -indecomposables contained in π° (and hence in \mathcal{C}°) are π° itself and the four single point centred permutations. Hence the generating function of the \boxplus -indecomposables in \mathcal{C}° is $g(z) = 4z + z^4$. Of course, as the only single-quadrant \boxplus -indecomposables in this list are the four single point permutations, the generating functions $g_i(z)$ of the single-quadrant \boxplus -indecomposables in \mathcal{C}° are given by $g_1(z) = g_2(z) = g_3(z) = g_4(z) = z$. Hence the generating function of \mathcal{C}° is given by:

$$\begin{aligned} f(z) &= \frac{1}{1 - [(4z + z^4) - 2z^2]} \\ &= \frac{1}{1 - 4z + 2z^2 - z^4} \end{aligned}$$

And so by Theorem 2.22 we can deduce that the growth rate of $\text{gr}(\mathcal{C}^\circ)$ is the reciprocal of the modulus of the smallest root of the denominator, which is ≈ 3.44372 .

Example 2.24 (The \mathcal{X} -class). Next, we turn to a class introduced by Waton [15] in the context of picture classes: the \mathcal{X} -class, consisting of all permutations which can be drawn on an X . Waton showed that this class has generating function

$$\frac{z(1 - 2z)}{1 - 4z + 2z^2}$$

and hence growth rate $2 + \sqrt{2} \approx 3.41421$

We shall show that this growth rate (but not the generating function) can be derived quickly by instead considering the *centred* permutation class \mathcal{X}° , defined to be the \boxplus -closure of the four single point centred permutations: $\mathcal{X}^\circ = \boxplus\{\cdot|\cdot, \cdot|\cdot, \cdot|\cdot, \cdot|\cdot\}$. Clearly \mathcal{X}° has \mathcal{X} as its underlying (uncentred) class: we can think of this as simply adding an origin point at the centre of the

X diagram used to define \mathcal{X} . But then for \mathcal{X}° we clearly have $g(z) = 4z$ and $g_1(z) = g_2(z) = g_3(z) = g_4(z) = z$; hence \mathcal{X}° has generating function:

$$\frac{1}{1 - 4z + 2z^2}$$

Note that this has the same denominator as the uncentred class \mathcal{X} , demonstrating, as expected, that these two classes have the same growth rate.

3 Pin Sequences and Pin Classes

3.1 Pin Sequences

We begin with a definition, following Brignall, Ruškuc and Vatter [9]:

Definition 3.1. A *pin sequence* is a word (finite or infinite) over the language

$$\mathcal{L}_p = \{1, 2, 3, 4\}(\{l, r\}\{u, d\})^* \cup \{1, 2, 3, 4\}(\{u, d\}\{l, r\})^*$$

We tend to refer to a finite pin sequence as a *pin word* and reserve the term ‘pin sequence’ for the infinite case. For example, $w = 3lrdrdlurdl$ is a pin word and $4(urul)^*$ is a pin sequence (we use the notation $(f)^*$ to denote the factor f recurring indefinitely, so $4(urul)^* = 4urulurulurulurul\dots$).

We think of the letters u, d, l, r as representing up, down, left, right, respectively. We then say that letters u and d have **vertical alignment** and letters l and r have **horizontal alignment**. We can then informally describe a pin sequence as consisting of an initial numeral from $\{1, 2, 3, 4\}$ followed by a sequence of letters from $\{u, l, d, r\}$ which must alternate between horizontal and vertical alignment. Note that this implies that there are 4 pin words of length 1 and 2^{n+2} pin words of every length $n \geq 2$.

We shall also have cause to refer to the language

$$\mathcal{L}_p^* = (\{l, r\}\{u, d\})^* \cup (\{u, d\}\{l, r\})^*$$

Note that a pin word consists of an initial numeral between 1 and 4 followed by a (possibly empty) word over \mathcal{L}_p^* .

A finite pin word w of length n can be converted into a centred permutation π_w° of length n by the following procedure (see Fig. 9 for an illustration of this process):

Definition 3.2 (The π -map and pin permutations). Let $n \in \mathbb{N}$; furthermore, let Π_n denote the set of pin words of length n , and S_n° denote the set of centred permutations of length n . The π -map is the map

$$\pi^\circ : \Pi_n \rightarrow S_n^\circ$$

defined by the following procedure: given $w \in \Pi_n$:

1. Place the initial origin point p_0 ; use this point to split the plane into four numbered quadrants, as in Fig. 2.
2. Place the first point p_1 in the quadrant specified by the initial numeral of the pin word w .
3. Once the first $k - 1$ points have been placed, place the point p_k either up, down, left or right (depending on the letter $u, d, l, \text{ or } r$ appearing in the k -th place in the pin sequence) of the bounding rectangle of all previous points $\{p_0, p_1, \dots, p_{k-1}\}$ at the end of a 'pin' which separates the last point p_{k-1} from all previous points.
4. Once all n points have been placed, read off the centred permutation π_w° , given by the points $\{p_0, p_1, \dots, p_n\}$, with p_0 as the origin.

We refer to π_w° as the (centred) pin permutation associated with w ; similarly, we write π_w for the underlying (uncentred) permutation of π_w° and refer to this as the (uncentred) pin permutation associated with w .

We refer to a centred permutation σ° as a (centred) **pin permutation** if there is some pin word w such that $\sigma^\circ \leq \pi_w$. Similarly, an uncentred permutation σ is an (uncentred) pin permutation if it is the underlying permutation of a centred pin permutation.

Finally, we refer to a centred permutation σ° as a **contiguous pin permutation** if there is some pin word w such that $\sigma^\circ = \pi_w$.




Figure 9: The (centred) pin permutation 31586427 (and its underlying uncentred permutation 3147526), constructed from the pin sequence 2lurdld.

Observation 3.3. *By definition the point p_k slices the bounding rectangle $\text{rec}(p_0, p_1, \dots, p_{k-1})$ of all previous points, including the origin. In fact, the definition implies that p_k is the **only** point after p_{k-1} that slices this rectangle; this means that if we remove p_k from the corresponding centred permutation $\text{rec}(p_0, p_1, \dots, p_{k-1})$ becomes a \circ -interval, and hence the resulting centred permutation is \boxplus -decomposable. This will be important in our later structure theorem for a pin class.*

We denote the class consisting of *all* centred pin permutations \mathcal{P}° , and its uncentred counterpart by \mathcal{P} . We call this the **complete pin class**. Bassino, Bouvel and Rossin [2] proved that \mathcal{P}

has rational generating function and growth rate $\omega_\infty \approx 5.24112$. We shall later find the generating function of \mathcal{P}° , which of course has the same growth rate as \mathcal{P} by Proposition 2.8. We are primarily interested in this paper in certain subclasses of \mathcal{P} , known as **pin classes**. These are defined somewhat analogously to pin permutations: just as we can convert a (finite) pin word into a permutation, we can also convert an (infinite) pin sequence into a pin class:

Definition 3.4 (Pin Classes). Suppose that w is an infinite pin sequence. Let π_n° be the pin permutation obtained by the above process from the initial contiguous subsequence of w of length n (which is itself a finite pin word), and consider the set $\Pi = \{\pi_1^\circ, \pi_2^\circ, \pi_3^\circ, \dots\}$. The downward closure of Π under the pattern containment order \leq forms a centred permutation class \mathcal{C}_w° , known as the centred *pin class* of w . We similarly define the uncentred pin class of w , \mathcal{C}_w , to be the underlying class of \mathcal{C}_w° .

Informally, we can think of this as using the infinite pin sequence w to draw an 'infinite diagram' by the same process as in the finite case; the pin class \mathcal{C}_w is then the class of all permutations that can be found somewhere within this infinite diagram (by picking a finite collection of points and 'forgetting' everything else).

Example 3.5. Taking $w = 1(\text{ldrdluru})^*$ we generate the pin diagram shown in Fig. 10. We could of course extend this diagram indefinitely. The class \mathcal{C}_w then consists of all permutations that can be found somewhere in this diagram (with \mathcal{C}_w° being the class of all *centred* permutations that can be found).




Figure 10: The pin class \mathcal{C}_w° generated by the pin sequence $w = 1(\text{ldrdluru})^*$.

3.1.1 Oscillations

One particular collection of pin permutations is already well-known and will play a key role in our later theory:

Definition 3.6 (Oscillations). Let w be a (finite) pin word that stays in its initial quadrant (for example, $1ururu$, $1rurur$, $3ldl$ and $4drdrd$). We call the permutation π_w° generated by w an **oscillation** (sometimes a **one-quadrant oscillation** for emphasis).

Oscillations have natural ‘staircase’ structures - see Fig. 11 for examples. Note that there are precisely two oscillations of each length $n \geq 3$ in each quadrant (eg. those generated by $1ururu$ and $1rurur$ of length 6 in the first quadrant), but only one of length 2 (as $1u$ and $1r$ in fact generate the same (centred) permutation $\begin{array}{c} \bullet \\ | \\ \bullet \end{array}$ - see Fig. 12 for an illustration).




Figure 11: Three oscillations and the pin words that generate them. Note the two distinct oscillations of length 6 in the first quadrant. In general, there are two distinct oscillations of each length $n \geq 2$ in each quadrant.




Figure 12: Whilst there are two distinct oscillations in each quadrant for all lengths $n \geq 3$, there is only one at length 2. This is because the pin words $1u$ and $1r$ in fact generate the *same* centred permutation $\underline{132}$. This is an example of a **collision** of pin factors, a phenomenon that we will study in detail later.

We can also consider the pin class \mathcal{O}° of **increasing oscillations**, defined to be \mathcal{C}_w° for the pin word $1(ru)^*$ (see Fig. 13). This class has been shown to have growth rate $\kappa \approx 2.20557$, a fact which we shall be able to prove later. We will also be able to prove that κ is in fact the *smallest* possible growth rate of a pin class, and is only achievable by pin classes which are ‘essentially’ \mathcal{O}° .

Oscillations have been studied extensively in the literature. In fact, to a significant extent pin permutations are worth studying because they generalise the oscillations and end up being interesting for much the same reasons. For example, Bevan [5] demonstrated that an infinite antichain contained in the two-point extension \mathcal{O}^{+2} can be used to construct permutation classes (all of which contain long oscillations) at *every* growth rate $\lambda \geq 2.35526$. This construction in fact generalises to pin classes in general, allowing us to construct genuinely distinct antichains at every real number which is the growth rate of a pin class.




Figure 13: The class \mathcal{O}° of **increasing oscillations**, generated by the pin sequence $1(\text{ru})^*$.

We thus have established a method for converting a pin sequence into a (centred or uncentred) permutation class. This allows us to describe an extraordinarily large number of classes which would be awkward to describe by other means (it would, for example, take a lot of work to describe the class in Fig. 10 in terms of its basis). There are two main reasons that this construction will prove useful. First, controlling the initial pin sequence allows us to control various properties of the permutation class produced (for example, the aforementioned connection with simple permutations allows us to control the number of simples in a given pin class). Second, and most importantly, we have a process for determining the growth rates of the pin permutation classes produced: this will depend on an in-built structure possessed by pin permutation classes, the \boxplus -**decomposition**.

3.2 Pin Factors

We will need to have a notion of a ‘subsequence’ of a pin sequence, one that should hopefully preserve pattern containment in both the centred and uncentred case:

Definition 3.7 (Pin Factor). Suppose that w is a (finite or infinite) pin word. A **pin factor** of w is either:

- an initial contiguous subsequence of w ; or
- a non-initial contiguous subsequence of w in which the first letter is replaced by the quadrant in which the corresponding point appears in π_w° .

Note that a pin factor of a given pin word is by definition itself a pin word. We write $w_{i,j}$ for the pin factor obtained by taking the contiguous subsequence of w between the i th and j th place.

Example 3.8. Consider the pin word $w = 2\text{ru}d\text{lur}d\text{ru}$. Then $w_{1,2} = 2\text{r}$, $w_{1,4} = 2\text{ru}l$, $w_{1,9} = 2\text{ru}d\text{lur}d\text{ru}$ and w itself are all pin factors of w as they are all (finite) initial subsequences. If instead we take out the contiguous subsequence from the fifth to ninth terms we obtain the word $d\text{lur}d$. This is a factor in the word sense, but it is not a pin factor as it does not begin with a numeral (and so is not a pin word). However, if we draw out the (centred) permutation π_w° generated by w (see Fig. 14) we can see that the fifth point placed is in the 3rd quadrant, so we replace the initial d in $d\text{lur}d$ with a 3 to obtain the pin word $3\text{lur}d$, which is now a pin factor of w ; specifically $w_{5,9} = 3\text{lur}d$. For an internal pin word we can always work out what this

initial numeral should be by looking at the previous letter: for example, if we wish to work out $w_{9,11}$ we first take out the subword dru (between the 9th and 11th places); to work out which numeral we replace the initial d with, note that it was preceded by an r and the d in any rd clearly must be placed in the 4th quadrant. Hence $w_{9,11} = 4ru$. This suggests a correspondence between pin factors of length n and \mathcal{L}^* -subwords of length $n + 1$, stated below in Lemma 3.11.




Figure 14: The construction of the (centred) permutation π_w° from the pin word $w = 2ruldurdr$. Note that the fifth-placed point, p_5 , corresponding to the first d in the word, is in the 3rd quadrant, so when we extract the word factor $durd$ from w we replace the initial d with a 3 to obtain the pin factor $\tilde{w} = 3lurd$. Note that the (centred) permutation $\sigma_{\tilde{w}}^\circ$ (highlighted in red) is a subpermutation of π_w° .

As Fig. 14 suggests, this notion preserves pattern containment:

Observation 3.9. *Suppose w_1, w_2 are (finite) pin words and that w_1 is a pin factor of w_2 . Then:*

$$\pi_{w_1}^\circ \leq \pi_{w_2}^\circ$$

It's also worth noting that, while pin factors of w of length n cannot in general be identified with subwords of w of length n (for example, the subword $ulur$ could instantiate either $1lur$ or $2lur$ depending on the context) they can *almost* be identified with subwords of length $n + 1$ (with the first letter determining the quadrant numeral): the only issue being initial pin factors, which may not reappear. In practice this will be only a minor inconvenience (which may be completely ignored in the recurrent case) as initial pin factors cannot affect the growth rate of a pin class, at worst only affecting the enumeration sequence.

We now have the terminology to define an important category of pin sequences, whose associated pin classes (we shall see) have a structure which will be particularly amenable to the enumeration methods developed in Section 1:

Definition 3.10 (Recurrent Pin Sequences). We refer to an infinite pin sequence w as **recurrent** if every pin factor that occurs in w occurs infinitely often; similarly, w is **eventually recurrent** if every pin factor that occurs after a certain point occurs infinitely often. We will often refer to the pin class \mathcal{C}_w° generated by a recurrent pin sequence as a **recurrent pin class**.

Note that this definition is subtly different from recurrence in the usual (subword) sense, due to complications introduced by initial subsequences (which begin with a numeral): these initial

subwords can never occur as subwords again in w by the definition of a pin sequence, but (in order for w to be recurrent) they *must* appear infinitely often a pin factors. This means it can often be difficult to tell whether a pin sequence is recurrent or not, even in the periodic case, until it is drawn out: for example, $2(\text{ul})^*$ is recurrent, but $1(\text{ul})^*$ is not, as $w_{1,2} = 1\text{u}$ never reoccurs as a pin factor.

We wish to ignore, as far as possible, this subtle and often irritating distinction between subwords and pin factors. First, we call a pin factor of w *fully internal* if it occurs in w starting after the second place; that is, fully internal pin factors are those of the form $w_{i,j}$ where $3 \leq i < j$. Note that fully internal pin factors can be found somewhere in w with a preceding letter (as opposed to numeral). This letter is enough to reconstruct the initial numeral of the pin factor, thus leading to the following:

Lemma 3.11. *Let w be a pin sequence. Then the number of fully internal pin factors of w of length $n \geq 2$ is equal to the number of \mathcal{L}^* -subwords of w of length $n + 1$.*

Of course, the fully internal restriction can be removed if we further assume that w is recurrent, as in this case any pin factor beginning in the first or second place can also be found later on; hence all pin factors are fully internal, giving us the following simplification:

Corollary 3.12. *Let w be a recurrent pin sequence. Then the number of pin factors of w of length n is equal to the number of \mathcal{L}^* -subwords of w of length $n + 1$.*

3.2.1 Left-truncations

We shall also require an infinite analogue of a pin factor of w , the (infinite) pin *sequence* obtained by starting w at a later point:

Definition 3.13 (Left-truncation of a pin sequence). Let w be a pin sequence and $n \in \mathbb{N}$. The n th **left-truncation** of w , $w_{\geq n}$, is the pin sequence obtained from w by replacing the symbol in the n th place with the number of the quadrant in which p_n is placed, and then removing all of the previous symbols.

Informally, $w_{\geq n}$ is ‘ w but started in the n th place’. For example, if $w = 3\text{rurdlurur}$ then $w_{\geq 4} = 1\text{dlurur}$ and $w_{\geq 7} = 2\text{rur}$. Again, we can always work out the numeral to replace the letter in the n th place with by looking at the previous symbol.

Crucially, left-truncating a pin sequence does not affect the asymptotics of its pin class:

Lemma 3.14 (Finite Prefix Lemma). *Let w be a pin sequence and $n \in \mathbb{N}$. Then $\overline{\text{gr}}(\mathcal{C}_w^\circ) = \overline{\text{gr}}(\mathcal{C}_{w_{\geq n}}^\circ)$ and $\underline{\text{gr}}(\mathcal{C}_w^\circ) = \underline{\text{gr}}(\mathcal{C}_{w_{\geq n}}^\circ)$.*

Proof. Recall that we write $\mathcal{C}^{\circ+k}$ for the k -point extension of \mathcal{C}° (that is, the class of centred permutations from \mathcal{C}° with at most k extra points added anywhere). Note that the (infinite) pin diagram generated by w is simply that of $w_{\geq n}$ with $n - 1$ extra points, hence:

$$\mathcal{C}_{w_{\geq n}}^\circ \subseteq \mathcal{C}_w^\circ \subseteq \mathcal{C}_{w_{\geq n}}^{\circ+(n-1)}$$

and by Lemma 2.9 the classes on the left and right have the same (upper and lower) growth rates. \square

We call this the **Finite Prefix Lemma**, as it tells us that any finite prefix of a pin sequence cannot affect the growth rate of the associated pin class, a fact that we shall use often.

3.3 The Pin Decomposition

The key idea we will use to enumerate pin classes comes from the following observation: when we take a centred permutation generated by a pin word and remove any interior point, we decompose the resulting permutation as the \boxplus -sum of two smaller pin permutations. Fig. 15 illustrates this process.




Figure 15: When we remove the point p_6 from the pin permutation generated by $1ldldruruld$ we remove the only point that slices the bounding rectangle of the first five points; we can thus contract this rectangle down to a single point and in doing so express the resulting centred permutation as the box sum of π_{1ldld}° and π_{1ruld}° .

We formalise this as follows:

Lemma 3.15 (Removing an interior point from a pin permutation).

Suppose that w is a pin word of length n , and that p_1, p_2, \dots, p_n are the corresponding points of the centred pin permutation π_w° . Let $k \in \{2, 3, \dots, n-1\}$. Then:

$$\pi_w^\circ - \{p_k\} = \pi_{w_{1,k-1}}^\circ \boxplus \pi_{w_{k+1,n}}^\circ$$

Proof. We note that by Observation 3.3, p_k is the *only* point after p_{k-1} that slices the bounding rectangle of p_0, p_1, \dots, p_{k-1} . Hence when p_k is removed, $\text{rec}(p_0, p_1, \dots, p_{k-1})$ becomes a \circ -interval \mathcal{J} in the resulting permutation, enclosing the permutation $\pi_{w_{1,k-1}}^\circ$. Hence, by Observation 2.15:

$$\pi_w^\circ - \{p_k\} = \pi_{w_{1,k-1}}^\circ \boxplus \tau^\circ$$

for some centred permutation τ° . That τ° is $\pi_{w_{k+1,n}}^\circ$ follows from Observation 3.9, as this corresponds to the remaining points. \square

This process equips pin classes with an in-built structure theorem:

Theorem 3.16 (The Pin Decomposition).

Suppose that w is an infinite pin sequence and that \mathcal{C}_w° is the pin class it generates. Then

$$\sigma^\circ \in \mathcal{C}_w^\circ \text{ iff } \sigma^\circ = \pi_{w_1}^\circ \boxplus \pi_{w_2}^\circ \boxplus \cdots \boxplus \pi_{w_k}^\circ$$

where w_1, w_2, \dots, w_k is a sequence of pin factors of w that occur *in that order, in non-overlapping instances and separated from each other by at least one letter in w .*

Proof. This is almost immediate from the process derived above: $\sigma^\circ \in \mathcal{C}_w^\circ$ must be contained in some centred permutation generated by an initial subsequence w^* of w , so we can obtain σ° by deleting letters from w^* ; but every time we do we split the resulting permutation into the \boxplus -sum of the pin factors directly before and after. Repeatedly doing this gives the required result. \square

We note one immediate consequence of this theorem:

Corollary 3.17 (\boxplus -indecomposables in a pin class).

Let w be a pin sequence and \mathcal{C}_w° the associated pin class. Suppose $\pi^\circ \in \mathcal{C}_w^\circ$ is a \boxplus -indecomposable pin permutation. Then $\pi^\circ = \pi_{\tilde{w}}^\circ$ for some pin factor \tilde{w} of w .

The structure theorem 3.16 is often awkward to apply due to the conditions on the pin factors w_i ; it becomes much easier however, if we assume that w is a *recurrent* pin sequence - that is, every pin factor of w occurs infinitely often. The theorem then becomes:

Theorem 3.18 (The Pin Decomposition - Recurrent Case).

Suppose that w is an recurrent infinite pin sequence and that \mathcal{C}_w° is the pin class it generates. Then:

$$\sigma^\circ \in \mathcal{C}_w^\circ \text{ iff } \sigma^\circ = \pi_{w_1}^\circ \boxplus \pi_{w_2}^\circ \boxplus \cdots \boxplus \pi_{w_k}^\circ$$

where w_1, w_2, \dots, w_k is a sequence of pin factors of w , and $\pi_{w_i}^\circ$ is the (centred) permutation generated from w_i .

This has the following crucial corollary:

Corollary 3.19 (Recurrent pin classes are \boxplus -closed).

Suppose that w is a recurrent pin sequence. Then the pin class \mathcal{C}_w° is \boxplus -closed.

Comment 3.20. We note now that the converse of Corollary 3.19 is true, but the proof involves results derived in the next section, specifically regarding the possibility of certain pin factors generating \boxplus -decomposable permutations (we know that every \boxplus -indecomposable in a pin class is of the form $\pi_{w_{i,j}}^\circ$, but, as we shall see, the converse is certainly not true). As we shall not require this converse we omit the proof.

Corollary 3.19 tells us that to enumerate a *recurrent* pin class \mathcal{C}_w° it will suffice to enumerate its \boxplus -indecomposables and then apply the generating function specification given in Theorem 2.20. It is thus important for us to know how to enumerate the \boxplus -indecomposables in a pin class; Corollary 3.17 suggests a way of doing this that we will study in a later section.

3.4 Recurrent Pin Classes have Growth Rates

We know that pin classes all have *upper* growth rates by Proposition 2.8. In this section we show that *recurrent* pin classes in fact have *proper* growth rates. In fact, as our eventual aim is to show that *all* pin classes have proper growth rates we will have to be slightly more general here and aim to prove that all \boxplus -closed subclasses of \mathcal{P}° (the class of all pin permutations) have proper growth rates.

Our strategy in proving this will be to follow Arratia [1] who proved that \oplus -closed permutation classes have proper growth rates. Arratia's strategy was to note that, in a \oplus -closed class \mathcal{C} , the map

$$(\sigma, \tau) \mapsto \sigma \oplus \tau$$

is an injection from $\mathcal{C}_m \times \mathcal{C}_n$ to \mathcal{C}_{m+n} , and so the enumeration sequence of \mathcal{C} satisfies the **supermultiplicative inequality**:

$$C_{m+n} \geq C_m C_n$$

Finally, Arratia applies the supermultiplicative form of Fekete's Lemma to deduce that $\text{gr}(C_n)$ exists.

We aim to apply this method to \boxplus -closed centred permutation classes, but we have a problem: as we have seen, the \boxplus -decomposition is certainly *not* unique and so the map

$$\begin{aligned} \mathcal{C}_m^\circ \times \mathcal{C}_n^\circ &\rightarrow \mathcal{C}_{m+n}^\circ \\ (\sigma^\circ, \tau^\circ) &\mapsto \sigma^\circ \boxplus \tau^\circ \end{aligned} \tag{1}$$

is not necessarily an injection and we will not, in general, be able to prove the supermultiplicative inequality for a \boxplus -closed subclass of \mathcal{P}° . We note, however, that even the weaker 'supermultiplicative-like' inequality $C_{m+n} \geq k C_m C_n$ (for some constant $k < 1$) would be sufficient to deduce the existence $\text{gr}(C_n)$, on applying Fekete's Lemma to $D_n = k C_n$ instead. Unfortunately, the enumeration sequence of a \boxplus -closed subclass of \mathcal{P}° does not in general satisfy even this weaker form of supermultiplicativity: for example, if we take \mathcal{C}° to be the \boxplus -closure of $\begin{smallmatrix} \cdot \\ | \\ \cdot \end{smallmatrix}$ and $\begin{smallmatrix} \cdot \\ | \\ \cdot \\ | \\ \cdot \end{smallmatrix}$ we can easily show that $\lim_{n \rightarrow \infty} |\mathcal{C}_{2n}^\circ| / |\mathcal{C}_n^\circ|^2 = 0$. The problem in this example is that there is *too much commutativity* due to the class only containing points in the first and third quadrants, which are opposite to each other. As it turns out, this is really the only thing that can go wrong in proving a supermultiplicative-like identity: if we have any points from a pair of adjacent quadrants then we already have enough pairs that don't commute to conclude that $C_{m+n} \geq k C_m C_n$ for some constant k . We thus exclude the bad case in which we have only points from opposite quadrants via the following definition:

Definition 3.21 (Adjacency Condition). Let μ_i° denote the centred permutation consisting of a single point in the i th quadrant (so $\mu_1^\circ = \begin{smallmatrix} \cdot \\ | \\ \cdot \end{smallmatrix}$, etc.). We say that a centred permutation class \mathcal{C}° satisfies the **adjacency condition** if it contains either precisely one of $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ, \mu_4^\circ$ or it contains a pair μ_i°, μ_j° from adjacent quadrants.

Assuming this condition we may now prove a weak form of supermultiplicativity for the enumeration sequence of a \boxplus -closed subclass of \mathcal{P}° :

Proposition 3.22. *Let \mathcal{C}° be a \boxplus -closed subclass of \mathcal{P}° which satisfies the adjacency condition. Let \mathcal{C}_n° denote the set of centred permutations of length n in \mathcal{C}° and write $C_n = |\mathcal{C}_n^\circ|$. Then, for all $m, n \in \mathbb{N}$:*

$$C_{m+n} \geq C_{m-1}C_{n-1}$$

Proof. Note first that certainly $\mu_i \in \mathcal{C}^\circ$ for some i : now, if $\sigma^\circ \in \mathcal{C}_{n-1}^\circ$ then $\mu_i \boxplus \sigma^\circ \in \mathcal{C}_n^\circ$, and by left-cancellation the implied map from \mathcal{C}_{n-1}° to \mathcal{C}_n° is injective. Hence:

$$C_n \geq C_{n-1}$$

We now split into cases based on how many of the one-point permutations $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ, \mu_4^\circ$ are contained in \mathcal{C}° :

- **Case 1:** Suppose \mathcal{C}° contains either one or two of $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ, \mu_4^\circ$. Then, by the Adjacency Condition, \mathcal{C}° is contained entirely within one half-plane. Hence there is no commutativity in \mathcal{C}° and so (by Theorem 2.19) the map

$$\begin{aligned} \mathcal{C}_m^\circ \times \mathcal{C}_n^\circ &\rightarrow \mathcal{C}_{m+n}^\circ \\ (\sigma^\circ, \tau^\circ) &\mapsto \sigma^\circ \boxplus \tau^\circ \end{aligned}$$

is an injection, hence

$$\begin{aligned} C_{m+n} &\geq C_m C_n \\ &\geq C_{m-1} C_{n-1} \end{aligned}$$

as required.

- **Case 2:** Suppose \mathcal{C}° contains precisely three of $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ, \mu_4^\circ$. Without loss of generality, assume that these are $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ$. Note that μ_2° commutes with nothing in \mathcal{C}° . Thus

$$\begin{aligned} \mathcal{C}_m^\circ \times \mathcal{C}_{n-1}^\circ &\rightarrow \mathcal{C}_{m+n}^\circ \\ (\sigma^\circ, \tau^\circ) &\mapsto \sigma^\circ \boxplus \mu_2^\circ \boxplus \tau^\circ \end{aligned}$$

is an injection. Hence:

$$\begin{aligned} C_{m+n} &\geq C_m C_{n-1} \\ &\geq C_{m-1} C_{n-1} \end{aligned}$$

as required.

- **Case 3:** Suppose \mathcal{C}° contains all four of $\mu_1^\circ, \mu_2^\circ, \mu_3^\circ, \mu_4^\circ$. We claim that

$$\begin{aligned} \mathcal{C}_{m-1}^\circ \times \mathcal{C}_{n-1}^\circ &\rightarrow \mathcal{C}_{m+n}^\circ \\ (\sigma^\circ, \tau^\circ) &\mapsto \sigma^\circ \boxplus \mu_2^\circ \boxplus \mu_3^\circ \boxplus \tau^\circ \end{aligned}$$

is an injection (as nothing can commute with both of the interior permutations). Hence:

$$C_{m+n} \geq C_{m-1} C_{n-1}$$

as required.

□

This, as it stands, is slightly weaker than a supermultiplicativity result (which should relate C_{m+n} with $C_m C_n$, not $C_{m-1} C_{n-1}$), but we could easily convert it into such if we could obtain a bound on the ratio between consecutive terms in the enumeration sequence of a \boxplus -closed subclass of \mathcal{P}° . To find such a bound we notice that all pin permutations of length $n + 1$ can be obtained as ‘extensions’ of pin permutations of length n , motivating the following definition:

Definition 3.23 (Pin Representations). Let σ° be a pin permutation. We call a k -tuple of pin words (w_1, w_2, \dots, w_k) a **pin representation** of σ° if

$$\sigma^\circ = \pi_{w_1}^\circ \boxplus \pi_{w_2}^\circ \boxplus \dots \boxplus \pi_{w_k}^\circ$$

where each w_i is a pin word.

Clearly, every pin permutation σ° has a pin representation (as every \boxplus -indecomposable is of the form π_w° ; but note that we do not require that each $\pi_{w_i}^\circ$ is \boxplus -indecomposable). Note, however, that pin representations are in general highly non-unique (it can, for example, easily be seen that $(1, 3, 1ul)$, $(3, 1, 1ul)$ and $(1, 1uld)$ are all representations of the same centred permutation, namely $51\bar{2}364$). We can now formalise the notion of one pin permutation being an extension of another:

Definition 3.24 (One-point Extensions). Let $\sigma^\circ, \tilde{\sigma}^\circ \in \mathcal{P}^\circ$ with $|\tilde{\sigma}^\circ| = |\sigma^\circ| + 1$. We say that $\tilde{\sigma}^\circ$ is a **one-point extension** of σ° if there is some pin representation (w_1, w_2, \dots, w_k) of σ° for which there exists **either**:

- some $L \in \{u, d, l, r\}$ such that the concatenation $w_k L$ is a valid pin word and

$$(w_1, w_2, \dots, w_k L)$$

is a pin representation of $\tilde{\sigma}^\circ$; **or**:

- some $Q \in \{1, 2, 3, 4\}$ such that

$$(w_1, w_2, \dots, w_k, Q)$$

is a pin representation of $\tilde{\sigma}^\circ$.

(Informally, $\tilde{\sigma}^\circ$ is a one-point extension of σ° if there is a pin representation of σ° which can be extended to a pin representation of $\tilde{\sigma}^\circ$ by the appendment of one final symbol, either as an extension of the final pin word in the representation, or as an additional pin word which consists of a single quadrant numeral.) We note the following crucial facts about one-point extensions:

Lemma 3.25. *Suppose \mathcal{C}° is a subclass of the complete pin class \mathcal{P}° . Then:*

1. *Every $\tilde{\sigma}^\circ \in \mathcal{C}^\circ$ of length $n + 1$ is a one-point extension of some $\sigma^\circ \in \mathcal{C}^\circ$ of length n .*
2. *Let $\sigma^\circ \in \mathcal{C}^\circ$. Then σ° has at most 12 one-point extensions in \mathcal{C}° .*

Comment 3.26. Note that 2. is a statement about *permutations* not representations: any given $\sigma^\circ \in \mathcal{C}^\circ$ of length n may have many distinct representations, which may extend to many more than 12 distinct representations of length $n + 1$; the lemma tells us that these representations represent at most 12 distinct permutations.

Proof. 1. Let $\tilde{\sigma}^\circ \in \mathcal{C}^\circ$ of length $n + 1$ and take a pin representation (w_1, w_2, \dots, w_k) of $\tilde{\sigma}^\circ$. Simply by deleting the final symbol of w_k (which may be all of w_k if w_k is simply a quadrant numeral) we obtain a pin representation of a permutation σ° which must also be in \mathcal{C}° (by Observation 3.9 and closure of \mathcal{C}° under the containment order \leq) and which has $\tilde{\sigma}^\circ$ as a one-point extension.

2. Let $\sigma^\circ \in \mathcal{C}^\circ$ and suppose that $\tilde{\sigma}^\circ$ is a one-point extension of σ° (which may or may not be in \mathcal{C}°). We claim that *regardless of the pin representation chosen for σ° , $\tilde{\sigma}^\circ$ must be obtained by adding a point in one of the 12 positions indicated in Fig. 16: if we append a numeral Q to the pin representation then $\tilde{\sigma}^\circ = \sigma^\circ \boxplus \mu_Q$ and we have added a point in one of the extreme corners; if, on the other hand, we append a letter L to the pin representation then we add a point which is the most extreme in the direction indicated by L and the second-most extreme in another (perpendicular) direction, determined by the final symbol of the pin representation of σ° .*

□




Figure 16: If σ° is contained in the square then *any* one-point extension of σ° in \mathcal{C}° is formed by adding a point in one of the 12 positions shown.

Lemma 3.25 immediately implies the following:

Proposition 3.27. *Let \mathcal{C}° be a subclass of the complete pin class \mathcal{P}° . Let \mathcal{C}_n° denote the set of centred permutations of length n in \mathcal{C}° and write $C_n = |\mathcal{C}_n^\circ|$. Then, for all $n \in \mathbb{N}$:*

$$C_n \leq 12C_{n-1}$$

Finally, we combine Propositions 3.22 and 3.27 to deduce a supermultiplicativity-like identity for \boxplus -closed subclasses of \mathcal{P}° satisfying the adjacency condition:

Corollary 3.28. *Let \mathcal{C}° be a \boxplus -closed subclass of \mathcal{P}° which satisfies the adjacency condition and let $C_n = |\mathcal{C}_n^\circ|$. Then, for all $m, n \in \mathbb{N}$:*

$$C_{m+n} \geq \frac{1}{144} C_m C_n$$

This is finally enough to prove our main result:

Theorem 3.29. *Suppose that \mathcal{C}° is a \boxplus -closed subclass of \mathcal{P}° which satisfies the adjacency condition. Then \mathcal{C}° has a proper growth rate.*

Proof. Let C_n be the enumeration sequence of \mathcal{C}° and set $D_n = \frac{1}{144} C_n$. Then, by Corollary 3.28,

$$D_{m+n} \geq D_m D_n,$$

so we may apply Fekete's Lemma to deduce that $\lim_{n \rightarrow \infty} \sqrt[n]{D_n}$ exists and is equal to $\sup_{n \in \mathbb{N}} \sqrt[n]{D_n}$, which is finite by Marcus-Tardos. Hence:

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{C_n} &= \lim_{n \rightarrow \infty} (\sqrt[n]{144}) (\sqrt[n]{D_n}) \\ &= \lim_{n \rightarrow \infty} \sqrt[n]{D_n} \\ &< \infty, \end{aligned}$$

which is to say that $\text{gr}(\mathcal{C}^\circ)$ exists. □

As any recurrent pin class is \boxplus -closed and satisfies the adjacency condition, we immediately obtain:

Corollary 3.30 (Recurrent pin classes have proper growth rates). *Let w be a recurrent pin word. Then the pin class \mathcal{C}_w° has a proper growth rate.*

Of course, Corollary 3.30 (along with Proposition 2.8) immediately implies that *uncentred* pin classes generated by recurrent pin sequences also have growth rates, which will equal that of their centred counterparts: $\text{gr}(\mathcal{C}_w) = \text{gr}(\mathcal{C}_w^\circ)$.

3.5 Classification of Collisions and \boxplus -decomposables

We now know that any recurrent pin class \mathcal{C}_w° has a proper growth rate. In order to find the growth rate of a given recurrent pin class in practice we shall have to enumerate the \boxplus -indecomposables contained in \mathcal{C}_w° : if we can do this then Theorem (2.20) will allow us to write down the generating function of the class, from which we can deduce the growth rate via the Exponential Growth Theorem. Accordingly, in this section we take up the problem of enumerating the \boxplus -indecomposables in a pin class.

We begin by noting that Corollary 3.17 immediately implies the following property of the restriction of the π -map to pin factors of length n :

Proposition 3.31 (Induced π -map contains all \boxplus -indecomposables in its image). *Let w be a pin sequence and \mathcal{C}_w° the associated (centred) pin class. Let P_n denote the set of pin factors of w of length n and \mathcal{B}_n° denote the set of \boxplus -indecomposables in \mathcal{C}_w° of length n . Then the induced map*

$$\pi_n : P_n \rightarrow \mathcal{C}_w^\circ$$

defined by

$$\pi_n : \tilde{w} \mapsto \pi_{\tilde{w}}^\circ$$

contains the set \mathcal{B}_n° in its image.

The study of this induced map will be vital for further progress as it suggests a natural method for counting the \boxplus -indecomposables in a given pin class \mathcal{C}_w° : if (for some n) we had the following two properties:

- the image $\pi_n(P_n)$ is equal to \mathcal{B}_n° (that is, $\pi_n(\tilde{w})$ is \boxplus -indecomposable for all pin factors \tilde{w} of w);
- π_n is injective

then π_n would be a bijection between P_n and \mathcal{B}_n° , thus reducing the problem of enumerating the \boxplus -indecomposables of \mathcal{C}_w° to the (much simpler) combinatorial problem of counting the pin factors of w .

We do in fact have some intuitive reasons for suspecting that these two properties should hold, at least for sufficiently large n : the definition of the π -map (specifically, the placing of each new point in order to intersect the bounding rectangle of all previous points) seems almost designed to ensure that $\pi_{\tilde{w}}$ can have no proper non-trivial \circ -interval, and pin permutations have so much internal structure that it seems unlikely that two distinct pin sequences could conspire to generate the exact same centred permutation. Alas, both turn out not to be true for any n : for every length $n \geq 2$ there are pin sequences which generate \boxplus -decomposable permutations, as well as pairs of pin sequences which generate the same (centred) permutation. Fortunately, as our intuition suggests, both of these ‘bad behaviours’ are somewhat pathological and easily classified, so with some minor adjustments we will be able to make our strategy for counting \boxplus -indecomposables in a pin class work.

We begin by introducing some terminology for these pathological behaviours. First, those pin words whose image under the induced π -map is not contained in \mathcal{B}_n° :





Definition 3.32 (\boxplus -decomposable pin words). We call a (finite) pin word w a **\boxplus -decomposable pin word** if the centred permutation it generates, π_w° , is \boxplus -decomposable.








And those pin words which make injectivity of the induced π -map fail:

Definition 3.33 (Collisions). We call a set $\{w_1, w_2, \dots, w_k\}$ of pin words which generate the same centred permutation π° a **collision** (sometimes a **k -collision** when we wish to emphasise the size of the set) of pin words.

It is fairly easy to find all \boxplus -decomposables and collisions of length $n \leq 5$ through an exhaustive search, especially on applying symmetries of the square. This search reveals families of \boxplus -decomposables and collisions which generalise to all longer lengths. We can then, with considerable effort, prove that *all* examples of lengths greater than 6 belong to these families, giving us a full classification of these confounding behaviours. This is the content of the following two theorems: the statements will be crucial and will be quoted in full here; the proofs are technical and long and will be relegated to the appendix.

Theorem 3.34 (Classification of \boxplus -decomposables and Collisions). *The following two tables form a complete list of \boxplus -decomposable pin words and collisions:*

List of \boxplus-decomposable pin words:					
Length	Representative	Total number	Full List		
2	 1l	8	1l, 1d, 2r, 2d, 3u, 3r, 4l, 4u		
3	 1ld	8	1ld, 1dl, 2dr, 2rd, 3ru, 3ur, 4ul, 4lu		
$n \geq 4$	Type 1:  $1(ur)^kuld$ or $1(ru)^kld$	8	$n \geq 4$ even: $1(ur)^kuld,$ $1(ru)^krdl,$ $2(ul)^kurd,$ $2(lu)^kldr,$ $3(dl)^kdru,$ $3(ld)^klur,$ $4(dr)^kdlu,$ $4(rd)^krul$	$n \geq 5$ odd: $1(ru)^kld,$ $1(ur)^kdl,$ $2(lu)^krd,$ $2(ul)^kdr,$ $3(ld)^kru,$ $3(dl)^kur,$ $4(rd)^klu,$ $4(dr)^kul$	
	Type 2:  $1l(dl)^k$ or $1l(dl)^kd$	8	$n \geq 4$ even: $1l(dl)^k,$ $1d(ld)^k,$ $2r(dr)^k,$ $2d(rd)^k,$ $3r(ur)^k,$ $3u(ru)^k,$ $4l(ul)^k,$ $4u(lu)^k$	$n \geq 5$ odd: $1(ld)^k,$ $1(dl)^k,$ $2(rd)^k,$ $2(dr)^k,$ $3(ru)^k,$ $3(ur)^k,$ $4(lu)^k,$ $4(ul)^k$	

List of collisions of pin factors:			
Length	Representative	Total number	Full List
2	 1u = 1r	4 pairs	{1u, 1r}, {2l, 2u}, {3d, 3l}, {4r, 4d}
3	 1ul = 2ru	8 pairs	{1ul, 2ru}, {1rd, 4ur}, {2ur, 1lu}, {2ld, 3ul}, {3lu, 2dl}, {3dr, 4ld}, {4ru, 1dr}, {4dl, 3rd}
4	 1ldr = 2dru = 3rul = 4uld	2 quad.s	{1ldr, 2dru, 3rul, 4uld}, {1dlu, 2rdl, 3urd, 4lur}
5	Pathological:  1uldl = 3luru	4 pairs	{1uldl, 3luru}, {1rldd, 3drur}, {2urdr, 4rulu}, {2ldrd, 4dlul}
	Regular:  1ldlu = 2dlur	8 pairs	{1ldlu, 2dlur}, {1dldr, 4ldr}, {2rdru, 1drul}, {2drdl, 3rdlu}, {3urul, 2rudl}, {3rurd, 4urd}, {4luld, 3uldr}, {4ulur, 1lurd}
$n \geq 6$ even:	 $1(ld)^k_r = 2(dl)^k_dru$	8 pairs	{1(ld) ^k _r , 2(dl) ^k _{dru} }, {1(dl) ^k _u , 4(ld) ^k _{lur} }, {2(dr) ^k _u , 3(rd) ^k _{rul} }, {2(rd) ^k _l , 1(dr) ^k _{dlu} }, {3(ru) ^k _l , 4(ur) ^k _{uld} }, {3(ur) ^k _d , 2(ru) ^k _{rdl} }, {4(ul) ^k _d , 1(lu) ^k _{ldr} }, {4(lu) ^k _r , 3(ul) ^k _{urd} }
$n \geq 7$ odd:	 $1(ld)^k_lu = 2(dl)^k_ur$	8 pairs	{1(ld) ^k _{lu} , 2(dl) ^k _{ur} }, {1(dl) ^k _{dr} , 4(ld) ^k _{ru} }, {2(dr) ^k _{dl} , 3(rd) ^k _{lu} }, {2(rd) ^k _{ru} , 1(dr) ^k _{ul} }, {3(ru) ^k _{rd} , 4(ur) ^k _{dl} }, {3(ur) ^k _{ul} , 2(ru) ^k _{ld} }, {4(ul) ^k _{ur} , 1(lu) ^k _{rd} }, {4(lu) ^k _{ld} , 3(ul) ^k _{dr} }

Note: in the previous table we write $(f)^k$ to denote the factor f repeating k times for some $k \geq 0$.

Observation 3.35. *We emphasise some noteworthy features of these lists:*

1. *There is no overlap between the two lists: no pin word which generates a \boxplus -decomposable is also involved in a collision.*
2. *All collisions are pairs except for the two quadruples at length 4.*
3. *There are two distinct families of collisions at length 5: the **regular** length 5 collisions, which generalise to a family of collisions for all $n \geq 6$; and the **pathological** length 5 collisions, which are a family unique to length 5.*
4. *All even collisions of length $n \geq 4$ require all four quadrants, whereas the odd collisions of length $n \geq 5$ require only three.*

3.6 Enumeration of Recurrent Pin Classes

We now have a general strategy for obtaining the generating function of the \boxplus -closure of a pin class:

Procedure 3.36 ($\boxplus(\mathcal{C}_w^\circ)$ Enumeration Procedure). *Suppose that w is a pin word. We can find the generating function of the centred pin class $\boxplus(\mathcal{C}_w^\circ)$ by the following procedure:*

1. *Find the generating function $h(z)$ of the pin factors of w ;*
2. *Enumerate the pin factors of w that generate \boxplus -decomposable pin permutations, as well as colliding sets of pin factors of w , by comparison with the lists in Theorem 3.34. Subtract these from $h(z)$ to obtain $g(z)$, the generating function of \boxplus -indecomposables in \mathcal{C}_w° ;*
3. *Amend for commutativity: find the generating function $g_i(z)$ of the one-quadrant \boxplus -indecomposables of \mathcal{C}_w° in the i th quadrant, for each $i \in \{1, 2, 3, 4\}$. (In other words: enumerate the one-quadrant oscillations.) Set:*

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z),$$

this is the amended G -sequence of \mathcal{C}_w° ;

4. *Then the generating function of $\boxplus(\mathcal{C}_w^\circ)$ is given by*

$$f(z) = \frac{1}{1 - G(z)}$$

and the growth rate of $\boxplus(\mathcal{C}_w^\circ)$ is the reciprocal of the radius of convergence of $f(z)$.

Of course, if w is recurrent then \mathcal{C}_w° is \boxplus -closed and so $f(z)$ is in fact the generating function of \mathcal{C}_w° .

We illustrate this procedure with a sequence of examples, beginning with the class of increasing oscillations \mathcal{O}° , already defined in Fig. 13:

Example 3.37 (The Increasing Oscillations). The growth rate of \mathcal{O}° is $\kappa \approx 2.20557$, defined to be the reciprocal of the smallest positive real root of

$$1 - 2z - z^3 = 0$$

Proof. The defining pin sequence of \mathcal{O}° is $w = 1(\text{ru})^*$, which has two pin factors of every length ≥ 2 (corresponding to the two distinct starting positions within the period) and one pin factor (namely 1 itself) of length 1, so

$$h(z) = z + 2z^2 + 2z^3 + 2z^4 + 2z^5 \dots$$

By comparison with the lists in Theorem 3.34, there are no \boxplus -decomposables amongst the pin factors of w and precisely one colliding pair, namely $\{1u, 1r\}$, at length 2. We thus subtract 1 at length 2 to obtain the generating function of the \boxplus -indecomposables in \mathcal{O}° :

$$\begin{aligned} g(z) &= z + z^2 + 2z^3 + 2z^4 + 2z^5 + \dots \\ &= \frac{z + z^3}{1 - z} \end{aligned}$$

Noting that $g_2(z) = g_3(z) = g_4(z) = 0$ we see that $G(z) = g(z)$ and the generating function of \mathcal{O}° is

$$\begin{aligned} f(z) &= \frac{1}{1 - g(z)} \\ &= \frac{1 - z}{1 - 2z - z^3} \end{aligned}$$

and we may now deduce the growth rate by the Exponential Growth Theorem 2.22. □

We shall see later that κ is in fact the smallest possible growth rate of a pin class. For now we give three more classes as examples:




Figure 17: The pin class \mathcal{V}° , defined by the pin sequence $w = 2(\text{urul})^*$

Example 3.38 (Pin Class \mathcal{V}). The pin class \mathcal{V} is the underlying (uncentred) permutation class of the centred class $\mathcal{V}^\circ = \mathcal{C}_w^\circ$, where $w = 2(\text{urul})^*$ - see Fig. 17. Writing this pin sequence out with letters subscripted by the corresponding quadrant number

$$w = 2u_2r_1u_1l_2u_2r_1u_1l_2u_2r_1u_1l_2u_2r_1u_1l_2 \dots$$

we note immediately that w has 2 pin factors of length 1 and 4 of every greater length. By comparison with our list of \boxplus -decomposables and collisions, we note that we have to remove 2 from this count for lengths 2 and 3 (due to the \boxplus -decomposables $1l$ and $2r$ at length 2 and the collisions $\{1ul, 2ru\}$ and $\{2ru, 1lu\}$ at length 3) to obtain the generating function of the \boxplus -indecomposables of \mathcal{V}° :

$$\begin{aligned} g(z) &= 2z + 2z^2 + 2z^3 + 4z^4 + 4z^5 + 4z^6 + 4z^7 + \dots \\ &= \frac{2z + 2z^4}{1 - z} \end{aligned}$$

As we are in the upper half-plane, $g_3(z) = g_4(z) = 0$ so the amended G-sequence is just $g(z)$ itself, and the generating function of \mathcal{V}° is given by

$$\begin{aligned} f(z) &= \frac{1}{1 - g(z)} \\ &= \frac{1 - z}{1 - 3z - 2z^4} \end{aligned}$$

We deduce that $\text{gr}(\mathcal{V})$, which we shall call ν , is the reciprocal of the smallest real root of

$$1 - 3z - 2z^4 = 0;$$

explicitly:

$$\nu = \text{gr}(\mathcal{V}) = \text{gr}(\mathcal{V}^\circ) \approx 3.06918$$

Next, a class in three quadrants which will force us to confront commutativity:

Example 3.39 (The class \mathcal{Y}). Let w be the pin sequence $1(\text{uldlur})^*$. We refer to \mathcal{C}_w , the associated pin class, as \mathcal{Y} - see Fig. (18). Then the growth rate of \mathcal{Y} is γ , the reciprocal of the smallest positive real root of

$$1 - 4z + 2z^2 + z^3 - z^4 - 2z^5 - 3z^6 = 0.$$

Explicitly, $\gamma \approx 3.36637$.

Proof. We shall find the generating function of the corresponding centred class $\mathcal{Y}^\circ = \mathcal{C}_w^\circ$. We note that w , being periodic, is recurrent and we can thus use Procedure 3.36. First, we enumerate the pin factors of $w = 1(\text{uldlur})^*$; this will be easier if we first write out w with letters subscripted by the corresponding quadrant number:

$$w = 1u_1l_2d_3l_3u_2r_1u_1l_2d_3l_3u_2r_1u_1l_2d_3l_3u_2r_1u_1l_2d_3l_3u_2r_1 \dots$$




Figure 18: The pin class \mathcal{Y}° , generated by the pin sequence $w = 1(\text{uldlur})^*$.

Clearly there are three pin factors of length 1, and for $n \geq 2$ the six distinct starting points in the period give six pin factors of length n . Hence the generating function of the pin factors of w is

$$h(z) = 3z + 6z^2 + 6z^3 + 6z^4 + 6z^5 + 6z^6 + 6z^7 + \dots$$

We now list the collisions and \boxplus -decomposables found amongst the pin factors of w by comparison with the lists given in Theorem 3.34:

- Collisions:** $\{1\text{ul}, 2\text{ru}\}, \{2\text{dl}, 3\text{lu}\}$ (length 3)
 $\{1\text{uldl}, 3\text{luru}\}, \{1\text{ldlu}, 2\text{dlur}\}, \{3\text{urul}, 2\text{ru ld}\}$ (length 5)
 \boxplus -dec.s: $1\text{l}, 2\text{d}, 3\text{u}, 2\text{r}$ (length 2)
 $1\text{ld}, 3\text{ur}$ (length 3)
 $1\text{uld}, 1\text{ldl}, 3\text{lur}, 3\text{uru}$ (length 4)

Hence we subtract these from the generating function of pin factors to obtain the generating function of \boxplus -indecomposables of \mathcal{Y}° :

$$\begin{aligned} g(z) &= h(z) - (2z^3 + 3z^5) - (4z^2 + 2z^3 + 4z^4) \\ &= 3z + 2z^2 + 2z^3 + 2z^4 + 3z^5 + 6z^6 + 6z^7 + 6z^8 + 6z^9 + \dots \end{aligned}$$

Next we must enumerate the one-quadrant \boxplus -indecomposables in each quadrant; this is easily done by looking at the diagram: $g_1(z) = g_3(z) = z + z^2$, $g_2(z) = z$ and $g_4(z) = 0$. Hence we can calculate the amended G-sequence:

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &= 3z + z^2 + z^4 + 3z^5 + 6z^6 + 6z^7 + 6z^8 + 6z^9 + \dots \\ &= \frac{3z - 2z^2 - z^3 + z^4 + 2z^5 + 3z^6}{1 - z} \end{aligned}$$

Finally, by the Generating Function Specification 2.20, the generating function of \mathcal{Y}° is given by:

$$\begin{aligned} f(z) &= \frac{1}{1 - G(z)} \\ &= \frac{1 - z}{1 - 4z + 2z^2 + z^3 - z^4 - 2z^5 - 3z^6} \end{aligned}$$

This is a rational function, whose singularities are precisely the roots of

$$1 - 4z + 2z^2 + z^3 - z^4 - 2z^5 - 3z^6 = 0$$

By computation, the smallest root of this equation is positive and real and has reciprocal $\gamma \approx 3.36637$, which by the Exponential Growth Theorem is therefore the growth rate of \mathcal{Y} . \square

And finally, a four-quadrant pin class that has in fact been studied before in the literature: the Widdershins Spiral, \mathcal{W}° , generated by the pin sequence $w = 1(\text{ldru})^*$ (see Fig. 19). This class was first introduced by Murphy [13], who gave the generating function of the uncentred class by an explicit enumeration. Using our specification we can now give an alternative derivation of its growth rate:




Figure 19: The Widdershins Spiral: this is the smallest pin class in four quadrants.

Example 3.40 (The Widdershins Spiral).

Let $\mathcal{W} = \mathcal{C}_w$ be the pin class generated by the pin sequence $w = 1(\text{ldru})^*$. Then the generating function of the corresponding centred class \mathcal{W}° is given by:

$$f(z) = \frac{1 - z}{1 - 5z + 6z^2 - 2z^3 - z^4 - 3z^5}$$

Hence $\omega_0 = \text{gr}(\mathcal{W}^\circ) \approx 3.48806$.

Proof. The defining pin sequence w is recurrent so we can enumerate \mathcal{W}° using our standard procedure. We first determine the generating function of pin factors of w : as w has period 4 and all four pin factors are already distinct at length 1 we have:

$$h(z) = 4z + 4z^2 + 4z^3 + 4z^4 + 4z^5 + 4z^6 + \dots$$

We must amend this for collisions and \boxplus -decomposables to obtain the generating function $g(z)$ of \boxplus -indecomposables in \mathcal{W}° . By comparison with the lists in Theorem 3.34 we see that the only collision amongst the pin factors of w is the colliding quadruple $\{1\text{ldr}, 2\text{dru}, 3\text{rul}, 4\text{uld}\}$

and that there are precisely four \boxplus -decomposables of length 2 (1l, 2d, 3r and 4u), four \boxplus -decomposables of length 3 (1ld, 2dr, 3ru and 4ul), and none of length $n \geq 4$. Hence the generating function of \boxplus -indecomposables in \mathcal{W}° is given by:

$$\begin{aligned} g(z) &= h(z) - 3z^4 - (4z^2 + 4z^3) \\ &= 4z + z^4 + 4z^5 + 4z^6 + 4z^7 + 4z^8 + \dots \\ &= \frac{4z - 4z^2 + z^4 + 3z^5}{1 - z} \end{aligned}$$

Next we note that there is precisely 1 one-quadrant \boxplus -indecomposable of length 1 in each quadrant and none of any greater length:

$$g_1(z) = g_2(z) = g_3(z) = g_4(z) = z$$

Hence the amended G-sequence of \mathcal{W}° is given by:

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &= \frac{4z - 4z^2 + z^4 + 3z^5}{1 - z} - 2z^2 \\ &= \frac{4z - 6z^2 + 2z^3 + z^4 + 3z^5}{1 - z} \end{aligned}$$

And so, by the Generating Function Specification 2.20, we can derive the generating function of \mathcal{W}° :

$$\begin{aligned} f(z) &= \frac{1}{1 - G(z)} \\ &= \frac{1 - z}{1 - 5z + 6z^2 - 2z^3 - z^4 - 3z^5} \end{aligned}$$

as required. Finally, we can calculate the growth rate ω_0 of \mathcal{W}° as the reciprocal of the smallest positive real root of the denominator: $\omega_0 \approx 3.48806$. \square

4 Non-recurrent Pin Classes

We have now established a general method for determining the growth rate and generating function of a pin class \mathcal{C}_w defined by a *recurrent* pin sequence w over the language \mathcal{L} . This also allows us to determine the growth rate of an *eventually* recurrent pin class, by the Finite Prefix Lemma 3.14. This leaves the non-eventually-recurrent case: these classes are not \boxplus -closed, so the methods outlined in the previous section will not enable us to determine their generating functions (in fact, the author does not know the generating function of *any* not-eventually-recurrent pin class). Somewhat surprisingly, however, we *can* at least determine the *growth rate* of a general non-recurrent pin class \mathcal{C}_w° . The key idea here is simple: the classes that we know how to enumerate are the \boxplus -closed classes, so we enumerate \mathcal{C}_w° by taking better and better \boxplus -closed approximations. The limiting behaviour of these approximations will give us the growth rate, but *not* the generating function, of \mathcal{C}_w° .

We begin by noting that we already know how to bound \mathcal{C}_w° by a \boxplus -closed class from above: Procedure 3.36 gives us the generating function of the \boxplus -closure of \mathcal{C}_w° ; in the non-recurrent case this is not equal to \mathcal{C}_w° but will function as an upper bound as $\mathcal{C}_w^\circ \subseteq \boxplus\mathcal{C}_w^\circ$. We also note that, like pin classes, \boxplus -closures of pin classes have proper growth rates:

Lemma 4.1. *Let w be a pin sequence. Then $\boxplus\mathcal{C}_w^\circ$ has a proper growth rate.*

Proof. This is an application of Theorem 3.29: $\boxplus\mathcal{C}_w^\circ$ is a \boxplus -closed class contained in the complete pin class \mathcal{P}° and satisfies that adjacency condition by the same reasoning as $\boxplus\mathcal{C}_w^\circ$. \square

Hence, for any pin sequence w , $\text{gr}(\mathcal{C}_w^\circ) \leq \text{gr}(\boxplus\mathcal{C}_w^\circ)$.

4.1 The \boxplus -interior of a Pin Class

We now wish to determine a *lower* bound on the growth rate of a pin class, which we again do by comparison with a \boxplus -closed class whose generating function we can find. We thus consider the ‘largest \boxplus -closed subclass contained in \mathcal{C}_w° ’, which we refer to as the \boxplus -interior of \mathcal{C}_w° . In order to define this rigorously, recall that if w is an infinite pin word over the language \mathcal{L} , we use the notation $w_{i,j}$ to denote the pin factor of w taken between the i th and j th places. Now we can define:

Definition 4.2 (Recurrent Pin Factors). Let w be a pin sequence and \tilde{w} a (finite) pin word. We say that \tilde{w} is a **recurrent pin factor** of w if for all $n \in \mathbb{N}$ there exist $j \geq i \geq n$ such that $w_{i,j} = \tilde{w}$.

We call a \boxplus -indecomposable permutation $\pi_{\tilde{w}}^\circ$ generated by a recurrent pin factor \tilde{w} of w a **recurrent \boxplus -indecomposable** in \mathcal{C}_w° .

That is, a recurrent pin factor of w is a pin factor that occurs in w infinitely-often. We may now define:

Definition 4.3 (The \boxplus -interior of a pin class).

Let w be a pin sequence with pin class \mathcal{C}_w° . We define the **\boxplus -interior**, \mathcal{C}_w^\boxplus , of \mathcal{C}_w° to be the \boxplus -closure of the set of pin permutations of the form $\pi_{\tilde{w}}^\circ$ where \tilde{w} is a *recurrent* pin factor of w .

We then have:

Observation 4.4 (Basic properties of the \boxplus -interior). *For any pin sequence w over \mathcal{L} :*

1. \mathcal{C}_w^\boxplus is a (non-empty) \boxplus -closed subclass of \mathcal{C}_w° ;
2. \mathcal{C}_w^\boxplus is the union of all \boxplus -closed subclasses of \mathcal{C}_w° ;
3. $\mathcal{C}_w^\boxplus = \mathcal{C}_w^\circ$ if and only if \mathcal{C}_w° is \boxplus -closed;
4. $\sigma^\circ \in \mathcal{C}_w^\boxplus$ if and only if for all $i \in \mathbb{N}$ there exists $j \geq i$ such that $\sigma^\circ \leq \pi_{w_{i,j}}^\circ$.
5. \mathcal{C}_w^\boxplus is the \boxplus -closure of the set of recurrent \boxplus -indecomposables in \mathcal{C}_w° .

Informally, the \boxplus -interior of \mathcal{C}_w° is the set of all centred permutations that can be found in the pin diagram of w in infinitely many (non-overlapping) instances.

As the \boxplus -interior of a pin class is \boxplus -closed, we can determine its generating function in terms of its G-sequence, which we can obtain from w as follows:

Lemma 4.5. *Suppose that $g(z)$ is the generating function of the **recurrent** \boxplus -indecomposables in \mathcal{C}_w° , and $g_1(z), g_2(z), g_3(z), g_4(z)$ are the generating functions of the **recurrent one-quadrant** \boxplus -indecomposables in quadrants 1, 2, 3, 4, respectively. Then*

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z)$$

is the amended G-sequence for \mathcal{C}_w^\boxplus , and

$$f(z) = \frac{1}{1 - G(z)}$$

is the generating function of \mathcal{C}_w^\boxplus .

Proof. Immediate from the Generating Function Specification on noting that \mathcal{C}_w° is a \boxplus -closed class and the recurrent \boxplus -indecomposables of \mathcal{C}_w° are its \boxplus -indecomposables. \square

As with the \boxplus -closure we briefly note the following:

Lemma 4.6. *Let w be a pin sequence. Then \mathcal{C}_w^\boxplus has a proper growth rate.*

Proof. If \mathcal{C}_w^\boxplus contains $\uparrow\bullet$ and $\bullet\downarrow$ then w contains 1 and 3 as recurrent pin factors, which is to say that w visits quadrants 1 and 3 infinitely-often. Every time w moves from quadrant 1 to quadrant 3 it must pass through either quadrant 2 or 4, and so must also visit at least one of these quadrants infinitely-often; hence \mathcal{C}_w^\boxplus also contains either $\bullet\downarrow$ or $\uparrow\bullet$ and so w satisfies the adjacency condition. The same reasoning works if \mathcal{C}_w^\boxplus contains $\downarrow\bullet$ or $\bullet\uparrow$, and so we conclude that \mathcal{C}_w^\boxplus satisfies the adjacency condition. As \mathcal{C}_w^\boxplus is \boxplus -closed we can now apply Theorem 3.29 to deduce that \mathcal{C}_w^\boxplus has a proper growth rate. \square

Recall that $\kappa \approx 2.20557$ is the growth rate of the class \mathcal{O}° of increasing oscillations. We shall require the following elementary bound:

Lemma 4.7. *Let w be a pin sequence. Then $\text{gr}(\mathcal{C}_w^\boxplus) \geq \kappa$.*

Proof. • If w visits only one quadrant recurrently, which by symmetry we may take to be the first quadrant, then $w = \tilde{w}(\text{ur})^*$, where \tilde{w} is some finite prefix. Hence \mathcal{C}_w^\boxplus is just \mathcal{O}° with a finite prefix, and so has growth rate κ .

- Suppose w visits precisely two (necessarily adjacent) quadrants recurrently - by symmetry we may take these to be quadrants 1 and 2. Then after some finite prefix, w stays in the upper half-plane, and moves between quadrants 1 and 2 infinitely-often. Hence w contains 1l and 2r as recurrent pin factors, and these must extend to recurrent pin factors

1lu and 2ru (as w stays in the upper half-plane from this point). Hence π_{1lu}° and π_{2ru}° are contained in \mathcal{C}_w^\boxplus and we note that $\bullet\mid$ and $\mid\bullet$ are contained in these two pin permutations, respectively. Hence \mathcal{C}_w^\boxplus contains the \boxplus -closure $\boxplus\{\bullet\mid, \mid\bullet\}$, which by calculation has growth rate $\approx 2.73205 > \kappa$.

- Suppose w visits three or four quadrants recurrently - by symmetry assume these include quadrants 1, 2 and 3. Then \mathcal{C}_w^\boxplus contains $\boxplus\{\mid\bullet, \bullet\mid, \bullet\mid\bullet\}$, which by calculation has growth rate $\approx 2.61803 > \kappa$.

□

We note that this now implies the following as a corollary:

Corollary 4.8. *Let w be a pin sequence. Then:*

- $\overline{gr}(\mathcal{C}_w) \geq \kappa$
- $\overline{gr}(\mathcal{C}_w) = \kappa$ if and only if $w = \tilde{w}(ur)^*$ or some symmetry of this form, where \tilde{w} is a finite prefix.

4.2 G-sequence properties

Let w be a pin sequence, and let \mathcal{C}° be either the pin class generated by w or its \boxplus -interior. We now know how to obtain the amended G-sequence $G(z)$ of \mathcal{C}° , by enumerating either the pin factors or recurrent pin factors of w . Then the generating function of $\boxplus(\mathcal{C}^\circ)$ (note that this is \mathcal{C}° itself in the \boxplus -interior case) is given by:

$$f(z) = \frac{1}{1 - G(z)}$$

We know that the growth rate ρ of $\boxplus(\mathcal{C}^\circ)$ is equal to the reciprocal of the radius of convergence of $f(z)$. We should like to be able to deduce from this that $G(\rho^{-1}) = 1$, as this will give us a strategy for studying growth rates of pin classes in terms of analytic properties of $G(z)$, but this will require a slightly more thorough study of the properties of $G(z)$:

Proposition 4.9 (G-sequence of pin classes and \boxplus -interiors). *Let w be a pin sequence and suppose that $G(z) = \sum_{n=1}^{\infty} a_n z^n$ ($a_n \in \mathbb{Z}$) is the amended G-sequence of a class \mathcal{C}° , which is **either** the pin class generated by w **or** its \boxplus -interior. Then:*

1. $a_1 \in \{1, 2, 3, 4\}$.
2. For all $n \geq 2$:

$$-8n \leq a_n < 2^{n+2}$$

3. $G(z)$ converges to a smooth function on the interval $[0, \frac{1}{2})$, with $G(0) = 0$.
4. $G(z) = 1$ has a solution in $[0, \frac{1}{\kappa}]$.

5. Let α be the smallest positive real solution of $G(z) = 1$. The growth rate of $\boxplus \mathcal{C}^\circ$ is equal to α^{-1} .

Proof. Recall that

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \quad (2)$$

where $g(z)$ counts the \boxplus -indecomposables in \mathcal{C}° and $g_i(z)$ counts the one-quadrant \boxplus -indecomposables contained in the i th quadrant.

1. This is clear from 2: the z -term of $G(z)$ must agree with the z -term of $g(z)$ (as the products $g_1(z)g_3(z)$ and $g_2(z)g_4(z)$ have no term of degree lower than z^2), and this simply counts the number of quadrants that \mathcal{C}° visits (either at all or recurrently), which is clearly either 1, 2, 3 or 4.
2. Let \leq denote coefficient-wise ordering on formal power series, so $p(z) \leq q(z)$ means that the z^n -coefficient of $p(z)$ is less than or equal to the z^n -coefficient of $q(z)$ for all $n \in \mathbb{N}$. Then, from 2 (and using the fact that the coefficients of $g(z), g_i(z)$ are all non-negative):

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &\leq g(z) \\ &\leq 4z + \sum_{n=2}^{\infty} 2^{n+2}z^n \end{aligned}$$

where the final inequality is simply the number of pin words of length n over \mathcal{L} .

On the other hand, using the fact that the generating function of *all* one-quadrant boxplus-indecomposable pin permutations in the i th quadrant is

$$g_i(z) = z + z^2 + 2z^3 + 2z^4 + 2z^5 + \dots = \frac{z + z^3}{1 - z},$$

we obtain:

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &\geq -g_1(z)g_3(z) - g_2(z)g_4(z) \\ &\geq -2 \frac{(z + z^3)^2}{(1 - z)^2} \\ &= -2z^2 - 4z^3 - 10z^4 - 16z^5 - 24z^6 - 32z^7 - 40z^8 - 48z^9 - \dots \\ &\geq \sum_{n=1}^{\infty} -8nz^n \end{aligned}$$

Combining these two inequalities gives the desired result.

3. This is from the Ratio Test, on noting that the coefficients of $G(z)$ have magnitudes bounded by powers of 2.

4. We know that the all the classes we are dealing with have growth rates greater than or equal to κ . Hence the generating function of $\boxplus \mathcal{C}^\circ$, namely

$$f(z) = \frac{1}{1 - G(z)},$$

has radius of convergence (in \mathbb{C}) $R \leq \frac{1}{\kappa} < \frac{1}{2}$. As $f(z)$ has non-negative coefficients (regardless of whether $G(z)$ does), we can apply Pringsheim's Theorem [11, Theorem IV.7] to deduce that $f(z)$ has a singularity at $z = R$. Singularities of $f(z)$ correspond either to singularities of $G(z)$ or solutions of $G(z) = 1$. But $G(z)$ converges on $[0, \frac{1}{2})$ which includes R , so $G(R) = 1$.

5. We saw that R solves $G(z) = 1$ in the proof of 4.; there cannot be a smaller solution to this equation as that would give a singularity with magnitude smaller than R . Now apply the Exponential Growth Theorem 2.22.

□

We require one more property of G -sequences in the \boxplus -interior case only:

Lemma 4.10. *Let $G(z)$ be the amended G -sequence of the \boxplus -interior \mathcal{C}_w^\boxplus of some pin class \mathcal{C}_w° . Let α be the smallest positive real root of the equation $G(z) = 1$, guaranteed to exist by Proposition 4.9. Then $G(z)$ is positive on the interval $(0, \alpha)$.*

Proof. We note that this property is obvious if $G(z)$ has non-negative coefficients, which immediately deals with the case in which w visits only two of the quadrants (either recurrently or otherwise). We next suppose that \mathcal{C}° has points in precisely three quadrants, which, without loss of generality, we take to be quadrants 1, 2 and 3. Then \mathcal{C}° must contain the permutations $\uparrow\bullet$, $\bullet\uparrow$, $\bullet\uparrow\bullet$, $\uparrow\bullet\uparrow$ and $\uparrow\bullet\uparrow\bullet$ (as the defining pin sequence w must turn around in the first and third quadrants infinitely-often), and so:

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) \\ &\geq (3z + 2z^2 + 2z^3) - \frac{(z+z)^2}{(1-z)^2} \\ &= \frac{3z - 5z^2 + z^3 - 4z^4 + 2z^5 - z^6}{(1-z)^2} \end{aligned}$$

Hence:

$$G(z) = \frac{3z - 5z^2 + z^3 - 4z^4 + 2z^5 - z^6}{(1-z)^2} + F(z)$$

where $F(z)$ has non-negative coefficients. By computation the function given here is positive on the desired range, and $F(z)$ is certainly positive here due to non-negativity of its coefficients. Hence $G(z)$ is positive on $(0, \alpha)$. Finally, if $G(z)$ visits all four quadrants then:

$$\begin{aligned} G(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &\geq 4z - 2\frac{(z+z)^2}{(1-z)^2} \\ &= \frac{4z - 10z^2 + 4z^3 - 4z^4 - 2z^6}{(1-z)^2} \end{aligned}$$

and as in the three quadrant case, this ‘worst case scenario’ function is nevertheless still positive in the desired range. \square

4.3 \mathcal{C}_w° has same growth rate as \mathcal{C}_w^\boxplus

The moral of the preceding sections is that we can sandwich *any* pin class (recurrent or otherwise) between two \boxplus -closed (centred) permutation classes that we know how to enumerate. If the pin class happens to be recurrent then these containments are in fact equalities and we have enumerated the pin class. If the pin class is *not* recurrent then we at least have bounds on the growth rate:

$$\text{gr}(\mathcal{C}_w^\boxplus) \leq \text{gr}(\mathcal{C}_w^\circ) \leq \text{gr}(\boxplus\mathcal{C}_w^\circ) \quad (3)$$

The aim of this section is to prove the remarkable result that the left-hand inequality in (3) is in fact an equality; that is, the growth rate of a pin class is always equal to that of its \boxplus -interior. This will enable us to conclude that *any* pin class (recurrent or otherwise) has a proper growth rate, and will enable us to determine this growth rate providing that we can enumerate the recurrent pin factors of the defining pin sequence w . The core idea in the proof is relatively simple to understand: if w is a pin sequence we know by the Finite Prefix Theorem that dfg and cvgbh have the same (upper) growth rate and we note that if we take n sufficiently large then the pin factors of dfg are precisely ...

Proposition 4.11. *Let \mathcal{C}_w° be a pin class generated by a pin sequence w . Let \mathcal{C}_w^\boxplus be the \boxplus -interior of \mathcal{C}_w° , and write $w_{\geq n}$ for the left-truncation of w starting in the n th position. Then:*

$$\lim_{n \rightarrow \infty} \text{gr}(\boxplus\mathcal{C}_{w_{\geq n}}^\circ) = \text{gr}(\mathcal{C}_w^\boxplus)$$

Proof. Let $g(z)$ be the generating function of the \boxplus -indecomposables of \mathcal{C}_w^\boxplus , and $g_i(z)$ be the generating function of one-quadrant \boxplus -indecomposables of \mathcal{C}_w^\boxplus in the i th quadrant. Then

$$G(z) = g(z) - g_1(z)g_3(z) - g_2(z)g_4(z)$$

is the amended G -sequence of \mathcal{C}_w^\boxplus , and $\alpha = \text{gr}(\mathcal{C}_w^\boxplus)^{-1}$ is the smallest positive real root of the equation $G(z) = 1$.

Now, let $t \in \mathbb{N}$, and consider the set \mathcal{W}_t^* of all **non**-recurrent pin factors of w of length $\leq t$. This is a finite set and each element of it occurs as a pin factor of w only a finite number of times, so there is some $n(t) \in \mathbb{N}$ such that no element of \mathcal{W}_k^* is contained as a pin factor of $w_{\geq n(t)}$ (for concreteness, we may take $n(t)$ to be the *smallest* positive integer with this property). But of course *all* of the recurrent pin factors of w of length $\leq t$ occur as pin factors of $w_{\geq n(t)}$, so we deduce that the pin factors of $w_{\geq n(t)}$ of length $\leq t$ are precisely the recurrent pin factors of w of length $\leq t$. Hence if we write $g_t(z)$ for the generating function of the \boxplus -indecomposables of $\mathcal{C}_{w_{\geq n(t)}}^\circ$, and $g_{t,i}(z)$ for generating function of the one-quadrant \boxplus -indecomposables of $\mathcal{C}_{w_{\geq n}}^\circ$ in the i th quadrant, then $g_t(z), g_{t,1}(z), g_{t,2}(z), g_{t,3}(z), g_{t,4}(z)$ will

agree with $g(z), g_1(z), g_2(z), g_3(z), g_4(z)$, respectively, up to and including the z^t -term. Hence the amended G -sequence of $\mathcal{C}_{w \geq n(t)}^\circ$, namely

$$G_t(z) = g_t(z) - g_{t,1}(z)g_{t,3}(z) - g_{t,2}(z)g_{t,4}(z),$$

agrees with $G(z)$ up to and including the z^t -term.

Fix a t : we now consider some basic analytic facts about the functions $G(z)$ and $G_t(z)$. First, note that by Proposition 4.9, $G(z)$ and $G_t(z)$ are smooth functions defined on the interval $[0, \frac{1}{\kappa}]$. Further, $G(z) = 1$ and $G_t(z) = 1$ have solutions in this interval; we call the smallest solution to these equations in this interval α and α_t , respectively. Then \mathcal{C}_w^{\boxplus} and $\boxplus \mathcal{C}_{w \geq n(t)}^\circ$ have growth rates $\rho = \alpha^{-1}$ and $\rho_t = \alpha_t^{-1}$, respectively. Note that we have:

$$\rho = \text{gr}(\mathcal{C}_w^{\boxplus}) \leq \text{gr}(\mathcal{C}_w^\circ) = \text{gr}(\mathcal{C}_{w \geq n(t)}^\circ) \leq \text{gr}(\boxplus \mathcal{C}_{w \geq n(t)}^\circ) = \rho_t$$

where the two inequalities follow from containment of the corresponding classes, and the middle equality follows from the Finite Prefix Lemma 3.14. Note that this implies that $\alpha_t \leq \alpha$.

Now, write $G_t(z) = \sum_{n=1}^{\infty} a_n z^n$ and $G(z) = \sum_{n=1}^{\infty} b_n z^n$: then each a_n, b_n is an integer (not necessarily positive) with magnitude bounded by 2^{n+2} and $a_n = b_n$ for all $n \leq t$. We combine these facts to deduce a bound on the difference between these two functions:

$$\begin{aligned} |G_t(z) - G(z)| &= \left| \sum_{n=t+1}^{\infty} (a_n - b_n) z^n \right| \\ &\leq \sum_{n=t+1}^{\infty} |a_n - b_n| z^n \\ &\leq \sum_{n=t+1}^{\infty} (2^{n+2} + 8n) z^n \\ &\leq \sum_{n=t+1}^{\infty} 2^{n+3} z^n \\ &= \frac{8(2z)^{t+1}}{1-2z} \end{aligned}$$

and so, for $z \in [0, \frac{1}{\kappa}]$:

$$\begin{aligned} |G_t(z) - G(z)| &\leq \sup_{z \in [0, \frac{1}{\kappa}]} \left\{ \frac{8(2z)^{t+1}}{1-2z} \right\} \\ &= \frac{8(\frac{2}{\kappa})^{t+1}}{1 - \frac{2}{\kappa}} \\ &= \frac{16}{\kappa - 2} \cdot \left(\frac{2}{\kappa} \right)^t \end{aligned}$$

Note, crucially, that (as $\kappa > 2$) this expression approaches 0 as $t \rightarrow \infty$. We claim that this fact implies that $\alpha_t \rightarrow \alpha$:

Let $\epsilon > 0$ and consider $G(z)$ on the interval $[0, \alpha - \epsilon]$. As a continuous function on a closed interval, $G(z)$ achieves a maximum value M on $[0, \alpha - \epsilon]$. Further, as $G(z)$ is positive on $(0, \alpha]$ (by Lemma 4.10) and $G(z) = 1$ does not have a root in $[0, \alpha - \epsilon]$ (as α is, by definition, the *smallest* positive real root of $G(z) = 1$), M must be a positive number smaller than 1. Now, choose a $K \in \mathbb{N}$ such that:

$$\frac{16}{\kappa - 2} \cdot \left(\frac{2}{\kappa}\right)^K < 1 - M$$

and take any $t \geq K$. Then, for $z \in [0, \alpha - \epsilon]$:

$$\begin{aligned} |G_t(z)| &= |(G_t(z) - G(z)) + G(z)| \\ &\leq |G_t(z) - G(z)| + |G(z)| \\ &\leq |G_t(z) - G(z)| + G(z) \\ &\leq \frac{16}{\kappa - 2} \cdot \left(\frac{2}{\kappa}\right)^t + M \\ &< (1 - M) + M \\ &= 1 \end{aligned}$$

Hence, in particular, $G_t(z) = 1$ does not have a root in $[0, \alpha - \epsilon]$. But $G_t(z) = 1$ certainly does have a root, namely α_t , which is smaller than α . Hence $\alpha_t \in (\alpha - \epsilon, \alpha]$.

We have thus proved that for every $\epsilon > 0$ there exists $K \in \mathbb{N}$ such that $\alpha_t \in (\alpha - \epsilon, \alpha]$ for all $t \geq K$; hence $\alpha_t \rightarrow \alpha$ as $t \rightarrow \infty$. Hence $\text{gr}(\boxplus_{\geq n(t)} \mathcal{C}_w^\circ) \rightarrow \text{gr}(\mathcal{C}_w^\boxplus)$ as $t \rightarrow \infty$, as required. \square

We may now finally deduce the main result of this paper:

Theorem 4.12. *Let w be a pin sequence. Then the associated pin class \mathcal{C}_w° (along with its uncentred counterpart \mathcal{C}_w) has a proper growth rate which is equal to that of its \boxplus -interior, \mathcal{C}_w^\boxplus*

Proof. Suppose w is a pin sequence. Then $\mathcal{C}_w^\boxplus \subseteq \mathcal{C}_w^\circ$ and so

$$\text{gr}(\mathcal{C}_w^\boxplus) = \underline{\text{gr}}(\mathcal{C}_w^\boxplus) \leq \underline{\text{gr}}(\mathcal{C}_w^\circ) \quad (4)$$

where the left-hand equality is due to the fact that \boxplus -interiors have (proper) growth rates. Now, letting $n \in \mathbb{N}$ we have

$$\overline{\text{gr}}(\mathcal{C}_w^\circ) = \overline{\text{gr}}(\mathcal{C}_{w_{\geq n}}^\circ) \quad (5)$$

by the Finite Prefix Lemma 3.14, and by containment we also have

$$\overline{\text{gr}}(\mathcal{C}_{w_{\geq n}}^\circ) \leq \overline{\text{gr}}(\boxplus \mathcal{C}_{w_{\geq n}}^\circ) = \text{gr}(\boxplus \mathcal{C}_{w_{\geq n}}^\circ) \quad (6)$$

where the right-hand equality follows from existence of proper growth rates of \boxplus -closures of pin classes.

Combining equations (4), (5) and (6) yields:

$$\text{gr}(\mathcal{C}_w^\boxplus) \leq \underline{\text{gr}}(\mathcal{C}_w^\circ) \leq \overline{\text{gr}}(\mathcal{C}_w^\circ) \leq \text{gr}(\boxplus \mathcal{C}_{w_{\geq n}}^\circ)$$

and taking limits (using Proposition 4.11) as $n \rightarrow \infty$ forces

$$\underline{\text{gr}}(\mathcal{C}_w^\circ) = \overline{\text{gr}}(\mathcal{C}_w^\circ)$$

as required. \square

4.4 The Complete Class \mathcal{P}_c°

We illustrate the use of this theory by explicitly enumerating the (centred) complete pin class \mathcal{P}_c° . Recall that this is the class of *all* pin permutations. It may not be immediately obvious, but \mathcal{P}_c° is itself a pin class, and is in fact the *only* pin class that achieves its growth rate:

Proposition 4.13 (Complete Pin Class).

1. There is a pin sequence w_c which contains every finite pin word as a (recurrent) pin factor.
2. For any such pin sequence w_c , $\mathcal{C}_w^\circ = \mathcal{P}_c^\circ$.
3. The complete pin class has growth rate $\omega_\infty \approx 5.24112$, where ω_∞ is defined to be the reciprocal of the smallest positive real root of the equation

$$1 - 8z + 19z^2 - 26z^3 + 14z^4 - 12z^5 - 8z^6 + 20z^7 - 8z^8 = 0$$

Proof. 1. Note that \mathcal{L}^* is countable, we can thus order all \mathcal{L}^* -words and concatenate them (after any initial numeral) in this order, possibly placing a letter in between consecutive words to ensure the alignments alternate. This resulting pin sequence w_c will contain all \mathcal{L}^* -words as subword factors, and will hence contain all possible finite pin words as pin factors.

2. This is clear by the definition of a pin permutation.
3. First, we enumerate the set of pin factors of w_c , which is to say the set of all finite pin words. Clearly, there are 4 pin words of length 1. For length $n \geq 2$, there are 4 choices for the initial numeral, 4 choices for the letter in second place, and then 2 choices for every subsequent letter as the alignments must now alternate. Hence there are 2^{n+2} pin words of length $n \geq 2$, and the generating function of the set of all finite pin words is given by:

$$h(z) = 4z + \sum_{n=2}^{\infty} 2^{n+2} z^n$$

This will be an overcount of the \boxplus -indecomposables in \mathcal{P}° due to collisions and \boxplus -decomposable pin words, but we have already classified all of these in Theorem 3.34. The generating function of the overcount due to collisions is

$$h_{\text{Col}}(z) = 4z^2 + 8z^3 + 6z^4 + 12z^5 + 8z^6 + 8z^7 + 8z^8 + 8z^9 + \dots$$

(Note: this is the generating function of the *overcount* due to collisions. At all $n \neq 4$ it is simply equal to the *number* of collisions, as these are all pairs so we need only take away one from the count for each collision. But at $n = 4$ there are two colliding *quadruples* so the overcount is 6.) And the generating function of the \boxplus -decomposable pin words is

$$h_{\boxplus\text{-Dec.}}(z) = 8z^2 + 8z^3 + 16z^4 + 16z^5 + 16z^6 + 16z^7 + 16z^8 + 16z^9 + \dots$$

Subtracting these from $h(z)$ gives us the generating function of the \boxplus -indecomposable pin permutations:

$$\begin{aligned} g(z) &= h(z) - h_{\text{Col}}(z) - h_{\boxplus\text{-Dec.}}(z) \\ &= 4z + 4z^2 + 16z^3 + 42z^4 + 100z^5 + \sum_{n=6}^{\infty} (2^{n+2} - 24)z^n \\ &= \frac{4z - 8z^2 + 12z^3 + 2z^4 + 6z^5 + 16z^6 - 8z^7}{(1-z)(1-2z)} \end{aligned}$$

Next, we shall require the generating functions $g_i(z)$ ($i \in \{1, 2, 3, 4\}$) of the \boxplus -indecomposable pin permutations entirely contained in the i th quadrant. Clearly, these are all equal to the generating function of the oscillations; namely:

$$\begin{aligned} g_1(z) = g_2(z) = g_3(z) = g_4(z) &= z + z^2 + 2z^3 + 2z^4 + 2z^5 + 2z^6 + \dots \\ &= \frac{z + z^3}{1 - z} \end{aligned}$$

Hence we obtain the amended G-sequence for the complete class of pin permutations:

$$\begin{aligned} G_{\infty}(z) &= g(z) - g_1(z)g_3(z) - g_2(z)g_4(z) \\ &= \frac{4z - 14z^2 + 24z^3 - 14z^4 + 12z^5 + 8z^6 - 20z^7 + 8z^8}{(1-z)^2(1-2z)} \end{aligned}$$

Thus, we can now apply the Generating Function Specification to obtain the generating function for the class of all (centred) pin permutations:

$$\begin{aligned} f(z) &= \frac{1}{1 - G_{\infty}(z)} \\ &= \frac{(1-z)^2(1-2z)}{1 - 8z + 19z^2 - 26z^3 + 14z^4 - 12z^5 - 8z^6 + 20z^7 - 8z^8} \end{aligned}$$

By the Exponential Growth Rate formula, and the fact that $f(z)$ has positive coefficients, the growth rate of \mathcal{P}° , and hence also the uncentred class \mathcal{P} , is the reciprocal of the smallest positive real root of the denominator, as required.

□

5 Concluding Remarks

We have proved that all pin classes have growth rates and established a procedure to determine them. We also have bounds on the possible growth rate of a pin class: for any pin sequence w ,

$$\kappa \leq \text{gr}(\mathcal{C}_w) \leq \omega_{\infty}$$

where $\kappa \approx 2.20557$ is the reciprocal of the smallest positive real root of

$$1 - 2z - z^3 = 0$$

and ω_∞ the smallest positive real root of

$$1 - 8z + 19z^2 - 26z^3 + 14z^4 - 12z^5 - 8z^6 + 20z^7 - 8z^8 = 0$$

A natural further question is what happens within these bounds: what are the possible growth rates of pin classes? We can also ask about bounds on growth rates of pin classes subject to certain characteristics: the number of quadrants visited (recurrently) by a pin class and the length of the longest oscillation contained in \mathcal{C}_w° are natural characteristics to consider. We can in fact state some answers in the former case already: the pin classes \mathcal{V} and \mathcal{Y} are in fact the smallest pin classes which visit two and three quadrants recurrently, respectively. It is also relatively easy to deduce *upper* bounds on the growth rates of pin classes in two and three quadrants by considering the **complete pin classes** in these bounded quadrants: for example, the complete class in two quadrants, $\mathcal{V}_c = \mathcal{V}_{w_c}$ is the pin class generated by a pin sequence w_c that contains *all* pin words in quadrants 1 and 2 as pin factors. Without too much difficulty (though we omit the proof here) we can calculate the growth rate of \mathcal{V}_c to be $\nu_c \approx 3.51205$, where ν_c is the reciprocal of the smallest positive real root of

$$1 - 2z - 4z^2 - 2z^3 - 8z^4 - 4z^5 = 0$$

In the sequel [6] we take up the question of what happens within this interval $[\nu, \nu_c]$: we move towards a classification of the growth rates of two-quadrant pin classes and observe some interesting structures in this set of growth rates. We show, for example, that there is a point $\nu_{\mathcal{L}} \approx 3.28277$ at which there are uncountably many distinct pin classes and that $\nu_{\mathcal{L}}$ is in fact an accumulation point in the set of pin class growth rates from both above and below. This has potential consequences for the study of well-quasi-ordered permutation classes because \mathcal{V} -classes (that is, two-quadrant pin classes) can be used to generate infinite antichains.

Potential further directions for study include:



- A systematic study of pin class growth rates in three and four quadrants;
- The possibility of conjecturing a classification of ‘small’ antichains (perhaps taking $\nu_{\mathcal{L}}$ as a cut-off) using \mathcal{V} -classes;
- The question of whether we can explicitly determine the generating function (not merely the growth rate) or *any* not-eventually-recurrent pin class, such as the Liouville \mathcal{V} , introduced in the sequel.

Appendix

In this appendix we prove Theorem 3.34, that the lists of collisions and \boxplus -decomposables given are in fact complete. We begin with collisions:

(A) Collisions Proof

We can verify that the list of collisions given in the table is complete for lengths $n \geq 5$ by an exhaustive search (which can be done fairly quickly on applying symmetries). We thus aim to prove that the table is complete at lengths $n \geq 6$. We repeat the relevant section of the table for reference:

List of collisions of pin factors:			
Length	Representative	Total number	Full List
$n \geq 6$ even:	 $1(ld)^k r = 2(dl)^k dru$	8 pairs	$\{1(ld)^k r, 2(dl)^k dru\}, \{1(dl)^k u, 4(ld)^k lur\},$ $\{2(dr)^k u, 3(rd)^k rul\}, \{2(rd)^k l, 1(dr)^k dlu\},$ $\{3(ru)^k l, 4(ur)^k uld\}, \{3(ur)^k d, 2(ru)^k rdl\},$ $\{4(ul)^k d, 1(lu)^k ldr\}, \{4(lu)^k r, 3(ul)^k urd\}$
$n \geq 7$ odd:	 $1(ld)^k lu = 2(dl)^k ur$	8 pairs	$\{1(ld)^k lu, 2(dl)^k ur\}, \{1(dl)^k dr, 4(ld)^k ru\},$ $\{2(dr)^k dl, 3(rd)^k lu\}, \{2(rd)^k ru, 1(dr)^k ul\},$ $\{3(ru)^k rd, 4(ur)^k dl\}, \{3(ur)^k ul, 2(ru)^k ld\},$ $\{4(ul)^k ur, 1(lu)^k rd\}, \{4(lu)^k ld, 3(ul)^k dr\}$

In order to prove this we shall first make the observation that in each of the pairs listed the two pin words end in different letters. We shall call a collision a *minimal collision* if all pin words in the tuple differ in their final letter. We shall first prove that the list above is a complete list of minimal collisions, and then deduce from this that there are no non-minimal collisions.

Theorem .1 (Classification of Minimal Collisions). *Any minimal collision of length $n \geq 6$ is one of the colliding pairs listed in the table above.*

Proof. In order to prove this we shall first apply symmetries to this list so that one pin word of each pair ends in dr:

We shall now proceed as follows: suppose we have a minimal collision of pin words w_1 and w_2 of length ≥ 6 and that w_1 ends in dr. Then the permutation π° is of the form given in Fig. 22. We shall use this to deduce facts about w_2 in order to show that this collision is in fact one of the pairs listed in Figs 20 and 21.

As in Fig 22, we shall call the lowest and second-lowest points of π° (A) and (B), respectively. These are generated by the final two letters dr of w_1 . We shall split into four cases based on which position the letter of w_2 which generates (A) is in: the final letter, the initial numeral, or an internal letter.




Figure 20: Odd collisions ending in dr of length ≥ 6




Figure 21: Even collisions ending in dr of length ≥ 6

(A).1 Case 1: (A) is the final point

We begin by deducing various facts about the pin word w_2 , using the fact that the permutation π looks like Fig. 23 (with all points other than (A) and (B) in the box), as well as the assumption that the final letter of w_2 corresponds to the point (A):

- By assumption, the final letter of w_2 corresponds to the point (A). The point corresponding to the final letter of a pin sequence must be the most extreme point in the direction indicated by that letter. Fig. 23 shows that (A) is the downmost point but not the most extreme point in any other direction (clearly, (B) is further up and to the right, and the fact that there must be at least one point in the box on each side of the dotted line implies that there is a point further to the left). Hence the final letter of w_2 *must* be a d .
- Consider the pin word w_2^- , formed by removing the final letter of w_2 . This must correspond to a permutation of the shape given in Fig. 24 (basically Fig. 23 without the point (A)). Note that the point (B) does not separate the bounding rectangle of all other points

$$w_1 = _ _ _ dr$$




Figure 22: The permutation π° generated by the pin word $w_1 = _ _ _ dr$ has this form. Note that (A) is in the lower half-plane and (B) is in the fourth quadrant. All points other than (A) and (B) (of which there are at least 4 by the assumption that the length of π° is ≥ 6) are in the box, with precisely one point on one side of the pin attached to (A) and all other points on the other.

$$w_2 = 4 _ _ _ d$$




Figure 23: The fact that the permutation π° can be generated by the pin word $w_1 = _ _ _ dr$ means that it has this form. Note that (A) is in the lower half-plane and (B) is in the fourth quadrant. All points other than (A) and (B) (of which there are at least 4 by the assumption that the length of π° is ≥ 6) are in the box, with precisely one point on one side of the dotted line attached to (A) and all other points on the other.

in the permutation. By the definition of a pin permutation, this can only happen if (B) was the first point placed: so (B) corresponds to the initial numeral of w_2 . But as (B) also corresponds to the final r in the pin word $w_1 = _ _ _ dr$ it must be in the fourth quadrant. Hence w_2 must begin with the numeral 4.

- Hence $w_2 = 4 _ _ _ d$, with the 4 corresponding to (B) and the final d corresponding to (A). Suppose that the blank space in the middle contained an r . Then this would correspond to a point to the right of all points placed before. But as (B) was the first point placed, this would imply the existence of a point to the right of (B). Fig. 23 shows that no such point exists, and so w_2 contains no r .
- Similarly, suppose that w_2 contained another d , in addition to the final one. Then this would correspond to a point below (B). But the only point below (B) is (A), already accounted for by the final d . Hence w_2 contains no d apart from its final letter.
- Combining these facts, we see that $w_2 = 4 _ _ _ d$, with the letters in the blank space alter-




Figure 24: The permutation generated by w_2^- has this form.

nating between u and l . Hence $w_2 = 4(ul)^{\geq 2}d$ or $w_2 = 4l(ul)^{\geq 2}d$, depending of whether the length of π° is even or odd, respectively. We thus now know what the permutation π looks like, as shown in Fig. 25.




Figure 25: Even and odd cases for the permutation π° , respectively

- We now return to $w_1 = _ _ dr$; as we now know what the permutation π looks like, we can deduce that the blank space here also does not contain a d or r : first, if the blank space contained a d then the points corresponding to both this and the next letter would be in the lower half-plane. But Fig. 25 shows that there is at most one point in the lower half-plane in addition to (A) and (B) (which are already accounted for by the final two letters). Similarly, if the blank space contained an r then the points corresponding to this and the next letter (neither of which can be the point (B) as this is accounted for by the final r) would be in the right half-plane. But Fig. 25 shows that there is at most one point in the right half-plane other than (B). Hence $w_1 = _ _ dr$ contains no d or r apart from the final two letters. This is now enough to deduce all of w_1 : $w_1 = 1(lu)^{\geq 1}ldr$ in the even case, and $w_1 = 3(ul)^{\geq 2}dr$ in the odd case.
- This means that there is only one (potential) collision of each length $n \geq 6$ in Case 1: $w_1 = 1(lu)^{\geq 1}ldr$ paired with $w_2 = 4(ul)^{\geq 2}d$ in the even case, and $w_1 = 3(ul)^{\geq 2}dr$ paired with $w_2 = 4l(ul)^{\geq 2}d$ in the odd case. These are the collisions on the right hand sides of Fig.s 20 and 21.

(A).2 Case 2: (A) is the first point

We now deal with the case in which (A) is the first point placed according to the pin word w_2 , corresponding to the initial numeral. We will again use the shape of the permutation π shown in Fig. 22 to deduce the form of w_2 .

- First note that, as (A) corresponds to the letter d in w_1 , (A) must be in the 3rd or 4th quadrant. Hence $w_2 = \{3/4\}____$
- Next, note that, as (B) is not the first point placed, it must correspond to a letter. If this letter were d or l then (B) would be below or to the left of (A) (as (A) has already been placed), which it is not (see Fig. 22). If the letter were u, then (B) would be the upmost of all points placed so far, and as Fig. 22 shows that all other points of π are above (B), this would mean that (B) would have to be the second point placed, so w_2 begins with either 3u or 4u: in the first case, both (A) and (B) would be in quadrant 3 and in the second case (B) would be to the left of (A) - Fig. 22 shows that neither of these is true, so (B) cannot correspond to a u in w_2 . Hence, by elimination, (B) corresponds to an r in w_2 .
- Hence (B) is at the end of a right-pin separating the previously placed point from the bounding rectangle of all other previously placed points and the origin. As the origin is above (B) (as we know (B) is in quadrant 4), the previously placed point must be the only point below (B), namely (A). Hence (B) corresponds to the first letter of w_2 after the numeral.
- Hence $w_2 = \{3/4\}r____$. We can now easily deduce that the blank space here contains no r or d: if it contained an r then this would correspond to a point to the right of (B) (as this has already been placed), and if it contained a d then this would correspond to a point below (A); but Fig. 22 clearly shows that no point of either type exists.
- Hence $w_2 = \{3/4\}rulul\dots$, and so the permutation π° looks like one of the permutations in Fig. 26.




Figure 26: The two possible cases for the permutation π° , as generated by the pin word $w_2 = \{3/4\}rulul\dots$. As $n \geq 6$ all the points shown must exist; any further points are a continuation of the oscillation in the second quadrant.

- We now return to $w_1 = ____dr$ which also generates π° , with the final two letters corresponding to (A) and (B). Note first that all the points in the upper half-plane of Fig. 26 have been placed before (A) in w_1 , and in either case there are at least three points to the

left of (A) which have been placed already: this means that the previous letter to the d corresponding to (A) must have been an r, and this must correspond to the point marked (C) (the only point other than (B) in the right half-plane). Thus $w_1 = _ _ _ r d r$, with the final three letters corresponding to (C), (A), (B), respectively. Note that the blank space here cannot contain a d, as (A) and (B) (already accounted for by the final two letters) are the only points in the lower half-plane. Thus $w_1 = _ _ _ u r d r$. But this implies that (C) is the second-upmost point, despite Fig. 26 showing that (C) has at least two points above it. This contradiction implies that there are in fact no collisions in Case 2.

(A).3 Case 3: (A) is an internal point

We now deal with the case where (A) is generated an internal letter of the pin word w_2 (ie., neither the initial numeral nor the final letter):

- First, we determine which letter represents (A) in w_2 . By looking at Fig. 22, we see that (A) is in the lower half-plane and is indeed the lowest point in the entire permutation. If (A) were represented by a u in w_2 it would be in the upper half-plane, so we can exclude this possibility. If (A) were represented by an l or r, on the other hand, it would have to follow either the numeral 3 or 4 or the letter d (as otherwise (A) would be in the upper half-plane); but in this case (A) would be above the previously-placed point, contradicting the fact that it is the lowest point in the entire permutation. Hence (A) must be represented by the letter d in w_2 .
- Next, note that the d representing (A) must in fact be the *final* d in w_2 : if there were a d after (A) it would place a point below (A), but (A) is the lowest point in the entire permutation. As (A) is internal to w_2 there is a letter immediately after the d corresponding to (A) in w_2 , either an l or an r. The point corresponding to this next letter, which we shall call (B'), will be the second-lowest point *so far* when it is placed. But since all points placed after (B') (if there are any) will be in the upper half-plane (as there is no further d in the word), (B') is in fact the second-lowest point in the *entire* permutation. Hence (B') must actually be (B), which must correspond to an r in w_2 as it is to the right of (A).
- We now know that the points (A) and (B) are generated by a consecutive dr in w_2 , and that there is no letter d in w_2 after this dr appears; by the same token there is also no further r after this dr, as this would generate a point to the right of (B). Hence, after the dr in w_2 corresponding to (A), (B), the pin word w_2 alternates between u and l for any remaining points. We now ask how many points are remaining in w_2 after this final dr.
- If there were *no* points after the dr in w_2 then w_1 and w_2 would both end in dr, so this would not be a pair of *minimal* collisions. Hence we can assume that there is at least one further letter after the final dr, which must be a u.
- Now suppose that there were at least two letters after the final dr; then w_2 would have the form $\dots drul\dots$, with any further letters at the end alternating between u and l. Thus π° would look like the permutation shown on the right-hand side of Fig. 27.




Figure 27: The permutation π° is (under the assumption that there are at least two letters after the r that generates (B) in w_2) generated by both the words $w_1 = \dots dr$ and $w_2 = \dots drul\dots$, so has both of these forms. The points marked (A) and (B) must match up.

- Hence the permutation π° is simultaneously of both forms represented in Fig. 27. On closer inspection, however, these forms contradict each other, which we can see by counting the points (other than (B)) to the left and right of (A): looking first at the diagram generated by w_1 , the box here is non-empty (as the length of π° is ≥ 6), and the pin attached to (A) must separate the previously-placed point from all others (including the origin). Hence, excluding (B), there is precisely one point on one side of (A) and all other points (including the origin) are on the other side. Looking at the diagram generated by w_2 , however, we see that the box here is also non-empty (as (A) is internal there is at least one non-origin point preceding it), with the previous point on one side and all other points (including the origin) on the other. Including the points marked (C) and (D), we now see that there are at least two points (including the origin but excluding (B)) on either side of (A), thus contradicting what the diagram generated by w_1 told us. We conclude that there cannot in fact be more than one point after the final dr in w_2 .
- Thus there is in fact precisely one point after the final dr , corresponding to (A), (B) in w_2 , and so $w_2 = \dots dru$. Hence our permutation π° is of both the forms shown in Fig. 28.




Figure 28: The permutation π° is generated by both the words $w_1 = \dots dr$ and $w_2 = \dots dru$, so has both of these forms. The points marked (A) and (B) must match up.

- But now consider the pin sequence w_1^{-1} , obtained by removing the final letter from w_1 . This must generate a pin permutation corresponding to the permutations shown in Fig.

28 with the point (B) removed; in particular, it contains the point (C) which (given the absence of (B)) will be both the highest and rightmost point in the corresponding pin permutation $\pi_{w_1}^\circ$. This can only happen if (C) was the *first* point placed in $\pi_{w_1}^\circ$ (otherwise it separates the previously placed point from all others in one direction, making it only the second-most point in that direction); as (C) is clearly in the first quadrant, this means that w_1 starts with a 1. Hence $w_1 = 1 \dots dr$. Further, given that there is no point above (C) in π° and the only point to the right is (B), accounted for by the final r , this means that there is no u in w_1 and no r except for the final letter. Hence w_1 is either $1l(dl)^{\geq 1}dr$ or $1(dl)^{\geq 2}dr$, depending on whether the length of π° is even or odd. Thus π° is one of the permutations on the left-hand sides of Figs 20 and 21, with w_1 being the upper word in the pair. Noting that $w_2 = \dots dru$ cannot contain an r or u apart from the final two letters (all points on the upper and right half-planes are already accounted for) allows us to conclude that w_2 must be the lower word in the pair.

□

We can now quickly deduce that *all* collisions are in fact minimal, and thus the list given above is complete:

Lemma .2. *Every colliding tuple $\{w_1, w_2, \dots, w_k\}$ of pin words is minimal (that is to say, $k \leq 4$ and each pair w_i, w_j differs in the final letter).*

Proof. Note that every collision of length $n \leq 5$ (which we have exhaustively listed) is minimal. Now suppose that $\{w_1, w_2\}$ is a non-minimal colliding pair of some length $n \geq 6$. By applying symmetries, we can assume that both w_1 and w_2 ends in a d . But then this d generates the lowest point in the shared generated pin permutation π° - in particular, it represents the same point in each pin word. Thus w_1^{-1} and w_2^{-1} (the pin words obtained by removing the final d from each word in the pair) must also form a collision, indeed a collision of shorter length. Given that w_1 and w_2 are not the same word, we can continue this process until we arrive at a minimal collision w_1^{-k}, w_2^{-k} . But we have already classified all minimal collisions, and can note that all minimal collisions are of different type: if one of the pair ends in d or u then the other must end in l or r , and therefore cannot be extended to a collision of longer length, forming a contradiction.

□

Thus all collisions are minimal and the list given in the table is complete.

(B) Classification of \boxplus -decomposables

We now aim to prove that the list of \boxplus -decomposables given in Theorem 3.34 is complete:

Proof. First, we list the \boxplus -decomposables of lengths 2 and 3: these are 1l and 1ld along with their eight symmetries, as can be confirmed by an exhaustive list.

We will now construct a \boxplus -decomposable pin permutation π° of length $n \geq 4$, generated by the pin word w , and show that it must be one of those listed in the statement of the theorem. We proceed by noting that every \boxplus -decomposable pin permutation can be drawn on the diagram in Fig. 29, with no points in the shaded region, and at least one point in each of the inner and outer regions.




Figure 29: The general shape of a \boxplus -decomposable permutation. The shaded region must be empty, whilst both the inner and outer unshaded regions must be non-empty.

By applying symmetries if necessary we can assume that the first point of π° placed was in the first quadrant. In fact, this point must be placed in the outer region of the first quadrant, as it is impossible to get out of the inner region once a point has been placed in there: if the point p_n has been placed in the inner region then the point p_{n+1} will be closer to the origin in the direction specified by the letter that placed p_n , and hence must also be in the inner region. Hence our permutation π° starts out like that in Fig. 30.




Figure 30: First point of π° is placed in the outer region of the first quadrant.

Next, note that it is impossible to get into the inner region of the first quadrant now: every

point placed in the first quadrant from now on will be either to the right of or above p_1 . Hence in order to construct a \boxplus -decomposable, we must at some point move to another quadrant. Again, by symmetry, we can assume that the next quadrant visited by the pin permutation π° is the second quadrant, and we now split into cases based on whether this first point in the second quadrant is in the inner or outer region:

(B).1 Case 1: First point in the second quadrant is in the outer region

In this case, we may (or may not) begin by oscillating in the first quadrant for any length: if there is an oscillation of length ≥ 2 this will ensure that the first point placed in the second quadrant will be in the outer region (it will be the second-highest point placed so far, and so will be above p_1 , the first point placed, which is in the outer region); if not, we shall simply make this assumption for now. We thus have a permutation that looks like that in Fig. 31.




Figure 31: The first point of π° outside the first quadrant can be assumed to be in the second quadrant by symmetry.

We note that the inner regions of the first, second and fourth quadrants (shaded in red in Fig. 31) are now inaccessible: there are now at least two points in the outer region of the upper half-plane, and because any new point in the upper half-plane must be either the highest or second-highest placed so far, it is now impossible to place any points below those two, hence the inner region of the upper half-plane will remain empty. A similar argument shows that the inner region of the right-half plane is now also inaccessible: if we had oscillated in the first quadrant initially then there are now at least two points to the right of the inner region in the right half-plane, and the argument goes through exactly as before. If we went directly to the second quadrant with our second point then there would be only one point to the right of the inner region so far, but in order to place anything in the right half-plane again we would need to take a right step, which would create a second, thus rendering the inner region of the right half-plane again inaccessible. In any case, the only part of the inner region which is now accessible is in the third quadrant, so we must be aiming to end up there.

If, after placing our first point in the second quadrant, we took either an upward step or a downward step into the outer region of the third quadrant, then the whole of the inner region

would be rendered inaccessible, by the same argument as in the previous paragraph. Hence we must now take a downstep into the inner region of the third quadrant, as in Fig. 32.




Figure 32: This is \boxplus -indecomposable, but will not be if any right or up step is taken.

Now note that we cannot now place any further points: either a left or right step here will place a point into the shaded region. Hence the only \boxplus -decomposable pin permutations in Case 1 are those shown in Fig. 32, which is to say those generated by a pin word of the form $1 \dots ld$, where the only letters in the ellipsis are u and r .

(B).2 Case 2: First point in the second quadrant is in the inner region

In this case, it is clear that we have to move into the second quadrant immediately after placing the first point in quadrant 1 (otherwise the first point in quadrant 2 will be above p_1 and hence not in the inner region). Hence we start off as in Fig. 33.




Figure 33: This is \boxplus -indecomposable, but will not be if any further point is added.

Note that we can now no longer take an up- or right-step: doing so would place a point above

or to the right of p_1 and therefore within the shaded region. Conversely, if we only take down- and left-steps we can stay in the inner region indefinitely, as shown in Fig. 34.




Figure 34: The general form of a \oplus -decomposable pin permutation in Case 2

Hence we see that we have obtained another family of \oplus -decomposable pin permutations, those generated by the pin words $1ldldl\dots$, and that these are exhaustive in Case 2.

Combining these two cases, we can now see that we have exhaustively obtained all possible \oplus -decomposable pin permutations that begin in the first quadrant and first visit the second. By applying all eight symmetries we thus obtain all \oplus -decomposable pin permutations and can see that these are precisely those described in the statement of the Theorem 3.34. \square

References

- [1] Richard Arratia. On the Stanley-Wilf conjecture for the number of permutations avoiding a given pattern. *Electron. J. Combin.*, 6:Note 1, 4 pp., 1999.
- [2] Frédérique Bassino, Mathilde Bouvel, and Dominique Rossin. Enumeration of pin-permutations. *Electron. J. Combin.*, 18(1):Paper 57, 39, 2011.
- [3] D. Bevan, R. Brignall, and N. Ruškuc. On cycles in monotone grid classes of permutations. Submitted.
- [4] David Bevan. Permutation patterns: basic definitions and notation. arXiv:1506.06673, 2015.
- [5] David Bevan. Intervals of permutation class growth rates. *Combinatorica*, 38(2):279–303, Apr 2018.
- [6] R. Brignall and B. Jarvis. Pin classes II: Small pin classes. Submitted.
- [7] R. Brignall and V. Vatter. Uncountably many enumerations of well-quasi-ordered permutation classes. Submitted.

- [8] Robert Brignall, Sophie Huczynska, and Vincent Vatter. Decomposing simple permutations, with enumerative consequences. *Combinatorica*, 28:385–400, 2008.
- [9] Robert Brignall, Nik Ruškuc, and Vincent Vatter. Simple permutations: decidability and unavoidable substructures. *Theoret. Comput. Sci.*, 391(1–2):150–163, 2008.
- [10] P. Cartier and D. Foata. *Problèmes combinatoires de commutation et réarrangements*. Lecture Notes in Mathematics, No. 85. Springer-Verlag, Berlin, 1969.
- [11] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University Press, Cambridge, 2009.
- [12] Adam Marcus and Gábor Tardos. Excluded permutation matrices and the Stanley-Wilf conjecture. *J. Combin. Theory Ser. A*, 107(1):153–160, 2004.
- [13] Maximilian M. Murphy. *Restricted permutations, antichains, atomic classes, and stack sorting*. PhD thesis, Univ. of St Andrews, 2002.
- [14] Vincent Vatter. Small permutation classes. *Proc. Lond. Math. Soc. (3)*, 103:879–921, 2011.
- [15] Steve Waton. *On Permutation Classes Defined by Token Passing Networks, Griding Matrices and Pictures: Three Flavours of Involvement*. PhD thesis, Univ. of St Andrews, 2007.