# On the Replica Symmetry of a Variant of the Sherrington-Kirkpatrick Spin Glass

Christian Brennecke<sup>\*</sup> Ad

Adrien Schertzer\*

December 10, 2024

#### Abstract

We consider N i.i.d. Ising spins with mean  $m \in (-1, 1)$  whose interactions are described by a Sherrington-Kirkpatrick Hamiltonian with a quartic correction. This model was recently introduced by Bolthausen in [3] as a toy model to understand whether a second moment argument can be used to derive the replica symmetric formula in the full high temperature regime if  $m \neq 0$ . In [3], Bolthausen suggested that a natural analogue of the de Almeida-Thouless condition for the toy model is

$$\beta^2 (1 - m^2)^2 \le 1. \tag{1}$$

Here,  $\beta \geq 0$  corresponds to the inverse temperature. While the second moment method implies replica symmetry for  $\beta$  sufficiently small, Bolthausen showed that the method fails to prove replica symmetry in the full region described by (1). A natural question that was left open in [3] is whether (1) correctly characterizes the high temperature phase of the toy model. In this note, we show that this is indeed not the case. We prove that if  $|m| \geq m_*$ , for some  $m_* \in (0, 1)$ , the limiting free energy of the toy model is negative for suitable  $\beta$  that satisfy (1).

### 1 Introduction and Main Result

For  $\beta \geq 0$  and  $m \in (-1, 1)$ , consider the mean field spin glass with partition function

$$Z_{N,\beta,m} := \sum_{\sigma \in \{-1,1\}^N} p_m(\sigma) \exp\left(\sum_{1 \le i,j \le N} \frac{\beta}{\sqrt{2}} g_{ij} \widehat{\sigma}_i \widehat{\sigma}_j - \frac{\beta^2}{4N} \sum_{1 \le i,j \le N} \widehat{\sigma}_i^2 \widehat{\sigma}_j^2\right).$$
(2)

Here,  $\sigma = (\sigma_1, \ldots, \sigma_N) \in \{-1, 1\}^N$ ,  $\widehat{\sigma}_i := \sigma_i - m$ ,  $p_m(\sigma) := \prod_{i=1}^N \frac{1}{2}(1 + m\sigma_i)$  and the couplings  $(g_{ij})_{1 \le i,j \le N}$  are i.i.d. Gaussian random variables with distribution  $\mathcal{N}(0, N^{-1})$ .

The model in (2) was recently introduced by Bolthausen in [3]. The main result of [3] is a new proof of the replica symmetric (RS) formula for the classical Sherrington-Kirkpatrick (SK) model [13] with external field strength  $h \in \mathbb{R}$  and for sufficiently small  $\beta \geq 0$ . The proof is based on a second moment argument, conditionally on an

<sup>\*</sup>Institute for Applied Mathematics, University of Bonn, Endenicher Allee 60, 53115 Bonn, Germany

approximate solution of the TAP equations [18] that was constructed in [2]. The RS formula is expected to be true under the de Almeida-Thouless [8] condition

$$\beta^2 E(1 - m_{\beta,h}^2)^2 \le 1, \text{ for } m_{\beta,h} = \tanh(h + \beta\sqrt{q}Z), q = E \tanh^2(h + \beta\sqrt{q}Z),$$
(3)

where Z is a standard Gaussian random variable. To date, this was verified in a very large subregion of the phase diagram in [19, 16, 17, 10], based on the analysis of the Parisi formula [11, 12] proved in [9, 15]. A complete proof that the RS phase is fully characterized by (3) still remains open, however. Note for a centered Gaussian external field, the RS phase indeed turns out to be characterized by an analogue of (3), see [6].

It is a natural question whether a second moment argument as in [3] can be improved to verify the replica symmetry in the expected region (3). Although the temperature range from [3] can be improved through basic restriction arguments [5], Bolthausen already suggested in [3] that understanding the full RS phase likely requires additional techniques. His argument is based on the analysis of the toy model defined in (2). For this model, the second moment method can be evaluated explicitly and it is straightforward to show that for  $\beta \geq 0$  small enough, one has that

$$\lim_{N \to \infty} \frac{1}{N} \log \mathbb{E} Z_{N,\beta,m} = \lim_{N \to \infty} \frac{1}{N} \log \mathbb{E} Z_{N,\beta,m}^2 = 0.$$
(4)

By Gaussian concentration, see e.g. [16, Th. 1.3.4.], this implies for small  $\beta$  that

$$\lim_{N \to \infty} \mathbb{E} \frac{1}{N} \log Z_{N,\beta,m} = 0.$$
(5)

By analogy to (3), Bolthausen then suggested that the high temperature region

$$\operatorname{HT}_{m} := \left\{ \beta \ge 0 : \lim_{N \to \infty} \mathbb{E} \frac{1}{N} \log Z_{N,\beta,m} = 0 \right\}$$
(6)

the set in which the quenched equals the annealed free energy, should correspond to

$$AT_m := \left\{ \beta \ge 0 : \beta^2 (1 - m^2)^2 \le 1 \right\}$$
(7)

and it was left open whether (5) indeed holds true for all  $\beta \in AT_m$  (cf. [3, Eq. (6.1)]).

Note that one might expect that  $\operatorname{AT}_m \subset \operatorname{HT}_m$  based on different arguments. For example, an application of the TAP heuristics suggests that  $\mathbf{m} = (m, m, \dots, m) \in$  $(-1, 1)^N$  solves an analogue of the TAP equations associated with (2). Alternatively, it is not difficult to see that, for  $\beta$  sufficiently small, the free energy fluctuations converge to a normal variable with a variance that contains the term  $\log (1 - \beta^2 (1 - m^2)^2)$ . This indicates a phase transition when  $\beta^2 (1 - m^2)^2 \to 1$ , similarly as in the SK model without external field analyzed in detail in [1] (in which case the corresponding divergence at  $\beta \to 1$  detects the RS-RSB transition temperature correctly).

It turns out<sup>1</sup> that a precise characterization of  $HT_m$  may be obtained by adapting the methods from [7] to the present setting. Related numerical simulations suggest that

<sup>&</sup>lt;sup>1</sup>We thank W.-K. Chen for this remark and for pointing out [7] to us. We thank W.-K. Chen and S. Tang for sharing some related numerical simulations on (6) and (7) with us.

for large |m|, one can find  $\beta \in AT_m$  with  $\beta \notin HT_m$ . Perhaps surprisingly, this fact admits an elementary proof which does neither require precise knowledge of  $HT_m$  nor of a Parisi formula for the limiting free energy of the model.

**Theorem 1.** Let  $I(m) := -\frac{1+m}{2} \log \frac{1+m}{2} - \frac{1-m}{2} \log \frac{1+m}{2}$  and define  $m_* > 0$  to be such that  $I(m_*) = \frac{1}{4}$ . Then, for every  $|m| > m_*$ , there exists  $\beta \in AT_m$  with  $\beta \notin HT_m$ .

## 2 Proof of Theorem 1

Our proof is elementary and adapts an argument from the analysis of the random energy model, see [4, p. 167], to the present setting. By symmetry, we assume in the rest of this section without loss of generality that m > 0.

*Proof of Theorem 1.* In the following, denote by  $E_m$  the expectation with regards to the base measure  $p_m$  and, for  $\epsilon > 0$ , define the set

$$S_{N,m,\epsilon} := \Big\{ \sigma \in \{-1,1\}^N : \frac{1}{N} \sum_{i=1}^N \sigma_i \in [m-\epsilon, m+\epsilon] \Big\}.$$

Moreover, denote by  $H_{N,\beta,m}$  the Hamiltonian

$$H_{N,\beta,m}(\sigma) := \sum_{1 \le i,j \le N} \frac{\beta}{\sqrt{2}} g_{ij} \widehat{\sigma}_i \widehat{\sigma}_j - \frac{\beta^2}{4N} \sum_{1 \le i,j \le N} \widehat{\sigma}_i^2 \widehat{\sigma}_j^2$$

so that  $Z_{N,\beta,m} = E_m \exp(H_{N,\beta,m})$ .

We first claim that for every  $\beta \in \operatorname{HT}_m$  and every  $\epsilon > 0$ , we have that

$$\lim_{N \to \infty} \left| \frac{1}{N} \log(Z_{N,\beta,m}) - \frac{1}{N} \log E_m \left( \mathbf{1}_{S_{N,m,\epsilon}} \exp(H_{N,\beta,m}) \right) \right| = 0$$
(8)

in the sense of probability. Here,  $\mathbf{1}_S : \{-1, 1\}^N \to \{0, 1\}$  denotes the characteristic function of the subset  $S \subset \{-1, 1\}^N$ . To prove (8), set

$$Z_{N,\beta,m}^{\leq\epsilon} := E_m(\mathbf{1}_{S_{N,m,\epsilon}} \exp H_{N,\beta,m}), \ Z_{N,\beta,m}^{>\epsilon} := E_m(\mathbf{1}_{S_{N,m,\epsilon}^{\mathsf{C}}} \exp H_{N,\beta,m})$$

so that  $Z_{N,\beta,m} = Z_{N,\beta,m}^{\leq \epsilon} + Z_{N,\beta,m}^{>\epsilon}$ . Then, for every  $\delta > 0$ , we have that

$$\mathbb{P}\left(\left|\frac{1}{N}\log(Z_{N,\beta,m}) - \frac{1}{N}\log\left(Z_{N,\beta,m}^{\leq\epsilon}\right)\right| \geq \delta\right) \\
= \mathbb{P}\left(Z_{N,\beta,m}^{\geq\epsilon}/Z_{N,\beta,m}^{\leq\epsilon} \geq e^{\delta N} - 1, \ Z_{N,\beta,m}^{\leq\epsilon} \leq e^{-\delta N}\right) \\
+ \mathbb{P}\left(Z_{N,\beta,m}^{\geq\epsilon}/Z_{N,\beta,m}^{\leq\epsilon} \geq e^{\delta N} - 1, \ Z_{N,\beta,m}^{\leq\epsilon} > e^{-\delta N}\right) \\
\leq \mathbb{P}\left(Z_{N,\beta,m}^{\leq\epsilon} \leq e^{-\delta N}\right) + \mathbb{P}\left(Z_{N,\beta,m}^{\geq\epsilon} \geq 1 - e^{-\delta N}\right) \\
\leq \mathbb{P}\left(Z_{N,\beta,m}^{\leq\epsilon} \leq e^{-\delta N}\right) + \left(1 - e^{-\delta N}\right)^{-1} p_m\left(S_{N,m,\epsilon}^{\complement}\right),$$
(9)

where in the last step, we applied Markov's inequality. Similarly, we have that

$$\mathbb{P}\left(Z_{N,\beta,m}^{\leq\epsilon} \leq e^{-\delta N}\right)$$
  
$$\leq \mathbb{P}\left(Z_{N,\beta,m}^{\leq\epsilon} \leq e^{-\delta N}, \ Z_{N,\beta,m}^{>\epsilon} \leq e^{-\delta N}\right) + \mathbb{P}\left(Z_{N,\beta,m}^{>\epsilon} > e^{-\delta N}\right)$$
  
$$\leq \mathbb{P}\left(\frac{1}{N}\log Z_{N,\beta,m} \leq -\frac{\delta}{2}\right) + e^{\delta N}p_m\left(S_{N,m,\epsilon}^{\complement}\right)$$

for N large enough so that

,

$$\mathbb{P}\left(\left|\frac{1}{N}\log(Z_{N,\beta,m}) - \frac{1}{N}\log\left(Z_{N,\beta,m}^{\leq\epsilon}\right)\right| \geq \delta\right)$$
$$\leq \mathbb{P}\left(\left|\frac{1}{N}\log Z_{N,\beta,m}\right| \geq \frac{\delta}{2}\right) + \frac{e^{\delta N}}{1 - e^{-\delta N}}p_m\left(S_{N,m,\epsilon}^{\complement}\right).$$

Markov's inequality and Gaussian concentration (see [16, Th. 1.3.4.]) imply that

$$\mathbb{P}\left(\left|\frac{1}{N}\log Z_{N,\beta,m}\right| \geq \frac{\delta}{2}\right) \leq \mathbb{P}\left(\left|\frac{1}{N}\log Z_{N,\beta,m} - \mathbb{E}\frac{1}{N}\log Z_{N,\beta,m}\right| \geq \frac{\delta}{4}\right) + \frac{4}{\delta}\left|\mathbb{E}\frac{1}{N}\log Z_{N,\beta,m}\right| \\ \leq Ce^{-cN\delta^2/\beta^2} + \frac{4}{\delta}\left|\mathbb{E}\frac{1}{N}\log Z_{N,\beta,m}\right|$$

for suitable constants C, c > 0 (independent of all parameters). On the other hand, setting  $h := \tanh^{-1}(m)$ , a standard exponential moment bound shows that

$$p_m(S_{N,m,\epsilon}^{\complement}) \leq \inf_{\lambda \geq 0} \exp\left(-N\epsilon\lambda - Nm\lambda + N(\log\cosh(h+\lambda) - \log\cosh(h))\right) \\ + \inf_{\lambda \geq 0} \exp\left(-N\epsilon\lambda + Nm\lambda + N(\log\cosh(h-\lambda) - \log\cosh(h))\right) \\ \leq 2\inf_{\lambda \geq 0} \exp\left(-N\epsilon\lambda + N\lambda^2/2\right) = 2\exp(-N\epsilon^2/2).$$

Now, for every  $\delta' > 0$ , we can find  $0 < \delta < \min(\delta', \epsilon^2/4)$  so that we arrive at

$$\begin{split} &\limsup_{N \to \infty} \mathbb{P}\Big(\Big|\frac{1}{N}\log(Z_{N,\beta,m}) - \frac{1}{N}\log\left(Z_{N,\beta,m}^{\leq \epsilon}\right)\Big| \ge \delta'\Big) \\ &\le \limsup_{N \to \infty} \mathbb{P}\Big(\Big|\frac{1}{N}\log(Z_{N,\beta,m}) - \frac{1}{N}\log\left(Z_{N,\beta,m}^{\leq \epsilon}\right)\Big| \ge \delta\Big) \\ &\le \limsup_{N \to \infty} \left(Ce^{-cN\delta^2/\beta^2} + Ce^{-N\epsilon^2/4} + \frac{4}{\delta}\Big|\mathbb{E}\frac{1}{N}\log Z_{N,\beta,m}\Big|\right) = \frac{4}{\delta}\limsup_{N \to \infty}\Big|\mathbb{E}\frac{1}{N}\log Z_{N,\beta,m}\Big|. \end{split}$$

By definition of  $HT_m$  in (6), the r.h.s. vanishes if  $\beta \in HT_m$  and this concludes (8).

Notice that (8) implies that for every  $\beta \in \operatorname{HT}_m$  and every  $\epsilon > 0$ , we have that

$$\lim_{N \to \infty} \frac{1}{N} \log Z_{N,\beta,m}^{\leq \epsilon} = \lim_{N \to \infty} \mathbb{E} \frac{1}{N} \log Z_{N,\beta,m}^{\leq \epsilon} = 0,$$
(10)

where the first equality holds true almost surely. Indeed, this follows from (8) combined with Gaussian concentration applied to both  $Z_{N,\beta,m}$  and  $Z_{N,\beta,m}^{\leq\epsilon}$ .

We now argue that for every  $m > m_*$ , we can find  $\beta \in AT_m$  such that for all  $\epsilon > 0$  sufficiently small, it holds true that

$$\limsup_{N \to \infty} \mathbb{E} \frac{1}{N} \log Z_{N,\beta,m}^{\leq \epsilon} < 0.$$
(11)

By (10), this implies  $\beta \notin \operatorname{HT}_m$  and it concludes that  $\operatorname{AT}_m \not\subset \operatorname{HT}_m$  whenever  $m > m_*$ . In the remainder, we abbreviate

$$F_{N,\beta,m}^{\leq \epsilon} := \mathbb{E}\frac{1}{N} \log Z_{N,\beta,m}^{\leq \epsilon}.$$

Then, we have that

$$\begin{split} \frac{d}{d\beta} F_{N,\beta,m}^{\leq\epsilon} &= \mathbb{E} \, E_m \bigg( \frac{1}{N} \sum_{1 \leq i,j \leq N} \frac{g_{ij}}{\sqrt{2}} \widehat{\sigma}_i \widehat{\sigma}_j - \frac{\beta}{2} \Big( \frac{1}{N} \sum_{i=1}^N \widehat{\sigma}_i^2 \Big)^2 \Big) \mathbf{1}_{S_{N,m,\epsilon}} \frac{e^{H_{N,\beta,m}}}{Z_{N,\beta,m}^{\leq\epsilon}} \\ &= \mathbb{E} \, E_m \, \frac{1}{N} \sum_{1 \leq i,j \leq N} \frac{g_{ij}}{\sqrt{2}} \widehat{\sigma}_i \widehat{\sigma}_j \, \mathbf{1}_{S_{N,m,\epsilon}} \frac{e^{H_{N,\beta,m}}}{Z_{N,\beta,m}^{\leq\epsilon}} - \frac{\beta}{2} (1-m^2)^2 + C\beta\epsilon \\ &\leq \frac{\mathbb{E} M_{N,m,\epsilon}}{\sqrt{N}} - \frac{\beta}{2} (1-m^2)^2 + C\beta\epsilon. \end{split}$$

Here, C > 0 is some universal constant and we defined

$$M_{N,m,\epsilon} := \max \left\{ X_{\sigma} : \sigma \in S_{N,m,\epsilon} \right\} \quad \text{for} \quad X_{\sigma} := \frac{1}{\sqrt{2N}} \sum_{1 \le i,j \le N} g_{ij} \widehat{\sigma}_i \widehat{\sigma}_j.$$

To control the maximum  $M_{N,m,\epsilon}$ , we apply Jensen's inequality so that for t > 0, we get

$$\frac{\mathbb{E}M_{N,m,\epsilon}}{\sqrt{N}} \leq \frac{1}{Nt} \mathbb{E}\log \exp\left(\sqrt{N}tM_{N,m,\epsilon}\right) \\
\leq \frac{1}{Nt}\log \mathbb{E}\exp\left(\sqrt{N}tM_{N,m,\epsilon}\right) \\
\leq \frac{1}{Nt}\log \mathbb{E}\sum_{\sigma \in S_{N,m,\epsilon}} \exp\left(\sqrt{N}tX_{\sigma}\right) = \frac{1}{Nt}\log \sum_{\sigma \in S_{N,m,\epsilon}} \exp\left(\frac{Nt^{2}}{2}\mathbb{E}X_{\sigma}^{2}\right).$$

Now, recall that for every  $\sigma \in S_{N,m,\epsilon}$ , we have that

$$\mathbb{E}X_{\sigma}^{2} = \frac{1}{2}(1+m^{2}-2ms_{\sigma})^{2} \text{ for some } s_{\sigma} \in \Delta_{N} \cap [m-\epsilon,m+\epsilon],$$

where  $\Delta_N := \{ -1 + 2k/N : k = 0, 1, ..., N \}$ . Moreover, by Stirling's approximation, we have for every  $s \in \Delta_N \cap [m - \epsilon, m + \epsilon]$  (and  $\epsilon > 0$  small enough) that

$$\left|\left\{\sigma \in \{-1,1\}^N : s_{\sigma} = s\right\}\right| = \binom{N}{\frac{1+s}{2}N} \le C \exp\left(NI(s)\right)$$

for N large enough, a universal constant C > 0 and where we recall that  $I : [-1, 1] \rightarrow [0, 1]$  denotes the binary entropy function, defined by

$$I(s) = -\frac{1+s}{2}\log\frac{1+s}{2} - \frac{1-s}{2}\log\frac{1-s}{2}$$

Recall that I is a strictly concave function, that it attains its unique maximum at s = 0, vanishes at  $s = \pm 1$  and that it is strictly decreasing in [0, 1]. Using that  $|\Delta_N \cap [m - \epsilon, m + \epsilon]| \leq N$ , we obtain for every t > 0 the upper bound

$$\frac{\mathbb{E}M_{N,m,\epsilon}}{\sqrt{N}} \le \frac{1}{t} \frac{\log(CN)}{N} + \frac{1}{t} \max_{s \in [m-\epsilon, m+\epsilon]} \Big( \frac{t^2}{4} (1+m^2-2ms)^2 + I(s) \Big).$$

Choosing  $t := 2(1-m^2)^{-1}\sqrt{I(m)}$  s.t. in particular t > 0 for  $m \in (0,1)$ , this implies

$$\frac{\mathbb{E}M_{N,m,\epsilon}}{\sqrt{N}} \leq \frac{(1-m^2)}{2\sqrt{I(m)}} \frac{\log(CN)}{N} \\ + \frac{(1-m^2)}{2\sqrt{I(m)}} \max_{s \in [m-\epsilon,m+\epsilon]} \left( I(m)(1-m^2)^{-2}(1+m^2-2ms)^2 + I(s) \right) \\ \leq \frac{(1-m^2)}{2\sqrt{I(m)}} \frac{\log(CN)}{N} + (1-m^2)\sqrt{I(m)} + C\epsilon$$

for some C > 0 (that is independent of  $\beta$ ) and for all  $\epsilon > 0$  small enough, by continuity.

Collecting the previous estimates, we conclude that for every  $m \in (0, 1)$ , every  $\beta \ge 0$ and for every sufficiently small  $\epsilon > 0$  it holds true that

$$\frac{d}{d\beta} F_{N,\beta,m}^{\leq \epsilon} \leq (1-m^2)\sqrt{I(m)} - \frac{\beta}{2}(1-m^2)^2 + C(1+\beta)\epsilon + O(\log N/N).$$

Integrating the last inequality in  $\beta \geq 0$  and combining this with  $F_{N,\beta,m}^{\leq \epsilon} \leq 0$  for all  $\beta \geq 0$ , which follows from Jensen's inequality, we obtain the upper bound

$$\limsup_{\epsilon \to 0} \limsup_{N \to \infty} F_{N,\beta,m}^{\leq \epsilon} \leq \inf_{\beta_0 \geq 0} \begin{cases} 0 & :\beta < \beta_0, \\ (\beta - \beta_0)(1 - m^2)\sqrt{I(m)} - \frac{\beta^2 - \beta_0^2}{4}(1 - m^2)^2 & :\beta \geq \beta_0. \end{cases}$$

Choosing  $\beta_c = 2(1-m^2)^{-1}\sqrt{I(m)} > 0$  to bound the r.h.s. from above, we arrive at

$$\limsup_{\epsilon \to 0} \limsup_{N \to \infty} F_{N,\beta,m}^{\leq \epsilon} \leq \begin{cases} 0 & : \beta < \beta_c, \\ -\frac{1}{4}(\beta - \beta_c)^2(1 - m^2)^2 & : \beta \geq \beta_c. \end{cases}$$
(12)

To conclude the proof, we finally notice that (12) implies that

$$\limsup_{\epsilon \to 0} \limsup_{N \to \infty} F_{N,\beta,m}^{\leq \epsilon} < 0$$

whenever

$$\beta > \beta_c \iff \beta^2 (1 - m^2)^2 > 4I(m).$$
 (13)

If  $m > m_*$ , for  $m_* > 0$  defined by  $I(m_*) = \frac{1}{4}$ , the monotonicity of I implies that 4I(m) < 1 so that, in particular, there exists some  $\beta \in AT_m$  that satisfies

$$4I(m) < \beta^2 (1-m^2)^2 \le 1 \iff \beta \in \left( (1-m^2)^{-1} \sqrt{4I(m)}, (1-m^2)^{-1} \right].$$

Choosing  $\epsilon > 0$  small enough, this implies (11) and concludes that  $AT_m \not\subset HT_m$ .

Acknowledgements. C. B. thanks G. Genovese for helpful discussions. C. B. and A. S. acknowledge support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – GZ 2047/1, Projekt-ID 390685813. A.S. is also funded by the Deutsche Forschungsgemeinschaft – project number 432176920.

#### References

- M. Aizenman, J. L. Lebowitz, D. Ruelle. Some rigorous results on the Sherrington-Kirkpatrick spin glass model. *Comm. Math. Phys.* 11, pp. 3-20 (1987).
- [2] E. Bolthausen. An Iterative Construction of Solutions of the TAP Equations for the Sherrington-Kirkpatrick Model. Comm. Math. Phys. 325, pp. 333-366 (2014).
- [3] E. Bolthausen. A Morita Type Proof of the replica-symmetric Formula for SK. In: Statistical Mechanics of Classical and Disordered Systems, Springer Proceedings in Mathematics & Statistics, pp. 63-93 (2018).
- [4] A. Bovier. Statistical Mechanics of Disordered Systems: A Mathematical Perspective. Cambridge Series in Statistical and Probabilistic Mathematics (2012).
- [5] C. Brennecke, H.-T. Yau. The Replica Symmetric Formula for the SK Model Revisited. J. Math. Phys. 63, 2022.
- [6] W.-K. Chen. On the Almeida-Thouless transition line in the SK model with centered Gaussian external field. *Electron. Commun. Probab.* 26, pp. 1-9 (2021).
- [7] W.-K. Chen, M. Handschy, G. Lerman. Phase transition in random tensors with multiple spikes. Ann. App. Probab. 31, No. 4, 1868-1913 (2021).
- [8] J. R. L. de Almeida, D. J. Thouless. Stability of the Sherrington-Kirkpatrick solution of a spin glass model. J. Phys. A: Math. Gen. 11, pp. 983-990 (1978).
- [9] F. Guerra. Broken Replica Symmetry Bounds in the Mean Field Spin Glass Model. Comm. Math. Phys. 233, pp. 1-12 (2003).
- [10] A. Jagannath, I. Tobasco. Some properties of the phase diagram for mixed p-spin glasses. Probab. Theory Relat. Fields 167, pp. 615-672 (2017).
- G. Parisi. Infinite number of order parameters for spin-glasses. *Phys. Rev. Lett.* 43, pp. 1754-1756 (1979).

- [12] G. Parisi. A sequence of approximate solutions to the S-K model for spin glasses. J. Phys. A 13, pp. L-115 (1980).
- [13] D. Sherrington, S. Kirkpatrick. Solvable model of a spin glass. Phys. Rev. Lett. 35, pp. 1792-1796 (1975).
- [14] D. Slepian. The one-sided barrier problem for Gaussian noise. Bell System Tech. J., 41, 463–501 (1962).
- [15] M. Talagrand. The Parisi formula. Ann. of Math. 163, no. 1, pp. 221-263 (2006).
- [16] M. Talagrand. Mean Field Models for Spin Glasses. Volume I: Basic Examples. A Series of Modern Surveys in Mathematics 54, Springer Berlin, Heidelberg (2010).
- [17] M. Talagrand. Mean Field Models for Spin Glasses. Volume II: Advanced Replica-Symmetry and Low Temperature. A Series of Modern Surveys in Mathematics 54, Springer Verlag Berlin-Heidelberg (2011).
- [18] D. J. Thouless, P. W. Anderson, R. G. Palmer. Solution of 'solvable model in spin glasses'. *Philos. Magazin* 35, pp. 593-601 (1977).
- [19] F. L. Toninelli. About the Almeida-Thouless transition line in the Sherrington-Kirkpatrick mean-field spin glass model. *Europhys. Lett.* 60 (5), p. 764 (2002).