

# A Proof of Exact Convergence Rate of Gradient Descent. Part II. Performance Criterion $(f(x_N) - f_*)/\|x_0 - x_*\|^2$

Jungbin Kim

## Abstract

We prove the exact worst-case convergence rate of gradient descent for smooth strongly convex optimization, with respect to the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$ . This rate was previously conjectured in [DT14; THG17].

## 1 Introduction

Consider the optimization problem

$$x_* \in \arg \min_{x \in \mathbb{R}^d} f(x), \quad (1)$$

where  $f$  is in  $\mathcal{F}_{\mu,L}$ , the space of  $L$ -smooth and  $\mu$ -strongly convex functions on  $\mathbb{R}^d$ . Consider the gradient descent method

$$x_{k+1} = x_k - \gamma \nabla f(x_k) \quad (\text{GD})$$

with a constant stepsize  $\gamma \in (0, 2/L)$ . Given the number  $N$  of iterations, the *convergence rate*, or *worst-case performance*, of GD is defined as follows:

$$\sup \left\{ \frac{f(x_N) - f(x_*)}{\|x_0 - x_*\|^2} : f \in \mathcal{F}_{\mu,L}, x_0 \in \mathbb{R}^d, x_k \text{ generated by GD} \right\}, \quad (2)$$

where the performance criterion is set as  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$ . Based on numerical evidence, The exact value of (2) was conjectured by Drori and Teboulle when  $\mu = 0$ , and by Taylor *et al.* when  $\mu > 0$ , as stated below.

**Theorem 1** ([DT14, Conj. 1; THG17, Conj. 2]). *The iterates of GD satisfy*

$$f(x_N) - f(x_*) \leq \max \left\{ \frac{\kappa}{(\kappa - 1) + (1 - \gamma\mu)^{-2N}}, (1 - \gamma L)^{2N} \right\} \frac{L}{2} \|x_0 - x_*\|^2, \quad (3)$$

where  $\kappa = \mu/L$ . Moreover, this convergence guarantee is tight.

Note that the statement of Theorem 1 consists of two parts: the first part claims an upper bound of (2), while the second part claims a lower bound of (2). To the best of our knowledge, the upper bound has only been proved for the case where  $\mu = 0$  and  $\gamma \in (0, 1/L]$  [DT14, Thm. 2; TV23, Thm. 2], while the lower bound was completely proved in [THG17, §4.1.2] by providing a simple worst-case example.

In contrast, the exact convergence rate of GD for another performance criterion  $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$  was recently proved by Rotaru *et al.*, as stated below. Another proof can be found in Part I of our paper.

**Theorem 2** ([RGP24, Thm. 2.2, Prop. 5.2; Kim24, Thm. 1]). *The iterates of GD satisfy*

$$\frac{1}{2L} \|\nabla f(x_N)\|^2 \leq \max \left\{ \frac{\kappa}{(\kappa - 1) + (1 - \gamma\mu)^{-2N}}, (1 - \gamma L)^{2N} \right\} (f(x_0) - f(x_*)), \quad (4)$$

where  $\kappa = \mu/L$ . Moreover, this convergence guarantee is tight.

In this work, we prove Theorem 1 by showing the upper bound (3) for any strong convexity parameter  $\mu \in [0, L)$  and any stepsize  $\gamma \in (0, 2/L)$ . Our proof strategy is based on the following phenomenon: some convergence analyses for the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$  have *dual* analyses for the performance criterion  $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$ . See, for example, [KY24, Thm. 3; Kim+24, Thms. 1, 2] for a recent theory on this phenomenon. In this correspondence, proving Theorem 2 will mirror the task of proving Theorem 1. In comparing our proofs of Theorem 1 and 2, the proof is significantly simpler for the mirrored problem than for the original problem: only  $O(N)$  inequalities are needed to prove Theorem 2, whereas  $O(N^2)$  number of inequalities are needed to prove Theorem 1. Throughout the paper, we assume that the readers are familiar with the background materials and notations in [Kim24, §1.1, 2.1].

## 2 Anti-transpose of first-order method

For an  $N \times N$  lower-triangular matrix  $\mathbf{H}$ , its anti-transpose  $\mathbf{H}^A$  is obtained by reflecting  $\mathbf{H}$  over its anti-diagonal. The first-order algorithm associated with  $\mathbf{H}^A$  can be written as follows:

$$\begin{bmatrix} x_1 - x_0 \\ \vdots \\ x_N - x_{N-1} \end{bmatrix} = -\frac{1}{L} \begin{bmatrix} \mathbf{H}_{N,N} & & \\ & \ddots & \\ \mathbf{H}_{N,1} & \cdots & \mathbf{H}_{1,1} \end{bmatrix} \begin{bmatrix} \nabla f(x_0) \\ \vdots \\ \nabla f(x_{N-1}) \end{bmatrix} \quad (5)$$

Perhaps, the observation of the anti-transpose relationship between first-order methods dates back to [KF21, §6.6]. The anti-transpose can also be defined using the language of operators as follows.

**Definition 1.** *Let  $H$  be an operator. Then, its anti-transpose is the operator  $H^A$  that satisfies*

$$\langle Hx, Ty \rangle = \langle Tx, H^A y \rangle,$$

for all  $x$  and  $y$ . Equivalently,  $H^A = TH^T T$ . Here,  $T$  is the time reversal operator on  $(\mathbb{R}^d)^N$ :

$$T \begin{bmatrix} x_1 \\ \vdots \\ x_N \end{bmatrix} = \begin{bmatrix} x_N \\ \vdots \\ x_1 \end{bmatrix}.$$

From the identity  $(H^T)^{-1} = (H^{-1})^T$ , it follows that

$$(H^A)^{-1} = (TH^T T)^{-1} = (T)^{-1} (H^T)^{-1} (T)^{-1} = T (H^{-1})^T T = (H^{-1})^A.$$

This justifies denoting this operator as  $H^{-A}$ . We also note that  $\widetilde{\mathbf{H}}^A = \tilde{\mathbf{H}}^A$ , where the tilde indicates that the  $H$ -matrix generates the iterates  $x_k^+$  instead of  $x_k$ .

Performance criterion	$\frac{f(x_N) - f(x_*)}{\ x_0 - x_*\ ^2}$	$\frac{\ \nabla f(y_N)\ ^2}{f(y_0) - f(y_*)}$
Algorithm	$\mathbf{H}$	$\mathbf{H}^A$
Iterates	$x_k$	$y_k$
Multipliers	$\lambda_{i,j}$	$\nu_{i,j}$

Table 1: Notational differences in Sections 3.1 and 3.2.

### 3 Equivalence between convergence analyses for performance criteria $(f(x_N) - f_*)/\|x_0 - x_*\|^2$ and $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$

In Sections 3.1 and 3.2, we present two PEP analyses, one for the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$ , and the other one for the performance criterion  $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$ . Then, we observe an equivalence between the two analyses. Recall that the convergence analysis using PEP is essentially the same as taking an appropriate weighted sum of the following inequalities:

$$0 \geq Q_{i,j} := f(x_j) - f(x_i) + \langle \nabla f(x_j), x_i - x_j \rangle + \frac{1}{2L} \|\nabla f(x_i) - \nabla f(x_j)\|^2 + \frac{\mu L}{2(L - \mu)} \left\| x_i - x_j - \frac{1}{L} (\nabla f(x_i) - \nabla f(x_j)) \right\|^2. \quad (6)$$

As in Part I of our paper, we formulate the PEP using the basis  $\{x_{k+1}^+ - x_k^+\}$  instead of  $\{\nabla f(x_k)\}$ . Although we will not formulate the positive semi-definite condition in terms of matrices, it can be written in such a form after substituting (8) or (11) and expressing the quantity  $S$  as a quadratic form. To avoid notational overlap, different notations are used for each case, as shown in Table 1. Unless otherwise specified, the indices  $i$  or  $j$  range over  $I = \{0, 1, \dots, N, *\}$ .

#### 3.1 PEP analysis for performance criterion $(f(x_N) - f_*)/\|x_0 - x_*\|^2$

In this subsection, we present a PEP analysis for the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$ .

**Theorem 3.** *Given a set of non-negative multipliers  $\{\lambda_{i,j}\}$ , the inequality*

$$f(x_N) - f(x_*) \leq \frac{\tau}{2} \|x_0 - x_*\|^2 \quad (7)$$

*can be obtained by taking the weighted sum of (6) with weights  $\lambda_{i,j}$ ,<sup>1</sup> if and only if the following conditions are satisfied:*

(F1)

$$\sum_{i \in I} \lambda_{i,k} - \sum_{j \in I} \lambda_{k,j} = \begin{cases} 1 & \text{if } k = N \\ -1 & \text{if } k = * \\ 0 & \text{otherwise.} \end{cases}$$

---

<sup>1</sup>This means that after taking the weighted sum, (7) can be derived using only algebraic manipulation, neglecting terms whose removal does not affect the validity of the inequality. See Appendix A for the precise meaning.

(F2) For any points  $x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_* \in \mathbb{R}^d$  satisfying

$$\begin{aligned} y_k^+ - y_{k-1}^+ &= \sum_{i \in I} \lambda_{i, N-k} (x_i^+ - x_{N-k}^+) \text{ for } k = 1, \dots, N \\ \frac{y_0^+ + y_*}{2} - y_N^+ &= \frac{x_N^+ - x_*}{2} + \sum_{i \in I} \lambda_{i, *} (x_i^+ - x_*), \end{aligned} \quad (8)$$

the following quantity is non-negative:

$$\begin{aligned} S &= -L \langle \tilde{\mathbf{H}}^{-1} \mathbf{x}, \mathbf{T} \mathbf{y} \rangle + \frac{L}{2} \left\| y_0 - \frac{y_0^+ + y_*}{2} - \frac{x_N^+ - x_*}{2} \right\|^2 \\ &\quad - \frac{\mu L}{L - \mu} \left( \sum_{j=1}^N \langle y_k^+ - y_{k-1}^+, x_{N-k}^+ \rangle + \left\langle \frac{y_0^+ + y_*}{2} - y_N^+, x_* \right\rangle + \left\langle \frac{y_0^+ - y_*}{2}, x_N^+ \right\rangle \right), \end{aligned} \quad (9)$$

where

$$\begin{aligned} \mathbf{x} &= [x_0^+ - x_0, x_1^+ - x_0^+, \dots, x_N^+ - x_{N-1}^+]^T \\ \mathbf{y} &= [y_0^+ - y_0, y_1^+ - y_0^+, \dots, y_N^+ - y_{N-1}^+]^T. \end{aligned}$$

The proof of Theorem 3 can be found in Appendix A. A pair  $(\tau, \{\lambda_{i,j}\})$  that satisfies the assumptions in Theorem 3 is referred to as a *dual feasible point*.

### 3.2 PEP analysis for performance criterion $\|\nabla f(x_N)\|^2 / (f(x_0) - f_*)$

In this subsection, we present a PEP analysis for the performance criterion  $\|\nabla f(x_N)\|^2 / (f(x_0) - f_*)$ .

**Theorem 4.** Given a set of non-negative multipliers  $\{\nu_{i,j}\}$ , the inequality

$$\frac{1}{2\tau} \|\nabla f(y_N)\|^2 \leq f(y_0) - f(y_*) + \frac{L}{2} \left\| \sum_{i \in I} \nu_{i,*} (y_i^+ - y_*) \right\|^2 \quad (10)$$

can be obtained by taking the weighted sum of (6) with weights  $\nu_{i,j}$ ,<sup>2</sup> if and only if the following conditions are satisfied:

(G1)

$$\sum_{i \in I} \nu_{i,k} - \sum_{j \in I} \nu_{k,j} = \begin{cases} 1 & \text{if } k = * \\ -1 & \text{if } k = 0 \\ 0 & \text{otherwise.} \end{cases}$$

---

<sup>2</sup>See Footnote 1.

(G2) For any points  $x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_* \in \mathbb{R}^d$  satisfying

$$\begin{aligned} x_k^+ - x_{k-1}^+ &= \sum_{i \in I} \nu_{i, N-k} (y_i^+ - y_{N-k}^+) \text{ for } k = 1, \dots, N-1 \\ x_0^+ - x_* &= \sum_{i \in I} \nu_{i, N} (y_i^+ - y_N^+) \\ \frac{x_* - x_N^+}{2} &= \frac{y_* - y_0^+}{2} + \sum_{i \in I} \nu_{i, *} (y_i^+ - y_*), \end{aligned} \tag{11}$$

the quantity (9) is non-negative.

The proof of Theorem 4 can be found in Appendix B. A pair  $(\tau, \{\nu_{i,j}\})$  that satisfies the assumptions in Theorem 4 is referred to as a *dual feasible point*. The presence of the squared distance term on the right-hand side of (10) is undesirable. However, the following proposition shows that this term can be removed when the point  $x_*$  is involved in the proof only through the inequality  $0 \geq Q_{N,*}$ .

**Proposition 1.** *If the set of multipliers  $\{\nu_{i,j}\}$  satisfy*

$$\nu_{i,*} = \begin{cases} 1 & \text{if } i = N \\ 0 & \text{otherwise,} \end{cases} \text{ and } \nu_{*,j} = 0 \text{ for } j = 0, \dots, N, \tag{12}$$

then the statement of Theorem 2 is valid with the inequality (10) replaced by

$$\frac{1}{2\tau} \|\nabla f(y_N)\|^2 \leq f(y_0) - f(y_*). \tag{13}$$

The proof of Proposition 1 can be found in Appendix B.1.

### 3.3 Correspondence between two PEP analyses

It is straightforward to observe the following fact.

**Proposition 2.** *If (8) and (11) are equivalent for the given multipliers  $\{\lambda_{i,j}\}$  and  $\{\nu_{i,j}\}$ , then the conditions (F2) and (G2) are equivalent.*

Thus, we obtained a natural strategy to translate a convergence analysis for the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$  into another one for performance criterion  $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$ : suppose we have a proof of the convergence guarantee (7) using Theorem 3 with the set of multipliers  $\{\lambda_{i,j}\}$ . Then, in most cases, we can find a set of multiplier  $\{\nu_{i,j}\}$  for which (11) is equivalent to (8),<sup>3</sup> and (G2) holds. We denote the map that transforms  $\{\lambda_{i,j}\}$  into  $\{\nu_{i,j}\}$  by this rule as  $\Phi$  (i.e.,  $\Phi(\{\lambda_{i,j}\}) = \{\nu_{i,j}\}$ ). The same reasoning applies when the roles of the two analyses are switched.

However, we note that this equivalence does not provide a complete one-to-one correspondence between the two types of convergence analyses. This is because some  $\{\lambda_{i,j}\} \geq 0$  (i.e., a valid proof) are mapped to  $\{\nu_{i,j}\} = \Phi(\{\lambda_{i,j}\}) \not\geq 0$  (i.e., an invalid proof).

---

<sup>3</sup>This is the case for almost every  $\{\lambda_{i,j}\}$ , similar to how almost every square matrix is invertible.

### 3.4 Exact correspondence between two relaxed PEP analyses

In the previous subsection, we provided a general technique for translating convergence analyses. In this subsection, we show that this provides a complete one-to-one correspondence between all valid proofs, if only a specific subset of interpolation inequalities can be used. Specifically, we restrict Theorem 3 by assuming that

$$\lambda_{i,j} = 0 \text{ for all } (i, j) \notin \{(k, k+1), (*, k)\}_{0 \leq k \leq N-1} \cup \{(*, N)\}, \quad (14)$$

and restrict Theorem 4 by assuming that

$$\nu_{i,j} = 0 \text{ for all } (i, j) \notin \{(k, k+1), (N, k)\}_{0 \leq k \leq N-1} \cup \{(N, *)\}. \quad (15)$$

The restriction (14) was adopted in [DT14], and (15) was adopted in [KF21]. The following proposition shows that the map  $\Phi$  gives a one-to-one correspondence between all valid  $\{\lambda_{i,j}\}$  and all valid  $\{\nu_{i,j}\}$ .

**Proposition 3.** *If the set of multipliers  $\{\nu_{i,j}\}$  satisfies  $\nu_{i,j} \geq 0$ , (G1), and (15), then  $\{\lambda_{i,j}\} = \Phi^{-1}(\{\nu_{i,j}\})$  satisfies  $\lambda_{i,j} \geq 0$ , (F1), and (14). The same result holds when switching the roles of  $\{\nu_{i,j}\}$  and  $\{\lambda_{i,j}\}$ .*

The proof of Proposition 3 can be found in Appendix C. Note that the convergence guarantee (13) can be obtained instead of the weaker one (10) by Proposition 1. Combined with Proposition 2, we conclude that any PEP analysis in [DT14] is equivalent to the one in [KF21], and vice versa. This conclusion is valid for any  $\mu \in [0, L)$ . See [Kim+24, Thm. 1] for the same conclusion in the non-strongly convex case ( $\mu = 0$ ). See [KY24, Thm. 3; Kim+24, Thm. 2] for the essentially same conclusion in the corresponding continuous-time analysis ( $\mu = 0$  and  $1/L \rightarrow 0$ ).

## 4 Proof of Theorem 1

In this section, we prove Theorem 1 using the correspondence result from the previous section. Note that the H-matrix representation of GD is given by

$$\mathbf{H}_{i,j} = \begin{cases} \gamma L & \text{if } i = j \\ 0 & \text{if } i > j. \end{cases} \quad (16)$$

In particular, the anti-transpose of GD is itself. Thus, one can expect an equivalence between proving Theorem 1 and proving Theorem 2. The restricted correspondence result in Section 3.4 is not sufficient for our purpose since the relaxed PEP in [DT14] cannot prove Theorem 1, as indicated by a numerical experiment in [THG17, §4.1.2]. Therefore, we need to use the general correspondence result in Section 3.3.

It turns out that the proof of Theorem 2 in Part I of our paper fits well with our correspondence framework for proving Theorem 1. We summarize its structure in the following lemma.

**Lemma 1** ([Kim24, Lem. 1]). *The convergence guarantee (4) can be proved by taking a weighted sum of the interpolation inequalities (6) with non-negative weights  $\nu_{i,j}$  satisfying*

$$\nu_{i,j} = 0 \text{ for all } (i, j) \notin \{(k, k+1), (k+1, k), (N, k)\}_{0 \leq k \leq N-1} \cup \{(N, *)\} \quad (17)$$

The following proposition generalizes one direction of Proposition 3, and provides a way to translate the proof of Theorem 2 into the one of Theorem 1.

**Proposition 4.** *If a set of multipliers  $\{\nu_{i,j}\}$  satisfy  $\nu_{i,j} \geq 0$ , (G1), and (17), then  $\{\lambda_{i,j}\} = \Phi^{-1}(\{\nu_{i,j}\})$  satisfies  $\lambda_{i,j} \geq 0$  and (F2).*

The proof of Proposition 4, and in particular the expression for  $\{\lambda_{i,j}\} = \Phi^{-1}(\{\nu_{i,j}\})$ , can be found in Appendix D. We can now prove Theorem 1.

*Proof of Theorem 1.* For given  $N$ ,  $\mu$ ,  $L$ , and  $\gamma$ , consider the dual feasible point  $(\tau, \{\nu_{i,j}\})$  given by Lemma 1 with Proposition 1. Then, by Proposition 4,  $(\tau, \Phi^{-1}(\{\nu_{i,j}\}))$  is a dual feasible point. Applying Theorem 3 to this dual feasible point yields the upper bound (3), which matches the lower bound proved in [THG17, §4.1.2]. This completes the proof.  $\square$

See Remark 1 in the appendix for a comparison of our findings with [GSW24, Conj. 2].

**Example 1.** *For  $N = 5$ ,  $\mu = 0.1$ ,  $L = 1$ , and  $\gamma = \gamma^*(N, \mu, L)$  (the optimal stepsize), the set of multipliers  $\{\lambda_{i,j}\}$  used in the proof of Theorem 1 is calculated as follows:*

$$[\lambda_{i,j}] = \begin{matrix} & * & 0 & 1 & 2 & 3 & 4 & 5 \\ * & \left[ \begin{array}{cccccc} 0 & 0.0384 & 0.0621 & 0.1063 & 0.1873 & 0.3342 & 0.2718 \\ 0 & 0 & 0.0182 & 0.0119 & 0.0060 & 0.0020 & 0.0003 \\ 1 & 0 & 0 & 0.0472 & 0.0237 & 0.0080 & 0.0013 \\ 2 & 0 & 0 & 0 & 0.1186 & 0.0401 & 0.0067 \\ 3 & 0 & 0 & 0 & 0 & 0.2876 & 0.0479 \\ 4 & 0 & 0 & 0 & 0 & 0 & 0.6719 \\ 5 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right. & \end{matrix}. \quad (18)$$

A MATLAB code for generating  $\{\lambda_{i,j}\}$  used in the proof of Theorem 1 is available at

<https://github.com/jungbinkim1/GD-Exact-Rate>.

## 5 Conclusion

In this paper, we identified the exact convergence rate of GD for the performance criterion  $(f(x_N) - f_*)/\|x_0 - x_*\|^2$  by proving [DT14, Conj. 1; THG17, Conj. 2]. Our proof is very elementary, although Part I relies on a heavy symbolic calculation. The problem is simple, but it was one of the central open questions in classical first-order convex optimization, possibly dating back to the early development of complexity theory in the 1980s (a standard reference is [NY83]). However, it is interesting that the anti-transpose relationship between first-order methods, which has not gained attention until recently, played a crucial role in obtaining our proof.

A drawback of our proof of Theorem 1 is that the proof itself — taking the weighted sum of inequalities (6) with weights (18), and then performing algebraic manipulations, yields the desired convergence guarantee (3) — seems not understandable without its derivation. This contrasts with the proofs of Theorem 2 in [RGP24; Kim24], which do not deviate much from the traditional proof structure. We hope that a simpler proof of Theorem 1 will be found in future work.

## A Proof of Theorem 3

The condition that the inequality (7) can be obtained by taking the weighted sum of (6) with weights  $\lambda_{i,j}$  means that the following inequality can be proved only using its algebraic structure:

$$\begin{aligned} & f_N^+ - f_* + \frac{1}{2L} \|g_N\|^2 - \frac{\tau}{2} \|x_0 - x_*\|^2 \\ & \leq \sum_{i,j \in I} \lambda_{ij} \left( f_j^+ - f_i^+ + \langle g_j, x_i^+ - x_j^+ \rangle + \frac{\mu L}{2(L-\mu)} \|x_i^+ - x_j^+\|^2 \right). \end{aligned} \quad (19)$$

If readers find the expression unfamiliar, they can see [Kim24, §2.2] before going through the proof. For (19) to hold for any  $\{x_k^+, g_k, f_k^+\}$ , the terms  $f_k^+$  in (19) should vanish, which is (F1). When (F1) holds, using  $\mathbf{g} = -L\tilde{\mathbf{H}}^{-1}\mathbf{x}$ , (19) can be written as follows:

$$\begin{aligned} 0 & \leq S_1(x_0, x_0^+, \dots, x_N^+, x_*) \\ & := \frac{\tau}{2} \|x_0 - x_*\|^2 - \frac{L}{2} \left\| (\tilde{\mathbf{H}}^{-1}\mathbf{x})_{N+1} \right\|^2 \\ & \quad - L \left\langle \tilde{\mathbf{H}}^{-1}\mathbf{x}, \begin{bmatrix} \sum_{i \in I} \lambda_{i,0} (x_i^+ - x_0^+) \\ \sum_{i \in I} \lambda_{i,1} (x_i^+ - x_1^+) \\ \vdots \\ \sum_{i \in I} \lambda_{i,N} (x_i^+ - x_N^+) \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \lambda_{ij} \|x_i^+ - x_j^+\|^2. \end{aligned}$$

Thus, proving the inequality (19) is equivalent to proving  $\inf S_1 \geq 0$ . We introduce dual iterates  $y_0^+, \dots, y_N^+, y_*$  by the rule (8). Using (F1) and (8), we have

$$\begin{aligned} \sum_{i \in I} \lambda_{i,N} (x_i^+ - x_N^+) & = \sum_{i,j \in I} \lambda_{i,j} (x_i^+ - x_j^+) - \sum_{j \in I \setminus \{N\}} \sum_{i \in I} \lambda_{i,j} (x_i^+ - x_j^+) \\ & = (x_* - x_N^+) - \sum_{k=1}^N (y_k^+ - y_{k-1}^+) - \left( \frac{y_0^+ + y_*}{2} - y_N^+ - \frac{x_N^+ - x_*}{2} \right) \\ & = y_0^+ - \mathbf{v}, \end{aligned} \quad (20)$$

where  $\mathbf{v} = \frac{y_0^+ + y_*}{2} + \frac{x_N^+ - x_*}{2}$ . We can now rewrite (19) as follows:

$$\begin{aligned} 0 & \leq S_2(x_0, x_0^+, \dots, x_N^+, x_*, y_0^+, \dots, y_N^+, y_*) \\ & := \frac{\tau}{2} \|x_0 - x_*\|^2 - \frac{L}{2} \left\| (\tilde{\mathbf{H}}^{-1}\mathbf{x})_{N+1} \right\|^2 \\ & \quad - L \left\langle \tilde{\mathbf{H}}^{-1}\mathbf{x}, \begin{bmatrix} y_N^+ - y_{N-1}^+ \\ \vdots \\ y_1^+ - y_0^+ \\ y_0^+ - \mathbf{v} \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \lambda_{ij} \|x_i^+ - x_j^+\|^2. \end{aligned}$$

Thus, proving (19) is equivalent to proving  $\inf S_2 \geq 0$ , where the infimum is taken over all points  $x_0, x_0^+, \dots, x_N^+, x_*, y_0^+, \dots, y_N^+, y_*$  that satisfy (8). Introduce an auxiliary iterate  $y_0$  and consider the



following quantity:

$$\begin{aligned}
& S_3(x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*) \\
& := \frac{L}{2} \|y_0 - \mathbf{v}\|^2 + \frac{\tau}{2} \|x_0 - x_*\|^2 \\
& \quad - L \langle \tilde{\mathbf{H}}^{-1} \mathbf{x}, \mathbf{T} \mathbf{y} \rangle + \frac{\mu L}{2(L - \mu)} \sum_{i,j \in I} \lambda_{ij} \|x_j^+ - x_i^+\|^2,
\end{aligned}$$

When other points are fixed, one can verify that  $S_2 = \inf_{y_0} S_3$ . Thus, proving (19) is equivalent to proving  $S_3 \geq 0$ , where the infimum is taken over all points  $x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*$  that satisfy (8). We proceed by rewriting the last term of  $S_3$ .

$$\begin{aligned}
\sum_{i,j \in I} \lambda_{i,j} \|x_i^+ - x_j^+\|^2 &= \sum_{k \in I} \left( \sum_{j \in I} \lambda_{k,j} + \sum_{i \in I} \lambda_{i,k} \right) \|x_k^+\|^2 - 2 \sum_{j \in I} \sum_{i \in I} \lambda_{i,j} \langle x_i^+, x_j^+ \rangle \\
&= \sum_{k \in I} \left( \sum_{j \in I} \lambda_{k,j} - \sum_{i \in I} \lambda_{i,k} \right) \|x_k^+\|^2 - 2 \sum_{j \in I} \left\langle \sum_{i \in I} \lambda_{i,j} (x_i^+ - x_j^+), x_j^+ \right\rangle.
\end{aligned}$$

We simplify the first term by using (F1) and the second term by using (8) and (20).

$$\begin{aligned}
\sum_{i,j \in I} \lambda_{i,j} \|x_i^+ - x_j^+\|^2 &= -\|x_N^+\|^2 + \|x_*\|^2 - 2 \sum_{k=1}^N \langle y_k^+ - y_{k-1}^+, x_{N-k}^+ \rangle \\
&\quad - 2 \left\langle \frac{y_0^+ + y_*}{2} - y_N^+ - \frac{x_N^+ - x_*}{2}, x_* \right\rangle - 2 \left\langle \frac{y_0^+ - y_*}{2} - \frac{x_N^+ - x_*}{2}, x_N^+ \right\rangle \\
&= -2 \left( \sum_{k=1}^N \langle y_k^+ - y_{k-1}^+, x_{N-k}^+ \rangle + \left\langle \frac{y_0^+ + y_*}{2} - y_N^+, x_* \right\rangle + \left\langle \frac{y_0^+ - y_*}{2}, x_N^+ \right\rangle \right).
\end{aligned}$$

This completes the proof.

## B Proof of Theorem 4

The condition that the inequality (10) can be obtained by taking the weighted sum of (6) with weights  $\nu_{i,j}$  means that the following inequality can be proved only using its algebraic structure.

$$\begin{aligned}
& \frac{1}{2\tau} \|g_N\|^2 + f_* - f_0^+ - \frac{1}{2L} \|g_0\|^2 - \frac{L}{2} \left\| \sum_{i \in I} \nu_{i,*} (y_i^+ - y_*) \right\|^2 \\
& \leq \sum_{i,j \in I} \nu_{ij} \left( f_j^+ - f_i^+ + \langle g_j, y_i^+ - y_j^+ \rangle + \frac{\mu L}{2(L - \mu)} \|y_i^+ - y_j^+\|^2 \right).
\end{aligned} \tag{21}$$

For (21) to hold for any  $\{x_k^+, g_k, f_k^+\}$ , the terms  $f_k^+$  in (21) should vanish, which is (G1). When (G1) holds, using  $\mathbf{g} = -L\tilde{\mathbf{H}}^{-1}\mathbf{x}$  (in particular,  $g_0 = -L(y_0^+ - y_0)$ ), (21) can be written as follows:

$$\begin{aligned} 0 &\leq S_1(y_0, y_0^+, \dots, y_N^+, y_*) \\ &:= \frac{L}{2} \|y_0^+ - y_0\|^2 - \frac{L^2}{2\tau} \left\| (\tilde{\mathbf{H}}^{-1}\mathbf{y})_{N+1} \right\|^2 + \frac{L}{2} \left\| \sum_{i \in I} \nu_{i,*} (y_i^+ - y_*) \right\|^2 \\ &\quad - L \left\langle \tilde{\mathbf{H}}^{-1}\mathbf{y}, \begin{bmatrix} \sum_{i \in I} \nu_{i,0} (y_i^+ - y_0^+) \\ \sum_{i \in I} \nu_{i,1} (y_i^+ - y_1^+) \\ \vdots \\ \sum_{i \in I} \nu_{i,N} (y_i^+ - y_N^+) \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \nu_{ij} \|y_i^+ - y_j^+\|^2. \end{aligned}$$

Thus, proving the inequality (21) is equivalent to proving  $\inf S_1 \geq 0$ . We introduce dual iterates  $x_0^+, \dots, x_N^+, x_*$  by the rule of (11). Using (G1) and (11), we have

$$\begin{aligned} \sum_{i \in I} \nu_{i,0} (y_i^+ - y_0^+) &= \sum_{i,j \in I} \nu_{i,j} (y_i^+ - y_j^+) - \sum_{j \in I \setminus \{0\}} \sum_{i \in I} \nu_{i,j} (y_i^+ - y_j^+) \\ &= (y_0^+ - y_*) - \sum_{k=1}^{N-1} (x_k^+ - x_{k-1}^+) - (x_0^+ - x_*) - \left( \frac{x_* - x_N^+}{2} - \frac{y_* - y_0^+}{2} \right) \quad (22) \\ &= x_N^+ - x_{N-1}^+ + y_0^+ - \mathbf{v}, \end{aligned}$$

where  $\mathbf{v} = \frac{y_0^+ + y_*}{2} + \frac{x_N^+ - x_*}{2}$ . Note that the last equality in (11) can be written as  $\sum_{i \in I} \nu_{i,*} (y_i^+ - y_*^+) = y_0^+ - \mathbf{v}$ . We can now rewrite (21) as follows:

$$\begin{aligned} 0 &\leq S_2(x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*) \\ &:= \frac{L}{2} \|y_0^+ - y_0\|^2 - \frac{L^2}{2\tau} \left\| (\tilde{\mathbf{H}}^{-1}\mathbf{y})_{N+1} \right\|^2 + \frac{L}{2} \|y_0^+ - \mathbf{v}\|^2 \\ &\quad - L \left\langle \tilde{\mathbf{H}}^{-1}\mathbf{y}, \begin{bmatrix} x_N^+ - x_{N-1}^+ + y_0^+ - \mathbf{v} \\ \vdots \\ x_1^+ - x_0^+ \\ x_0^+ - x_* \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \nu_{ij} \|y_i^+ - y_j^+\|^2 \\ &= \frac{L}{2} \|y_0 - \mathbf{v}\|^2 - \frac{L^2}{2\tau} \left\| (\tilde{\mathbf{H}}^{-1}\mathbf{y})_{N+1} \right\|^2 \\ &\quad - L \left\langle \tilde{\mathbf{H}}^{-1}\mathbf{y}, \begin{bmatrix} x_N^+ - x_{N-1}^+ \\ \vdots \\ x_1^+ - x_0^+ \\ x_0^+ - x_* \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \nu_{ij} \|y_i^+ - y_j^+\|^2. \end{aligned}$$

Thus, proving (21) is equivalent to proving  $\inf S_2 \geq 0$ , where the infimum is taken over all points  $x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*$  that satisfy (11). Introduce an auxiliary iterate  $x_0$  and consider

the following quantity:

$$\begin{aligned}
0 &\leq S_3(x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*) \\
&:= \frac{L}{2} \|y_0 - \mathbf{v}\|^2 + \frac{\tau}{2} \|x_0 - x_*\|^2 \\
&\quad - L \langle \tilde{\mathbf{H}}^{-1} \mathbf{y}, \mathbf{T} \mathbf{x} \rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j} \nu_{ij} \|y_i^+ - y_j^+\|^2.
\end{aligned}$$

When other points are fixed, one can check that  $S_2 = \inf_{x_0} S_3$ . Thus, proving (21) is equivalent to proving  $S_3 \geq 0$ , where the infimum is taken over all points  $x_0, x_0^+, \dots, x_N^+, x_*, y_0, y_0^+, \dots, y_N^+, y_*$  that satisfy (11). Using Definition 1, we have  $\langle \tilde{\mathbf{H}}^{-1} \mathbf{y}, \mathbf{T} \mathbf{x} \rangle = \langle \mathbf{T} \mathbf{y}, \tilde{\mathbf{H}}^{-\mathbf{A}} \mathbf{x} \rangle$ . We proceed by rewriting the last term of  $S_3$ .

$$\begin{aligned}
\sum_{i,j \in I} \nu_{i,j} \|y_i^+ - y_j^+\|^2 &= \sum_{k \in I} \left( \sum_{j \in I} \nu_{k,j} + \sum_{i \in I} \nu_{i,k} \right) \|y_k^+\|^2 - 2 \sum_{j \in I} \sum_{i \in I} \nu_{i,j} \langle y_i^+, y_j^+ \rangle \\
&= \sum_{k \in I} \left( \sum_{j \in I} \nu_{k,j} - \sum_{i \in I} \nu_{i,k} \right) \|y_k^+\|^2 - 2 \sum_{j \in I} \left\langle \sum_{i \in I} \nu_{i,j} (y_i^+ - y_j^+), y_j^+ \right\rangle.
\end{aligned}$$

We simplify the first term by using (G1) and the second term by using (11) and (22).

$$\begin{aligned}
\sum_{i,j \in I} \nu_{i,j} \|y_i^+ - y_j^+\|^2 &= -\|y_*\|^2 + \|y_0^+\|^2 - 2 \sum_{k=1}^{N-1} \langle x_k^+ - x_{k-1}^+, y_{N-k}^+ \rangle - 2 \langle x_0^+ - x_*, y_N^+ \rangle \\
&\quad - 2 \left\langle \frac{x_* - x_N^+}{2} - \frac{y_* - y_0^+}{2}, y_* \right\rangle - 2 \left\langle \frac{x_N^+ + x_*}{2} - x_{N-1}^+ + \frac{y_0^+ - y_*}{2}, y_0^+ \right\rangle \\
&= -2 \left( \sum_{k=1}^N \langle y_k^+ - y_{k-1}^+, x_{N-k}^+ \rangle + \left\langle \frac{y_0^+ + y_*}{2} - y_N^+, x_* \right\rangle + \left\langle \frac{y_0^+ - y_*}{2}, x_N^+ \right\rangle \right).
\end{aligned}$$

This completes the proof.

## B.1 Proof of Proposition 1

Assume that (12) holds. Then, the quantity  $S_1$  in Appendix B is the sum of  $(\frac{L}{2} + \frac{\mu L}{2(L-\mu)}) \|y_* - y_N^+\|^2$  and other terms independent of  $y_*$ , and is thus minimized at  $y_* = y_N^+$  when other points  $y_0, y_0^+, \dots, y_N^+$  are fixed. When the inequality (10) is replaced by (13) in the statement of Theorem 4, mimicking the proof in Appendix B yields the following quantity instead of  $S_1$ :

$$\begin{aligned}
&S_1'(y_0, y_0^+, \dots, y_N^+, y_*) \\
&:= \frac{L}{2} \|y_0^+ - y_0\|^2 - \frac{L^2}{2\tau} \left\| (\tilde{\mathbf{H}}^{-1} \mathbf{y})_{N+1} \right\|^2 \\
&\quad - L \left\langle \tilde{\mathbf{H}}^{-1} \mathbf{y}, \begin{bmatrix} \sum_i \nu_{i,0} (y_i^+ - y_0^+) \\ \sum_i \nu_{i,1} (y_i^+ - y_1^+) \\ \vdots \\ \sum_i \nu_{i,N} (y_i^+ - y_N^+) \end{bmatrix} \right\rangle + \frac{\mu L}{2(L-\mu)} \sum_{i,j \in I} \nu_{ij} \|y_i^+ - y_j^+\|^2.
\end{aligned}$$

Since  $S'_1$  is the sum of  $\frac{\mu L}{2(L-\mu)}\|y_* - y_N^+\|^2$  and other terms independent of  $y_*$ , it is minimized at  $y_* = y_N^+$  when other points  $y_0, y_0^+, \dots, y_N^+$  are fixed. Thus, we have

$$\begin{aligned} \inf_{y_0, y_0^+, \dots, y_N^+, y_*} S_1(y_0, y_0^+, \dots, y_N^+, y_*) &= \inf_{y_0, y_0^+, \dots, y_N^+} S_1(y_0, y_0^+, \dots, y_N^+, y_N^+) \\ &= \inf_{y_0, y_0^+, \dots, y_N^+} S'_1(y_0, y_0^+, \dots, y_N^+, y_N^+) \\ &= \inf_{y_0, y_0^+, \dots, y_N^+, y_*} S'_1(y_0, y_0^+, \dots, y_N^+, y_*). \end{aligned}$$

Therefore, proving (13) is equivalent to proving  $\inf S_1 \geq 0$ . This completes the proof.

## C Proof of Proposition 3

We first prove the direction stated in the proposition, and then provide a proof sketch for the opposite direction. Although not explicitly stated in the proposition, we assume that all multipliers which are allowed to be non-zero, are indeed non-zero. This assumption prevents the cases where the maps (8) or (11) have no inverse.

**Translating  $\{\nu_{i,j}\}$  into  $\{\lambda_{i,j}\}$ .** Let  $A_k = 1 + \sum_{j=0}^{k-1} \nu_{N,j}$  for  $k = 1, \dots, N$ . Then,  $\{A_k\}$  is an increasing sequence in  $(1, \infty)$ . Using (15) and (G1), we can express all the multipliers  $\{\nu_{i,j}\}$  in terms of  $A_k$  as follows:

$$\begin{aligned} \nu_{k-1,k} &= A_k \text{ for } k = 1, \dots, N \\ \nu_{N,0} &= A_1 - 1 \\ \nu_{N,k} &= A_{k+1} - A_k \text{ for } k = 1, \dots, N-1 \\ \nu_{N,*} &= 1. \end{aligned} \tag{23}$$

Thus, (11) without the last equality can be written as follows:

$$\begin{bmatrix} x_{N-1}^+ - x_{N-2}^+ \\ \vdots \\ x_1^+ - x_0^+ \\ x_0^+ - x_* \end{bmatrix} = - \begin{bmatrix} \mathbf{M}_{1,1} & \cdots & \mathbf{M}_{1,N-1} & \mathbf{M}_{1,N} \\ & \ddots & \vdots & \vdots \\ & & \mathbf{M}_{N-1,N-1} & \mathbf{M}_{N-1,N} \\ & & & \mathbf{M}_{N,N} \end{bmatrix} \begin{bmatrix} y_1^+ - y_0^+ \\ \vdots \\ y_{N-1}^+ - y_{N-2}^+ \\ y_N^+ - y_{N-1}^+ \end{bmatrix}, \tag{24}$$

where

$$\mathbf{M}_{i,j} = \begin{cases} A_i & \text{if } j = i \\ A_i - A_{i+1} & \text{if } j \geq i + 1. \end{cases} \tag{25}$$

The last equality in (11) can be written as follows:

$$\frac{x_* - x_N^+}{2} = y_N^+ - \frac{y_* + y_0^+}{2}. \tag{26}$$

We rewrite (24) using the inverse of  $\mathbf{M}$  as follows:

$$\begin{bmatrix} y_1^+ - y_0^+ \\ \vdots \\ y_{N-1}^+ - y_{N-2}^+ \\ y_N^+ - y_{N-1}^+ \end{bmatrix} = - \begin{bmatrix} \mathbf{M}_{1,1}^{-1} & \cdots & \mathbf{M}_{1,N-1}^{-1} & \mathbf{M}_{1,N}^{-1} \\ & \ddots & \vdots & \vdots \\ & & \mathbf{M}_{N-1,N-1}^{-1} & \mathbf{M}_{N-1,N}^{-1} \\ & & & \mathbf{M}_{N,N}^{-1} \end{bmatrix} \begin{bmatrix} x_{N-1}^+ - x_{N-2}^+ \\ \vdots \\ x_1^+ - x_0^+ \\ x_0^+ - x_* \end{bmatrix}, \tag{27}$$

where the entries of the inverse matrix are given by

$$\mathbf{M}_{i,j}^{-1} = \begin{cases} \frac{1}{A_i} & \text{if } j = i \\ \frac{1}{A_i} - \frac{1}{A_{i+1}} & \text{if } j \geq i + 1. \end{cases} \quad (28)$$

We set the multipliers  $\{\lambda_{i,j}\}$  as follows:

$$\begin{aligned} \lambda_{k-1,k} &= \frac{1}{A_{N-k+1}} \text{ for } k = 1, \dots, N \\ \lambda_{*,0} &= \frac{1}{A_N} \\ \lambda_{*,k} &= \frac{1}{A_{N-k}} - \frac{1}{A_{N-k+1}} \text{ for } k = 1, \dots, N-1 \\ \lambda_{*,N} &= 1 - \frac{1}{A_1}. \end{aligned} \quad (29)$$

Then, (27) with (28) holds, and (26) holds. Thus, (8) and (11) are equivalent for these  $\{\lambda_{i,j}\}$  and  $\{\nu_{i,j}\}$ . Furthermore, the obtained convergence analysis using  $\{\lambda_{i,j}\}$  is valid since (F1) holds and  $\lambda_{i,j} \geq 0$  for all  $i, j \in I$ .

**Translating  $\{\lambda_{i,j}\}$  into  $\{\nu_{i,j}\}$ .** Let  $A_k = 1/\sum_{j=0}^{N-k} \lambda_{*,j}$  for  $k = 1, \dots, N$ . Then, one can show that  $\{A_k\}$  is an increasing sequence in  $(1, \infty)$ . We can express all the multipliers  $\{\lambda_{i,j}\}$  in terms of  $A_k$  by (29). Following the procedure above in the opposite direction, we arrive at the multipliers  $\{\nu_{i,j}\}$  given by (23), which satisfy all the desired conditions.

## D Proof of Proposition 4

We assume that the readers are familiar with the proof in Appendix C. Although not explicitly stated in the proposition, we assume that all multipliers which are allowed to be non-zero, are indeed non-zero. Let  $A_k = 1 + \sum_{j=0}^{k-1} \nu_{N,j}$  for  $k = 1, \dots, N$ , and  $B_k = \nu_{k,k-1}$  for  $k = 1, \dots, N-1$ . Then,  $\{A_k\}$  is an increasing sequence in  $(1, \infty)$ , and  $\{B_k\}$  is a sequence in  $(0, \infty)$ . For convenience, let  $B_N = 0$ . Using (15) and (G1), we can express all the multipliers  $\{\nu_{i,j}\}$  in terms of  $A_k$  as follows:

$$\begin{aligned} \nu_{k-1,k} &= A_k + B_k \text{ for } k = 1, \dots, N \\ \nu_{N,0} &= A_1 - 1 \\ \nu_{N,k} &= A_{k+1} - A_k \text{ for } k = 1, \dots, N-1 \\ \nu_{k,k-1} &= B_k \text{ for } k = 1, \dots, N-1 \\ \nu_{N,*} &= 1. \end{aligned}$$

Thus, (11) without the last equality can be expressed as (24) with

$$\mathbf{M}_{i,j} = \begin{cases} A_i + B_i & \text{if } j = i \\ A_i - A_{i+1} - B_{i+1} & \text{if } j = i + 1. \\ A_i - A_{i+1} & \text{if } j \geq i + 2 \end{cases}$$

The last equality in (11) can be expressed as (26). The entries of  $\mathbf{M}^{-1}$  can be computed as follows:

$$\mathbf{M}_{i,j}^{-1} = \begin{cases} \frac{1}{A_i+B_i} & \text{if } j = i \\ \mathbf{M}_{i,i}^{-1} - \frac{1}{A_i+B_i} \frac{A_i}{A_{i+1}+B_{i+1}} & \text{if } j = i + 1 \\ \mathbf{M}_{i,j-1}^{-1} - \frac{1}{A_i+B_i} \frac{A_i}{A_{i+1}+B_{i+1}} \left( \prod_{k=i+1}^{j-1} \frac{B_k}{A_{k+1}+B_{k+1}} \right) & \text{if } j \geq i + 2 \end{cases} \quad (30)$$

A MATLAB code for symbolically verifying (30) is available at

<https://github.com/jungbinkim1/GD-Exact-Rate>.

We set the multipliers  $\{\lambda_{i,j}\}$  as follows:

$$\lambda_{i,j} = \begin{cases} \mathbf{M}_{N-j,N-i-1}^{-1} - \mathbf{M}_{N-j,N-i}^{-1} & \text{if } 0 \leq i < j \leq N - 1 \\ \mathbf{M}_{N-j,N}^{-1} & \text{if } i = *, j \leq N - 1 \\ \sum_{k=*,0,\dots,i-1} \lambda_{k,i} - \sum_{k=i+1,\dots,N-1} \lambda_{i,k} & \text{if } 0 \leq i < j = N \\ 1 - \sum_{k=0,\dots,N-1} \lambda_{*,k} & \text{if } i = *, j = N \\ 0 & \text{otherwise,} \end{cases} \quad (31)$$

or equivalently,

$$\lambda_{i,j} = \begin{cases} \frac{1}{A_{N-j}+B_{N-j}} \frac{A_{N-j}}{A_{N-j+1}+B_{N-j+1}} \left( \prod_{k=N-j+1}^{N-i-1} \frac{B_k}{A_{k+1}+B_{k+1}} \right) & \text{if } 0 \leq i < j \leq N - 1 \\ \frac{1}{A_{N-j}+B_{N-j}} - \sum_{k=0}^{j-1} \lambda_{k,j} & \text{if } i = *, j \leq N - 1 \\ \sum_{k=*,0,\dots,i-1} \lambda_{k,i} - \sum_{k=i+1,\dots,N-1} \lambda_{i,k} & \text{if } 0 \leq i < j = N \\ 1 - \sum_{k=0,\dots,N-1} \lambda_{*,k} & \text{if } i = *, j = N \\ 0 & \text{otherwise.} \end{cases} \quad (32)$$

Then, one can easily see that (27) with (30) holds, and that (26) and (F1) hold. One can also show that  $\lambda_{i,j} \geq 0$  for all  $i, j \in I$ , for which we omit the verification, as it can be derived from algebraic manipulation with that  $\{A_k\}$  is an increasing sequence in  $(1, \infty)$ , and that  $\{B_k\}$  is a sequence in  $(0, \infty)$ . This completes the proof.

**Remark 1.** For the set of multipliers  $\{\lambda_{i,j}\}$  we obtained, one can observe that

$$\frac{\lambda_{i,j}}{\lambda_{i',j}} = \frac{\lambda_{i,j'}}{\lambda_{i',j'}}$$

holds when  $0 \leq i, i' \leq j - 1, j' - 1$ . This property can be also observed in [GSW24, Conj. 2]. However, we note that our  $\{\lambda_{i,j}\}$  does not follow their full structure. For example, the equality above does not hold when  $i = *$ .

## References

- [NY83] Arkadij Semenovič Nemirovskij and David Borisovich Yudin. “Problem complexity and method efficiency in optimization”. In: (1983).

- [DT14] Yoel Drori and Marc Teboulle. “Performance of first-order methods for smooth convex minimization: a novel approach”. In: *Mathematical Programming* 145.1 (2014), pp. 451–482.
- [THG17] Adrien B Taylor, Julien M Hendrickx, and François Glineur. “Smooth strongly convex interpolation and exact worst-case performance of first-order methods”. In: *Mathematical Programming* 161 (2017), pp. 307–345.
- [KF21] Donghwan Kim and Jeffrey A Fessler. “Optimizing the efficiency of first-order methods for decreasing the gradient of smooth convex functions”. In: *Journal of optimization theory and applications* 188.1 (2021), pp. 192–219.
- [TV23] Marc Teboulle and Yakov Vaisbourd. “An elementary approach to tight worst case complexity analysis of gradient based methods”. In: *Mathematical Programming* 201.1 (2023), pp. 63–96.
- [GSW24] Benjamin Grimmer, Kevin Shu, and Alex L Wang. “A strengthened conjecture on the minimax optimal constant stepsize for gradient descent”. In: *arXiv preprint arXiv:2407.11739* (2024).
- [Kim+24] Jaeyeon Kim et al. “Time-reversed dissipation induces duality between minimizing gradient norm and function value”. In: *Advances in Neural Information Processing Systems* 36 (2024).
- [Kim24] Jungbin Kim. *A Proof of Exact Convergence Rate of Gradient Descent. Part I. Performance Criterion  $\|\nabla f(x_N)\|^2/(f(x_0) - f_*)$* . 2024.
- [KY24] Jungbin Kim and Insoon Yang. “Convergence analysis of ODE models for accelerated first-order methods via positive semidefinite kernels”. In: *Advances in Neural Information Processing Systems* 36 (2024).
- [RGP24] Teodor Rotaru, François Glineur, and Panagiotis Patrinos. “Exact worst-case convergence rates of gradient descent: a complete analysis for all constant stepsizes over non-convex and convex functions”. In: *arXiv preprint arXiv:2406.17506* (2024).