WOLF-RAYET STARS – WHAT WE KNOW AND WHAT WE DON'T

O. V. Maryeva a^*

^a Astronomical Institute of the Czech Academy of Sciences, Fričova 298, 25165 Ondřejov, Czech Republic

Today, we have a sufficiently complete picture of what the Wolf-Rayet (WR) stars are. Predictions of stellar evolution theory are in a good agreement with their parameters, estimated from observational data using stellar atmospheres codes; predictions of population synthesis also agree well with number of known WR stars. This article provides an overview of the main historical milestones in the studies of WR stars, showing how we came to this understanding, and what questions are still unanswered.

Keywords: Stars: Wolf-Rayet – Stars: evolution – Stars: atmospheres – General: history and philosophy of astronomy

1. INTRODUCTION

Wolf-Rayet (WR) stars are a class of objects identified on the basis of their spectral features. WR spectra show strong emission lines of helium, nitrogen, carbon and oxygen in different stages of ionization. The width of these lines reaches tens of angstroms, the central intensities are sometimes 10-20 times the intensity of the continuum spectrum. A prominent feature of WRs are two bumps at 4650^{11} and 5808 Å, which are groups of closely located lines of N II, N III, C III and C IV, which are visible even in low-resolution spectra. The lines are formed in an extended atmosphere – in the stellar wind, expelling at velocities of $10^2 - 10^3$ km s⁻¹. WR stars are characterized by a high mass-loss rate of several 10^{-5} M_{\odot}yr⁻¹ and by high temperatures ($T_{\rm eff} \gtrsim 30000$ K). WR are the final stage in the evolution of massive stars (stars with an initial mass $\gtrsim 25$ M_{\odot}) before the core-collapse supernova explosion.

^{*} E-mail: olga.maryeva@asu.cas.cz

¹⁾ Sometimes blue WR bump is also called Bowen blend.

With accumulation of observational data it became clear that emission spectra and WR bumps are intrinsic to objects of different natures. Therefore nowadays there is a clear distinction between WR stars and WR phenomenon, which occurs when fast-moving, hot plasma is expanding around a hot star (Gräfener et al., 2011; Vink, 2015). WR phenomenon may occur in the low mass stars after ejecting their outer layers in the planetary nebula phase and exposing their hot cores prior to the white-dwarf phase (Marcolino et al., 2007; Todt, 2009; van der Hucht et al., 1981). Central stars of planetary nebulae showing WR phenomenon in their spectra are denoted as [WR] (van der Hucht et al., 1981). Rare and still unique case of low mass object with WR phenomenon and the absence of a clearly detected circumstellar nebula is LAMOST J040901.83+323955.6 (Maryeva et al., 2024). It was selected as a WR star by Skoda et al. (2020), but Maryeva et al. (2024) showed that this object is 0.9 M_{\odot} star caught in a rare transitional phase from post-AGB to CSPN. Another unique object with WR phenomenon is IRAS 00500+6713, formed as a result of a merger of two white dwarfs (Gvaramadze et al., 2019). The WR phenomenon is also observed in young supernovae (Gal-Yam et al., 2014).

As we know now, massive stars showing WR spectra are also not a homogeneous class. They are split into two groups: very massive stars and classical WR stars. Very massive stars ($M_{\rm ini} \ge 100 M_{\odot}$) experience strong stellar winds and show WR spectra already during core H-burning phase, when they are located on main sequence. All of them show signs of hydrogen on their surface and are classified as WNh objects (Crowther et al., 2010; Martins et al., 2023; Smith et al., 1996). Classical WR stars (usually just named WR stars) stars are hydrogen-depleted objects that have evolved off the main sequence and suffered intense mass loss (Crowther, 2007; Shenar, 2024). Classical WR stars will be the subject of this review article.

The field of contemporary studies of WR stars is enormous and cannot be fully covered in a single small review like this one, so I'd like to recommend two other detailed reviews – Crowther (2007) and Shenar (2024) – for better understanding what WR stars are. On the other hand, here I will concentrate more on the historical aspects of WR studies, on how we have arrived to our present day understanding of them. Therefore Section 2 will cover the history of studies of WR stars, while Section 3 will highlight several topical problems and open questions about them.

2. FROM DISCOVERY TO UNDERSTANDING

In 1867 year astronomers from Paris observatory Charles Wolf and Georges Rayet discovered three stars in Cygnus constellation those spectra significantly differ from other stars. In contrast with other stars, where usually absorption lines are dominant in spectra, discovered objects showed strong emission lines (Wolf & Rayet, 1867). The class of objects received its name in honor of the discoverers and gradually began to be replenished with new members. One hundred years after the discovery of WRs, by the end of 1960-s the number of known WRs in the Galaxy had reached over a hundred, totalling 127 (Roberts, 1962; Smith, 1968), while now is ~ 700 (Rosslowe & Crowther, 2015). At the same time, in the late 1950-s middle 1960-s, work began to search for extragalactic WRs – WRs in the Large Magellanic Cloud (Westerlund & Rodgers, 1959; Westerlund & Smith, 1964).

Significant contribution to the understanding of the physics of WR stars was given by Carlyle S. Beals (1929). Beals found that some lines in the spectra of WR stars have a P Cygni profile and based on comparison with P Cygni profile lines in spectra of novae he suggested that WRs are surrounded by expanding envelopes (Beals, 1929). Beals also found a difference between WR stars and novae: the P Cygni profiles in the WR stars do not change over time. This allowed to propose that an outflow from WR stars occurs continuously (Beals, 1929). This was confirmed by Chandrasekhar (1934), who developed a solid footing for interpreting P Cygni profiles as arising in expanding atmospheres. Kosirev (1934) used the diagnostics developed by Chandrasekhar to estimate the mass loss and maximum outflow velocity of a WR star and found, respectively, ~ $10^{-5} M_{\odot} yr^{-1}$ and ~ 1000 km s⁻¹.

At the beginning of the 20th century, stellar spectroscopy developed in close cooperation with atomic physics. Investigating spectra of the star ζ Puppis Edward Pickering paid his attention to previously unknown lines 5411, 4541, 4200, 4100, 4026, 3924, 3858, 3813, 3782 Å (Pickering, 1897). He interpreted them as another series of hydrogen (besides Balmer) (Pickering, 1901). Line 4686 Å, first discovered during a solar eclipse and often found in WR spectra, was also considered to be a hydrogen line (Fowler, 1912). Although Fowler (1912) was able to obtain these spectral lines in the laboratory in 1912, during an experiment with a helium-filled tube, he considered their appearance to be a contribution from hydrogen impurity. It was not until 1913 that Niels Bohr's work (Bohr, 1913) explained the nature of these lines as ionized helium He II, confirming the atomic model (see Robotti (1983) for historical review).

In 1890 Pickering drew attention to the similarity of the spectra of WR stars and planetary nebula (Pickering, 1891). Due to this, in papers devoted to emission spectra of nebulae it is possible to find descriptions of spectra of WR stars. The lines at 4647, 4650, 5696, 5801 and 5812 Å were identified with the carbon by Wright (1918). Ira S. Bowen, who found an interpretation of the "nebulium" lines as [O III], gave a detailed list of emission lines observed in nebulae, which includes an identification of almost all lines visible in the WRs (Bowen, 1928). In a 1934 paper Bowen (1934) suggests a significant role of the fluorescence for formation of the NIII 4634, 46340 lines belonging to the WR bump.

Already the spectra of the first discovered WR stars showed a difference: the broad emission bands are located in different places (Huggins & Huggins, 1890; Wolf & Rayet, 1867). After the identification of all the main spectral lines, Beals (1933) proposed the splitting of WR into two groups. This classification was approved in 1938, the International Astronomical Union (IAU) divided the spectra of WR stars into types WN and WC, depending on whether the spectrum was dominated by lines of nitrogen or carbon-oxygen respectively (Beals, 1933; Swings, 1942). WN stars are believed to show the hydrogen burning products via the CNO cycle, while WC stars reveal the helium burning products via the triple- α cycle. Based on the strength of the emission lines and line ratios, WN stars into the spectral subtypes WC4 to WC9 (Crowther et al., 1998; Smith, 1968; Smith et al., 1990, 1994, 1996). Also, there are transition types from WN to WC, which are called WN/C, whose spectra show strong emission lines of carbon and nitrogen simultaneously (Conti & Massey, 1989).

In 1933 Victor A. Ambartsumian (1933) estimated He/H ratio for WR stars using intensities of H β and He II 4686 lines and found He/H \gtrsim 1.8. Subsequent studies (Rublev (1972a) and references therein) confirmed excess of helium. Gamow (1943) suggested that the anomalous composition of WR stars was the result of nuclear processed material being visible on their surfaces. Despite these arguments in favor of far evolved status of WR stars, debates about it continued until the mid-1970s. There was alternative hypothesis, that WR are young and more massive than T Tau objects, before main sequence (Sahade, 1958; Underhill, 1968). It is important to mention the studies of Sergej V. Rublev, in which he developed methods for determining the temperatures and luminosities of WR stars and estimated the hydrogen abundances (Rublev, 1965, 1970, 1972b, 1975). These works played a significant role in the formation of the modern understanding of WR stars as far evolved massive objects.

2.1. Development of numerical methods

Contemporary progress in our understanding of hot stars and especially WR stars has been significantly defined by the progress of numerical modelling of stellar atmospheres, in turn enabled by the development of both physical approximations and numerical methods.

Victor V. Sobolev developed the theory of radiative transfer in moving media and applied it to the determination of physical conditions in the envelopes of WR, Be, novae, etc. type stars (Sobolev, 1947, 1960). He showed that due to the presence of a velocity gradient, photons of spectral lines are able to leave the deep layers of the atmosphere directly, avoiding the usual diffusion process; this significantly simplifies the study of the radiative equilibrium of moving atmospheres. This theory became the basis for the transition to modern quantitative calculations of the stellar wind.

WR stars are essentially hot stellar cores surrounded by extended, low-density atmospheres. Due to their high luminosity the radiation in the stellar wind dominates over collisional processes, and the models of WR's atmospheres should be calculated taking into account the deviations from local thermodynamic equilibrium (non-LTE). Straightforward iterative procedure for the solution of radiative transfer equations – Λ iteration – fails to converge in typical non-LTE conditions, as in scattering dominated media of large optical thickness, its convergence is extremely slow (see overview by Atanackovic-Vukmanovic (2004) and references therein).

Several approaches have been proposed in early 1970-s as a way to numerically solve radiative transfer (RT) problems in non-LTE case. The first fully self-consistent solution of RT problems for non-LTE case was achieved by Auer & Mihalas (1969) who developed the complete linearization method based on the Newton–Raphson scheme to solve the set of discretized structural equations in a robust manner that opened a way for significant progress in the modelling of stellar atmospheres.

A different approach is the "core saturation method", introduced by Rybicki (1972) who implemented a modification of the Λ iteration method that made it practically usable for solving the RT and statistical equilibrium equations. In order to eliminate the cause of the slow convergence of simple Λ iteration (poor conditioning of the equations arising from a large number of scatterings), Rybicki proposed the elimination of scattering events in the line core as they do not contribute much to the transfer process.

Another approach is "perturbation technique" proposed by Cannon (1973a,b) who was the first to use the idea of 'operator splitting' in radiative transfer computations. He replaced the full (exact) Λ operator by a simplified one (so called approximate Lambda operator, or ALO) and computed a small error term made by this approximation by using a perturbation technique. Cannon's operator perturbation technique was an efficient way to combine computational advantages of approximate solutions with the accuracy of exact methods.

In 1980-s computational approximations to accelerate the Λ iteration started to be used together with the operator perturbation technique introduced by Cannon (1973a,b) to simplify the direct solution, making the basis for the class of methods known as Accelerated Lambda Iteration. Wolf-Rainer Hamann (1985, 1986) adopted perturbation technique to approximate lambda operator, based on idea of "core saturation" and presented test calculations for a typical WR star atmosphere. This effort became the foundation for the PoWR code (Hamann & Gräfener, 2003) that is still being actively developed and successfully applied for numerical modelling of various types of astrophysical objects. Independently, D. John Hillier (1990) presented a method based on tridiagonal Newton-Raphson operator, complete linearization scheme and perturbation technique for calculations of WN and WC atmosphere models. Over time, this method has grown into CMFGEN code (Hillier & Miller, 1998).

Since the mid-1990s, CMFGEN and PoWR have become a reliable instruments for studying hot stars with high mass-loss rates. Systematic study of Galactic WR stars started with a series of works by Paul Crowther titled "Fundamental parameters of WR stars" and devoted to the analysis of properties of nitrogen-rich WR stars (WN) based on optical and ultraviolet (UV) data and numerical modelling using CMFGEN code (Crowther et al., 1995a,b,c,d,e). Advances in infrared (IR) spectroscopy also enabled studies of WR stars near the Galactic center (Liermann et al., 2010; Martins et al., 2007). Modeling of large sample of Galctic WN (Hamann et al., 2006) and WC (Sander et al., 2012) stars (≈ 126 objects in total) allowed to obtain a homogeneous set of stellar and atmospheric parameters for them, to determine the correct location of WR stars in the Hertzsprung–Russell (HR) diagram.

PoWR code was widely used for studying extragalactic objects. Among WR stars in nearby galaxies, the ones in Magellanic Clouds are most studied (Bestenlehner et al., 2014; Hainich et al., 2014, 2015). Sample of 74 WR stars belonged to M33 galaxy was analysed by Pritzkuleit (2020) (Pritzkuleit, 2020). Sander et al. (2014) investigated late-type WN stars in M31 galaxy (Sander et al., 2014). These studies extended our understanding of properties of WR stars in different galactic environments and metallicity-dependence of massive star winds.

2.2. Evolution

Modern view on WRs as products of evolution of massive stars was finally formed only in 1980s. Although by the 1980s WR stars were already recognised as descendants of massive OB stars, their exact evolutionary status with respect to the main sequence and other evolved massive stars was still an open question. WR stars were being proposed as possible progenitors of supernovae, and particularly the newly-discovered type Ib supernovae that lack hydrogen but are apparently associated with young massive stars.

The start of systematic X-ray observations with space-borne X-ray telescopes in 1960s significantly advanced the theory of close binary stars and their interactions, and it led Bohdan Paczyński (1967) to the suggestion that WR stars are the products of binary evolution – namely, mass exchange in a binary system. In his scenario, a more massive star in binary system evolves faster and, upon exhaustion of hydrogen in the core, fills the Roche lobe and a rapidly loses its hydrogen envelope that outflows towards the companion. As a result, when the envelope is fully transferred, its hot helium core remains as a WR star with little hydrogen in the outermost layers (Paczyński, 1973).

The hypothesis of Paczyński was well accepted and it naturally explained the anomalous chemical composition of WR stars as naked stellar cores enriched with products of nuclear reactions. On the other hand, based on the similarity of Of and WN stars (Figure 1) and on the first estimates of the mass loss rate for hot massive stars based on UV observations Peter Conti (1975) suggested that the stellar wind of O stars is strong enough to remove the outer layers and uncover the stellar core, thus proposing a single star mass loss channel to form WR stars. In addition, brightest WR star, γ Vel, was inconsistent with Paczyński's hypothesis as, while being a binary, its orbit was clearly elliptical and its second component lacks any signs of previous mass exchange, both suggesting that there was no interaction between the components (Conti, 2015).

In 1984 Peter Conti (1984) introduced a new class of objects. In his talk at the IAU symposium, Peter Conti grouped S Doradus variables, Hubble–Sandage variables, η Carinae, P Cygni, and other similar stars together under the term "luminous blue variables" (LBV). LBV are characterized by strong mass loss (around $10^{-5} \text{ M}_{\odot} \text{yr}^{-1}$) and occasional giant eruptive events (Humphreys & Davidson, 1994). Conti suggested that massive O stars transit to WR stars through this short-duration LBV phase. This idea is consolidated as the "Conti scenario" and becomes the main channel for WR star formation. Modern stellar evolution calculations are consistent with the "Conti scenario" (Groh et al., 2014; Sander et al., 2019). According to numerical calculations the stars with initial mass of 60 M_☉ spend 2×10^5 yr in the LBV phase — only 5% of their lifetime (Groh et al., 2014). Recently observed transition from LBV to WN8 spectral type in Romano' star is also a direct confirmation of "Conti scenario" (Maryeva et al., 2019; Polcaro et al., 2016).

Similar scenario for WR formation was also proposed by Bisnovatyi-Kogan & Nadyozhin (1972), with significant mass loss on the red supergiant phase. They show that large inverse density gradient appears in the outer layers of a 30 M_{\odot} star in this phase due to supercritical luminosity, resulting in powerful and rapid mass loss and formation of a stripped helium core.

In the last decade, Paczyński's hypothesis has again been discussed as a mechanism for the formation of WR stars. Analysing the spatial positions of LBV and WR stars, Nathan Smith (Smith, 2016, 2019; Smith & Tombleson, 2015) con-




Fig. 1: Comparison of spectra of J013340.19+303134.5 (WC4), J013352.43+304351.7 (WN8) and HD 14947 (O4.5If) as illustration of significant differences in intensities of spectral lines for O stars and different sub-types of WR stars.

cluded that both LBVs and WRs could be the products of binary evolution, LBVs being mass-gainers, and WRs – mass-donors, or stripped components. This idea was critically discussed by Humphreys et al. (2016) and Davidson et al. (2016). Now we may confidently conclude only that we do not have enough data for statistical analysis. Modern evolutionary codes allow detailed calculations of binary system evolution (see for example Götberg et al. (2018); Laplace et al. (2021)), thus making it possible to compare theoretical predictions for stripped stars with properties of real WR stars (Götberg et al., 2018). Our current understanding is that binary interaction scenario cannot properly describe all observed WR stars, but most probably takes place in a part of them (see Shenar (2024) and references therein).

Improving the statistics of binarity among WR stars is a crucial task. Among the first researchers who started to work on it was Virpi S. Niemelä, who determined the stellar masses in binary systems and discovered and analyzed the spectroscopic orbits of many binary systems with WR components (see for example her works Niemela (1995, 2001)). Nowadays this work is being continued, both through spectroscopy (Chené et al., 2022) and using high-resolution imaging (Deshmukh et al., 2024).

3. CURRENT PROBLEMS

3.1. Search for new WR stars

In our Galaxy there are 679 WR stars currently discovered²⁾ (Rosslowe & Crowther, 2015), while the theory based on current Milky Way star formation rate and duration of WR phase predicts the numbers of WR stars ~ 1200 (Rosslowe & Crowther, 2015). It means that significant part of WR stars are still not discovered. WR stars in the Galaxy align with spiral arms and the regions of star formation. Thus, the discovery of new WR stars through optical observations has been significantly limited due to the presence of dust extinction. Although the probability of finding a WR star during optical spectroscopic surveys remains (Maíz Apellániz et al., 2016; Zhang et al., 2020), the majority of discoveries of new WR stars now happens through the observations in IR range. For example, Mauerhan et al. (2011) identified 60 Galactic WR stars: candidates were selected using the photometry from *Spitzer* and Two Micron All Sky Survey (2MASS) surveys and confirmed by near-IR spectroscopy. Shara et al. (2012) expanded the list by adding 71 more stars, also preliminary selected from J and K band 2MASS photometry.

Moreover, the search for new WR stars is complicated by the effect of spectral mimicry – low mass [WR] objects masquerading as classical WR stars. Thanks to the results of *Gaia* mission (Gaia Collaboration et al., 2016) we now have reliable estimations of distances, and it helps to quickly understand the nature of individual objects and separate low-mass evolved stars from massive stars, and, moreover, to find the objects of unusual origin among low mass stars. For example, based on *Gaia* distance estimation Gvaramadze et al. (2019) demonstrated that IRAS 00500+6713 is a product of merging of two white dwarfs, while the star has WO-type spectrum. Counterexample is PMR 5 star classified as [WN6] by Morgan et al. (2003) that was recently reclassified as a classical WN.

Systematic search for WR stars in nearby galaxies is ongoing since 1970-s. In 1972 first catalog of WR candidate stars located in M33 galaxy was published (Wray & Corso, 1972). Twenty-five objects were selected based on photometry in three narrow-band filters (one filter is centered on He II 4686 – the strongest emission line in WN-type WRs, one is centered on C III-IV ≈ 4650 – the strongest

²⁾ Galactic Wolf-Rayet Catalogue accessible at




Fig. 2: HR diagram and evolutionary tracks for the low metallicity massive stars (Z = 0.002) from Georgy et al. (2013) (left panel) and Grasha et al. (2021) (right panel) evolutionary models. Geneva tracks do not produce massive WR stars at such low metallicities, while Stromlo tracks show rather different evolutionary path and produce WR stars with $M_* > 70 M_{\odot}$.

line in WC-type WRs, and one is centered on continuum). This technique of selection of WR candidates for following spectroscopy has proven itself successfully and is still widely used (Massey & Conti, 1983; Massey & Johnson, 1998; Neugent & Massey, 2011; Neugent et al., 2012). As of today, 211 WR are known in M33 galaxy (Massey et al., 2016) and 173 are in M31 (Neugent & Massey, 2023). Search for new WR and their following investigations are important for understanding the impact of metallicity on mass-loss rate.

3.2. Wolf-Rayet stars in low metallicity environment

As the opacity of stellar wind depends on the metallicity, mass-loss rates of massive stars decrease with Z. It means that in low metallicity environment single massive stars are unable to lose enough material through the stellar wind to become WR star. Thus, the mass loss on pre-WR stages is of fundamental importance for the final fate of massive stars, and may influence e.g. the formation of long-duration Gamma Ray Bursts (GRBs) or the yields from early stellar generations.

Metallicity impacts overall evolutionary behaviour of massive stars in a complicated way. Georgy et al. (2013) calculated evolutionary tracks for stars in a wide range of initial masses at low metallicity. According to their calculations, metal-poor stars become colder after the end of hydrogen burning in the core and move along the track to the right side of the diagram. Unlike solar metallicity stars, metal-poor ones no longer return to the left side of the diagram (see Figure 2), even when the effects of stellar rotation are taken into account in the models. On the other hand, evolutionary tracks behave differently if the models implement Galactic Concordance abundances instead of solar, scaled-solar, or alpha-element enhanced abundances (Grasha et al., 2021) – the stars with the initial mass of $M_* > 70 M_{\odot}$ with rotation $V/V_{\rm crit} = 0.2$ after the main sequence move to the right and then return to the left to the WR stars region, in contrast to predictions of Georgy et al. (2013).

Yarovova et al. (2023) used evolutionary tracks from Georgy et al. (2013) and Grasha et al. (2021) for the study of emission-line stellar-like object in the nearby low-metallicity ($Z \sim 0.1 Z_{\odot}$) dwarf galaxy NGC 4068. They found the best agreement between the modelled and observed spectra for the model assuming ionization by low-metallicity WR star of $M_* \approx 80 M_{\odot}$ mass, ionizing the nebula through the strong wind and enriching the interstellar medium with nitrogen. Thus, parameters of the object favor Grasha et al. (2021) predictions. Therefore, understanding of mass loss rate requires complex approach that includes both studies of massive stars in different metallicites, and studies of present-day chemical abundances.

3.3. Wolf–Rayet stars and red supergiants

Isolated massive stars with initial masses in 8 $M_{\odot} \lesssim M_{*} \lesssim 20 M_{\odot}$ interval undergo the transition to Red Supergiants (RSG) after hydrogen in the core is exhausted. For them RSG is the final stage before Type-II supernova (SNII) explosion. More massive stars (40 $M_{\odot} \lesssim M_* < 90 M_{\odot}$) evolve through LBV phase before reaching WR phase, and finally explode as SN Type Ib/c. The evolution of stars with 25 $M_{\odot} \lesssim M_* \lesssim 35 M_{\odot}$ initial masses is more complicated. Observed population of supernovae Type II progentions in the local Universe shows that they are produced by the stars with initial masses $M_* \lesssim 18 M_{\odot}$ (Smartt, 2015), and it suggests that more massive stars should either directly collapse to black holes, or evolve leftwards in HR diagram, thus becoming Wolf-Rayet stars. It is supported by the luminosity distributions of cool supergiants in Large and Small Magellanic Clouds (Davies et al., 2018) that overlaps with those of apparently-single WR stars in both galaxies, thus suggesting a changing evolutionary sequence of massive stars with increasing initial mass. At the same time, between the RSG and WR regions in HR diagram lies the region of *low luminosity LBVs* and yellow hypergiants (YHGs) (Figure 3). Therefore, the evolution of 25 $M_{\odot} \leq M_{*} \leq 35 M_{\odot}$




Fig. 3: Hertzsprung–Russell (HR) diagram (luminosities versus effective temperatures). Color horizontal lines show evolutionary tracks, with solid parts showing the phase when hydrogen burns in the stellar core. The tracks are taken from Ekström et al. (2012) (Ekström et al., 2012). ZAMS is zero age Main Sequence.

initial mass may look like that:

$$O \to RSG \to YHG(?) \to low \ LBV \to WR \ (for \ M_* \approx 25 - 35 \ M_{\odot})$$

It means that the star with initial mass about 30 M_{\odot} should pass through all four stages – do such objects really exist?

Groh et al. (2013) demonstrated that the stars with $20-25 \text{ M}_{\odot}$ initial masses are unstable on *low luminosity LBV* stage, and explode as core-collapse supernovae (SN Type IIL/b). Prior to the explosion they show the spectrum similar to S Doradus during its minimum of brightness (WN11h spectral type). Maryeva et al. (2020) have recently found a Galactic object, Wray 15-906, with characteristics typical of a low luminosity LBV and the spectrum looking like WN11h. It was identified via the detection of its infrared circumstellar shell (of ≈ 2 pc in diameter) with the Wide-field Infrared Survey Explorer (WISE) and the Herschel Space Observatory. Maryeva et al. (2020) found that Wray 15-906 is a relatively low-luminosity, $\log(L * / L_{\odot}) \approx 5.4$, star with a temperature of 25 ± 2 kK, with position in HR diagram corresponding to post-red supergiant with initial mass of $\approx 25 M_{\odot}$. Its spectrum and properties are consistent with theoretical predictions of Groh et al. (2013). Thus, the monitoring of this object, and the search for similar ones, may improve our understanding of the link between WRs and RSGs.

4. CONCLUSION

Wolf-Rayet stars were discovered more than 150 years ago, and since then they were targets of numerous studies that combined astrophysical insights with advancements in atomic physics, development of numerical methods, and instrumental progress. Nowadays we understand quite well what Wolf-Rayet stars are, and they are no longer a hot topic in astrophysics. However they still are keystones for many directions of research, from theory of stellar atmospheres up to stellar populations in other galaxies and gravitational waves. Through the existence of the Wolf-Rayet phenomenon WR stars are linked, besides massive stars, with low mass evolved stars.

We expect new progress in understanding the Wolf-Rayet stars and massloss of massive stars in general from the upcoming advances in the instrumentation. For example, SITELLE imaging Fourier transform spectrometer at Canada-France-Hawaii telescope (Drissen et al., 2019) may provide spatially resolved line ratios and kinematics of WR nebulae thus hinting us on ionizing flux from the stars, and their mass loss history. James Webb Space Telescope will allow studying dust shells around WR stars (Lau et al., 2022), and to assess their role in enrichment of interstellar medium with organic compounds and carbonaceous dust. Spectropolarimetric surveys of WR and monitoring of variability of linear polarization are also still topical.

Acknowledgements I would like to thank the organizers of the conference for their kind invitation to present this overview. This research received funding from the European Union's Framework Programme for Research and Innovation Horizon 2020 (2014–2020) under the Marie Skłodowska-Curie Grant Agreement No. 823734 (POEMS project). The Astronomical Institute in Ondřejov is supported by the project RVO:67985815.

REFERENCES

Ambartsumian V. A., 1933, Izvestiya Glavnoj Astronomicheskoj Observatorii v Pulkove, 13

Atanackovic-Vukmanovic O., 2004, Serbian Astronomical Journal, 169, 1

- Auer L. H., Mihalas D., 1969, ApJ, 158, 641
- Beals C. S., 1929, MNRAS, 90, 202
- Beals C. S., 1933, The Observatory, 56, 196
- Bestenlehner J. M., et al., 2014, A&A, 570, A38
- Bisnovatyi-Kogan G. S., Nadyozhin D. K., 1972, Ap&SS, 15, 353
- Bohr N., 1913, Nature, 92, 231
- Bowen I. S., 1928, ApJ, 67, 1
- Bowen I. S., 1934, PASP, 46, 146
- Cannon C. J., 1973a, J. Quant. Spec. Radiat. Transf., 13, 627
- Cannon C. J., 1973b, ApJ, 185, 621
- Chandrasekhar S., 1934, MNRAS, 94, 522
- Chené A.-N., Mahy L., Gosset E., St-Louis N., Dsilva K., Manick R., 2022, MN-RAS, 516, 1022
- Conti P. S., 1975, Memoires of the Societe Royale des Sciences de Liege, 9, 193
- Conti P. S., 1984, in Maeder A., Renzini A., eds, IAU Symposium Vol. 105, Observational Tests of the Stellar Evolution Theory. p. 233
- Conti P. S., 2015, in Hamann W.-R., Sander A., Todt H., eds, Wolf-Rayet Stars. pp 347–350
- Conti P. S., Massey P., 1989, ApJ, 337, 251
- Crowther P. A., 2007, ARA&A, 45, 177
- Crowther P. A., Hillier D. J., Smith L. J., 1995a, A&A, 293, 172
- Crowther P. A., Hillier D. J., Smith L. J., 1995b, A&A, 293, 403
- Crowther P. A., Smith L. J., Hillier D. J., Schmutz W., 1995c, A&A, 293, 427
- Crowther P. A., Smith L. J., Hillier D. J., 1995d, A&A, 302, 457
- Crowther P. A., Smith L. J., Willis A. J., 1995e, A&A, 304, 269
- Crowther P. A., De Marco O., Barlow M. J., 1998, MNRAS, 296, 367

- Crowther P. A., Schnurr O., Hirschi R., Yusof N., Parker R. J., Goodwin S. P., Kassim H. A., 2010, MNRAS, 408, 731
- Davidson K., Humphreys R. M., Weis K., 2016, arXiv e-prints, p. arXiv:1608.02007
- Davies B., Crowther P. A., Beasor E. R., 2018, MNRAS, 478, 3138
- Deshmukh K., et al., 2024, arXiv e-prints, p. arXiv:2409.15212
- Drissen L., et al., 2019, MNRAS, 485, 3930
- Ekström S., et al., 2012, A&A, 537, A146
- Fowler A., 1912, MNRAS, 73, 62
- Gaia Collaboration et al., 2016, A&A, 595, A1
- Gal-Yam A., et al., 2014, Nature, 509, 471
- Gamow G., 1943, ApJ, 98, 500
- Georgy C., et al., 2013, A&A, 558, A103
- Götberg Y., de Mink S. E., Groh J. H., Kupfer T., Crowther P. A., Zapartas E., Renzo M., 2018, A&A, 615, A78
- Gräfener G., Vink J. S., de Koter A., Langer N., 2011, A&A, 535, A56
- Grasha K., Roy A., Sutherland R. S., Kewley L. J., 2021, arXiv e-prints, p. arXiv:2101.01197
- Groh J. H., Meynet G., Ekström S., 2013, A&A, 550, L7
- Groh J. H., Meynet G., Ekström S., Georgy C., 2014, A&A, 564, A30
- Gvaramadze V. V., Gräfener G., Langer N., Maryeva O. V., Kniazev A. Y., Moskvitin A. S., Spiridonova O. I., 2019, Nature, 569, 684
- Hainich R., et al., 2014, A&A, 565, A27
- Hainich R., Pasemann D., Todt H., Shenar T., Sander A., Hamann W. R., 2015, A&A, 581, A21
- Hamann W. R., 1985, A&A, 148, 364
- Hamann W. R., 1986, A&A, 160, 347

- Hamann W. R., Gräfener G., 2003, A&A, 410, 993
- Hamann W. R., Gräfener G., Liermann A., 2006, A&A, 457, 1015
- Hillier D. J., 1990, A&A, 231, 116
- Hillier D. J., Miller D. L., 1998, ApJ, 496, 407
- Huggins W., Huggins M., 1890, Proceedings of the Royal Society of London Series I, 49, 33
- Humphreys R. M., Davidson K., 1994, PASP, 106, 1025
- Humphreys R. M., Weis K., Davidson K., Gordon M. S., 2016, ApJ, 825, 64
- Kosirev N. A., 1934, MNRAS, 94, 430
- Laplace E., Justham S., Renzo M., Götberg Y., Farmer R., Vartanyan D., de Mink S. E., 2021, A&A, 656, A58
- Lau R. M., et al., 2022, Nature Astronomy, 6, 1308
- Liermann A., Hamann W. R., Oskinova L. M., Todt H., Butler K., 2010, A&A, 524, A82
- Maíz Apellániz J., et al., 2016, ApJS, 224, 4
- Marcolino W. L. F., Hillier D. J., de Araujo F. X., Pereira C. B., 2007, ApJ, 654, 1068
- Martins F., Genzel R., Hillier D. J., Eisenhauer F., Paumard T., Gillessen S., Ott T., Trippe S., 2007, A&A, 468, 233
- Martins F., Schaerer D., Marques-Chaves R., Upadhyaya A., 2023, A&A, 678, A159
- Maryeva O., Viotti R. F., Koenigsberger G., Calabresi M., Rossi C., Gualandi R., 2019, Galaxies, 7, 79
- Maryeva O. V., Gvaramadze V. V., Kniazev A. Y., Berdnikov L. N., 2020, MN-RAS, 498, 5093
- Maryeva O., Abdulkarimova A., Karpov S., Moiseev A., Oparin D., 2024, MN-RAS, 527, 11925
- Massey P., Conti P. S., 1983, ApJ, 273, 576

- Massey P., Johnson O., 1998, ApJ, 505, 793
- Massey P., Neugent K. F., Smart B. M., 2016, AJ, 152, 62
- Mauerhan J. C., Van Dyk S. D., Morris P. W., 2011, AJ, 142, 40
- Morgan D. H., Parker Q. A., Cohen M., 2003, MNRAS, 346, 719
- Neugent K. F., Massey P., 2011, ApJ, 733, 123
- Neugent K. F., Massey P., 2023, AJ, 166, 68
- Neugent K. F., Massey P., Georgy C., 2012, ApJ, 759, 11
- Niemela V. S., 1995, in van der Hucht K. A., Williams P. M., eds, IAU Symposium Vol. 163, Wolf-Rayet Stars: Binaries; Colliding Winds; Evolution. p. 223
- Niemela V., 2001, in Revista Mexicana de Astronomia y Astrofísica Conference Series. pp 23–26
- Paczyński B., 1967, Acta Astron., 17, 355
- Paczyński B., 1973, in Bappu M. K. V., Sahade J., eds, IAU Symposium Vol. 49, Wolf-Rayet and High-Temperature Stars. p. 143
- Pickering E. C., 1891, Astronomische Nachrichten, 127, 1
- Pickering E. C., 1897, ApJ, 5, 92
- Pickering E. C., 1901, Harvard College Observatory Circular, 55, 1
- Polcaro V. F., et al., 2016, AJ, 151, 149
- Pritzkuleit M., 2020, Master's thesis, University of Potsdam, Germany
- Roberts M. S., 1962, AJ, 67, 79
- Robotti N., 1983, Historical Studies in the Physical Sciences, 14, 123
- Rosslowe C. K., Crowther P. A., 2015, MNRAS, 447, 2322
- Rublev S. V., 1965, Soviet Ast., 9, 274
- Rublev S. V., 1970, Astrofizicheskie Issledovaniia Izvestiya Spetsial'noj Astrofizicheskoj Observatorii, 1, 25
- Rublev S. V., 1972a, Astrofizicheskie Issledovaniia Izvestiya Spetsial'noj Astrofizicheskoj Observatorii, 4, 3

- Rublev S. V., 1972b, Astrofizicheskie Issledovaniia Izvestiya Spetsial'noj Astrofizicheskoj Observatorii, 4, 18
- Rublev S. V., 1975, in Sherwood V. E., Plaut L., eds, IAU Symposium Vol. 67, Variable Stars and Stellar Evolution. p. 259
- Rybicki G. B., 1972, in Line Formation in the Presence of Magnetic Fields. p. 145
- Sahade J., 1958, The Observatory, 78, 79
- Sander A., Hamann W. R., Todt H., 2012, A&A, 540, A144
- Sander A., Todt H., Hainich R., Hamann W. R., 2014, A&A, 563, A89
- Sander A. A. C., Hamann W. R., Todt H., Hainich R., Shenar T., Ramachandran V., Oskinova L. M., 2019, A&A, 621, A92
- Shara M. M., Faherty J. K., Zurek D., Moffat A. F. J., Gerke J., Doyon R., Artigau E., Drissen L., 2012, AJ, 143, 149
- Shenar T., 2024, arXiv e-prints, p. arXiv:2410.04436
- Smartt S. J., 2015, PASA, 32, e016
- Smith L. F., 1968, MNRAS, 138, 109
- Smith N., 2016, MNRAS, 461, 3353
- Smith N., 2019, MNRAS, 489, 4378
- Smith N., Tombleson R., 2015, MNRAS, 447, 598
- Smith L. F., Shara M. M., Moffat A. F. J., 1990, ApJ, 358, 229
- Smith L. J., Crowther P. A., Prinja R. K., 1994, A&A, 281, 833
- Smith L. F., Shara M. M., Moffat A. F. J., 1996, MNRAS, 281, 163
- Sobolev V. V., 1947, Moving Envelopes of Stars.. Leningrad: Leningrad State University Press; (in Russian), doi:10.4159/harvard.9780674864658
- Sobolev V. V., 1960, Moving Envelopes of Stars. English edition: Cambridge, MA: Harvard University Pres, doi:10.4159/harvard.9780674864658
- Swings P., 1942, ApJ, 95, 112
- Todt H., 2009, PhD thesis, University of Potsdam, Germany

- Underhill A. B., 1968, ARA&A, 6, 39
- Vink J. S., 2015, in Hamann W.-R., Sander A., Todt H., eds, Wolf-Rayet Stars. pp 133–138 (arXiv:1510.00227), doi:10.48550/arXiv.1510.00227
- Westerlund B. E., Rodgers A. W., 1959, The Observatory, 79, 132
- Westerlund B. E., Smith L. F., 1964, MNRAS, 128, 311
- Wolf C. J. E., Rayet G., 1867, Academie des Sciences Paris Comptes Rendus, 65, 292
- Wray J. D., Corso G. J., 1972, ApJ, 172, 577
- Wright W. H., 1918, Publications of Lick Observatory, 13, 191
- Yarovova A. D., Egorov O. V., Moiseev A. V., Maryeva O. V., 2023, MNRAS, 518, 2256
- Zhang W., et al., 2020, ApJ, 902, 62
- Skoda P., Podsztavek O., Tvrdík P., 2020, A&A, 643, A122
- van der Hucht K. A., Conti P. S., Lundstrom I., Stenholm B., 1981, Space Sci. Rev., 28, 227