

Inertial primal-dual dynamics with Hessian-driven damping and Tikhonov regularization for convex-concave bilinear saddle point problems

Xiangkai Sun¹ · Liang He¹ · Xian-Jun Long¹

Received: date / Accepted: date

Abstract This paper deals with a second-order primal-dual dynamical system with Hessian-driven damping and Tikhonov regularization terms in connection with a convex-concave bilinear saddle point problem. We first obtain a fast convergence rate of the primal-dual gap along the trajectory generated by the dynamical system, and provide some integral estimates. Then, based on the setting of the parameters involved, we demonstrate that both the convergence rate of the primal-dual gap and the strong convergence of the trajectory can be achieved simultaneously. Furthermore, we evaluate the performance of the proposed system using two numerical examples.

Keywords Saddle point problems · Primal-dual dynamical system · Tikhonov regularization · Strong convergence · Hessian-driven damping

Mathematics Subject Classification (2000) 34D05 · 37N40 · 46N10 · 90C25

1 Introduction

In recent years, there has been substantial interest in studying the close link between second-order continuous-time dynamical systems and optimization algorithms. On the one hand, by developing the asymptotic properties of second-order dynamical systems, one can efficiently solve optimization problems. On the other hand, it provides thoughtful insights to understand the existing optimization algorithms from a continuous-time perspective. Furthermore, an appropriate time discretization of second-order dynamical systems can potentially lead to inertial optimization algorithms that inherit the convergence properties of the original dynamical systems. We refer the readers to [1–7] for more details.

Xiangkai Sun (✉)
sunxk@ctbu.edu.cn

Liang He
liangheee@126.com

Xian-Jun Long
xianjunlong@ctbu.edu.cn

¹Chongqing Key Laboratory of Statistical Intelligent Computing and Monitoring, College of Mathematics and Statistics, Chongqing Technology and Business University, Chongqing 400067, China.

For most of the studies in this area, we observe that second-order dynamical systems only controlled by viscous damping may exhibit undesired oscillations when it is used to solving optimization problems. However, the oscillations of trajectories can be mitigated by virtue of a geometric damping that is driven by the Hessian of the objective functions. Therefore, for solving optimization problems, the research directions have focused on investigating second-order dynamical systems with Hessian-driven damping. Nowadays, there have been a wide variety of works devoted to second-order dynamical systems with Hessian-driven damping for finding the solution set of the unconstrained convex optimization problem:

$$\min_{x \in \mathbb{R}^n} f(x), \quad (1)$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a twice continuously differentiable convex function.

In order to solve problem (1), Alvarez et al. [8] first introduce the second-order dynamical inertial system with constant Hessian-driven damping:

$$\ddot{x}(t) + \alpha \dot{x}(t) + \gamma \nabla^2 f(x(t)) \dot{x}(t) + \nabla f(x(t)) = 0, \quad (2)$$

where α and γ are positive parameters. This system (2) involves a constant viscous damping coefficient α and a Hessian-driven damping coefficient γ , which effectively reduce the oscillations associated with the Heavy-ball with friction system. Subsequently, Attouch et al. [9] propose a more general second-order dynamical system as follows:

$$\ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \gamma \nabla^2 f(x(t)) \dot{x}(t) + \nabla f(x(t)) = 0, \quad (3)$$

where $\frac{\alpha}{t}$ is the asymptotic vanishing viscous damping coefficient, and $\gamma > 0$ is the constant Hessian-driven damping coefficient. They show that as time t approaches infinity, the fast convergence rate of the objective function value along the trajectory is $\mathcal{O}\left(\frac{1}{t^2}\right)$. To further accelerate the fast convergence rate, by using a time scaling technique, Attouch et al. [10] propose the following second-order dynamical system with a more general Hessian-driven damping coefficient:

$$\ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \gamma(t) \nabla^2 f(x(t)) \dot{x}(t) + \beta(t) \nabla f(x(t)) = 0, \quad (4)$$

where $t_0 > 0$, $\gamma : [t_0, +\infty) \rightarrow [0, +\infty)$ is the Hessian-driven damping function and $\beta : [t_0, +\infty) \rightarrow [0, +\infty)$ is the time scaling function. They show that the convergence rate of the objective function value along the trajectory is $\mathcal{O}\left(\frac{1}{t^2 \beta(t)}\right)$.

In the quest for obtaining strong convergence results of solution trajectories, Boţ et al. [11] consider the following Tikhonov regularized second-order dynamical system with asymptotic vanishing viscous damping and constant Hessian-driven damping:

$$\ddot{x}(t) + \frac{\alpha}{t} \dot{x}(t) + \gamma \nabla^2 f(x(t)) \dot{x}(t) + \nabla f(x(t)) + \epsilon(t)x(t) = 0, \quad (5)$$

where $\epsilon : [t_0, +\infty) \rightarrow [0, +\infty)$ is the Tikhonov regularization function. They show that it inherits the fast convergence properties of the dynamical system (3). Furthermore, they establish the strong convergence of trajectory generated by the dynamical system (5) to the minimum norm solution of problem (1). Moreover, inspired by the fast convergence rates attached to the heavy ball method in the strongly convex case [1], Attouch et al. [12] propose the following Tikhonov regularized second-order dynamical system where the viscous damping coefficient is proportional to the square root of the Tikhonov regularization parameter:

$$\ddot{x}(t) + \delta \sqrt{\epsilon(t)} \dot{x}(t) + \gamma \nabla^2 f(x(t)) \dot{x}(t) + \nabla f(x(t)) + \epsilon(t)x(t) = 0, \quad (6)$$

where $\delta > 0$. They establish the fast convergence rate of the objective function along the trajectory generated by the system (6), and show that the trajectory converges strongly to the minimal norm solution of problem (1). For a bibliographical review of the development of Tikhonov regularized second-order dynamical systems with Hessian-driven damping for unconstrained optimization problem (1), we refer the readers to the survey paper [13] and some recent papers [14–18] for references.

It is important to mention that all the aforementioned papers concentrate on studying the unconstrained optimization problem (1). Recently, more researchers have extended second-order primal-dual dynamical systems from unconstrained optimization problem (1) to the following constrained optimization problem with linear equality constraints:

$$\begin{cases} \min_{x \in \mathbb{R}^n} f(x) \\ \text{s.t. } Kx = b, \end{cases} \quad (7)$$

where $K : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a continuous linear operator and $b \in \mathbb{R}^m$. In order to solve problem (7), many successful treatment of second-order primal-dual dynamical systems with viscous damping, time scaling, and extrapolation coefficients have been obtained from several different perspectives, see, e.g. [7, 19–25].

However, in contrast to unconstrained optimization problem (1), we observe that there exist only few papers devoting to the investigation of second-order primal-dual dynamical systems with Hessian driven damping for solving problem (7). He et al. [26] propose the following second-order plus first-order primal-dual dynamical system with general Hessian-driven damping:

$$\begin{cases} \ddot{x}(t) + \frac{\alpha}{t}\dot{x}(t) + \gamma(t)\frac{d}{dt}(\nabla_x \widehat{\mathcal{L}}(x(t), \lambda(t))) + \beta(t)\nabla_x \widehat{\mathcal{L}}(x(t), \lambda(t)) = 0, \\ \dot{\lambda}(t) - \eta(t)\nabla_\lambda \widehat{\mathcal{L}}\left(x(t) + \frac{t}{\alpha-1}\dot{x}(t), \lambda(t)\right) = 0, \end{cases} \quad (8)$$

where $\alpha > 1$, $\frac{\alpha}{t}$ is the asymptotic vanishing viscous damping coefficient, $\gamma : [t_0, +\infty) \rightarrow [0, +\infty)$ is the Hessian-driven damping function, $\beta, \eta : [t_0, +\infty) \rightarrow [0, +\infty)$ are the time scaling functions, and $\frac{t}{\alpha-1}$ is the extrapolation parameter. Here, $\widehat{\mathcal{L}}$ is the known Lagrangian function defined by

$$\widehat{\mathcal{L}}(x, \lambda) := f(x) + \langle \lambda, Kx - b \rangle.$$

They show that as time t approaches infinity, the fast convergence rates of the Lagrangian residual, the objective residual and the feasibility violation along the trajectory generated by the dynamical system (8) are $\mathcal{O}\left(\frac{1}{t\eta(t)}\right)$. More recently, Csetnek and László [27] associate to (7) the Tikhonov regularized second-order primal-dual dynamical system:

$$\begin{cases} \ddot{x}(t) + \frac{\alpha}{t^q}\dot{x}(t) + \beta t^q \frac{d}{dt}(\nabla_x \widehat{\mathcal{L}}(x(t), y(t))) + (1 - \frac{\gamma}{t^s})\nabla_x \widehat{\mathcal{L}}(x(t), y(t)) + \frac{c}{t^{2q+s}}x(t) = 0, \\ \ddot{y}(t) + \frac{\alpha}{t^q}\dot{y}(t) - \beta t^q \frac{d}{dt}(\nabla_y \widehat{\mathcal{L}}(x(t), y(t))) - (1 - \frac{\gamma}{t^s})\nabla_y \widehat{\mathcal{L}}(x(t), y(t)) + \frac{c}{t^{2q+s}}y(t) = 0, \end{cases} \quad (9)$$

where $\alpha > 1$, $q, \beta, \gamma, s, c > 0$ and $0 < q + s < 1$. Under mild conditions on the parameters involved, they obtain strong convergence of the trajectories to the minimal norm primal-dual solution, as well as fast convergence rates for the feasibility measure, velocities and objective function values along the trajectory generated by the dynamical system (9).

On the other hand, the problem (7) is closely related to the convex-concave saddle-point problem, it is therefore natural to employ the second-order primal-dual dynamical system approaches to solve convex-concave saddle point problems. However, it is worth mentioning that there is no research on Tikhonov regularized second-order primal-dual dynamical

systems with Hessian-driven damping for solving convex-concave saddle point problems although there are some preliminary results available in [28–32] on some characterizations of second-order primal-dual dynamical systems with viscous damping, time scaling and extrapolation coefficients. This is also the main motivation for the study of second-order primal-dual dynamical systems with Hessian-driven damping and Tikhonov regularization for solving convex-concave saddle point problems.

Motivated by the works reported in [17, 29, 30], in this paper, we consider the following bilinearly coupled convex-concave saddle point problem:

$$\min_{x \in \mathbb{R}^n} \max_{y \in \mathbb{R}^m} \mathcal{L}(x, y) := f(x) + \langle Kx, y \rangle - g(y), \quad (10)$$

where $K : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a continuous linear operator, $\langle \cdot, \cdot \rangle$ represents the standard inner product of vectors, and both $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^m \rightarrow \mathbb{R}$ are twice continuously differentiable convex functions. For problem (10), we introduce the following Tikhonov regularized second-order primal-dual dynamical system, which consists of slow viscous damping, extrapolation, Hessian-driven damping and time scaling,

$$\begin{cases} \ddot{x}(t) + \frac{\alpha}{t^q} \dot{x}(t) + \gamma \frac{d}{dt} (\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))) + t^s \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)) = 0, \\ \ddot{y}(t) + \frac{\alpha}{t^q} \dot{y}(t) - \gamma \frac{d}{dt} (\nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))) - t^s \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)) = 0, \end{cases} \quad (11)$$

where $\alpha > 1$, $0 < q < 1$, $\gamma > 0$ and $s > 0$, $\mathcal{L}_t : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ is the augmented Lagrangian saddle function (see (15) for details), $\frac{\alpha}{t^q}$ is the slow viscous damping parameter, γ is the constant Hessian-driven damping parameter, t^s is the time scaling parameter, and $\theta(t)$ is the extrapolation parameter defined as

$$\theta(t) := \frac{t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q}{(\alpha - 1)(t^{q+s} - \gamma qt^{q-1})}. \quad (12)$$

The contributions of this paper can be more specifically stated as follows:

- (i) We propose a new second-order primal-dual dynamical system (11) with Hessian-driven damping and Tikhonov regularization for solving convex-concave saddle point problems. In comparison to the dynamical systems introduced in [29, 30, 32], the system (11) incorporates slow viscous damping, extrapolation, time scaling, constant Hessian-driven damping and Tikhonov regularization.
- (ii) Under mild assumptions on the parameters, we show that the primal-dual gap along the trajectories generated by the dynamical system (11) enjoys $\mathcal{O}\left(\frac{1}{t^{2q+s}}\right)$ convergence rate.
- (iii) By appropriately adjusting these parameters, we show that there exists a single regime where both the convergence rate of the primal-dual gap and the strong convergence of trajectories towards the element of minimal norm of the saddle point set can be simultaneously ensured.
- (iv) Through numerical experiments, we demonstrate that the dynamical system (11), controlled by Hessian-driven damping, not only significantly neutralizes the oscillations that occur during iterations, but also may help to accelerate the convergence of values.

The rest of this paper is organized as follows. In Section 2, we recall some basic notations and present some preliminary results. In Section 3, we obtain the fast convergence properties of the primal-dual gap, and give some integral estimate results. In Section 4, we establish a simultaneous result concerning the convergence rate of the augmented primal-dual gap and the strong convergence of the trajectory generated by the dynamical system (11) to the minimal norm solution of the problem (10). In Section 5, we conduct some numerical experiments to illustrate the theoretical results.

2 Preliminaries

Throughout this paper, \mathbb{R}^n is the n -dimensional Euclidean space with inner product and Euclidean norm denoted by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$, respectively. The norm of the Cartesian product $\mathbb{R}^n \times \mathbb{R}^m$ is defined as

$$\|(x, y)\| = \sqrt{\|x\|^2 + \|y\|^2}, \quad \forall x \in \mathbb{R}^n \text{ and } y \in \mathbb{R}^m.$$

For the continuous linear operator $K : \mathbb{R}^n \rightarrow \mathbb{R}^m$. We denote that

$$\|K\| := \max\{\|Kx\| : x \in \mathbb{R}^n \text{ with } \|x\| \leq 1\},$$

and $K^* : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is the adjoint operator of K . For a set $D \subseteq \mathbb{R}^n$, $\text{Proj}_D 0$ denotes the set of points in D that are closest to the origin, where Proj is the projection operator. If D is a closed convex set, $\text{Proj}_D 0$ represents the one with minimal norm in D .

Let $h : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuously differentiable convex function. We say that the gradient of h is Lipschitz continuous on \mathbb{R}^n iff there exists $0 < l < +\infty$ such that

$$\|\nabla h(x_1) - \nabla h(x_2)\| \leq l\|x_1 - x_2\|, \quad \forall x_1, x_2 \in \mathbb{R}^n.$$

Moreover, we say that h is ϵ -strongly convex function with a strong convexity parameter $\epsilon \geq 0$ iff $h - \frac{\epsilon}{2}\|\cdot\|^2$ is a convex function. Clearly, the following gradient inequality holds:

$$\langle \nabla h(x_1), x_2 - x_1 \rangle \leq h(x_2) - h(x_1) - \frac{\epsilon}{2}\|x_1 - x_2\|^2, \quad \forall x_1, x_2 \in \mathbb{R}^n. \quad (13)$$

Now, consider the saddle point problem (10). A pair $(x^*, y^*) \in \mathbb{R}^n \times \mathbb{R}^m$ is said to be a saddle point of the Lagrangian function \mathcal{L} iff

$$\mathcal{L}(x^*, y) \leq \mathcal{L}(x^*, y^*) \leq \mathcal{L}(x, y^*), \quad \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^m. \quad (14)$$

In the sequel, the set of saddle points of \mathcal{L} is denoted by Ω . Moreover, we assume that $\Omega \neq \emptyset$.

For $c > 0$ and $p > 0$, associated with the Lagrangian function \mathcal{L} , we introduce the augmented Lagrangian function $\mathcal{L}_t : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ which defined as

$$\mathcal{L}_t(x, y) := \mathcal{L}(x, y) + \frac{c}{2t^p} (\|x\|^2 - \|y\|^2) = f(x) + \langle Kx, y \rangle - g(y) + \frac{c}{2t^p} (\|x\|^2 - \|y\|^2). \quad (15)$$

Obviously, $\mathcal{L}_t(\cdot, y)$ is $\frac{c}{t^p}$ -strongly convex for any $y \in \mathbb{R}^m$, and $\mathcal{L}_t(x, \cdot)$ is $\frac{c}{t^p}$ -strongly concave for any $x \in \mathbb{R}^n$. This means that \mathcal{L}_t admits a unique saddle point $(x_t, y_t) \in \mathbb{R}^n \times \mathbb{R}^m$, i.e.,

$$\mathcal{L}_t(x_t, y) \leq \mathcal{L}_t(x_t, y_t) \leq \mathcal{L}_t(x, y_t), \quad \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^m. \quad (16)$$

Naturally, the system of primal-dual optimality conditions reads

$$\begin{cases} 0 = \nabla_x \mathcal{L}_t(x_t, y_t) = \nabla f(x_t) + \frac{c}{t^p} x_t + K^* y_t, \\ 0 = \nabla_y \mathcal{L}_t(x_t, y_t) = -\nabla g(y_t) - \frac{c}{t^p} y_t + K x_t. \end{cases} \quad (17)$$

Here, $\nabla_x \mathcal{L}_t$ and $\nabla_y \mathcal{L}_t$ denote the gradients of \mathcal{L}_t with respect to the first argument and the second argument, respectively.

The following important property will be used in the sequel.

Lemma 2.1 [7, Lemma 2.3] *Let $(\bar{x}^*, \bar{y}^*) \in \text{Proj}_\Omega 0$ and (x_t, y_t) be the saddle point of \mathcal{L}_t . Then,*

- (i) $\lim_{t \rightarrow +\infty} \|(x_t, y_t) - (\bar{x}^*, \bar{y}^*)\| = 0$ and $\|(x_t, y_t)\| \leq \|(\bar{x}^*, \bar{y}^*)\|$ for all $t \geq t_0$.
- (ii) $\|(\dot{x}_t, \dot{y}_t)\| \leq \frac{2}{t}\|(x_t, y_t)\| \leq \frac{2}{t}\|(\bar{x}^*, \bar{y}^*)\|$ for all $t \geq t_0$.

Remark 2.1 Let $z_t = (x_t, y_t)$ and $\bar{z}^* = (\bar{x}^*, \bar{y}^*)$. By Lemma 2.1, it is easy to see that

$$\max \{ \|x_t\|^2, \|y_t\|^2 \} \leq \|\bar{z}^*\|^2, \quad (18)$$

and

$$\max \{ \|\dot{x}_t\|^2, \|\dot{y}_t\|^2 \} \leq \|\dot{z}_t\|^2 \leq \frac{p^2}{t^2} \|\bar{z}^*\|^2. \quad (19)$$

3 Fast convergence of the values

In this section, by using the tools and techniques provided by the Lyapunov analysis, we establish the fast convergence rates of primal-dual gap along the trajectory generated by the dynamical system (11). Moreover, we also give some integral estimates.

Theorem 3.1 *Let $(x, y) : [t_0, +\infty) \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ be a global solution of the dynamical system (11). Suppose that $s > \max\{0, p-2\}$ and $p-q-s-1 > 0$. Then, for any $(x^*, y^*) \in \Omega$, the trajectory $(x(t), y(t))_{t \geq t_0}$ is bounded and*

$$\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{2q+s}}\right), \quad \text{as } t \rightarrow +\infty,$$

$$\|\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty,$$

$$\|\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty,$$

$$\int_{t_0}^{+\infty} t^q (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) dt < +\infty,$$

and

$$\int_{t_0}^{+\infty} t^{2q+s} (\|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 + \|\nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\|^2) dt < +\infty.$$

Proof For any fixed $(x^*, y^*) \in \Omega$, we introduce the energy function $\mathcal{E} : [t_0, +\infty) \rightarrow \mathbb{R}$ as follows:

$$\mathcal{E}(t) = \mathcal{E}_1(t) + \mathcal{E}_2(t) + \mathcal{E}_3(t), \quad (20)$$

where

$$\mathcal{E}_1(t) = (t^{2q+s} - 2\gamma q t^{2q-1} + \gamma t^q) (\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t)) + \frac{c}{2t^p} (\|x(t)\|^2 + \|y(t)\|^2)),$$

$$\mathcal{E}_2(t) = \frac{1}{2} \|(\alpha - 1)(x(t) - x^*) + t^q (\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|^2 + \frac{\alpha-1}{2} (1 - qt^{q-1}) \|x(t) - x^*\|^2,$$

and

$$\mathcal{E}_3(t) = \frac{1}{2} \|(\alpha - 1)(y(t) - y^*) + t^q (\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)))\|^2 + \frac{\alpha-1}{2} (1 - qt^{q-1}) \|y(t) - y^*\|^2.$$

By $0 < q < 1$ and $s > 0$, there exists $t_1 \geq t_0$ such that $t^{2q+s} - 2\gamma q t^{2q-1} + \gamma t^q \geq 0$ and $1 - qt^{q-1} \geq 0$, for all $t \geq t_1$. Thus, $\mathcal{E}(t) \geq 0$, for all $t \geq t_1$.

We now analyze the time derivative of $\mathcal{E}(t)$.

Clearly,

$$\begin{aligned} \dot{\mathcal{E}}_1(t) = & ((2q+s)t^{2q+s-1} - 2\gamma q(2q-1)t^{2q-2} + \gamma qt^{q-1}) \left(\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t)) \right. \\ & \left. + \frac{c}{2t^p} (\|x(t)\|^2 + \|y(t)\|^2) \right) + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \left(\langle \nabla_x \mathcal{L}(x(t), y^*), \dot{x}(t) \rangle \right. \\ & \left. - \langle \nabla_y \mathcal{L}(x^*, y(t)), \dot{y}(t) \rangle - \frac{cp}{2t^{p+1}} (\|x(t)\|^2 + \|y(t)\|^2) \right. \\ & \left. + \frac{c}{t^p} (\langle x(t), \dot{x}(t) \rangle + \langle y(t), \dot{y}(t) \rangle) \right). \end{aligned} \quad (21)$$

Now, set

$$\mu(t) := (\alpha - 1)(x(t) - x^*) + t^q \left(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)) \right).$$

Then,

$$\begin{aligned} \dot{\mu}(t) = & (\alpha - 1)\dot{x}(t) + qt^{q-1} \left(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)) \right) \\ & + t^q \left(\ddot{x}(t) + \gamma \frac{d}{dt} \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)) \right). \end{aligned}$$

This, together with the first equality of (11), follows that

$$\dot{\mu}(t) = (qt^{q-1} - 1)\dot{x}(t) + (\gamma qt^{q-1} - t^{q+s}) \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)). \quad (22)$$

Therefore,

$$\begin{aligned} \langle \mu(t), \dot{\mu}(t) \rangle = & (\alpha - 1)(qt^{q-1} - 1) \langle x(t) - x^*, \dot{x}(t) \rangle + t^q (qt^{q-1} - 1) \|\dot{x}(t)\|^2 \\ & + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), x(t) - x^* \rangle \\ & - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\ & + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 \\ = & (\alpha - 1)(qt^{q-1} - 1) \langle x(t) - x^*, \dot{x}(t) \rangle + t^q (qt^{q-1} - 1) \|\dot{x}(t)\|^2 \\ & + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle \nabla_x \mathcal{L}(x(t), y^*) + \frac{c}{t^p} x(t), x(t) - x^* \rangle \\ & + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), x(t) - x^* \rangle \\ & - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}(x(t), y^*) + \frac{c}{t^p} x(t), \dot{x}(t) \rangle \\ & - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\ & + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2, \end{aligned} \quad (23)$$

where the second equality follows from

$$\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)) = \nabla_x \mathcal{L}(x(t), y^*) + K^*(y(t) - y^* + \theta(t)\dot{y}(t)) + \frac{c}{t^p} x(t).$$

Since $\mathcal{L}(\cdot, y^*) + \frac{c}{2t^p} \|\cdot\|^2$ is ϵ -strong convex function with strong convexity parameter $\epsilon = \frac{c}{t^p}$, it follows from (13) that

$$\begin{aligned} & \langle \nabla_x \mathcal{L}(x(t), y^*) + \frac{c}{t^p} x(t), x^* - x(t) \rangle \\ \leq & \mathcal{L}(x^*, y^*) - \mathcal{L}(x(t), y^*) + \frac{c}{2t^p} (\|x^*\|^2 - \|x(t)\|^2 - \|x^* - x(t)\|^2). \end{aligned} \quad (24)$$

Note that there exists $t_2 \geq t_1$ such that

$$\gamma qt^{q-1} - t^{q+s} \leq 0, \quad \forall t \geq t_2. \quad (25)$$

Combining (25) with (23) and (24), we have

$$\begin{aligned}
& \langle \mu(t), \dot{\mu}(t) \rangle \\
& \leq (\alpha - 1)(qt^{q-1} - 1) \langle x(t) - x^*, \dot{x}(t) \rangle + t^q(qt^{q-1} - 1) \|\dot{x}(t)\|^2 \\
& \quad + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y^*) + \frac{c}{2t^p} (\|x(t)\|^2 - \|x^*\|^2 \right. \\
& \quad \left. + \|x(t) - x^*\|^2) \right) + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), x(t) - x^* \rangle \\
& \quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}(x(t), y^*) + \frac{c}{t^p} x(t), \dot{x}(t) \rangle \\
& \quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\
& \quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2,
\end{aligned} \tag{26}$$

for all $t \geq t_2$. Moreover,

$$\begin{aligned}
\frac{d}{dt} \left(\frac{\alpha-1}{2} (1 - qt^{q-1}) \|x(t) - x^*\|^2 \right) &= \frac{(\alpha-1)q(1-q)}{2} t^{q-2} \|x(t) - x^*\|^2 \\
& \quad + (\alpha - 1)(1 - qt^{q-1}) \langle x(t) - x^*, \dot{x}(t) \rangle.
\end{aligned} \tag{27}$$

By (26) and (27), we obtain

$$\begin{aligned}
\dot{\mathcal{E}}_2(t) &= \langle \mu(t), \dot{\mu}(t) \rangle + \frac{d}{dt} \left(\frac{\alpha-1}{2} (1 - qt^{q-1}) \|x(t) - x^*\|^2 \right) \\
& \leq t^q(qt^{q-1} - 1) \|\dot{x}(t)\|^2 + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}(x(t), y^*) \right. \\
& \quad \left. - \mathcal{L}(x^*, y^*) + \frac{c}{2t^p} (\|x(t)\|^2 - \|x^*\|^2) \right) + \left(\frac{(\alpha-1)q(1-q)}{2} t^{q-2} \right. \\
& \quad \left. + \frac{(\alpha-1)c}{2} (\gamma qt^{q-1-p} - t^{q+s-p}) \right) \|x(t) - x^*\|^2 \\
& \quad + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), x(t) - x^* \rangle \\
& \quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}(x(t), y^*) + \frac{c}{t^p} x(t), \dot{x}(t) \rangle \\
& \quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K^*(y(t) - y^* + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\
& \quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2,
\end{aligned} \tag{28}$$

for all $t \geq t_2$.

Similarly, for all $t \geq t_2$,

$$\begin{aligned}
\dot{\mathcal{E}}_3(t) & \leq t^q(qt^{q-1} - 1) \|\dot{y}(t)\|^2 + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}(x^*, y(t)) \right. \\
& \quad \left. - \mathcal{L}(x^*, y^*) + \frac{c}{2t^p} (\|y(t)\|^2 - \|y^*\|^2) \right) + \left(\frac{(\alpha-1)q(1-q)}{2} t^{q-2} \right. \\
& \quad \left. + \frac{(\alpha-1)c}{2} (\gamma qt^{q-1-p} - t^{q+s-p}) \right) \|y(t) - y^*\|^2 \\
& \quad - (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle K(x(t) - x^* + \theta(t)\dot{x}(t)), y(t) - y^* \rangle \\
& \quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_y \mathcal{L}(x^*, y(t)) - \frac{c}{t^p} y(t), \dot{y}(t) \rangle \\
& \quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K(x(t) - x^* + \theta(t)\dot{x}(t)), \dot{y}(t) \rangle \\
& \quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\|^2.
\end{aligned} \tag{29}$$

Combining (12), (21), (28) and (29), we get

$$\begin{aligned}
\dot{\mathcal{E}}(t) & \leq a(t)(\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t))) + b(t) (\|x(t)\|^2 + \|y(t)\|^2) \\
& \quad + \frac{\alpha-1}{2} (q(1-q)t^{q-2} + c\gamma qt^{q-1-p} - ct^{q+s-p}) (\|x(t) - x^*\|^2 + \|y(t) - y^*\|^2) \\
& \quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \Delta(t) + t^q(qt^{q-1} - 1) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) \\
& \quad + \frac{(\alpha-1)c}{2} (t^{q+s-p} - \gamma qt^{q-1-p}) (\|x^*\|^2 + \|y^*\|^2),
\end{aligned} \tag{30}$$

for all $t \geq t_2$, where

$$a(t) := (2q + s)t^{2q+s-1} - 2\gamma q(2q - 1)t^{2q-2} + \gamma qt^{q-1} + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}), \tag{31}$$

$$b(t) := \frac{c}{2} \left((2q + s - p)t^{2q+s-p-1} - 2\gamma q(2q - 1 - p)t^{2q-p-2} + \gamma(q - p)t^{q-p-1} + (\alpha - 1)(\gamma qt^{q-p-1} - t^{q+s-p}) \right), \quad (32)$$

and

$$\Delta(t) := \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 + \|\nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\|^2. \quad (33)$$

Now, we analyze the coefficients on the right hand side of (30).

Firstly, from $0 < q < 1$ and $s > 0$, we have

$$q + s > 2q + s - 1 > q - 1 > 2q - 2. \quad (34)$$

This, together with $\alpha > 1$, yields that there exists $t_3 \geq t_2$ such that

$$a(t) \leq 0 \text{ and } b(t) \leq 0, \quad \forall t \geq t_3. \quad (35)$$

Secondly, it follows from $s > \max\{0, p - 2\}$ that there exists $t_4 \geq t_3$ such that

$$\frac{\alpha - 1}{2} (q(1 - q)t^{q-2} + c\gamma qt^{q-1-p} - ct^{q+s-p}) \leq 0, \quad \forall t \geq t_4. \quad (36)$$

Thirdly, we can deduce that there exists $t_5 \geq t_4$ such that

$$t^q(qt^{q-1} - 1) \geq 0, \quad \forall t \geq t_5. \quad (37)$$

Finally, by (14), we have $\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t)) \geq 0$. This, together with (25), (35), (36), and (37), follows that

$$\dot{\mathcal{E}}(t) \leq \frac{(\alpha - 1)c}{2} t^{q+s-p} (\|x^*\|^2 + \|y^*\|^2), \quad \forall t \geq t_5.$$

Integrating it from t_5 to t , we obtain

$$\mathcal{E}(t) \leq \mathcal{E}(t_5) + \int_{t_5}^t \frac{(\alpha - 1)c}{2} \omega^{q+s-p} (\|x^*\|^2 + \|y^*\|^2) d\omega, \quad \forall t \geq t_5.$$

Since $p - q - s - 1 > 0$, there exists $M_1 \geq 0$ such that

$$\mathcal{E}(t) \leq M_1, \quad \forall t \geq t_5.$$

Taking (20) into account, it is clear that the trajectory $(x(t), y(t))$,

$$\|(\alpha - 1)(x(t) - x^*) + t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|$$

and

$$\|(\alpha - 1)(y(t) - y^*) + t^q(\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)))\|$$

are bounded for all $t \geq t_5$. Note that

$$\begin{aligned} & \|t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|^2 \\ & \leq 2\|(\alpha - 1)(x(t) - x^*)\|^2 + 2\|(\alpha - 1)(x(t) - x^*) + t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|^2. \end{aligned}$$

Then, from the boundedness of $\|(\alpha - 1)(x(t) - x^*) + t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|$ and the trajectory $(x(t), y(t))$, we can deduce that

$$\|\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty.$$

Similarly, we can show that

$$\|\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty.$$

Moreover, by (20), we have

$$(t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) (\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t))) \leq \mathcal{E}(t) \leq M_1, \quad \forall t \geq t_5,$$

which implies

$$\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{2q+s}}\right), \quad \text{as } t \rightarrow +\infty.$$

On the other hand, it is clear from (30) that

$$\begin{aligned} & \dot{\mathcal{E}}(t) + t^q(1 - qt^{q-1})(\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \gamma t^q(t^{q+s} - \gamma qt^{q-1})\Delta(t) \\ & \leq \frac{(\alpha-1)c}{2} t^{q+s-p} (\|x^*\|^2 + \|y^*\|^2), \quad \forall t \geq t_5. \end{aligned}$$

Since $0 < q < 1$, there exists $t_6 \geq t_5$ such that

$$\frac{1}{2}t^q \leq t^q(1 - qt^{q-1}) \quad \text{and} \quad \frac{1}{2}\gamma t^{2q+s} \leq \gamma t^q(t^{q+s} - \gamma qt^{q-1}), \quad \forall t \geq t_6.$$

Then, for any $t \geq t_6$,

$$\dot{\mathcal{E}}(t) + \frac{1}{2}t^q(\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \frac{1}{2}\gamma t^{2q+s}\Delta(t) \leq \frac{(\alpha-1)c}{2} t^{q+s-p} (\|x^*\|^2 + \|y^*\|^2),$$

Integrating it from t_6 to t , we have

$$\begin{aligned} & \mathcal{E}(t) + \int_{t_6}^t \frac{1}{2}\omega^q(\|\dot{x}(\omega)\|^2 + \|\dot{y}(\omega)\|^2)d\omega + \int_{t_6}^t \frac{1}{2}\gamma\omega^{2q+s}\Delta(\omega)d\omega \\ & \leq \mathcal{E}(t_6) + \int_{t_6}^t \frac{(\alpha-1)c}{2}\omega^{q+s-p} (\|x^*\|^2 + \|y^*\|^2) d\omega, \quad \forall t \geq t_6. \end{aligned}$$

Combining this with $p - q - s - 1 > 0$ and noting that $\mathcal{E}(t) \geq 0$, for all $t \geq t_6$, we have

$$\int_{t_0}^{+\infty} t^q (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) dt < +\infty \quad \text{and} \quad \int_{t_0}^{+\infty} t^{2q+s}\Delta(t)dt < +\infty.$$

The proof is complete. \square

Remark 3.1 In [6, Theorem 4], László provided a Tikhonov regularized second-order dynamical system with slow vanishing damping for solving the problem (1), and obtained that

$$f(x(t)) - \min f = \mathcal{O}\left(\frac{1}{t^{2q}}\right), \quad \text{and} \quad \|\dot{x}(t)\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty.$$

However, by taking $K \equiv 0$, $g \equiv 0$, $\gamma \equiv 0$ and $\theta \equiv 0$ in the primal-dual dynamical system (11), it follows from Theorem 3.1 that

$$f(x(t)) - \min f = \mathcal{O}\left(\frac{1}{t^{2q+s}}\right), \quad \text{and} \quad \|\dot{x}(t)\| = \mathcal{O}\left(\frac{1}{t^q}\right), \quad \text{as } t \rightarrow +\infty.$$

Therefore, Theorem 3.1 can be viewed as an extension of [6, Theorem 4], which deals with the unconstrained convex optimization problem (1), to the convex-concave bilinear saddle point problem (10).

4 Strong convergence of trajectory to the minimal norm solution

In this section, by using Tikhonov regularization techniques, we establish a simultaneous result concerning both the convergence rate of the primal-dual gap and the strong convergence of trajectory generated by dynamical system (11). In order to do this, we first state the following lemma, which gives the estimates on a newly defined energy function.

Lemma 4.1 *Let $(x, y) : [t_0, +\infty) \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ be a global solution of the dynamical system (11) and (x_t, y_t) be the saddle point of \mathcal{L}_t . Consider the energy function $\hat{\mathcal{E}} : [t_0, +\infty) \rightarrow \mathbb{R}$ defined as*

$$\hat{\mathcal{E}}(t) = \hat{\mathcal{E}}_1(t) + \hat{\mathcal{E}}_2(t) + \hat{\mathcal{E}}_3(t), \quad (38)$$

where

$$\begin{cases} \hat{\mathcal{E}}_1(t) = (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))), \\ \hat{\mathcal{E}}_2(t) = \frac{1}{2} \|(\alpha - 1)(x(t) - x_t) + t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|^2 \\ \quad + \frac{\alpha-1}{2} (1 - qt^{q-1}) \|x(t) - x_t\|^2, \\ \hat{\mathcal{E}}_3(t) = \frac{1}{2} \|(\alpha - 1)(y(t) - y_t) + t^q(\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)))\|^2 \\ \quad + \frac{\alpha-1}{2} (1 - qt^{q-1}) \|y(t) - y_t\|^2. \end{cases}$$

Then, there exists $t_1 \geq t_0$ such that

$$\begin{aligned} & \frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \\ & \leq \left(M(t^{q+s} - 2\gamma qt^{q-1} + \gamma) + a(t) \right) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))) \\ & \quad - \frac{c\gamma}{2} (t^{2q+s-p-1} - 2\gamma qt^{2q-p-2} + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2) - (\|x_t\|^2 + \|y_t\|^2) \\ & \quad + l(t) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) + \left(\frac{3M-1}{2} t^q + qt^{2q-1} \right) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) \\ & \quad + \left(\frac{3M}{2} \gamma^2 t^q + \gamma^2 qt^{2q-1} - \frac{1}{2} \gamma t^{2q+s} \right) \Delta(t) \\ & \quad + \left(\frac{2}{(\alpha-1)c} (\|K\|^2 + \|K^*\|^2) (t^{3q+s+p} - 2\gamma qt^{3q-1+p} + \gamma t^{2q+p}) \right. \\ & \quad \left. + \frac{2(\alpha-1)\alpha}{c} (\alpha t^{p-s-q} - qt^{p-s-1}) + \frac{1}{2} (\alpha-1)^2 t^q + \frac{1}{2} \gamma (\alpha-1)^2 t^{-s} \right) (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2), \end{aligned}$$

for all $t \geq t_1$. Here, M is arbitrarily positive constant,

$$l(t) := \left(\frac{M(\alpha-1)(3\alpha-2)}{2} + \frac{(\alpha-1)c\gamma}{8} \right) t^{-q} - \frac{M(\alpha-1)q}{2} t^{-1} - \frac{(\alpha-1)c}{4} t^{q+s-p} + \frac{(\alpha-1)c\gamma q}{4} t^{q-1-p} \\ - \frac{(\alpha-1)qc}{8\alpha} t^{2q+s-1-p} + \frac{(\alpha-1)q(1-q)}{2} t^{q-2},$$

$a(t)$ and $\Delta(t)$ are defined as (31) and (33), respectively.

Proof Obviously,

$$\begin{aligned} & \frac{1}{2} \|(\alpha - 1)(x(t) - x_t) + t^q(\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)))\|^2 \\ & \leq \frac{3}{2} (\alpha - 1)^2 \|x(t) - x_t\|^2 + \frac{3}{2} t^{2q} \|\dot{x}(t)\|^2 + \frac{3}{2} \gamma^2 t^{2q} \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{2} \|(\alpha - 1)(y(t) - y_t) + t^q(\dot{y}(t) - \gamma \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)))\|^2 \\ & \leq \frac{3}{2} (\alpha - 1)^2 \|y(t) - y_t\|^2 + \frac{3}{2} t^{2q} \|\dot{y}(t)\|^2 + \frac{3}{2} \gamma^2 t^{2q} \|\nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t))\|^2. \end{aligned}$$

Therefore, for any $M > 0$,

$$\begin{aligned} \frac{M}{t^q} \hat{\mathcal{E}}(t) & \leq M(t^{q+s} - 2\gamma qt^{q-1} + \gamma) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))) \\ & \quad + \frac{M}{2} (\alpha - 1) ((3\alpha - 2)t^{-q} - qt^{-1}) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\ & \quad + \frac{3M}{2} t^q (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \frac{3M}{2} \gamma^2 t^q \Delta(t). \end{aligned} \quad (39)$$

Now, we analyze the time derivative of $\hat{\mathcal{E}}(t)$.

Firstly, from (15), it is easy to see that

$$\begin{aligned} \frac{d}{dt}\mathcal{L}_t(x(t), y_t) &= \langle \nabla f(x(t)), \dot{x}(t) \rangle + \frac{c}{t^p} \langle x(t), \dot{x}(t) \rangle - \frac{cp}{2t^{p+1}} \|x(t)\|^2 + \langle K^* y_t, \dot{x}(t) \rangle \\ &\quad + \langle Kx(t), \dot{y}_t \rangle - \langle \nabla g(y_t), \dot{y}_t \rangle - \frac{c}{t^p} \langle y_t, \dot{y}_t \rangle + \frac{cp}{2t^{p+1}} \|y_t\|^2 \\ &= \langle \nabla_x \mathcal{L}_t(x(t), y_t), \dot{x}(t) \rangle + \langle \nabla_y \mathcal{L}_t(x(t), y_t), \dot{y}_t \rangle - \frac{cp}{2t^{p+1}} (\|x(t)\|^2 - \|y_t\|^2). \end{aligned}$$

Similarly, we can show that

$$\frac{d}{dt}\mathcal{L}_t(x_t, y(t)) = \langle \nabla_x \mathcal{L}_t(x_t, y(t)), \dot{x}_t \rangle + \langle \nabla_y \mathcal{L}_t(x_t, y(t)), \dot{y}(t) \rangle - \frac{cp}{2t^{p+1}} (\|x_t\|^2 - \|y(t)\|^2).$$

Consequently,

$$\begin{aligned} \dot{\hat{\mathcal{E}}}_1(t) &= ((2q+s)t^{2q+s-1} - 2\gamma q(2q-1)t^{2q-2} + \gamma qt^{q-1}) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))) \\ &\quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \left(\langle \nabla_x \mathcal{L}_t(x(t), y_t), \dot{x}(t) \rangle + \langle \nabla_y \mathcal{L}_t(x(t), y_t), \dot{y}_t \rangle \right. \\ &\quad \left. - \frac{cp}{2t^{p+1}} (\|x(t)\|^2 - \|y_t\|^2) - \left(\langle \nabla_x \mathcal{L}_t(x_t, y(t)), \dot{x}_t \rangle + \langle \nabla_y \mathcal{L}_t(x_t, y(t)), \dot{y}(t) \rangle \right. \right. \\ &\quad \left. \left. - \frac{cp}{2t^{p+1}} (\|x_t\|^2 - \|y(t)\|^2) \right) \right) \end{aligned} \quad (40)$$

Note that

$$\nabla_y \mathcal{L}_t(x(t), y_t) = \nabla_y \mathcal{L}_t(x_t, y_t) + K(x(t) - x_t) = K(x(t) - x_t) \quad (41)$$

and

$$\nabla_x \mathcal{L}_t(x_t, y(t)) = \nabla_x \mathcal{L}_t(x_t, y_t) + K^*(y(t) - y_t) = K^*(y(t) - y_t), \quad (42)$$

where the last two equalities of (41) and (42) hold due to (17). Therefore, combining (40), (41), and (42), and rearranging the terms, we get

$$\begin{aligned} \dot{\hat{\mathcal{E}}}_1(t) &= ((2q+s)t^{2q+s-1} - 2\gamma q(2q-1)t^{2q-2} + \gamma qt^{q-1}) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))) \\ &\quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \left(\langle \nabla_x \mathcal{L}_t(x(t), y_t), \dot{x}(t) \rangle - \langle \nabla_y \mathcal{L}_t(x_t, y(t)), \dot{y}(t) \rangle \right. \\ &\quad \left. + \langle K(x(t) - x_t), \dot{y}_t \rangle - \langle K^*(y(t) - y_t), \dot{x}_t \rangle \right) - \frac{cp}{2} (t^{2q+s-p-1} - 2\gamma qt^{2q-p-2} \\ &\quad + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2 - (\|x_t\|^2 + \|y_t\|^2)). \end{aligned} \quad (43)$$

Secondly, let

$$\bar{\mu}(t) := (\alpha - 1)(x(t) - x_t) + t^q (\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))).$$

Together with the first equality of (11), it follows that

$$\dot{\bar{\mu}}(t) = (qt^{q-1} - 1)\dot{x}(t) - (\alpha - 1)\dot{x}_t + (\gamma qt^{q-1} - t^{q+s})\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)).$$

Moreover, using a similar argument as given in Theorem 3.1, we can show that there exists $t_1 \geq t_0$ such that for any $t \geq t_1$,

$$\begin{aligned} &\langle \bar{\mu}(t), \dot{\bar{\mu}}(t) \rangle \\ &\leq (\alpha - 1)(qt^{q-1} - 1) \langle x(t) - x_t, \dot{x}(t) \rangle + t^q (qt^{q-1} - 1) \|\dot{x}(t)\|^2 \\ &\quad + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y_t) + \frac{c}{2t^p} \|x(t) - x_t\|^2 \right) \\ &\quad + (\alpha - 1)(\gamma qt^{q-1} - t^{q+s}) \langle K^*(y(t) - y_t + \theta(t)\dot{y}(t)), x(t) - x_t \rangle \\ &\quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}_t(x(t), y_t), \dot{x}(t) \rangle \\ &\quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K^*(y(t) - y_t + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\ &\quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 \\ &\quad + \langle (\alpha - 1)(x(t) - x_t) + t^q (\dot{x}(t) + \gamma \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))), -(\alpha - 1)\dot{x}_t \rangle \end{aligned}$$

and

$$\begin{aligned} & \frac{d}{dt} \left(\frac{\alpha-1}{2} (1-qt^{q-1}) \|x(t) - x_t\|^2 \right) \\ &= \frac{(\alpha-1)q(1-q)}{2} t^{q-2} \|x(t) - x_t\|^2 \\ & \quad + (\alpha-1)(1-qt^{q-1}) (\langle x(t) - x_t, \dot{x}(t) \rangle - \langle x(t) - x_t, \dot{x}_t \rangle). \end{aligned}$$

Therefore, for any $t \geq t_1$,

$$\begin{aligned} \dot{\mathcal{E}}_2(t) &= \langle \dot{\mu}(t), \dot{\mu}(t) \rangle + \frac{d}{dt} \left(\frac{\alpha-1}{2} (1-qt^{q-1}) \|x(t) - x_t\|^2 \right) \\ &\leq t^q (qt^{q-1} - 1) \|\dot{x}(t)\|^2 + (\alpha-1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y_t) \right) \\ &\quad + \left(\frac{(\alpha-1)q(1-q)}{2} t^{q-2} + \frac{(\alpha-1)c}{2} (\gamma qt^{q-1-p} - t^{q+s-p}) \right) \|x(t) - x_t\|^2 \\ &\quad + (\alpha-1)(\gamma qt^{q-1} - t^{q+s}) \langle K^*(y(t) - y_t + \theta(t)\dot{y}(t)), x(t) - x_t \rangle \\ &\quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_x \mathcal{L}_t(x(t), y_t), \dot{x}(t) \rangle \\ &\quad - (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K^*(y(t) - y_t + \theta(t)\dot{y}(t)), \dot{x}(t) \rangle \\ &\quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t))\|^2 \\ &\quad - (\alpha-1)(\alpha - qt^{q-1}) \langle x(t) - x_t, \dot{x}_t \rangle - (\alpha-1)t^q \langle \dot{x}(t), \dot{x}_t \rangle \\ &\quad - (\alpha-1)\gamma t^q \langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), \dot{x}_t \rangle. \end{aligned} \tag{44}$$

Similarly, we can deduce that

$$\begin{aligned} \dot{\mathcal{E}}_3(t) &\leq t^q (qt^{q-1} - 1) \|\dot{y}(t)\|^2 + (\alpha-1)(\gamma qt^{q-1} - t^{q+s}) \left(\mathcal{L}_t(x_t, y_t) - \mathcal{L}_t(x_t, y(t)) \right) \\ &\quad + \left(\frac{(\alpha-1)q(1-q)}{2} t^{q-2} + \frac{(\alpha-1)c}{2} (\gamma qt^{q-1-p} - t^{q+s-p}) \right) \|y(t) - y_t\|^2 \\ &\quad - (\alpha-1)(\gamma qt^{q-1} - t^{q+s}) \langle K(x(t) - x_t + \theta(t)\dot{x}(t)), y(t) - y_t \rangle \\ &\quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle \nabla_y \mathcal{L}_t(x_t, y(t)), \dot{y}(t) \rangle \\ &\quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \langle K(x(t) - x_t + \theta(t)\dot{x}(t)), \dot{y}(t) \rangle \\ &\quad + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \|\nabla_y \mathcal{L}_t((x(t) + \theta(t)\dot{x}(t)), y(t))\|^2 \\ &\quad - (\alpha-1)(\alpha - qt^{q-1}) \langle y(t) - y_t, \dot{y}_t \rangle - (\alpha-1)t^q \langle \dot{y}(t), \dot{y}_t \rangle \\ &\quad + (\alpha-1)\gamma t^q \langle \nabla_y \mathcal{L}_t((x(t) + \theta(t)\dot{x}(t)), y(t)), \dot{y}_t \rangle, \end{aligned} \tag{45}$$

for any $t \geq t_1$.

Finally, combining (12), (43), (44) and (45), we have

$$\begin{aligned} \dot{\mathcal{E}}(t) &\leq a(t) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t))) \\ &\quad - \frac{cp}{2} (t^{2q+s-p-1} - 2\gamma qt^{2q-p-2} + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2) - (\|x_t\|^2 + \|y_t\|^2) \\ &\quad + \frac{\alpha-1}{2} (q(1-q)t^{q-2} + c\gamma qt^{q-1-p} - ct^{q+s-p}) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\ &\quad + t^q (qt^{q-1} - 1) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \gamma t^q (\gamma qt^{q-1} - t^{q+s}) \Delta(t) \\ &\quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) (\langle K(x(t) - x_t), \dot{y}_t \rangle - \langle K^*(y(t) - y_t), \dot{x}_t \rangle) \\ &\quad - (\alpha-1)(\alpha - qt^{q-1}) (\langle x(t) - x_t, \dot{x}_t \rangle + \langle y(t) - y_t, \dot{y}_t \rangle) \\ &\quad - (\alpha-1)t^q (\langle \dot{x}(t), \dot{x}_t \rangle + \langle \dot{y}(t), \dot{y}_t \rangle) \\ &\quad - (\alpha-1)\gamma t^q (\langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), \dot{x}_t \rangle - \langle \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)), \dot{y}_t \rangle), \end{aligned}$$

for any $t \geq t_1$. Here, $a(t)$ and $\Delta(t)$ are defined as (31) and (33), respectively. This, together with (39), yields

$$\begin{aligned}
& \frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \\
& \leq \left(M(t^{q+s} - 2\gamma qt^{q-1} + \gamma) + a(t) \right) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}(x_t, y(t))) \\
& \quad - \frac{c^p}{2} (t^{2q+s-p-1} - 2\gamma qt^{2q-p-2} + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2) - (\|x_t\|^2 + \|y_t\|^2) \\
& \quad + \left(\frac{M}{2}(\alpha - 1) ((3\alpha - 2)t^{-q} - qt^{-1}) + \frac{\alpha-1}{2}(q(1-q)t^{q-2} \right. \\
& \quad \left. + c\gamma qt^{q-1-p} - ct^{q+s-p}) \right) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\
& \quad + \left(\left(\frac{3M}{2} - 1 \right) t^q + qt^{2q-1} \right) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) \\
& \quad + \left(\frac{3M}{2} \gamma^2 t^q + \gamma^2 qt^{2q-1} - \gamma t^{2q+s} \right) \Delta(t) \\
& \quad + (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) (\langle K(x(t) - x_t), \dot{y}_t \rangle - \langle K^*(y(t) - y_t), \dot{x}_t \rangle) \\
& \quad - (\alpha - 1)(\alpha - qt^{q-1}) (\langle x(t) - x_t, \dot{x}_t \rangle + \langle y(t) - y_t, \dot{y}_t \rangle) \\
& \quad - (\alpha - 1)t^q (\langle \dot{x}(t), \dot{x}_t \rangle + \langle \dot{y}(t), \dot{y}_t \rangle) \\
& \quad - (\alpha - 1)\gamma t^q (\langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), \dot{x}_t \rangle - \langle \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)), \dot{y}_t \rangle).
\end{aligned} \tag{46}$$

On the other hand, note that

$$\langle K(x(t) - x_t), \dot{y}_t \rangle \leq \frac{1}{2} \left(\frac{1}{\frac{4}{(\alpha-1)c} t^{q+p} \|K\|^2} \|K\|^2 \|x(t) - x_t\|^2 + \frac{4}{(\alpha-1)c} t^{q+p} \|K\|^2 \|\dot{y}_t\|^2 \right)$$

and

$$\langle K^*(y(t) - y_t), \dot{x}_t \rangle \leq \frac{1}{2} \left(\frac{1}{\frac{4}{(\alpha-1)c} t^{q+p} \|K^*\|^2} \|K^*\|^2 \|y(t) - y_t\|^2 + \frac{4}{(\alpha-1)c} t^{q+p} \|K^*\|^2 \|\dot{x}_t\|^2 \right).$$

Thus,

$$\begin{aligned}
& (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) (\langle K(x(t) - x_t), \dot{y}_t \rangle - \langle K^*(y(t) - y_t), \dot{x}_t \rangle) \\
& \leq \frac{1}{2} (t^{2q+s} - 2\gamma qt^{2q-1} + \gamma t^q) \left(\frac{1}{\frac{4}{(\alpha-1)c} t^{q+p}} (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \right. \\
& \quad \left. + \frac{4}{(\alpha-1)c} t^{q+p} (\|K\|^2 + \|K^*\|^2) (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2) \right) \\
& = \frac{(\alpha-1)c}{8} (t^{q+s-p} - 2\gamma qt^{q-1-p} + \gamma t^{-p}) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\
& \quad + \frac{2}{(\alpha-1)c} (t^{3q+s+p} - 2\gamma qt^{3q-1+p} + \gamma t^{2q+p}) (\|K\|^2 + \|K^*\|^2) (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2).
\end{aligned} \tag{47}$$

In a same manner as above, we can deduce that

$$\begin{aligned}
& -(\alpha - 1)(\alpha - qt^{q-1}) (\langle x(t) - x_t, \dot{x}_t \rangle + \langle y(t) - y_t, \dot{y}_t \rangle) \\
& \leq \frac{1}{2} (\alpha - 1)(\alpha - qt^{q-1}) \left(\frac{1}{4\frac{\alpha}{c} t^{p-s-q}} (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \right. \\
& \quad \left. + 4\frac{\alpha}{c} t^{p-s-q} (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2) \right) \\
& = \frac{(\alpha-1)c}{8} (t^{q+s-p} - \frac{\alpha}{c} t^{2q+s-1-p}) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\
& \quad + \frac{2(\alpha-1)\alpha}{c} (\alpha t^{p-s-q} - qt^{p-s-1}) (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2),
\end{aligned} \tag{48}$$

$$\begin{aligned}
& -(\alpha - 1)t^q (\langle \dot{x}(t), \dot{x}_t \rangle + \langle \dot{y}(t), \dot{y}_t \rangle) \\
& \leq \frac{1}{2} t^q (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \frac{1}{2} (\alpha - 1)^2 t^q (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2),
\end{aligned} \tag{49}$$

and

$$\begin{aligned}
& -(\alpha - 1)\gamma t^q (\langle \nabla_x \mathcal{L}_t(x(t), y(t) + \theta(t)\dot{y}(t)), \dot{x}_t \rangle - \langle \nabla_y \mathcal{L}_t(x(t) + \theta(t)\dot{x}(t), y(t)), \dot{y}_t \rangle) \\
& \leq \frac{1}{2} \gamma t^{2q+s} \Delta(t) + \frac{1}{2} \gamma (\alpha - 1)^2 t^{-s} (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2).
\end{aligned} \tag{50}$$

Combining (47), (48), (49) and (50) with (46), we have

$$\begin{aligned}
& \frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \\
& \leq \left(M(t^{q+s} - 2\gamma q t^{q-1} + \gamma) + a(t) \right) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}(x_t, y(t))) \\
& \quad - \frac{c\beta}{2} (t^{2q+s-p-1} - 2\gamma q t^{2q-p-2} + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2) - (\|x_t\|^2 + \|y_t\|^2) \\
& \quad + \left(\left(\frac{M(\alpha-1)(3\alpha-2)}{2} + \frac{(\alpha-1)c\gamma}{8} \right) t^{-q} - \frac{M(\alpha-1)q}{2} t^{-1} - \frac{(\alpha-1)c}{4} t^{q+s-p} + \frac{(\alpha-1)c\gamma q}{4} t^{q-1-p} \right. \\
& \quad \left. - \frac{(\alpha-1)qc}{8\alpha} t^{2q+s-1-p} + \frac{(\alpha-1)q(1-q)}{2} t^{q-2} \right) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) \\
& \quad + \left(\frac{3M-1}{2} t^q + q t^{2q-1} \right) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) + \left(\frac{3M}{2} \gamma^2 t^q + \gamma^2 q t^{2q-1} - \frac{1}{2} \gamma t^{2q+s} \right) \Delta(t) \\
& \quad + \left(\frac{2}{(\alpha-1)c} (\|K\|^2 + \|K^*\|^2) (t^{3q+s+p} - 2\gamma q t^{3q-1+p} + \gamma t^{2q+p}) \right. \\
& \quad \left. + \frac{2(\alpha-1)\alpha}{c} (\alpha t^{p-s-q} - q t^{p-s-1}) + \frac{1}{2} (\alpha-1)^2 t^q + \frac{1}{2} \gamma (\alpha-1)^2 t^{-s} \right) (\|\dot{x}_t\|^2 + \|\dot{y}_t\|^2),
\end{aligned}$$

for any $t \geq t_1$. The proof is complete. \square

Now, we establish the main result of this section, which demonstrates that both the convergence rate of the primal-dual gap and the strong convergence of the trajectory can be achieved simultaneously.

Theorem 4.1 *Let $(x, y) : [t_0, +\infty) \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ be a global solution of the dynamical system (11) and let (x_t, y_t) be the saddle point of \mathcal{L}_t . Suppose that*

$$p - 2q - s < 0, 4q + s + p - 2 < 0, \text{ and } 3q + s - p - 1 < 0.$$

Then, for $(\bar{x}^*, \bar{y}^*) = \text{Proj}_{\Omega} 0$,

$$\lim_{t \rightarrow +\infty} \|(x(t), y(t)) - (\bar{x}^*, \bar{y}^*)\| = 0,$$

and the following statements hold:

(i) If $q + 2p \geq 1$, one has

$$\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) = \mathcal{O}\left(\frac{1}{t^{2-2q-p}}\right), \text{ as } t \rightarrow +\infty,$$

and

$$\|x(t) - x_t\| + \|y(t) - y_t\| = \mathcal{O}\left(\frac{1}{t^{1-2q-\frac{1}{2}(s+p)}}\right), \text{ as } t \rightarrow +\infty.$$

(ii) If $q + 2p < 1$, one has

$$\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) = \mathcal{O}\left(\frac{1}{t^{1-q+p}}\right), \text{ as } t \rightarrow +\infty,$$

and

$$\|x(t) - x_t\| + \|y(t) - y_t\| = \mathcal{O}\left(\frac{1}{t^{\frac{1}{2}(1+p-s)-\frac{3}{2}q}}\right), \text{ as } t \rightarrow +\infty.$$

Proof We start with the energy function $\hat{\mathcal{E}} : [t_0, +\infty) \rightarrow \mathbb{R}$ defined as (38). It is easy to verify that there exists $t'_1 \geq t_0$ such that $\hat{\mathcal{E}}(t) \geq 0$, for all $t \geq t'_1$.

Let $\bar{z}^* = (\bar{x}^*, \bar{y}^*)$. By virtue of (18) and (19), we have

$$\begin{aligned}
& \frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \\
& \leq \left(M (t^{q+s} - 2\gamma q t^{q-1} + \gamma) + a(t) \right) (\mathcal{L}_t(x(t), y_t) - \mathcal{L}(x_t, y(t))) \\
& \quad - \frac{cp}{2} (t^{2q+s-p-1} - 2\gamma q t^{2q-p-2} + \gamma t^{q-p-1}) (\|x(t)\|^2 + \|y(t)\|^2) \\
& \quad + l(t) (\|x(t) - x_t\|^2 + \|y(t) - y_t\|^2) + \left(\frac{3M-1}{2} t^q + q t^{2q-1} \right) (\|\dot{x}(t)\|^2 + \|\dot{y}(t)\|^2) \\
& \quad + \left(\frac{3M}{2} \gamma^2 t^q + \gamma^2 q t^{2q-1} - \frac{1}{2} \gamma t^{2q+s} \right) \Delta(t) + m(t) \|\bar{z}^*\|^2,
\end{aligned} \tag{51}$$

where

$$\begin{aligned}
m(t) &= cp (t^{2q+s-p-1} - 2\gamma q t^{2q-p-2} + \gamma t^{q-p-1}) \\
& \quad + \frac{4p^2}{(\alpha-1)c} (\|K\|^2 + \|K^*\|^2) (t^{3q+s+p-2} - 2\gamma q t^{3q+p-3} + \gamma t^{2q+p-2}) \\
& \quad + \frac{4(\alpha-1)\alpha p^2}{c} (\alpha t^{p-s-q-2} - q t^{p-s-3}) + (\alpha-1)^2 p^2 (t^{q-2} + \gamma t^{-s-2}).
\end{aligned}$$

We analyze the coefficients on the right hand side of the inequality (51).

Clearly,

$$\begin{aligned}
M (t^{q+s} - 2\gamma q t^{q-1} + \gamma) + a(t) &= (M - (\alpha-1)) t^{q+s} + (\alpha - 2M) \gamma q t^{q-1} + (2q+s) t^{2q+s-1} \\
& \quad - 2\gamma q (2q-1) t^{2q-2} + M\gamma.
\end{aligned}$$

Now, set $M < \min\{\frac{1}{3}, \alpha-1\}$. Together with (34), $\alpha > 1$ and $M < \alpha-1$, there exists $t'_2 \geq t'_1$ such that

$$M (t^{q+s} - 2\gamma q t^{q-1} + \gamma) + a(t) \leq 0 \tag{52}$$

and

$$-\frac{cp}{2} (t^{2q+s-p-1} - 2\gamma q t^{2q-p-2} + \gamma t^{q-p-1}) \leq 0, \tag{53}$$

for all $t \geq t'_2$.

Using a similar argument as above, there exists $t'_3 \geq t'_2$ such that

$$\frac{3M-1}{2} t^q + q t^{2q-1} \leq 0 \text{ and } \frac{3M}{2} \gamma^2 t^q + \gamma^2 q t^{2q-1} - \frac{1}{2} \gamma t^{2q+s} \leq 0, \quad \forall t \geq t'_3. \tag{54}$$

Now, we consider the coefficient $l(t)$. From $0 < q < 1$, $s > 0$, $p > 0$, and $p - 2q - s < 0$, we have

$$q + s - p > -q > -1 > q - 2$$

and

$$q + s - p > 2q + s - 1 - p > q - 1 - p.$$

Since $-\frac{(\alpha-1)c}{4} < 0$, it follows that there exists $t'_4 \geq t'_3$ such that

$$l(t) \leq 0, \quad \forall t \geq t'_4. \tag{55}$$

Further, according to (16), we have $\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) \geq 0$. Therefore, combining (52), (53), (54) and (55), and neglecting the nonpositive terms of (51), we obtain that

$$\frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \leq m(t) \|\bar{z}^*\|^2, \quad \forall t \geq t'_4. \tag{56}$$

Now, we analyze the coefficient $m(t)$ of (56). It is easy to verify that

$$\begin{cases} 2q + s - p - 1 > q - p - 1 > 2q - p - 2 \\ 3q + s + p - 2 > -s - 2, \\ 3q + s + p - 2 > 3q + p - 3, \\ 3q + s + p - 2 > 2q + p - 2 > p - s - q - 2 > p - s - 3. \end{cases}$$

Thus, we can deduce that there exists $t'_5 \geq t'_4$ and a constant $C_0 > 0$ such that

$$m(t) \|\bar{z}^*\|^2 \leq C_0(t^{3q+s+p-2} + t^{2q+s-p-1}), \quad \forall t \geq t'_5.$$

Consequently, (56) leads to

$$\frac{M}{t^q} \hat{\mathcal{E}}(t) + \dot{\hat{\mathcal{E}}}(t) \leq C_0(t^{3q+s+p-2} + t^{2q+s-p-1}), \quad \forall t \geq t'_5. \quad (57)$$

Multiplying $e^{\frac{M}{1-q}t^{1-q}}$ on both sides of (57), we have

$$\frac{d}{dt} \left(e^{\frac{M}{1-q}t^{1-q}} \hat{\mathcal{E}}(t) \right) \leq C_0(t^{3q+s+p-2} + t^{2q+s-p-1}) e^{\frac{M}{1-q}t^{1-q}}, \quad \forall t \geq t'_5.$$

Then, integrating it from t'_5 to t , we have

$$e^{\frac{M}{1-q}t^{1-q}} \hat{\mathcal{E}}(t) \leq C_1 + C_0 \int_{t'_5}^t (\omega^{3q+s+p-2} + \omega^{2q+s-p-1}) e^{\frac{M}{1-q}\omega^{1-q}} d\omega, \quad \forall t \geq t'_5, \quad (58)$$

where $C_1 := e^{\frac{M}{1-q}t'^{1-q}} \hat{\mathcal{E}}(t'_5)$.

In what follows, we consider the right hand side of (58). Using the equation (60) introduced in [17], there exist $t'_6 \geq t'_5$, $C_2 > 0$ and $C_3 > 0$ such that

$$t^{3q+s+p-2} e^{\frac{M}{1-q}t^{1-q}} \leq C_2 \frac{d}{dt} \left(t^{4q+s+p-2} e^{\frac{M}{1-q}t^{1-q}} \right) \quad (59)$$

and

$$t^{2q+s-p-1} e^{\frac{M}{1-q}t^{1-q}} \leq C_3 \frac{d}{dt} \left(t^{3q+s-p-1} e^{\frac{M}{1-q}t^{1-q}} \right). \quad (60)$$

for all $t \geq t'_6$.

Obviously, from (59), we have

$$\begin{aligned} \int_{t'_5}^t \omega^{3q+s+p-2} e^{\frac{M}{1-q}\omega^{1-q}} d\omega &\leq c'_1 + C_2 \int_{t'_6}^t \frac{d}{d\omega} \left(\omega^{4q+s+p-2} e^{\frac{M}{1-q}\omega^{1-q}} \right) d\omega \\ &= c'_1 + C_2 \left(t^{4q+s+p-2} e^{\frac{M}{1-q}t^{1-q}} - C_4 \right), \end{aligned} \quad (61)$$

where $c'_1 := \int_{t'_5}^{t'_6} \omega^{3q+s+p-2} e^{\frac{M}{1-q}\omega^{1-q}} d\omega$ and $C_4 := t'^{4q+s+p-2} e^{\frac{M}{1-q}t'^{1-q}}$.

Similarly, we can deduce from (60) that

$$\int_{t'_5}^t \omega^{2q+s-p-1} e^{\frac{M}{1-q}\omega^{1-q}} d\omega \leq c'_2 + C_3 \left(t^{3q+s-p-1} e^{\frac{M}{1-q}t^{1-q}} - C_5 \right), \quad (62)$$

where $c'_2 := \int_{t'_5}^{t'_6} \omega^{2q+s-p-1} e^{\frac{M}{1-q}\omega^{1-q}} d\omega$ and $C_5 := t'^{3q+s-p-1} e^{\frac{M}{1-q}t'^{1-q}}$.

By using (61) and (62), (58) becomes

$$\begin{aligned} &e^{\frac{M}{1-q}t^{1-q}} \hat{\mathcal{E}}(t) \\ &\leq C_1 + C_0 \left(c'_1 + C_2 \left(t^{4q+s+p-2} e^{\frac{M}{1-q}t^{1-q}} - C_4 \right) + c'_2 + C_3 \left(t^{3q+s-p-1} e^{\frac{M}{1-q}t^{1-q}} - C_5 \right) \right) \\ &= C + C_0 \left(C_2 t^{4q+s+p-2} + C_3 t^{3q+s-p-1} \right) e^{\frac{M}{1-q}t^{1-q}}, \end{aligned}$$

where $C := C_1 + C_0 c'_1 + C_0 c'_2$. This means that

$$\hat{\mathcal{E}}(t) \leq \frac{C}{e^{\frac{M}{1-q}t^{1-q}}} + C_6 \left(t^{4q+s+p-2} + t^{3q+s-p-1} \right), \quad \forall t \geq t'_6, \quad (63)$$

where $C_6 := \max\{C_0 C_2, C_0 C_3\}$.

Now, we consider the following two cases:

Case I: $q + 2p \geq 1$. In this case, $0 > 4q + s + p - 2 \geq 3q + s - p - 1$. Consequently, from (63), there exists $t'_7 \geq t'_6$ such that

$$\hat{\mathcal{E}}(t) \leq \frac{C}{e^{\frac{M}{1-q}t^{1-q}}} + C_6 t^{4q+s+p-2}, \quad \forall t \geq t'_7.$$

Taking into account (38), it is easy to show that

$$\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) = \mathcal{O}\left(\frac{1}{t^{2-2q-p}}\right), \quad \text{as } t \rightarrow +\infty,$$

and

$$\|x(t) - x_t\| + \|y(t) - y_t\| = \mathcal{O}\left(\frac{1}{t^{1-2q-\frac{1}{2}(s+p)}}\right), \quad \text{as } t \rightarrow +\infty. \quad (64)$$

Case II: $q + 2p < 1$. In this case, $0 > 3q + s - p - 1 > 4q + s + p - 2$. Consequently, from (63), there exists $t''_7 \geq t'_6$ such that

$$\hat{\mathcal{E}}(t) \leq \frac{C}{e^{\frac{M}{1-q}t^{1-q}}} + C_6 t^{3q+s-p-1}, \quad \forall t \geq t''_7.$$

Taking into account (38) again, it is easy to show that

$$\mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) = \mathcal{O}\left(\frac{1}{t^{1-q+p}}\right), \quad \text{as } t \rightarrow +\infty,$$

and

$$\|x(t) - x_t\| + \|y(t) - y_t\| = \mathcal{O}\left(\frac{1}{t^{\frac{1}{2}(1+p-s)-\frac{3}{2}q}}\right), \quad \text{as } t \rightarrow +\infty. \quad (65)$$

Finally, by using Lemma 2.1, we have $\lim_{t \rightarrow +\infty} \|(x_t, y_t) - (\bar{x}^*, \bar{y}^*)\| = 0$. Together this with (64) and (65), we can verify that

$$\lim_{t \rightarrow +\infty} \|(x(t), y(t)) - (\bar{x}^*, \bar{y}^*)\| = 0.$$

The proof is complete. \square

By Theorem 4.1, we have the following result.

Corollary 4.1 *Let $(x, y) : [t_0, +\infty) \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ be a global solution of the dynamical system (11). Assume that the basic hypotheses of Theorem 4.1 are fulfilled. Then, for $(\bar{x}^*, \bar{y}^*) = \text{Proj}_\Omega 0$, the following results hold.*

(i) *If $q + 2p \geq 1$, one has*

$$\mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{1-2q-\frac{1}{2}(s+p)}} + \frac{1}{t^p}\right), \quad \text{as } t \rightarrow +\infty.$$

(ii) *If $q + 2p < 1$, one has*

$$\mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{\frac{1}{2}(1+p-s)-\frac{3}{2}q}} + \frac{1}{t^p}\right), \quad \text{as } t \rightarrow +\infty.$$

Proof Let (x_t, y_t) be the saddle point of \mathcal{L}_t . Clearly

$$\begin{aligned} \mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) &= \mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y_t) + \mathcal{L}_t(x_t, y_t) - \mathcal{L}_t(x_t, y(t)) \\ &\geq \mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(\bar{x}^*, y_t) + \mathcal{L}_t(x_t, \bar{y}^*) - \mathcal{L}_t(x_t, y(t)) \\ &\geq \mathcal{L}(x(t), y_t) - \mathcal{L}(\bar{x}^*, y_t) + \mathcal{L}(x_t, \bar{y}^*) - \mathcal{L}(x_t, y(t)) \\ &\quad - \frac{c}{2t^p}(\|\bar{x}^*\|^2 + \|\bar{y}^*\|^2), \end{aligned} \quad (66)$$

where the two inequalities are deduced from (16) and (15), respectively. Note that

$$\begin{aligned}
& \mathcal{L}(x(t), y_t) - \mathcal{L}(\bar{x}^*, y_t) + \mathcal{L}(x_t, \bar{y}^*) - \mathcal{L}(x_t, y(t)) \\
&= \mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) + \langle K(x(t) - x_t), y_t - \bar{y}^* \rangle \\
&\quad + \langle K(\bar{x}^* - x_t), y(t) - y_t \rangle \\
&\geq \mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) - \|K\| \|x(t) - x_t\| \|y_t - \bar{y}^*\| \\
&\quad - \|K\| \|\bar{x}^* - x_t\| \|y(t) - y_t\|
\end{aligned} \tag{67}$$

By combining (66) and (67), we have

$$\begin{aligned}
\mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) &\leq \mathcal{L}_t(x(t), y_t) - \mathcal{L}_t(x_t, y(t)) + \frac{c}{2t^p} (\|\bar{x}^*\|^2 + \|\bar{y}^*\|^2) \\
&\quad + \|K\| (\|x(t) - x_t\| \|y_t - \bar{y}^*\| + \|x_t - \bar{x}^*\| \|y(t) - y_t\|)
\end{aligned} \tag{68}$$

Now, we consider the following two cases:

Case I: $q + 2p \geq 1$. In this case, by $4q + s + p - 2 < 0$, we have

$$0 < 1 - 2q - \frac{1}{2}(p + s) < 2 - 4q - s - p < 2 - 2q - p.$$

Combining this with (68), Theorem 4.1 (i) and $\lim_{t \rightarrow +\infty} \|(x_t, y_t) - (\bar{x}^*, \bar{y}^*)\| = 0$, we have

$$\mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{1-2q-\frac{1}{2}(s+p)}} + \frac{1}{t^p}\right), \quad \text{as } t \rightarrow +\infty.$$

Case II: $q + 2p < 1$. In this case, by $3q + s - p - 1 < 0$, we get

$$0 < \frac{1}{2}(1 + p - s) - \frac{3}{2}q < 1 + p - s - 3q < 1 + p - q.$$

Combining this with (68), Theorem 4.1 (ii) and $\lim_{t \rightarrow +\infty} \|(x_t, y_t) - (\bar{x}^*, \bar{y}^*)\| = 0$, it follows that

$$\mathcal{L}(x(t), \bar{y}^*) - \mathcal{L}(\bar{x}^*, y(t)) = \mathcal{O}\left(\frac{1}{t^{\frac{1}{2}(1+p-s)-\frac{3}{2}q}} + \frac{1}{t^p}\right), \quad \text{as } t \rightarrow +\infty.$$

The proof is complete. \square

Remark 4.1 Note that Corollary 4.1 extends [6, Theorem 4], [12, Theorem 2] and [17, Theorem 10] for the unconstrained convex optimization problem (1) to the convex-concave bilinear saddle point problem (10).

5 Numerical experiments

In this section, we validate the obtained theoretical results by two numerical examples. In these experiments, the dynamical system (11) is solved numerically with a Runge-Kutta adaptive method (ode45 in MATLAB version R2019b). All codes are performed on a PC (with 2.30GHz Intel Core i5-8300H and 8GB memory).

Example 5.1 Let $x := (x_1, x_2) \in \mathbb{R}^2$ and $y := (y_1, y_2) \in \mathbb{R}^2$.

$$\min_{x \in \mathbb{R}^2} \max_{y \in \mathbb{R}^2} (mx_1 + nx_2)^2 + (mx_1 + nx_2)(jy_1 + ky_2) - (jy_1 + ky_2)^2, \tag{69}$$

where $m, n, j, k \geq 0$, $f(x) = (mx_1 + nx_2)^2$, $g(y) = (jy_1 + ky_2)^2$, and $K = [mj, mk; nj, nk]$. It is easy to show that the solution set of problem (69) is

$$\{(x, y) \in \mathbb{R}^2 \times \mathbb{R}^2 \mid mx_1 + nx_2 = 0 \text{ and } jy_1 + ky_2 = 0\}.$$

Clearly, $(\bar{x}^*, \bar{y}^*) := (0, 0, 0, 0)^\top$ is the minimal norm solution of problem (69).

In following experiments, the dynamical system (11) is solved on the time interval [1, 20]. We also consider the following two cases:

- $m = 1, n = 2, j = 5, k = 4$.
- $m = 2, n = 3, j = 1, k = 5$.

In the first experiment, for any $(x^*, y^*) \in \Omega$, we consider the influence of Hessian-driven damping on the convergence of the primal-dual gap.

Consider the initial time $t_0 = 1$ and take the following initial conditions:

$$x(t_0) = \begin{bmatrix} 1 \\ 1.5 \end{bmatrix}, \quad y(t_0) = \begin{bmatrix} 1 \\ 1.5 \end{bmatrix}, \quad \dot{x}(t_0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \text{and} \quad \dot{y}(t_0) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}. \quad (70)$$

We set $\alpha = 2.5, q = 0.42, s = 0.05, p = 0.268, c = 10$, which are satisfied with the assumptions in Theorem 4.1.

Figure 1 displays the behavior of $\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t))$ along the trajectory $(x(t), y(t))$ generated by the dynamical system (11) under different settings on $\gamma \in \{0, 0.1, 0.2, 0.3, 0.4\}$.




Fig. 1: Convergence analysis of the dynamical system (11) with different parameter γ

As shown in Figure 1, the numerical results are in agreement with the theoretical claims. In each case, $\mathcal{L}(x(t), y^*) - \mathcal{L}(x^*, y(t))$ converges more smoothly when the system is controlled by Hessian-driven damping. Interestingly, it shows that Hessian-driven damping may help to accelerate the convergence.

In the second experiment, we investigate the strong convergence of the trajectory to the minimal norm solution $(\bar{x}^*, \bar{y}^*) := (0, 0, 0, 0)^\top$ as well as the influence of Hessian-driven damping on the behaviors of trajectories.

Take $\alpha = 2.5, q = 0.42, s = 0.05$ and $p = 0.268$. Under the initial conditions (70), we plot the trajectory $(x(t), y(t))$ generated by the dynamical system (11) in the following three cases:

- $c = 0, \gamma = 0$ (i.e., (11) without Tikhonov regularization and Hessian-driven damping).
- $c = 0, \gamma = 0.2$ (i.e., (11) without Tikhonov regularization).
- $c = 10, \gamma = 0.2$.




Fig. 2: Convergence of trajectories when $m = 1, n = 2, j = 5, k = 4$




Fig. 3: Convergence of trajectories when $m = 2, n = 3, j = 1, k = 5$

As shown in Figures 2 and 3, we can observe that:

- (i) Only when the dynamical system (11) is controlled by Tikhonov regularization, does the trajectory $(x(t), y(t))$ converge to the minimal norm solution $(\bar{x}^*, \bar{y}^*) = (0, 0, 0, 0)^\top$.
- (ii) When the dynamical system (11) is controlled by Hessian-driven damping (i.e., with $\gamma = 0.2$), the trajectory $(x(t), y(t))$ converges more smoothly compared to the case where $\gamma = 0$.

Example 5.2 Let $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$. Consider the following ℓ_2 -regularized problem:

$$\min_{x \in \mathbb{R}^n} \Phi(x) = \frac{1}{2} \|Kx - b\|^2 + \eta \|x\|^2, \quad (71)$$

where $K \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. Equivalently, its saddle point formulation reads as:

$$\min_{x \in \mathbb{R}^n} \max_{y \in \mathbb{R}^m} \eta \|x\|^2 + \langle Kx, y \rangle - \left(\frac{1}{2} \|y\|^2 + \langle b, y \rangle \right).$$

In the following numerical experiment, we are interested in the influence of Hessian-driven damping on the convergence of objective function value of problem (71).

The dynamical system (11) is solved on the time interval $[1, 100]$. All entries of initial conditions, K and b are generated from the standard Gaussian distribution. Moreover, three different dimensional settings are considered:

- $m = 20, n = 50$.
- $m = 50, n = 100$.
- $m = 100, n = 200$.

Take $\eta = 1, \alpha = 3, c = 5, s = 0.4$ and $p = 2.3$. We test the dynamical system (11) under the following settings on parameters q and γ :

- $q = 0.6, \gamma = 0$.

- $q = 0.7, \gamma = 0.$
- $q = 0.8, \gamma = 0.$
- $q = 0.6, \gamma = 0.2.$
- $q = 0.7, \gamma = 0.2.$
- $q = 0.8, \gamma = 0.2.$

Figure 4 depicts the behaviors of the objective error $\Phi(x(t)) - \Phi(x^*)$ along the trajectory $(x(t), y(t))$ generated by the dynamical system (11) in different dimensions.




Fig. 4: Convergence analysis of the dynamical system (11) in different dimensional settings

As shown in Figure 4, in different dimensions, the dynamical system (11) controlled by Hessian-driven damping can significantly reduce oscillations.

6 Conclusion

In this paper, we introduce a Tikhonov regularized second-order primal-dual dynamical system (11) to solve the convex-concave bilinear saddle point problem (10). Compared with the dynamical systems introduced in [29, 30, 32] for solving the convex-concave bilinear saddle point problem (10), the system (11) not only encompasses slow viscous damping, extrapolation and time scaling, but is also controlled by a constant Hessian-driven damping. By using appropriate damping, extrapolation, scaling and Tikhonov regularization coefficients, we establish the fast convergence rates of the values and the strong convergence property of the generated trajectory. Through numerical experiments, we observe that the dynamical system involved in Hessian-driven damping can neutralize the oscillations occurring during iterations.

In the future, a natural direction for research will be to investigate the explicit discretization of the system (11), which leads to an inertial algorithm of gradient type that incorporates a Tikhonov regularization term for solving the convex-concave bilinear saddle point problem (10). Furthermore, based on the results obtained in the present paper, we can also explore the strong convergence properties of the corresponding algorithm.

Funding

This research is supported by the Natural Science Foundation of Chongqing (CSTB2024NSCQ-MSX0651 and CSTB2024NSCQ-MSX1282) and the Team Building Project for Graduate Tutors in Chongqing (yds223010).

Data availability

The authors confirm that all data generated or analysed during this study are included in this article.

Declaration

Conflict of interest No potential conflict of interest was reported by the authors.

References

1. Polyak, B.T.: Some methods of speeding up the convergence of iteration methods. U.S.S.R. Comput. Math. Math. Phys. 4: 1-17 (1964)
2. Su, W., Boyd, S., Candès, E.J.: A differential equation for modeling Nesterov’s accelerated gradient method: theory and insights. J. Mach. Learn. Res. 17, 1-43 (2016)
3. Attouch, H., Chbani, Z., Peypouquet, J., Redont, P.: Fast convergence of inertial dynamics and algorithms with asymptotic vanishing viscosity. Math. Program. 168: 123-175 (2018)
4. Attouch, H., Balhag, A., Chbani, Z., Riahi, H.: Damped inertial dynamics with vanishing Tikhonov regularization: strong asymptotic convergence towards the minimum norm solution. J. Differ. Equ. 311: 29-58 (2022)
5. Attouch, H., Chbani, Z., Fadili, J., Riahi, H.: Fast convergence of dynamical ADMM via time scaling of damped inertial dynamics. J. Optim. Theory Appl. 193: 704-736 (2022)
6. László, S.C.: On the strong convergence of the trajectories of a Tikhonov regularized second order dynamical system with asymptotically vanishing damping. J. Differ. Equ. 362: 355-381 (2023)
7. Chbani, Z., Riahi, H., Battahi, F.: On the simultaneous convergence of values and trajectories of continuous inertial dynamics with Tikhonov regularization to solve convex minimization with affine constraints. Preprint HAL (2024) <https://hal.science/hal-04511296>.
8. Alvarez, F., Attouch, H., Bolte, J., Redont, P.: A second-order gradient-like dissipative dynamical system with Hessian-driven damping: application to optimization and mechanics. Journal de Mathématiques Pures et Appliquées 81: 747-779 (2002)
9. Attouch, H., Peypouquet, J., Redont, P.: Fast convex optimization via inertial dynamics with Hessian driven damping. J. Differ. Equ. 261: 5734-5783 (2016)
10. Attouch, H., Chbani, Z., Fadili, J., Riahi, H.: First-order optimization algorithms via inertial systems with Hessian driven damping. Math. Program. 193: 113-155 (2022)
11. Boţ, R.I., Csetnek, E.R., László, S.C.: Tikhonov regularization of a second order dynamical system with Hessian driven damping. Math. Program. 189: 151-186 (2021)
12. Attouch, H., Balhag, A., Chbani, Z., Riahi, H.: Accelerated gradient methods combining Tikhonov regularization with geometric damping driven by the Hessian. Appl. Math. Optim. 88: 29 (2023)
13. Attouch, H., Boţ, R.I., Csetnek, E.R.: Fast optimization via inertial dynamics with closed-loop damping. J. Eur. Math. Soc. 25: 1985-2056 (2023)
14. Alecsa, C.D., László, S.C.: Tikhonov regularization of a perturbed heavy ball system with vanishing damping. SIAM J. Optim. 31: 2921-2954 (2021)
15. Bagy, A.C., Chbani, Z., Riahi, H.: Strong convergence of trajectories via inertial dynamics combining Hessian driven damping and Tikhonov regularization for general convex minimizations. Numer. Funct. Anal. Optim. 44: 1481-1509 (2023),
16. Csetnek, E.R., Karapetyants, M.A.: Second-order dynamics featuring Tikhonov regularization and time scaling. J. Optim. Theory Appl. 202: 1385-1420 (2024)
17. László, S.C.: Solving convex optimization problems via a second-order dynamical system with implicit Hessian damping and Tikhonov regularization. Comput. Optim. Appl. (2024) <https://doi.org/10.1007/s10589-024-00620-5>
18. Zhong, G.F., Hu, X.Z., Tang, M., Zhong, L.Q.: Fast convex optimization via differential equation with Hessian-driven damping and Tikhonov regularization. J. Optim. Theory Appl. 203: 42-82 (2024)
19. Boţ, R.I., Nguyen, D.K.: Improved convergence rates and trajectory convergence for primal-dual dynamical systems with vanishing damping. J. Differ. Equ. 303: 369-406 (2021)
20. He, X., Hu, R., Fang, Y.P.: Convergence rates of inertial primal-dual dynamical methods for separable convex optimization problems. SIAM J. Control Optim. 59: 3278-3301 (2021)
21. He, X., Hu, R., Fang, Y.P.: “Second-order primal” + “first-order dual” dynamical systems with time scaling for linear equality constrained convex optimization problems. IEEE Trans. Automat. Control. 67: 4377-4383 (2022)
22. He, X., Hu, R., Fang, Y.P.: Inertial primal-dual dynamics with damping and scaling for linearly constrained convex optimization problems. Appl. Anal. 102: 4114-4139 (2023)
23. Hulett, D.A., Nguyen, D.K.: Time rescaling of a primal-dual dynamical system with asymptotically vanishing damping. Appl. Math. Optim. 88: 27 (2023)

24. Zeng, X.L., Lei, J., Chen, J.: Dynamical primal-dual Nesterov accelerated method and its application to network optimization. *IEEE Trans. Autom. Control.* 68: 1760-1767 (2023)
25. Zhu, T.T., Hu, R., Fang, Y.P.: Tikhonov regularized second-order plus first-order primal-dual dynamical systems with asymptotically vanishing damping for linear equality constrained convex optimization problems. *Optimization.* (2024) <https://doi.org/10.1080/02331934.2024.2407515>
26. He, X., Tian, F., Li, A.Q., Fang, Y.P.: Convergence rates of mixed primal-dual dynamical systems with Hessian driven damping. *Optimization.* (2023) <https://doi.org/10.1080/02331934.2023.2253813>
27. Csetnek, E.R., László, S.C.: Strong convergence and fast rates for systems with Tikhonov regularization. *arXiv.2411.17329* (2024)
28. Zeng, X., Dou, L., Chen, J.: Accelerated first-order continuous-time algorithm for solving convex-concave bilinear saddle point problem. *IFAC-PapersOnLine* 53, 7362-7367 (2020)
29. Ding, K.W., Fliege, J., Vuong, P.T.: Fast convergence of the primal-dual dynamical system and corresponding algorithms for a nonsmooth bilinearly coupled saddle point problem. *Comput. Optim. Appl.* (2024) <https://doi.org/10.1007/s10589-024-00626-z>
30. He, X., Hu, R., Fang, Y.P.: A second-order primal-dual dynamical system for a convex-concave bilinear saddle point problem. *Appl. Math. Optim.* 89: 30 (2024)
31. Luo, H.: A continuous perspective on the inertial corrected primal-dual proximal splitting. *arXiv:2405.14098* (2024)
32. Sun, X.K., He, L., Long, X.J.: Tikhonov regularized inertial primal-dual dynamics for convex-concave bilinear saddle point problems. *arXiv:2409.05301* (2024)