

Quantitative particle approximation of nonlinear stochastic Fokker-Planck equations with singular kernel

Josué Knorst, Christian Olivera and Alexandre B. de Souza

Abstract

We derive quantitative estimates for large stochastic systems of interacting particles perturbed by both idiosyncratic and environmental noises, as well as singular kernels. We prove that the (mollified) empirical process converges to the solution of the nonlinear stochastic Fokker-Planck equation. The proof is based on Itô's formula for H_q^1 -valued process, commutator estimates, and some estimations for the regularization of the empirical measure. Moreover, we show that the aforementioned equation admits a unique strong solution in the probabilistic sense. The approach applies to repulsive and attractive kernels.

MSC2010 subject classification: 49N90, 60H30, 60K35.

1 Introduction

The aim of this article is to derive a quantitative convergence result for the following nonlinear stochastic Fokker-Planck equation

$$d\rho_t = \frac{1}{2} \sum_{i,j=1}^d \partial_{ij} \left(\rho_t \sum_{k=1}^d \left(\nu_t^{ik} \nu_t^{jk} + \sigma_t^{ik} \sigma_t^{jk} \right) \right) dt - \nabla \cdot (\rho_t F(\cdot, K * \rho_t)) dt - \nabla \rho_t \cdot \sigma_t dB_t \quad (1)$$

from a stochastic moderately interacting particle system. The equation (1) is considered for arbitrary dimensions $d \geq 1$. We present a method that allows us to derive a fairly general class of such systems as limit dynamics of interacting stochastic many-particle systems. This procedure has been methodically employed in the literature, from the seminal papers [40, 41, 42], (see also [33] and [38]) to the recent works [23] and [43]. Concerning motivations for the model with transport noise, let us mention model reduction, see [37], in addition to other motivations like [17] and [30]. Our results cover several classical models such as the stochastic $2d$ Navier-Stokes equation, which in vorticity form can

be written as in (1) with the Biot-Savart kernel, the stochastic Burgers equation and the parabolic-elliptic Keller-Segel PDE in any dimension $d \geq 1$, which models the phenomenon of chemotaxis.

The derivation of macroscopic models from interacting particle systems is a fascinating and active research topic in mathematics. The basic idea is that, as the number of particles increases to infinity, the macroscopic model can effectively describe the universal properties of particles. In this article, we derive the stochastic PDE from the stochastic moderately interacting particle system on \mathbb{T}^d given by

$$\begin{aligned} dX_t^{i,N} = & F \left(X_t^{i,N}, \frac{1}{N} \sum_{k=1}^N (K * V^N) (X_t^{i,N} - X_t^{k,N}) \right) dt + \nu_t (X_t^{i,N}) dW_t^i \\ & + \sigma_t (X_t^{i,N}) dB_t \end{aligned} \quad (2)$$

where W_t^i and B_t are independent standard \mathbb{T}^d -valued Brownian motions, defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$, the interaction kernel V^N depends on the number of particles $N \in \mathbb{N}$ via the moderate interaction parameter β . Systems of the form (2) are commonly employed in the field of mean-field games, see [6], [7], [18] and [36].

The microscopic *empirical process* of this N -particle system, which is a probability measure on the ambient space \mathbb{T}^d , is given as usual by

$$S_t^N := \frac{1}{N} \sum_{i=1}^N \delta_{X_t^{i,N}}, \quad t \geq 0, \quad (3)$$

where δ_a is the delta Dirac measure concentrated at $a \in \mathbb{T}^d$. Then, $(S_t^N)_{t \geq 0}$ is a measure-valued process associated to the \mathbb{T}^d -valued processes $\{t \mapsto X_t^{i,N}\}_{i=1, \dots, N}$.

The first contribution of this paper is to investigate the large N limit of the dynamical process $(S_t^N)_{t \geq 0}$ in the common noise setting with singular kernels. For that purpose, we introduce the *mollified empirical measure*

$$\rho_t^N := V^N * S_t^N = \int_{\mathbb{T}^d} V^N(\cdot - y) S_t^N(dy),$$

which is more regular than S_t^N , where $V^N = N^\beta V(N^{\frac{\beta}{d}} \cdot)$, $\beta \in (0, 1)$. We quantify the distance between the empirical measure (and the regularized version) and the unique solution to the SPDE (1). The proof is based on Itô's formula for H_q^1 -valued processes, commutator estimates, and stochastic analysis techniques.

The second contribution is to show the well-posedness of the nonlinear stochastic Fokker-Planck equation (1). The main challenge in proving existence and uniqueness results for (1) is the nonlinear term $K * \rho_t$ since this prevents us from directly applying known results in the existing literature on SPDEs, such as those found in [35]. We take advantage of the specific SPDE structure to employ a fixed point argument, e.g., the recent work [31].

1.1 Related literatures

In contrast to interacting particle system, which are driven by idiosyncratic noise, see [3, 8, 9, 16, 25, 24, 26, 27, 32, 39, 43, 47, 52], the literature on common noise and singular kernels remains limited. For systems with uniformly Lipschitz interaction coefficients, [14] established conditional propagation of chaos. The entropy method has recently been explored for systems with common noise, as shown in [49] for the Navier–Stokes equations, [13] for the Hegselmann–Krause model, and [46] for mean-field systems with bounded kernels.

In the following, we will focus only on the results of moderately interacting systems: in [40], Oelschläger introduced and studied moderately interacting particle systems, which are used to obtain a local nonlinear partial differential equation. The idea is to take a radially symmetric function V and then consider the interaction potential $V^N = N^\beta V(N^{\frac{\beta}{a}} \cdot)$, with $\beta \in (0, 1)$, which converges to the Dirac delta at 0 in the sense of distributions. The article [40] is part of a series of works by the author on this subject (see also [41, 42]). The convergence results were improved in [33] and [38]. In the recent paper [19] it was developed a semigroup approach which enables them to show uniform convergence of mollified empirical measures, see [20, 21, 23, 18, 50, 28] for additional applications of this method. Recently, in [43] and [44], the authors developed new quantitative estimates for a class of moderately interacting particle systems using a semigroup approach. This technique was further improved in [29], [34], and [45]. About more advances in moderate particle systems, see, for instance, [10], [11], [15], [12], and [51].

The problem of deriving models represented by SPDE with transport noise and singular kernels from microscopic models of stochastic interacting particles has not been highly investigated in the literature, just with the contributions in [13], [46] and [49]. As we will see in more detail in Section 1.5, our result may be applied to the following well-known models: the stochastic $2d$ Navier-Stokes equation, the stochastic Burgers equation and the stochastic parabolic-elliptic Keller-Segel PDE in any dimension $d \geq 1$.

1.2 Notations

For $d \geq 1$, we denote by $C^k(\mathbb{T}^d)$ the space of k -times differentiable functions on the torus $\mathbb{T}^d \doteq [-\frac{1}{2}, \frac{1}{2}]^d$, where $k \in \mathbb{N} \cup \{\infty\}$. The set of density functions defined on \mathbb{T}^d is denoted by $\mathcal{P}(\mathbb{T}^d)$. The corresponding spaces on \mathbb{R}^d are defined by $C^k(\mathbb{R}^d)$ and $\mathcal{P}(\mathbb{R}^d)$, respectively.

For a measurable space (X, \mathcal{M}, μ) and f a measurable function in it (which we denote by $f \in \mathcal{M}$) we write

$$\langle \mu, f \rangle \doteq \int_X f d\mu$$

for the duality pair involving the measure μ and the function f in this space.

Also, for $a \in [1, \infty)$ the Lebesgue's space is given by

$$L^a = L^a(X) = \left\{ f \in \mathcal{M} \mid \|f\|_a \doteq \left(\int_X |f|^a d\mu \right)^{\frac{1}{a}} < \infty \right\}$$

and if $a = \infty$

$$L^\infty = L^\infty(X) = \left\{ f \in \mathcal{M} \mid \|f\|_\infty \doteq \operatorname{ess\,sup}_{x \in X} |f(x)| < \infty \right\}.$$

In some contexts we will write $\|\cdot\|_a = \|\cdot\|_{L^a(X)}$.

Given $(U, \|\cdot\|_U)$ a Banach space, we denote by U^* its dual and the associated duality pair by $(\cdot, \cdot)_{U, U^*}$. With this notation, for $T > 0$ and denoting \mathcal{L} the set of Bochner measurable functions, the Bochner's space is denoted by

$$L^a U = L^a([0, T]; U) = \left\{ f : [0, T] \rightarrow U, f \in \mathcal{L} \mid \|f\|_{L^a U} \doteq \left(\int_0^T \|f(t)\|_U^a dt \right)^{\frac{1}{a}} < \infty \right\},$$

for $a \in [1, \infty)$ and if $a = \infty$ by

$$L^\infty U = L^\infty([0, T]; U) = \left\{ f : [0, T] \rightarrow U, f \in \mathcal{L} \mid \|f\|_{T, U} \doteq \operatorname{ess\,sup}_{t \in [0, T]} \|f(t)\|_U < \infty \right\}.$$

For the space of tempered distributions on \mathbb{T}^d , we write \mathcal{S}' . So, for $q > 1, n \in \mathbb{R}$, the Bessel potential space reads

$$H_q^n = H_q^n(\mathbb{T}^d) = \left\{ f \in \mathcal{S}' \mid \|f\|_{n, q} \doteq \|(I - \Delta)^{\frac{n}{2}} f\|_q < \infty \right\}.$$

For $\gamma \in (0, 1]$ the Hölder's space on \mathbb{T}^d is given by

$$C^\gamma = C^\gamma(\mathbb{T}^d) = \left\{ f : \mathbb{T}^d \rightarrow \mathbb{R} \mid \|f\|_\gamma \doteq \|f\|_\infty + \sup_{x \neq y \in \mathbb{T}^d} \frac{|f(x) - f(y)|}{|x - y|^\gamma} < \infty \right\}.$$

Let a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$, a Banach space $(U, \|\cdot\|_U)$, a real number $q \in [1, \infty]$, and a stopping time $0 < \tau \leq T$. We denote by \mathcal{X} the set of U -valued, $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted and continuous processes $X = \{X_s\}_{s \in [0, \tau]}$. We define

$$S_{\mathcal{F}}^q([0, \tau]; U) = \left\{ X \in \mathcal{X} \mid \left\| \|X\|_{\tau, U} \right\|_{L^q(\Omega)} < \infty \right\}$$

Also, let \mathcal{Y} denote the set of U -valued predictable processes $Y = \{Y_s\}_{s \in [0, \tau]}$. Then, for $q \in [1, \infty)$ we define

$$L_{\mathcal{F}}^q([0, \tau]; U) = \left\{ Y \in \mathcal{Y} \mid \left\| \|Y\|_{L^q U} \right\|_{L^q(\Omega)} < \infty \right\}.$$

These are Banach spaces endowed with their respective norms.

1.3 Assumptions

(\mathbf{A}^V) The mollifier $V \in C^2(\mathbb{R}^d) \cap \mathcal{P}(\mathbb{R}^d)$ is such that $\text{supp } V \subset \{x \in \mathbb{R}^d; |x| < \frac{1}{2}\}$.

(\mathbf{A}^F) The vector field $F : \mathbb{T}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$ is a bounded Lipschitz continuous function, i.e., there exists $L > 0$ such that

$$\begin{aligned} |F(x, u)| &\leq L, \quad \forall x \in \mathbb{T}^d, u \in \mathbb{R}. \\ |F(x, u) - F(y, v)| &\leq L(|x - y| + |u - v|), \quad \forall x, y \in \mathbb{T}^d, u, v \in \mathbb{R}. \end{aligned}$$

(\mathbf{A}^I) The vector field $F : \mathbb{T}^d \times \mathbb{R} \rightarrow \mathbb{R}$ is given by $F(x, u) = u$.

(\mathbf{A}^K) There exists $q \geq 2$ such that $q > d$, and $\gamma \in (0, 1]$ and $C_K > 0$ such that for any $f \in L^q(\mathbb{T}^d)$, it holds

$$\|K * f\|_\gamma \leq C_K \|f\|_q.$$

(\mathbf{A}^c) The diffusion coefficients ν and $\sigma : [0, T] \times \mathbb{T}^d \rightarrow \mathbb{R}^{d \times d}$ are measurable, and for every $t \in [0, T]$ satisfy

(\mathbf{A}_i^c) There exists $M > 0$ such that

$$\nu_t^{jk}(\cdot), \sigma_t^{jk}(\cdot) \in (C^2(\mathbb{T}^d), \|\cdot\|_{C^2}), \quad \text{for } j, k = 1, \dots, d.$$

$$\text{and} \quad \sum_{j,k=1}^d \left(\|\nu_t^{jk}(\cdot)\|_{C^2} + \|\sigma_t^{jk}(\cdot)\|_{C^2} \right) \leq M.$$

$$(\mathbf{A}_{ii}^c) \quad \sum_{j=1}^d \partial_j \sigma_t^{jk}(x) = 0, \quad \text{for } k = 1, \dots, d; x \in \mathbb{T}^d.$$

(\mathbf{A}_{iii}^c) There exist $\Lambda, \lambda > 0$ such that, for all $x \in \mathbb{T}^d$, $\xi = (\xi^i) \in \mathbb{R}^d$,

$$\Lambda |\xi|^2 \geq \sum_{i,j,k=1}^d \nu_t^{ik}(x) \nu_t^{jk}(x) \xi^i \xi^j \geq \lambda |\xi|^2.$$

1.4 Main theorems.

Regarding the limiting equation (1), we will prove the following existence and uniqueness theorem, based on Krylov's L^q -theory of SPDEs, see [35].

Theorem 1. *Assume (\mathbf{A}^c) and one of the two regimes:*

1. (\mathbf{A}^F) and (\mathbf{A}^K).
2. (\mathbf{A}^I) and (\mathbf{A}^K).

Let $0 \leq \rho_0 \in L^1 \cap H_q^{1-\frac{2}{q}}(\mathbb{T}^d)$, with $\|\rho_0\|_1 = 1$. For each $\kappa > 0$, there exists a $T > 0$ depending on $\kappa, \lambda, q, C_K, L, M$ and d such that if $\|\rho_0\|_q = \kappa$, the SPDE (1) admits a unique nonnegative solution in

$$\mathbb{M} \doteq L_{\mathcal{F}^B}^q([0, T]; H_q^1(\mathbb{T}^d)) \cap S_{\mathcal{F}^B}^\infty([0, T]; L^1 \cap L^q(\mathbb{T}^d)).$$

The proof will be presented in Appendix B.

Theorem 2. Assume (\mathbf{A}^V) , (\mathbf{A}^c) , (\mathbf{A}^F) , and (\mathbf{A}^K) , let T_{max} be the maximal existence time for (1) and fix $T \in (0, T_{max})$. In addition, let the dynamics of the particle system be given by (2) with $\nu = Id$, σ independent of the space and for any $m \geq 1$,

$$\sup_{N \in \mathbb{N}} \left\| \|\rho_0^N\|_q \right\|_{L^m(\Omega)} < \infty$$

with $d \geq 1$, $q > d$ given by (\mathbf{A}^K) and $\beta \in \left(0, \frac{1}{2[1+\frac{1}{d}-\frac{1}{q}]}\right)$. Then

$$\left\| \|\rho - \rho^N\|_{T,q} \right\|_{L^m(\Omega)} \leq C \left\| \|\rho_0 - \rho_0^N\|_q \right\|_{L^m(\Omega)} + CN^{-\varkappa}$$

where

$$\varkappa = \min\left(\frac{\beta}{d}\gamma, \frac{1}{2} - \beta\left(1 + \frac{1}{d} - \frac{1}{q}\right)\right)$$

and ρ is a solution of SPDE (1) with initial condition ρ_0 .

Theorem 3. Assume (\mathbf{A}^V) , (\mathbf{A}^c) , (\mathbf{A}^F) , $K = \delta_0$, let T_{max} be the maximal existence time for (1) and fix $T \in (0, T_{max})$. In addition, let the dynamics of the particle system be given by (12) with $\nu = Id$, σ independent of the space and for any $m \geq 1$,

$$\sup_{N \in \mathbb{N}} \left\| \|\rho_0^N\|_2 \right\|_{L^m(\Omega)} < \infty$$

with $d = 1$ and $\beta \in (0, \frac{1}{3})$. Then

$$\left\| \|\rho - \rho^N\|_{T,2} \right\|_{L^m(\Omega)} \leq C \left\| \|\rho_0 - \rho_0^N\|_2 \right\|_{L^m(\Omega)} + CN^{-\varkappa}$$

where

$$\varkappa = \min\left(\frac{\beta}{2}, \frac{1}{2} - \frac{3}{2}\beta\right)$$

and ρ is a solution of SPDE (1) with initial condition ρ_0 .

In view of the previous result, we obtain a rate of convergence for the genuine empirical measure, which can be interpreted as a propagation of chaos for the marginals of the empirical measure of the particle system. Following [2, Section 8.3], let us introduce the Kantorovich-Rubinstein metric which reads, for any two probability measures μ and ν on \mathbb{T}^d ,

$$\|\mu - \nu\|_0 = \sup \left\{ \int_{\mathbb{T}^d} \phi d(\mu - \nu); \phi \text{ Lipschitz with } \|\phi\|_{L^\infty(\mathbb{T}^d)} \leq 1 \text{ and } \|\phi\|_{\text{Lip}} \leq 1 \right\}. \quad (4)$$

Corollary 1. *Let the same assumptions as in Theorem 2 or Theorem 3 be in place. Then for any $\varepsilon \in (0, \varkappa)$, there exists $C > 0$ such that, for any $N \in \mathbb{N}$,*

$$\left\| \sup_{t \in [0, T]} \|S_t^N - \rho_t\|_0 \right\|_{L^m(\Omega)} \leq C \left(\left\| \|S_0^N - \rho_0\|_0 \right\|_{L^m(\Omega)} + N^{-\varkappa + \varepsilon} \right).$$

Proof. Let $t \in (0, T_{max})$. Let us observe first that there exists $C > 0$ such that for any Lipschitz continuous function ϕ on \mathbb{T}^d , one has

$$|\langle \rho_t^N, \phi \rangle - \langle S_t^N, \phi \rangle| \leq \frac{C \|\phi\|_{\text{Lip}}}{N^{\frac{\beta}{\alpha}}} \quad a.s. \quad (5)$$

Indeed,

$$\begin{aligned} |\langle S_t^N, \phi \rangle - \langle \rho_t^N, \phi \rangle| &= |\langle S_t^N, (\phi - \phi * V^N) \rangle| \\ &\leq \left\langle S_t^N, \int_{\mathbb{T}^d} V(y) |\phi(\cdot) - \phi\left(\cdot - \frac{y}{N^\beta}\right)| dy \right\rangle \\ &\leq \frac{C \|\phi\|_{\text{Lip}}}{N^{\frac{\beta}{\alpha}}}. \end{aligned}$$

Recalling the definition (4) of the Kantorovich-Rubinstein distance, it comes

$$\begin{aligned} \left\| \sup_{t \in [0, T]} \|S_t^N - \rho_t\|_0 \right\|_m &\leq \left\| \sup_{t \in [0, T]} \|S_t^N - \rho_t^N\|_0 \right\|_m + \left\| \sup_{t \in [0, T]} \sup_{\|\phi\|_{L^\infty} \leq 1} \langle \rho_t^N - \rho_t, \phi \rangle \right\|_m \\ &\leq \left\| \sup_{t \in [0, T]} \|S_t^N - \rho_t^N\|_0 \right\|_m + \left\| \sup_{t \in [0, T]} \|\rho_t^N - \rho_t\|_{L^1(\mathbb{T}^d)} \right\|_m, \end{aligned}$$

where we are using the following notation for the norm in $L^m(\Omega)$: $\|\cdot\|_{L^m(\Omega)} = \|\cdot\|_m$. Now applying inequality (5) to the first term on the right-hand side of the above inequality, and Theorem 2 (or Theorem 3) to the second term, we obtain the inequality of Corollary 1. \square

1.5 Application for some stochastic PDE

Let $A > 0$ and $f_A : \mathbb{R} \rightarrow \mathbb{R}$ be a $C^2(\mathbb{R})$ bounded function such that

1. $f_A(x) = x$, for $x \in [-A, A]$,
2. $f_A(x) = A$, for $x > A + 1$ and $f_A(x) = -A$, for $x < -(A + 1)$,
3. $\|f'_A\|_\infty \leq 1$ and $\|f''_A\|_\infty < \infty$.

As a consequence, $\|f_A\|_\infty \leq A + 1$. Now F_A is given by

$$F_A : (x_1, \dots, x_d)^\top \mapsto (f_A(x_1), \dots, f_A(x_d))^\top. \quad (6)$$

When $A = \infty$ then $F_A = Id$. Now, we consider the stochastic PDE

$$d\rho_t = \frac{1}{2} \Delta \rho_t dt - \nabla \cdot (\rho_t (K * \rho_t)) dt - \nabla \rho_t \cdot \sigma_t \circ dB_t \quad (7)$$

where the integral is in the Stratonovich sense, or equivalently in Itô formulation

$$d\rho_t = \frac{1}{2} \left(\Delta \rho_t + \sum_{i,j=1}^d \partial_{ij} \rho_t \sum_{k=1}^d \sigma_t^{ik} \sigma_t^{jk} \right) dt - \nabla \cdot (\rho_t (K * \rho_t)) dt - \nabla \rho_t \cdot \sigma_t dB_t. \quad (8)$$

We consider the associated stochastic moderate particle system

$$dX_t^{i,N} = F_A \left(K * \rho_t^N \left(X_t^{i,N} \right) \right) dt + dW_t^i + \sigma_t dB_t \quad (9)$$

and also the particle system without cutoff:

$$d\tilde{X}_t^{i,N} = \frac{1}{N} \sum_{k=1}^N (K * V^N) (\tilde{X}_t^{i,N} - \tilde{X}_t^{k,N}) dt + dW_t^i + \sigma_t dB_t, \quad t \leq T, \quad 1 \leq i \leq N. \quad (10)$$

For fixed $N \in \mathbb{N}^*$, the drift term is smooth and bounded. Hence, this particle system has a unique strong solution. We denote by \tilde{S}_t^N its empirical distribution, and define $\tilde{\rho}^N = V^N * \tilde{S}^N$. We denote $A_T := C_K \left\| \|\rho\|_{T,q} \right\|_{L^\infty(\Omega)}$ where C_K is given by (\mathbf{A}^K) and ρ is the unique solution of equation (8). Then ρ is also the unique solution of the equation

$$d\rho_t = \frac{1}{2} \left(\Delta \rho_t + \sum_{i,j=1}^d \partial_{ij} \rho_t \sum_{k=1}^d \sigma_t^{ik} \sigma_t^{jk} \right) dt - \nabla \cdot (\rho_t F_A(K * \rho_t)) dt - \nabla \rho_t \cdot \sigma_t dB_t \quad (11)$$

The proof of the following result is based on Corollary 2.3 in [43].

Corollary 2. *Assume (\mathbf{A}^c) , (\mathbf{A}^V) , (\mathbf{A}^I) and (\mathbf{A}^K) and let ρ the unique solution of the SPDE (7) in $L^q_{\mathcal{F}_B}([0, T]; H^1_q(\mathbb{T}^d)) \cap S^\infty_{\mathcal{F}_B}([0, T]; L^1 \cap L^q(\mathbb{T}^d))$, given by Theorem 1. In addition, let the dynamics of the particle system be given by (10) and for any $m \geq 1$,*

$$\sup_{N \in \mathbb{N}} \left\| \|\tilde{\rho}_0^N\|_q \right\|_{L^m(\Omega)} < \infty$$

with $d \geq 1$, $q > d$ given by (\mathbf{A}^K) and $\beta \in \left(0, \frac{1}{2[1+\frac{1}{d}-\frac{1}{q}]}\right)$. Then we have

$$\mathbb{P} \left(\|\tilde{\rho}_t^N - \rho_t\|_{T,q} \geq \eta \right) \leq \frac{C}{\eta^m} \left(\left\| \|\tilde{\rho}_0^N - \rho_0\|_{T,q} \right\|_{L^m(\Omega)} + CN^{-\alpha} \right)^m.$$

Proof. Let us introduce some notations to distinguish the particle systems with and without cutoff. We fix $A > A_T$ to be precisely chosen later. We consider the particles $(X^{i,N})_{1 \leq i \leq N}$ and $(\tilde{X}^{i,N})_{1 \leq i \leq N}$ defined respectively by (9) and (10) with the same initial conditions and driven by the same family of independent \mathbb{T}^d -valued Brownian motions $(W^i)_{i \in \mathbb{N}^*}$ and B_t . For any $N \in \mathbb{N}^*$ and $A > 0$, define

$$\Omega_{N,A} = \left\{ \sup_{\substack{t \in [0,T] \\ i \in \{1, \dots, N\}}} \frac{1}{N} \left| \sum_{k=1}^N (K * V^N) (X_t^{i,N} - X_t^{k,N}) \right| \leq A \right\}$$

and observe that on $\Omega_{N,A}$, we have $X_t^{i,N} = \tilde{X}_t^{i,N}$ for all $t \in [0, T]$ and all $i \in \{1, \dots, N\}$. Since $\frac{1}{N} \sum_{k=1}^N (K * V^N) (X_t^{i,N} - X_t^{k,N}) = K * \rho_t^N(X_t^{i,N})$, we also get that on $\Omega_{N,A}$, $\rho_t^N = \tilde{\rho}_t^N$ for all $t \in [0, T]$ and all $i \in \{1, \dots, N\}$. Hence

$$\begin{aligned} \mathbb{P} \left(\|\tilde{\rho}^N - \rho\|_{T,q} \geq \eta \right) &= \mathbb{P} \left(\Omega_{N,A}^c \cap \left\{ \|\tilde{\rho}^N - \rho\|_{T,q} \geq \eta \right\} \right) \\ &\quad + \mathbb{P} \left(\Omega_{N,A} \cap \left\{ \|\tilde{\rho}^N - \rho\|_{T,q} \geq \eta \right\} \right) \\ &\leq \mathbb{P} \left(\Omega_{N,A}^c \right) + \mathbb{P} \left(\|\rho^N - \rho\|_{T,q} \geq \eta \right). \end{aligned}$$

Now by hypothesis we have that $|K * \rho_t^N(X_t^{i,N})| \leq C_K \|\rho^N\|_{T,q}$. Thus we get that for $A = C_K \left(\eta + \left\| \|\rho\|_{T,q} \right\|_{L^\infty(\Omega)} \right)$,

$$\begin{aligned} \mathbb{P} \left(\Omega_{N,A}^c \right) &\leq \mathbb{P} \left(\|\rho^N\|_{T,q} > \frac{A}{C_K} \right) \\ &\leq \mathbb{P} \left(\|\rho\|_{T,q} + \|\rho^N - \rho\|_{T,q} > \frac{A}{C_K} \right) \\ &\leq \mathbb{P} \left(\left\| \|\rho\|_{T,q} \right\|_{L^\infty(\Omega)} + \|\rho^N - \rho\|_{T,q} > \frac{A}{C_K} \right) \\ &\leq \mathbb{P} \left(\|\rho^N - \rho\|_{T,q} > \eta \right). \end{aligned}$$

Hence

$$\mathbb{P}\left(\|\tilde{\rho}^N - \rho\|_{T,q} \geq \eta\right) \leq 2\mathbb{P}\left(\|\rho^N - \rho\|_{T,q} > \eta\right).$$

Now using Markov's inequality and Theorem 2 with $F(x, u) = F_A(u)$, we obtain the desired result. \square

Biot-Savart kernel and the 2d Navier-Stokes equation. By considering the vorticity field ξ_t associated to the stochastic incompressible two-dimensional Navier-Stokes solution ρ_t , one gets equation (8) with the Biot-Savart kernel $K(x) = \frac{i}{2\pi} \sum_{k \in \mathbb{Z}^2, k \neq 0} \exp(ik \cdot x) \frac{k^\perp}{k^2}$. The original Navier-Stokes solution is then recovered thanks to the formula $\rho_t = K * \xi_t$. The kernel $K \in L^p(\mathbb{T}^2)$ with $p < 2$ and $\|\nabla K * f\|_q \leq C\|f\|_q$ with $1 < q < \infty$. Then $\|K * f\|_{1,q} \leq C\|f\|_q$ with $q > 2$, thus by Sobolev embedding $\|K * f\|_{C^{1-\frac{2}{q}}(\mathbb{T}^2)} \leq C\|f\|_q$. This model arises naturally in fluid mechanics; see, for instance, [4] and [22]. This case is covered by Corollary 2, also we can apply theorem 2 with cutoff $A > C_K \left\| \|\rho\|_{T,q} \right\|_{L^\infty(\Omega)}$ in the SPDE (11) where C_K is given by (\mathbf{A}^K) .

Parabolic-elliptic Keller-Segel models in any dimension. An important and tricky example covered by this paper is the parabolic-elliptic Keller-Segel PDE, which takes the form (8) where the kernel K is the periodization of the function defined on \mathbb{T}^d by $K_0(x) = -\chi_d \frac{x}{|x|^d}$ for some $\chi_d > 0$. The K verifies for all $f \in L^q(\mathbb{T}^d)$, $\|K * f\|_{C^{1-\frac{d}{q}}(\mathbb{T}^d)} \leq C\|f\|_{L^q(\mathbb{T}^d)}$, see Lemma 41 in [44]. This case is also covered by Corollary 2, also we can apply theorem 2 with cutoff $A > C_K \left\| \|\rho\|_{T,q} \right\|_{L^\infty(\Omega)}$ in the SPDE (11) where C_K is given by (\mathbf{A}^K) .

Burgers Equation. When $d = 1$, $K = \delta_0$ the equation (7)-(8) is the well-known stochastic scalar Burgers equation. If the initial condition $\rho_0 \in H^2_2$ then by lemma 4.10 in [1] the solution verifies the maximum principle, that is, $\|\rho\|_\infty \leq C_T \|\rho_0\|_\infty$. Taking $A > C_T \|\rho_0\|_\infty$ the solution of the equation (8) is also solution of the equation (11). The Theorem 3 covers this case.

1.6 Heuristic deduction

For the ease of reading, we may write X_t^i in place of $X_t^{i,N}$.

The starting $X_0^{i,N}$ are i.i.d. with common law μ_0 with density ρ_0 .

Let $x \in \mathbb{T}^d$. Applying Itô's formula with the smooth function $V^N(x - \cdot)$ for

each $i \in \{1, \dots, N\}$ we have

$$\begin{aligned}
V^N(x - X_t^i) &= V^N(x - X_0^i) \\
&\quad - \sum_{j=1}^d \int_0^t \partial_j V^N(x - X_s^i) F_j(X_s^i, K * \rho_s^N(X_s^i)) ds \\
&\quad + \frac{1}{2} \sum_{j,k=1}^d \int_0^t \partial_{jk} V^N(x - X_s^i) (\sigma \sigma^\top)_s^{jk}(X_s^i) ds \\
&\quad - \sum_{j,k=1}^d \int_0^t \partial_j V^N(x - X_s^i) \sigma_s^{jk}(X_s^i) dB_s^k \\
&\quad - \sum_{j,k=1}^d \int_0^t \partial_j V^N(x - X_s^i) \nu_s^{jk}(X_s^i) dW_s^{i,k} \\
&\quad + \frac{1}{2} \sum_{j,k=1}^d \int_0^t \partial_{jk} V^N(x - X_s^i) (\nu \nu^\top)_s^{jk}(X_s^i) ds.
\end{aligned}$$

with $(\sigma \sigma^\top)_s^{jk} \doteq \sum_{l=1}^d \sigma_s^{jl} \sigma_s^{kl}$ and $(\nu \nu^\top)_s^{jk} \doteq \sum_{l=1}^d \nu_s^{jl} \nu_s^{kl}$. Then since $\rho^N = V^N * S^N$ we have

$$\begin{aligned}
\rho_t^N(x) &= \rho_0^N(x) \\
&\quad - \sum_{j=1}^d \int_0^t \langle S_s^N, \partial_j V^N(x - \cdot) F_j(\cdot, K * \rho_s^N(\cdot)) \rangle ds \\
&\quad + \frac{1}{2} \sum_{j,k=1}^d \int_0^t \langle S_s^N, \partial_{jk} V^N(x - \cdot) (\sigma \sigma^\top)_s^{jk}(\cdot) \rangle ds \\
&\quad - \sum_{j,k=1}^d \int_0^t \langle S_s^N, \partial_j V^N(x - \cdot) \sigma_s^{jk}(\cdot) \rangle dB_s^k \\
&\quad - \frac{1}{N} \sum_{i=1}^N \sum_{j,k=1}^d \int_0^t \partial_j V^N(x - X_s^i) \nu_s^{jk}(X_s^i) dW_s^{i,k} \\
&\quad + \frac{1}{2} \sum_{j,k=1}^d \int_0^t \langle S_s^N, \partial_{jk} V^N(x - \cdot) (\nu \nu^\top)_s^{jk}(\cdot) \rangle ds. \tag{12}
\end{aligned}$$

If σ and ν do not depend on the spatial variable we obtain

$$\begin{aligned}
\rho_t^N(x) &= \rho_0^N(x) \\
&- \sum_{j=1}^d \int_0^t \langle S_s^N, \partial_j V^N(x - \cdot) F_j(\cdot, K * \rho_s^N(\cdot)) \rangle ds \\
&+ \frac{1}{2} \sum_{j,k=1}^d \int_0^t \partial_{jk} \rho_s^N (\sigma \sigma^\top)_s^{jk} ds \\
&- \sum_{j,k=1}^d \int_0^t \partial_j \rho_s^N \sigma_s^{jk} dB_s^k \quad (= \sigma_s^\top \nabla \rho_s^N dB_s) \\
&- \frac{1}{N} \sum_{i=1}^N \sum_{j,k=1}^d \int_0^t \partial_j V^N(x - X_s^i) \nu_s^{jk} dW_s^{i,k} \quad (= \nu_s^\top \nabla V^N(x - X_s^i) dW_s^i) \\
&+ \frac{1}{2} \sum_{j,k=1}^d \int_0^t \partial_{jk} \rho_s^N (\nu \nu^\top)_s^{jk} ds. \tag{13}
\end{aligned}$$

Under these conditions, we may conjecture that a limiting measure-valued deterministic process ρ_t exists and its evolution equation (in weak form) is given by

$$\begin{aligned}
d\rho_t &= \frac{1}{2} \sum_{i,j=1}^d \partial_{ij} \left(\rho_t \sum_{k=1}^d (\nu_t^{ik} \nu_t^{jk} + \sigma_t^{ik} \sigma_t^{jk}) \right) dt \\
&- \nabla \cdot (\rho_t F(\cdot, K * \rho_t)) dt - \nabla \rho_t \cdot \sigma_t dB_t. \tag{14}
\end{aligned}$$

1.7 Definition of solution

Definition 1.1. A family of random functions $\{\rho_t(\omega) : t \geq 0, \omega \in \Omega\}$ lying in $S_{\mathcal{F}^B}^\infty([0, T]; L^1 \cap L^q(\mathbb{T}^d))$ is a solution to (1) if ρ_t satisfies the following stochastic integral equation for all $\phi \in C^2(\mathbb{T}^d)$,

$$\begin{aligned}
\langle \rho_t, \phi \rangle &= \langle \rho_0, \phi \rangle + \int_0^t \langle \rho_s F(\cdot, K * \rho_s), \nabla \phi \rangle ds \\
&+ \frac{1}{2} \int_0^t \left\langle \rho_s, \sum_{i,j=1}^d \partial_{ij} \phi \sum_{k=1}^d (\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk}) \right\rangle ds \\
&+ \int_0^t \left\langle \rho_s, \sum_{i=1}^d \partial_i \phi \sum_{k=1}^d \sigma_s^{ik} \right\rangle dB_s^k. \tag{15}
\end{aligned}$$

Remark 1. Let $q \geq 2$. The following will be helpful throughout the text.

$$(i) \quad \nabla(|\rho(x)|^q) = q|\rho(x)|^{q-2}\rho(x)\nabla\rho(x).$$

$$(ii) \quad \nabla(q|\rho(x)|^{q-2}\rho(x)) = q(q-1)|\rho(x)|^{q-2}\nabla\rho(x).$$

2 Proof of the main theorems

2.1 Proof of Theorem 2

Proof. Applying to $(\rho - \rho^N)$ the Itô's formula for the L^q -norm of a H_q^1 -valued process in [35] we have

$$\begin{aligned} \|\rho_t - \rho_t^N\|_q^q &= \|\rho_0 - \rho_0^N\|_q^q \\ &\quad - \sum_{j,k=1}^d \int_0^t \int_{\mathbb{T}^d} q|\rho_s - \rho_s^N|^{q-2}(\rho_s - \rho_s^N) \partial_j(\rho_s - \rho_s^N) \sigma_s^{jk} dx dB_s^k \\ &\quad - \frac{1}{N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q|\rho_s - \rho_s^N|^{q-2}(\rho_s - \rho_s^N) (\nabla V^N)(x - X_s^i) dx dW_s^i \\ &\quad + q(q-1) \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^{q-2} \nabla(\rho_s - \rho_s^N) \left[\rho_s F(x, K * \rho_s) \right. \\ &\quad \quad \left. - \langle S_s^N, V^N(x - \cdot) \cdot F(\cdot, K * \rho_s^N(\cdot)) \rangle \right] dx ds \\ &\quad + \frac{1}{2} \sum_{j,k=1}^d \int_0^t \int_{\mathbb{T}^d} q|\rho_s - \rho_s^N|^{q-2}(\rho_s - \rho_s^N) \partial_{jk}(\rho_s - \rho_s^N) (\sigma \sigma^T)_s^{jk} dx ds \\ &\quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\nabla(\rho_s - \rho_s^N)|^2 dx ds \\ &\quad + \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\sigma_s^T \nabla(\rho_s - \rho_s^N)|^2 dx ds \\ &\quad + \frac{1}{2N^2} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |(\nabla V^N)(x - X_s^i)|^2 dx ds. \end{aligned}$$

Integration by parts gives us (recall that σ does not depend on the spatial

variable)

$$\begin{aligned} & \frac{1}{2} \sum_{j,k=1}^d \int_0^t \int_{\mathbb{T}^d} q |\rho_s - \rho_s^N|^{q-2} (\rho_s - \rho_s^N) \partial_{jk} (\rho_s - \rho_s^N) (\sigma \sigma^T)_s^{jk} dx ds \\ &= -\frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\sigma_s^T \nabla (\rho_s - \rho_s^N)|^2 dx ds, \end{aligned}$$

hence we get

$$\begin{aligned} \|\rho_t - \rho_t^N\|_q^q &\leq \|\rho_0 - \rho_0^N\|_q^q + I_1 + I_2 \\ &\quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\nabla (\rho_s - \rho_s^N)|^2 dx ds \\ &\quad - \sum_{j,k=1}^d \int_0^t \sigma_s^{jk} \underbrace{\int_{\mathbb{T}^d} \partial_j |\rho_s - \rho_s^N|^q dx}_{=0} dB_s^k \\ &\quad - M_t^N, \end{aligned} \tag{16}$$

where

$$\begin{aligned} I_1 &\doteq q(q-1) \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^{q-2} \nabla (\rho_s - \rho_s^N) \left[\rho_s F(x, K * \rho_s) \right. \\ &\quad \left. - \langle S_s^N, V^N(x - \cdot) F(\cdot, K * \rho_s^N) \rangle \right] dx ds, \end{aligned}$$

$$I_2 \doteq \frac{1}{2N^2} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |(\nabla V^N)(x - X_s^i)|^2 dx ds,$$

and

$$M_t^N \doteq \frac{1}{N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q |\rho_s - \rho_s^N|^{q-2} (\rho_s - \rho_s^N) (\nabla V^N)(x - X_s^i) dx dW_s^i.$$

If $q > 2$, by Young inequality with $\frac{1}{\frac{q}{q-2}} + \frac{1}{2} = 1$ we obtain

$$\begin{aligned} I_2 &= \frac{q(q-1)}{2N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^{q-2} \frac{1}{N} |(\nabla V^N)(x - X_s^i)|^2 dx ds \\ &\leq \frac{q(q-1)}{2} \int_0^t \|\rho_s - \rho_s^N\|_q^q ds + \frac{q(q-1)}{2N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} \frac{1}{N^{\frac{q}{2}}} |(\nabla V^N)(x - X_s^i)|^q dx ds \end{aligned}$$

Now by (22),

$$I_2 \leq C_2 \int_0^t \|\rho_s - \rho_s^N\|_q^q ds + \frac{q(q-1)}{2} TN^{(-\frac{q}{2} + q\beta(1 + \frac{1}{d}) - \beta)} \|\nabla V\|_q^q$$

where $C_2 = \frac{q(q-1)}{2}$, if $q > 2$ and $C_2 = 0$, if $q = 2$.

Now, we will estimate the first term of I_1 . Since F is Lipschitz, K satisfies (\mathbf{A}^K) , and $V \geq 0$ we have

$$\begin{aligned}
& |\langle S_s^N, V^N(x - \cdot) (F(x, K * \rho_s^N(x)) - F(\cdot, K * \rho_s^N(\cdot))) \rangle| \\
& \leq \langle S_s^N, V^N(x - \cdot) |F(x, K * \rho_s^N(x)) - F(\cdot, K * \rho_s^N(\cdot))| \rangle \\
& \leq L \langle S_s^N, V^N(x - \cdot) (|x - \cdot| + |K * (\rho_s^N(x) - \rho_s^N(\cdot))|) \rangle \\
& \leq L \langle S_s^N, V^N(x - \cdot) (|x - \cdot| + \|\rho_s^N\|_q |x - \cdot|^\gamma) \rangle \\
& \leq L \left(N^{-\frac{\beta}{d}} + \|\rho_s^N\|_q N^{-\frac{\beta}{d}\gamma} \right) |\rho_s^N(x)|,
\end{aligned}$$

where in the last inequality we are using that $\text{supp } V \subset \{x \in \mathbb{R}^d \mid |x| < \frac{1}{2}\}$. We add and subtract the terms $F(x, K * \rho_s^N)\rho_s$ and $F(x, K * \rho_s^N)\rho_s^N$ to deduce

$$\begin{aligned}
& |F(x, K * \rho_s)\rho_s - \langle S_s^N, V^N(x - \cdot)F(\cdot, K * \rho_s^N) \rangle| \\
& \leq |F(x, K * \rho_s)\rho_s - F(x, K * \rho_s^N)\rho_s| \\
& \quad + |F(x, K * \rho_s^N)\rho_s - F(x, K * \rho_s^N)\rho_s^N| \\
& \quad + \langle S_s^N, V^N(x - \cdot) |F(x, K * \rho_s^N) - F(\cdot, K * \rho_s^N)| \rangle \\
& \leq L |K * (\rho_s - \rho_s^N)| |\rho_s| \\
& \quad + L |\rho_s - \rho_s^N| \\
& \quad + L \left(N^{-\frac{\beta}{d}} + \|\rho_s^N\|_q N^{-\frac{\beta}{d}\gamma} \right) |\rho_s^N(x)| \\
& \doteq (a + b + c).
\end{aligned}$$

Now, using ϵ -Young inequality for $\frac{1}{2} + \frac{1}{2} = 1$, we get

$$\begin{aligned}
& |\nabla(\rho_s - \rho_s^N)|(a + b + c) \\
& \leq \epsilon |\nabla(\rho_s - \rho_s^N)|^2 + C_\epsilon (a + b + c)^2 \\
& \leq \epsilon |\nabla(\rho_s - \rho_s^N)|^2 + C_\epsilon 3L^2 \left(|K * (\rho_s - \rho_s^N)|^2 |\rho_s|^2 + |\rho_s - \rho_s^N|^2 + C_N^2 |\rho_s^N|^2 \right)
\end{aligned}$$

where $C_N \doteq N^{-\frac{\beta}{d}} + \|\rho_s^N\|_q N^{-\frac{\beta}{d}\gamma}$. For the other constants, we will merge them into C , which may change from line to line. Thus we have

$$\begin{aligned}
|I_1| & \leq \epsilon \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\nabla(\rho_s - \rho_s^N)|^2 dx ds \\
& \quad + C \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^q dx ds. \\
& \quad + C \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^{q-2} (|K * (\rho_s - \rho_s^N)|^2 |\rho_s|^2 + C_N^2 |\rho_s^N|^2) dx ds.
\end{aligned}$$

We continue with the last term above. For $q > 2$, we use 1-Young inequality for $\frac{1}{q-2} + \frac{1}{2} = 1$,

$$\begin{aligned} & \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^{q-2} (|K * (\rho_s - \rho_s^N)|^2 |\rho_s|^2 + C_N^2 |\rho_s^N|^2) dx ds \\ & \leq C \int_0^t \int_{\mathbb{T}^d} |\rho_s - \rho_s^N|^q dx ds + C \int_0^t \int_{\mathbb{T}^d} \|K * (\rho_s - \rho_s^N)\|_\infty^q |\rho_s|^q + C_N^q |\rho_s^N|^q dx ds \\ & \leq C \int_0^t \|\rho_s - \rho_s^N\|_q^q ds + C \|\rho\|_{T,q}^q \int_0^t \|\rho_s - \rho_s^N\|_q^q ds + CT C_N^q \|\rho^N\|_{T,q}^q \end{aligned}$$

where in the last inequality we are using (\mathbf{A}^K) . Now, using that $\|\rho\|_{T,q} \leq C$ a.s. due to Theorem 1, we arrive at

$$\begin{aligned} |I_1| & \leq \epsilon \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s - \rho_s^N|^{q-2} |\nabla(\rho_s - \rho_s^N)|^2 ds \\ & \quad + C \int_0^t \|\rho_s - \rho_s^N\|_q^q ds + C C_N^q \|\rho^N\|_{T,q}^q. \end{aligned}$$

Finally, we join the estimates with $\epsilon < \frac{1}{2}$ and continue from (16) to get

$$\begin{aligned} \|\rho_t - \rho_t^N\|_q^q & \leq \|\rho_0 - \rho_0^N\|_q^q + C \int_0^t \|\rho_s - \rho_s^N\|_q^q ds \\ & \quad + CN^{(-\frac{q}{2} + q\beta(1 + \frac{1}{d}) - \beta)} \\ & \quad + C \left(N^{-\frac{\beta}{d}} + \|\rho^N\|_{T,q} N^{-\frac{\beta}{d}\gamma} \right)^q \|\rho^N\|_{T,q}^q \\ & \quad + |M_t^N|. \end{aligned}$$

By Jensen's inequality with $|\cdot|^{\frac{m}{q}}$, $\frac{m}{q} > 1$ we have for all $r \in [0, T]$

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in [0, r]} \|\rho_t - \rho_t^N\|_q \right)^m & \leq C \mathbb{E} \left(\|\rho_0 - \rho_0^N\|_q \right)^m \\ & \quad + C \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t - \rho_t^N\|_q \right)^m ds \\ & \quad + CN^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m} \\ & \quad + C \left(N^{-\frac{\beta}{d}} \right)^m \mathbb{E} \|\rho^N\|_{T,q}^m \\ & \quad + C \left(N^{-\frac{\beta}{d}\gamma} \right)^m \mathbb{E} \left(\|\rho^N\|_{T,q} \|\rho^N\|_{T,q} \right)^m \\ & \quad + C \mathbb{E} \left(\sup_{t \in [0, r]} |M_t^N| \right)^{\frac{m}{q}}. \end{aligned} \tag{17}$$

Now we need a uniform in $N \in \mathbb{N}$ estimate for the regularized empirical measure, for which the proof is given in the Appendix A .

Lemma 4. *Under the hypotheses of Theorem 2 or 3, we have the following:*

$$\sup_{N \in \mathbb{N}} \left\| \left\| \rho^N \right\|_{T,q} \right\|_m < \infty.$$

Finally, similar computations as those of Lemma 6 on Appendix A yields

$$\mathbb{E} \left(\sup_{t \in [0,r]} |M_t^N| \right)^{\frac{m}{q}} \lesssim \int_0^r \mathbb{E} \left(\sup_{t \in [0,s]} \|\rho_t - \rho_t^N\|_q \right)^m ds + N^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m}.$$

The above with Lemma 4 in (17) leads to

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in [0,r]} \|\rho_t - \rho_t^N\|_q \right)^m &\leq C \mathbb{E} \left(\|\rho_0 - \rho_0^N\|_q \right)^m \\ &\quad + C \int_0^r \mathbb{E} \left(\sup_{t \in [0,s]} \|\rho_t - \rho_t^N\|_q \right)^m ds \\ &\quad + CN^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m} \\ &\quad + C \left(N^{-\frac{\beta}{d}} \right)^m \\ &\quad + C \left(N^{-\frac{\beta}{d}\gamma} \right)^m \\ &\quad + CN^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m}. \end{aligned}$$

Now take $r = T$. Grönwall's Lemma implies

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in [0,T]} \|\rho_t - \rho_t^N\|_q \right)^m &\leq C \mathbb{E} \left(\|\rho_0 - \rho_0^N\|_q \right)^m \\ &\quad + CN^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m} \\ &\quad + C \left(N^{-\frac{\beta}{d}} \right)^m \\ &\quad + C \left(N^{-\frac{\beta}{d}\gamma} \right)^m \end{aligned}$$

and since $\gamma \leq 1$, we end up with

$$\left\| \|\rho - \rho^N\|_{T,q} \right\|_{L^m(\Omega)} \leq C \left\| \|\rho_0 - \rho_0^N\|_q \right\|_{L^m(\Omega)} + CN^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))} + CN^{-\frac{\beta}{d}\gamma}.$$

□

2.2 Proof of Theorem 3

Proof. Applying the Itô's formula for the L^q -norm of a H_q^1 -valued process in [35] as in the proof of Theorem 2 we get

$$\begin{aligned}
\|\rho_t - \rho_t^N\|_2^2 &= \|\rho_0 - \rho_0^N\|_2^2 + I_1 + I_2 \\
&\quad - \int_0^t \int_{\mathbb{T}} |\nabla(\rho_s - \rho_s^N)|^2 dx ds \\
&\quad - \sum_{j,k=1}^d \int_0^t \sigma_s^{jk} \underbrace{\int_{\mathbb{T}} \nabla|\rho_s - \rho_s^N|^2 dx}_{=0} dB_s^k \\
&\quad - M_t^N,
\end{aligned} \tag{18}$$

where

$$\begin{aligned}
I_1 &\doteq 2 \int_0^t \int_{\mathbb{T}} \nabla(\rho_s - \rho_s^N) [\rho_s F(x, \rho_s) - \langle S_s^N, V^N(x - \cdot) F(\cdot, \rho_s^N(\cdot)) \rangle] dx ds, \\
I_2 &\doteq \frac{1}{N^2} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}} |(\nabla V^N)(x - X_s^i)|^2 dx ds,
\end{aligned}$$

and

$$M_t^N \doteq \frac{2}{N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}} (\rho_s - \rho_s^N) (\nabla V^N)(x - X_s^i) dx dW_s^i.$$

By a change of variables in I_2 and the estimate (22) we obtain

$$I_2 \leq CN^{-(1+3\beta)} \|\nabla V\|_2^2.$$

Now, for I_1 , we use that F is Lipschitz (\mathbf{A}^F) and the estimate provided by Lemma 5 on ρ_s^N combined with the Sobolev embedding $H_2^1(\mathbb{T}) \hookrightarrow C^{\frac{1}{2}}(\mathbb{T})$.

$$\begin{aligned}
&|\langle S_s^N, V^N(x - \cdot) (F(x, \rho_s^N(x)) - F(\cdot, \rho_s^N(\cdot))) \rangle| \\
&\leq \langle S_s^N, V^N(x - \cdot) |F(x, \rho_s^N(x)) - F(\cdot, \rho_s^N(\cdot))| \rangle \\
&\leq L \langle S_s^N, V^N(x - \cdot) (|x - \cdot| + |\rho_s^N(x) - \rho_s^N(\cdot)|) \rangle \\
&\leq L \langle S_s^N, V^N(x - \cdot) (|x - \cdot| + \|\rho_s^N\|_{\frac{1}{2}} |x - \cdot|^{\frac{1}{2}}) \rangle \\
&\leq L \left(N^{-\beta} + \|\rho_s^N\|_{\frac{1}{2}} N^{-\frac{\beta}{2}} \right) |\rho_s^N(x)|
\end{aligned}$$

where in the last inequality we are using that $\text{supp } V \subset \{x \in \mathbb{R}^d \mid |x| < \frac{1}{2}\}$.

In the same way, we add and subtract the terms $F(x, \rho_s^N)\rho_s$ and $F(x, \rho_s^N)\rho_s^N$ to obtain

$$\begin{aligned}
&|F(x, \rho_s)\rho_s - \langle S_s^N, V^N(x - \cdot) F(\cdot, \rho_s^N) \rangle| \\
&\leq |F(x, \rho_s)\rho_s - F(x, \rho_s^N)\rho_s| + |F(x, \rho_s^N)\rho_s - F(x, \rho_s^N)\rho_s^N| \\
&\quad + \langle S_s^N, V^N(x - \cdot) |F(x, \rho_s^N(x)) - F(\cdot, \rho_s^N(\cdot))| \rangle \\
&\leq L |\rho_s - \rho_s^N| |\rho_s| + L |\rho_s - \rho_s^N| + L \left(N^{-\beta} + \|\rho_s^N\|_{\frac{1}{2}} N^{-\frac{\beta}{2}} \right) |\rho_s^N|.
\end{aligned}$$

Now by Theorem 8.8 in [5], for all $p \geq 1$,

$$\|\cdot\|_\infty \leq \|\cdot\|_p + \|\nabla \cdot\|_p.$$

We observe that $\rho \in L^2 H_2^1(\mathbb{T}) \hookrightarrow L^2 C^{\frac{1}{2}}(\mathbb{T})$, a.s.. Then a.e. in $s \in [0, T]$, taking $p = 1$, we have, a.s.

$$\|(\rho_s - \rho_s^N)^2\|_\infty \leq \|(\rho_s - \rho_s^N)^2\|_1 + 2\|(\rho_s - \rho_s^N)\nabla(\rho_s - \rho_s^N)\|_1$$

and thus by Holder's inequality,

$$\|\rho_s - \rho_s^N\|_\infty^2 \leq \|\rho_s - \rho_s^N\|_2^2 + 2\|\rho_s - \rho_s^N\|_2 \|\nabla(\rho_s - \rho_s^N)\|_2.$$

By ϵ -Young inequality we have

$$\begin{aligned} & \int_{\mathbb{T}} |\nabla(\rho_s - \rho_s^N)| |\rho_s - \rho_s^N| |\rho_s| dx \\ & \leq \epsilon \|\nabla(\rho_s - \rho_s^N)\|_2^2 + C_\epsilon \|\rho_s\|_2^2 \|\rho_s - \rho_s^N\|_\infty^2 \\ & \leq 2\epsilon \|\nabla(\rho_s - \rho_s^N)\|_2^2 + C_\epsilon (\|\rho_s\|_2^2 + 4C_\epsilon \|\rho_s\|_2^4) \|\rho_s - \rho_s^N\|_2^2. \end{aligned}$$

By denoting $\hat{C}_N \doteq N^{-\beta} + \|\rho_s^N\|_{\frac{1}{2}} N^{-\frac{\beta}{2}}$, the ϵ -Young inequality yields

$$\begin{aligned} & |\nabla(\rho_s - \rho_s^N)| |\rho_s - \rho_s^N| + \hat{C}_N |\nabla(\rho_s - \rho_s^N)| |\rho_s^N| \\ & \leq 2\epsilon |\nabla(\rho_s - \rho_s^N)|^2 + C_\epsilon |\rho_s - \rho_s^N|^2 + C_\epsilon \hat{C}_N^2 |\rho_s^N|^2. \end{aligned}$$

Therefore, we obtain an estimate for I_1 :

$$\begin{aligned} |I_1| & \leq 2L \int_0^t \int_{\mathbb{T}} |\nabla(\rho_s - \rho_s^N)| (|\rho_s - \rho_s^N| |\rho_s| + |\rho_s - \rho_s^N| + \hat{C}_N |\rho_s^N|) dx ds \\ & \leq \epsilon 2L \int_0^t \|\nabla(\rho_s - \rho_s^N)\|_2^2 ds + C_\epsilon 2L \int_0^t (\|\rho_s\|_2^2 + 4C_\epsilon \|\rho_s\|_2^4) \|\rho_s - \rho_s^N\|_2^2 ds \\ & \quad + 4\epsilon L \int_0^t \|\nabla(\rho_s - \rho_s^N)\|_2^2 ds + C_\epsilon 2L \int_0^t \|\rho_s - \rho_s^N\|_2^2 ds \\ & \quad + C_\epsilon 4L \left(N^{-2\beta} + \int_0^T \|\rho_s^N\|_{\frac{1}{2}}^2 N^{-2\frac{\beta}{2}} ds \right) T \|\rho^N\|_{T,2}^2. \end{aligned}$$

Finally, taking $\epsilon < \frac{1}{6L}$, the previous estimates lead to

$$\begin{aligned} \|\rho_t - \rho_t^N\|_2^2 & \leq \|\rho_0 - \rho_0^N\|_2^2 + C \int_0^t (C + \|\rho\|_{T,2}^2 + \|\rho\|_{T,2}^4) \|\rho_s - \rho_s^N\|_2^2 ds \\ & \quad + CN^{(-1+3\beta)} \\ & \quad + C \left(N^{-2\beta} + \int_0^T \|\rho_s^N\|_{\frac{1}{2}}^2 N^{-\beta} ds \right) \|\rho^N\|_{T,2}^2 \\ & \quad + |M_t^N|. \end{aligned}$$

Since $\left\| \|\rho\|_{T,2} \right\|_{L^\infty(\Omega)} < \infty$, by Jensen's inequality with $|\cdot|^{\frac{m}{2}}$, $\frac{m}{2} > 1$ we have for all $r \in [0, T]$

$$\begin{aligned}
\mathbb{E} \left(\sup_{t \in [0,r]} \|\rho_t - \rho_t^N\|_2 \right)^m &\leq C \mathbb{E} (\|\rho_0 - \rho_0^N\|_2)^m \\
&+ C \int_0^r \mathbb{E} \left(\sup_{t \in [0,s]} \|\rho_t - \rho_t^N\|_2 \right)^m ds \\
&+ CN^{(-1+3\beta)\frac{m}{2}} \\
&+ C (N^{-\beta})^m \mathbb{E} \|\rho^N\|_{T,2}^m \\
&+ C \left(N^{-\frac{\beta}{2}} \right)^m \mathbb{E} \left(\|\rho^N\|_{T,2} \|\rho^N\|_{L^2 C^{\frac{1}{2}}} \right)^m \\
&+ E \left(\sup_{t \in [0,r]} |M_t^N| \right)^{\frac{m}{2}}. \tag{19}
\end{aligned}$$

Note that for all $m \geq 1$,

$$\sup_N \left\| \|\rho^N\|_{L^2 C^{\frac{1}{2}}} \right\|_m \lesssim \sup_N \left\| \|\rho^N\|_{L^2 H_2^1} \right\|_m$$

where we use that $L^2 H_2^1(\mathbb{T}) \hookrightarrow L^2 C^{\frac{1}{2}}(\mathbb{T})$. We now need a uniform in $N \in \mathbb{N}$ estimate for the regularized empirical measure, whose proof is given in the Appendix A.

Lemma 5. *Under the hypotheses of Theorem 3, we have the following:*

$$\sup_{N \in \mathbb{N}} \left\| \|\rho^N\|_{L^2 H_2^1} \right\|_m < \infty.$$

Finally, similar computations as those of Lemma 6 on Appendix A yield

$$\mathbb{E} \left(\sup_{t \in [0,r]} |M_t^N| \right)^{\frac{m}{2}} \lesssim \int_0^r \mathbb{E} \left(\sup_{t \in [0,s]} \|\rho_t - \rho_t^N\|_2 \right)^m ds + N^{(-1+3\beta)\frac{m}{2}}.$$

From (19), and Lemmas 4 and 5 we conclude

$$\begin{aligned}
\mathbb{E} \left(\sup_{t \in [0,r]} \|\rho_t - \rho_t^N\|_2 \right)^m &\leq C \mathbb{E} (\|\rho_0 - \rho_0^N\|_2)^m \\
&+ C \int_0^r \mathbb{E} \left(\sup_{t \in [0,s]} \|\rho_t - \rho_t^N\|_2 \right)^m ds \\
&+ CN^{(-1+3\beta)\frac{m}{2}} \\
&+ C (N^{-\beta})^m \\
&+ C \left(N^{-\frac{\beta}{2}} \right)^m \\
&+ CN^{(-1+3\beta)\frac{m}{2}}.
\end{aligned}$$

Then taking $r = T$, Grönwall's Lemma implies

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in [0, T]} \|\rho_t - \rho_t^N\|_2 \right)^m &\leq C \mathbb{E} (\|\rho_0 - \rho_0^N\|_2)^m \\ &\quad + CN^{(-1+3\beta)\frac{m}{2}} + C \left(N^{-\frac{\beta}{2}} \right)^m \end{aligned}$$

and

$$\begin{aligned} \left\| \|\rho - \rho^N\|_{T,2} \right\|_m &\leq C \left\| \|\rho_0 - \rho_0^N\|_2 \right\|_m \\ &\quad + CN^{(-1+3\beta)\frac{1}{2}} + CN^{-\frac{\beta}{2}}. \end{aligned}$$

□

A Uniform in $N \in \mathbb{N}$ estimates for the regularized empirical measure

This section presents the proofs of Lemmas 4 and 5.

Proof of Lemmas 4 and 5. Applying the Itô's formula on ρ_t^N for the function $x \mapsto |x|^q$, $q \geq 2$, (essentially (16) with $\rho = 0$), Fubini's theorem and its stochastic version, we get

$$\begin{aligned} \|\rho_t^N\|_q^q &= \|\rho_0^N\|_q^q + I_1 + I_2 \\ &\quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s^N|^{q-2} |\nabla \rho_s^N|^2 dx ds \\ &\quad - M_t^N \end{aligned} \tag{20}$$

where

$$\begin{aligned} I_1 &\doteq q(q-1) \int_0^t \int_{\mathbb{T}^d} |\rho_s^N|^{q-2} \nabla \rho_s^N \langle S_s^N, V^N(x - \cdot) F(\cdot, K * \rho_s^N(\cdot)) \rangle dx ds, \\ I_2 &\doteq \frac{1}{2N^2} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s^N|^{q-2} |(\nabla V^N)(x - X_s^i)|^2 dx ds, \end{aligned}$$

and

$$M_t^N \doteq \frac{1}{N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} q |\rho_s^N|^{q-1} (\nabla V^N)(x - X_s^i) dx dW_s^i.$$

If $q > 2$ by Young inequality, with $\frac{1}{\frac{q}{q-2}} + \frac{1}{2} = 1$, we obtain

$$\begin{aligned} I_2 &= \frac{q(q-1)}{2N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} |\rho_s^N|^{q-2} \frac{1}{N} |(\nabla V^N)(x - X_s^i)|^2 dx ds \\ &\lesssim \frac{q(q-1)}{2N} \sum_{i=1}^N \int_0^t \|\rho_s^N\|_q^q ds + \frac{q(q-1)}{2N} \sum_{i=1}^N \int_0^t \int_{\mathbb{T}^d} \frac{1}{N^{\frac{q}{2}}} |(\nabla V^N)(x - X_s^i)|^q dx ds. \end{aligned} \tag{21}$$

Now for $q \geq 2$, by a change of variables we have

$$\begin{aligned}
\int_0^t \int_{\mathbb{T}^d} |(\nabla V^N)(x)|^q dx ds &\leq T \int_{\mathbb{R}^d} \left| \nabla \left[N^\beta V \left(N^{\beta/d} \cdot \right) \right] (x) \right|^q dx \\
&= T \int_{\mathbb{R}^d} \left| N^{\beta(1+\frac{1}{d})} (\nabla V) \left(N^{\beta/d} x \right) \right|^q dx \\
&= T N^{q\beta(1+\frac{1}{d})} \int_{\mathbb{R}^d} \left| (\nabla V) \left(N^{\beta/d} x \right) \right|^q dx \\
&= T N^{q\beta(1+\frac{1}{d})} \int_{\mathbb{R}^d} \left| (\nabla V) \left(N^{\beta/d} x \right) \right|^q \frac{N^\beta}{N^\beta} dx \\
&= T N^{q\beta(1+\frac{1}{d})} \int_{\mathbb{R}^d} |(\nabla V)(y)|^q \frac{1}{N^\beta} dy \\
&= T N^{(q\beta(1+\frac{1}{d})-\beta)} \int_{\mathbb{R}^d} |\nabla V(y)|^q dy \\
&= T N^{(q\beta(1+\frac{1}{d})-\beta)} \|\nabla V\|_q^q. \tag{22}
\end{aligned}$$

Then

$$I_2 \leq C_2 \int_0^t \|\rho_s^N\|_q^q ds + \frac{q(q-1)}{2} T N^{(-\frac{q}{2}+q\beta(1+\frac{1}{d})-\beta)} \|\nabla V\|_q^q$$

where $C_2 = \frac{q(q-1)}{2}$, if $q > 2$ and $C_2 = 0$, if $q = 2$.

Now, we observe

$$\begin{aligned}
|\langle S_s^N, V^N(x - \cdot) F(\cdot, K * \rho_s^N) \rangle| &\leq \langle S_s^N, |V^N(x - \cdot) F(\cdot, K * \rho_s^N)| \rangle \\
&\stackrel{V \geq 0}{\leq} \|F\|_\infty |\langle S_s^N, V^N(x - \cdot) \rangle| \\
&= \|F\|_\infty |\rho_s^N|.
\end{aligned}$$

Then we deduce

$$\begin{aligned}
I_1 &= \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s^N|^{q-2} \nabla \rho_s^N \langle S_s^N, V^N(x - \cdot) F(\cdot, K * \rho_s^N) \rangle dx ds \\
&\leq \|F\|_\infty \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s^N|^{q-2} |\nabla \rho_s^N| |\rho_s^N| dx ds.
\end{aligned}$$

By ϵ -Young inequality with $\frac{1}{2} + \frac{1}{2} = 1$ we get

$$\begin{aligned}
I_1 &\leq C_\epsilon q(q-1) \|F\|_\infty^2 \int_0^t \|\rho_s^N\|_q^q ds \\
&\quad + \epsilon \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s^N|^{q-2} |\nabla \rho_s^N|^2 dx ds. \tag{23}
\end{aligned}$$

Thus, we join the estimates with $\epsilon < \frac{1}{2}$ and continue from (20):

$$\begin{aligned}
\|\rho_t^N\|_q^q &\leq \|\rho_0^N\|_q^q \\
&+ \left(C_\epsilon q(q-1)\|F\|_\infty^2 + \frac{q(q-1)}{2}1_{[q>2]} \right) \int_0^t \|\rho_s^N\|_q^q ds \\
&+ \frac{q(q-1)}{2} TN^{(-\frac{q}{2}+q\beta(1+\frac{1}{d})-\beta)} \|\nabla V\|_q^q \\
&+ |M_t^N|. \tag{24}
\end{aligned}$$

Taking the supremum, applying Jensen's inequality, and then taking the expectation, we obtain:

$$\begin{aligned}
\mathbb{E} \left(\sup_{t \in [0, r]} \|\rho_t^N\|_q \right)^m &\leq C \mathbb{E} \left(\|\rho_0^N\|_q \right)^m \\
&+ C \left(C_\epsilon q(q-1)\|F\|_\infty^2 + \frac{q(q-1)}{2}1_{[q>2]} \right)^{\frac{m}{q}} \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t^N\|_q \right)^m ds \\
&+ C \left(\frac{q(q-1)}{2} T \right)^{\frac{m}{q}} N^{(-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q}))m} \|\nabla V\|_q^m \\
&+ C \mathbb{E} \left(\sup_{t \in [0, r]} |M_t^N| \right)^{\frac{m}{q}}. \tag{25}
\end{aligned}$$

Finally, from (25) and lemma 6 we deduce

$$\begin{aligned}
\mathbb{E} \left(\sup_{t \in [0, r]} \|\rho_t^N\|_q \right)^m &\leq C \mathbb{E} \left(\|\rho_0^N\|_q \right)^m + C \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t^N\|_q \right)^m ds \\
&+ C \left(N^{(-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q}))m} + N^{(-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q}))m} \right).
\end{aligned}$$

Taking $r = T$ and Grönwall's Lemma, we conclude that

$$\mathbb{E} \left(\sup_{t \in [0, T]} \|\rho_t^N\|_q \right)^m \leq C \mathbb{E} \left(\|\rho_0^N\|_q \right)^m + CN^{(-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q}))m}$$

and

$$\left\| \|\rho^N\|_{T, q} \right\|_m \leq C \left\| \|\rho_0^N\|_q \right\|_m + CN^{-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q})}.$$

This proves Lemma 4. In order to get a proof of Lemma 5, considering the above estimates, and continuing from (20) with $\epsilon = \frac{1}{4}$, instead of (24), we obtain

$$\begin{aligned}
\|\rho_t^N\|_2^2 + \frac{1}{2} \int_0^t \|\nabla \rho_s^N\|_2^2 ds &\leq \|\rho_0^N\|_2^2 + C \int_0^t \|\rho_s^N\|_2^2 ds \\
&+ TN^{(-1+2\beta(1+\frac{1}{d})-\beta)} \|\nabla V\|_2^2 \\
&+ |M_t^N|.
\end{aligned}$$

From Lemma 4 we have $\sup_{N \in \mathbb{N}} \mathbb{E} \|\rho^N\|_{L^2 L^2} < \infty$. Now we estimate $\|\nabla \rho^N\|_2$. By Jensen's inequality, with $|\cdot|^{\frac{m}{2}}$ and $m > 2$, we have

$$\begin{aligned} \mathbb{E} \left(\int_0^r \|\nabla \rho_s^N\|_2^2 ds \right)^{\frac{m}{2}} &\leq C \left(1 + E(\|\rho_0^N\|_2)^m \right) \\ &+ C \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t^N\|_2 \right)^m ds \\ &+ C \mathbb{E} \left(\sup_{t \in [0, r]} |M_t^N| \right)^{\frac{m}{2}}. \end{aligned}$$

From Lemma 6 we conclude

$$\begin{aligned} \sup_N \left(\mathbb{E} \left(\int_0^T \|\nabla \rho_s^N\|_2^2 ds \right)^{\frac{m}{2}} \right)^{\frac{1}{m}} &\lesssim 1 + \sup_N \left\| \|\rho_0^N\|_2 \right\|_m \\ &+ T \sup_N \left\| \|\rho^N\|_{T, 2} \right\|_m < \infty. \end{aligned}$$

□

Lemma 6. *Under the assumptions of Theorem 2 or 3, it holds that*

$$\mathbb{E} \left(\sup_{t \in [0, r]} |M_t^N| \right)^{\frac{m}{q}} \lesssim \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t^N\|_q \right)^m ds + N^{(-\frac{1}{2} + \beta(1 + \frac{1}{d} - \frac{1}{q}))m}.$$

Proof. First, notice that

$$M_t^N \doteq \int_0^t \int_{\mathbb{T}^d} q |\rho_s^N|^{q-1} \frac{1}{N} \sum_{i=1}^N (\nabla V^N)(x - X_s^i) dx dW_s^i$$

is a martingale, and $|M_t^N|$ is a positive sub-martingale. So, by Doob's maximal inequality

$$\mathbb{E} \left(\sup_{t \in [0, r]} |M_t^N| \right)^{\frac{m}{q}} \leq \left(\frac{\frac{m}{q}}{\frac{m}{q} - 1} \right)^{\frac{m}{q}} \mathbb{E} |M_r^N|^{\frac{m}{q}}.$$

Now, by Burkholder-Davis-Gundy inequality

$$\begin{aligned} \mathbb{E} |M_r^N|^{\frac{m}{q}} &\doteq \mathbb{E} \left| \frac{1}{N} \sum_{i=1}^N \int_0^r \int_{\mathbb{T}^d} q |\rho_s^N|^{q-1} (\nabla V^N)(x - X_s^i) dx dW_s^i \right|^{\frac{m}{q}} \\ &\lesssim \mathbb{E} \left| \frac{1}{N^2} \sum_{i=1}^N \int_0^r \left(\int_{\mathbb{T}^d} q |\rho_s^N|^{q-1} (\nabla V^N)(x - X_s^i) dx \right)^2 ds \right|^{\frac{m}{2q}} \\ &= \mathbb{E} \left| \frac{1}{N} \sum_{i=1}^N \int_0^r \left(\int_{\mathbb{T}^d} q |\rho_s^N|^{q-1} \frac{1}{N^{\frac{1}{2}}} (\nabla V^N)(x - X_s^i) dx \right)^2 ds \right|^{\frac{m}{2q}} \end{aligned}$$

where in the last equality we moved a power of N inside the integral. Now by Young's inequality with $\frac{1}{q-1} + \frac{1}{q}$ and Jensen's inequality,

$$\begin{aligned}
E|M_r^N|^{\frac{m}{q}} &\lesssim \mathbb{E} \left| \frac{1}{N} \sum_{i=1}^N \int_0^r \left(\int_{\mathbb{T}^d} |\rho_s^N|^q dx \right)^2 ds \right|^{\frac{m}{2q}} \\
&\quad + \mathbb{E} \left| \frac{1}{N} \sum_{i=1}^N \int_0^r \left(\int_{\mathbb{T}^d} \frac{1}{N^{q/2}} |(\nabla V^N)(x - X_s^i)|^q dx \right)^2 ds \right|^{\frac{m}{2q}} \\
&\lesssim \mathbb{E} \int_0^r \|\rho_s^N\|_q^m ds \\
&\quad + \frac{1}{N^{\frac{m}{2}}} \left| \int_0^r \left(\int_{\mathbb{T}^d} |(\nabla V^N)(x)|^q dx \right)^2 ds \right|^{\frac{m}{2q}}
\end{aligned}$$

where in the last inequality, we have applied a change of variables. Finally, by (22)

$$\begin{aligned}
E|M_r^N|^{\frac{m}{q}} &\lesssim \mathbb{E} \int_0^r \|\rho_s^N\|_q^m ds \\
&\quad + N^{-\frac{m}{2}} \left(TN^{(2q\beta(1+\frac{1}{d})-2\beta)} \|\nabla V\|_q^{2q} \right)^{\frac{m}{2q}} \\
&\lesssim \mathbb{E} \int_0^r \|\rho_s^N\|_q^m ds \\
&\quad + N^{(-q+2q\beta(1+\frac{1}{d})-2\beta)\frac{m}{2q}} \|\nabla V\|_q^m \\
&\lesssim \int_0^r \mathbb{E} \left(\sup_{t \in [0, s]} \|\rho_t^N\|_q \right)^m ds \\
&\quad + N^{(-\frac{1}{2}+\beta(1+\frac{1}{d}-\frac{1}{q}))m} \|\nabla V\|_q^m.
\end{aligned}$$

□

B Well-posedness of limiting equation (1)

Here, we will prove the well posedness of the limiting equation for the system of particles in a suitable space. We will adapt the strategy of [31].

Proof of Theorem 1. The proof is divided into two parts. First, we solve the linearized problem using the L^q -Theory of SPDEs. Next, we make a contraction argument to yield the solution of the original equation as a fixed point of the solution map.

Let

$$\mathbb{B} \doteq \left\{ \rho \in S_{\mathcal{FB}}^\infty([0, T]; L^q(\mathbb{T}^d)) \mid \|\rho\|_{T, q} \Big\|_{L^\infty(\Omega)} \leq \kappa \ell \right\}$$

with metric $d(\rho, \rho') \doteq \left\| \|\rho - \rho'\|_{T,q} \right\|_{L^\infty(\Omega)}$, $\kappa = \|\rho_0\|_q$ and ℓ is a positive constant to be determined.

We define the operator $\mathcal{T} : \mathbb{B} \rightarrow S_{\mathcal{F}B}^\infty([0, T]; L^q(\mathbb{T}^d))$ as follows: for each $\xi \in \mathbb{B}$, let $\mathcal{T}(\xi) := \rho^\xi$ be the solution to the following linear SPDE:

$$\begin{cases} d\rho_t = \frac{1}{2} \sum_{i,j=1}^d \partial_{ij} \left(\rho_t \sum_{k=1}^d \left(\nu_t^{ik} \nu_t^{jk} + \sigma_t^{ik} \sigma_t^{jk} \right) \right) dt - \nabla \cdot (\rho_t F(\cdot, K * \xi_t)) dt \\ \quad - \nabla \rho_t \cdot \sigma_t dB_t \\ \rho_0 \text{ is given.} \end{cases} \quad (26)$$

Now by Assumption (\mathbf{A}^c) , we rewrite (26) as follows:

$$\begin{cases} d\rho_t = \frac{1}{2} \sum_{i,j=1}^d \partial_{ij} \rho_t \sum_{k=1}^d \left(\nu_t^{ik} \nu_t^{jk} + \sigma_t^{ik} \sigma_t^{jk} \right) dt + f_t(\rho) dt \\ \quad - \nabla \rho_t \cdot \sigma_t dB_t \\ \rho_0 \text{ is given.} \end{cases} \quad (27)$$

where

$$\begin{aligned} f_t(\rho) &\doteq -\nabla \cdot (\rho_t F(\cdot, K * \xi_t)) + \sum_{i,j=1}^d \partial_j \rho_t \partial_i \left((\nu \nu^\top)_t^{ij} + (\sigma \sigma^\top)_t^{ij} \right) \\ &\quad + \frac{1}{2} \rho_t \sum_{i,j=1}^d \partial_{ij} \left((\nu \nu^\top)_t^{ij} + (\sigma \sigma^\top)_t^{ij} \right) \\ &= -\nabla \cdot (\rho_t F(\cdot, K * \xi_t)) + 2 \sum_{j=1}^d \partial_j \rho_t \sum_{i=1}^d \partial_i a_t^{ij} + \rho_t \sum_{i,j=1}^d \partial_{ij} a_t^{ij} \end{aligned}$$

and

$$a_t^{ij} \doteq \frac{1}{2} \sum_{k=1}^d \left(\nu_t^{ik} \nu_t^{jk} + \sigma_t^{ik} \sigma_t^{jk} \right).$$

We now proceed to verify the conditions 5.1–6 of Theorem 5.1 in [35]. For the reader's convenience, we restate these conditions below using our notations:

5.1 There exist $\Lambda, \lambda > 0$ such that

$$\Lambda |\xi|^2 \geq \sum_{i,j,k=1}^d \nu_t^{ik}(x) \nu_t^{jk}(x) \xi^i \xi^j \geq \lambda |\xi|^2,$$

for any $t \geq 0$, $x \in \mathbb{T}^d$, $\xi \in \mathbb{R}^d$.

5.2 For any $i, j \in \{1, \dots, d\}$ and $\epsilon > 0$, there exists $\delta > 0$, such that

$$|a_t^{ij}(x) - a_t^{ij}(y)| + |\sigma_t^i(x) - \sigma_t^i(y)| < \epsilon$$

whenever $|x - y| < \delta$, $t \geq 0$.

5.3 For any $i, j \in \{1, \dots, d\}$ and $t \geq 0$, the functions a_t^{ij} and σ_t^i are continuously differentiable.

5.4 For any $u \in H_q^1$, $f_t(u)$ is predictable as a function taking values in H_q^{-1} .

5.5 For any $i, j \in \{1, \dots, d\}$ and $t \geq 0$,

$$\|a_t^{ij}\|_{C^1} + \|\sigma_t^i\|_\infty \leq M.$$

5.6 For any $\epsilon > 0$, there exists $C_\epsilon > 0$, such that, for any $t \geq 0$ and $u_t, v_t \in H_q^1$, we have

$$\|f_t(u) - f_t(v)\|_{-1,q} \leq \epsilon \|u_t - v_t\|_{1,q} + C_\epsilon \|u_t - v_t\|_{-1,q}.$$

Now note that the condition 5.1 is exactly (\mathbf{A}_{iii}^c) . The conditions 5.2, 5.3 and 5.5 are verified by assumption (\mathbf{A}_t^c) . For condition 5.4, note that $f_t(u)$ is a composition of predictable functions.

For the remaining one, 5.6, we notice that if F is given by (\mathbf{A}^F) (thus bounded) and K by (\mathbf{A}^K) , we have for all $u_t, v_t \in H_q^1$,

$$\begin{aligned} \|\nabla \cdot (F(\cdot, K * \xi_t)(u_t - v_t))\|_{-1,q} &= \left\| (I - \Delta)^{-\frac{1}{2}} \nabla \cdot (F(\cdot, K * \xi_t)(u_t - v_t)) \right\|_q \\ &\leq C \|F(\cdot, K * \xi_t)(u_t - v_t)\|_q \\ &\leq C \|u_t - v_t\|_q \\ &\leq C \|u_t - v_t\|_{1,q}^{\frac{1}{2}} \|u_t - v_t\|_{-1,q}^{\frac{1}{2}} \\ &\leq \epsilon \|u_t - v_t\|_{1,q} + \epsilon^{-1} C^2 \|u_t - v_t\|_{-1,q}, \end{aligned}$$

by the interpolation inequality. On the other hand, if we assume (\mathbf{A}^I) and (\mathbf{A}^K) we have for all $u_t, v_t \in H_q^1$,

$$\begin{aligned} \|\nabla \cdot (F(\cdot, K * \xi_t)(u_t - v_t))\|_{-1,q} &= \left\| (I - \Delta)^{-\frac{1}{2}} \nabla \cdot (K * \xi_t)(u_t - v_t) \right\|_q \\ &\leq C \|(K * \xi_t)(u_t - v_t)\|_q \\ &\leq CC_K \|\xi_t\|_q \|u_t - v_t\|_q \\ &\leq CC_K \kappa l \|u_t - v_t\|_{0,q} \\ &\leq c \|u_t - v_t\|_{1,q}^{\frac{1}{2}} \|u_t - v_t\|_{-1,q}^{\frac{1}{2}} \\ &\leq \epsilon \|u_t - v_t\|_{1,q} + \epsilon^{-1} c^2 \|u_t - v_t\|_{-1,q}, \end{aligned}$$

by the interpolation inequality and using $\xi \in \mathbb{B}$. Recall that $\|K * f\|_\infty \leq C_K \|f\|_q$, for q given by (\mathbf{A}^K) .

For the remaining terms in $f_t(\rho)$, we have

$$\|(u_t - v_t) \partial_{ij} a_t^{ij}\|_{-1,q} \leq \|(u_t - v_t) \partial_{ij} a_t^{ij}\|_q \leq \|a_t^{ij}\|_{C^2} \|u_t - v_t\|_q$$

and then it follows by interpolation inequality as before, while for the remaining one, we use the dual space $(H_q^{-1})^* = H_q^1$ with the duality pair given by $(\cdot, \cdot)_{-1,1}$. Indeed, by applying the Leibniz's rule, we have

$$\begin{aligned} & \|\partial_j(u_t - v_t) \partial_i a_t^{ij}\|_{-1,q} \\ &= \sup_{\|\phi\|_{1,q'}=1} (\partial_j(u_t - v_t) \partial_i a_t^{ij}, \phi)_{-1,1} \\ &= \sup_{\|\phi\|_{1,q'}=1} ((u_t - v_t) \partial_i a_t^{ij}, -\partial_j \phi)_{-1,1} + \sup_{\|\phi\|_{1,q'}=1} ((u_t - v_t) \partial_{ji} a_t^{ij}, -\phi)_{-1,1} \\ &\leq \sup_{\|\phi\|_{1,q'}=1} \left(\|\partial_j \phi\|_{q'} \|(u_t - v_t) \partial_i a_t^{ij}\|_q + \|\phi\|_{1,q'} \|(u_t - v_t) \partial_{ji} a_t^{ij}\|_q \right) \\ &\leq 2 \|a_t^{ij}\|_{C^2} \|u_t - v_t\|_q \end{aligned}$$

where the Hölder's inequality was applied in the second-to-last inequality. Then again, interpolation inequality yields the estimate that verifies assumption 5.6.

Notice that the initial condition $\rho_0 \in H_q^{1-\frac{2}{q}}(\mathbb{T}^d)$ is deterministic, therefore, we have by Theorem 5.1 in [35] that the linear SPDE (26) admits a unique solution $\rho^\xi \in L^q_{\mathcal{F}^B}([0, T]; H_q^1(\mathbb{T}^d))$ (see Definition 3.1 in [35]). In addition, since Theorem 7.1 (iii) in [35] holds for $q \geq 2$ (by Theorem 4.2 in [35]) and $\|\rho_t\|_1 = \|\rho_0\|_1 = 1$, a.s., we have $\rho^\xi \in S^q_{\mathcal{F}^B}([0, T]; L^1 \cap L^q(\mathbb{T}^d))$. Moreover, by the maximum principle (Theorem 5.12 in [35]) $\rho^\xi(t, \cdot) \geq 0$, a.s..

We now check that ρ^ξ is also in $S^\infty_{\mathcal{F}^B}([0, T]; L^1 \cap L^q(\mathbb{T}^d))$, and we drop the superscript ξ to ease the computations. For the following, if not mentioned, we are assuming (\mathbf{A}^F) and (\mathbf{A}^K) , and we make observations about the other sets of assumptions. Applying the Itô's formula for the L^q -norm of a H_q^1 -valued process in [35],

$$\begin{aligned} \|\rho_t\|_q^q &= \|\rho_0\|_q^q \\ &+ \frac{1}{2} \int_{\mathbb{T}^d} \int_0^t q |\rho_s(x)|^{q-2} \rho_s(x) (\nu_s(x) \nu_s^\top(x) + \sigma_s(x) \sigma_s^\top(x)) \cdot D^2 \rho_s(x) ds dx \\ &- \int_{\mathbb{T}^d} \int_0^t q |\rho_s(x)|^{q-2} \rho_s(x) f_s(\rho)(x) ds dx \\ &- \int_{\mathbb{T}^d} \int_0^t q |\rho_s(x)|^{q-2} \rho_s(x) \nabla \rho_s(x) \cdot \sigma_s(x) dB_s dx \\ &+ \frac{1}{2} \int_{\mathbb{T}^d} \int_0^t q(q-1) |\rho_s(x)|^{q-2} |\sigma_s^\top(x) \nabla \rho_s(x)|^2 ds dx \\ &= \text{(I)} + \text{(II)} + \text{(III)} + \text{(IV)}. \end{aligned} \tag{28}$$

Above, $A \cdot B \doteq \sum_{i,j} A_{ij} B_{ij}$. Using stochastic Fubini and integration by parts, we handle each term separately. Notice that under assumption (\mathbf{A}_{ii}^c) , i.e., $\sum_i \partial_i \sigma_t^{ik} = 0$ we have

$$\begin{aligned}
(\text{III}) &= - \int_0^t \int_{\mathbb{T}^d} \sum_{ik} q |\rho_s(x)|^{q-2} \rho_s(x) \partial_i \rho_s(x) \sigma_s^{ik}(x) dx dB_s^k \\
&= - \int_0^t \int_{\mathbb{T}^d} \sum_{ik} \partial_i (|\rho_s(x)|^q) \sigma_s^{ik}(x) dx dB_s^k \\
&= \int_0^t \int_{\mathbb{T}^d} |\rho_s(x)|^q \sum_{ik} \partial_i \sigma_s^{ik}(x) dx dB_s^k = 0.
\end{aligned}$$

For (II), we have two parts

$$\begin{aligned}
(\text{II}) &= - \int_0^t \int_{\mathbb{T}^d} q |\rho_s(x)|^{q-2} \rho_s(x) \nabla \cdot [\rho_s(x) F(x, K * \xi_s)] dx ds \\
&\quad - \int_0^t \int_{\mathbb{T}^d} q |\rho_s(x)|^{q-2} \rho_s(x) \left(2 \sum_{ij} \partial_j \rho_s \partial_i a_s^{ij} + \rho_s \sum_{ij} \partial_{ij} a_s^{ij} \right) dx ds
\end{aligned}$$

For the first one, after integration by parts, we use ϵ -Young inequality with $\frac{1}{2} + \frac{1}{2} = 1$

$$\begin{aligned}
&= q(q-1) \int_0^t \int_{\mathbb{T}^d} |\rho_s(x)|^{q-2} \rho_s(x) \nabla \rho_s(x) \cdot F(x, K * \xi_s) dx ds \\
&\leq q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\rho_s(x)|^{q-2} \rho_s(x) |\nabla \rho_s(x)| dx ds \tag{29} \\
&\leq q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\rho_s(x)|^{q-2} (C_\epsilon \rho_s(x)^2 + \epsilon |\nabla \rho_s(x)|^2) dx ds \\
&= q(q-1)LC_\epsilon \int_0^t \|\rho_s\|_q^q ds + \epsilon q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds
\end{aligned}$$

Above, we used (\mathbf{A}^F) so that F is bounded by L . In fact, if one assumes (\mathbf{A}^I) and (\mathbf{A}^K) , (29) implies that the first term in (II) reads

$$\leq C_\epsilon q(q-1)C_K^2 \kappa^2 \ell^2 \int_0^t \|\rho_s\|_q^q ds + \epsilon q(q-1) \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds,$$

since $\|K * \xi_s\|_{L^\infty} \leq C_K \|\xi_s\|_q \leq C_K \kappa \ell$. For the second (II),

$$\begin{aligned}
& - \int_0^t \int_{\mathbb{T}^d} q |\rho_s(x)|^{q-2} \rho_s(x) \left(2 \sum_{ij} \partial_j \rho_s \partial_i a_s^{ij} + \rho_s \sum_{ij} \partial_{ij} a_s^{ij} \right) dx ds \\
& \leq \int_0^t \int_{\mathbb{T}^d} 2q |\rho_s|^{q-2} \rho_s \left| \sum_{ij} (\partial_j \rho_s \partial_i a_s^{ij}) \right| dx ds + q \int_0^t \sum_{ij} \|a_s^{ij}\|_{C^2} \|\rho_s\|_q^q ds \\
& \leq \int_0^t \int_{\mathbb{T}^d} 2q |\rho_s|^{q-2} \rho_s |\nabla \rho_s| \left(\sum_j \left(\sum_i \partial_i a_s^{ij} \right)^2 \right)^{\frac{1}{2}} dx ds + qM \int_0^t \|\rho_s\|_q^q ds
\end{aligned}$$

where we used Cauchy-Schwarz inequality. In order to apply the Grönwall's lemma, the second term above is already suitable, while the first one, after bounding the a^{ij} term with its L^∞ norm provided in (\mathbf{A}^c) , is a multiple of (29), from which we get an analogous estimate. In fact, due to assumption (\mathbf{A}_i^c) , we have

$$\begin{aligned}
& \left| \sum_{ik} \partial_i \nu_s^{ik} \nu_s^{jk} + \nu_s^{ik} \partial_i \nu_s^{jk} \right| \\
& \leq \sum_{ik} \left(\|\partial_i \nu_s^{ik}\|_\infty + \|\nu_s^{ik}\|_\infty \right) \left(\|\nu_s^{jk}\|_\infty + \|\nabla \nu_s^{jk}\|_\infty \right) \leq 2M^2.
\end{aligned}$$

and analogously for σ . Thus,

$$(\text{II}) \leq C \int_0^t \|\rho_s\|_q^q ds + \epsilon (q(q-1)L + q\sqrt{d}4M^2) \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds.$$

Also, if (\mathbf{A}^I) and (\mathbf{A}^K) are in place

$$\begin{aligned}
(\text{II}) & \leq C_\epsilon \left(q(q-1)C_K^2 \kappa^2 \ell^2 + q\sqrt{d}4M^2 + qM \right) \int_0^t \|\rho_s\|_q^q ds \\
& \quad + \epsilon (q(q-1) + q\sqrt{d}4M^2) \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds.
\end{aligned}$$

Now for (I), again by integration by parts,

$$\begin{aligned}
(\text{I}) & \doteq \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} \sum_{ijk} q |\rho_s|^{q-2} \rho_s (\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk}) \partial_{ij} \rho_s dx ds \\
& = -\frac{1}{2} \int_0^t \int_{\mathbb{T}^d} \sum_{ijk} q(q-1) |\rho_s|^{q-2} \partial_i \rho_s (\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk}) \partial_j \rho_s dx ds \\
& \quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} \sum_{ijk} q |\rho_s|^{q-2} \rho_s \partial_i \left(\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk} \right) \partial_j \rho_s dx ds.
\end{aligned}$$

Notice that $\sum_{ijk} \nu_s^{ik} \nu_s^{jk} \partial_i \rho_s \partial_j \rho_s \geq \lambda |\nabla \rho_s|^2$, by assumption (\mathbf{A}_{iii}^c) . So by multiplying this inequality by -1 and taking into account that

$$\sum_k \left| \sum_i \partial_i \rho_s \sigma_s^{ik} \right|^2 = \sum_{ijk} \partial_i \rho_s (\sigma_s^{ik} \sigma_s^{jk}) \partial_j \rho_s,$$

we get

$$\begin{aligned}
(\text{I}) &\leq -\frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s|^{q-2} \lambda |\nabla \rho_s|^2 dx ds \\
&\quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} \sum_k q(q-1) |\rho_s|^{q-2} \left| \sum_i \partial_i \rho_s \sigma_s^{ik} \right|^2 dx ds \\
&\quad - \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q |\rho_s|^{q-2} \rho_s \sum_j \partial_j \rho_s \partial_i \left(\sum_{ik} \nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk} \right) dx ds \\
&\leq -\frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s|^{q-2} \left(\lambda |\nabla \rho_s|^2 + |\sigma_s^T \nabla \rho_s|^2 \right) dx ds \\
&\quad + \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q |\rho_s|^{q-1} |\nabla \rho_s| \left(\sum_j \left(\partial_i \sum_{ik} (\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk}) \right)^2 \right)^{\frac{1}{2}} dx ds
\end{aligned}$$

where the last inequality follows by Cauchy-Schwarz inequality. Notice that the second-to-last line contains $-(\text{IV})$. Recall

$$\left(\sum_j \left(\partial_i \sum_{ik} (\nu_s^{ik} \nu_s^{jk} + \sigma_s^{ik} \sigma_s^{jk}) \right)^2 \right)^{\frac{1}{2}} \leq \left(\sum_j (4M^2)^2 \right)^{\frac{1}{2}} = \sqrt{d} 4M^2.$$

Thus, we have

$$\begin{aligned}
(\text{I}) &\leq -(\text{IV}) - \frac{\lambda}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds \\
&\quad + \frac{1}{2} \int_0^t \int_{\mathbb{T}^d} q |\rho_s|^{q-1} |\nabla \rho_s| 4M^2 \sqrt{d} dx ds \\
&\leq -(\text{IV}) - \frac{\lambda}{2} \int_0^t \int_{\mathbb{T}^d} q(q-1) |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds \\
&\quad + C_\epsilon \int_0^t \|\rho_s\|_q^q ds + \epsilon q 4M^4 d \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds
\end{aligned}$$

by ϵ -Young inequality. Finally,

$$\begin{aligned}
\|\rho_t\|_q^q &= \|\rho_0\|_q^q + (\text{I}) + (\text{II}) + (\text{III}) + (\text{IV}) \\
&\leq \|\rho_0\|_q^q + C \int_0^t \|\rho_s\|_q^q ds \\
&\quad + \left(\epsilon L q(q-1) + \epsilon q 4M^4 d - \frac{\lambda}{2} q(q-1) \right) \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds.
\end{aligned}$$

Again, if (\mathbf{A}^I) and (\mathbf{A}^K) are in place

$$\begin{aligned}
\|\rho_t\|_q^q &= \|\rho_0\|_q^q + (\text{I}) + (\text{II}) + (\text{III}) + (\text{IV}) \\
&\leq \|\rho_0\|_q^q + C_\epsilon \left(q(q-1) C_K^2 \kappa^2 \ell^2 + q \sqrt{d} 4M^2 + qM + 1 \right) \int_0^t \|\rho_s\|_q^q ds \\
&\quad + \left(\epsilon q(q-1) + \epsilon q 4M^4 d - \frac{\lambda}{2} q(q-1) \right) \int_0^t \int_{\mathbb{T}^d} |\rho_s|^{q-2} |\nabla \rho_s|^2 dx ds.
\end{aligned}$$

Finally, under the assumption (\mathbf{A}^F) , by taking ϵ sufficiently small and $\ell \doteq e^{\frac{CT}{q}}$, an application of Grönwall's Lemma yields

$$\sup_{t \in [0, T]} \|\rho_t\|_q \leq \|\rho_0\|_q (e^{CT})^{\frac{1}{q}} \leq \kappa \ell \quad \text{a.s.}$$

which means $\rho^\xi \in \mathbb{B}$.

Now, under the assumptions (\mathbf{A}^I) and (\mathbf{A}^K) , and by taking T and ϵ sufficiently small and $\ell > 1$, an application of Grönwall's Lemma yields

$$\sup_{t \in [0, T]} \|\rho_t\|_q \leq \|\rho_0\|_q (e^{CT})^{\frac{1}{q}} \leq \kappa \ell \quad \text{a.s.}$$

where

$$C \doteq C_\epsilon \left(q(q-1)C_K^2 \kappa^2 \ell^2 + q\sqrt{d}4M^2 + qM + 1 \right).$$

This means $\rho^\xi \in \mathbb{B}$.

We now show that the map \mathcal{T} is a contraction.

For any $\bar{\xi}, \xi \in \mathbb{B}$, set $\delta\rho = \rho^{\bar{\xi}} - \rho^\xi$ and $\delta\xi = \bar{\xi} - \xi$. As before, we apply Itô's formula for the L^q -norm of a $H^{1,q}$ -valued process in [35] to $\delta\rho$ and obtain

$$\begin{aligned} & \|\delta\rho_t\|_q^q \\ &= \frac{1}{2} \int_{\mathbb{T}^d} \int_0^t q |\delta\rho_s(x)|^{q-2} \delta\rho_s(x) (\nu_s(x) \nu_s^\top(x) + \sigma_s(x) \sigma_s^\top(x)) \cdot D^2 \delta\rho_s(x) \, ds \, dx \\ & - \int_{\mathbb{T}^d} \int_0^t q |\delta\rho_s(x)|^{q-2} \delta\rho_s(x) [f_{\bar{\xi}}(\rho^{\bar{\xi}}, s, x) - f_\xi(\rho^\xi, s, x)] \, ds \, dx \\ & - \int_{\mathbb{T}^d} \int_0^t q |\delta\rho_s(x)|^{q-2} \delta\rho_s(x) \nabla \delta\rho_s(x) \cdot \sigma_s(x) \, dB_s \, dx \\ & + \frac{1}{2} \int_{\mathbb{T}^d} \int_0^t q(q-1) |\delta\rho_s(x)|^{q-2} |\sigma_s^\top(x) \nabla \delta\rho_s(x)|^2 \, ds \, dx \\ &= \text{(I)} + \text{(II)} + \text{(III)} + \text{(IV)}, \end{aligned} \tag{30}$$

where the numbered terms are analogous to before with $\delta\rho$ in place of ρ , except (II), which has a nonlinear part

$$\begin{aligned} \text{(II)} & \doteq - \int_{\mathbb{T}^d} \int_0^t q |\delta\rho_s(x)|^{q-2} \delta\rho_s \nabla \cdot [\rho_s^{\bar{\xi}}(x) F(x, K * \bar{\xi}_s)] \, ds \, dx \\ & + \int_{\mathbb{T}^d} \int_0^t q |\delta\rho_s(x)|^{q-2} \delta\rho_s \nabla \cdot [\rho_s^\xi(x) F(x, K * \xi_s)] \, ds \, dx \\ & - \int_0^t \int_{\mathbb{T}^d} q |\delta\rho_s(x)|^{q-2} \delta\rho_s(x) \left(2 \sum_{ij} \partial_j \delta\rho_s \partial_i a_s^{ij} + \delta\rho_s \sum_{ij} \partial_{ij} a_s^{ij} \right) \, dx \, ds \end{aligned}$$

The above linear part has estimates on $\delta\rho$ following from the same reasoning as before, while for the nonlinear (let us denote it by (II)') we subtract and add

$\rho_s^\xi(x)$ to get

$$\begin{aligned}
(\text{II})' &= - \int_0^t \int_{\mathbb{T}^d} q |\delta \rho_s|^{q-2} \delta \rho_s \nabla \cdot [\delta \rho_s(x) F(x, K * \bar{\xi}_s)] dx ds \\
&\quad - \int_0^t \int_{\mathbb{T}^d} q |\delta \rho_s|^{q-2} \delta \rho_s \nabla \cdot [\rho_s^\xi(x) F(x, K * \bar{\xi}_s)] dx ds \\
&\quad + \int_{\mathbb{T}^d} \int_0^t q |\delta \rho_s|^{q-2} \delta \rho_s \nabla \cdot [\rho_s^\xi(x) F(x, K * \xi_s)] ds dx
\end{aligned}$$

So integration by parts gives

$$\begin{aligned}
(\text{II})' &\leq q(q-1) \int_0^t \int_{\mathbb{T}^d} |\delta \rho_s|^{q-2} \left(|\nabla \delta \rho_s| |\delta \rho_s| L + |\nabla \delta \rho_s| |\rho_s^\xi| L |K * \delta \xi_s| \right) dx ds \\
&\leq q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\delta \rho_s|^{q-2} \left(|\nabla \delta \rho_s| |\delta \rho_s| + |\nabla \delta \rho_s| |\rho_s^\xi| \|\delta \xi_s\|_q \right) dx ds \quad (31)
\end{aligned}$$

Above, we also used (\mathbf{A}^K) and that F is bounded and Lipschitz (\mathbf{A}^F) . In fact, one can show similar bounds assuming (\mathbf{A}^K) and (\mathbf{A}^I) , noticing that $F(x, u) = u$ is also Lipschitz, and the boundedness of F being replaced by that of $K * \bar{\xi}_s$, since $\|K * \bar{\xi}_s\|_\infty \leq C_K \|\bar{\xi}_s\|_q \leq C_K \kappa \ell$.

Next, by ϵ -Young inequality, we have

$$|\delta \rho_s|^{q-1} |\nabla \delta \rho_s| = |\delta \rho_s|^{q-2} |\delta \rho_s| |\nabla \delta \rho_s| \leq C_\epsilon |\delta \rho_s|^q + \epsilon |\delta \rho_s|^{q-2} |\nabla \delta \rho_s|^2$$

and also

$$\begin{aligned}
&\| |\delta \rho_s|^{q-2} |\nabla \delta \rho_s| \rho_s^\xi \|\delta \xi_s\|_q \|_1 \\
&\leq \| |\delta \rho_s|^{\frac{q-2}{2}} |\nabla \delta \rho_s| \|_2 \| |\delta \rho_s|^{\frac{q-2}{2}} \rho_s^\xi \|\delta \xi_s\|_q \|_2 \\
&\leq \epsilon \| |\delta \rho_s|^{\frac{q-2}{2}} |\nabla \delta \rho_s| \|_2^2 + C_\epsilon \| |\delta \rho_s|^{\frac{q-2}{2}} \rho_s^\xi \|\delta \xi_s\|_q \|_2^2
\end{aligned}$$

Again, by Young's inequality, we have

$$\begin{aligned}
\| |\delta \rho_s|^{\frac{q-2}{2}} \rho_s^\xi \|\delta \xi_s\|_q \|_2^2 &= \| |\delta \rho_s|^{q-2} (\rho_s^\xi)^2 \|\delta \xi_s\|_q^2 \|_1 \\
&\leq \| \delta \rho_s \|_q^q + \| \rho_s^\xi \|\delta \xi_s\|_q \|_q^q \\
&= \| \delta \rho_s \|_q^q + \| \rho_s^\xi \|_q^q \|\delta \xi_s\|_q^q
\end{aligned}$$

Using the above in (31), we get

$$\begin{aligned}
(\text{II})' &\leq 2C_\epsilon q(q-1)L \int_0^t \| \delta \rho_s \|_q^q ds + 2\epsilon q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\delta \rho_s|^{q-2} |\nabla \delta \rho_s|^2 dx ds \\
&\quad + C_\epsilon q(q-1)L \int_0^t \| \rho_s^\xi \|_q^q \|\delta \xi_s\|_q^q ds \\
&\leq 2C_\epsilon q(q-1)L \int_0^t \| \delta \rho_s \|_q^q ds + 2\epsilon q(q-1)L \int_0^t \int_{\mathbb{T}^d} |\delta \rho_s|^{q-2} |\nabla \delta \rho_s|^2 dx ds \\
&\quad + C_\epsilon q(q-1)L \left\| \| \rho_s^\xi \|_{T,q} \right\|_\infty^q \int_0^t \| \delta \xi_s \|_q^q ds
\end{aligned}$$

Finally, similarly to before, by choosing $\epsilon > 0$ sufficiently small, we obtain

$$\begin{aligned} \|\delta\rho_t\|_q^q &= \text{(I)} + \text{(II)} + \text{(III)} + \text{(IV)} \\ &\leq C \int_0^t \|\delta\rho_s\|_q^q ds + C(\kappa\ell)^q \int_0^t \|\delta\xi_s\|_q^q ds \end{aligned}$$

Thus, an application of Grönwall's lemma yields

$$\left\| \|\delta\rho\|_{T,q} \right\|_\infty \leq e^{\frac{CT}{q}} (C)^{\frac{1}{q}} (\kappa\ell) T^{\frac{1}{q}} \left\| \|\delta\xi\|_{T,q} \right\|_\infty$$

and taking T small, depending on all given data $\kappa, \lambda, C_K, q, L, M$ and d , we obtain that \mathcal{T} is a contraction.

□

Acknowledgements

C. Olivera is partially supported by FAPESP-ANR by the grant 2022/03379-0, by FAPESP by the grant 2020/04426-6, and CNPq by the grant 422145/2023-8. A. B. de Souza is partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001. J. Knorst is partially supported by FAPESP by the grants 2022/13413-0, BEPE 2023/14629-0.

References

- [1] D. Alonso-Oran, D., A. Bethencourt de León and S. Takao, *The Burgers' equation with stochastic transport: shock formation, local and global existence of smooth solutions*, Nonlinear Differ. Equ. Appl. 26, 2019.
- [2] V. I. Bogachev, *Measure Theory, Vol. II*, Springer-Verlag, Berlin, 2007.
- [3] D. Bresch, P.-E. Jabin, and Z. Wang. *Mean-field limit and quantitative estimates with singular attractive kernels*, Duke Mathematical Journal, 2023.
- [4] Z. Brzezniak and G. Dhariwal, *Stochastic constrained Navier–Stokes equations on T^2* Journal of Differential Equations, 2021.
- [5] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2011.
- [6] R. Carmona and F. Delarue, *Probabilistic Theory of Mean Field Games with Applications. I: Mean Field FBSDEs, Control, and Games*. vol. 84 of Probability Theory and Stochastic Modelling, Springer, Cham, 2018.
- [7] ———, *Probabilistic theory of Mean Field Games with Applications II: Mean Field Games with Common Noise and Master Equations*, vol. 84 of Probability Theory and Stochastic Modelling, Springer, Cham, 2018.

- [8] J. A. Carrillo, F. Xuanrui Feng, S. Guo, P. E. Jabin, and Z. Wang. *Relative entropy method for particle approximation of the Landau equation for Maxwellian molecules*, Preprint arXiv:2408.15035, 2024.
- [9] P. Cattiaux, *Entropy on the Path Space and Application to Singular Diffusions and Mean-field Models*, Preprint arXiv:2404.09552, 2024.
- [10] L. Chen, E. S. Daus, A. Holzinger A. Jüngel, *Rigorous Derivation of Population Cross-Diffusion Systems from Moderately Interacting Particle Systems*, Journal of Nonlinear Science, 31, 2021.
- [11] L. Chen, A. Holzinger and X. Huo, *Quantitative convergence in relative entropy for a moderately interacting particle system on \mathbb{R}^d* , Preprint arXiv:2311.01980, 2023.
- [12] L. Chen, A. Holzinger, A. Jungel, N. Zamponi, *Analysis and mean-field derivation of a porous-medium equation with fractional diffusion*, Communications in Partial Differential Equations, 47, 2022.
- [13] L. Chen, P. Nikolaev and D.J. Promel, *Hegselmann-Krause model with environmental noise*, Trans. Amer. Math. Soc., DOI: <https://doi.org/10.1090/tran/9289>, 2024.
- [14] M. Coghi and F. Flandoli, *Propagation of chaos for interacting particles subject to environmental noise*, Ann. Appl. Probab. 26, 2016.
- [15] J. Correa and C. Olivera, *From Stochastic Hamiltonian Systems to Stochastic Compressible Euler Equation*, Preprint arXiv:2301.08101, 2023.
- [16] R. Cortez and J. Fontbona. *Quantitative uniform propagation of chaos for Maxwell molecules*. Comm. Math. Phys., 357, 2018.
- [17] F. Flandoli, *Random perturbation of PDEs and fluid dynamic models*. Lectures from the 40th Probability Summer School held in Saint-Flour, 2010. Lecture Notes in Mathematics, 2015. Springer, Heidelberg, 2011.
- [18] F. Flandoli, M. Ghio and G. Livieri, *N-player games and mean field games of moderate interactions*, Appl. Math. Optim., 85, 2022.
- [19] F. Flandoli, M. Leimbach and C. Olivera, *Uniform convergence of proliferating particles to the FKPP equation*, J. Math. Anal. Appl., 473, 2019.
- [20] F. Flandoli and M. Leocata, *A particle system approach to aggregation phenomena*, J. Appl. Probab., 56, 2019.
- [21] F. Flandoli and M. Leocata, *The Navier-Stokes-Vlasov-Fokker-Planck system as a scaling limit of particles in a fluid*, J. Math. Fluid Mech., 23, 2021.

- [22] F. Flandoli and E. Luongo, *Stochastic Navier–Stokes Equations and State-Dependent Noise*. In: *Stochastic Partial Differential Equations in Fluid Mechanics*, Lecture Notes in Mathematics, Springer, 2330, 2023.
- [23] F. Flandoli, C. Olivera and M. Simon, *Uniform approximation of 2 dimensional Navier-Stokes equation by stochastic interacting particle systems*, SIAM J. Math. Anal., 52, 2020.
- [24] N. Fournier, M. Hauray and S. Mischler. *Propagation of chaos for the 2D viscous vortex model*. J. Eur. Math. Soc. (JEMS), 16(7):1423–1466, 2014.
- [25] N. Fournier and B. Jourdain. *Stochastic particle approximation of the Keller–Segel equation and two-dimensional generalization of Bessel processes*, *Ann. Appl. Probab.*, 27, 2017.
- [26] N. Fournier, Y. Tardy, *Collisions of the supercritical Keller–Segel particle system*, Journal of the European Mathematical Society, 2024.
- [27] A. Guillin, P. Le Bris and P. Monmarche, *Uniform in time propagation of chaos for the 2D vortex model and other singular stochastic systems*, J. Eur. Math. Soc, 2024.
- [28] S. Guo and D. Luo, *Scaling limit of moderately interacting particle systems with singular interaction and environmental noise*, The Annals of Applied Probability, 33, 2023.
- [29] Z. Hao, J.F. Jabir, S. Menozzi and M. Rockner, *Propagation of chaos for moderately interacting particle systems related to singular kinetic McKean-Vlasov SDEs*, Preprint arXiv:2405.09195, 2024.
- [30] D.D. Holm, *Variational principles for stochastic fluid dynamics*, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences, 471, 2016.
- [31] H. Huang and J. Qiu, *The microscopic derivation and well-posedness of the stochastic Keller-Segel equation*, J. Nonlin. Sci., 31, 2021.
- [32] P.-E. Jabin and Z. Wang. *Quantitative estimates of propagation of chaos for stochastic systems with $W^{-1,\infty}$ kernels*. Invent. Math., 214, 2018.
- [33] B. Jourdain and S. Méléard, *Propagation of chaos and fluctuations for a moderate model with smooth initial data*, Ann. Inst. H. Poincaré Probab. Statist., 34, 1998.
- [34] J. Knorst, C. Olivera and A. B. de Souza, *Convergence Rate for Moderate Interaction particles and Application to Mean Field Games*, Preprint arXiv:2402.13167, 2024.
- [35] N. V. Krylov, *An analytic approach to SPDEs*. In *Stochastic partial differential equations: six perspectives*, Math. Surveys Monogr., Amer. Math. Soc., Providence, RI, 1999.

- [36] D. Lacker and L. Le Flem, *n-Closed-loop convergence for mean field games with common noise*, The Annals of Applied Probability, 2023.
- [37] A.J. Majda, I. Timofeyev and E. Vanden-Eijnden, *A mathematical framework for stochastic climate models*, Communications on Pure and Applied Mathematics, 54, 2001.
- [38] S. Méléard and S. Roelly-Coppoletta, *A propagation of chaos result for a system of particles with moderate interaction*, Stochastic Process. Appl., 26, 1987.
- [39] Q.H. Nguyen, M. Rosenzweig, S. Serfaty, *Mean-field limits of Riesz-type singular flows*, Ars Inveniendi Analytica, July 2022.
- [40] K. Oelschläger, *A martingale approach to the law of large numbers for weakly interacting stochastic processes*, Ann. Probab., 12, 1984.
- [41] K. Oelschläger, *A law of large numbers for moderately interacting diffusion processes*, Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete, 69, 1985.
- [42] K. Oelschläger, *A fluctuation theorem for moderately interacting diffusion processes*, Probab. Theory Related Fields, 74, 1987.
- [43] C. Olivera, A. Richard and M. Tomašević, *Quantitative particle approximation of nonlinear Fokker-Planck equations with singular kernel*, Ann. Sc. Norm. Super. Pisa Cl. Sci., 5, 2021.
- [44] C. Olivera, A. Richard and M. Tomašević, *Quantitative approximation of the Burgers and Keller-Segel equations by moderately interacting particles*, Preprint arXiv:2004.03177, 2023.
- [45] C. Olivera and M. Simon, *Microscopic derivation of non-local models with anomalous diffusions from stochastic particle systems*, Preprint arXiv:2404.03772, to appear Nonlinear Analysis, 2024.
- [46] P. Nikolaev, *Quantitative relative entropy estimates for interacting particle systems with common noise*, Preprint arXiv:2407.01217, 2024.
- [47] S. Serfaty, *Mean field limit for Coulomb-type flows*. Duke Math. J., 169, 2020.
- [48] H.-J. Schmeisser and H. Triebel, *Topics in Fourier analysis and function spaces*, A Wiley-Interscience Publication, John Wiley & Sons, Ltd., Chichester, 1987.
- [49] Y. Shao and X. Zhao, *Quantitative particle approximations of stochastic 2d navier-stokes equation*, Preprint arXiv:2402.02336, 2024.
- [50] M. Simon and C. Olivera, *Non-local conservation law from stochastic particle systems*, J. Dynam. Differential Equations, 30, 2018.

- [51] A. Stevens, *The derivation of chemotaxis equations as limit dynamics of moderately interacting stochastic many-particle systems*, SIAM Journal on Applied Mathematics, 61, 2000.
- [52] M. Tomasevic, D Talay, *A new McKean-Vlasov stochastic interpretation of the parabolic-parabolic Keller-Segel model: The one-dimensional case*, Bernoulli 26, 2020.
- [53] H. Triebel, *Interpolation theory, function spaces, differential operators*, Johann Ambrosius Barth, Heidelberg, second ed., 1995.