# Model-independent constraints on Lorentz Invariance Violation with update observations of Gamma-Ray Bursts

Yu Pan<sup>1</sup> \*, Jun Tian<sup>1,2</sup>, Shuo Cao<sup>3,4</sup> †, Qing-Quan Jiang<sup>2</sup>, Wei-Liang Qian<sup>5,6</sup> 1. School of Science, Chongqing University of Posts and Telecommunications, Chongqing 400065, China;

2. School of Physics and Astronomy,

China West Normal University, Nanchong 637002, China;

- 3. Institute for Frontiers in Astronomy and Astrophysics, Beijing Normal University, Beijing 102206, China;
  - 4. School of Physics and Astronomy, Beijing Normal University, Beijing 100875, China;
- 5. Escola de Engenharia de Lorena, Universidade de São Paulo 12602-810, Lorena, SP, Brazil;
  - 6. Center for Gravitation and Cosmology, College of Physical Science and Technology, Yangzhou University, Yangzhou 225009, China

# Abstract

Searching the possible Lorentz Invariance Violation (LIV) from astrophysical sources such as gamma-ray bursts (GRBs) is essential for finding evidences of new theories of quantum gravity. However, the effect of the underlying cosmological model is still understudied in the previous analysis. We take a novel approach using artificial neural networks to reconstruct the expansion history of the universe, thereby eliminating the influence of potential cosmological models to constrain LIV. 74 time delays from GRBs are considered to obtain stringent results on LIV, including 37 time delays measurements from GRB 160625B across various energy bands at redshift z=1.41, and 37 additional GRBs with time delays spanning redshifts  $0.117 \le z \le 1.99$ . Our analysis yields stringent constraints on both linear and quadratic LIV, with  $E_{QG,1} \ge 2.63 \times 10^{15}$  GeV and  $E_{QG,2} \ge 1.19 \times 10^{10}$  GeV that are four orders of magnitude beneath the Planck energy scale, and shows the positive intrinsic time delay in GRBs. Our results demonstrate that such combination would significantly improve the precision and robustness of final results. Taking this into account may be an important contribution in the case of possible LIV detection in the future.

<sup>\*</sup> Corresponding author: panyu@cqupt.edu.cn

<sup>†</sup> Corresponding author: caoshuo@bnu.edu.cn

## I. INTRODUCTION

The Lorentz Invariance holds crucial significance in physics and is a fundamental assumption of Relativity. However, several quantum gravity (QG) theories suggest that Lorentz Invariance Violation (LIV) may occur at the Planck energy scale,  $E_{QG} \approx E_{P1} = \sqrt{\hbar c^5/G} \simeq 1.22 \times 10^{19}$  GeV [1, 2]. From an astronomical point of view, the observational Gamma-Ray Bursts (GRBs) can capture this microcosmic quantum gravity. The GRBs are the brightest astrophysical events in the Universe, characterized by extensive distances, brief spectral lags, and energetic emissions [2–4]. They stand out as the most promising candidates for conducting LIV tests [2]. Over the past two decades, GRBs have served as the primary source for a majority of LIV tests [5, 6].

GRBs are divided into two types based on their duration: Short Gamma-ray bursts (SGRBs), lasting less than two seconds, and Long Gamma-ray bursts (LGRBs), lasting more than two seconds [7]. SGRBs are commonly linked to neutron star mergers [8], while LGRBs are typically associated with core-collapse supernovae [9]. LGRBs typically exhibit more pronounced positive or negative spectral lags compared to SGRBs [10–12], whereas SGRBs generally display smaller spectral lags [5, 12]. Spectral lag, a widely recognized phenomenon in GRBs, is characterized by a time delay between low-energy photons (i.e., light curves) and high-energy photons [13]. The primary approach for searching LIV with GRBs is analyzing these spectral lags [5].

The lower limits of  $E_{QG}$  in some LIV tests have approached the Planck energy scale, determined through the observation of spectral lag in extremely high-energy photons from individual GRBs [14–16]. However, spectral lag includes both the time delay associated with LIV and an unknown intrinsic time delay. Using a single burst to constrain LIV makes it challenging to disentangle them. Ellis et al. [3] addressed this issue by stacking a sample of GRBs, primarily LGRBs, and replacing the intrinsic time delay in the rest frame with an assumed constant for all GRBs. Nevertheless, it's unlikely that every GRB shares an identical intrinsic spectral lag. Wei et al. [17] employed a more realistic power-law model using multiple lags from GRB 106625B, explaining the transition in spectral lags from positive to negative. However, the data fitting in their study was not entirely satisfactory. Agrawal et al. [18] employed a similar approach, stacking a larger number of GRBs, which improved the precision of the results but also faced challenges related to the goodness of the data fitting.

There were a lot of work that choosed the cosmological models to constrain LIV such as the  $\Lambda$ CDM model. However, these models are based on the general relativity framework without considering LIV [19]. In a recent study, an investigation into the effects of different cosmological models on LIV constraints revealed the complexity of this relationship, adding more uncertainty to the results [20]. To address this issue, Pan et al. [19] first employed a model-independent method using the Gaussian Process (GP) to reconstruct cosmic expansion history (represented by the Hubble parameter H(z)) for constraints on LIV, their results are more robust and conservative. For more work on the use of GP, we refer to Refs [21–25]. Nevertheless, GP comes with various disadvantages, such as overfitting and a heavy dependence on the selection of covariance functions [26]. Recent research has also emphasized the need for caution when using GP for the reconstruction of the Hubble parameter [27]. On the other hand, the Artificial Neural Network (ANN) is a fully data-driven method that accurately captures input data distribution with a suitable network architecture [28]. Recent studies have increasingly utilized this method to reconstruct the Hubble parameter [26, 28–30].

In this paper, we propose a novel, model-independent method by using an Artificial Neural Network (ANN) to reconstruct the cosmic expansion history to constrain LIV, effectively circumvents the potential influence of cosmological models. The outline for methodology of LIV-induced time delay and the observational datasets are given in Section II. We show the results and discussions in Section III. Finally, the main conclusions are summarized in Section IV.

#### II. METHODOLOGY AND DATASETS

#### A. Lorentz Invariance Violation

In some QG models, the Lorentz invariance may be violated around the Planck energy scale  $E_{QG} \sim 10^{19} GeV$ , leading the speed of light to depend on the energy of photons [2, 31]. This phenomenon, which can be described through a modified dispersion relation (MDR), although it does not specify a particular theoretical framework, can prompt various experimental follow-up tests [32, 33]. In our work, we use an empirical formula for the MDR of massless particles, widely recognized for its robustness across various theoretical contexts [33]. This relation can be expressed as follows:

$$E^{2} = c^{2}p^{2} \left[ 1 + f(E/E_{QG}) \right], \tag{1}$$

Where  $E_{QG}$  denotes the Planck energy scale, f is the model dependent function of the dimensionless ratio of  $E/E_{QG}$ . The dispersion relation can be expanded as a Taylor series [34]:

$$E^2 \simeq c^2 p^2 \left[ 1 - s_{\pm} \sum_{n=1}^{\infty} \left( \frac{E}{E_{QG,n}} \right)^n \right], \tag{2}$$

For  $E_{QG,n} \gg E$ , the lowest order term in the series is expected to dominate the sum, so we keep only the lowest-order dominant term:

$$E^2 \simeq c^2 p^2 \left[ 1 - s_{\pm} \left( \frac{E}{E_{QG,n}} \right)^n \right], \tag{3}$$

Assuming the relation  $v(E) = \partial E/\partial p$  holds at least approximately, the energy-dependent velocity of light is [31]:

$$v(E) = c \left[ 1 - s_{\pm} \frac{n+1}{2} \left( \frac{E}{E_{QG}} \right)^n \right]. \tag{4}$$

Here,  $s_{\pm}=+1$  (or -1) indicates whether the photon velocity decreases (or increases) with the growth of photon energy, and n is a term dependent on the model, with n=1 (or 2) corresponding to a linear (or quadratic) term. Considering that the lower-energy photons are generally expected to travel faster than higher-energy ones due to LIV effects [17], we only focus on  $s_{\pm}=+1$  in this work.

Due to the relationship between the velocity of light and energy, photons of varying energies (represented by E and  $E_0$ , with  $E > E_0$ ) emitted by the identical origin will reach Earth at distinct moments. In addition, based on the recent astronomical observations, particularly of the observations of type Ia supernovae (SNe Ia), have shown that our universe contains a component with negative pressure that drives the accelerated expansion of the universe [35]. With the effects

of cosmological expansion, the time delay between photons with different energy levels induced by LIV is [4, 36]:

$$\Delta t_{LIV} = -\frac{1+n}{2H_0} \frac{E^n - E_0^n}{E_{QG,n}^n} \int_0^z \frac{(1+z')^n dz'}{h(z')},\tag{5}$$

where  $H_0$  stands for the Hubble constant, while  $h(z) = H(z)/H_0$  denotes the dimensionless Hubble expansion H(z) rate at redshift z. The complete time delay across various energy bands should be included five terms:

$$\Delta t = \Delta_{LIV} + \Delta_{int} + \Delta_{spe} + \Delta_{DM} + \Delta_{gra}, \tag{6}$$

where  $\Delta_{LIV}$  represents the potential time delay due to LIV via an energy-dependent velocity of light,  $\Delta_{int}$  is the intrinsic time delay between two test photons, and this has the largest uncertainty since it is impossible to determine it based on the observational data alone,  $\Delta_{spe}$  is the potential time delay due to special relativistic effects when the photons have a non-zero rest mass,  $\Delta_{DM}$  denote the time delay contribution from the dispersion by the line-of-sight free electron content, and  $\Delta_{gra}$  is the gravitational potential along the propagation path of photons if the Einstein equivalence principle is violated. However, recent works pointed that, for cosmic transient sources (like GRB), the last three terms have not significantly influence on the total time delay  $\Delta t$ , and can be ignored [37, 38]. Thus, we assume the  $\Delta t$  is the sum of  $\Delta_{LIV}$  and  $\Delta_{int}$ .

Specifically, we still have a limited understanding of the detailed emission mechanism of GRBs, so choosing the appropriate form of intrinsic time delay is very important. Intrinsic time delay is commonly represented by the parameter b, but considering the cosmological expansion  $\Delta_{int} = b(1+z)$  [3], this form implies that GRBs exhibit a uniform intrinsic time delay. For SGRBs, using b to represent intrinsic time delay may be reasonable due to small spectral lags [13]. However, LGRBs have more long spectral lags and are very likely to have different intrinsic time lags, the format of a power-law function is more reasonable [17]:

$$\Delta_{int}(E) = \tau \left[ \left( \frac{E}{keV} \right)^{\alpha} - \left( \frac{E_0}{keV} \right)^{\alpha} \right], \tag{7}$$

where  $\tau$  and  $\alpha$  are treated as free parameters and have the positive value, with  $E_0$  representing the midpoint of the lowest energy band. When both  $\tau$  and  $\alpha$  are greater than zero, the intrinsic time delay is positive. Finally, the theoretical time delay between different energy bands is expressed as follows:

$$\Delta t_{th} = \Delta_{LIV} + \Delta_{int}(1+z)$$

$$= -\frac{1+n}{2H_0} \frac{E^n - E_0^n}{E_{QG,n}^n} \int_0^z \frac{(1+z')^n dz'}{h(z')} + \tau \left[ \left( \frac{E}{keV} \right)^\alpha - \left( \frac{E_0}{keV} \right)^\alpha \right] (1+z). \tag{8}$$

Traditionally, considerable work has chosen a specific cosmological model like the flat  $\Lambda$ CDM model to obtain the function h(z). However, this is cosmological model-dependent. It is shown that the dimensionless Hubble parameter functions h(z) of different cosmological models differ significantly at high redshift, which will significantly affect our study of LIV. The test of LIV by using cosmological model independent method is particularly important.

#### **B.** Observational Data

In this paper, we have selected a sample of 74 time delay measurements for GRBs (mostly LGRBs), which comprises three datasets analyzed in [3, 13, 17]. These time delays were obtained by measuring the time lags between various energy bands within their respective light curves.

The first dataset is presented in Table 1 of Ellis et al. [3]. This table contains 35 time delay GRBs for various redshift, primarily LGRBs, within the redshift range of  $0.168 \sim 6.29$ . Among these GRBs, there are 9 GRBs from BATSE data (time resolution: 64 ms), 15 GRBs from HETE data (time resolution: 164 ms), and 11 GRBs from SWIFT data (time resolution: 64 ms). Spectral lags were calculated by analyzing the light curves, comparing the 115-350 keV energy band with the 25-55 keV energy band. However, it should be noted that in our analysis, we have only selected the data with redshifts less than 2. The reason for this choice will be explained in the next subsection. Therefore, the first dataset consists of 21 time delays with a redshift range of  $0.168 \sim 1.99$ .

The second dataset consists of 37 time delays from GRB 160625B at various energy bands (all with a common redshift value of z=1.41) is presented in Table 1 of Wei et al. [17] GRB 160625B was first identified by the Fermi Gamma-Ray Burst Monitor (GBM) and subsequently confirmed by the Fermi Large Area Telescope (LAT). GRB 160625B's light curve displays three distinct sub-pulses and the total duration of approximately  $T_{90}=770\mathrm{s}$  [39], which means this burst is LGRB. Specifically, the sub-pulse number two is exceptionally luminous, enabling straightforward extraction of the light curves in various energy bands. By analyzing the light curves, the spectral lags were calculated using the CCF method in the lowest energy band (10-12~keV) relative to other higher energy bands (15-350~keV).

The last dataset is presented in Table 3 of Xiao et al. [13] This table contains data on 21 time delay LGRBs, with redshifts covering the range from z = 0.117 to z = 2.938 (redshifts obtained Swift Burst Analyser [40], Fermi GBM Burst Catalog [41], and GCN circulars of the Swift and GBM teams). These LGRBs are observed by GMB, BAT, and GECAMB detectors, but only BAT observed all LGRBs, so we chose to utilize BAT data for our analysis. These spectral lags were extracted using the Li-CCF method, which calculates the spectral lags in light curves recorded in energy bands ranging from  $15 - 70 \ keV$  to  $120 - 250 \ keV$ . As previously mentioned, we limited our selection to data with redshifts less than 2, resulting in a dataset of 16 time delays with redshifts in the range of  $0.117 \sim 1.937$ .

Finally, the combined dataset consists of 37 time delays from GRB 160625B at various energy bands, all sharing a common redshift value of z=1.41 [17], as well as 37 time delays from various other GRBs with redshift values ranging from z=0.117 to z=1.99 [3, 13].

# C. Artificial Neural Network Method

Artificial Neural Network (ANN) is a fully data-driven approach, providing a model-independent method. We employ an ANN to reconstruct the cosmic expansion history, represented by the Hubble parameter H(z). The Python code employed for this task is ReFANN, developed by Wang et al. [28], a single hidden layer comprising a total of 4096 neurons. ReFANN is constructed using PyTorch, a widely recognized and well-optimized open-source tensor library designed for




FIG. 1: The reconstructed H(z) (shown as a red line) using ANN, along with the corresponding  $1\sigma$  error (depicted as a sky-blue area), and the CC H(z) data (represented by black dots with error bars).

machine learning. To ensure the complete model independence of the H(z) functions, it is essential to carefully select the data, avoiding any influence from the cosmological model. The values of H(z) can be acquired using two distinct methods. The first method is the cosmic chronometer method, which calculates the age differences among galaxies undergoing passive evolution at various redshifts [42]. The Hubble parameter of the cosmic chronometer is denoted as CC H(z), and they offer model-independent measurements of H(z) [42]. Another one is through the detection of radial BAO features [29, 43, 44]. However, the values acquired by this method depend on a presupposed fiducial cosmological model. Given the requirement for model-independent H(z) reconstruction, we choose the CC H(z) data for this purpose. We utilized the latest collection of 32 data points of CC H(z) [45–49], spanning the redshift range extending to approximately z=2.

In Figure 1, we show the outcomes derived from the ANN method, where the reconstructed Hubble parameter is denoted as ANN H(z). In this figure, black dots with error bars represent CC H(z) data, and the red lines and sky-blue areas indicate the ANN H(z)'s best values and  $1\sigma$  errors. It's important to note that all CC H(z) data fall within the sky-blue areas, taking into account the  $1\sigma$  error, which indicates the success of ANN H(z). The reconstructed Hubble constant is determined as  $H_0 = 68.48 \pm 16.61 \ km \ s^{-1} \ Mpc^{-1}$ , a value very close to the Planck CMB result:  $H_0 = 67.4 \ km \ s^{-1} \ Mpc^{-1}$  [50]. This similarity further validates the accuracy of the reconstructed Hubble parameter. Therefore, the dimensionless Hubble expansion rate  $h(z) = \text{ANN } H(z)/H_0$ . However, please be aware that the CC H(z) data is limited to a redshift of 2. Therefore, we limited ANN H(z) to redshifts around 2 [26]. Consequently, we exclusively selected GRB data with redshifts less than 2 for our analysis. For more work on the data reconstruction of cosmological-independent models by using ANN method, we refer readers to Refs [51–56].




FIG. 2: The 1D probability distribution of each parameter and the 2D confidence contours for the parameters  $E_{QG}$ ,  $\tau$ , and  $\alpha$  (the linear LIV case, i.e., n = 1).

## III. RESULTS AND DISCUSSION

We have applied the ANN method to exclude the effects of the cosmological model. Next, we employ the Monte Carlo Markov Chain method [57] to determine the LIV parameters ( $E_{QG}$ ,  $\tau$ , and  $\alpha$ ) by minimizing the  $\chi^2$  objective function:

$$\chi^2 = \sum_{i=1}^{N_{GRB}} \left[ \frac{\Delta t_{th} \left( E_{QG}, \tau, \alpha \right) - \Delta t_{obs}}{\sigma_{tot}} \right]^2, \tag{9}$$

where  $\Delta t_{th}$  represents the theoretical time delay,  $\Delta t_{obs}$  stands for the observed time delay, and  $\sigma_{tot}$  denotes the total uncertainty.

$$\sigma_{tot}^2 = \sigma_t^2 + (\frac{\partial f}{\partial E})^2 \sigma_E^2 + (\frac{\partial f}{\partial E_0})^2 \sigma_{E_0}^2, \tag{10}$$

where f is Eq. 5,  $\sigma_t$  is the observed time delay uncertainty, and  $\sigma_{E_0}$  and  $\sigma_E$  represent the half-width of the lower and higher energy bands, respectively. The posterior probability distribution for each parameters and the corresponding two-dimensional confidence contours are plotted in Figures 2 and 3.

We begin our discussion with the linear LIV case (i.e., n=1). The constraints for  $\tau$ ,  $\alpha$ , and  $E_{QG,1}$  are depicted in Figure 2. The best-fit parameter values along with their corresponding  $1\sigma$ 

uncertainties are  $\log_{10} E_{QG,1} = 15.660^{+0.110}_{-0.240}~GeV,~\tau = 0.441^{+0.057}_{-0.270},~{\rm and}~\alpha = 0.208^{+0.054}_{-0.054}.$  The lower bound for LIV at the  $1\sigma$  confidence level is set at  $E_{OG,1} \geq 2.63 \times 10^{15} \ GeV$ , a value that is four orders of magnitude beneath the Planck energy scale. Although it is well below the Planck energy scale, it is similar to that of Xiao et al. [13], who obtained  $E_{QG,1} \ge 2.5 \times 10^{15} \ GeV$  through an analysis of 46 SGRBs. The best-fit values,  $\tau \sim 0.441$  and  $\alpha \sim 0.208$ , and both parameters lie within the  $1\sigma$  range as positive values, indicating a positive intrinsic lag in our analysis. Most of the GRB pulses are characterized by positive lags [58, 59], our results align with the observational data. On the other hand, these results are similar to the analyses conducted by Wei et al. [17] and Du et al. [60]. From the Eq. 8, we see that as the photon's energy increases, the LIV lag decreases while the intrinsic lag increases. The relationship between them is key to studying LIV and the spectral lag of GRB. At low energy levels, intrinsic lag dominates the time lag. However, with increasing energy, the LIV lag becomes dominant, leading to a transition in spectral lag. Wei et al. [17] used the power-law model and LIV to explain the spectral lag transition of GRB 160625B from positive to negative (Du et al. [60] explained the transition of GRB 190114C using similar methods), which the constant model cannot achieve. However, the GRB they chose is special, which exhibited a transition in the spectral lags from positive to negative, mostly GRBs don't exhibit this transition. In our analysis, we chose a large sample of various GRBs. Evidently, the spectral lags of different GRBs provide more robust limits on LIV. Compared to the earlier findings from the single special source GRB 160625B [17] and GRB 190114C [60], our results exhibit higher precision and impose better constraints on the parameters.

In the quadratic LIV case (i.e., n=2), the best-fit parameter values along with their corresponding  $1\sigma$  uncertainties are  $\log_{10}E_{QG,2}=10.153^{+0.047}_{-0.077}~GeV$ ,  $\tau=0.710^{+0.100}_{-0.380}$ , and  $\alpha=0.150^{+0.036}_{-0.036}$ . The 1D probability distributions and 2D contours with  $1\sigma$  and  $2\sigma$  confidence levels for  $\tau$ ,  $\alpha$ , and  $E_{QG,2}$  are also illustrated in Figure 3. The  $1\sigma$  confidence-level lower boundary for LIV is  $E_{QG,2}\geq 1.19\times 10^{10}~GeV$ , a value within the same order of magnitude as the obtained from the single high energy GRB 190114C [16]. The two parameters  $\tau$  and  $\alpha$  largely align with the corresponding values in both cases. Additionally, a significant correlation is observed between the parameters  $\tau$  and  $\alpha$  in both cases of the power-law model. The parameters  $E_{QG,2}$ ,  $\tau$  and  $\alpha$  still have higher precision than the single GRB [17, 60]. Meanwhile, compared with the previous work using the GP method [19], the results of ANN don't exhibit better precision. It is not surprising, to ensure a fully model-independence, we only utilize 32 CC H(z), without taking into the H(z) obtained through the detection of radial BAO features. So, we hope that more H(z) values can be measured to expand the redshift range of ANN H(z) in the future. However, the parameters  $\tau$  and  $\alpha$  exhibit a more general range of values compared to the results of GP, not being excessively large or too small.

These are the main reasons for our choice of the power-law model and the majority of time delay measurements of LGRBs. On one hand, it is not reasonable to assume the intrinsic lag as an unknown constant, especially when considering the fact that LGRBs exhibit long spectral lags. Although some researchers exclusively use the constant model for a sample of SGRBs, which may be reasonable, primarily due to the negligible spectral lags of SGRBs [5]. However, considering the wide range of variations observed in GRB light curves [61], assuming same time lags appears implausible and could potentially introduce systematic effects [5]. The power-law model was initially employed by Wei et al. [17] to explain the transition in the spectral lag of GRBs, it is a function of energy. Clearly, this model is more reasonable and provides a better description of




FIG. 3: The 1D probability distribution of each parameter and the 2D confidence contours for the parameters  $E_{QG}$ ,  $\tau$ , and  $\alpha$  (the linear LIV case, i.e., n = 2).

intrinsic lags when compared to the constant model [17, 19]. On the other hand, the power-law model was deduced empirically by examining spectral lag characteristics in a dataset of 50 single-pulsed GRBs [62]. Please note that the dataset consists of only one short GRB, while the others are long GRBs. Lastly, a recent study by Desai et al. [5] utilized this power-law model in 46 SGRBs to constrain LIV, but they did not achieve a favorable outcome. The values of  $\tau$  and  $\alpha$  are centered around zero within a  $1\sigma$  range, along with the inability to obtain a closed contour for  $E_{QG}$ , possibly due to SGRBs' short spectral lag and different emission mechanism.

## IV. CONCLUSIONS

The search of possible Lorentz Invariance Violation (LIV) from astrophysical sources such as gamma-ray bursts (GRBs) is essential for finding evidences of new theories of quantum gravity. We analyze 74 time delays from GRBs, including 37 measurements from GRB 160625B across various energy bands at redshift z=1.41, and 37 additional GRBs with time delays spanning redshifts from z=0.117 to z=1.99. However, the effect of the underlying cosmological model is still understudied in the previous analysis. In this work, we adopt a novel approach by utilizing an Artificial Neural Network (ANN) to reconstruct the cosmic expansion history. Specially, the newest observations of cosmic chronometers acting as the training set ensures a strict model-independent methodology. Our choice of training data encompasses 32 data points of the cosmic

chronometer H(z) (CC H(z)). This selection ensures that the resulting reconstruction of the cosmic expansion history remains independent of any specific theoretical model. Furthermore, we include the power-law model and discuss the searches for LIV in both linear and quadratic cases.

By using the power-law model and stacking of a large sample, we have established constraints for both linear and quadratic cases of LIV:  $E_{QG,1} \geq 2.63 \times 10^{15}~GeV$  and  $E_{QG,2} \geq 1.19 \times 10^{10}~GeV$ . Notably, in the quadratic case, the value of  $E_{QG,2}$  within the same order of magnitude as that obtained from high energy photon[16]. The parameters  $\tau$  and  $\alpha$  of the power-law model are similar to the results of GRB 160625B [17]. Compared to using the time delays of a single special GRB, our findings present higher precision and improved constraints, different GRBs provide more robust limits on LIV. However, compared with our previous work using the GP method [19], the results of ANN have a lower precision. It is not surprising, because we reconstruct the cosmic expansion history only based on 32 CC H(z), which ensures it is fully cosmological-model independent. Meanwhile, we also explain the main reasons why we choose the power-law model and the choice GRBs are almost LGRBs. On the one hand, the power-law model is more reasonable than the constant model, it is a function of energy that can explain the transition of spectral lags [17, 60]. On the other hand, some researchers have used this model in a sample of SGRBs, but their results didn't provide good constraints on LIV [5].

Finally, the GRBs often have high redshifts and the reconstructed H(z) is constrained within a range up to 2, we must exclude some data, which could introduce unforeseen effects on the results. In the future, we hope to obtain more H(z) values to extend the redshift range of ANN H(z). More importantly, we have observed a significant correlation between  $\tau$  and  $\alpha$  in the power-law model, which may indicate a potential issue with this model.

# Acknowledgments

We thank Shao-Qi Hou and Marek Biesiada for the helpful discussion on this paper. This work was supported by National Natural Science Foundation of China Grant Nos. 12105032, 12203009, 12021003, 11690023, and 11920101003; the Natural Science Foundation of Chongqing 18 No. cstc2021jcyj-msxmX0481; the Strategic Priority Research Program of the Chinese Academy of Sciences, Grant No. XDB23000000; the Central Guidance on Local Science and Technology Development Fund of Sichuan Province (24ZYZYTS0188); the Interdiscipline Research Funds of Beijing Normal University.

- [1] David Mattingly, Living Reviews in Relativity 8 (2005) 5.
- [2] Giovanni Amelino-Camelia, Living Reviews in Relativity 16 (2013) 5.
- [3] John Ellis, N. E. Mavromatos, D. V. Nanopoulos, A. S. Sakharov, and E. K. G. Sarkisyan, Astroparticle Physics 25 (2006) 402.
- [4] Uri Jacob and Tsvi Piran, JCAP2008 (2008) 031.
- [5] Shantanu Desai, arXiv:2303.10643.
- [6] Kai Liao, Marek Biesiada, and Zong-Hong Zhu, Chinese Physics Letters 39 (2022) 119801.

- [7] Chryssa Kouveliotou, Charles A. Meegan, Gerald J. Fishman, Narayana P. Bhat, Michael S. Briggs, Thomas M. Koshut, William S. Paciesas, and Geoffrey N. Pendleton, Astrophysical Journal Letters 413 (1993) L101.
- [8] Ehud Nakar, Physical Review Letters 442 (2007) 166.
- [9] S. E. Woosley and J. S. Bloom, Annual Review of Astronomy and Astrophysics 44 (2006) 507.
- [10] Li Chen, Yu-Qing Lou, Mei Wu, Jin-Lu Qu, Shu-Mei Jia, and Xue-Juan Yang, Astrophys. J. 619 (2005) 983.
- [11] T. N. Ukwatta, K. S. Dhuga, M. Stamatikos, C. D. Dermer, T. Sakamoto, E. Sonbas, W. C. Parke, L. C. Maximon, J. T. Linnemann, P. N. Bhat, A. Eskandarian, N. Gehrels, A. U. Abeysekara, K. Tollefson, and J. P. Norris, Mon. Not. Roy. Astron. Soc.419 (2012) 614.
- [12] M. G. Bernardini, G. Ghirlanda, S. Campana, S. Covino, R. Salvaterra, J. L. Atteia, D. Burlon, G. Calderone, P. D'Avanzo, V. D'Elia, G. Ghisellini, V. Heussaff, D. Lazzati, A. Melandri, L. Nava, S. D. Vergani, and G. Tagliaferri, Mon. Not. Roy. Astron. Soc.446 (2015) 1129.
- [13] Shuo Xiao, Shao-Lin Xiong, Yue Wang, Shuang-Nan Zhang, He Gao, Zhen Zhang, Ce Cai, Qi-Bin Yi, Yi Zhao, You-Li Tuo, Xin-Qiao Li, Xiang-Yang Wen, Zheng-Hua An, Wen-Xi Peng, Shi-Jie Zheng, Fan Zhang, Li-Ming Song, and Ti-Pei Li, Astrophysical Journal Letters 924 (2022) L29.
- [14] A. A. Abdo, M. Ackermann, M. Ajello, K. Asano, Fermi LAT Collaboration, et al., Nature (London) 462 (2009) 331.
- [15] A. A. Abdo, M. Ackermann, M. Arimoto, Fermi LAT Collaboration, Fermi GBM Collaboration, et al., Science 323 (2009) 1688.
- [16] V. A. Acciari, S. Ansoldi, L. A. Antonelli, MAGIC Collaboration, et al., Phys. Rev. Lett. 125 (2020) 021301.
- [17] Jun-Jie Wei, Bin-Bin Zhang, Lang Shao, Xue-Feng Wu, and Peter Mészáros, Astrophysical Journal Letters 834 (2017) L13.
- [18] Rajdeep Agrawal, Haveesh Singirikonda, and Shantanu Desai, JCAP2021 (2021) 029.
- [19] Yu Pan, Jingzhao Qi, Shuo Cao, Tonghua Liu, Yuting Liu, Shuaibo Geng, Yujie Lian, and Zong-Hong Zhu, Astrophys. J. 890 (2020) 169.
- [20] Denitsa Staicova, Classical and Quantum Gravity 40 (2023) 195012.
- [21] Suhail Dhawan, Justin Alsing, and Sunny Vagnozzi, Mon. Not. Roy. Astron. Soc. 506 (2021) L1.
- [22] Hai Yu, Bharat Ratra, and Fa-Yin Wang, Astrophys. J. 856 (2018) 3.
- [23] Tonghua Liu, Shuo Cao, Jia Zhang, Shuaibo Geng, Yuting Liu, Xuan Ji, and Zong-Hong Zhu, Astrophys. J. 886 (2019) 94.
- [24] Tonghua Liu and Kai Liao, Mon. Not. Roy. Astron. Soc. 528 (2024) 1354.
- [25] Arianna Favale, Adrià Gómez-Valent, and Marina Migliaccio, Mon. Not. Roy. Astron. Soc.523 (2023) 3406.

- [26] Purba Mukherjee, Jackson Levi Said, and Jurgen Mifsud, JCAP2022 (2022) 029.
- [27] Huan Zhou and Zhengxiang Li, Chinese Physics C 43 (2019) 035103.
- [28] Guo-Jian Wang, Xiao-Jiao Ma, Si-Yao Li, and Jun-Qing Xia, Astrophys. J., Suppl. Ser.246 (2020) 13.
- [29] Isidro Gómez-Vargas, Ricardo Medel-Esquivel, Ricardo García-Salcedo, and J. Alberto Vázquez, European Physical Journal C 83 (2023) 304.
- [30] Jing-Zhao Qi, Ping Meng, Jing-Fei Zhang, and Xin Zhang, Phys. Rev. D 108 (2023) 063522.
- [31] G. Amelino-Camelia, John Ellis, N. E. Mavromatos, D. V. Nanopoulos, and Subir Sarkar, Nature (London) 395 (1998) 525.
- [32] V. Alan Kostelecký and Matthew Mewes, Astrophys. J. Lett. 689 (2008) L1.
- [33] A. Addazi, J. Alvarez-Muniz, R. Alves Batista, G. Amelino-Camelia, V. Antonelli, M. Arzano, M. Asorey, J. L. Atteia, S. Bahamonde, F. Bajardi, A. Ballesteros, B. Baret, D. M. Barreiros, S. Basilakos, D. Benisty, O. Birnholtz, J. J. Blanco-Pillado, D. Blas, J. Bolmont, D. Boncioli, P. Bosso, G. Calcagni, S. Capozziello, J. M. Carmona, S. Cerci, M. Chernyakova, S. Clesse, J. A. B. Coelho, S. M. Colak, J. L. Cortes, S. Das, V. D'Esposito, M. Demirci, M. G. Di Luca, A. di Matteo, D. Dimitrijevic, G. Djordjevic, D. Dominis Prester, A. Eichhorn, J. Ellis, C. Escamilla-Rivera, G. Fabiano, S. A. Franchino-Viñas, A. M. Frassino, D. Frattulillo, S. Funk, A. Fuster, J. Gamboa, A. Gent, L. Á. Gergely, M. Giammarchi, K. Giesel, J. F. Glicenstein, J. Gracia-Bondía, R. Gracia-Ruiz, G. Gubitosi, E. I. Guendelman, I. Gutierrez-Sagredo, L. Haegel, S. Heefer, A. Held, F. J. Herranz, T. Hinderer, J. I. Illana, A. Ioannisian, P. Jetzer, F. R. Joaquim, K. H. Kampert, A. Karasu Uysal, T. Katori, N. Kazarian, D. Kerszberg, J. Kowalski-Glikman, S. Kuroyanagi, C. Lämmerzahl, J. Levi Said, S. Liberati, E. Lim, I. P. Lobo, M. López-Moya, G. G. Luciano, M. Manganaro, A. Marcianò, P. Martín-Moruno, Manel Martinez, Mario Martinez, H. Martínez-Huerta, P. Martínez-Miravé, M. Masip, D. Mattingly, N. Mavromatos, A. Mazumdar, F. Méndez, F. Mercati, S. Micanovic, J. Mielczarek, A. L. Miller, M. Milosevic, D. Minic, L. Miramonti, V. A. Mitsou, P. Moniz, S. Mukherjee, G. Nardini, S. Navas, M. Niechciol, A. B. Nielsen, N. A. Obers, F. Oikonomou, D. Oriti, C. F. Paganini, S. Palomares-Ruiz, R. Pasechnik, V. Pasic, C. Pérez de los Heros, C. Pfeifer, M. Pieroni, T. Piran, A. Platania, S. Rastgoo, J. J. Relancio, M. A. Reyes, A. Ricciardone, M. Risse, M. D. Rodriguez Frias, G. Rosati, D. Rubiera-Garcia, H. Sahlmann, M. Sakellariadou, F. Salamida, E. N. Saridakis, P. Satunin, M. Schiffer, F. Schüssler, G. Sigl, J. Sitarek, J. Solà Peracaula, C. F. Sopuerta, T. P. Sotiriou, M. Spurio, D. Staicova, N. Stergioulas, S. Stoica, J. Strišković, T. Stuttard, D. Sunar Cerci, Y. Tavakoli, C. A. Ternes, T. Terzić, T. Thiemann, P. Tinyakov, M. D. C. Torri, M. Tórtola, C. Trimarelli, T. Trześniewski, A. Tureanu, F. R. Urban, E. C. Vagenas, D. Vernieri, V. Vitagliano, J. C. Wallet, and J. D. Zornoza, Progress in Particle and Nuclear Physics 125 (2022) 103948.
- [34] V. Vasileiou, A. Jacholkowska, F. Piron, J. Bolmont, C. Couturier, J. Granot, F. W. Stecker, J. Cohen-Tanugi, and F. Longo, Phys. Rev. D 87 (2013) 122001.

- [35] Adam G. Riess, Alexei V. Filippenko, Peter Challis, Alejandro Clocchiatti, Alan Diercks, Peter M. Garnavich, Ron L. Gilliland, Craig J. Hogan, Saurabh Jha, Robert P. Kirshner, B. Leibundgut, M. M. Phillips, David Reiss, Brian P. Schmidt, Robert A. Schommer, R. Chris Smith, J. Spyromilio, Christopher Stubbs, Nicholas B. Suntzeff, and John Tonry, Astron. J.116 (1998) 1009.
- [36] Shu Zhang and Bo-Qiang Ma, Astroparticle Physics 61 (2015) 108.
- [37] He Gao, Xue-Feng Wu, and Peter Mészáros, Astrophys. J. 810 (2015) 121.
- [38] Jun-Jie Wei, He Gao, Xue-Feng Wu, and Peter Mészáros, Phys. Rev. Lett. 115 (2015) 261101.
- [39] B. B. Zhang, B. Zhang, A. J. Castro-Tirado, et al., Nature Astronomy 2 (2018) 69.
- [40] P. A. Evans, A. P. Beardmore, K. L. Page, et al., Mon. Not. Roy. Astron. Soc. 397 (2009) 1177.
- [41] A. von Kienlin, C. A. Meegan, W. S. Paciesas, et al., Astrophys. J. 893 (2020) 46.
- [42] Raul Jimenez and Abraham Loeb, Astrophys. J. 573 (2002) 37.
- [43] Enrique Gaztañaga, Anna Cabré, and Lam Hui, Mon. Not. Roy. Astron. Soc.399 (2009) 1663.
- [44] Lado Samushia, Beth A. Reid, Martin White, et al., Mon. Not. Roy. Astron. Soc.429 (2013) 1514.
- [45] Cong Zhang, Han Zhang, Shuo Yuan, Siqi Liu, Tong-Jie Zhang, and Yan-Chun Sun, Research in Astronomy and Astrophysics 14 (2014) 1221.
- [46] Daniel Stern, Raul Jimenez, Licia Verde, Marc Kamionkowski, and S. Adam Stanford, JCAP2010 (2010) 008.
- [47] M. Moresco, A. Cimatti, R. Jimenez, et al., JCAP2012 (2012) 006.
- [48] Michele Moresco, Lucia Pozzetti, Andrea Cimatti, Raul Jimenez, Claudia Maraston, Licia Verde, Daniel Thomas, Annalisa Citro, Rita Tojeiro, and David Wilkinson, JCAP2016 (2016) 014.
- [49] A. L. Ratsimbazafy, S. I. Loubser, S. M. Crawford, C. M. Cress, B. A. Bassett, R. C. Nichol, and P. Väisänen, Mon. Not. Roy. Astron. Soc.467 (2017) 3239.
- [50] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, et al., Astron. Astrophys.641 (2020) A6.
- [51] Tonghua Liu, Shuo Cao, Sixuan Zhang, Xiaolong Gong, Wuzheng Guo, and Chenfa Zheng, European Physical Journal C 81 (2021) 903.
- [52] Bin Zhang, Xiaoyao Xie, Xiaodong Nong, Guangzhen Wang, Zhiguo Xiong, Puxun Wu, and Nan Liang, arXiv e-prints.
- [53] Guo-Jian Wang, Xiao-Jiao Ma, and Jun-Qing Xia, Mon. Not. Roy. Astron. Soc. 501 (2021) 5714.
- [54] Jing-Zhao Qi, Ping Meng, Jing-Fei Zhang, and Xin Zhang, Phys. Rev. D 108 (2023) 063522.
- [55] Tonghua Liu, Shuo Cao, Shuai Ma, Yuting Liu, Chenfa Zheng, and Jieci Wang, Physics Letters B 838 (2023) 137687.
- [56] Tonghua Liu, Xiyan Yang, Zisheng Zhang, Jieci Wang, and Marek Biesiada, Physics Letters B 845 (2023) 138166.
- [57] Antony Lewis and Sarah Bridle, Phys. Rev. D 66 (2002) 103511.
- [58] J. P. Norris, G. F. Marani, and J. T. Bonnell, Astrophys. J. 534 (2000) 248.

- [59] Tingfeng Yi, Enwei Liang, Yiping Qin, and Ruijing Lu, Mon. Not. Roy. Astron. Soc. 367 (2006) 1751.
- [60] Shen-Shi Du, Lin Lan, Jun-Jie Wei, Zi-Min Zhou, He Gao, Lu-Yao Jiang, Bin-Bin Zhang, Zi-Ke Liu, Xue-Feng Wu, En-Wei Liang, and Zong-Hong Zhu, Astrophys. J. 906 (2021) 8.
- [61] Gerald J. Fishman, Publications of the Astronomical Society of the Pacific 107 (1995) 1145.
- [62] Lang Shao, Bin-Bin Zhang, Fu-Ri Wang, et al., Astrophys. J. 844 (2017) 126.