

Sign pattern matrices associated with cycle graphs that require algebraic positivity

Sunil Das *

Department of Mathematics, Faculty of Science and Technology (IcfaiTech), ICFAI Foundation for Higher Education, Hyderabad, Telangana - 501203, India

Abstract

A real matrix is said to be positive if its every entry is positive, and a real square matrix A is algebraically positive if there exists a real polynomial f such that $f(A)$ is a positive matrix. A sign pattern matrix A is said to require a property if all matrices having sign pattern as A have that property. In this paper, we characterize all sign pattern matrices associated with cycle graphs having only negative 2-cycles that require algebraic positivity. We also give some necessary conditions for a sign pattern matrix to require algebraic positivity, and some sufficient conditions for a sign pattern matrix with cycle graph to require algebraic positivity. Finally, we describe all 4-by-4 sign pattern matrices associated with cycle graphs that require algebraic positivity.

Keywords: Algebraically positive, Sign pattern, Require algebraic positivity, Sign non-singular, Signed digraph.

AMS Subject Classifications: 15B35, 15B48, 05C50.

1 Introduction

A real matrix is said to be positive if its every entry is positive. Kirkland, Qiao, and Zhan introduced the concept of algebraically positive matrices in 2016. A real square matrix A is algebraically positive if there exists a real polynomial f such that $f(A)$ is a positive matrix. The characterization of algebraically positive matrices was given by the following result.

Theorem 1.1 ([7]). *A real square matrix is algebraically positive if and only if it has a simple real eigenvalue and corresponding left and right positive eigenvectors.*

A sign pattern matrix is a matrix whose every entry is a symbol from the set $\{+, -, 0\}$. The qualitative class of a matrix A , denoted by $Q(A)$, is defined by the set of all real matrices obtained from A by replacing ‘+’ by some positive number, ‘-’ by some negative number and ‘0’ by the zero number. A sign pattern matrix A allows a property P if at least one matrix in $Q(A)$ has the property P and requires property P if all matrices in $Q(A)$ have the property P .

Let $\mathbb{T} = \{+, -, 0, +_0, -_0, \#\}$. Any matrix with entries from \mathbb{T} is said to be a \mathbb{T} -pattern matrix, and the qualitative class of a \mathbb{T} -pattern matrix A is the set of all real matrices obtained from A replacing $+, -, 0, +_0, -_0, \#$ by a positive, negative, zero, nonnegative, non-positive, and an arbitrary real number respectively. We may describe a list of sign pattern matrices through a \mathbb{T} -pattern matrix. If a \mathbb{T} -pattern matrix requires a property P , then all sign pattern matrices described by it also requires the property P .

The problem of identifying sign patterns requiring algebraic positivity is widely open. Path sign patterns requiring algebraic positivity is described in [4], and list of all 3-by-3 sign patterns requiring algebraic positivity are presented in [1]. Characterization of sign patterns allowing algebraic positivity is still not solved completely, but many results to identify those patterns can be observed in [1, 3, 4, 7]. In this paper, we give several results on sign patterns associated with cycle graphs that require algebraic positivity.

Let us recall from [2, 6] that the digraph of a sign pattern matrix A of order n , denoted by $D(A)$, is defined by a directed graph with vertex set $\{1, 2, \dots, n\}$, where (\vec{i}, \vec{j}) is an arc of $D(A)$ if and only if $a_{ij} \neq 0$. The graph of a sign pattern matrix A of order n , denoted by $G(A)$, is defined by an undirected graph with vertex set $\{1, 2, \dots, n\}$, where $[i, j]$ is an edge of $G(A)$ if and only if either $a_{ij} \neq 0$ or $a_{ji} \neq 0$.

*Email address: sunilmath92@gmail.com

A directed path α from vertex u to vertex v in $D(A)$ is a sequence $(u = i_0, i_1, \dots, i_k = v)$ of distinct vertices such that $(\overrightarrow{i_{t-1}, i_t})$ is an arc in $D(A)$ for $t = 1, 2, \dots, k$. If all vertices are not distinct, then it is called a directed walk. The length of a directed path is the number of arcs lying on the path. Here length of the directed path α is k . Sign of the path α is defined by $\text{sign}(\alpha) = \text{sign}(a_{i_0 i_1} a_{i_1 i_2} \cdots a_{i_{k-1} i_k})$. A directed cycle γ of length k or a directed k -cycle in $D(A)$ is a sequence $(i_1, i_2, \dots, i_k, i_1)$ of vertices such that i_1, i_2, \dots, i_k are distinct and $(\overrightarrow{i_1, i_2}), (\overrightarrow{i_2, i_3}), \dots, (\overrightarrow{i_k, i_1})$ are arcs in $D(A)$. Sign of the directed cycle γ is defined by $\text{sign}(\gamma) = \text{sign}(a_{i_1 i_2} a_{i_2 i_3} \cdots a_{i_k i_1})$. A square matrix A is said to be irreducible if $D(A)$ is strongly connected, that is, for any ordered pair (i, j) of vertices there is a directed path from i to j in $D(A)$. In $G(A)$, an (undirected) cycle of length k is a sequence $(i_1, i_2, \dots, i_k, i_1)$ of vertices such that i_1, i_2, \dots, i_k are distinct and $[i_1, i_2], [i_2, i_3], \dots, [i_k, i_1]$ are edges in $G(A)$.

Throughout this paper, \mathbb{R} denotes the set of all real numbers, \mathbb{R}^n is the n -dimensional Euclidean space, and A^T is the transpose of the matrix A . A vector in \mathbb{R}^n is denoted by \bar{x} and its i -th entry is denoted by x_i . A zero vector is denoted by $\bar{0}$ and an identity matrix is denoted by I , whose sizes will be clear from the context. We denote the adjugate of a square matrix A by $\text{adj}(A)$.

In Section 2, we provide some necessary conditions for a sign pattern matrix to require algebraic positivity. In Section 3, we generalize a known sufficient condition to identify sign patterns A such that $\text{adj}(B)$ has the same sign pattern for all $B \in Q(A)$. We use this result in Section 4 to provide some sufficient conditions for a sign pattern matrix to require algebraic positivity. We describe all sign pattern matrices with cycle graphs having only negative 2-cycles that require algebraic positivity, and all 4-by-4 sign pattern matrices with cycle graphs that require algebraic positivity in Section 5.

2 Necessary conditions for a sign pattern to require algebraic positivity

The following lemma from [5, Theorem 1.4.7] gives a relation between left and right eigenvectors corresponding to two distinct eigenvalues of a matrix.

Lemma 2.1 ([5]). *Let A be a real square matrix and $\bar{x}, \bar{y} \in \mathbb{R}^n \setminus \{\bar{0}\}$ such that $A\bar{x} = \lambda\bar{x}$ and $\bar{y}^T A = \mu\bar{y}^T$ for some $\lambda, \mu \in \mathbb{R}$. If $\lambda \neq \mu$, then $\bar{y}^T \bar{x} = 0$.*

The following lemma shows that any non-negative left or right eigenvector of an algebraically positive matrix must be a scalar multiple of its left or right positive eigenvector, whose existence is given by Theorem 1.1.

Lemma 2.2. *If a matrix A has a non-negative left or right eigenvector \bar{x} with some $x_i = 0$, then A is not algebraically positive.*

Proof. Suppose that $A\bar{x} = \lambda\bar{x}$, where \bar{x} is non-negative right eigenvector of A with some $x_i = 0$. If A is algebraically positive, then by Theorem 1.1, it has a simple eigenvalue μ such that $A\bar{z} = \mu\bar{z}$ and $\bar{y}^T A = \mu\bar{y}^T$ for some positive vectors \bar{y} and \bar{z} . Since \bar{x} and \bar{z} are independent, and μ is simple eigenvalue, $\lambda \neq \mu$. But $\bar{y}^T \bar{x} \neq 0$, a contradiction to Lemma 2.1. Therefore A is not algebraically positive. Similarly, if A has a non-negative left eigenvector \bar{x} with some $x_i = 0$, then A is not algebraically positive. \square

Let $A(i, :)$ be the matrix obtained from A by deleting i -th row, and $A(:, j)$ be the matrix obtained from A by deleting j -th column. The following lemma gives a sufficient condition to identify sign patterns that do not require algebraic positivity.

Lemma 2.3. *Let $B \in Q(A)$. If there exist $t, \lambda \in \mathbb{R}$ such that each column of the matrix $(B - \lambda I)(t, :)$ is either a zero vector or has both positive and negative entries, then A does not require algebraic positivity.*

Proof. Suppose that each column of the matrix $(B - \lambda I)(t, :)$ is either a zero vector or has both positive and negative entries. Keeping λ fixed, we can choose the entries of B in such a way that each column sum of $(B - \lambda I)(t, :)$ is zero so that $\bar{x}^T B = \lambda\bar{x}^T$, where

$$x_i = \begin{cases} 1, & \text{if } i \neq t; \\ 0, & \text{if } i = t. \end{cases}$$

So by Lemma 2.2, B is not algebraically positive, and thus A does not require algebraic positivity. \square

If $B \in Q(A)$ and there exist $t, \lambda \in \mathbb{R}$ such that each row of the matrix $(B - \lambda I)(:, t)$ is either a zero vector or has both positive and negative entries, then we can similarly show that A does not require algebraic positivity. Consequently, we have the following result. It will be used numerous in later part of this paper.

Corollary 2.4. *Let A be a sign pattern matrix. If there exists t such that the t -th column of $A(t, :)$ is either a zero vector or has both $+$ and $-$, and all other columns has a $+$, then A does not require algebraic positivity.*

Let A be a sign pattern matrix, and let the matrices A_+ , A_- and B_A be defined as follows ([1]):

$$(A_+)_{ij} = \begin{cases} +, & \text{if } a_{ij} = +; \\ 0, & \text{if } a_{ij} \neq +. \end{cases} \quad (A_-)_{ij} = \begin{cases} -, & \text{if } a_{ij} = -; \\ 0, & \text{if } a_{ij} \neq -. \end{cases} \quad B_A = A_+ - A_-^T$$

Recall from [3] that a sign pattern matrix A is AP-irreducible if every row and column of A contains a $+$, and both A and B_A are irreducible. If a sign pattern matrix requires some property, then it also allows that property. So we have the following result from [1, 7].

Lemma 2.5. *If a sign pattern matrix requires algebraic positivity, then it is AP-irreducible.*

We have the following results from [7].

Lemma 2.6 ([7]). *The following statements are equivalent for a square matrix A .*

1. A is algebraically positive.
2. $-A$ is algebraically positive.
3. $P^T A P$ is algebraically positive for every permutation matrix P .
4. $A - \alpha I$ is algebraically positive for every real number α .

We shall focus on the sign patterns whose graphs are cycles. If $G(A)$ is a cycle for some sign pattern matrix A of order n , we can assume that the cycle is given by the sequence $(1, 2, 3, \dots, n, 1)$ because of Lemma 2.6.3. So we consider A to be of the form

$$A = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & \cdots & 0 & a_{1n} \\ a_{21} & a_{22} & a_{23} & 0 & \cdots & \cdots & 0 \\ 0 & \ddots & \ddots & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & a_{n-2,n-1} & a_{n-1,n-1} & a_{n-1,n} \\ a_{n1} & 0 & \cdots & \cdots & 0 & a_{n,n-1} & a_{nn} \end{bmatrix}, \quad (1)$$

where $a_{ij} \in \{+, -, 0\}$ for all i, j , and either $a_{ij} \neq 0$ or $a_{ji} \neq 0$ for $j = i + 1 \pmod n$.

Lemma 2.7. *Let A be a sign pattern matrix of the form (1). If A requires algebraic positivity, then we have the following:*

1. *There do not exist i, j such that $a_{i,i+1} = 0$ and $a_{j+1,j} = 0$.*
2. *There do not exist i, j such that $a_{i,i+1} = -0$, $a_{i+1,i} = +0$ and $a_{j+1,j} = -0$, $a_{j,j+1} = +0$.*

Proof. Since A requires algebraic positivity, by Lemma 2.5, A is AP-irreducible. Let us say that $(\overrightarrow{i, j})$ is a forward arc if $j = i + 1 \pmod n$, and $(\overleftarrow{j, i})$ is a backward arc if $j = i + 1 \pmod n$.

1. Suppose that there exist i, j such that $a_{i, i+1} = 0$ and $a_{j+1, j} = 0$. Since A is of the form (1), $i \neq j$. Now any directed path in $D(A)$ to vertex j can be traversed via forward arcs only, and any directed path in $D(A)$ from vertex i can be traversed via backward arcs only. So there is no directed path from i to j in $D(A)$, and thus A is reducible. So A is not AP-irreducible, a contradiction.
2. Suppose that there exist i, j such that $a_{i, i+1} = -0$, $a_{i+1, i} = +0$ and $a_{j+1, j} = -0$, $a_{j, j+1} = +0$. So $(B_A)_{i, i+1} = 0$ and $(B_A)_{j+1, j} = 0$. Now any directed path in $D(B_A)$ to vertex j can be traversed via forward arcs only, and any directed path in $D(B_A)$ from vertex i can be traversed via backward arcs only. So there is no directed path from i to j in $D(B_A)$, and thus B_A is reducible. So A is not AP-irreducible, a contradiction.

□

The following result shows that for a sign pattern matrix A with cycle graph to require algebraic positivity, we must have a directed n -cycle in $D(A)$ either with positive arcs only or with negative arcs only.

Lemma 2.8. *Let A be a sign pattern matrix of order n of the form (1). If A requires algebraic positivity, then either $D(A_+)$ or $D(A_-)$ contains a directed cycle of length n .*

Proof. Since A requires algebraic positivity, due to Lemma 2.5, we can assume that every row and column of A contains a $+$.

Suppose that neither $D(A_+)$ nor $D(A_-)$ contains a directed cycle of length n . We first prove the following three statements successively.

1. $a_{ij} = a_{ji}$ for all i, j .
2. There exist no three distinct vertices i, j, k such that $a_{ij}a_{jk} = +$.
3. $a_{ii} = 0$ for all i .

Our goal is to arrive at a contradiction with the help of these three statements. Since A is of the form (1), and neither $D(A_+)$ nor $D(A_-)$ contains a directed cycle of length n , we can assume that there exist $i, j \in \{1, 2, \dots, n-1\}$ such that $a_{i, i+1} = +$ and $a_{j, j+1} = -$.

1. Suppose that $a_{i+1, i} = -0$. Because of Lemma 2.6.3, we can consider that $a_{12} = +$, and $a_{k+1, k} = -0$ for some $k < t = \min\{i : a_{i, i+1} = -0\}$. Then $a_{k, k+1} = +$, and thus by Lemma 2.7.2, $a_{t+1, t} = -$.

Since every column of A contains a $+$, and $a_{k+1, k} = -0$ for some $k < t = \min\{i : a_{i, i+1} = -0\}$, the t -th column of $A(t, :)$ has both $+$ and $-$, and all other columns of $A(t, :)$ has a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction. Thus $a_{i+1, i} = +$.

Therefore by Lemma 2.6, we can conclude that $a_{ij} = +$ implies $a_{ji} = +$, and $a_{ij} = -$ implies $a_{ji} = -$.

2. Since A has the form (1), $a_{ij} \neq 0$ whenever $|i - j| = 1$. Suppose that there exist three distinct vertices i, j, k such that $a_{ij}a_{jk} = +$. Because of Lemma 2.6, we can assume that $a_{12}, a_{23} = +$ and $a_{34} = -$. So $a_{43} = -$.

Since every column of A contains a $+$, the third column of $A(3, :)$ has a $+$ and a $-$, and all other columns has a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction. So there exist no three distinct vertices i, j, k such that $a_{ij}a_{jk} = +$. Consequently, n is even.

3. We can assume that $a_{12} = +$ and $a_{23} = -$. Then $a_{21} = +$ and $a_{32} = -$.
 - (a) If $a_{11} = +$, then the second column of $A(2, :)$ has both $+$ and $-$, and all other columns has a $+$, because every column of A contains a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction.
 - (b) If $a_{22} = +$, then the first column of $A(1, :)$ has both $+$ and $-$, and all other columns has a $+$, because every column of A contains a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction.

Thus $a_{ii} \neq +$ for all i . Again by Lemma 2.6, we have $a_{ii} \neq -$ for all i . Therefore $a_{ii} = 0$ for all i .

Using the above three statements, we can assume that n is even, $a_{1n} = a_{n1} = -$, $a_{i,i+1} = a_{i+1,i} = +$ for all odd i , and $a_{i,i+1} = a_{i+1,i} = -$ for all even i . Consider $B \in Q(A)$ by replacing $+$ with 1 and $-$ with -1 . Then $B\bar{x} = 0$, where

$$x_i = \begin{cases} 1, & \text{if } i \text{ is odd;} \\ 0, & \text{if } i \text{ is even.} \end{cases}$$

So by Lemma 2.2, B is not algebraically positive. Therefore A does not require algebraic positivity, a contradiction. Hence either $D(A_+)$ or $D(A_-)$ contains a directed cycle of length n . \square

The following two results successively shows that for a sign pattern matrix A to require algebraic positivity, no vertex is covered by both positive and negative directed 2-cycles in $D(A)$, and $a_{ii} = -0$ if vertex i is covered by a negative directed 2-cycle in $D(A)$.

Lemma 2.9. *Let A be a sign pattern matrix of order n such that it has the form (1). If $D(A_+)$ contains a directed cycle of length n , and A requires algebraic positivity, then no vertex is covered by both positive and negative directed 2-cycles in $D(A)$.*

Proof. Let us assume that A requires algebraic positivity. Since $D(A_+)$ contains a directed cycle of length n , assume that $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$. Suppose that we have some vertex covered by both positive and negative directed 2-cycles. Then we can assume that either $a_{1n} = +$ and $a_{21} = -$ or $a_{21} = -$ and $a_{32} = +$, because of Lemma 2.6.

1. Assume that $a_{1n} = +$ and $a_{21} = -$. Now the second row of $A(:, 2)$ has both $+$ and $-$, and all other rows have a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction.
2. Assume that $a_{21} = -$ and $a_{32} = +$. Now the first column of $A(1, :)$ has both $+$ and $-$, and all other columns have a $+$. So by Corollary 2.4, A does not require algebraic positivity, a contradiction.

\square

Lemma 2.10. *Let A be a sign pattern matrix of order n having the form (1), and $D(A_+)$ contains a directed cycle of length n . If $a_{ij}a_{ji} = -$ for some $i \neq j$ with $a_{ii} = +$, then A does not require algebraic positivity.*

Proof. Since $D(A_+)$ contains a directed cycle of length n , assume that $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$. Due to Lemma 2.6, we can consider that $a_{21} = -$ and $a_{22} = +$. Now the first column of $A(1, :)$ has both $+$ and $-$, and all other columns has a $+$. So by Corollary 2.4, A does not require algebraic positivity. \square

A square sign pattern matrix A is said to be sign non-singular (SNS) if every matrix in $Q(A)$ is non-singular. For a square matrix A of order n , the standard determinant expansion of A is given by

$$\det A = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}, \quad (2)$$

where the sum is taken over the set S_n of all permutations of the set $\{1, 2, \dots, n\}$, and $\text{sgn}(\sigma)$ is the sign of the permutation σ .

Lemma 2.11 ([2]). *Let $A = [a_{ij}]$ be a matrix of order n . Then the following are equivalent:*

1. A is an SNS matrix.
2. There is a nonzero term in the standard determinant expansion of A and every nonzero term has the same sign.

The following lemma gives a necessary condition for a sign pattern matrix associated with cycle graphs to require algebraic positivity if no diagonal entries are positive and no 2-cycles are positive.

Lemma 2.12. *Let A be a sign pattern matrix of order n of the form (1) such that $a_{ii} = -0$ for all i and all directed 2-cycles in $D(A)$ are negative. Let $D(A_+)$ contains a directed n -cycle, $D(A_-)$ does not contain a directed n -cycle and $a_{ij}a_{ji} = -$ for some $i \neq j$. If A is not SNS and A requires algebraic positivity, then $a_{ii} = -$ and $a_{jj} = -$ whenever $a_{ij} = +$ and $a_{ji} = 0$ for some $i \neq j$.*

Proof. Since $D(A_+)$ contains a directed n -cycle, let $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$. Since $D(A_-)$ does not contain a directed n -cycle, assume that $a_{1n} = 0$.

Now $D(A)$ has no directed k -cycle for $2 < k < n$ and only one directed n -cycle with sign $+$. The standard determinant expansion of every matrix in $Q(A)$ has a term with sign $(-)^{n-1}$ corresponding to the n -cycle. Since A is not SNS, the standard determinant expansion of every matrix in $Q(A)$ has a term with sign $(-)^n$. So we can choose $B \in Q(A)$ such that $\det B = (-1)^n$. Moreover, all 2-cycles in $D(A)$ are negative. So the coefficients of x^i in the characteristic polynomial of B are non-negative for $i = 1, 2, \dots, n$. Consequently, the characteristic equation $p(B) = 0$ has no positive real root, by Descartes' rule of signs. Thus all eigenvalues of B are negative, as $\det B \neq 0$.

If $a_{nn} = 0$, then for any $\lambda < 0$, the n -th column of $B - \lambda I$ is a nonzero, non-negative vector so that $B - \lambda I$ does not have a positive left eigenvector. Similarly, if $a_{11} = 0$, then for any $\lambda < 0$, the first row of $B - \lambda I$ is a nonzero, non-negative vector so that $B - \lambda I$ does not have a positive right eigenvector. Thus if either $a_{11} = 0$ or $a_{nn} = 0$, then B is not algebraically positive so that A does not require algebraic positivity, a contradiction. Therefore $a_{11} = -$ and $a_{nn} = -$. Hence the result follows. \square

Lemma 2.13. *A real skew-symmetric matrix of even order cannot be algebraically positive.*

Proof. Eigenvalues of a real skew-symmetric matrix are either 0 or purely imaginary. So an even order real skew-symmetric matrix cannot have a simple real eigenvalue. Therefore, by Theorem 1.1, a real skew-symmetric matrix of even order cannot be algebraically positive. \square

The following result is a necessary condition for a sign pattern of the form (1) to require algebraic positivity when there is no positive directed 2-cycle in its digraph.

Theorem 2.14. *Let A be a sign pattern matrix of order n of the form (1) and all directed 2-cycles in $D(A)$ are negative. Let $D(A_+)$ contains a directed n -cycle, $a_{ij}a_{ji} = -$ for some $i \neq j$, and A requires algebraic positivity.*

1. *If $D(A_-)$ does not contain an n -cycle and $a_{ii} = -_0$ for all i , then A is SNS.*
2. *If $D(A_-)$ contains an n -cycle, then n is odd and $a_{ii} = 0$ for all i .*

Proof. Assume that A requires algebraic positivity. Since $D(A_+)$ contains a directed n -cycle, let $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$.

1. Suppose that A is not SNS. Since $a_{ij}a_{ji} = -$ for some $i \neq j$ and $D(A_-)$ does not contain a directed n -cycle, assume that $a_{21} = -$ and $a_{1n} = 0$. Further, by Lemma 2.12, $a_{ii} = -$ and $a_{jj} = -$ provided that $a_{ji} = 0$ for some $j = i + 1 \pmod n$. So for every k -th column of A , either $a_{kk} = -$ or $a_{k+1,k} = -$ for $1 \leq k \leq n - 1$, and $a_{nn} = -$.

Now we can choose $B \in Q(A)$ and $\lambda < 0$ such that the second column of $(B - \lambda I)(1, :)$ is zero vector or has both $+$ and $-$, and all other columns of $(B - \lambda I)(t, :)$ has both $+$ and $-$. So by Lemma 2.3, A does not require algebraic positivity, a contradiction. Therefore A is SNS.

2. Suppose that $D(A_-)$ contains an n -cycle. Then $a_{ij}a_{ji} = -$ for all i, j with $j = i + 1 \pmod n$. So by Lemma 2.6 and 2.10, $a_{ii} = 0$ for all i . If n is even, we can choose $B \in Q(A)$ by replacing $+$ with 1 and $-$ by -1 so that B is skew-symmetric. Then by Lemma 2.13, B is not algebraically positive, a contradiction to the assumption that A requires algebraic positivity. Therefore n is odd. \square

3 Identifying sign patterns to require a fixed sign at each adjugate entry

Given an SNS matrix A with negative main diagonal, Theorem 3.2.5 in [2] provides a necessary and sufficient condition for a non-diagonal entry of B^{-1} to have the same sign for all $B \in Q(A)$. We generalize this result as follows.

Theorem 3.1. *Let $A = [a_{ij}]$ be an SNS matrix of order n , and let r and s be distinct integers with $1 \leq r, s \leq n$. The entries in the (s, r) -position of matrices in $\{\text{adj}(B) : B \in Q(A)\}$ are of the same sign if and only if for every directed path α from s to r in the signed digraph $D(A)$, the sign $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta)$ is the same for all nonzero term β in the standard determinant expansion of the matrix $A(\alpha)$ obtained from A by deleting rows and columns corresponding to the vertices covered by α , where l is the length of α .*

Proof. Given a matrix $B \in Q(A)$, the entry in the (s, r) -position of $\text{adj}(B)$ is $(-1)^{r+s}$ times the determinant of the matrix obtained from B by deleting the r -th row and the s -th column. Let \bar{e}_r be the $n \times 1$ column vector having only one nonzero entry 1 at position r . So the entry in the (s, r) -position of $\text{adj}(B)$ is $\det B(s \leftarrow \bar{e}_r)$, where the matrix $\tilde{B} = B(s \leftarrow \bar{e}_r)$ obtained from B by replacing s -th column of B with \bar{e}_r . Then $\tilde{b}_{rs} = 1$. Every nonzero term in $\det B(s \leftarrow \bar{e}_r)$ can be expressed as

$$(-1)^l (\tilde{b}_{rs} b_{si_1} b_{i_1 i_2} \cdots b_{i_k r}) \beta = (-1)^l (b_{si_1} b_{i_1 i_2} \cdots b_{i_k r}) \beta,$$

where $\alpha = (s, i_1, i_2, \dots, i_k, r)$ is a directed path in $D(A)$, l is the length of α , and β is a nonzero term in the determinant expansion of $B(\alpha)$ obtained from B by deleting rows and columns corresponding to the vertices covered by α . Since A is SNS, $A(\alpha)$ is SNS provided that $\text{sign}(\alpha) \neq 0$. Therefore the result follows from Lemma 2.11. \square

4 Sufficient conditions for a sign pattern to require algebraic positivity

The following three lemmas from [4, 7] show some sufficient conditions for any sign pattern matrix to require algebraic positivity.

Lemma 4.1 ([7]). *If A is a real matrix and there is a positive integer k such that A^k is algebraically positive, then A is algebraically positive.*

Lemma 4.2 ([4]). *If $\text{adj}(A)$ is algebraically positive for some real square matrix A , then A is algebraically positive.*

Lemma 4.3 ([4]). *If all nonzero off-diagonal entries of an irreducible sign pattern matrix are the same sign (+ or -), then that sign pattern matrix requires algebraic positivity.*

The following three results give sufficient conditions for a sign pattern matrix of the form (1) to require algebraic positivity if there are no positive directed 2-cycles.

Theorem 4.4. *Let A be a sign pattern matrix of odd order n of the form (1). If $D(A_+)$ contains a directed n -cycle, $D(A_-)$ contains a directed n -cycle and $a_{ii} = 0$ for all i , then A requires algebraic positivity.*

Proof. Since $D(A_+)$ contains a directed n -cycle and $D(A_-)$ contains a directed n -cycle, assume that $a_{ij} = +$ and $a_{ji} = -$ for $j = i + 1 \pmod n$. Since $a_{ii} = 0$ for all i , there is a directed walk of length 2 in $D(A)$ from vertex i to vertex j if and only if $j = i + 2 \pmod n$ or $j = i - 2 \pmod n$. Each directed walk of length 2 consists of either two positive arcs or two negative arcs, and thus it has sign $+$. Therefore

$$(A^2)_{ij} = \begin{cases} +, & \text{if } j = i + 2 \pmod n \text{ or } j = i - 2 \pmod n; \\ 0, & \text{otherwise.} \end{cases}$$

So A^2 is irreducible and all nonzero off-diagonal entries are $+$. Therefore A^2 requires algebraic positivity, by Lemma 4.3, and hence A requires algebraic positivity, by Lemma 4.1. \square

Theorem 4.5. *Let A be a sign pattern matrix of order n of the form (1), all directed 2-cycles in $D(A)$ are negative and $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$. If A is an SNS matrix such that $a_{ii} = -0$ for all i , then A requires algebraic positivity.*

Proof. Let $B \in Q(A)$. Since all directed 2-cycles in $D(A)$ are negative and $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$, every path in $D(A)$ consists of either all positive edges or all negative edges.

Given two distinct indices r and s , we wish to determine the sign of the (s, r) -entry of $\text{adj}(B)$, that is, $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta)$, where $\alpha = (s, i_1, i_2, \dots, i_k, r)$ is a directed path from s to r in $D(A)$, l is the length of α , and β is a nonzero term in the determinant expansion of $A(\alpha)$ obtained from A by deleting rows and columns corresponding to the vertices covered by α .

1. Let n be odd. Since A is SNS, and $D(A)$ contains the cycle $\gamma = (1, 2, 3, \dots, n, 1)$ with $\text{sign} \gamma = +$, by Lemma 2.11, every nonzero term in the determinant expansion of B has sign $(-)^{n-1} \cdot + = +$.

- (a) Assume that the directed path α consists of all positive edges.

- i. If l is even, total number of vertices covered in α is odd. So $B(\alpha)$ is of even order, and thus $\text{sign}(\beta) = +_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = + \cdot + \cdot +_0 = +_0$. In particular, if $l = n - 1$, then α covers all vertices. So the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) = + \cdot + = +$. Consequently, the (s, r) -entry of $\text{adj}(B)$ has a contributing term with sign $+$ for all $(s, r) \in \{(2, 1), (3, 2), \dots, (n, n-1), (1, n)\}$.
 - ii. If l is odd, total number of vertices covered in α is even. So $B(\alpha)$ is of odd order, and thus $\text{sign}(\beta) = -_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = - \cdot + \cdot -_0 = +_0$.
- (b) Assume that the directed path α consists of all negative edges.
- i. If l is even, total number of vertices covered in α is odd. So $B(\alpha)$ is of even order, and thus $\text{sign}(\beta) = +_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = + \cdot + \cdot +_0 = +_0$.
 - ii. If l is odd, total number of vertices covered in α is even. So $\text{sign}(\alpha) = +$. Let $B[\alpha]$ be the submatrix of B having rows and columns corresponding to the indices in α . Since α consists of only negative arcs and covers even number of vertices, $B[\alpha]$ has a nonzero term δ in its determinant expansion with sign $+$. Now $B(\alpha)$ is of odd order. If β is a nonzero term in the determinant expansion of $B(\alpha)$, then $\text{sign}(\beta) = -$ so that $\det B$ has a term with sign $\text{sign}(\delta) \cdot \text{sign}(\beta) = -$, a contradiction. So $\text{sign}(\beta) = 0$. Thus the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = - \cdot + \cdot 0 = 0$.

Thus all nonzero off-diagonal entries of $\text{adj}(B)$ have the sign $+$. Further, $\text{adj}(B)$ is irreducible because the (s, r) -entry of $\text{adj}(B)$ has the sign $+$ for all $(s, r) \in \{(2, 1), (3, 2), \dots, (n, n-1), (1, n)\}$. Therefore, by Lemma 4.2, B is algebraically positive. Hence A requires algebraic positivity.

2. Let n be even. Since A is SNS, and $D(A)$ contains the cycle $\gamma = (1, 2, 3, \dots, n, 1)$ with $\text{sign } \gamma = +$, by Lemma 2.11, every nonzero term in the determinant expansion of B has sign $(-)^{n-1} \cdot + = -$.

- (a) Assume that the directed path α consists of all positive edges.
- i. If l is even, total number of vertices covered in α is odd. So $B(\alpha)$ is of odd order, and thus $\text{sign}(\beta) = -_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = + \cdot + \cdot -_0 = -_0$.
 - ii. If l is odd, total number of vertices covered in α is even. So $B(\alpha)$ is of even order, and thus $\text{sign}(\beta) = +_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = - \cdot + \cdot +_0 = -_0$. In particular, if $l = n - 1$, then α covers all vertices. So the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) = - \cdot + = -$. Consequently, the (s, r) -entry of $\text{adj}(B)$ has a contributing term with sign $-$ for all $(s, r) \in \{(2, 1), (3, 2), \dots, (n, n-1), (1, n)\}$.
- (b) Assume that the directed path α consists of all negative edges.
- i. If l is even, total number of vertices covered in α is odd. So $B(\alpha)$ is of odd order, and thus $\text{sign}(\beta) = -_0$. Moreover, $\text{sign}(\alpha) = +$. In this case, the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = + \cdot + \cdot -_0 = -_0$.
 - ii. If l is odd, total number of vertices covered in α is even. So $\text{sign}(\alpha) = +$. Let $B[\alpha]$ be the submatrix of B having rows and columns corresponding to the indices in α . Since α consists of only negative arcs and covers even number of vertices, $B[\alpha]$ has a nonzero term δ in its determinant expansion with sign $+$. Now $B(\alpha)$ is of even order. If β is a nonzero term in the determinant expansion of $B(\alpha)$, then $\text{sign}(\beta) = +$ so that $\det B$ has a term with sign $\text{sign}(\delta) \cdot \text{sign}(\beta) = +$, a contradiction. So $\text{sign}(\beta) = 0$. Thus the sign of the contributing term to the (s, r) -entry of $\text{adj}(B)$ is $(-)^l \text{sign}(\alpha) \cdot \text{sign}(\beta) = - \cdot + \cdot 0 = 0$.

Thus all nonzero off-diagonal entries of $\text{adj}(B)$ have the sign $-$. Further, $\text{adj}(B)$ is irreducible because the (s, r) -entry of $\text{adj}(B)$ has the sign $-$ for all $(s, r) \in \{(2, 1), (3, 2), \dots, (n, n-1), (1, n)\}$. Therefore, by Lemma 4.2, B is algebraically positive. Hence A requires algebraic positivity. \square

Corollary 4.6. *Let A be a sign pattern matrix of order n of the form (1), all directed 2-cycles in $D(A)$ are negative and $a_{ij} = +$ for all i, j with $j = i + 1 \pmod n$. If $a_{ii} = -_0$ for all i covered by negative directed 2-cycles in $D(A)$ and $a_{ii} = +$ for some i , then A requires algebraic positivity.*

Proof. Let $B \in Q(A)$, and let $a_{ii} = +$ for some vertex i covered by no negative directed 2-cycles in $D(A)$. If $k = \arg \max_i b_{ii}$, then k is covered by no negative directed 2-cycles in $D(A)$. Consider the pattern of $B - b_{kk}I$, which is SNS and all its diagonal entries are from $\{-, 0\}$. Then by Theorem 4.5, $B - b_{kk}I$ is algebraically positive. So by Lemma 2.6, B is algebraically positive, and hence A requires algebraic positivity. \square

If a sign pattern matrix of the form (1) has both positive and negative directed 2-cycles, then a sufficient condition for it to require algebraic positivity is given by the following result.

Theorem 4.7. *Let A be a sign pattern matrix of order n of the form (1) such that $D(A_+)$ contains a directed n -cycle, and no vertex is covered by both positive and negative directed 2-cycles. If $a_{ii} = 0$ for all vertices i covered by negative directed 2-cycles, and $a_{ii} = +$ for all vertices i covered by positive directed 2-cycles, then A requires algebraic positivity.*

Proof. Since $D(A_+)$ contains a directed n -cycle, assume that $a_{ij} = +$ for $j = i + 1 \pmod n$. Since no vertex is covered by both positive and negative directed 2-cycles, no directed walk of length 2 between two distinct vertices consists of both positive and negative arcs. Each directed walk of length 2 between two distinct vertices consists of either two positive arcs, or two negative arcs, or a positive loop and a positive arc. There is a directed walk of length 2 in $D(A)$ from vertex i to vertex j with sign $+$ if $j = i + 2 \pmod n$. Therefore

$$(A^2)_{ij} = \begin{cases} +, & \text{if } j = i + 2 \pmod n; \\ +0, & \text{if } i \neq j \text{ and } j \neq i + 2 \pmod n. \end{cases}$$

So A^2 is irreducible and all nonzero off-diagonal entries are $+$. Therefore A^2 requires algebraic positivity, by Lemma 4.3, and hence A requires algebraic positivity, by Lemma 4.1. \square

5 Main results

We characterize all sign pattern matrices A that require algebraic positivity, where $G(A)$ is a cycle and all directed 2-cycles in $D(A)$ are negative, through the following result.

Theorem 5.1. *Let A be an irreducible sign pattern matrix of order n such that $G(A)$ is a cycle. If all directed 2-cycles in $D(A)$ are negative, then A requires algebraic positivity if and only if either A or $-A$ satisfies one of the following conditions:*

1. All nonzero off-diagonal entries have the sign $+$.
2. n is odd, $D(A_+)$ contains a directed n -cycle and $D(A_-)$ contains a directed n -cycle such that $a_{ii} = 0$ for all i .
3. $D(A_+)$ contains a directed n -cycle and $D(A_-)$ does not contain a directed n -cycle such that A is SNS and $a_{ii} = -$ for all i .
4. $D(A_+)$ contains a directed n -cycle and $D(A_-)$ does not contain a directed n -cycle such that $a_{ii} = +$ for some i and no vertex k with $a_{kk} = +$ is covered by any directed negative 2-cycles in $D(A)$.

Proof. Suppose that A requires algebraic positivity. Since $G(A)$ is a cycle, we can assume that A has the form (1). Then by Lemma 2.8, either $D(A_+)$ or $D(A_-)$ contains a directed n -cycle. Using Lemma 2.6, we can assume that $D(A_+)$ contains a directed n -cycle.

Suppose that A has $-$ as a off-diagonal entry. Since $a_{ij}a_{ji} = -$ for all $i \neq j$, by Lemma 2.10, no vertex k with $a_{kk} = +$ is covered by any directed negative 2-cycles in $D(A)$. Further, by Theorem 2.14, if $D(A_-)$ contains a directed n -cycle, then n is odd and $a_{ii} = 0$ for all i , and if $D(A_-)$ does not contain a directed n -cycle and $a_{ii} = -$ for all i , then A is SNS. Therefore conditions 1, 2, 3, 4 are necessary for A to require algebraic positivity.

Lemma 4.3, Theorem 4.4, Theorem 4.5, Corollary 4.6 respectively show that conditions 1, 2, 3, 4 are sufficient for A to require algebraic positivity. \square

Since we have considered that $G(A)$ is a cycle, there are no directed k -cycles in $D(A)$ for $2 < k < n$. Lemma 2.8 shows that a sign pattern matrix of the form (1) must have a directed n -cycle with positive arcs only for A to require algebraic positivity. If all directed 2-cycles are negative, then the characterization of sign pattern requiring algebraic positivity is described by Theorem 5.1. The proof is done by using sign patterns of adjugates and the patterns of A^2 . But we may have both positive and negative directed

2-cycles. In that case, adjugate patterns have both $+$ and $-$ on off-diagonal. Further, for a fixed k , A^k has both $+$ and $-$ on off-diagonal. So we have to find suitable polynomial for matrices $B \in Q(A)$ to show that A requires algebraic positivity, and it gets very difficult when order of A is high. We conclude the paper describing all 4-by-4 sign pattern matrices A requiring algebraic positivity such that $G(A)$ is cycle and $D(A)$ has both positive and negative directed 2-cycles.

Lemma 5.2. *All sign pattern matrices described by the \mathbb{T} -patterns*

$$P_1 = \begin{bmatrix} -_0 & + & 0 & 0 \\ - & -_0 & + & 0 \\ 0 & 0 & +_0 & + \\ + & 0 & + & +_0 \end{bmatrix}, \quad P_2 = \begin{bmatrix} -_0 & + & 0 & 0 \\ - & -_0 & + & 0 \\ 0 & 0 & 0 & + \\ + & 0 & + & - \end{bmatrix}, \quad P_3 = \begin{bmatrix} -_0 & + & 0 & 0 \\ - & -_0 & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & 0 \end{bmatrix},$$

$$P_4 = \begin{bmatrix} -_0 & + & 0 & 0 \\ - & -_0 & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & + \end{bmatrix}, \quad P_5 = \begin{bmatrix} -_0 & + & 0 & 0 \\ - & -_0 & + & 0 \\ 0 & 0 & + & + \\ + & 0 & + & - \end{bmatrix}$$

require algebraic positivity.

Proof. Every matrix in qualitative classes of P_1, P_2, P_3 has the form

$$B = \begin{bmatrix} -b_{11} & b_{12} & 0 & 0 \\ -b_{21} & -b_{22} & b_{23} & 0 \\ 0 & 0 & b_{33} & b_{34} \\ b_{41} & 0 & b_{43} & b_{44} \end{bmatrix},$$

where $b_{11}, b_{22} \geq 0$ and $b_{12}, b_{21}, b_{23}, b_{34}, b_{41}, b_{43} > 0$. The off-diagonal entries of the matrix $M = \alpha B + \beta B^2 + B^3$ can be expressed as follows:

$$\begin{aligned} m_{14} &= b_{12}b_{23}b_{34} \\ m_{32} &= b_{34}b_{41}b_{12} \\ m_{13} &= b_{12}b_{23}(\beta - b_{11} - b_{22} + b_{33}) \\ m_{24} &= b_{23}b_{34}(\beta - b_{22} + b_{33} + b_{44}) \\ m_{31} &= b_{34}b_{41}(\beta + b_{33} + b_{44} - b_{11}) \\ m_{42} &= b_{41}b_{12}(\beta + b_{44} - b_{11} - b_{22}) \\ m_{34} &= b_{34}[\alpha + \beta(b_{33} + b_{44}) + b_{33}^2 + b_{44}^2 + b_{33}b_{44} + b_{34}b_{43}] \\ m_{43} &= b_{43}[\alpha + \beta(b_{33} + b_{44}) + b_{33}^2 + b_{44}^2 + b_{34}b_{43} + b_{33}b_{44}] + b_{41}b_{12}b_{23} \\ m_{12} &= b_{12}[\alpha - \beta(b_{11} + b_{22}) + b_{11}^2 + b_{22}^2 + b_{11}b_{22} - b_{12}b_{21}] \\ m_{23} &= b_{23}[\alpha - \beta(b_{22} - b_{33}) + b_{22}^2 + b_{33}^2 - b_{22}b_{33} - b_{12}b_{21}] \\ m_{41} &= b_{41}[\alpha - \beta(b_{11} - b_{44}) + b_{11}^2 + b_{44}^2 - b_{11}b_{44} - b_{12}b_{21} + b_{43}b_{34}] \\ m_{21} &= -b_{21}[\alpha - \beta(b_{11} + b_{22}) + b_{11}^2 + b_{22}^2 + b_{11}b_{22} - b_{12}b_{21}] + b_{23}b_{34}b_{41} \end{aligned}$$

Let $B \in Q(P_1)$. Then $b_{33}, b_{44} \geq 0$. Choose $\alpha = b_{12}b_{21} + b_{11}b_{22} + \frac{b_{23}b_{34}b_{41}}{b_{21}}$ and $\beta = b_{11} + b_{22}$. Then all off-diagonal entries of $M = \alpha B + \beta B^2 + B^3$ are non-negative, and $m_{14}, m_{41}, m_{34}, m_{43}, m_{23}, m_{32} > 0$. So M is irreducible. Thus by Lemma 4.3, B is algebraically positive, and hence P_1 requires algebraic positivity.

Let $B \in Q(P_2)$. Then $b_{33} = 0$ and $b_{44} < 0$. Choose $\alpha = b_{12}b_{21} + b_{11}b_{22} - b_{11}b_{44} - b_{22}b_{44} + \frac{b_{23}b_{34}b_{41}}{b_{21}}$ and $\beta = b_{11} + b_{22} - b_{44}$. Then all off-diagonal entries of $M = \alpha B + \beta B^2 + B^3$ are non-negative, and $m_{14}, m_{41}, m_{34}, m_{43}, m_{23}, m_{32} > 0$. So M is irreducible. Thus by Lemma 4.3, B is algebraically positive, and hence P_2 requires algebraic positivity. Similarly, P_3 requires algebraic positivity.

For every $B \in Q(P_4)$, there exists $\lambda > 0$ such that $B - \lambda I \in Q(P_3)$. Since P_3 requires algebraic positivity, so is P_4 . Similarly, P_5 requires algebraic positivity. \square

Lemma 5.3. *No sign pattern matrix among*

$$R_1 = \begin{bmatrix} - & + & 0 & 0 \\ - & - & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & - \end{bmatrix}, \quad R_2 = \begin{bmatrix} 0 & + & 0 & 0 \\ - & - & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & - \end{bmatrix}, \quad R_3 = \begin{bmatrix} - & + & 0 & 0 \\ - & 0 & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & - \end{bmatrix}, \quad R_4 = \begin{bmatrix} 0 & + & 0 & 0 \\ - & 0 & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & - \end{bmatrix}$$

requires algebraic positivity.

Proof. Let $B \in Q(S)$ for some sign pattern matrix

$$S = \begin{bmatrix} + & + & 0 & 0 \\ - & - & + & 0 \\ 0 & 0 & - & + \\ + & 0 & + & - \end{bmatrix}.$$

For suitable $\lambda > 0$, $B - \lambda I \in Q(R_1)$. If R_1 requires algebraic positivity, then S requires algebraic positivity, a contradiction to Lemma 2.9. So R_1 does not require algebraic positivity.

Now the matrices

$$B_2 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \\ 1 & 0 & 1 & -2 \end{bmatrix}, \quad B_3 = \begin{bmatrix} -1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & -2 & 1 \\ 1 & 0 & 1 & -1 \end{bmatrix}, \quad B_4 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \\ 1 & 0 & 1 & -2 \end{bmatrix}$$

are in $Q(R_2), Q(R_3), Q(R_4)$, respectively. Further, $B_2\bar{x} = B_3^T\bar{y} = B_4\bar{x} = \bar{0}$, where

$$\bar{x} = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \bar{y} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

So by Lemma 2.2, none among B_2, B_3, B_4 is algebraically positive. Therefore none among R_2, R_3, R_4 requires algebraic positivity. \square

Theorem 5.4. Let A be a 4×4 sign pattern matrix given by

$$A = \begin{bmatrix} a_{11} & a_{12} & 0 & a_{14} \\ a_{21} & a_{22} & a_{23} & 0 \\ 0 & a_{32} & a_{33} & a_{34} \\ a_{41} & 0 & a_{43} & a_{44} \end{bmatrix}, \quad (3)$$

where $a_{11}, a_{22}, a_{33}, a_{44}, a_{21}, a_{32}, a_{43}, a_{14} \in \{+, -, 0\}$, $a_{12}, a_{23}, a_{34}, a_{41} = +$ such that $a_{ij}a_{ji} = -$ for some $i \neq j$ and $a_{kl}a_{lk} = +$ for some $k \neq l$. Then A requires algebraic positivity if and only if A is permutationally similar to one of P_1, P_2, P_3, P_4, P_5 in Lemma 5.2.

Proof. If A is permutationally similar to one of P_1, P_2, P_3, P_4, P_5 , then A requires algebraic positivity by Lemma 5.2.

If A is of the form (3), by Lemma 2.9, no vertex in $D(A)$ is covered by both positive and negative directed 2-cycles. So we can assume that $a_{21} = -, a_{32} = 0, a_{43} = +, a_{14} = 0$. Then by Lemma 2.10, $a_{11} = -, a_{22} = -$. By Lemma 5.3, both a_{33}, a_{44} cannot be $-$. Therefore A is permutationally similar to one of P_1, P_2, P_3, P_4, P_5 in Lemma 5.2. \square

Theorem 5.1 and 5.4 together describes all 4-by-4 sign pattern matrices associated with cycle graphs that require algebraic positivity.

Declaration of competing interest

There is no competing interest associated with this work.

Data availability

No data was used for the research described in the article.

References

- [1] J. Abagat and D. Pelejo. On sign pattern matrices that allow or require algebraic positivity. *Electron. J. Linear Algebra*, 35: 331–356, 2019.
- [2] R.A. Brualdi and B.L. Shader. Matrices of sign-solvable linear systems. Cambridge University Press, Cambridge, 1995.
- [3] S. Das . Sign patterns that allow algebraic positivity. *Linear Algebra Appl.*, 653: 91–122, 2022.
- [4] S. Das and S. Bandopadhyay. On some sign patterns of algebraically positive matrices. *Linear Algebra Appl.*, 562: 91–122, 2019.
- [5] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, Second edition, 2013.
- [6] F J. Hall and Z. Li. *Sign pattern matrices*, in: Handbook of linear algebra, Edited by Leslie Hogben. Second edition, CRC Press, Boca Raton, FL, 2014.
- [7] S. Kirkland, P. Qiao, and X. Zhan. Algebraically positive matrices. *Linear Algebra Appl.*, 504: 14–26, 2016.