CONTACT LIE ALGEBRAS, GENERIC STABILISERS, AND AFFINE SEAWEEDS

OKSANA S. YAKIMOVA

ABSTRACT. Let q = Lie Q be a Lie algebra of index 1, i.e., a generic Q-orbit on q^* has codimension 1. We show that the following conditions are equivalent: q is contact; a generic Q-orbit on q^* is not conical; there is a generic stabiliser for the coadjoint action of q. In addition, if q is contact, then the subalgebra $S(q)_{si} \subset S(q)$ generated by symmetric semi-invariants of q is a polynomial ring. We study also affine seaweed Lie algebras of type A and find some contact as well as non-contact examples among them.

INTRODUCTION

Let \mathfrak{q} be a Lie algebra over a field \Bbbk of characteristic zero. For $\alpha \in \mathfrak{q}^*$, let $d_{\mathfrak{q}}\alpha \in \wedge^2 \mathfrak{q}^*$ be the image of α under the Chevalley–Eilenberg differential. Suppose that $\dim \mathfrak{q} = 2n + 1$. Then \mathfrak{q} is said to be *contact*, if there is $\alpha \in \mathfrak{q}^*$ such that $(\wedge^n d_{\mathfrak{q}}\alpha) \wedge \alpha \neq 0$. This definition originated from geometric constructions.

A contact structure on a smooth manifold M of dimension 2n+1 is a differential 1-form v such that $(dv)^n \wedge v \neq 0$ at each point of M, where dv is the de Rham differential of v. For information about contact geometry or topology, see e.g. [9]. According to Gromov [10], there is a contact structure on every odd-dimensional connected non-compact real Lie group Q. In general, such contact structures are not invariant under left translations by the group elements. Furthermore, Q admits an invariant contact form if and only if its Lie algebra Lie Q is contact, see e.g. [6, Sect. 2].

Let $q_{\alpha} \subset q$ be the kernel of the skew-symmetric bilinear form $d_{q}\alpha$. Then q_{α} is also the stabiliser of α w.r.t. the coadjoint action. We define the *index of* q by

(0.1)
$$\operatorname{ind} \mathfrak{q} = \min_{\gamma \in \mathfrak{q}^*} \dim \mathfrak{q}_{\gamma}$$

If $\alpha \in \mathfrak{q}^*$ is a contact linear function, i.e., $(\wedge^n d_\mathfrak{q} \alpha) \wedge \alpha \neq 0$, then dim $\mathfrak{q}_\alpha = 1$. Thereby each contact Lie algebra is of index 1. But not any \mathfrak{q} with ind $\mathfrak{q} = 1$ is contact. However, there are classes of Lie algebras, where these two properties are equivalent. For instance, this is

²⁰²⁰ Mathematics Subject Classification. 17B05, 17B08, 17B45, 53D10.

Key words and phrases. coadjoint action, conical orbit, generic stabiliser, seaweed, quasi-reductive.

This work is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) — project number 404144169.

O. YAKIMOVA

true for algebraic Lie algebras whose radicals consist of ad-nilpotent elements, see Corollary 3.4. This class includes all nilpotent Lie algebras, see also Example 2.4 for special feature of the nilpotent case.

Until the end of the Introduction assume that \Bbbk is algebraically closed and that \mathfrak{q} is an algebraic Lie algebra, i.e., $\mathfrak{q} = \operatorname{Lie} Q$, where Q is a connected affine algebraic group. In Section 2, we observe first that \mathfrak{q} with $\operatorname{ind} \mathfrak{q} = 1$ is contact if and only if a generic Q-orbit in \mathfrak{q}^* is not conical or, equivalently, if the extended group $\widetilde{Q} = Q \times \Bbbk^{\times}$ acts on \mathfrak{q}^* with an open orbit. This implies that a contact form is unique up to conjugation by elements of Q and scalar multiplication.

A linear form $\gamma \in \mathfrak{q}^*$ is *stable* in the terminology of [27] and its stabiliser $\mathfrak{q}_{\gamma} \subset \mathfrak{q}$ is a *generic stabiliser for the coadjoint action*, if there is a non-empty open subset $U \subset \mathfrak{q}^*$ such that \mathfrak{q}_{γ} and \mathfrak{q}_{β} are conjugate by an element of Q for each $\beta \in U$.

Our main result, Theorem 2.2, states that a Lie algebra q of index 1 is contact if and only if there is a generic stabiliser for the coadjoint action. In particular, if $\alpha \in q^*$ is contact, then it is stable.

In [4], it is shown that a seaweed subalgebra of $\mathfrak{sl}_r(\mathbb{C})$ or $\mathfrak{sp}_{2r}(\mathbb{C})$ that has index 1 is contact. To be more precise, the main theorem of [4] states that an index-one seaweed of a complex simple Lie algebra is contact if and only if it is quasi-reductive. By a result of Panyushev [18] each seaweed in \mathfrak{sl}_r or \mathfrak{sp}_{2r} is quasi-reductive. The equivalence of Theorem 2.2 explains, simplifies, and generalises arguments of [4], see Section 2.1.

The authors of [4] claim that their result provides a classification of contact seaweeds. Unfortunately, this is very far from the truth. In spite of many formulas for the index of a seaweed [5, 17, 11, 20], no one knows how to list all seaweeds of index 1 in \mathfrak{sl}_r or in \mathfrak{sp}_{2r} . In this article, we are not interested in classification aspects, but rather in properties of the coadjoint action of a contact Lie algebra.

In Section 3, we consider the subring $S(q)^q$ of symmetric invariants and the ring $S(q)_{si} \subset S(q)$ generated by semi-invariants of q. If q is contact, then $S(q)_{si} \subset S(q)$ is a finitely generated polynomial ring, see Proposition 3.6. Being contact is essential for our conclusion. There are non-contact Lie algebras of index 1 such that $S(q)_{si}$ is not a polynomial ring, see Example 3.12. Our proof relies on an old result of Sato–Kimura [24]. Their method applies also to Lie algebras of index zero and leads to a similar conclusion, see [21, Sect. 3.2]. A different approach to $S(q)_{si}$ in case ind q = 0 is developed in [15].

An important feature is that $S(\mathfrak{q}_{tr})^{\mathfrak{q}_{tr}}$ is a polynomial ring in $m = \operatorname{ind} \mathfrak{q}_{tr} = \operatorname{tr.deg} S(\mathfrak{q})_{si}$ variables for *the canonical truncation* $\mathfrak{q}_{tr} \subset \mathfrak{q}$ of a contact \mathfrak{q} ; furthermore, this property extends to each finite-dimensional quotient $\mathfrak{q}_{tr}[t]/(t^k)$ of the current algebra $\mathfrak{q}_{tr}[t]$, see Section 3.4. Non-reductive Lie algebras \mathfrak{s} such that $S(\mathfrak{s})^{\mathfrak{s}}$ is a polynomial ring with $\operatorname{ind} \mathfrak{s}$ generators attract a lot of attention, see e.g. [26, 11, 19, 3, 21]. A quest for this type of algebras continues. Our results provide another source of them.

The ring $S(\mathfrak{q})^{\mathfrak{q}}$ itself is less spectacular. If $\operatorname{ind} \mathfrak{q} = 1$, then either $S(\mathfrak{q})^{\mathfrak{q}} = \Bbbk$ or $S(\mathfrak{q})^{\mathfrak{q}}$ is generated by one homogeneous polynomial, see Section 3.1. If $\operatorname{ind} \mathfrak{q} = 1$ and $S(\mathfrak{q})^{\mathfrak{q}} \neq \Bbbk$, then \mathfrak{q} is contact by Proposition 3.3.

Section 4 is devoted to *affine seaweeds*. These Lie algebras are analogues of the usual seaweeds in the setting of loop algebras, see Section 2.1 for details on seaweeds in reductive Lie algebras. For one particular series of examples in type A_r , we derive an explicit formula for the index.

Let $\mathfrak{p} \subset \mathfrak{sl}_{r+1}$ be a maximal parabolic with the blocks of sizes a and b. Write $\mathfrak{p} = \mathfrak{l} \ltimes \mathfrak{n}$, where $\mathfrak{n} \cong \Bbbk^a \otimes (\Bbbk^b)^*$ is the nilpotent radical, which is Abelian. Add another copy of \mathfrak{n} obtaining $\mathfrak{q} = \overline{\mathfrak{q}}(a, b) = \mathfrak{l} \ltimes (\mathfrak{n} \oplus \mathfrak{n})$. Then \mathfrak{q} is an affine seaweed and ind $\mathfrak{q} = \gcd(2a, a+b) - 1$ by Theorem 4.1. If $\gcd(2a, a+b) = 2$, then $\operatorname{ind} \overline{\mathfrak{q}}(a, b) = 1$. A natural question is whether this Lie algebra is contact or not. As it turns out, the question is rather difficult.

In order to obtain an answer, we deal with semi-direct products $Q = L \ltimes \exp(V)$, where $\exp(V)$ is a normal Abelian unipotent subgroup, L acts on V^* with an open orbit $L\gamma$, and ind Lie Q = 1. Let L_{γ} be the stabiliser of γ and $L_{<\gamma>}$ the normaliser of $\Bbbk\gamma$. Certain properties of $\mathfrak{l}_{\gamma} = \operatorname{Lie} L_{\gamma}$ and $\mathfrak{l}_{<\gamma>} = \operatorname{Lie} L_{<\gamma>}$ are decisive factors here. For instance, if \mathfrak{l}_{γ} is not contact and ind $\mathfrak{l}_{<\gamma>} = 0$, then Lie Q is contact. A reduction from Q to L_{γ} is rather intricate, see Theorem 4.2 and Prposition 4.3. If a and b are even and $\operatorname{ind} \overline{\mathfrak{q}}(a, b) = 1$, then $\mathfrak{q}(a, b)$ is quasi-reductive and contact, see Theorem 4.4. On the contrary, $\overline{\mathfrak{q}}(1, b)$ is not contact for any odd b, see Example 4.5. There is an ample opportunity for further investigation of the coadjoint action of an affine seaweed.

1. PRELIMINARIES AND NOTATION

For an irreducible affine variety Y over \Bbbk , we let $\Bbbk[Y]$ stand for the ring of regular functions on Y and $\Bbbk(Y) = \text{Quot } \Bbbk[Y]$ for the field of rational functions on Y. A statement that a certain assertion holds for *generic points* of Y (or for generic orbits on Y) means that this assertion holds for all points of a non-empty open subset $U \subset Y$ (for all orbits intersecting U). If an algebraic group Q acts on Y, then $\Bbbk[Y]^Q$ is the ring of Q-invariant regular functions and $\Bbbk(Y)^Q$ is the field of Q-invariant rational functions on Y.

Suppose $\alpha \in \mathfrak{q}^*$. The kernel $\mathfrak{q}_\alpha \subset \mathfrak{q}$ of the skew-symmetric form $d_{\mathfrak{q}}\alpha$ is defined by

$$\mathfrak{q}_{\alpha} = \{\xi \in \mathfrak{q} \mid \alpha([\xi, \mathfrak{q}]) = 0\}.$$

It is the stabiliser of α in \mathfrak{q} . If $\mathfrak{q} = \text{Lie } Q$ and $Q_{\alpha} \subset Q$ is the stabiliser of α for the coadjoint action, then $\mathfrak{q}_{\alpha} = \text{Lie } Q_{\alpha}$. Therefore $\dim Q\alpha = \dim \mathfrak{q} - \dim \mathfrak{q}_{\alpha}$. Hence $\inf \mathfrak{q}$ is the minimal

codimension of a *Q*-orbit in q^* , see (0.1) for the definition of index. If *Q* is an algebraic group, then

(1.1)
$$\operatorname{ind} \mathfrak{q} = \operatorname{tr.deg} \mathbb{k}(\mathfrak{q}^*)^Q$$

by the Rosenlicht theorem, see [25, IV.2].

Set $\mathfrak{q}_{\mathsf{reg}}^* = \{\gamma \in \mathfrak{q}^* \mid \dim \mathfrak{q}_{\gamma} = \operatorname{ind} \mathfrak{q} \}$ and $\mathfrak{q}_{\mathsf{sing}}^* = \mathfrak{q}^* \setminus \mathfrak{q}_{\mathsf{reg}}^*$. We say that \mathfrak{q} has the *codim*-*n* property if $\dim \mathfrak{q}_{\mathsf{sing}}^* \leq \dim \mathfrak{q} - n$.

Let *V* be a finite-dimensional vector space over \Bbbk and *V*^{*} the dual space. Then $(V^*)^*$ is canonically isomorphic to *V*. For a subspace $W \subset V$, let $Ann(W) \subset V^*$ be the *annihilator* of *W*.

Suppose dim $\mathfrak{q} = 2n + 1$, while ind $\mathfrak{q} = 1$, and $\alpha \in \mathfrak{q}_{\mathsf{reg}}^*$. Then \mathfrak{q}^* has a basis $\{\xi_1, \ldots, \xi_{2n+1}\}$ such that $d_{\mathfrak{q}}\alpha = \xi_1 \wedge \xi_2 + \ldots + \xi_{2n-1} \wedge \xi_{2n}$. Here \mathfrak{q}_α is equal to $\operatorname{Ann}(\langle \xi_1, \ldots, \xi_{2n} \rangle_{\Bbbk}) \subset \mathfrak{q}$. Note that $(\wedge^n d_{\mathfrak{q}}\alpha) \wedge \alpha = 0$ if and only if $\alpha \in \langle \xi_1, \ldots, \xi_{2n} \rangle_{\Bbbk}$. The point α is contained in the subspace $\langle \xi_1, \ldots, \xi_{2n} \rangle_{\Bbbk}$ if and only if $\alpha \in \operatorname{Ann}(\mathfrak{q}_\alpha)$, i.e., if $\alpha(\mathfrak{q}_\alpha) = 0$. In other word, α is a contact form if and only if $\alpha(\mathfrak{q}_\alpha) \neq 0$. For a Lie algebra \mathfrak{q} of index 1, we have also

(1.2) \mathfrak{q} is contact $\Leftrightarrow \alpha(\mathfrak{q}_{\alpha}) \neq 0$ for a generic point $\alpha \in \mathfrak{q}_{\mathsf{reg}}^*$.

Over an algebraically closed field, an orbit Qy of a group Q is said to be *conical*, if $\Bbbk^{\times} y \subset Qy$. For the coadjoint representation

$$\mathrm{ad}^* \colon \mathfrak{q} \to \mathfrak{gl}(\mathfrak{q}^*)$$

and $\alpha \in \mathfrak{q}^*$, an equivalent condition is $\alpha \in \mathrm{ad}^*(\mathfrak{q}) \cdot \alpha$.

Lemma 1.1. For any finite-dimensional Lie algebra \mathfrak{q} and any $\alpha \in \mathfrak{q}^*$, there is an equivalence: $\alpha \in \mathrm{ad}^*(\mathfrak{q}) \cdot \alpha \Leftrightarrow \alpha(\mathfrak{q}_\alpha) = 0.$

Proof. If $\alpha \in ad^*(\mathfrak{q}) \cdot \alpha$, then $\alpha = ad^*(\xi)(\alpha)$ for some $\xi \in \mathfrak{q}$ and

$$\alpha(\mathfrak{q}_{\alpha}) = \mathrm{ad}^*(\xi)(\alpha)(\mathfrak{q}_{\alpha}) = -\alpha([\xi, \mathfrak{q}_{\alpha}]) = 0.$$

Suppose now that $\alpha(\mathfrak{q}_{\alpha}) = 0$. Then $\alpha \in \operatorname{Ann}(\mathfrak{q}_{\alpha})$. A standard fact is that $\operatorname{Ann}(\mathfrak{q}_{\alpha}) = \operatorname{ad}^{*}(\mathfrak{q}) \cdot \alpha$, it follows from dimension reasons. Hence $\alpha \in \operatorname{ad}^{*}(\mathfrak{q}) \cdot \alpha$.

In most of the paper, we assume that $k = \overline{k}$. This is a natural assumption, since the property of being contact does not change under field extensions.

Whenever dealing with classical Lie algebras, we assume that E_{ij} are elementary matrices (matrix units).

If *V* is a vector space and *k* a natural number, then kV is a direct sum of *k* copies of *V*.

2. LIE ALGEBRAS OF INDEX 1

Proposition 2.1. Suppose that $\operatorname{ind} \mathfrak{q} = 1$. Then \mathfrak{q} is contact if and only if $\alpha \notin \operatorname{ad}^*(\mathfrak{q}) \cdot \alpha$ for a generic point $\alpha \in \mathfrak{q}^*$. For an algebraic \mathfrak{q} over an algebraically closed field, this condition means that a generic *Q*-orbit in \mathfrak{q}^* is not conical.

Proof. By Lemma 1.1, α is contained in $\operatorname{ad}^*(\mathfrak{q}) \cdot \alpha$ if and only if $\alpha(\mathfrak{q}_{\alpha}) = 0$. The desired equivalence follows now from (1.2).

Note that in case $\mathbb{k} = \overline{\mathbb{k}}$, an orbit $Q\alpha \subset \mathfrak{q}^*$ is conical if and only if $\operatorname{ad}^*(\mathfrak{q}) \cdot \alpha$ contains $\mathbb{k}\alpha$.

Unless otherwise stated, we assume from now on that $\mathbb{k} = \overline{\mathbb{k}}$, Q is a connected affine algebraic group, and $\mathfrak{q} = \text{Lie } Q$.

Theorem 2.2. *A Lie algebra* **q** *of index* 1 *is contact if and only if there is a generic stabiliser for the coadjoint action.*

Proof. Suppose first that q is contact. Choose $\alpha \in \mathfrak{q}_{\mathsf{reg}}^*$ such that $\alpha(\mathfrak{q}_\alpha) \neq 0$. Then the orbit $Q\alpha$ is not conical by Lemma 1.1 and Proposition 2.1. Thereby $Y = \Bbbk^{\times}Q\alpha$ is a dense open subset of \mathfrak{q}^* . Clearly \mathfrak{q}_α and \mathfrak{q}_γ are conjugate by an element of Q for each $\gamma \in Y$. Thus, \mathfrak{q}_α is a generic stabiliser for the coadjoint action.

Suppose now that \mathfrak{q} is not contact, but \mathfrak{q}_{α} with $\alpha \in \mathfrak{q}_{\mathsf{reg}}^*$ is a generic stabiliser for the coadjoint action. Let $y \in \mathfrak{q}_{\alpha}$ be a non-zero vector. We show that $[\mathfrak{q}, y]$ contains y.

It suffices to prove that $\gamma(y) = 0$ for every $\gamma \in Ann([\mathfrak{q}, y])$. Note that

Ann(
$$[\mathfrak{q}, y]$$
) = { $\beta \in \mathfrak{q}^* \mid y \in \mathfrak{q}_{\beta}$ }.

In particular, $\alpha \in \operatorname{Ann}([\mathfrak{q}, y])$. Also α is a regular point of \mathfrak{q}^* , thereby $R = \mathfrak{q}^*_{\mathsf{reg}} \cap \operatorname{Ann}([\mathfrak{q}, y])$ is a dense open subset of $\operatorname{Ann}([\mathfrak{q}, y])$. For each point $\gamma \in R$, we have $\gamma(\mathfrak{q}_{\gamma}) = 0$ by (1·2). Since here $y \in \mathfrak{q}_{\gamma}$ and dim $\mathfrak{q}_{\gamma} = 1$, the equality $\gamma(y) = 0$ holds. Clearly, it extends from R to all points of $\operatorname{Ann}([\mathfrak{q}, y])$.

By a criterion [27, Corollaire 1.8] of Tauvel and Yu, $q_{\alpha} \cap [q, q_{\alpha}] = 0$, because $q_{\alpha} = ky$ is a generic stabiliser. This provides a contradiction, since $y \in q_{\alpha} \cap [q, q_{\alpha}]$.

2.1. Seaweeds and quasi-reductive Lie algebras. Let $\mathfrak{g} = \text{Lie } G$ be a simple Lie algebra and $\mathfrak{p}_1, \mathfrak{p}_2 \subset \mathfrak{g}$ two parabolic subalgebras such that $\mathfrak{p}_1 + \mathfrak{p}_2 = \mathfrak{g}$. Then $\mathfrak{q} = \mathfrak{p}_1 \cap \mathfrak{p}_2$ is a Lie algebra of *seaweed type* or just a seaweed also called a *bi-parabolic*. Each seaweed \mathfrak{q} is an algebraic Lie algebra and $\mathfrak{q} = \text{Lie } Q$, where $Q \subset G$ is the intersection of two parabolic subgroups. In [18], Panyushev conjectured that if \mathfrak{q}_γ with $\gamma \in \mathfrak{q}^*$ is a generic stabiliser for the coadjoint action of a seaweed $\mathfrak{q} \subset \mathfrak{g}$, then Q_γ is reductive. If there is a stable point $\alpha \in q^*$, then q is called *stable* as well. This terminology is not standard, but it is used, for example, in [4].

Following [7], say that q is *quasi-reductive* if there is $\beta \in q^*$ such that the quotient Q_β/Z by the centre $Z \subset Q$ is a reductive subgroup of $GL(q^*)$. For more information on these Lie algebras see e.g. [7, 14].

Suppose Q_{β}/Z is a reductive subgroup of $GL(\mathfrak{q}^*)$. Then there is a generic stabilliser for the action of Q/Z on \mathfrak{q}^* , which is equal to a generic stabiliser for the action of Q_{β}/Z on \mathfrak{q}^*_{β} , see e.g. [14, Lemma 2.3.]. Thus, a quasi-reductive \mathfrak{q} is stable and its generic stabiliser \mathfrak{q}_{α} is a sum Lie $Z \oplus \text{Lie}(\mathbb{k}^{\times})^{\ell}$, where ℓ is the rank of Q_{β}/Z .

Since the centre of a seaweed $q \subset g$ consists of semisimple elements, we may reformulate Panyushev's conjecture as follows: q is stable if and only if it is quasi-reductive. The conjecture is proven by Ammari [1] on a case-by-case basis. Combining this with Theorem 2.2, we derive the equivalence: a seaweed of index 1 is contact if and only if it is quasi-reductive, which is the main result of [4].

Note that the argument in [4] goes as follows:

stable
$$\stackrel{\text{Ammari}}{\Longrightarrow}$$
 quasi-reductive \implies contact \implies stable

The first paragraph of the proof of Theorem 2.2 shows that the last implication is quite straightforward. The implications "quasi-reductive \implies stable \implies contact" are true for any Lie algebra of index 1. It is possible to see directly, why a quasi-reductive \mathfrak{q} with ind $\mathfrak{q} = 1$ has to be contact. An ingredient here is the fact that for a generic point $\alpha \in \mathfrak{q}^*$, its stabiliser \mathfrak{q}_{α} consists of ad-nilpotent elements, if \mathfrak{q} is of index 1, but not contact.

There is no equivalence between "contact" and "quiasi-reductive". There are many example such that Q_{α} is unipotent and not central for a contact form $\alpha \in q^*$. Below is one of them.

Example 2.3. Let $Q = SL_2 \ltimes \exp(\mathbb{k}^2)$ be a semi-direct product of SL_2 and a two-dimensional Abelian unipotent group. Then $\operatorname{ind} \mathfrak{q} = 1$ and $Z = \{e\}$. Let $\{x, y\}$ be a basis of the nilpotent radical of \mathfrak{q} such that $[E_{12}, x] = 0$ and $[E_{12}, y] = x$. Take $\alpha \in \mathfrak{q}^*$ such that $\alpha(y) = 1 = \alpha(E_{12})$ and $\alpha(x) = \alpha(E_{21}) = \alpha(E_{11} - E_{22}) = 0$. Then \mathfrak{q}_α is spanned by $u = E_{12} + 2y$. Since $\alpha(u) = 3 \neq 0$, the orbit $Q\alpha$ is not conical and $\mathbb{k}u$ is a generic stabiliser for the coadjoint action.

We end this section by addressing a very special type of Lie algebras, namely, nilpotent.

Example 2.4. Let $\mathfrak{n} = \text{Lie } N$ be a nilpotent Lie algebra of index 1. For instance, \mathfrak{n} may be a 3-dimensional Heisenberg Lie algebra with a basis $\{x, y, z\}$ such that z is central and [x, y] = z. The centre \mathfrak{z} of \mathfrak{n} is non-zero and $\mathfrak{z} \subset \mathfrak{n}_{\alpha}$ for any $\alpha \in \mathfrak{n}^*$. Thus, $\mathfrak{n}_{\alpha} = \mathfrak{z}$ for any $\alpha \in \mathfrak{n}^*_{\text{reg}}$. Then \mathfrak{z} is the generic stabiliser for the coadjoint action of \mathfrak{n} . Clearly $\alpha(\mathfrak{z}) \neq 0$ for

a generic α and \mathfrak{n} is contact. Since N_{α} is connected, it is equal to the centre $Z \subset N$. Hence $N_{\alpha}/Z = \{e\}$ and \mathfrak{n} is quasi-reductive.

3. INVARIANTS AND SEMI-INVARIANTS

Symmetric algebra $S(\mathfrak{q})$ is identified with the graded ring $\Bbbk[\mathfrak{q}^*]$ of polynomial functions on \mathfrak{q}^* . The adjoint action of \mathfrak{q} extends to a representation of \mathfrak{q} on $S(\mathfrak{q})$. Since Q is connected, we have $\Bbbk[\mathfrak{q}^*]^Q = S(\mathfrak{q})^Q = S(\mathfrak{q})^{\mathfrak{q}}$. The elements of this ring are called *symmetric invariants* of \mathfrak{q} . For $F \in S(\mathfrak{q})$, we have

$$F \in \mathfrak{S}(\mathfrak{q})^{\mathfrak{q}} \Leftrightarrow \xi \cdot F = 0 \; \forall \xi \in \mathfrak{q}.$$

Then $F \in S(\mathfrak{q})$ is called a *semi-invariant of* \mathfrak{q} if $QF \subset \Bbbk F$ or equivalently if $\xi \cdot F \in \Bbbk F$ for each $\xi \in \mathfrak{q}$. Let $S(\mathfrak{q})_{si} \subset S(\mathfrak{q})$ be the subring generated by the semi-invariants of \mathfrak{q} . Note that both $S(\mathfrak{q})^{\mathfrak{q}}$ and $S(\mathfrak{q})_{si}$ are graded subalgebras.

3.1. **General facts.** Let $f \in \mathbb{k}(\mathfrak{q}^*)^Q$ be a rational *Q*-invariant. Then f = u/v for some coprime polynomials $u, v \in \mathbb{k}[\mathfrak{q}^*]$. Such a presentation is unique up to scalar multiples. Therefore *u* and *v* are semi-invariants of *Q*.

Suppose ind $\mathfrak{q} = 1$. Then $\operatorname{tr.deg} \mathbb{k}(\mathfrak{q}^*)^Q = 1$ by (1·1). Since $\operatorname{Quot}(\mathbb{k}[\mathfrak{q}^*]^Q)$ is contained in $\mathbb{k}(\mathfrak{q}^*)^Q$, we obtain $\operatorname{tr.deg} S(\mathfrak{q})^{\mathfrak{q}} \leq 1$. It may happen that $S(\mathfrak{q})^{\mathfrak{q}} = \mathbb{k}$. Suppose $S(\mathfrak{q})^{\mathfrak{q}} \neq \mathbb{k}$. Then $S(\mathfrak{q})^{\mathfrak{q}}$ is generated by one element. Below we give an explanation of this standard fact.

Let $F, F_1 \in S(\mathfrak{q})^{\mathfrak{q}}$ be two non-constant homogeneous polynomials. Then they are algebraically dependent and hence $F^a = sF_1^b$ for some coprime natural numbers a, b and $s \in \Bbbk^{\times}$. Thereby $H = \sqrt[b]{F} \in \Bbbk[\mathfrak{q}^*]$ and the orbit QH has at most b elements. Since the group Q is connected, $H \in \Bbbk[\mathfrak{q}^*]^Q$. If we take F to be an invariant of the minimal degree, then $F_1 \in \Bbbk[F]$. Thus $\Bbbk[\mathfrak{q}^*] = \Bbbk[F]$.

In any case, $\Bbbk(\mathfrak{q}^*)^Q \neq \Bbbk$. There is at least one rational invariant $f = u/v \notin \Bbbk$, where $u, v \in S(\mathfrak{q})_{si}$. Hence $S(\mathfrak{q})_{si} \neq \Bbbk$. In view of classical results of Lüroth and Castelnouvo, $\Bbbk(\mathfrak{q}^*)^Q = \Bbbk(f)$ for some f.

Let $\mathcal{W} = \bigoplus_{i=0}^{\dim \mathfrak{q}} \mathcal{W}^i$ be the graded skew-symmetric algebra of polynomial vector fields on \mathfrak{q}^* . Over $\mathcal{S}(\mathfrak{q})$ it is generated by $\{\partial_i = \partial_{x_i} | 1 \leq i \leq N\}$, where $\{x_i | 1 \leq i \leq N\}$ is a basis of \mathfrak{q} and $N = \dim \mathfrak{q}$. Let $\pi = \sum_{i < j} [x_i, x_j] \partial_i \wedge \partial_j \in \mathcal{W}^2$ be the *Poisson tensor* of \mathfrak{q} . Evaluating π at a point $\gamma \in \mathfrak{q}^*$, we obtain $\pi(\gamma) = d_{\mathfrak{q}}\gamma$. Then $d = \frac{1}{2}(\dim \mathfrak{q} - \operatorname{ind} \mathfrak{q})$ is the largest number such that $\bigwedge^d \pi \neq 0$.

Definition 3.1. A polynomial p is a *fundamental semi-invariant* of q, if $\wedge^d \pi = p \Re$ with $\Re \in W^{2d}$ and the zero set of \Re in q^* has codimension grater than or equal to 2.

The zero set of p is the union of divisors contained in q_{sing}^* . Thereby p is indeed a semiinvariant of q. By the construction, q has the codim–2 property if and only if p = 1. *Remark* 3.2. For any $\mathfrak{q} \neq 0$, we have $\mathfrak{S}(\mathfrak{q})_{\mathfrak{s}\mathfrak{i}} \neq \Bbbk$. If $\operatorname{ind} \mathfrak{q} \geq 1$, then the argument with the inclusion $\Bbbk(\mathfrak{q}^*)^Q \subset \operatorname{Quot} \mathfrak{S}(\mathfrak{q})_{\mathfrak{s}\mathfrak{i}}$ works. If $\operatorname{ind} \mathfrak{q} = 0$, then N = 2d and $\deg p = d \geq 1$ for a non-zero \mathfrak{q} . Lie algebras of index zero are called *Frobenius*.

3.2. Importance of regular invariants.

Proposition 3.3. If ind q = 1 and $S(q)^q = k[F]$ for a non-constant F, then q is contact.

Proof. Since *F* is constant on each orbit $Q\alpha \subset \mathfrak{q}^*$, the orbit cannot be conical, if $F(\alpha) \neq 0$. The rest follows from Proposition 2.1.

Corollary 3.4. If ind q = 1 and the radical of q consists of ad-nilpotent elements, then q is contact.

Proof. Let $\operatorname{Rad}(\mathfrak{q}) \lhd \mathfrak{q}$ be the radical of \mathfrak{q} . Each $x \in \operatorname{Rad}(\mathfrak{q})$ is ad-nilpotent, hence it acts as a nilpotent element on every $S^k(\mathfrak{q})$. Therefore each semi-invariant of \mathfrak{q} in $S(\mathfrak{q})$ is invariant under x. In other words, $\operatorname{Rad}(\mathfrak{q})$ acts trivially on each \mathfrak{q} -stable line $\Bbbk u \subset S(\mathfrak{q})$. Then $\Bbbk u$ gives rise to a 1-dimensional representation of the semisimple algebra $\mathfrak{q}/\operatorname{Rad}(\mathfrak{q})$. Thus, each semi-invariant of \mathfrak{q} is an invariant and $S(\mathfrak{q})^{\mathfrak{q}} \neq \Bbbk$.

Extending the ground field we do not violate the conditions of Corollary 3.4. Hence its statement holds over any field.

There are contact Lie algebras with no regular invariants.

Example 3.5. (a) A Borel subalgebra $\mathfrak{b} \subset \mathfrak{sl}_3$ is contact. A generic stabiliser for \mathfrak{b}^* is a subspace of the Cartan subalgebra that commutes with the centre of the nilpotent radical. One sees easily that $\mathfrak{S}(\mathfrak{b})^{\mathfrak{b}} = \Bbbk$.

(b) Consider a maximal parabolic subalgebra $q \subset \mathfrak{sp}_4$ such that $q = \mathfrak{gl}_2 \ltimes \mathfrak{n}$, where the nilpotent radical \mathfrak{n} is a 3-dimensional Heisenberg Lie algebra. It is contact and a generic stabiliser for q^* is a Cartan subalgebra of \mathfrak{sl}_2 . Proper parabolic subalgebras of simple Lie algebras have only trivial symmetric invariants, see e.g. [18, Corollary 2.11].

3.3. A description of semi-invariants.

Proposition 3.6. Let \mathfrak{q} be a contact Lie algebra. Then $S(\mathfrak{q})_{si}$ is a polynomial algebra freely generated by irreducible homogeneous elements H_1, \ldots, H_m . Furthermore, the fundamental semiinvariant p of \mathfrak{q} is equal to $\prod_{i=1}^m H_i^{a_i}$ with $a_i \ge 0$.

Proof. Set $\tilde{Q} = Q \times \Bbbk^{\times}$. By Proposition 2.1, this group acts on \mathfrak{q}^* with an open orbit. By [24, §4], the algebra of \tilde{Q} -semi-invariants in $\Bbbk[\mathfrak{q}^*]$ is polynomial. More precisely, let $\tilde{Q}\alpha \subset \mathfrak{q}^*$ be the open \tilde{Q} -orbit, it appeared in the proof of Theorem 2.2 as the subset Y. Let D_1, \ldots, D_m be all simple divisors in $\mathfrak{q}^* \setminus \tilde{Q}\alpha$, we do not need the irreducible components of codimension ≥ 2 in \mathfrak{q}^* . If $D_i = \{H_i = 0\}$ and $H_i \in \Bbbk[\mathfrak{q}^*]$ is irreducible, then H_i is a semi-invariant of \tilde{Q} and H_1, \ldots, H_m freely generate the algebra of \tilde{Q} -semi-invariants in $\mathbb{k}[\mathfrak{q}^*]$.

Clearly each semi-invariant of \widetilde{Q} is also a semi-invariant of Q. Suppose that $QF \subset \Bbbk^{\times} F$ for some $F \in S(\mathfrak{q})$. Then each homogeneous component of F is also a semi-invariant of Q, and hence of \widetilde{Q} . This leads to the equality $S(\mathfrak{q})_{si} = \Bbbk[H_1, \ldots, H_m]$.

Note that $\widetilde{Q}\alpha \subset \mathfrak{q}^*_{reg}$. Let p_i be a prime factor of p. Then the zero set of p_i is a simple divisor in the complement of $\widetilde{Q}\alpha$. Hence p_i is equal to some H_j up to a non-zero scalar. \Box

Remark 3.7. If $\alpha \in \mathfrak{q}^*$ is a contact linear form, then $\widetilde{Q}\alpha$ is a dense open subset of \mathfrak{q}^* . This implies that a contact form is unique up to conjugation by elements of Q and scalar multiplication. Over a non-closed field, for example, over \mathbb{R} , the situation is different. Take $Q = \operatorname{SL}_2(\mathbb{R}) \ltimes \exp(2\mathbb{R}^2)$. Then \mathfrak{q} is contact. The group $\widetilde{Q} = Q \times \mathbb{R}^\times$ has two different orbits on \mathfrak{q}^* that consists of contact points. Studying contact orbits over \mathbb{R} is an excellent direction for further research.

Unlike a Frobenius Lie algebra, a contact one may have the codim–2 property. For instance, $\mathfrak{g} = \mathfrak{sl}_2$ is contact. The only singular point in $\mathfrak{g}^* \cong \mathfrak{g}$ is zero. A non-zero semisimple element defines a contact form, while the nilpotent orbit is conical, but has dimension 2 as well.

Remark 3.8. A contact Lie algebra q has another very important semi-invariant, which is never a constant. It is described in matrix terms in [23, Sect. 4]. Set $v = \sum_{i=1}^{\dim \mathfrak{q}} x_i \partial_i \in \mathcal{W}^1$. Then $v(\gamma) = \gamma$ for any $\gamma \in \mathfrak{q}^*$. Recall that $\pi(\gamma) = d_{\mathfrak{q}}\gamma$. Suppose dim $\mathfrak{q} = 2n+1$. The exterior product $(\bigwedge^n \pi) \land v$ is non-zero at $\alpha \in \mathfrak{q}^*$ if and only if α is contact. We have also

$$(\wedge^n \pi) \wedge v = \mathbf{f} \partial_1 \wedge \ldots \wedge \partial_{2n+1},$$

where f is a homogeneous polynomial of degree n + 1. The zero set of f is exactly the complement $q^* \setminus \tilde{Q}\alpha$, where α is a contact form. Then $f = c \prod_{i=1}^m H_i^{a_i} \in S(q)_{s_i}$ with $a_i \ge 1$ and $c \in \Bbbk^{\times}$. By the construction, p|f.

Example 3.9. (1) If $\mathfrak{g} = \mathfrak{sl}_2$, then *f* is the determinant (up to a non-zero scalar).

(2) For a Heisenberg Lie algebra $k^{2n} \oplus kz$ of dimension 2n+1, we have $p = z^n$ and $f = z^{n+1}$.

Example 3.10. Consider Lie algebras of Example 3.5. For $\mathfrak{b} = \text{Lie } B$, we choose a basis consisting of

$$h = \text{diag}(1, -2, 1), h_1 = \text{diag}(1, 0, -1), x = E_{12}, y = E_{23}, z = [x, y].$$

Then $\mathcal{S}(\mathfrak{b})_{si} = \Bbbk[z, H_2]$ with $H_2 = hz + 3xy$. If $\gamma(z) \neq 0$ for $\gamma \in \mathfrak{b}^*$, then $B\gamma$ contains a point $\tilde{\gamma}$ such that $\tilde{\gamma}(z) = \gamma(z)$ and $\tilde{\gamma}(x) = \tilde{\gamma}(y) = 0$. Clearly $\mathfrak{b}_{\tilde{\gamma}} = \Bbbk h$ and $\gamma \in \mathfrak{b}_{reg}^*$. Therefore \mathfrak{b}_{sing}^* is contained in $D_1 = \{z = 0\}$. However, $\mathfrak{b}_{\beta} = \Bbbk z$ for any $\beta \in D_1$ such that $\beta(x) \neq 0$ and $\beta(y) \neq 0$. Thus, p = 1. Since deg $\mathbf{f} = 3$, we have $\mathbf{f} \in \Bbbk z H_2$, actually $\mathbf{f} = 2zH_2$.

O. YAKIMOVA

For $\mathfrak{q} \subset \mathfrak{sp}_4$ from part (b), we have $S(\mathfrak{q})_{\mathfrak{si}} = \Bbbk[z, H_2]$, where z is again a non-zero element in the centre of the nilpotent radical and deg $H_2 = 3$. Here $\mathbf{f} = zH_2$ up to a scalar and deg $p \leq 3$, since dim $\mathfrak{b} = 7$. The semi-direct product $\mathfrak{gl}_2 \ltimes \Bbbk^2$ is a Frobenius Lie algebra. This implies $\mathfrak{q}_\beta = \Bbbk z$ for a generic point of $D_1 = \{z = 0\}$. In a generic $Q\gamma \subset D_2$, there is a point $\tilde{\gamma}$ such that $\tilde{\gamma}(z) = 1$, $\tilde{\gamma}(\Bbbk^2) = 0$, and $\bar{\gamma} = \tilde{\gamma}|_{\mathfrak{sl}_2}$ is a non-zero nilpotent element. Then $\mathfrak{q}_{\tilde{\gamma}} = (\mathfrak{sl}_2)_{\tilde{\gamma}}$ is of dimension 1. Therefore D_1 and D_2 are not contained in \mathfrak{q}^*_{sing} and p = 1.

Suppose \mathfrak{q} is contact and let H_1, \ldots, H_m be the generators of $\mathfrak{S}(\mathfrak{q})_{\mathfrak{s}i}$ described in the proof of Proposition 3.6. Let $\chi_i : \widetilde{Q} \to \Bbbk^{\times}$ be the \widetilde{Q} -character corresponding to H_i . We have $g \cdot H_i = \chi_i(g)H_i$ for all $g \in \widetilde{Q}$. Then the differentials $d\chi_i : (\mathfrak{q} \oplus \Bbbk) \to \Bbbk$ with $1 \leq i \leq m$ are linearly independent, because \widetilde{Q} has only trivial rational invariants on \mathfrak{q}^* , see also [24, §4].

Let $\bar{\chi}_i$ be the restriction of $d\chi_i$ to q. Then $\dim \langle \bar{\chi}_1, \ldots, \bar{\chi}_m \rangle_{\Bbbk} = m - 1$. Since the character group $\mathfrak{X}(\tilde{Q})$ is a lattice, any relation between $\bar{\chi}_i$ has integer coefficients up to scalar multiples. Up to a suitable numeration, the minimal relation looks as

(3.1)
$$\sum_{i=1}^{a} c_i \bar{\chi}_i = \sum_{i=a+1}^{b} c_i \bar{\chi}_i,$$

where $c_i \ge 1$ for each i and $gcd(c_1, \ldots, c_b) = 1$. Assume that $a \ge (b-a)$. If b = a, then $F = \prod_{i=1}^{a} H_i^{c_i} \in \Bbbk[\mathfrak{q}^*]^Q$ and $\Bbbk(\mathfrak{q}^*)^Q = \Bbbk(F) = Quot(\Bbbk[\mathfrak{q}^*]^Q)$. If b > a, then $f = (\prod_{i=1}^{a} H_i^{c_i})(\prod_{i=a+1}^{b} H_i^{-c_i})$ is a non-regular generator of $\Bbbk(\mathfrak{q}^*)^Q$.

Remark 3.11. Let $H \in S(q)_{si}$ be homogeneous. Then the zero set D of H is a \tilde{Q} -stable proper closed subset of q^* . Hence $D \cap \tilde{Q}\alpha = \emptyset$ for the open orbit $\tilde{Q}\alpha \subset q^*$ and H is a product of the polynomials H_i . There are also non-homogeneous semi-invariants. For instance, $z + H_2$ for the Lie algebra from Example 3.5(a), see Example 3.10.

Example 3.12. There are Lie algebras of index 1 such that $S(\mathfrak{q})_{si}$ is not a polynomial ring. Consider first a semi-direct product $\mathfrak{s} = \mathfrak{sl}_2 \ltimes 4\mathbb{k}^2$, where $4\mathbb{k}^2$ is an Abelian ideal. This Lie algebra is of dimension 11. Generic SL₂-orbits on $4\mathbb{k}^2$ are of dimension 3. By a formula of Raïs [22] for the index of a semi-direct product, $\operatorname{ind} \mathfrak{s} = 8 - 3 = 5$. The ring $S(\mathfrak{s})^{\mathfrak{s}} = \mathbb{k}[(4\mathbb{k}^2)^*]^{\operatorname{SL}_2}$ is generated by $\binom{4}{2} = 6$ invariants of degree 2. There is one relation among them. We extend \mathfrak{s} to $\tilde{\mathfrak{s}}$ by adding four copies of \mathbb{k} each acting on one \mathbb{k}^2 . A generic stabiliser for the obtained action of $(\mathbb{k}^{\times})^4 \times \operatorname{SL}_2$ on $4\mathbb{k}^2$ is still trivial. Hence $\operatorname{ind} \tilde{\mathfrak{s}} = 8 - 7 = 1$. Now $S(\tilde{\mathfrak{s}})_{\mathfrak{s}\mathfrak{i}} = S(\mathfrak{s})^{\mathfrak{s}}$ is not a polynomial ring. We can extend this example to any semi-direct product $(\mathfrak{sl}_{2n+2} \oplus 4\mathbb{k}) \ltimes 4\mathbb{k}^{2n+2}$ with $n \ge 1$ arguing in the spirit of [28].

The following example shows that $S(q)_{si}$ may be a polynomial ring for a non-contact Lie algebra of index 1, there is no equivalence in Proposition 3.6.

Example 3.13. Consider $q = \Bbbk \ltimes V$, where [V, V] = 0, dim V = 2, and \Bbbk acts on V with two non-zero characters α and β . Then $S(q)_{si} = S(V)$ is always a polynomial ring. However, q is contact if and only if $\alpha \neq \beta$.

3.4. The canonical truncation. Let $q_{tr} \subset q$ be the intersection of all kernels ker $\bar{\chi}_i$ of the characters $\bar{\chi}_i$ defined above. Then dim $q_{tr} = \dim q - m + 1$. This subalgebra is called *the canonical truncation of* q. Its crucial feature is that $S(q)_{si} \subset S(q_{tr})$, see [2, Kap. II, §6]. Actually, $S(q)_{si} \subset S(q_{tr})^{q_{tr}}$. Since q acts diagonalisably on $S(q_{tr})_{si}$, we have

$$S(\mathfrak{q}_{\mathrm{tr}})_{\mathsf{si}} \subset S(\mathfrak{q})_{\mathsf{si}} \subset S(\mathfrak{q}_{\mathrm{tr}})^{\mathfrak{q}_{\mathrm{tr}}}.$$

Hence $\mathcal{S}(\mathfrak{q}_{tr})_{si} = \mathcal{S}(\mathfrak{q}_{tr})^{\mathfrak{q}_{tr}} = \mathcal{S}(\mathfrak{q})_{si} = \mathbb{k}[H_1, \ldots, H_m]$, which implies $\mathbb{k}(\mathfrak{q}_{tr}^*)^{\mathfrak{q}_{tr}} = \operatorname{Quot}(\mathbb{k}[\mathfrak{q}_{tr}^*]^{\mathfrak{q}_{tr}})$ and $m = \operatorname{tr.deg} \mathbb{k}[\mathfrak{q}_{tr}^*]^{\mathfrak{q}_{tr}} = \operatorname{ind} \mathfrak{q}_{tr}$. The equality $\operatorname{ind} \mathfrak{q}_{tr} = \operatorname{ind} \mathfrak{q} + (m-1) = m$ follows also from a general formula obtained in [16, Lemma 3.7].

We have seen that if q is contact, then q_{tr} has a remarkable property. That is, the ring $S(q_{tr})^{q_{tr}}$ is freely generated by $m = ind q_{tr}$ elements. Repeating the argument of [21, Remark 3.1] one obtains further results as follows. Note that q_{tr} has no proper semiinvariants in $S(q_{tr})$. Then by [13, Prop. 5.2] the differentials dH_i are linearly independent in codimension 2, i.e.,

$$\dim\{\xi \in \mathfrak{q}_{\mathrm{tr}}^* \mid d_{\xi}H_1 \wedge \ldots \wedge d_{\xi}H_m = 0\} \leqslant \dim \mathfrak{q}_{\mathrm{tr}} - 2.$$

By [21, Thm. 2.2], each finite-dimensional quotient $\mathfrak{w}_k := \mathfrak{q}_{tr}[t]/(t^k)$ of the current algebra $\mathfrak{q}_{tr}[t]$ has the same property: the ring of symmetric invariants of \mathfrak{w}_k is freely generated by $k \cdot \operatorname{ind} \mathfrak{q}_{tr} = \operatorname{ind} \mathfrak{w}_k$ elements.

4. AFFINE SEAWEED SUBALGEBRAS

Let \mathfrak{g} be a simple finite-dimensional non-Abelian Lie algebra. We fix a Borel subalgebra $\mathfrak{b} \subset \mathfrak{g}$ and a Cartan subalgebra $\mathfrak{t} \subset \mathfrak{b}$. Let $\Pi = \{\alpha_1, \ldots, \alpha_r\}$, where $r = \mathsf{rk}\,\mathfrak{g}$, be the set of simple roots associated with $(\mathfrak{b}, \mathfrak{t})$. For any root α of $(\mathfrak{g}, \mathfrak{t})$, let $e_\alpha \in \mathfrak{g}$ be a root vector. To a subset $S \subset \Pi$ one associates a standard parabolic subalgebra $\mathfrak{p}(S)$, which is generated by \mathfrak{b} and $\{e_{-\alpha} \mid \alpha \in S\}$. Then let $\mathfrak{p}(S)^-$ be the opposite parabolic, which is generated by \mathfrak{t} together with $\{e_\alpha \mid \alpha \in S\}$ and $\{e_{-\alpha} \mid \alpha \in \Pi\}$. A seaweed in \mathfrak{g} is called *standard* if it is of the form $\mathfrak{p}(S) \cap \mathfrak{p}(T)^-$ for two subsets $S, T \subset \Pi$. By [17], any seaweed in \mathfrak{g} is conjugate to a standard one. We will consider standard seaweeds of the loop algebra $\mathfrak{g}[t, t^{-1}]$.

Let $\widetilde{\Pi} = \{\alpha_0\} \sqcup \Pi$ be the affine root system associated with Π and δ the highest root of \mathfrak{g} . Set $\widehat{\mathfrak{b}} = \mathfrak{b} \oplus t\mathfrak{g}[t]$ and $e_{-\alpha_0} = e_{\delta}t^{-1} \in \mathfrak{g}[t^{-1}]$. To a subset $S \subset \widetilde{\Pi}$, one associates a standard parabolic $\mathfrak{p} = \mathfrak{p}(S) \subset \mathfrak{g}[t, t^{-1}]$, which is generated by $\widehat{\mathfrak{b}}$ and $\{e_{-\beta} \mid \beta \in S\}$.

Let $\mathfrak{p}(S \cap \Pi)$ be the standard parabolic of \mathfrak{g} associated with $S \cap \Pi$. For any Lie algebra \mathfrak{q} , let $\mathfrak{U}(\mathfrak{q})$ be its enveloping algebra. We have

$$\mathfrak{p} = t\mathfrak{g}[t] \oplus \mathfrak{p}(S \cap \Pi) \oplus \mathfrak{p}_{-1} \oplus [\mathfrak{p}_{-1}, \mathfrak{p}_{-1}] \oplus [[\mathfrak{p}_{-1}, \mathfrak{p}_{-1}], \mathfrak{p}_{-1}] \oplus \dots,$$

where $\mathfrak{p}_{-1} = 0$ if $\alpha_0 \notin S$ and $\mathfrak{p}_{-1} = \mathfrak{U}(\mathfrak{p}(S \cap \Pi))e_{-\alpha_0}$ otherwise. The opposite parabolic \mathfrak{p}^- is defined by $\mathfrak{p}^- = \omega(\mathfrak{p})$, where ω is an involution of $\mathfrak{g}[t, t^{-1}]$ such that $\omega|_{\mathfrak{t}} = -\mathrm{id}_{\mathfrak{t}}$ and $\omega(\mathfrak{g}t^k) = \mathfrak{g}t^{-k}$ for all $k \in \mathbb{Z}$.

If $S = \tilde{\Pi}$, then $\mathfrak{p}(S) = \mathfrak{g}[t, t^{-1}]$. Suppose that $\alpha_0 \in S$ and $|S| \leq r$. Let \mathfrak{n} be the nilpotent radical of $\mathfrak{p}(S \cap \Pi)$. Then $\mathfrak{p}_{-1} = \mathfrak{z}(\mathfrak{n})t^{-1}$, where $\mathfrak{z}(\mathfrak{n})$ is the centre of \mathfrak{n} . Hence $[\mathfrak{p}_{-1}, \mathfrak{p}_{-1}] = 0$.

Having two standard parabolics \mathfrak{p} and \mathfrak{r} , we define an *affine seaweed subalgebra* $\mathfrak{q} = \mathfrak{p} \cap \mathfrak{r}^-$. An interesting situation occurs if \mathfrak{q} is finite-dimensional. This is the case if and only if both $\mathfrak{p}, \mathfrak{r}$ are smaller than $\mathfrak{g}[t, t^{-1}]$.

In a slightly more general setting of affine Kac–Moody algebras, a combinatorial formula for the index of a finite-dimensional affine seaweed subalgebra is obtained in [12]. We are interested in very particular examples of contact seaweeds.

Suppose that \mathfrak{g} is of type A_r . Then the extended Dynkin diagram of \mathfrak{sl}_{r+1} is a circle. We consider $\mathfrak{q} = \mathfrak{p} \cap \mathfrak{r}^-$, where $\mathfrak{p} = \mathfrak{p}(S)$, $\mathfrak{r} = \mathfrak{p}(T)$, and $S, T \subset \tilde{\Pi}$ are proper subsets. Therefore it is safe to assume that $\alpha_0 \notin T$. If also $\alpha_0 \notin S$, then \mathfrak{q} is a seaweed subalgebra of \mathfrak{g} . Such subalgebras have been discussed already.

4.1. Intersections of two maximal parabolics. We will treat one instance, namely, where $\mathfrak{g} = \mathfrak{sl}_{r+1}$, $T = \Pi$, |S| = r, and $\alpha_0 \in S$. Here $\mathfrak{q}(S, T)$ is a semi-direct product

$$\mathfrak{q}(S,T) \cong \mathfrak{s}(\mathfrak{gl}_a \oplus \mathfrak{gl}_b) \ltimes 2\Bbbk^a \otimes \Bbbk^b,$$

where a + b = r + 1, the subspace $2\Bbbk^a \otimes \Bbbk^b$ is an Abelian ideal, and the 1-dimensional centre of the Levi subalgebra $\mathfrak{s}(\mathfrak{gl}_a \oplus \mathfrak{gl}_b)$ acts on both copies of $\Bbbk^a \otimes \Bbbk^b$ with one and the same non-trivial character.

First we compute the index of \mathfrak{q} . Here a formula of Raïs [22] will be frequently used. Let $H = L \ltimes \exp(V)$ be a semi-direct product of a group L and an Abelian unipotent group $\exp(V)$. Suppose that L acts on V^* with an open orbit. Then

(4.1)
$$\operatorname{ind} \operatorname{Lie} H = \operatorname{ind} \operatorname{Lie} L_{\gamma}$$

for a generic point $\gamma \in V^*$, see [22].

It is convenient to extend q by a 1-dimensional central toral subalgebra and to consider another semi-direct product, *R*. Set

$$Q = Q(a, b) = (\mathrm{GL}_a \times \mathrm{GL}_b) \ltimes \exp(2\Bbbk^a \otimes \Bbbk^b)$$

and also

$$R = R(a, b) = (\mathrm{GL}_a \times \mathrm{GL}_b) \ltimes \exp((\mathfrak{gl}_a^{\mathsf{ab}} \oplus W) \oplus V),$$

where $[\mathfrak{gl}_a^{ab}, \mathfrak{gl}_a^{ab}] = [W, W] = 0$, $W \cong V \cong \Bbbk^a \otimes \Bbbk^b$, and $[\mathfrak{gl}_a^{ab}, W] = V$. In both cases, GL_a and GL_b act on \Bbbk^a and \Bbbk^b in the natural way, also GL_a acts on \mathfrak{gl}_a via the adjoint representation. Note that Q(a, b) = Q(b, a), but in the case of R(a, b), the numbers a and b play different roles. Set $\mathfrak{q} = \mathfrak{q}(a, b) = \operatorname{Lie} Q(a, b)$ and $\mathfrak{r} = \mathfrak{r}(a, b) = \operatorname{Lie} R(a, b)$. Assume that $\mathfrak{q} \neq 0$ and $\mathfrak{r} \neq 0$.

Theorem 4.1. We have ind $q = \gcd(2a, a + b)$ and ind $\mathfrak{r} = \gcd(2a, b)$.

Proof. We argue by induction on a + b dealing with both statements simultaneously. If a = 0, then $Q(0, b) = R(0, b) = GL_b$ and $ind \mathfrak{q} = b = gcd(0, b)$.

If b = 0, then $R(a, 0) = \operatorname{GL}_a \ltimes \exp(\mathfrak{gl}_a^{\mathsf{ab}})$ is a *Takiff* Lie algebra (truncated current algebra) and $\operatorname{ind} \mathfrak{r} = 2a = \operatorname{gcd}(2a, 0)$ [26]. In particular, both statements hold, if a + b = 1. Now suppose that ab > 0.

Set $H = \operatorname{GL}_a \times \operatorname{GL}_b$. Consider first Q. Assume that $a \leq b$. If $2a \leq b$, then H acts on $(2\Bbbk^a \otimes \Bbbk^b)^*$ with an open orbit, say $H\xi$. Furthermore,

$$H_{\xi} = (\mathrm{GL}_a \times \mathrm{GL}_{b-2a}) \ltimes \exp(2\Bbbk^a \otimes \Bbbk^{b-2a}) = Q(a, b-2a).$$

By (4.1) and induction, $\operatorname{ind} \mathfrak{q} = \operatorname{ind} \operatorname{Lie} H_{\xi} = \operatorname{gcd}(2a, b - a) = \operatorname{gcd}(2a, a + b).$

Keep the assumption $a \leq b$. We can decompose Q as $Q = (H \ltimes \exp(V_1)) \ltimes \exp(V_2)$ with $V_1 \cong V_2 \cong \Bbbk^a \otimes \Bbbk^b$. Then $L = H \ltimes \exp(V_1)$ acts on V_2^* with an open orbit, say $L\gamma$. Therefore ind $\mathfrak{q} = \operatorname{ind} \operatorname{Lie} L_{\gamma}$ by (4.1). We have $L_{\gamma} = H_{\gamma} \ltimes \exp(V_1)$. By a direct calculation,

$$L_{\gamma} = \mathrm{GL}_{a} \times \mathrm{GL}_{b-a} \ltimes \exp((\mathfrak{gl}_{a}^{\mathsf{ab}} \oplus \widehat{W}) \oplus \widehat{V})$$

with $[\mathfrak{gl}_a^{\mathsf{ab}}, \mathfrak{gl}_a^{\mathsf{ab}}] = [\tilde{W}, \tilde{W}] = 0$, $[\mathfrak{gl}_a^{\mathsf{ab}}, \tilde{W}] = \tilde{V}$, and $\tilde{W} \cong \tilde{V} \cong \Bbbk^a \otimes \Bbbk^{b-a}$, i.e., $L_{\gamma} = R(a, b-a)$. In case $2a \leq b$, we can conclude that

$$\operatorname{ind} \mathfrak{r}(a, b-a) = \operatorname{ind} \mathfrak{q}(a, b) = \operatorname{ind} \mathfrak{q}(a, b-2a) = \operatorname{gcd}(2a, b-a).$$

Therefore it remains to perform the induction step for $\mathfrak{r}(a, b)$ in the case, where 2a > a + b, i.e., for a > b.

Consider now $\mathfrak{r} = \mathfrak{r}(a, b)$ with a > b. Let $\eta \in V^*$ be a generic point. We regard it is an element of \mathfrak{r}^* . There is a more suitable decomposition of \mathfrak{r} , namely, $\mathfrak{r} = \tilde{\mathfrak{l}} \ltimes (W \oplus V)$, where $\tilde{\mathfrak{l}} = \operatorname{Lie} \tilde{L}$ with $\tilde{L} = H \ltimes \exp(\mathfrak{gl}_a^{\mathsf{ab}})$. Note that $[W \oplus V, W \oplus V] = 0$. The inequality a > b implies that $\operatorname{ad}^*(\mathfrak{gl}_a^{\mathsf{ab}}) \cdot \eta = W^*$ and that $\tilde{L}\eta$ is open in $(W \oplus V)^*$. Therefore $\operatorname{ind} \mathfrak{r} = \operatorname{ind} \tilde{\mathfrak{l}}_\eta$ by (4·1). By a straightforward computation, $\tilde{\mathfrak{l}}_\eta = \mathfrak{r}(a - b, b)$. Since b > 0, we have (a - b) + b < a + b and $\operatorname{ind} \tilde{\mathfrak{l}}_\eta = \operatorname{gcd}(2(a - b), b) = \operatorname{gcd}(2a, b)$ by the inductive hypothesis. \Box

If gcd(2a, a + b) = 2, then removing the centre of q(a, b) we obtain an affine seaweed of index 1. A natural question is whether it is contact or not.

4.2. Contact semi-direct products. Consider a semi-direct product $Q = L \ltimes \exp(V)$, where $\exp(V)$ is a normal Abelian unipotent subgroup. Suppose that $\operatorname{ind} \mathfrak{q} = 1$. Then by [22] there are two possibilities. Either (A): $\dim L\gamma = \dim V - 1$ and $\operatorname{ind} \operatorname{Lie} L_{\gamma} = 0$ or (B): $\dim L\gamma = \dim V$ and $\operatorname{ind} \operatorname{Lie} L_{\gamma} = 1$ for a generic point $\gamma \in V^*$. We want to check, whether \mathfrak{q} is contact or not. Regard γ as a function on \mathfrak{q} such that $\gamma(\mathfrak{l}) = 0$ for $\mathfrak{l} = \operatorname{Lie} L$.

In case (A), $Q\gamma = l^* \times L\gamma$. Therefore q is contact if and only if a generic *L*-orbit on V^* is not conical. Both instances, conical or not, take places, cf. Example 3.13.

The situation (B) is more interesting. Suppose that we are in it. A reduction from Q to L_{γ} works well for the index computation. The property of an orbit to be conical is more subtle. Let $L_{\langle\gamma\rangle}$ be the normaliser of the line $\Bbbk\gamma$ and set $\mathfrak{l}_{\langle\gamma\rangle} = \operatorname{Lie} L_{\langle\gamma\rangle}$. We identify \mathfrak{l}^* with $\operatorname{Ann}(V) \subset \mathfrak{q}^*$.

Theorem 4.2. Let $L\gamma \subset V^*$ be the open orbit.

(i) Suppose that l_{γ} is not contact. Then q is contact if and only if $\operatorname{ind} l_{\langle\gamma\rangle} = 0$. (ii) If l_{γ} is contact, but q is not contact, then $\operatorname{ind} l_{\langle\gamma\rangle} = 0$.

Proof. In each generic Q orbit in \mathfrak{q}^* , there is a point α such that $\alpha|_V = \gamma$. Set $\beta = \alpha|_{\mathfrak{l}_{\gamma}}$. Without loss of generality assume that β is a generic point of \mathfrak{l}^*_{γ} . For case (ii), assume that β is contact. Since $\overline{L\gamma} = V$, there is $x \in \mathfrak{l}$ such that $\mathrm{ad}^*(x)(\gamma) = \gamma$. Clearly $\mathfrak{l}_{\langle\gamma\rangle} = \Bbbk x \oplus \mathfrak{l}_{\gamma}$ and $[x, \mathfrak{l}_{\gamma}] \subset \mathfrak{l}_{\gamma}$. We say that x acts on \mathfrak{l}_{γ} and on \mathfrak{l}^*_{γ} .

Note that by dimension reasons, $\operatorname{ad}^*(V) \cdot \gamma = \operatorname{Ann}(\mathfrak{l}_{\gamma}) \subset \mathfrak{l}^*$. Therefore $Q\alpha$ is conical if and only if $x \cdot \beta \in \beta + \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$.

(i) Since \mathfrak{l}_{γ} is not contact, $L_{\gamma}\beta \subset \mathfrak{l}_{\gamma}^{*}$ is a conical orbit, see Proposition 2.1. Thus, $Q\alpha$ is conical if and only if $x \cdot \beta \in \mathrm{ad}^{*}(\mathfrak{l}_{\gamma}) \cdot \beta$.

If $x \cdot \beta \notin \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$, then $\dim L_{\langle \gamma \rangle} \beta \geq 1 + \dim L\beta$. Since each coadjoint orbit is evendimensional, $\operatorname{ind} \mathfrak{l}_{\langle \gamma \rangle} \leq (1 + \dim \mathfrak{l}) - (2 + \dim L\beta) \leq \operatorname{ind} \mathfrak{l} - 1 = 0$.

Suppose that $x \cdot \beta \in \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$. Let $\tilde{\beta} \in \mathfrak{l}^*_{<\gamma>}$ be an extension of β . Since [x, x] = 0, we may regard $\operatorname{ad}^*(\mathfrak{l}_{<\gamma>}) \cdot \tilde{\beta}$ is a subspace of \mathfrak{l}^*_{γ} and then it is equal to $x \cdot \beta + \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta = \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$. Thus, $\operatorname{ind} \mathfrak{l}_{<\gamma>} = (1 + \dim \mathfrak{l}) - \dim L_{\gamma}\beta = 2$.

(ii) Since q is not contact, $Q\alpha$ is conical by Proposition 2.1. Hence $x \cdot \beta \in \beta + \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$. If $x \cdot \beta \in \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$, then also $\beta \in \operatorname{ad}^*(\mathfrak{l}_{\gamma}) \cdot \beta$, which is a contradiction. Thereby again $\dim L_{<\gamma>\beta} \geq 1 + \dim L\beta$.

Following [14], we say that $\mathfrak{q} = \operatorname{Lie} Q$ is *strongly quasi-reductive*, if \mathfrak{q} is quasi-reductive and the centre $Z \subset Q$ consists of semisimple elements, in other words, if Q_{β} is reductive for some $\beta \in \mathfrak{q}^*$. In certain cases, this definition depends on the group Q. Although $\operatorname{Lie}(\mathbb{k}, +) \cong \operatorname{Lie}(\mathbb{k}^{\times}, \times)$, the first one is not strongly quasi-reductive, while the second is. Recall from Section 2.1 that if Q_{β} is reductive for one β , then Q_{α} is reductive for a generic point $\alpha \in \mathfrak{q}^*$.

Proposition 4.3 (cf. [18, Corollary 2.9.], [14, Lemma 4.3]). Let $Q = L \ltimes \exp(V)$ be such that $L\gamma \subset V^*$ is an open orbit. Then q is strongly quasi-reductive if and only if $l_{\gamma} = \text{Lie } L_{\gamma}$ is strongly quasi-reductive.

Proof. In each generic Q orbit in \mathfrak{q}^* , there is a point α such that $\alpha|_V = \gamma$. Clearly $Q_\alpha \subset L_\gamma \ltimes \exp(V)$. We have

$$\exp(V)\alpha = \alpha + \operatorname{ad}^*(V)\cdot\gamma, \text{ where } \operatorname{ad}^*(V)\cdot\gamma|_V = 0.$$

Furthermore, $\operatorname{ad}^*(V) \cdot \gamma = \operatorname{Ann}(\mathfrak{l}_{\gamma}) \subset \mathfrak{l}^*$ is of dimension $\dim V$ and $\operatorname{ad}^*(v)(\gamma) \neq 0$ for each non-zero $v \in V$. Hence $Q_{\alpha} = Q_{\alpha}/(Q_{\alpha} \cap \exp(V)) \cong (L_{\gamma})_{\beta}$ for the restriction $\beta = \alpha|_{\mathfrak{l}}$. Since β is a generic point of \mathfrak{l}^* , we are done. \Box

Recall that a (strongly) quasi-reductive Lie algebra of index 1 is contact, see Section 2.1. Therefore Theorrem 4.2 and Proposition 4.3 can be employed for obtaining a classification of contact semi-direct products $L \ltimes \exp(V)$. Similar results have been obtained before. The semi-direct products $L \ltimes \exp(V)$ of index zero, where *L* is reductive, are classified in [8] under the assumption that *V* is a simple *L*-module or the commutator group of *L* is simple.

Now we return to affine seaweeds. Recall that we have obtained q(a, b) as a central extension of a standard affine seaweed q(S, T). Therefore set $\bar{q} = \bar{q}(a, b) = q(S, T)$. We have

$$\operatorname{ind} \bar{q} = \operatorname{ind} \mathfrak{q}(a, b) - 1 = \operatorname{gcd}(2a, a + b) - 1,$$

see Theorem 4.1. If $\operatorname{ind} \overline{\mathfrak{q}} = 1$, then either $a \in 2\mathbb{Z}$ and $a + b \in 2 + 4\mathbb{Z}$ or both a and b are odd.

Let $\overline{R}(a, b)$ be the quotient of R(a, b) by the central toral subgroup.

The centre of $\bar{\mathfrak{q}}(a,b)$ is trivial. Thereby $\bar{\mathfrak{q}}(a,b)$ is strongly quasi-reductive if and only if it is quasi-reductive. If $b \neq 0$, then the centre of $\bar{\mathfrak{r}}(a,b)$ is trivial. In case of $\bar{R}(a,0) =$ $\operatorname{SL}_a \ltimes \exp(\mathfrak{gl}_a^{\operatorname{ab}})$ with $a \neq 0$, the centre is 1-dimensional and consists of unipotent elements. The corresponding Lie algebra $\bar{\mathfrak{r}}(a,b) := \operatorname{Lie} \bar{R}(a,b)$ is not quasi-reductive for a > 1 and is quasi-reductive, but not strongly quasi-reductive, if a = 1.

Theorem 4.4. If *a* and *b* are even and $\operatorname{ind} \bar{\mathfrak{q}}(a, b) = \operatorname{ind} \bar{\mathfrak{r}}(a, b) = 1$, then $\bar{\mathfrak{q}}(a, b)$ and $\bar{\mathfrak{r}}(a, b)$ are strongly quasi-reductive and hence contact.

Proof. We argue by induction on a + b. The case a + b = 0 does not take place. Suppose that a + b = 2. Then either a or b is equal to zero. Note that $\bar{\mathfrak{q}}(0,2) = \bar{\mathfrak{q}}(2,0) = \mathfrak{sl}_2$ is

reductive. Since $\operatorname{ind} \overline{\mathfrak{r}}(a, b) = 1$, the only possibility for $\overline{\mathfrak{r}}(a, b)$ is $\overline{\mathfrak{r}}(0, 2) = \mathfrak{sl}_2$. Now we have the induction base and assume that a + b > 2. This assumption implies that ab > 0.

Induction step for \bar{Q} . The group $\bar{Q}(a, b)$ is a semi-direct product of $L = S(GL_a \times GL_b) \ltimes \exp(\mathbb{k}^a \otimes \mathbb{k}^b)$ and $\exp(V)$ for $V = \mathbb{k}^a \otimes \mathbb{k}^b$. As was mentioned in the proof of Theorem 4.1, L acts on V^* with an open orbit $L\gamma$, where $L_{\gamma} = \bar{R}(a, b - a)$. Since a > 0, we have a + (b - a) < a + b.

Because *a* is even, $\bar{\mathfrak{r}}(a, b-a)$ is strongly quasi-reductive and hence so is $\bar{q}(a, b)$ by Proposition 4.3.

Induction step for \overline{R} . The group $\overline{R}(a, b)$ has two different decompositions into semidirect products $L \ltimes \exp(V)$. Suppose $a \leq b$. Then $L = S(\operatorname{GL}_a \times \operatorname{GL}_b) \ltimes \exp(W)$ and $V = \mathfrak{gl}_a^{\mathsf{ab}} \oplus V_0$, where $V_0 \cong W \cong \Bbbk^a \otimes \Bbbk^b$ and $[W, \mathfrak{gl}_a^{\mathsf{ab}}] = V_0$. If a > b, then

$$L = \mathrm{S}(\mathrm{GL}_a \times \mathrm{GL}_b) \ltimes \exp(\mathfrak{gl}_a^{\mathsf{ab}})$$

and $V = W \oplus V_0$. In both cases, *L* acts on V^* with an open orbit and a generic point $\gamma \in V^*$ can be chosen in V_0^* .

If
$$a \leq b$$
, then $L_{\gamma} = \overline{Q}(a, b - a)$;
if $a > b$, then $L_{\gamma} = \overline{R}(a - b, b)$.

Since *a* is even, we conclude that $\mathfrak{r}(a, b)$ is strongly quasi-reductive using Proposition 4.3 and the inductive hypothesis.

Keep the assumption $\operatorname{ind} \overline{\mathfrak{q}}(a, b) = \operatorname{ind} \overline{\mathfrak{r}}(a, b) = 1$. If *a* is odd, then neither $\overline{\mathfrak{q}}(a, b)$ nor $\overline{\mathfrak{r}}(a, b)$ is strongly quasi-reductive, since the above reductions end with $\overline{\mathfrak{r}}(1, 0)$. At each reduction step $\overline{\mathfrak{q}}(a, b) \rightsquigarrow \overline{\mathfrak{r}}(a, b-a)$, $\overline{\mathfrak{r}}(a, b) \rightsquigarrow \overline{\mathfrak{q}}(a, b-a)$ or $\overline{\mathfrak{r}}(a, b) \rightsquigarrow \overline{\mathfrak{r}}(a-b, b)$, the normaliser $\mathfrak{l}_{<\gamma>}$ is Frobenius. This makes inductive arguments difficult.

Example 4.5. The Lie algebra $\bar{\mathfrak{q}}(1,1) = \Bbbk \ltimes 2\Bbbk$ is not contact. We can decompose any $\bar{Q}(1,b)$ with an odd b as $L \ltimes \exp(V)$, where $L = \operatorname{GL}_b$ and $V = 2\Bbbk^b$. Then $L_{<\gamma>} = Q(1,b-2)$ for $\gamma \in V^*$ belonging to the open L-orbit. The Lie algebra of this normaliser is of index 2, see Theorem 4.1. Thereby we can use Theorem 4.2 and conclude by induction on b that $\bar{\mathfrak{q}}(1,b)$ is never contact.

Consider the chain of reductions $\bar{\mathfrak{q}}(1,3) \rightsquigarrow \bar{\mathfrak{r}}(1,2) \rightsquigarrow \bar{\mathfrak{q}}(1,1)$, where the first and the last algebras are not contact. By Theorem 4.2(i), $\bar{\mathfrak{r}}(1,2)$ is contact.

Consider another chain of reductions

$$\overline{\mathfrak{q}}(3,5) \rightsquigarrow \overline{\mathfrak{r}}(3,2) \rightsquigarrow \overline{\mathfrak{r}}(1,2) \rightsquigarrow \overline{\mathfrak{q}}(1,1).$$

Since $\bar{\mathfrak{r}}(1,2)$ is contact, the previous item, $\bar{\mathfrak{r}}(3,2)$, may be contact or not. In order to decide this, one needs a better understanding of the action of x on \mathfrak{l}^*_{γ} used in the proof of Theorem 4.2. The same applies to $\bar{\mathfrak{q}}(3,5)$. We suspect that $\bar{\mathfrak{q}}(a,b)$ with odd a and b is never contact. The matter deserves further investigations.

REFERENCES

- K. AMMARI, Stabilité des sous-algèbres biparaboliques des algèbres de Lie simples, J. Lie Theory, 32, no. 1 (2022), 239–260.
- [2] W. BORHO, P. GABRIEL, and R. RENTSCHLER, "Primideale in einhüllenden auflösbarer Lie-Algebren", Lecture Notes in Math., Bd. 357, Springer-Verlag, 1973.
- [3] J.-Y. CHARBONNEL and A. MOREAU, The symmetric invariants of centralizers and Slodowy grading, *Math. Z.*, **282** (2016), no. 1-2, 273–339.
- [4] V. E. COLL JR. and N. RUSSONIELLO, Classification of contact seaweeds, J. Algebra, 659 (2024), 811–817.
- [5] V. DERGACHEV and A.A. KIRILLOV, Index of Lie algebras of seaweed type, J. Lie Theory, **10** (2000), 331–343.
- [6] ANDRÉ DIATTA, Left invariant contact structures on Lie groups, Diff. Geometry and its Applications, 26 (2008), 544–552.
- [7] M. DUFLO, M.S. KHALGUI, and P. TORASSO, Algèbres de Lie quasi-réductives, *Transform. Groups*, 17 (2012), 417–470.
- [8] A. G. ELASHVILI, Frobenius Lie algebras, Funct. Anal. Appl., 16, no. 4 (1982), 326–328.
- [9] H. GEIGES, An Introduction to Contact Topology, Cambridge Studies in Advanced Mathematics, **109**, Cambridge University Press 2008.
- [10] M. GROMOV, Stable maps of foliations in manifolds, Izv. Akad. Nauk SSSR, 33 (1969), 707–734.
- [11] A. JOSEPH, On semi-invariants and index for biparabolic (seaweed) algebras, I, J. Algebra, **305** (2006), 487–515.
- [12] A. JOSEPH, On semi-invariants and index for biparabolic (seaweed) algebras II, J. Algebra, 312 (2007), 158–193.
- [13] A. JOSEPH and D. SHAFRIR, Polynomiality of invariants, unimodularity and adapted pairs, *Transform. Groups*, **15**, no. 4 (2010), 851–882.
- [14] A. MOREAU and O. YAKIMOVA, Coadjoint orbits of reductive type of parabolic and seaweed Lie subalgebras, *Int. Math. Res. Notices*, **2012**, no. 19 (2012), 4475–4519.
- [15] A. I. OOMS, On Frobenius Lie algebras, Comm. Algebra, 8, no. 1 (1980), 13–52.
- [16] A. OOMS and M. VAN DEN BERGH, A degree inequality for Lie algebras with a regular Poisson semicenter, J. Algebra, **323**, no. 2 (2010), 305–322.
- [17] D. PANYUSHEV, Inductive formulas for the index of seaweed Lie algebras, *Moscow Math. J.*, **1**(2001), 221–241.
- [18] D. PANYUSHEV, An extension of Raïs' theorem and seaweed subalgebras of simple Lie algebras, Ann. Inst. Fourier (Grenoble), 55 (2005), no. 3, 693–715.
- [19] D. PANYUSHEV, Periodic automorphisms of Takiff algebras, contractions, and θ-groups, *Transformation Groups*, 14, no. 2 (2009), 463–482.
- [20] D. PANYUSHEV and O. YAKIMOVA, On seaweed subalgebras and meander graphs in type C, *Pacific J. Math.*, 285, no. 2 (2016), 485–499.

O. YAKIMOVA

- [21] D. PANYUSHEV and O. YAKIMOVA, Takiff algebras with polynomial rings of symmetric invariants, *Transformation Groups*, **25** (2020), 609–624.
- [22] M. RAÏS, L'indice des produits semi-directs $E \times_{\rho} \mathfrak{g}$, C. R. Acad. Sci. Paris Ser. A, **287** (1978), 195–197.
- [23] G. SALGADO-GONZÁLEZ, Invariants of contact Lie algebras, J. Geometry and Physics, 144 (2019), 388– 396.
- [24] M. SATO and T. KIMURA, A classification of irreducible prehomogeneous vector spaces and their relative invariants, *Nagoya Math. J.*, 65 (1977), 1–155.
- [25] T. A. SPRINGER, Aktionen reduktiver Gruppen auf Varietäten, in: "Algebraische Transformationsgruppen und Invariantentheorie", DMV-Seminar, Bd. 13, Basel–Boston–Berlin: Birkhäuser 1989, pp. 3– 39.
- [26] S. J. TAKIFF, Rings of invariant polynomials for a class of Lie algebras, *Trans. Amer. Math. Soc.* 160 (1971), 249–262.
- [27] P. TAUVEL and R. W. T. YU, Indice et formes linéaires stables dans les algèbres de Lie, J. Algebra, 273 (2004), 507–516.
- [28] O. YAKIMOVA, Some semi-direct products with free algebras of symmetric invariants, F. Callegaro et al. (eds.), *Perspectives in Lie Theory*, Springer INdAM Series **19** (2017), 267–279.

(O. Yakimova) Institut für Mathematik, Friedrich-Schiller-Universität Jena, Jena, 07737, Deutschland

Email address: oksana.yakimova@uni-jena.de