

The Pristine Inner Galaxy Survey (PIGS)

XI: Revealing the chemical evolution of the interacting Sagittarius dwarf galaxy

Sara Vitali^{1,2*}, Alvaro Rojas-Arriagada^{4,2,3,5}, Paula Jofré^{1,2}, Federico Sestito⁶, Joshua Povick^{1,2}, Vanessa Hill⁷, Emma Fernández-Alvar^{8,9}, Anke Ardern-Arentsen¹⁰, Pascale Jablonka¹¹, Nicolas F. Martin^{12,13}, Else Starkenburg¹⁴, and David Aguado^{8,9}

¹ Instituto de Estudios Astrofísicos, Facultad de Ingeniería y Ciencias, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile, e-mail: sara.vitali@mail.udp.cl

² Millenium Nucleus ERIS

³ Millenium Institute of Astrophysics, Av. Vicuña Mackenna 4860, 82-0436 Macul, Santiago, Chile

⁴ Departamento de Física, Universidad de Santiago de Chile, Av. Victor Jara 3659, Santiago, Chile

⁵ Center for Interdisciplinary Research in Astrophysics and Space Exploration (CIRAS), Universidad de Santiago de Chile, Santiago, Chile

⁶ Centre for Astrophysics Research, Department of Physics, Astronomy and Mathematics, University of Hertfordshire, Hatfield, AL10 9AB, UK

⁷ Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, Nice, France

⁸ Instituto de Astrofísica de Canarias (IAC), Vía Láctea, 38205 La Laguna, Tenerife, Spain

⁹ Universidad de La Laguna, Dept. Astrofísica, 38206 La Laguna, Tenerife, Spain

¹⁰ Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

¹¹ Laboratoire d'astrophysique, École Polytechnique Fédérale de Lausanne (EPFL), CH-1290 Versoix, Switzerland

¹² Université de Strasbourg, CNRS, Observatoire astronomique de Strasbourg, UMR 7550, F-67000 Strasbourg, France

¹³ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

¹⁴ Kapteyn Astronomical Institute, University of Groningen, Landleven 12, 9747 AD Groningen, The Netherlands

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Context. The Sagittarius dwarf spheroidal galaxy (Sgr dSph) is a satellite orbiting the Milky Way that has experienced multiple stripping events due to tidal interactions with our Galaxy. Its accretion history has led to a distinct stellar overdensity, which is the remnant of the core of the progenitor.

Aims. We present a complete chemical analysis of 111 giant stars in the core of Sgr dSph to investigate the chemical evolution and enrichment history of this satellite.

Methods. Employing the metallicity-sensitive Ca H&K photometry from the Pristine Inner Galaxy Survey, we selected stars spanning a wide metallicity range and obtained high-resolution spectra with the ESO FLAMES/GIRAFFE multi-object spectrograph. For the stellar sample covering $-2.13 < [\text{Fe}/\text{H}] < -0.35$, we derived abundances for up to 14 chemical elements with average uncertainties of ~ 0.09 dex and a set of stellar ages which allowed us to build an age-metallicity relation (AMR) for the entire sample.

Results. With the most comprehensive set of chemical species measured for the core of Sgr (Na, Mg, Al, Si, Ca, Sc, Ti, V, Cr, Co, Ba, La and Eu), we studied several $[\text{X}/\text{Fe}]$ ratios. Most trends align closely with Galactic chemical trends, but notable differences emerge in the heavy n -capture elements, which offer independent insights into the star formation history of a stellar population.

Conclusions. The deficiency in the α -elements with respect the Milky Way suggests a slower, less efficient early star formation history, similar to other massive satellites. S -process element patterns indicate significant enrichment from AGB stars over time. The AMR and chemical ratios point to an extended star formation history, with a rapid early phase in the first Gyr, followed by declining activity and later star-forming episodes. These findings are consistent with Sgr hosting multiple stellar populations, from young (~ 4 Gyr) to old, metal-poor stars (~ 10 Gyr).

Key words. Galaxies: individual: Sagittarius dwarf galaxy - Galaxies: dwarf - Galaxies: abundances - Stars: abundances - Stars: Population II

1. Introduction

The Lambda Cold Dark Matter (Λ CDM) model provides the most up-to-date understanding of galaxy formation, suggesting that galaxies form through the merging and

* analysis of data taken as part of ESO program ID 109.234B.001

subsequent accretion of smaller systems. This hierarchical model posits that galaxies enclose dust, gas, and stars, all embedded within massive dark matter halos (Searle & Zinn 1978; White & Rees 1978; White & Frenk 1991). In this context, the Milky Way (MW) serves as an ideal laboratory, as its spatial resolution allows for the study of individual stars and the acquisition of detailed information to explore its star formation history. At smaller scales, the accretion events that built up our Galaxy can be better understood by studying the surrounding dwarf galaxies which comprise the majority of the MW's satellite populations (e.g., Venn et al. 2004; Tolstoy et al. 2009; Bland-Hawthorn & Gerhard 2016).

Of such galaxies, one of the biggest and most luminous known around the MW is the Sagittarius (Sgr) dwarf spheroidal (dSph) galaxy, discovered by Ibata et al. (1994). With an estimated total mass of $\sim 4 \times 10^8$ (McConnachie 2012; Vasiliev & Belokurov 2020) and a luminosity of $M \sim -13.5$ (McConnachie 2012), Sgr is an ideal "test case" for investigating chemical evolution theories within the framework of hierarchical accretion and learn about the early evolution of the MW. In addition, Sgr is currently undergoing a merger with the MW, having experienced its first infall approximately 5 Gyr ago (Ruiz-Lara et al. 2020). The long tidal interaction with our Galaxy has profoundly shaped Sgr, leaving behind a remnant core located on the opposite side of the Galactic center at a heliocentric distance $d \sim 26.5$ kpc (Ibata et al. 1994; Majewski et al. 2003; Law & Majewski 2010; Vasiliev & Belokurov 2020), along with two wide stellar streams.

Numerous studies, both photometric (e.g., Bellazzini et al. 1999a; Layden & Sarajedini 2000; Siegel et al. 2007; Vitali et al. 2022) and spectroscopic (e.g., Bonifacio et al. 2000; Monaco et al. 2005; McWilliam et al. 2013; Hasselquist et al. 2017; Hansen et al. 2018; Hasselquist et al. 2021; Sestito et al. 2024b,a), targeting the Sgr core in recent years, have revealed an extended star formation history (SFH) heavily influenced by gravitational interactions with the MW. During Sgr's orbital passages around the MW, multiple star formation episodes occurred, leading to the formation of distinct stellar populations: a predominant intermediate-age population (4-8 Gyr, $-0.6 \lesssim [\text{Fe}/\text{H}] \lesssim -0.4$), a young population (~ 2.5 Gyr and $[\text{Fe}/\text{H}] \sim 0$), and a very young population of a few Myr with super-solar metallicities (age < 2 Gyr and $[\text{Fe}/\text{H}] \sim +0.5$). These populations are hosted in the core of Sgr alongside the older, more metal-poor population (age > 10 Gyr, $[\text{Fe}/\text{H}] \lesssim -1.0$). Part of the most metal-poor population has been gradually stripped away from the core of Sgr during its history of interaction with the MW and deposited in the two long stellar streams that wrap around it (Ibata et al. 2001; Majewski et al. 2003). These streams are known to be ~ 1 dex more metal-poor than the core (de Boer et al. 2015).

Despite a fair understanding of the evolution of this dissolving system, its SFH and how its evolution has been affected by interactions with the MW remain open questions. To address this topic, the exquisite data releases of *Gaia* (Gaia Collaboration et al. 2016a, 2021, 2023), which have collected astrometric and photometric data for 1.8 billion sources in the MW and its satellites, can be efficiently combined with photometric surveys such as *Pristine* (Starkenburg et al. 2017; Martin et al. 2023), which is specifically dedicated to tracing the history of the MW and its satellites through the study of metal-poor stars. This

latter survey, primarily focused on the MW halo, has developed a particular focus on the inner Galactic region thanks to the *Pristine* Inner Galaxy Survey (PIGS, Arentsen et al. 2020b; Ardern-Arentsen et al. 2024). Both surveys rely on a narrow-band filter centered around the CaH&K region to identify metal-poor candidates. Their efficiency has led to numerous studies focused on both our Galaxy and the Sgr region (Arentsen et al. 2020a; Vitali et al. 2022; Sestito et al. 2024b,a), which is included in the coverage of PIGS. Using this dataset, Vitali et al. (2022) performed a photometric analysis and characterized the metallicity distribution of approximately 50,000 Sgr members, finding a negative photometric metallicity gradient extending up to 5.5 kpc along the main body of Sgr, indicative of an outside-in formation process. Sestito et al. (2024b) chemically characterized 12 very metal-poor (VMP) stars spectroscopically, providing new insights into the early evolution of Sgr by assembling the largest high-resolution study in the VMP regime and significantly expanding the existing sample of VMPs (Hansen et al. 2018; Chiti & Frebel 2019; Chiti et al. 2020). In a recent study, Sestito et al. (2024a) extended the investigation of the ancient chemical evolution of Sgr by analyzing carbon abundances, using low-/medium-resolution spectroscopic follow-up data from PIGS (Arentsen et al. 2020a; Ardern-Arentsen et al. 2024).

Despite these detailed analyses, a statistically robust characterization of Sgr's evolution needs to be extended to a broader metallicity range to encompass not only the early stages of this satellite. To achieve this, it is crucial to continue gathering precise and accurate measurements of various chemical species across a wider range of metallicities and for a larger sample of stars. For instance, constraining the trend of α -elements with metallicity allows for a deeper understanding of the initial mass function and the star formation history of a system (e.g., Tinsley 1968; McWilliam 1997). Simultaneously, obtaining additional measurements of heavier elements (atomic number $Z > 30$), which are produced in both the slow (s) and rapid (r) neutron-capture processes (e.g., Travaglio et al. 2004; Karakas & Lattanzio 2014; Bisterzo et al. 2014; Limongi & Chieffi 2018), is crucial for probing the astrophysical sites of these processes, unraveling their distinct chemical signatures, and exploring the timescales associated with the SFH of the galaxy. By adding high-resolution chemical abundances to the already existing dataset on Sgr (e.g., McWilliam et al. 2013; Monaco et al. 2005; Sbordone et al. 2007; Carretta et al. 2010; Hansen et al. 2018), this analysis enables comparative studies of chemical evolution between Sgr and other MW satellites, such as the Sculptor and Fornax dSphs, the Large Magellanic Cloud, and the accreted Gaia-Enceladus/Sausage system.

In this work, we present a sample of 111 Sgr giant stars for which we obtained high-resolution spectra thanks to observations performed with the ESO FLAMES/GIRAFFE multi-object spectrograph (Pasquini et al. 2002). Through a comprehensive spectroscopic analysis, we intend to study the evolution of the Sgr system and compare it with other massive MW satellites that have different masses and SFHs. Additionally, this spectroscopic sample represents, to our knowledge, the most comprehensive high-resolution dataset in terms of chemical elements available derived in homogeneous way. The target selection and data reduction are detailed in Section 2. Section 3 describes the spectroscopic analysis, while Section 4 discusses the main conclusions that




Fig. 1. Target selection. **Top:** Zoom in equatorial coordinates on the PIGS-Sgr photometric footprint after all the cuts (grey points are the Sgr candidate members). The black cross identifies the nuclear globular cluster M54, and the contour ellipse marks its spatial extent. The FLAMES targets are colored according to their radial velocities. **Bottom:** Density contour plot for the proper motion space (with mean values subtracted) for the Sgr-PIGS selection and the FLAMES targets.

can be drawn from the chemical abundance analysis regarding the chemical evolution of Sgr, comparing it with other known MW systems. The conclusions are summarized in the final section 5.

2. Data

2.1. Target selection

The targets are located in the fields observed by PIGS (Arentsen et al. 2020b), an extension of the *Pristine* survey. A specific region of PIGS is focused on the Sagittarius system, resulting in the collection of metallicity-sensitive CaH&K photometry for the remaining core of the galaxy up to the onset of the galaxy’s tidal stellar streams (Arentsen et al. 2020b; Vitali et al. 2022).

From these photometric measurements, the very metal-poor (VMP) candidates ($[\text{Fe}/\text{H}] < -2.0$ dex) identified by PIGS were followed up using the AAOmega+2dF facility mounted on the Anglo-Australian Telescope (AAT, Saunders et al. 2004; Lewis et al. 2002; Sharp et al. 2006). From the collected low-/medium-resolution spectra ($R \sim 1300 - 11\,000$), effective temperatures, surface gravities, metallicities, and carbon abundances were derived for $\approx 13\,000$ VMP




Fig. 2. Color-color diagram for the PIGS-Sgr sample. The four metallicity bins are identified with four different shades of gray and labeled according to their ranges. The FLAMES targets are overlaid as circles with black edges, colored by their spectroscopic metallicities.

stars (see Arentsen et al. 2020b; Ardern-Arentsen et al. 2024 for details). Four AAT pointings were dedicated to collecting spectra for about 400 members of the Sagittarius system (AAT-Sgr), which have already been used in various works (see for instance Vitali et al. 2022; Sestito et al. 2024b,a). Overall, these spectroscopic follow-up efforts have demonstrated the power of PIGS photometry as a metallicity predictor.

For this work the Sgr targets were selected from the PIGS photometric catalog, as there is not corresponding AAT spectroscopy. Their properties are displayed in Fig. 1. This central field (colored points) lies in the core of the galaxy but is far enough ($\sim 1^\circ$ in both RA and DEC) from the nuclear cluster M54/NGC 6715, marked by a black cross and an ellipse indicating its spatial extent, to exclude stars from the cluster in our selection. To include only Sgr members and exclude MW foreground stars (where the Sgr stars are shown as grey points in Fig. 1) from the PIGS photometric catalog (hereafter the PIGS-Sgr), we applied the membership criterion based on *Gaia* photometry and astrometry presented in Vitali et al. (2022) and Sestito et al. (2024a). In addition to the quality cuts on *Gaia* photometry and astrometry adopted in these previous studies, we used a cut on the parallax and parallax error ($|\varpi| \leq 2\epsilon_\varpi$) to remove nearby MW stars, as well as a cut of $b < -10^\circ$. We limited stars taking into account the change in average proper motions along the body of Sg within a radius of 0.6 mas yr^{-1} (Vasiliev & Belokurov 2020) as depicted on the bottom panel of Fig. 1. Finally, to ensure a high SNR for the spectra in the observations, the limiting magnitude was set to $G = 17$. As shown in section 3.1 the radial velocities of our targets fall within the typical range reported in the literature for Sgr, reinforcing the membership of these stars to the dwarf system.

Among the selected PIGS-Sgr members, we used the PIGS photometry to build a stellar sample with an extended metallicity range. This approach allowed us to include representatives of the various stellar populations within the system for a comprehensive chemical study. To do so, we relied on the same method featured in Vitali et al. (2022). This strategy combined the spectroscopic training sample from the main *Pristine* halo survey (Starkenburg et al. 2017) with the *Pristine* Ca H&K catalog (Martin

Filter	$\lambda_{\text{in}} - \lambda_{\text{fin}}$ (nm)	R	$\langle \text{SNR} \rangle$	Exp _{time} (h)
HR8	491.7 - 516.3	23500	40	6.67
HR11	559.7 - 584	29500	33	3.42
HR15N	647 - 679	19200	53	1.65

Table 1. Wavelength coverage, resolution, average signal-to-noise ratio and exposure time of the GIRAFFE setups used for this work. The signal-to-noise ratio is given per pixel (1 pixel = 0.005 nm).

et al. 2023) to derive iso-metallicity lines in the Ca H&K-*Gaia* color-color space. Consistent with the approach of Vitali et al. (2022), we also applied an extinction correction to the PIGS photometry.

Limiting the selection to giant stars ($\log g < 3.98$, $T_{\text{eff}} < 5700$ K and $-4.0 < [\text{Fe}/\text{H}] < -0.5$) we obtained the diagram shown in Fig. 2. The colored areas are delimited by the iso-metallicity lines, which divide the PIGS-Sgr color-color space into four different metallicity bins (different grey points): $[\text{Fe}/\text{H}] < -2.0$; $-2.0 < [\text{Fe}/\text{H}] < -1.5$; $-1.5 < [\text{Fe}/\text{H}] < -1.0$; and $-1.0 < [\text{Fe}/\text{H}] < -0.5$. As described in Vitali et al. (2022) an offset (~ 0.52 mag) was applied to the PIGS-Sgr photometry to align it with the training sample, which was calibrated against the main survey where the Ca H&K photometry has a constant offset. This offset arises because, at the time, the "main" *Pristine* fields were calibrated using a part of the Ca H&K-SDSS color-color space populated mainly by nearby dwarfs (Starkenburg et al. 2017), for which the adopted absolute scale was different from that acting on the regions of the PIGS-Sgr photometry. The final Sgr candidates were selected from this color diagram after applying the offset, ensuring the inclusion of stars across various metallicity bins, while also considering the configuration and positional constraints of the FLAMES/GIRAFFE 130 instrument fibers, which collectively cover a field of view 25 arcmin in diameter (for more details see Sect. 2.2).

The final targets, hereafter PIGS-HR-Sgr, are displayed in the same color-color diagram (Fig. 2), color-coded according to their spectroscopic metallicities (derived as explained in Section 3.3.1). Their positions on the diagram confirm the sensitivity of the PIGS photometry to metallicity and the accuracy of our measurements, as the majority of stars fall into the correct metallicity bins based on their spectroscopic $[\text{Fe}/\text{H}]$ values. However, there is an exception for a group of stars with $[\text{Fe}/\text{H}]_{\text{spec}} \sim -1.3/-1.4$, that the Ca H&K photometry classifies as more MP. For these stars, it is possible that the Ca H&K uncertainties (0.2-0.8 mag) played a role in their selection. On the other hand, considering the spectroscopic $\sigma_{[\text{Fe}/\text{H}]}$, which are around ~ 0.1 dex, these stars might be placed in the adjacent metallicity bin. Finally, another factor to consider is that Sgr has lower $[\alpha/\text{Fe}]$ than the training sample from *Pristine*, which is used to define the iso-metallicity lines. Indeed, the training sample is representative of the MW, with higher $[\alpha/\text{Fe}]$ than is typical for Sgr (Hasselquist et al. 2017), as already discussed in Vitali et al. (2022).

2.2. Observations

The observations were carried out between April-May 2022 using the ESO FLAMES/GIRAFFE multi-object spectrograph (Pasquini et al. 2002) mounted on the Very Large




Fig. 3. Examples of spectral lines from a FLAMES spectrum in three different setups. **Top:** The HR8 setup includes Sc II (503.1 nm), Fe I (505. nm), Y II (508.7 nm) and Ti I lines (512.2 nm), as well as the La II line at 511.4 nm, all marked with colored shaded regions. **Central:** For the HR11 filter, we display the following lines: Co I (564.7 nm), Na I (568.2 nm), Si I (569.04 nm), Cu I (570.02 nm), Ca I (570.1 nm), one line of the Mg I *b* Triplet region (571.1 nm), V I (570.1 nm), Ni I (574.8 nm) and Cr I (578.1 nm). **Bottom:** For the HR15 setup, the colored areas highlight the Ba line (649.6 nm) plus an adjacent Fe line (649.6 nm).

Telescope (VLT). For the selected field, the combination of HR8, HR11 and HR15N setups (see Table 1) were employed to observe 111 giant stars with a total wavelength coverage of 80.9 nm. Their spectra were taken in MEDUSA mode, and their signal-to-noise ratios (SNR) per pixel (1 pixel = 0.005 nm) span from 20-80 for HR8, 20-66 for HR11 and 29-99 for HR15. Table 1 reports the mean SNR for each setup, while the SNR values for the individual stars are provided in a complete table available as online material, along with comprehensive target information.

2.3. Spectra preparation

We retrieved the reduced data from the GIRAFFE ESO Phase 3 Data Stream reduction¹. The data were reduced, extracted, cleaned for cosmics and wavelength calibrated thanks to the GIRAFFE Base Line Data Reduction Software (BLDRS)².

We built an automatic multi-step pipeline that processes all of the individual exposures of the spectra to produce a final set of co-added spectra for each star. First, instrument signatures and sky emission lines are detected and removed, and the affected pixels are flagged, as their SNR is lower and some degree of residuals may persist. Next, it performs continuum normalization by iteratively (10 iterations) fitting a cubic spline while rejecting pixels using the standard deviation of the residuals as a metric. Finally, the multiple exposures of a given star are resampled to a common wavelength and co-added using mean. In the combination, pixels flagged in previous steps are ignored. Examples of reduced and normalised spectra are reported in Fig. 3. The panels display one target observed with three different grating se-

¹ http://www.eso.org/sci/observing/phase3/data_streams.html

² <http://girbldrs.sourceforge.net>




Fig. 4. Normalised distribution of RVs for Sgr samples. The target stars presented in this work appear in blue. The Sgr selections from Sestito et al. (2024a); Minelli et al. (2023); Abdurro’uf et al. (2022) are colored according to the legend. The continuous lines represent KDE curves, colored to match the corresponding Gaussian distributions. The average RV values occur at 143.9 km s^{-1} marked with the vertical dashed line.

tups. These examples have been reduced, normalized, and velocity-corrected. The corresponding SNR and $[\text{Fe}/\text{H}]$ values are indicated in the top figure, along with examples of absorption lines marked in yellow.

3. Spectroscopic analysis

3.1. Radial velocities

The radial velocities (RVs) were computed using a custom-created grid of synthetic models as templates over a range of atmospheric parameters and metallicity representative of the observed sample. For each spectrum, a first cross-correlation is performed with a randomly selected template to determine a preliminary RV value. The corrected observed spectrum is then compared to the whole grid, and the template with the lower chi-square is selected to infer the final velocity value. When calculating the cross-correlation function (CCF), the algorithm refines the maximum value by fitting a Gaussian curve. It is worth mentioning that, when computing the individual abundances, a small correction in RV was allowed to the spectrum in the rest frame to improve the fit to the specific lines used to measure the single abundances. Moreover, the RVs were computed separately for each setup, with a mean dispersion of 0.6 km s^{-1} , which falls within the average RV uncertainties of the order of $\sim 0.9\text{--}1 \text{ km s}^{-1}$ (see Appendix A), slightly higher than those of the Sgr candidate sample selected from APOGEE DR17 (Abdurro’uf et al. 2022), which are on the order of $\sim 0.6 \text{ km s}^{-1}$.

The final distribution of the Sgr RVs is illustrated in Fig. 4, which shows the RV values averaged over the three setups. Our FLAMES targets (in blue) are compared with the AAT-Sgr sample (in orange), the FLAMES high-resolution sample (light purple) from Minelli et al. (2023), a selection of Sgr members from APOGEE DR17 (in grey, Abdurro’uf et al. 2022), following the same selection (in coordinates: $b < -10^\circ$, proper motion space and radial velocity) criteria adopted in this work. These latter fall within the Sgr velocity distribution found in other stud-

ies, specifically $100\text{--}200 \text{ km s}^{-1}$. The FLAMES distribution skews towards lower RVs, possibly due to the observations being concentrated on a specific, localized field. Adopting the same method described in Sestito et al. (2024a), we inferred the systemic RV and RV dispersion using a Monte Carlo Markov Chain (MCMC) approach. A step function was employed as the prior probability distribution, considering both the system’s velocity dispersion and the uncertainties associated with individual RV measurements. By applying the same limits for the RV and its dispersion (i.e. $90 \leq \text{RV} \leq 230 \text{ km s}^{-1}$ and $\sigma_{\text{RV}} \leq 40 \text{ km s}^{-1}$), we obtained a systemic $\langle \text{RV} \rangle$ of $\sim 143.9 \pm 1.1 \text{ km s}^{-1}$ and a velocity dispersion σ_{RV} of $\sim 11.93 \pm 0.8 \text{ km s}^{-1}$, finding consistency with the values in the literature (e.g., Ibata et al. 1994; del Pino et al. 2021; Minelli et al. 2023; Sestito et al. 2024a; An et al. 2024). Specifically, examining the systematic RV values and corresponding dispersions from Sestito et al. (2024a), who divided their Sgr samples based on metallicity and on elliptical distance from the Sgr’s centre, our values are more closely aligned with those from their MR population ($[\text{Fe}/\text{H}] > -0.6$), in particular with both their total MR sample and the one at outer distances, defined as projected elliptical distances $\geq 0.25r_h$. Given that the average elliptical distance for our FLAMES field is $\approx 0.34r_h$, computed adopting the same values (coordinates of the system center, ellipticity, half-light radius, and position angle) as in Vitali et al. (2022) and Sestito et al. (2024a), our findings remain consistent with the outer MR group identified by Sestito et al. (2024a). In the case of Minelli et al. (2023), the similarity is found with their total population rather than the values for their MP and MR samples, which can be attributed to the different metallicity cuts employed in these works.

3.2. Atmospheric parameters

We followed the methodology described in Sestito et al. (2023) who employed an iterative process between the color-temperature relation from Mucciarelli et al. (2021) and the Stefan-Boltzmann equation to determine the atmospheric parameters. This strategy has proven effective in reliably deriving stellar parameters when the wavelength coverage is limited, provided the photometric quality is comparable to that of the spectra (Karovicova et al. 2020; Mucciarelli & Bonifacio 2020; Sestito et al. 2023). Besides, this approach is less affected by non-LTE effects (Frebel et al. 2013; Ezzedine et al. 2020).

The sets of effective temperatures and surface gravities were determined through an iterative process, consisting of the following steps: T_{eff} was derived using the color-temperature relation with the line-by-line spectroscopic metallicities used as input (described in Sect. 3.3.1). The surface gravity was calculated with the Stefan-Boltzmann equation (e.g., Venn et al. 2017). This latter step requires temperatures (inferred in the previous step), the heliocentric distances, *Gaia* magnitudes and bolometric corrections (Andrae et al. 2018). The necessary extinction correction was done with the 2D dust map from Schlegel et al. (1998) and updated by Schlafly & Finkbeiner (2011). The *Gaia* magnitudes were de-reddened using the extinction coefficients $A_V/E(B - V) = 3.1$ (Schultz & Wiemer 1975) and $A_G/A_V = 0.85926$, $A_{BP}/A_V = 1.06794$, $A_{RP}/A_V = 0.65199$ from Marigo et al. (2008); Evans et al. (2018). Absolute magnitudes were derived assuming a fixed distance




Fig. 5. Kiel diagram and parameter uncertainties of our stellar sample. **Left:** Kiel diagram with the parameters derived through photometry. All points are color-coded by their spectroscopic metallicities. Two examples of BaSTI isochrones are shown and zoomed in around the evolutionary stage of our giant stars. **Central and right:** The T_{eff} relative uncertainties and $\log g$ absolute uncertainties vs. effective temperatures for the entire sample following the same color-coding of the Kiel diagram.

of 26.5 kpc for the Sgr stars (Vasiliev & Belokurov 2020). The errors on the distance were assumed to be 5 kpc, which should reasonably incorporate the physical thickness of the Sgr as it is estimated to range from 0.7 to 2.6 kpc (Vasiliev & Belokurov 2020). The masses computing the surface gravities were estimated assuming a uniform distribution between 0.5 and $0.8M_{\odot}$ as a prior. These mass limits should be reasonable, as the majority of the stars are expected to be low-mass with ages of $\gtrsim 4$ Gyr. To verify the robustness of this assumption, we repeated these steps and recomputed the atmospheric parameters using a modified mass range of $0.6\text{--}1M_{\odot}$. The resulting differences in the surface gravities remained within the error bars.

For assessing the uncertainties on the measured stellar parameters we employed a Monte Carlo simulation repeated 1000 times on the input parameters. This step accounts for uncertainties in *Gaia* magnitudes, bolometric corrections, the heliocentric distance of Sgr (26.5 kpc), reddening (computed using the aforementioned extinction coefficients), and spectroscopic input metallicities, which are perturbed assuming a Gaussian distribution. The input mass values, however, are assumed to follow a flat distribution. The final mean uncertainties on T_{eff} is 75 K, while on $\log g$ is 0.11 dex.

As mentioned above, the color-temperature relation requires metallicities as input. These initial estimates were initially derived using the public spectroscopic software *iSpec* (Blanco-Cuaresma et al. 2014a; Blanco-Cuaresma 2019). At this stage, metallicities, microturbulence velocity (v_{mic}) and broadening parameters (resolution R and macroturbulence velocity v_{mac}) were inferred with *iSpec* via spectral synthesis. We adopted the radiative transfer code Turbospectrum (Alvarez & Plez 1998; Plez 2012) in local thermodynamic equilibrium, and the one-dimensional spherical MARCS model atmospheres (Gustafsson et al. 2008). Atomic data were taken from the line-list designed for the *Gaia*-ESO Survey (GES, Heiter et al. 2021). The v_{mac} was estimated using an empirical relation from the *Gaia*-ESO survey. Building on these results, we then computed the photometric T_{eff} and $\log g$. Our final parameters are thus those that combine both photometric effective temperatures, surface gravities, and spectroscopic v_{mic} and Fe values.

The Kiel diagram for the entire sample is shown in the left panel of Fig. 5, together with their relative uncertainties or absolute uncertainty in the case of $\log g$. The stars are all in the giant evolutionary stage and present a metallicity from ≈ -2.11 to -0.49 dex. To guide the eye, we compare our parameters with two BaSTI isochrones (Hidalgo et al. 2018; Pietrinferni et al. 2024) selected with the closest values to the mean $[\text{Fe}/\text{H}]$ value of -0.96 dex found with spectroscopy. The ages of 9 and 10 Gyr of the isochrones are compatible with intermediate-age and old MP populations, with metallicities of ~ -1 , as found in the study by Siegel et al. (2007). Looking at the central and right panels of Fig. 5 it is noticeable that the relative uncertainties in temperature increase for hotter stars and for the more MP ones, reaching $\sim 2.2\%$. The precision in $\log g$ shows no direct correlation with T_{eff} as it is most influenced by reddening effects, which are homogeneous within the FLAMES field. These trends might impact the derivation of elemental abundances for the colder and hotter targets, as well as for the most MP ones. The complete set of atmospheric parameters is provided as supplementary online material, following the format of Table 2.

3.3. Chemical abundances

We derived the chemical abundances for the following families of elements: α -capture (Mg, Si, Ca, Ti); odd-Z (Na, Sc, V); iron-peak (Cr, Fe, Co); n -capture (slow-process: Ba, La; rapid-process: Eu). The atmospheric parameters presented in Sect. 3.2 (T_{eff} , $\log g$) were fixed to measure line-by-line chemical abundances. To obtain the various chemical abundances we followed the strategy of Sect. 3.2, i.e. using the Turbospectrum code, the MARCS model atmosphere, and the GES line-list. During the synthesis of line-by-line abundances, we adjusted the line fitting to the different resolutions of the three setups and accounted for broadening effects by fitting the v_{mic} for each star and determining the R parameters individually for each spectrum, allowing for a local re-normalization. Although the R values for the three GIRAFFE filters are known, we allowed this parameter to vary because, in the *iSpec* fitting procedure, the R optimizes the combined effects of the star's rotational velocity and macroturbulence (Blanco-Cuaresma et al. 2014b),

Table 2. Example of the online target information table, including *Gaia* identifiers, the average SNR from the three observed setups, coordinates, G magnitudes, and our derived stellar photospheric parameters and radial velocities for two sample target stars. The complete table is provided as online material.

Target information and spectroscopic parameters									
Gaia DR3 ID	SNR	RA(deg)	DEC(deg)	G _{mag}	RV (kms ⁻¹)	T _{eff} (K)	log <i>g</i> (dex)	[Fe/H]	<i>v</i> _{mic} (kms ⁻¹)
6760361266479328000	43	284.431	-30.967	16.397	169.59 ± 0.49	4481 ± 82	1.47 ± 0.11	-0.92 ± 0.09	1.53 ± 0.07
6760381500040022528	45	284.312	-30.698	16.7227	140.58 ± 0.56	4582 ± 76	1.67 ± 0.10	-0.90 ± 0.10	1.46 ± 0.07

Table 3. Example of two rows from the online tables, presenting the line selection and the final chemical abundances. **Top:** Atomic lines used for deriving the abundance ratios presented in this work. Their logarithm values of the oscillator strength (log *gf*) and excitation potentials (*E*_{low}) are sourced from the atomic database presented in the study by Heiter et al. (2021). **Middle:** Final results from the averaged line-by-line abundances, along with the corresponding uncertainties and the number of lines used for each element. **Bottom:** Derived absolute solar abundances used in this work to calculate the final bracket abundance ratios, $[\frac{X}{Y}]$ abundance ratios.

Line selection			
Element	λ(nm)	<i>E</i> _{low} (eV)	log <i>gf</i>
Fe I	491.899	2.865	-0.342
Mg I	571.109	4.346	-1.724
Chemical abundances			
star	Element	[X/Fe]	N _{lines}
6760361266479328000	Mg I	-0.82 ± 0.09	1
6760381500040022528	Ca I	-0.61 ± 0.15	7
Solar abundances			
star	Element	A(X)	N _{lines}
Sun	Mg I	7.53 ± 0.03	1
Sun	Ca I	6.36 ± 0.06	7

while accounting for the fact that the three setups possess different resolving powers (listed in Table 1).

The line-by-line abundance measurements were averaged for each element and then expressed relative to solar values. For each setup, the solar values of each element were derived from a solar spectrum obtained from the public library of Blanco-Cuaresma et al. (2014b), which was cut and degraded to match the wavelength range and resolutions of the three different setups. The derived solar abundances are published as online material. By adding the absolute solar $[\frac{X}{Y}]_{\odot}$ ratios to our element ratios, we derived the final $[\frac{X}{Y}]$ abundance measurements with improved internal accuracy.

The combination of the three GIRAFFE setups allows us to measure different chemical species across three wavelength windows of approximately $\sim 240-320$ Å. Due to the limited wavelength coverage and varying resolutions, the line selection was customized independently for each setup. Initially, we relied on the quality flags *synflag* (indicating blending properties) and *gf_flag* (indicating the accuracy of the log *gf* value) associated to the GES line-list (Heiter et al. 2021). However, due to the low SNR of some of the targets, we performed a subsequent visual inspection to discard lines with problematic fits, ultimately retaining only one or a few lines per element. The only exception was for Fe I lines, where we removed those across the three setups that showed systematically different results (see sections below for further details). Furthermore, due to the quality of our spectra, some elements could only be measured for a

sub-sample of our FLAMES dataset. This restrictive strategy ensures that we obtain trustworthy results for studying the chemical evolution of Sgr.

The spectroscopic analysis was done assuming local thermodynamic equilibrium (LTE). Hyperfine structure (HFS) and isotopic splitting were taken into account in the analysis using the atomic data compiled from Heiter et al. (2021) and the references therein.

Concerning the uncertainties in the abundances, iSpec employs two internally consistent and homogeneous methods (Blanco-Cuaresma et al. 2014a). When sufficient spectral lines are available, uncertainties are calculated using the standard error of the mean of the individual line-by-line measurements. If there are not enough lines, the uncertainties were instead determined from the covariance matrix generated by the non-linear least-squares fitting algorithm. This method accounts for how the synthetic spectrum changes when the best-fit parameters are adjusted, as well as the residuals between the synthetic and observed spectra. It is important to emphasize that for many elements (e.g., Mg I, Cu I, Co I, Cr I, La II, Eu II), we were able to find only a few lines, or in some cases, just one line, suitable for abundance determination. The derivation of abundances for these species, which have fewer detected lines, can be more challenging if the only measurable lines are weak, exhibit blends, or are affected by uncertainties in atomic or molecular data. However, the internal uncertainties are not necessarily larger compared to abundances measured from multiple lines (e.g., N_{lines} $\gtrsim 3$). In fact, if the selected lines originate from different setups (with varying *R* and SNR), this can increase the internal abundance spread, affecting the precision of the measurement.

The line selection used in the analysis to determine the final average abundances for the entire sample is available as online material, with the first lines presented as an example in Table 3.

3.3.1. Stellar metallicities

The iron abundances were derived for the entire stellar sample, with values spanning from -2.13 to -0.35 dex. The typical metallicity uncertainties are approximately 0.1 dex. The final analysis employed eighteen Fe lines, sourced from the blue setup HR8 and the intermediate setup HR11. Lines from HR15 were excluded after it was found they exhibited an offset compared to the other two setups, likely due to their lower spectral resolution (see Sect. B).

The metallicity distribution, plotted against RV values, is shown in Fig. 6 and compared with the same literature samples already reported in Fig. 4. The literature [Fe/H] distribution is bimodal, reflecting the selection biases of various surveys and samples. Our targets are positioned at the tail of the more MR peak, with the majority of stars having [Fe/H] ~ -0.7 dex. Although our sample includes




Fig. 6. Metallicities and RVs for the Sgr members observed in this study are compared with those from three literature samples. These are the same samples shown in Fig. 4, and the selection criteria applied are similar to those used for the FLAMES targets. **Top:** Histogram of RVs, with literature data shown in light-blue and our targets in blue. **Bottom right:** Histogram of metallicities, using the same color scheme as the RV histogram.

fewer stars in the more MP range (77 stars with $[\text{Fe}/\text{H}] \geq -1.0$ dex), it contains 34 stars with $[\text{Fe}/\text{H}] < -1.0$ dex among which 13 with $[\text{Fe}/\text{H}] < -1.5$ dex.

3.3.2. α -elements: Mg, Si, Ca, Ti

The α -elements are synthesized in the cores of massive stars during the late stages of evolution. Specifically, they are produced and ejected to the ISM in core-collapse supernovae (CCSNe) via different processes (Heger & Woosley 2002; Kobayashi et al. 2006; Nomoto et al. 2013). These processes are expected to enrich the ISM on short timescale (Matteucci 2016).

Among this group, we were able to measure four different species. Magnesium (Mg) was derived for 82 stars, relying on a single line at 571.10 nm (HR11) that is not affected by blending, resulting in an average abundance of $\langle [\text{Mg I}/\text{Fe}] \rangle = 0.12 \pm 0.09$ dex. Silicon (Si) measurements were more challenging to derive due to the low SNR of the spectral lines. Final Si I abundances are available for only 17 stars, inferred from two weak lines at 569.04 nm and 579.30 nm (HR11). An attempt was made to use the line at 672.18 nm in the redder setup, but it turned out to be too weak for reliable fitting. The average abundance ratio is given by $\langle [\text{Si I}/\text{Fe}] \rangle = 0.18 \pm 0.06$ dex. For calcium (Ca) more absorption lines were available throughout the HR15 setup and was derived for 95 targets. The $\langle [\text{Ca I}/\text{Fe}] \rangle = 0.17 \pm 0.11$ dex is based on seven Ca I lines from $\approx 646 - 671$ nm. Some of these lines are slightly affected by blending effects, which could be contributing to the higher uncertainties. Finally, titanium (Ti) was measured for two different ionization stages. For Ti I, abundances were computed for 78

stars using five lines from three different setups, resulting in an average value of $\langle [\text{Ti I}/\text{Fe}] \rangle = 0.11 \pm 0.11$ dex. We were able to fit the ionized Ti II line at 649.15 nm, but we discarded it due to poor flags in the GES line list.

3.3.3. Fe-peak elements: Cr, Co

Due to the broad range of atomic numbers covered by iron-peak elements ($21 \leq Z \leq 30$), various complex nucleosynthesis processes have been identified as contributing to their production. In our analysis, the Fe-peak elements examined are those elements close to iron in the periodic table, with their primary contribution coming from Type Ia supernovae (SN Ia). These explosions typically occur in binary systems with a white dwarf as a companion (Woosley & Weaver 1995; Seitzzahl et al. 2013) and synthesize these elements on a longer timescale compared to the α -elements (Matteucci et al. 2009).

For chromium (Cr), the final results are based on the spectral line at 493.63 nm in HR8 with good GES flags are $\langle [\text{Cr I}/\text{Fe}] \rangle = 0.03 \pm 0.11$ dex and are averaged on 73 Sgr stars. Cobalt (Co) abundances were derived for 83 stars using two lines at 677.04 nm and 564.72 nm, both accounting for HFS splitting (Pickering 1996). The resulting average abundance is $\langle [\text{Co I}/\text{Fe}] \rangle = 0.02 \pm 0.06$ dex.

3.3.4. Odd-Z elements: Na, Al, Sc, V

Among the odd-Z group, the light elements sodium (Na) originates in massive stars via hydrogen burning or carbon burning (Cristallo et al. 2015; Kobayashi et al. 2020). Aluminum (Al) is noted to be an element whose yields are metallicity-dependent (Nomoto et al. 2013), and at the same time, it is mostly produced in CCSNe at rates comparable to α -capture elements (Kobayashi et al. 2020). Heavier elements from this group, specifically scandium (Sc) and vanadium (V), were analyzed in this study. Sc shares similar origins with Na, as it is synthesized during the carbon and neon burning phases in massive stars, while V is primarily produced through silicon burning in Type II supernovae (SNe II, Woosley & Weaver 1995; Battistini & Bensby 2015).

For the light element Na, we relied on the HR11 setup. We identified two Na I lines at 568.26 nm and 568.82 nm and measured sodium abundances for sixty targets. However, both lines are affected by blending features, and for roughly half of the sample, we were unable to achieve a satisfactory fit. This may be reflected in the overall average uncertainty of the final abundance measurements $\langle [\text{Na I}/\text{Fe}] \rangle = -0.18 \pm 0.13$ dex. Concerning Al, its derivation posed challenges as we could only use for most of the stars a single absorption line in the redder part of the spectrum (HR15) at 669.86 nm. For a minority of stars, we were also able to measure abundances from a line at 669.60 nm. According to the GES flag only the redder Al line (669.86 nm) is unblended. However, this line was quite weak in our spectra and could only be fitted for a small minority of the sample, specifically 17 stars. Their average abundance was found to be $\langle [\text{Al I}/\text{Fe}] \rangle = 0.06 \pm 0.06$ dex.

Moving to the heavier elements in this group, Sc was inferred for 62 stars relying on four lines from the HR11 (between 565 and 566 nm) and HR15 (660 nm) setups, resulting in $\langle [\text{Sc II}/\text{Fe}] \rangle = -0.04 \pm 0.06$. For V, six lines in the

central part of the wavelength range (562–573 nm) were fitted for 75 stars providing the mean abundance ratio is $\langle [VI/Fe] \rangle = 0.06 \pm 0.1$ dex. The HFS data were taken from Childs et al. (1979) and Cochrane et al. (1998).

3.3.5. N-capture elements: Ba, La, Eu

Neutron (n)-capture elements are those heavier than zinc ($Z > 30$), formed through the capture of neutrons by a seed nucleus, which subsequently undergoes β -decays (Burbidge et al. 1957; Sneden et al. 2008). Based on the density of the neutron flux, which determines the rate of nucleosynthesis, these processes are classified as slow (s), intermediate (i), and rapid (r) processes (Cowan & Rose 1977; Busso et al. 1999; Kratz et al. 2007). The s -process is known to take place in low- and intermediate-mass stars during their asymptotic giant branch (AGB) phase in longer timescales (see for instance Bisterzo et al. 2014) differently to r -process that is believed to occur in CCSNe and neutron-star mergers (e.g., Côté et al. 2018).

From the s -process elements, we derived barium (Ba) abundances using the redder line at 649.68 nm (HR15), which is generally strong in stellar spectra, except in VMP stars, but with reliable $\log gf$ data. However, due to the low resolution of this spectral window for the majority of the targets, we were only able to fit the line correctly for 20 stars, resulting in an average abundance of $\langle [Ba II/Fe] \rangle = 0.39 \pm 0.06$ dex. We then measured the heavy s -process element lanthanum (La) in 73 stars, more than double the number of stars compared to Ba. For these lines, corrections for HFS and isotopic splitting are the ones measured by Villemoes et al. (1993). The average abundance $\langle [La, II/Fe] \rangle = 0.67 \pm 0.10$ dex was computed using a line in the HR8 region at 492.17 nm, which is only slightly affected by blending. As with the Ba line, HFS corrections were necessary in the spectral analysis (Lawler et al. 2001). The only r -process element derived for our sample is europium (Eu). Although Eu abundances are typically measured from the bluer part of stellar spectra ($\lesssim 420$ nm), the Eu II line at 664.51 nm in the HR15 region can be measured in giant stars with higher SNR. Despite the inclusion of HFS (Villemoes et al. 1993) and isotopic corrections (Lawler et al. 2001), which reduce the impact of blending effects, Eu measurements remained difficult and were only possible for 29 stars, with an average abundance of $\langle [Eu, II/Fe] \rangle = 0.40 \pm 0.07$ dex.

We attempted to measure some of the light elements belonging to the s -process family, such as yttrium II at 508.74 nm and zirconium I and II at 507.82 nm and 511.22 nm, respectively. However, no successful spectral synthesis yielded satisfactory results.

4. Results & Discussion

4.1. Comparison with other Sgr samples

Figure 7 displays our PIGS-HR-Sgr element abundances (blue points) compared with the Sgr core selection performed by Hayes et al. (2020) on the near-infrared (NIR) APOGEE data (light blue triangles). The comparison is further extended to include five additional high-resolution optical spectroscopic samples and one medium-resolution sample, all focusing on the Sgr core.

The studies by Monaco et al. (2005) and Sbordone et al. (2007) conducted with FLAMES-UVES and UVES respectively, include 15 and 12 RGB Sgr stars with $-1.0 \lesssim [Fe/H] \lesssim 0.0$, with one star at $[Fe/H] = -1.52$ dex (Monaco et al. 2005). Both catalogs include abundances for α , Fe-peak and heavy elements. Again with FLAMES, Carretta et al. (2010) inferred the abundances of various elements in 27 giant stars ranging from solar metallicity to -1.2 dex in the core of the Sagittarius galaxy. The mid-resolution sample of Mucciarelli et al. (2017) comprises 235 Sgr giant stars (light grey crosses) from both the main body and the nuclear cluster M54. Iron and α -element abundances are provided across a wide metallicity range ($-1.8 \lesssim [Fe/H] \lesssim -0.1$). Lower metallicity regimes are covered by the studies of Hansen et al. (2018) and Sestito et al. (2024b). The former includes 13 different chemical species for 13 giant stars with $-2.97 < [Fe/H] < -1.07$ dex, while the latter analyzed 12 low-metallicity Sgr stars with $[Fe/H]$ down to -3.26 dex, providing a detailed chemical analysis. Lastly, we included the studies by Smecker-Hane & McWilliam (2002) and McWilliam et al. (2013). From Smecker-Hane & McWilliam (2002), we obtained Na, Al, La and Eu measurements for 14 RGB stars in Sgr, while from the McWilliam et al. (2013), we retrieved 3 RGB stars in the M54 cluster for the remaining elements (pink pentagons in Fig. 7). In this panorama, the PIGS-HR-Sgr sample, with $-2.13 < [Fe/H] < -0.5$ dex, represents the most complete high-resolution dataset.

Examining the different panels of Fig. 7, our data align with most literature trends. The α -elements exhibit the expected behavior with respect to metallicity. In fact, both our results and those from the literature show the typical decline in α abundances as metallicity increases, which is attributed to the time delay between prompt SNe II and SNe Ia, acting on longer timescales. The most noticeable differences are the ones with the APOGEE data, particularly for elements such as Si I, and less for Ti I. The NIR wavelength coverage of APOGEE (as opposed to the optical range used for the other samples) partly explains these discrepancies, as the offset in NIR Ti I abundances may be influenced by problematic lines or their dependence on uncalibrated T_{eff} values (Jönsson et al. 2020). This discrepancy with APOGEE, which becomes larger for the most iron depleted stars, is also observed in the samples from Monaco et al. (2005); Carretta et al. (2010) and Mucciarelli et al. (2017), while the Ti abundances reported by Sbordone et al. (2007) are lower and more comparable to those of APOGEE, though they cover the most MR end of the distribution. The results of Sestito et al. (2024b) account for NLTE corrections which range from 0.1 ($[Fe/H] \lesssim -0.5$ dex) to 0.3 dex for the most MP targets ($[Fe/H] \lesssim -1.5$ dex) according to Bergemann (2011). The other samples do not.

Si also shows an offset with the APOGEE measurements, which is expected to be very precise (Hegedűs et al. 2023). We therefore believe that our measurements, based on two weak absorption lines, may exhibit a systematic offset compared to the APOGEE measurements. This offset (~ 0.24 dex) could be due to differences between optical and infrared observations or line-list systematics. We consulted the MPIA database³ to gather NLTE corrections for the Si I lines used (Bergemann et al. 2013). However, the

³ <https://nlte.mpia.de/>




Fig. 7. Chemical abundances $[X/Fe]$ as a function of $[Fe/H]$ for different Sgr samples. The FLAMES targets analysed in this work are represented as blue circles. Different literature samples, represented by distinct colors and markers, provide additional $[X/Fe]$ data. These include the APOGEE dataset for the Sgr main body from Hayes et al. (2020), and optical data for the Sgr core from Monaco et al. (2005), Sbordone et al. (2007), Carretta et al. (2010) and Mucciarelli et al. (2017), with the most metal-poor stars reported by Hansen et al. (2018) and Sestito et al. (2024b). The studies by Smecker-Hane & McWilliam (2002) and McWilliam et al. (2013) contain data for both the nuclear cluster M54 and the main body of the galaxy.

maximum correction of 0.03 dex is insufficient to account for the offset observed in our results. On the other hand, our trend aligns much more with the optical samples from Sbordone et al. (2007) and Carretta et al. (2010) and all show the typical decline of α -elements with metallicity.

Mg and Ca show the most prominent drop with metallicity (up to ~ 1 dex) and the trends for our FLAMES targets are well aligned with those of the other optical Sgr samples but deviate more from the Hayes sample. Since APOGEE applies NLTE corrections for Mg and Ca (Osorio et al. 2020; Jönsson et al. 2020), we verified that ap-

plying corrections from the MPIA database (Bergemann et al. 2015) of ~ -0.2 dex for the Mg line at 571.10 nm would align our results with the APOGEE measurements. Regarding Ca, we were unable to find NLTE corrections for the lines used in this study.

Analyzing the overall trend of Na from the FLAMES sample along with three other optical samples, it appears that Sgr shows a decline in $[Na/Fe]$ with increasing metallicity. This trend is expected, as Na is predominantly produced by massive stars (Woosley & Weaver 1995). Our values do not display such strong decline nevertheless they

are comparable to the compilation reported by McWilliam et al. 2013 (see their Fig. 4). The measurements from Carretta et al. (2010) exhibit a different behavior, suggesting an increase for more metal-rich (MR) stars, making the overall trend harder to interpret. NLTE corrections that could be applied to our lines might lead to adjustments of -0.05 to -0.1 dex (Lind et al. 2022), which would bring our results closer to those of Sbordone et al. (2007). The results for Al, derived using LTE for all samples, are presented in the first row. Despite a slight spread at lower metallicity, a clear, tight trend highlights the primary origin of this element, predominantly from CCSNe (Kobayashi et al. 2020). However, the large variation of nearly 1 dex in $[\text{Al}/\text{Fe}]$ could be attributed to the strong metallicity dependence of this element (Nomoto et al. 2013). Nevertheless, a clear trend remains, reflecting the primary source of this element, which is mostly CCSNe (Kobayashi et al. 2020). AGB stars may also contribute to Al production, which aligns with the previously studied scenario that these stars played a role in the chemical enrichment of Sagittarius (Hasselquist et al. 2017; Hansen et al. 2018; Sestito et al. 2024b).

The heavier odd-Z elements, Sc and V, follow different trends with metallicity. Sc shows a clear decrease in abundance with increasing $[\text{Fe}/\text{H}]$, a trend consistent with the findings of Sbordone et al. (2007) and Carretta et al. (2010). This behavior aligns with Sc's primary production by CCSNe. In contrast, V exhibits a less pronounced decrease, though there is a general downward trend when considering both the FLAMES sample and the work of Sbordone et al. (2007). For this element, APOGEE is known to have low reliability in its abundance measurements (Buder et al. 2021; Hegedűs et al. 2023).

The two Fe-peak elements align well with the literature. The increase in Cr with metallicity matches expectations, as its primary source is SN Ia (Kobayashi et al. 2020). In contrast, Co shows a more variable trend due to its more inhomogeneous origins, which include both CCSNe and, to a lesser extent, AGBs (Nomoto et al. 2013; Kobayashi et al. 2020). Although NLTE corrections can affect the abundances of Cr and Co lines by 0.1 to 0.2 dex (Bergemann & Cescutti 2010; Bergemann et al. 2010), we did not apply these corrections because the comparison samples also assume LTE in their analyses.

The three last elements presented in Fig 7 belong to the n -capture elements. As previously mentioned, the heavy s -process elements (or second-peak elements), being released over longer time scale by AGB stars, exhibit an increase over time and with metallicity. This trend is evident for both Ba and La and is consistent across all samples reported in the figures, collectively showing an increase of nearly 2 dex for both elements. It is important to note that for Ba, we used the line at 649.68 nm, which is known to be affected by NLTE effects (see, e.g., Korotin et al. 2015). These effects can cause a variation in $A(\text{Ba})$ of about -0.2 dex for our atmospheric parameter space (Eitner et al. 2019). However, this variation does not eliminate the overall increasing trend with $[\text{Fe}/\text{H}]$. Since only Sestito et al. (2024b) accounts for these corrections, we retain the abundances on the LTE scale for consistency with other samples. The PIGS-HR-Sgr La measurements, which cover a broad range of metallicities, show a sharper increase compared to the results of Hansen et al. (2018), which only begin to rise after leveling off for $[\text{Fe}/\text{H}] < -2.0$ dex. In addition, the

samples from Sbordone et al. (2007) and Smecker-Hane & McWilliam (2002) display a trend very similar to ours.

One star in our sample stands out with significantly higher La and Ba abundances, enhanced by $\approx 1-1.5$ dex relative to the sample average. This star shows no distinct characteristics from the rest of the Sgr sample in the parameter or other chemical spaces. To confirm its membership in the Sgr core, we examined its RV and proper motion values, finding them to validate the Sgr origin of the target. These checks, combined with a visual inspection of the best fits to the La II and Ba II abundances, confirmed the genuine enhancement of heavier elements in this star, though this enhancement does not extend to the r -process element Eu. Generally, these overabundances of elements synthesized by the slow n -capture process in stars such as giants, which cannot produce s -process elements internally at their evolutionary stage, are thought to result from binary interactions (Burbidge et al. 1957; Käppeler et al. 2011). We examined the RV variability of this target by comparing the RV values derived from three different GIRAFFE setups and comparing these results to a star with "normal" heavy element abundances. No RV variability was detected beyond the precision limits of GIRAFFE, and the RV curves for both the potential binary target and the comparison star were consistent. However, this is not sufficient to exclude the possibility of a binary nature for the star, as the orbital period could be very long, potentially reaching up to 30,000 days (e.g., Escorza et al. 2019; Jorissen et al. 2019) and thus not detected by our few RV measurements. To further investigate the potential binarity of this Sgr giant and achieve a more accurate and complete chemical characterization, a broader wavelength coverage would be required.

The spread in Europium (Eu) abundances highlights the challenges in accurately measuring this element. Although the data from Sestito et al. (2024b) do not reveal a clear trend with $[\text{Fe}/\text{H}]$, when considering the entire optical sample, there is an observable increase of approximately 0.5 dex. If we exclude the literature measurements from Sestito et al. (2024b) and Hansen et al. (2018), which cover the most metal-poor regime, our FLAMES data, along with those of Smecker-Hane & McWilliam (2002) (pink pentagons), level off around $[\text{Eu}/\text{Fe}] \sim 0.5$ dex. This rise in Eu abundances is probably an indication of a prolonged and later active star-forming period, during which massive stars may have continued enriching the ISM with r -process elements (Reichert et al. 2021). Other exotic scenarios have also been proposed to explain the increase in r -process elements at higher metallicities, such as neutron star mergers (NSMs) contributing with a delayed production of Eu (Côté et al. 2018; Skúladóttir & Salvadori 2020) or massive stars that formed and exploded as magneto-rotational supernovae (MR-SNe, Reichert et al. 2021). This scenario will be further investigated in Sect. 4.4.

4.2. α -knee

In Fig. 8, we further compare the well-known α -knee diagram, or Tinsley diagram (Tinsley 1979), which is often used as an effective tracer of early star formation efficiency in galaxies. The dominance of the yields of CCSNe during the early chemical enrichment results in the production of α -elements. This is followed, after a time delay, by the metal production from SN Ia, which predominantly generates iron-peak elements. Therefore, the metallicity at which




Fig. 8. Trend of the α -elements vs. $[\text{Fe}/\text{H}]$ for the FLAMES targets (blue circles), compared with literature samples from Monaco et al. (2005); Sbordone et al. (2007); Mucciarelli et al. (2017); Sestito et al. (2024a). For all the samples, Mg and Ca abundances, propagating their corresponding uncertainties, were averaged and then fitted with two linear functions connected by a smoothing function. The grey shading represents the fit uncertainties, which are also displayed as residuals in the bottom plot. As explained in the text the data from Sestito et al. (2024a) were excluded from the fitting procedure. The grey dashed line marks our results for the knee position at $[\text{Fe}/\text{H}] = -1.05$, while the black dotted line at $[\text{Fe}/\text{H}] = -1.3$ represents an approximation of results from Carretta et al. (2010) and de Boer et al. (2014)

the turnover in the $[\alpha/\text{Fe}] - [\text{Fe}/\text{H}]$ occurs is linked to the star formation rate of a galaxy. A lower metallicity suggests a galaxy with a less efficient SFH as fewer SNe II events balanced the delayed contribution from SN Ia. In contrast, galaxies that build up and retain metals and gas quickly before the contribution of SN Ia to chemical enrichment will reach a higher metallicity at the α -knee (e.g. Kobayashi et al. 2006; Kirby et al. 2020).

For our data the fitting was performed using the `scipy` Python function `curve_fit`. To increase the statistical significance, we complemented our sample with three literature samples from Monaco et al. (2005); Sbordone et al. (2007); Mucciarelli et al. (2017) covering similar metallicity range. As in the comparison studies, we average the Mg and Ca as a proxy for α abundances. We applied a regression model consisting of two linear functions connected by a sigmoid smoothing function to fit the knee point at $[\text{Fe}/\text{H}]_{\alpha\text{-knee}} = -1.05 \pm 0.13$. The residuals of the fit are shown in the bottom panel of Fig. 8. The data from Sestito et al. (2024b) are displayed on the same figure, covering a different metallicity range. Although not statistically sufficient for fitting, this dataset serves as a visual guide to the average trend for metallicities below $[\text{Fe}/\text{H}] = -1.0$. If SN Ia were already influencing at this metallicity regime, we would expect a distinct slope, which is hinted at by the data from Mucciarelli et al. (2017). In the more metal-poor range observed by Sestito et al. (2024b), the data appear to level off, suggesting a low contribution of SN Ia to the chemical enrichment in this metallicity regime. Moreover, when examining our FLAMES data (shown in blue in Fig. 8), we




Fig. 9. Ratio of the average of the explosive α -elements (Ca and Ti) to the hydrostatic α -elements (Mg) as a function of $[\text{Mg}/\text{Fe}]$. The color code encodes the metallicity of the Sgr stars (filled circles). The square symbols denote the theoretical models from Nomoto et al. (2013), which incorporate yields from both SNe II and HNe. The dash-dot line delineates the region of chemical space where stars could have originated from gas enriched by both CCSNe and HNe.

can distinguish between the main stellar population, characterized by $[\text{Fe}/\text{H}] \sim -0.8$, and the remaining older population with $[\text{Fe}/\text{H}] < -1.0$. This metal-poor population shows a plateau at $[\alpha/\text{Fe}] \approx 0.3$, consistent with the trend observed in the more metal-poor stars studied by Sestito et al. (2024b). This behavior suggests the absence of significant Type Ia SNe contributions in this metallicity regime. Additional data for $[\text{Fe}/\text{H}] < -1.0$ and the comparison with chemical evolution models would help clarify the timing of SN Ia onset in this regime.

It is interesting to note that, in the third panel of the first row of Fig. 7, the $[\text{Al}/\text{Fe}]$ values begin to flatten at $[\text{Fe}/\text{H}]_{\alpha\text{-knee}} = -1.05$, corroborating the occurrence of the knee around this metallicity. However, the lack of higher statistics makes it impossible to exclude the possibility of a shift toward slightly lower metallicities, as suggested by Mucciarelli et al. (2017). Additionally, we acknowledge that the scatter in the abundance data makes the shape of the α -knee highly sensitive to the smoothing parameter and that the paucity of data below $[\text{Fe}/\text{H}] < -1.0$ affects the precise determination of the knee position. Another contributing factor is that the compilation of data-set presented in Fig. 8 is inhomogeneous in terms of resolution and instrumentation, which can introduce systematics into the comparison.

Despite these limitations, our result lines up with the finding of de Boer et al. (2014), who however determined the α -knee at $[\text{Fe}/\text{H}]_{\alpha\text{-knee}} = -1.27$ for the stellar streams of Sgr. We also compared our result with that obtained using the formula presented by Reichert et al. (2020) (equation 6), which posits a dependence of the position of the α -knee on the luminosity (and thus the mass) of the galaxy. By applying this formula, we found comparable results within the uncertainties: $[\text{Fe}/\text{H}]_{\text{knee, estimates}} = -1.65 \pm 0.48$ (using the absolute visual magnitude $M = -13.50$ from McConnachie 2012). This corroborates the idea that smaller systems with lower initial amount of Fe exhibit the knee position at lower metallicities compared to more massive galaxies. Carretta

et al. (2010) confirmed the knee at $[\text{Fe}/\text{H}] \sim -1.3$ dex, occurring at a lower metallicity than our result, which can be attributed to their use of data from M54 and the averaging of Si, Ca, and Ti to represent α -element abundances. The extensive work of Hasselquist et al. (2021) compares abundance trends for different MW satellites using chemical evolution models. Among these, they analyzed Sgr and found that α abundances flatten at $-1.0 \lesssim [\text{Fe}/\text{H}] \lesssim -0.8$, a result that is slightly more MR compared to our findings. However, according to both our results and theirs, Sgr differs from the LMC and SMC. Despite being ~ 100 times more massive, the knee of the LMC is found at lower $[\text{Fe}/\text{H}]$ values ($-2 \lesssim [\text{Fe}/\text{H}] \lesssim -1.8$ Nidever et al. 2020; Hasselquist et al. 2021; Chiti et al. 2024) due to its isolated existence. On the other hand, the literature results for the GES system indicate that $[\text{Fe}/\text{H}]_{\alpha\text{-knee}} \sim -1.2$ (e.g. Mackereth et al. 2019; Fernandes et al. 2023), suggesting that Sgr and GES share similar early star formation efficiencies both influenced by the proximity and the encounter with the MW.

Another insightful diagnostic diagram involving α -abundances is shown in Fig. 9, based on the recent study by Bandyopadhyay et al. (2024). The main idea is to use the ratio between hydrostatic burning (in our case Mg) and explosive nucleosynthesis (in our case Ca and Ti, we do not include Si as it is computed only for a minor part of the sample) α -elements and no Fe dependence. The underlying concept is that the mass of the CCSNe progenitor influences this ratio. Specifically, more massive stars ($M > 35M_{\odot}$) contribute to the abundance of hydrostatic elements like Mg, while less massive stars ($M < 25M_{\odot}$) are responsible for explosive elements like Ca and Ti (Woosley & Weaver 1995; Kobayashi et al. 2020). This suggests that the observed trends are not solely due to the onset of SN Ia, which would reduce the Mg abundance in more MR stars, but also reflect the contribution of slightly less massive CCSNe, which could influence these trends (McWilliam et al. 2018). Moreover, Bandyopadhyay et al. (2024) proposed a limit of $\Delta\alpha < -0.2$, suggesting that stars with high abundances of hydrostatic elements could originate from gas enriched by hypernovae (HNe) and are represented in Fig. 9 as filled squares. These yields overlap with the locus of the few FLAMES targets exhibiting high Mg abundances, reinforcing the possibility that some of the metal-poor stars in Sgr were enriched by HNe.

Carlin et al. (2018) investigated this same α -ratio for stars located in the Sgr stellar streams. Their analysis revealed a lower abundance of hydrostatic elements compared to explosive elements. By comparing these results with those from the LMC and the Fornax dSph, they concluded that the observed pattern suggests a dearth of massive stars, indicating a top-light IMF. The reported $\Delta\alpha$ values appear to be comparable with ours, which is expected given that the streams and the core are components of the same progenitor. These hypotheses could still accommodate the presence of rare hypernovae, which would primarily affect the more MP (and possibly older) stars (Lee et al. 2024). According to the detailed analysis by Skúladóttir et al. (2024) on Sculptor, the contribution of HNe becomes significant at much lower metallicities. Similarly, contributions from HNe have also been detected in the low-metallicity regime of Sagittarius from the pattern of the lighter elements (Sestito et al. 2024b) and from its

average $[\text{C}/\text{Fe}]$, which is lower in this system than in the MW (Sestito et al. 2024a).

4.3. Chemical evolution of the biggest Milky Way satellites

To gain deeper insights into the chemical enrichment of Sgr, we examine the same chemical trends shown in Fig. 7 this time comparing them with samples from various MW satellites/structures and MW halo stars. Specifically, for the MW stars we used the Stellar Abundances for Galactic Archaeology database⁴ (SAGA, Suda et al. 2008). From this database, we queried stars with atmospheric parameters similar to our sample, restricting our selection to giant stars to minimize potential systematic effects in the comparison. To ensure a sufficient number of data points for the comparative analysis, we set the maximum queried resolution to 40,000, which is higher than that of GIRAFFE.

We chose to use two massive systems for comparison: 62 GES members from the work of Carrillo et al. (2022), and 126 LMC stars presented in the studies of Van der Swaelmen et al. (2013) and Pompéia et al. (2008). Although these systems are among the most massive structures that have interacted or will interact with the MW, their estimated masses and evolutionary histories differ significantly. Indeed, GES (with an estimated total mass of $\sim 10^9 M_{\odot}$, e.g., Mackereth & Bovy 2020; Feuillet et al. 2020) was fully disrupted following its merger with the MW $\sim 8 - 11$ Gyr ago (Belokurov et al. 2018; Helmi et al. 2018; Das et al. 2020). In contrast, the irregular and more massive LMC galaxy (total mass of $\sim 10^{10-11} M_{\odot}$, Erkal et al. 2019; Shipp et al. 2021) has evolved in almost near isolation and is likely encountering the MW for the first time (Besla et al. 2007).

The other two objects of comparison, Fornax and Sculptor, are dSph galaxies. Fornax is the second most luminous and massive dSph galaxy in the Local Group (after Sgr), with an estimated total mass of $\sim 1.6 \times 10^8 M_{\odot}$ (de Boer et al. 2012b; Lokas 2009), showing a metallicity range comparable to that of Sgr. On the other hand, Sculptor, is less massive ($\sim 3 \times 10^7 M_{\odot}$ Lokas 2009) and significantly more metal-poor ($\langle [\text{Fe}/\text{H}] \rangle \sim -1.8$ Bettinelli et al. 2019; de Boer et al. 2012a).

The different $[\text{X}/\text{Fe}]$ vs $[\text{Fe}/\text{H}]$ trends are displayed in Fig. 10. It is important to note that, as with the comparison discussed in Sect. 4.1, there are inherent caveats when confronting results from different studies. These stem from the use of different instruments for observations and diverse analytical approaches (line selection, solar abundances, etc.). Hence, systematic offsets might be present in these trends. Finally, since no measurements for Al were available from the comparison samples, this element was excluded from the figure. From the various panels, we found that for most elements, Sgr exhibits similarities to the MW halo sample and the more massive systems, GES and LMC. As expected, Sculptor shows the most significant differences, likely due to its mass and the lower metallicities of its stellar populations. Although less pronounced than in Sculptor, Fornax also displays some differences. However, we were not able to obtain measurements for all abundances in this system.

The abundance trends analyzed in this study do not show signs of multiple sequences, indicating that the Sgr stars were formed from the galaxy's own gas, without significant contamination from the MW. This conclusion is

⁴ <http://sagadatabase.jp/>




Fig. 10. Chemical abundances $[X/Fe]$ as a function of $[Fe/H]$ for the Sgr sample (blue filled dots) are compared with MW halo stars and members of various MW satellites. Specifically, data for the MW halo (grey empty dots) are queried from the SAGA database (Suda et al. 2008). The GES stars (orange) are taken from Carrillo et al. (2022), LMC members (pink) from Van der Swaelmen et al. (2013), and Na measurements from Pompéia et al. (2008). Finally, stars from the Fornax galaxy (purple) are from Letarte et al. (2010), while Sculptor members (green) are from Hill et al. (2019).

consistent with the spatial distribution of the targets, located near the nuclear cluster in a region where the gravitational potential well of Sgr makes it unlikely for external gas to have infallen. Consequently, the chemical enrichment revealed by the analysis reflects the undisturbed chemical history of stellar populations that evolved within the system, corresponding to the various formation periods of this galaxy.

Regarding the α -elements, as noted in Carrillo et al. (2022), we find that all the satellites exhibit lower abundances compared to the MW sample. This suggests a lower star formation rate or a slower process of chemical evolution and enrichment. The deficiency relative to the MW is particularly evident in the hydrostatic element Mg, which may indicate a lack of more massive SNe II enriching the ISM of the satellites compared to that of the MW, and, in the case of Sgr, it was connected to a top-light IMF by different studies (e.g., McWilliam et al. 2013; Hasselquist et al. 2017). It is worth emphasizing that differences in the levels

of Mg and Ca are challenging to interpret, as they may result from the system’s star formation rate or the yields of SN Ia (Matteucci et al. 2009; Kobayashi & Nomoto 2009; Kobayashi et al. 2020). Consequently, linking the different enrichment levels between Sgr and the MW solely to differences in the IMF is not straightforward.

Among the satellites, there are also differences, with Sculptor and Fornax displaying the lowest α -abundances, particularly under-abundances in Mg, Si, and Ca. This difference is less pronounced when examining the GES dataset. According to Hasselquist et al. (2021) this can be explained by the fact that, compared to more isolated dSphs, which exhibit weaker earlier star formation periods, the GES showed enrichment to higher levels before the onset of SN Ia.

Unlike Sbordone et al. (2007) and Hasselquist et al. (2017), but similar to McWilliam et al. (2013), we find that for Sgr, the light odd-Z elements (Na, Sc, V) are not particularly different from the MW, GES, and LMC trends, ex-

cept for V in the LMC. In this case, [Van der Swaelmen et al. \(2013\)](#) noted that accounting for HFS corrections lowers their V abundances by about 0.2 dex in their LMC sample, which can explain the difference displayed in Fig. 10. Moreover, we do observe a slight depletion in Na abundances between Sgr and the MW. Since Na is synthesized during the hydrostatic stages of massive stars in a mass-dependent process, its abundance may suggest a relative lack of more massive stars compared to the MW or different mass ranges of the type of SNe responsible of the Na production. For Sc, compared to Sculptor, the $[X/Fe]$ ratio at the same metallicity is higher in the more massive systems, including Sgr. This suggests a relatively higher contribution of SN Ia than SNe II in this less massive dSph.

For the Fe-peak element Cr, the increasing trend with $[Fe/H]$ observed across all samples indicates the significant role of SN Ia in producing this element. In contrast, Co, belonging to the same category of Cr, shows a much more moderate contribution from this type of supernova, both in all systems and in the MW.

Finally, concerning the neutron-capture, specifically the s -process elements Ba and La, we observe a sharp enhancement ($\sim 0.4 - 0.5$ dex) in the Sgr abundances compared to those in the MW. This is particularly notable for La, which is synthesised both in s - and r -process ([Bisterzo et al. 2014](#)). This feature has been previously noted and discussed as a clear indication of substantial AGB star contributions ([McWilliam et al. 2013](#); [Hasselquist et al. 2017, 2021](#); [Sestito et al. 2024b](#)) with respect to the MW. In the trends of the s -process elements, Sgr bears similarities to Fornax and the LMC, but it differs from GES and, to an even greater extent, from Sculptor. This difference may be attributable to metallicity effects. Indeed, at lower $[Fe/H]$, heavy element production is dominated by the r -process, while AGB stars (usually with $M < 4M_{\odot}$) contribute to s -process enrichment with a time delay of ~ 1 Gyr after the onset of the star formation activity ([Travaglio et al. 2004](#)). This can also suggest that GES and Sculptor show less extended star formation with respect to the other systems. The r -process level of Sgr seems to be comparable with the one of the MW but less enhanced than GES (~ 0.3 dex difference) and Fornax and LMC (~ 0.2 and ~ 0.1 dex respectively). Despite the limited data points we have for Sgr, we observe a clear downward trend in metallicity, similar to what is seen in the LMC and Fornax in comparable metallicity regimes. This suggests that these systems experienced prolonged periods of star formation, allowing AGB stars to significantly contribute to their chemical enrichment ([Letarte et al. 2010](#); [Carrillo et al. 2022](#)). Their masses were likely sufficient to retain gas, which was crucial for reaching the stage where AGB stars could play a substantial role in the enrichment process. In contrast, GES appears to have had a shorter star formation process (extended up to ~ 3 Gyr, [Das et al. 2020](#); [Ernandes et al. 2024](#)) that ceased before AGB stars could influence its chemical composition, likely due to its early accretion into the MW, as also suggested by recent studies (e.g., [Hasselquist et al. \(2021\)](#); [Ernandes et al. \(2024\)](#)). Within this perspective, the gradual enrichment observed in the dSph Sgr, Fornax, Sculptor or the massive LMC contrasts with the more exotic and episodic events that could account for an earlier and higher levels of r -process pollution seen in GES. Sculptor exhibits only a mild enhancement in Ba, which may suggest that during its estimated 7 Gyr of SFH ([de Boer et al. 2012a](#)), metal-poor AGB stars

had sufficient time to enrich the system, albeit to a lower level compared to Sgr or Fornax.

4.4. The chemical enrichment histories revealed with heavy-elements

In Fig. 11 we explore five different chemical planes involving $[n/\alpha]$ -elements and $[s/r]$ -elements for Sgr, comparing them with the other satellites discussed in Sect. 4.3. These chemical ratios are widely used in the Galactic Archaeology field as their decoupled origins are known to help trace the chemical evolution of the galaxy and provide clues to the various nucleosynthesis channels which have chemically enriched the system (e.g., [Snedden et al. 2008](#); [Nomoto et al. 2013](#); [Skúladóttir et al. 2019](#); [Reichert et al. 2020](#)).

The $[Eu/Mg]$ ratio as a function of metallicity is a good proxy for estimating the contribution of delayed sources to Eu production, alongside prompt nucleosynthetic channels like SNe II, which produce both Eu and Mg ([Côté et al. 2018](#); [Skúladóttir & Salvadori 2020](#)). As previously mentioned, these delayed sources have been linked to exotic scenarios involving NSMs or MR-SNe ([Côté et al. 2018](#); [Skúladóttir & Salvadori 2020](#)). [Sestito et al. \(2024b\)](#) interpreted the spread observed in $[Eu/Mg]$ within a more limited metallicity range as evidence of a partial contribution from delayed r -process sources, such as compact binary mergers. However, it is important to consider that these scenarios may not necessarily be responsible for the increasing $[Eu/Mg]$ trend. It is possible that in dSphs with prolonged SFHs, another source contributing to the rise in r -process levels could involve massive stars formed during a later period of star formation, while gas was still available in the galaxy ([Reichert et al. 2021](#)).

In Sgr the $[Eu/Mg]$ trend and super-solar enrichment has been previously suggested in the literature ([McWilliam et al. 2013](#); [Skúladóttir et al. 2019](#); [Skúladóttir & Salvadori 2020](#); [Reichert et al. 2020](#)) and analyzed by [Sestito et al. \(2024b\)](#) in the VMP regime of Sgr. Our PIGS-HR-Sgr data confirm an increase in $[Eu/Mg]$, which appears to be consistent with more massive systems that experienced prolonged star formation SFHs, though much less pronounced than in Fornax and LMC. Similarly, Sculptor, despite occupying a different metallicity regime, exhibits a comparable rise in $[Eu/Mg]$ to that observed in Sgr. While both systems likely experienced star SFHs long enough to allow delayed r -process sources—estimated by [Skúladóttir & Salvadori \(2020\)](#) to require $\gtrsim 4$ Gyr to enrich the ISM, attributing this enrichment solely to such exotic scenarios is not straightforward.

For GES, although the data from [Carrillo et al. \(2022\)](#) do not reveal a clear trend, the work of [Matsuno et al. \(2021\)](#) identified an r -process enrichment linked to NSMs with a delay time contribution. However, compared to systems like Sgr, Fornax, and the LMC, [Matsuno et al. \(2021\)](#) highlighted a distinct difference in the extent of this r -enrichment. In fact, the weaker increasing trends in this chemical space for GES may suggest a shorter early star formation period, which is thought to be truncated $\sim 9-10$ Gyr following ~ 3 Gyr of star formation ([Belokurov et al. 2020](#); [Das et al. 2020](#); [Giribaldi & Smiljanic 2023](#); [Ernandes et al. 2024](#)). This would imply that a source with a delay time of about 4 Gyr, as proposed by [Skúladóttir & Salvadori \(2020\)](#), may not have had sufficient time to enrich GES as significantly as other massive systems (Sgr, Fornax, LMC).




Fig. 11. Different chemical combinations of n -capture elements (La, Ba, Eu) and α -element (Mg) for Sgr, compared to the systems shown in Fig 10 (illustrated with the same colors). In the top row, the chemical ratios are examined as a function of metallicity, while in the bottom row as a function of Mg abundance.

Alternatively, it is more likely that the violently truncated star formation in GES prevented a more prolonged SFH with subsequent star formation episodes which would rise the r -process level in the galaxy.

McWilliam et al. (2013), in explaining the [Eu/Mg] trend in Sgr, suggested that a top-light IMF, lacking the more massive SNe II that contribute hydrostatic elements like Mg to the ISM, played a role. From this perspective, the difference observed in GES could also suggest that CC-SNe continued to enrich this system in Mg to higher [Fe/H] values, unlike what happened in Sgr. This implies that the IMF of GES may have been richer in more massive stars, but that star formation was quenched earlier due to its violent infall into the MW.

Without need of advocating to the IMF, the differences between GES and Sgr can be explained by the fact that this latter might have had more time to continue forming stars before its infall into the MW, allowing for a later star forming episodes to increase its Eu levels. It is also possible that gas stripping was less effective in the central regions of the main body of Sgr, preventing a complete quenching of star formation. This could have left residual gas that was exhausted only later, after the galaxy's first pericentric passage (~ 5 Gyr Ruiz-Lara et al. 2020). Moreover, due to the spread in the Eu measurements in our analysis, it is not possible to rule out the contribution to the r -process enrichment from later massive stars and their subsequent SNe

explosions. Disentangling these scenarios in Sgr is challenging. However, considering the overall analysis, we believe that rise in the [Eu/Mg] panel most likely reflects the extended SFH of Sgr, characterized by multiple star-forming episodes that enriched the ISM in both r - and α -elements without producing a pronounced [Eu/Mg] trend. We also examined the Eu abundances as a function of age, but the resulting trend did not provide a clear timescale to constrain the timing of Eu enrichment.

The second and third panels of the top row in Fig. 11 provide valuable diagnostics for examining the contribution of the s -process channel. As noted in the study by Sestito et al. (2024b), Sgr exhibits an increasing trend of approximately 1 dex in [Ba/Eu] versus [Fe/H], a pattern also observed in the [La/Eu] ratio. This upward trend suggests a substantial contribution from s -process production channels relative to the r -process ($\sim 96\%$ of Eu is produced through r -process Travaglio et al., 2004, Bisterzo et al., 2014). This reinforces evidence for a significant AGB star contribution in Sgr and indicates a combination of ($r + s$) formation sites, as also proposed by Hansen et al. (2018) for the metal-poor sample analyzed in their study. This tendency is also observed in the LMC and Fornax, though it is less pronounced and more gradual in GES. This suggests that the progenitor of GES was dominated by different nucleosynthetic processes compared to these other systems. By combining the information from these panels with that

of the first, we can conclude that the dSphs Sgr, Fornax, and the LMC show enrichment in heavy elements from both prompt and delayed sources of r -process elements, balanced by a delayed onset of AGB star contributions (as found by Letarte et al. 2010; Lemasle et al. 2014; Hansen et al. 2018; Reichert et al. 2020), beginning around $[\text{Fe}/\text{H}] \sim -1.2$ for Sgr, similar to its α -knee, indicating a similar time delay (the same behaviour is shown by Sculptor where this phenomenon happens around -2.0 Hill et al. 2019). In contrast, GES experienced intense early enrichment in Eu with a smaller s -process contribution (as also evinced by Matsuono et al. 2021), likely due to a shorter and more abruptly quenched SFH. Finally, Sculptor exhibits a very high initial Eu abundance, which might be caused by compact merger events, followed by a steep rise in Ba at higher metallicities which can witness the s -enrichment from metal-poor AGBs.

The trends illustrated in the last two panels of the bottom rows serve as further tools for assessing the contribution of AGB stars in the closed system of Sgr. The rise in $[\text{La}/\text{Mg}]$ of ~ 2 dex excludes a scenario where only the yields from SNe II contribute to the enrichment of the system, as this would result in a flat $[\text{La}/\text{Mg}]$ or $[\text{Ba}/\text{Mg}]$ distribution (e.g., Pignatari et al. 2008; Cowan et al. 2021). This is because the s -processes occurring in AGB stars increase the levels of La and Ba relative to Mg, which is primarily produced by SNe II (Pignatari et al. 2008; Cristallo et al. 2015). Typically, this pattern is associated with the mass of these systems, which is high enough to retain their gas, allowing them to evolve to the point where AGB stars become the dominant contributors. As shown in Fig 11 the observations in Fornax and the LMC exhibit similar behaviors, which are supported by the literature (Pompéia et al. 2008; Letarte et al. 2010; Reichert et al. 2020). Sculptor also demonstrates a rapid increase in the $[\text{Ba}/\text{Mg}]$ vs. $[\text{Mg}/\text{H}]$ relation; however, the behavior for La is less distinct due to its more even distribution between s - and r -processes (with 75% of La produced by the s -process Bisterzo et al., 2014) contrary to Ba (85 % produced by s -process Bisterzo et al., 2014). Hill et al. (2019) also discussed the potential biases in La measurements, as they are derived from weak lines. The GES data show a similar trend, where the increase in Ba may indicate an additional contribution from s -process elements occurring on timescales comparable to the onset SN Ia. Ernandes et al. (2024) found similar results and concluded that the absence of an increasing $[\text{Ba}/\text{Mg}]$ ratio at higher metallicities (in our case, $[\text{Mg}/\text{H}]$) is indicative of a significant halt in star formation in GES, which can also lead to a lack of substantial s -process contribution as metallicity increases. Due to the limited Ba measurements in Sgr, it is challenging to discern this difference in their trends. The low ratios of $[\text{La}/\text{Mg}]$ and $[\text{Ba}/\text{Mg}]$ at lower values of $[\text{Mg}/\text{H}]$ in Sgr reflect a period of strong initial star formation in the galaxy, characterized by a rapid contribution from CCSNe (Mg) compared to the delayed yields from AGB stars (Ba). Hence, even if CCSNe occur at later times from the remaining gas in Sgr (before its complete quenching), contributing to delayed r -process enrichment and α -elements, the stronger contribution of s -process elements from AGB stars counteracts this effect.

Without measurements of the ratio of heavy (hs , e.g., La, Ba) to light (ls , e.g., Sr, Y, Zr) n -capture elements, constraining the mass of the AGB stars that enriched the ISM of Sgr becomes challenging. The $[(ls)/(hs)]$ ratio is




Fig. 12. Age-metallicity relation for the Sgr stellar sample (blue circles). The pink dots indicate five GCs associated to the Sgr streams reported by Garro et al. (2021), while the orange pentagons represent both GCs and the Sgr field population derived with isochrones fitting by Layden & Sarajedini (2000).

crucial for this purpose (Travaglio et al. 2004; Pignatari et al. 2010). However, based on the findings of McWilliam et al. (2013), MP and low-mass AGB ($\lesssim 2M_{\odot}$) are invoked to explain the chemical properties of Sgr. This view contrasts with Hansen et al. (2018), who find that their best fit for the chemical abundances of their VMP Sgr sample is provided by intermediate-mass AGBs ($\sim 5M_{\odot}$).

4.5. Age-metallicity relation

The SFH of Sgr has been deeply investigated over the years (e.g. Bellazzini et al. 1999a,b; Ruiz-Lara et al. 2020). The star formation episodes have been dated and compared them with the kinematic and orbital history of this dwarf galaxy, which has been modeled in various works (e.g. Peñarrubia et al. 2010; Vasiliev & Belokurov 2020; Davies et al. 2024). In this context, accurate age estimates are crucial for constructing a detailed evolutionary history of the system. However, deriving this information is not without its challenges due to the indirect methods and techniques used for age determination. Using isochrone techniques in combination with deep-band photometry, Layden & Sarajedini (2000) provided a detailed SFH for both M54 and Sgr field stars. Their study identified multiple episodes of star formation over a span of 0.5 to 14 Gyr, concluding a protracted SFH, a finding later validated by Siegel et al. (2007). Through photometric data from the HST and employing isochrone and CMD fitting, Siegel et al. (2007) identified several stellar populations within the main core, ranging from an old MP population (~ 10 -12 Gyr) to a very young population of only a few Myr. Stellar clusters are often used to determine ages more easily.

For the PIGS-HR-Sgr sample we derived ages following Povick et al. in prep. Using a finely spaced grid of PARSEC (Girardi et al. 2002; Bressan et al. 2012; Marigo et al. 2017) isochrones, all points within 7σ of each star are selected in a 10-dimensional parameter space, which includes opti-

cal/NIR multiband photometry from Gaia DR3 (Gaia Collaboration et al. 2016b, 2022) and 2MASS (Skrutskie et al. 2006) as well as T_{eff} , $[M/H]$, and $\log g$. For each isochrone point, the χ^2 value is calculated incorporating an additional term that functions as a prior based on the Kroupa IMF (Kroupa 2001, 2002). The age, extinction, and mass values for each star are then obtained with a weighted mean of the isochrone points, with the weights determined by the modified χ^2 values.

When deriving ages for the stars, their $[\alpha/\text{Fe}]$ values were fixed to solar. It was found that incorporating the measured $[\text{Mg}/\text{Fe}]$ as a proxy for overall $[\alpha/\text{Fe}]$ resulted in many more spurious ages, particularly for α -enhanced stars, leading to ages less consistent with a dominant population of ~ 10 Gyr as found by Layden & Sarajedini (1997). The uncertainties in the abundance measurements also contribute to the scatter observed when using α -abundances. Furthermore, the vast majority of stars in this sample have $[\text{Mg}/\text{Fe}] \sim 0.0$ dex, so assuming solar $[\alpha/\text{Fe}]$ was determined to be reasonable.

The resulting AMR for our targets (blue circles) appears in Fig. 12. To guide the eyes, we also present the age estimates from Layden & Sarajedini (2000) for the four GCs associated with the Sgr remnant (NGC 6715/M54, Terzan 7, Terzan 8, and Arp 2) as well as for three field populations. Additionally, we report the ages inferred by Garro et al. (2021) for five more GCs recently identified as part of the Sgr streams (Bellazzini et al. 2020). Despite the large scatter in our AMR, a general agreement in the trends can still be observed across different studies. The PIGS-HR-Sgr AMR is fairly flat for ages between 6 and 10 Gyr, with a more pronounced increase in metallicity evident among younger stars. The peak at ~ 6 -8 Gyr in our data corresponds to an intermediate-age population with average $[\text{Fe}/\text{H}] \sim -0.85$. This population has been previously identified as having undergone intense and rapid star formation activity and constitutes the predominant population (Bellazzini et al. 1999b; Siegel et al. 2007). Stars in this population likely formed before Sgr's first infall into the MW's potential (~ 5 Gyr ago; Ruiz-Lara et al. 2020), an event that may have triggered a burst of star formation in the MW (Ruiz-Lara et al. 2020) and likely led to the subsequent quenching of star formation in Sgr due to gas stripping, along with a natural decline in the dwarf's star formation history.

However, as noted in other studies, residual star formation activity can be identified with stars younger than 6 Gyr. These residual star-forming episodes could be related to interactions with the MW. While ram-pressure stripping likely removed gas from Sgr, it may also have compressed the remaining gas, triggering new star formation episodes—a scenario explored in the recent work by Yang et al. (2024) in other dSphs, but not in Sgr. Supporting evidence for these later star formation activities comes from our identification of a few younger stars with ages of ~ 4 Gyr, likely reflecting the lower metallicity limit of our PIGS-HR-Sgr sample ($[\text{Fe}/\text{H}] \gtrsim -0.5$ dex). The presence of younger populations with ages of ~ 3 -4 Gyr further supports the idea that, despite gas quenching, Sgr retained enough gas in its central regions to form stars from its remaining gas reservoir.

Looking at the older stars (> 8 Gyr) there is considerable scatter in metallicity between ~ 8 - 10 Gyr, which corresponds to the age range with the largest uncertainties

(~ 2.5 - 3 Gyr). We can assume that this old stellar populations may have potentially originated from the gas of the nuclear center of the galaxy and later displaced during Sgr's orbital history. Three stars with age > 12 Gyr show surprisingly high metallicity values (i.e. $-0.5 \lesssim [\text{Fe}/\text{H}] \lesssim -1.0$) and low age uncertainties ($\lesssim 0.3$ Gyr). We reviewed the input atmospheric parameters and metallicities and found no anomalies. It appears that the three stars are approaching the isochrone grid edge in age. Additionally, the density of isochrone points in metallicity, T_{eff} , and $\log g$ space drops near the three outliers. These two factors most likely explain what is happening with these three stars.

Studies such as the ones from Revaz et al. (2009); Revaz & Jablonka (2012) demonstrate that the star formation in massive dSphs (above $\sim 3 \times 10^8 M_{\odot}$) is naturally linked to the mass of the system. In such cases, star formation occurs in more homogeneous and less complex scenarios, where external factors such as gas removal or ram pressure are not necessarily required or would leave a noticeable impact if present. This contrasts with lower-mass dwarf galaxies, which exhibit more diverse SFHs that are strongly influenced by quenching mechanisms, including SNe feedback or external processes. For Sgr, its mass suggests a reduced impact from such mechanisms. Indeed, despite its merging history with the MW, the AMR observed in our central PIGS-HR-Sgr sample indicates that star formation has not been entirely halted. Consequently, the overall shape of its AMR likely reflects its intrinsic star formation history, which is closely linked to its initial mass.

Lastly, due to the scatter in ages around ~ 8 -10 Gyr, it is not trivial to determine the age at which our estimate of the knee occurred. In the study by de Boer et al. (2014) on the Sgr stellar stream, the knee was constrained to an age of ~ 11 Gyr. Based on our value of $[\text{Fe}/\text{H}]_{\text{knee}} = -1.05$, the knee appears to occur at around ~ 8 -9 Gyr, which is slightly later than the age found by de Boer et al. (2014). This discrepancy is reasonable, as they reported a lower metallicity for the knee, $[\text{Fe}/\text{H}] = -1.27$ dex, which was calculated for stream stars using different methodologies. Nevertheless, considering the time scales in both studies, this would place the occurrence of the knee around 1-3 Gyr after the formation of the system, as also found in de Boer et al. (2015). This timescale is consistent with the delay in the onset of Type Ia supernovae (Strolger et al. 2004; Sullivan et al. 2006; Wiseman et al. 2021).

5. Summary & conclusions

The history and chemical evolution of the interacting Sagittarius dwarf galaxy has been studied using both photometric and spectroscopic techniques. In this work, we have assembled a large high-resolution dataset, providing the most comprehensive set of chemical species measured for giants in the core of Sgr. Thanks to a prior selection based on metallicity using PIGS photometry, we conducted a spectroscopic follow-up of 111 Sagittarius stars with FLAMES/GIRAFFE. By inferring 14 chemical elements with average uncertainty of ~ 0.09 dex, we investigated the chemical enrichment history of this galaxy, placing it in the context of its interaction with the MW by comparing it to other satellites and structures.

By inferring RVs for the entire stellar sample (see Fig. 4) and finding a good match with the distribution from the literature, we can confirm the Sgr membership selection

performed using *Gaia* astrometry. Although we observe a slightly higher proportion of MR stars, particularly with $[\text{Fe}/\text{H}] > -1.0$ (see Fig. 2), the resulting spectroscopic metallicities confirm the effectiveness of the PIGS photometry in pre-selecting candidates based on their iron content. This leads to a $[\text{Fe}/\text{H}]$ coverage ranging from -2.13 to -0.35 , peaking at $[\text{Fe}/\text{H}] \sim -0.68$.

We compared the various $[\text{X}/\text{Fe}]$ trends explored in our analysis with different intermediate and high-resolution samples of the Sgr core from the literature (Fig. 7). Our Sgr sample aligns with the expected Galactic chemical trends: α -elements decrease with increasing metallicity, while n -capture elements exhibit the opposite trend, reflecting their distinct astrophysical production sites. This pattern is also evident for the two Fe-peak elements, whose enhancements, though at different levels, are influenced by SN Ia and therefore show different trends with $[\text{Fe}/\text{H}]$.

This chemical investigation was extended to compare the chemical trends of the MW with those of its most massive satellite systems (Fig. 10). Consistent with the literature (Carrillo et al. 2022), we found that the satellites are deficient in hydrostatic elements relative to the MW, a difference potentially linked and to a less efficient early star formation period, or connected as in McWilliam et al. (2013); Hasselquist et al. (2017) to a top-light IMF that lacks the most massive stars. However, we acknowledge that the observed differences in elements such as Mg and Ca are challenging to disentangle, as they may arise from the system's star formation rate or the yields of SN Ia. Further investigation, including the use of chemical evolution models, is therefore needed to confirm the true nature of Sgr's IMF.

Trends in Fe-peak and odd-Z elements appear similar among the more massive systems (Sgr, LMC, GES, and Fornax) and the MW. However, an insightful distinction arises with the n -capture elements, especially the s -process La and Ba, which exhibit a marked enhancement in Sgr compared to the MW—a trend that aligns with Fornax and the LMC but diverges from the GES. The latter also shows an enhanced r -process abundance compared to the MW and the other systems, among which Fornax and the LMC follow GES. Taking into account the $[\text{Fe}/\text{H}]$ dependence, which affects the trends of the systems that cover different metallicity ranges, it is yet possible to conclude that the dSph Sgr, Fornax, and the LMC show a larger contribution from AGB stars (possibly compared to SNe) relative to the MW. This result has been found in many studies concerning dSph galaxies (e.g., McWilliam et al. 2013; Letarte et al. 2010; Hasselquist et al. 2021). This is less true for GES, which is expected since this system only evolved over the course of 2–3 Gyr, a time frame insufficient for AGB stars to enrich the system. However, AGB contributions have been found to be part of its evolution (Hill et al. 2019; Erandes et al. 2024) hence, it is possible that the lower values of Ba and La reflect a metallicity effect on the AGB yields.

A more in depth investigation of the chemical enrichment histories of these MW satellites was carried throughout the exploration of heavy elements, as shown in Fig. 11. Sgr and Sculptor show a mild increase in the $[\text{Eu}/\text{Mg}]$ trend, much less pronounced than in Fornax and the LMC. A stronger increase in $[\text{Eu}/\text{Mg}]$ may suggest extended SFHs with contributions from delayed r -process sources, such as NSM (Skúladóttir & Salvadori 2020). These appear to be absent or not prominent in Sculptor and Sgr, and occur

on timescales of $\lesssim 4$ Gyr in GES, according to Matsuno et al. (2021). The most evident difference is between systems with long SFHs, such as Sgr (or Fornax and the LMC), compared to GES, which had halted and truncated star formation but, by contrast, an IMF rich in massive stars that continued to enrich in Mg to higher metallicities.

Looking at the $[s/r]$ ratio vs $[\text{Fe}/\text{H}]$ (i.e. $[\text{La}/\text{Eu}]$ and $[\text{Ba}/\text{Eu}]$) Sgr shows an increasing trend of ≈ 1 dex, slightly higher than those of Fornax and the LMC, suggesting enrichment in heavy elements due to a delayed onset of AGB stars, balancing both prompt and delayed sources of r -process elements (as noted in previous studies, e.g., Letarte et al. 2010; Hansen et al. 2018). Both GES and Sculptor exhibit a high level of r -process enrichment; however, it remains steady in GES, indicating a halted SFH, while it increases in Sculptor, reflecting ongoing AGB contributions. This information was further supported by the study of $[s/\alpha]$ ratios, which confirm a significant contribution from AGB stars in Sgr beginning at higher $[\text{Mg}/\text{H}]$. This suggests that early Mg production was driven by CCSNe, later superseded by delayed yields from AGB stars, a pattern also observed in the LMC and Fornax. In conclusion, even without detailed chemical evolution models, these chemical trends related to n -capture serve as powerful diagnostics for understanding the evolution of these systems, being closely tied to their SFH, as evinced in our analysis.

We derived $[\text{Fe}/\text{H}]_{\alpha\text{-knee}} = -1.05$ for Sgr (see Fig. 8), incorporating literature data sets (e.g., Monaco et al. 2005; Sbordone et al. 2007; Mucciarelli et al. 2017). Despite a non-homogeneous metallicity coverage, the occurrence of the knee suggests that Sgr experienced slower early enrichment compared to the MW, but was still able to utilize its gas to continue star formation to higher $[\text{Fe}/\text{H}]$ values than the LMC, which followed an isolated evolution. To answer a question already raised by Sestito et al. (2024b), no strong evidence of SN Ia was found at $[\text{Fe}/\text{H}] \lesssim -1.0$. Relying on the measured α -abundances, we examine the ratios between hydrostatic (Mg) and explosive elements (Ca and Ti) in Fig. 9 and propose the presence of HNe enriching the more metal-poor stars in Sgr, as also suggested by Sestito et al. (2024b).

Finally, the AMR presented in Fig. 12 aligns with the photometric studies of the SFH by Layden & Sarajedini (2000); Siegel et al. (2007). Despite significant scatter, we identify the main populations within Sgr: an older, metal-poor population aged 8–10 Gyr, which is also the period during which the knee can be placed. This is followed by a dominant intermediate-age population peaking around 6–8 Gyr, likely formed before Sgr's first infall onto the MW during a strong and rapid star formation period. Following this, Sgr continued to form stars by compressing the remaining gas, as evidenced by stars with ages $\sim 3 - 5$ Gyr, despite the effects of gas stripping.

To summarize, we propose that Sgr exhibits a chemical evolution characteristic of a massive dwarf galaxy. Its stellar populations, sampled from the galaxy's central regions and thus largely unaffected by contamination from MW gas, reflect Sgr's independent chemical enrichment history. This history includes multiple episodes of star formation, peaking within the first few billion years after its formation and subsequently declining. The results for heavier elements reveal an enrichment in n -capture elements, indicative of a prolonged star formation history, further supported by the AMR. The various chemical trends suggest

contributions from different enrichment sources, including various types of SNe, AGBs, while showing no evidence of sequences attributable to MW gas infall. When comparing Sgr with other significant satellites of the MW and its halo, we observe that many of its chemical trends resemble those of dwarf galaxies capable of sustaining longer star.

Despite the diagnostic power of chemical abundances, in future work, we aim to use chemical evolution models to further disentangle the history of the core of Sgr and constrain the contributions of SNe across different timescales. With precise and accurate models, the chemical evolution and SFH can be better connected and used to trace the precise history of this galaxy, from its progenitor through the long interaction period with our Galaxy.

Additionally, upcoming spectroscopic surveys like 4DWARFS (Skúladóttir et al. 2023) or MOONS (Gonzalez et al. 2020) will supply more data to enhance our study of the Sgr stellar populations within both the core and the stream. These surveys will allow us to investigate their common progenitor, trace the SFH, and conduct chemodynamical studies, expanding our understanding of Sgr as a satellite of the MW.

Acknowledgements. SV thanks ANID (Beca Doctorado Nacional, folio 21220489) and Universidad Diego Portales for the financial support provided. SV and PJ acknowledge the Millennium Nucleus ERIS (ERIS NCN2021017) and FONDECYT (Regular number 1231057) for the funding. A. R. A. acknowledges support from DICYT through grant 062319RA. ES acknowledges funding through VIDI grant "Pushing Galactic Archaeology to its limits" (with project number VI.Vidi.193.093) which is funded by the Dutch Research Council (NWO). This research has been partially funded from a Spinoza award by NWO (SPI 78-411). This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

Author contribution statement. SV led the analysis and various discussions in this work, contributed to writing most of the draft, and created all the figures presented in this work. AAR assisted with data reduction, analysis, and discussions related to the interpretation of the results. PJ and FS contributed significantly to the interpretation of the draft and the discussion of the results. JP helped revise the draft and provided the results for the stellar ages. AAA contributed the PIGS photometric data, which were used for target selection and largely contributed to the scientific discussion. VH and EF assisted with revising the draft and preparing the proposal. PJ, NM, ES, and DA provided insightful scientific and editorial comments on the manuscript.

References

Abdurro'uf, Accetta, K., Aerts, C., et al. 2022, *ApJS*, 259, 35
 Alvarez, R. & Plez, B. 1998, *A&A*, 330, 1109
 An, Z., Walker, M. G., & Pace, A. B. 2024, *MNRAS*[arXiv:2404.16184]
 Andrae, R., Fouesneau, M., Creevey, O., et al. 2018, *A&A*, 616, A8
 Ardern-Arentsen, A., Monari, G., Queiroz, A. B. A., et al. 2024, *MNRAS*, 530, 3391
 Arentsen, A., Starkenburg, E., Martin, N. F., et al. 2020a, *MNRAS*, 496, 4964
 Arentsen, A., Starkenburg, E., Martin, N. F., et al. 2020b, *MNRAS*, 491, L11
 Bandyopadhyay, A., Ezzeddine, R., Allende Prieto, C., et al. 2024, arXiv e-prints, arXiv:2408.03731
 Battistini, C. & Bensby, T. 2015, *A&A*, 577, A9
 Bellazzini, M., Ferraro, F. R., & Buonanno, R. 1999a, *MNRAS*, 304, 633

Bellazzini, M., Ferraro, F. R., & Buonanno, R. 1999b, *MNRAS*, 307, 619
 Bellazzini, M., Ibata, R., Malhan, K., et al. 2020, *A&A*, 636, A107
 Belokurov, V., Erkal, D., Evans, N. W., Koposov, S. E., & Deason, A. J. 2018, *MNRAS*, 478, 611
 Belokurov, V., Sanders, J. L., Fattahi, A., et al. 2020, *MNRAS*, 494, 3880
 Bergemann, M. 2011, *MNRAS*, 413, 2184
 Bergemann, M. & Cescutti, G. 2010, *A&A*, 522, A9
 Bergemann, M., Kudritzki, R.-P., Gazak, Z., Davies, B., & Plez, B. 2015, *ApJ*, 804, 113
 Bergemann, M., Kudritzki, R.-P., Würl, M., et al. 2013, *ApJ*, 764, 115
 Bergemann, M., Pickering, J. C., & Gehren, T. 2010, *MNRAS*, 401, 1334
 Besla, G., Kallivayalil, N., Hernquist, L., et al. 2007, *ApJ*, 668, 949
 Bettinelli, M., Hidalgo, S. L., Cassisi, S., et al. 2019, *MNRAS*, 487, 5862
 Bisterzo, S., Travaglio, C., Gallino, R., Wiescher, M., & Käppeler, F. 2014, *ApJ*, 787, 10
 Blanco-Cuaresma, S. 2019, *MNRAS*, 486, 2075
 Blanco-Cuaresma, S., Soubiran, C., Heiter, U., & Jofré, P. 2014a, *A&A*, 569, A111
 Blanco-Cuaresma, S., Soubiran, C., Jofré, P., & Heiter, U. 2014b, *A&A*, 566, A98
 Bland-Hawthorn, J. & Gerhard, O. 2016, *ARA&A*, 54, 529
 Bonifacio, P., Hill, V., Molaro, P., et al. 2000, *A&A*, 359, 663
 Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, 427, 127
 Buder, S., Sharma, S., Kos, J., et al. 2021, *MNRAS*, 506, 150
 Burbidge, E. M., Burbidge, G. R., Fowler, W. A., & Hoyle, F. 1957, *Reviews of Modern Physics*, 29, 547
 Busso, M., Gallino, R., & Wasserburg, G. J. 1999, *ARA&A*, 37, 239
 Carlin, J. L., Sheffield, A. A., Cunha, K., & Smith, V. V. 2018, *ApJ*, 859, L10
 Carretta, E., Bragaglia, A., Gratton, R. G., et al. 2010, *A&A*, 520, A95
 Carrillo, A., Hawkins, K., Jofré, P., et al. 2022, *MNRAS*, 513, 1557
 Childs, W., Poulsen, O., Goodman, L., & Crosswhite, H. 1979, *Physical Review A*, 19, 168
 Chiti, A. & Frebel, A. 2019, *ApJ*, 875, 112
 Chiti, A., Hansen, K. Y., & Frebel, A. 2020, *ApJ*, 901, 164
 Chiti, A., Mardini, M., Limberg, G., et al. 2024, *Nature Astronomy*, 8, 637
 Cochrane, E. C. A., Benton, D. M., Forest, D. H., & Griffith, J. A. R. 1998, *Journal of Physics B Atomic Molecular Physics*, 31, 2203
 Côté, B., Fryer, C. L., Belczynski, K., et al. 2018, *ApJ*, 855, 99
 Cowan, J. J. & Rose, W. K. 1977, *ApJ*, 212, 149
 Cowan, J. J., Sneden, C., Lawler, J. E., et al. 2021, *Reviews of Modern Physics*, 93, 015002
 Cristallo, S., Straniero, O., Piersanti, L., & Gobrecht, D. 2015, *ApJS*, 219, 40
 Das, P., Hawkins, K., & Jofré, P. 2020, *MNRAS*, 493, 5195
 Davies, E. Y., Monty, S., Belokurov, V., & Dillamore, A. M. 2024, *MNRAS*, 529, 772
 de Boer, T. J. L., Belokurov, V., Beers, T. C., & Lee, Y. S. 2014, *MNRAS*, 443, 658
 de Boer, T. J. L., Belokurov, V., & Koposov, S. 2015, *MNRAS*, 451, 3489
 de Boer, T. J. L., Tolstoy, E., Hill, V., et al. 2012a, *A&A*, 539, A103
 de Boer, T. J. L., Tolstoy, E., Hill, V., et al. 2012b, *A&A*, 544, A73
 del Pino, A., Fardal, M. A., van der Marel, R. P., et al. 2021, *ApJ*, 908, 244
 Eitner, P., Bergemann, M., & Larsen, S. 2019, *A&A*, 627, A40
 Erkal, D., Belokurov, V., Laporte, C. F. P., et al. 2019, *MNRAS*, 487, 2685
 Ernandes, H., Feuillet, D., Feltzing, S., & Skúladóttir, Á. 2024, arXiv e-prints, arXiv:2405.13641
 Escorza, A., Karinkuzhi, D., Jorissen, A., et al. 2019, *A&A*, 626, A128
 Evans, D. W., Riello, M., De Angeli, F., et al. 2018, *A&A*, 616, A4
 Ezzeddine, R., Rasmussen, K., Frebel, A., et al. 2020, *ApJ*, 898, 150
 Fernandes, L., Mason, A. C., Horta, D., et al. 2023, *MNRAS*, 519, 3611
 Feuillet, D. K., Feltzing, S., Sahlholdt, C. L., & Casagrande, L. 2020, *MNRAS*, 497, 109
 Frebel, A., Casey, A. R., Jacobson, H. R., & Yu, Q. 2013, *ApJ*, 769, 57
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2021, *A&A*, 649, A1
 Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016a, *A&A*, 595, A1

- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016b, *A&A*, 595, A1
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, *A&A*, 674, A1
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2022, arXiv e-prints, arXiv:2208.00211
- Garro, E. R., Minniti, D., Gómez, M., & Alonso-García, J. 2021, *A&A*, 654, A23
- Girardi, L., Bertelli, G., Bressan, A., et al. 2002, *A&A*, 391, 195
- Giribaldi, R. E. & Smiljanic, R. 2023, *A&A*, 673, A18
- Gonzalez, O. A., Mucciarelli, A., Origlia, L., et al. 2020, *The Messenger*, 180, 18
- Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951
- Hansen, C. J., El-Souri, M., Monaco, L., et al. 2018, *ApJ*, 855, 83
- Hasselquist, S., Hayes, C. R., Lian, J., et al. 2021, *ApJ*, 923, 172
- Hasselquist, S., Shetrone, M., Smith, V., et al. 2017, *ApJ*, 845, 162
- Hayes, C. R., Majewski, S. R., Hasselquist, S., et al. 2020, *ApJ*, 889, 63
- Hegedűs, V., Mészáros, S., Jofré, P., et al. 2023, *A&A*, 670, A107
- Heger, A. & Woosley, S. E. 2002, *ApJ*, 567, 532
- Heiter, U., Lind, K., Bergemann, M., et al. 2021, *A&A*, 645, A106
- Helmi, A., Babusiaux, C., Koppelman, H. H., et al. 2018, *Nature*, 563, 85
- Hidalgo, S. L., Pietrinferni, A., Cassisi, S., et al. 2018, *ApJ*, 856, 125
- Hill, V., Skúladóttir, Á., Tolstoy, E., et al. 2019, *A&A*, 626, A15
- Ibata, R., Lewis, G. F., Irwin, M., Totten, E., & Quinn, T. 2001, *ApJ*, 551, 294
- Ibata, R. A., Gilmore, G., & Irwin, M. J. 1994, *Nature*, 370, 194
- Jönsson, H., Holtzman, J. A., Allende Prieto, C., et al. 2020, *AJ*, 160, 120
- Jorissen, A., Boffin, H. M. J., Karinkuzhi, D., et al. 2019, *A&A*, 626, A127
- Käppeler, F., Gallino, R., Bisterzo, S., & Aoki, W. 2011, *Reviews of Modern Physics*, 83, 157
- Karakas, A. I. & Lattanzio, J. C. 2014, *Publications of the Astronomical Society of Australia*, 31, e030
- Karovicova, I., White, T. R., Nordlander, T., et al. 2020, *A&A*, 640, A25
- Kirby, E. N., Gilbert, K. M., Escala, I., et al. 2020, *AJ*, 159, 46
- Kobayashi, C., Leung, S.-C., & Nomoto, K. 2020, *ApJ*, 895, 138
- Kobayashi, C. & Nomoto, K. 2009, *ApJ*, 707, 1466
- Kobayashi, C., Umeda, H., Nomoto, K., Tominaga, N., & Ohkubo, T. 2006, *ApJ*, 653, 1145
- Korotin, S. A., Andrievsky, S. M., Hansen, C. J., et al. 2015, *A&A*, 581, A70
- Kratz, K. L., Farouqi, K., & Pfeiffer, B. 2007, *Progress in Particle and Nuclear Physics*, 59, 147
- Kroupa, P. 2001, *MNRAS*, 322, 231
- Kroupa, P. 2002, *Science*, 295, 82
- Law, D. R. & Majewski, S. R. 2010, *ApJ*, 714, 229
- Lawler, J. E., Bonvallet, G., & Sneden, C. 2001, *ApJ*, 556, 452
- Layden, A. C. & Sarajedini, A. 1997, *ApJ*, 486, L107
- Layden, A. C. & Sarajedini, A. 2000, *AJ*, 119, 1760
- Lee, T., Jeon, M., & Bromm, V. 2024, *MNRAS*, 527, 1257
- Lemasle, B., de Boer, T. J. L., Hill, V., et al. 2014, *A&A*, 572, A88
- Letarte, B., Hill, V., Tolstoy, E., et al. 2010, *A&A*, 523, A17
- Lewis, I. J., Cannon, R. D., Taylor, K., et al. 2002, *MNRAS*, 333, 279
- Limongi, M. & Chieffi, A. 2018, *The Astrophysical Journal Supplement Series*, 237, 13
- Lind, K., Nordlander, T., Wehrhahn, A., et al. 2022, *A&A*, 665, A33
- Łokas, E. L. 2009, *MNRAS*, 394, L102
- Mackereth, J. T. & Bovy, J. 2020, *MNRAS*, 492, 3631
- Mackereth, J. T., Schiavon, R. P., Pfeffer, J., et al. 2019, *MNRAS*, 482, 3426
- Majewski, S. R., Skrutskie, M. F., Weinberg, M. D., & Ostheimer, J. C. 2003, *ApJ*, 599, 1082
- Marigo, P., Girardi, L., Bressan, A., et al. 2008, *A&A*, 482, 883
- Marigo, P., Girardi, L., Bressan, A., et al. 2017, *ApJ*, 835, 77
- Martin, N. F., Starkenburg, E., Yuan, Z., et al. 2023, arXiv e-prints, arXiv:2308.01344
- Matsuno, T., Hirai, Y., Tarumi, Y., et al. 2021, *A&A*, 650, A110
- Matteucci, F. 2016, in *Journal of Physics Conference Series*, Vol. 703, *Journal of Physics Conference Series*, 012004
- Matteucci, F., Spitoni, E., Recchi, S., & Valiante, R. 2009, *A&A*, 501, 531
- McConnachie, A. W. 2012, *AJ*, 144, 4
- McWilliam, A. 1997, *Annual Review of Astronomy and Astrophysics*, 35, 503
- McWilliam, A., Piro, A. L., Badenes, C., & Bravo, E. 2018, *ApJ*, 857, 97
- McWilliam, A., Wallerstein, G., & Mottini, M. 2013, *ApJ*, 778, 149
- Minelli, A., Bellazzini, M., Mucciarelli, A., et al. 2023, *A&A*, 669, A54
- Monaco, L., Bellazzini, M., Bonifacio, P., et al. 2005, *A&A*, 441, 141
- Mucciarelli, A., Bellazzini, M., Ibata, R., et al. 2017, *A&A*, 605, A46
- Mucciarelli, A., Bellazzini, M., & Massari, D. 2021, *A&A*, 653, A90
- Mucciarelli, A. & Bonifacio, P. 2020, *A&A*, 640, A87
- Nidever, D. L., Hasselquist, S., Hayes, C. R., et al. 2020, *ApJ*, 895, 88
- Nomoto, K., Kobayashi, C., & Tominaga, N. 2013, *ARA&A*, 51, 457
- Osorio, Y., Allende Prieto, C., Hubeny, I., Mészáros, S., & Shetrone, M. 2020, *A&A*, 637, A80
- Pasquini, L., Avila, G., Blecha, A., et al. 2002, *The Messenger*, 110, 1
- Peñarrubia, J., Belokurov, V., Evans, N. W., et al. 2010, *MNRAS*, 408, L26
- Pickering, J. C. 1996, *ApJS*, 107, 811
- Pietrinferni, A., Salaris, M., Cassisi, S., et al. 2024, *MNRAS*, 527, 2065
- Pignatari, M., Gallino, R., Heil, M., et al. 2010, *ApJ*, 710, 1557
- Pignatari, M., Gallino, R., Meynet, G., et al. 2008, *ApJ*, 687, L95
- Plez, B. 2012, *Turbospectrum: Code for spectral synthesis*, *Astrophysics Source Code Library*, record ascl:1205.004
- Pompéia, L., Hill, V., Spite, M., et al. 2008, *A&A*, 480, 379
- Reichert, M., Hansen, C. J., & Arcones, A. 2021, *ApJ*, 912, 157
- Reichert, M., Hansen, C. J., Hanke, M., et al. 2020, *A&A*, 641, A127
- Revaz, Y. & Jablonka, P. 2012, *A&A*, 538, A82
- Revaz, Y., Jablonka, P., Sawala, T., et al. 2009, *A&A*, 501, 189
- Ruiz-Lara, T., Gallart, C., Bernard, E. J., & Cassisi, S. 2020, *Nature Astronomy*, 4, 965
- Saunders, W., Bridges, T., Gillingham, P., et al. 2004, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 5492, *Ground-based Instrumentation for Astronomy*, ed. A. F. M. Moorwood & M. Iye, 389–400
- Sbordone, L., Bonifacio, P., Buonanno, R., et al. 2007, *A&A*, 465, 815
- Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schultz, G. V. & Wiemer, W. 1975, *A&A*, 43, 133
- Searle, L. & Zinn, R. 1978, *ApJ*, 225, 357
- Seitzzahl, I. R., Cescutti, G., Röpkke, F. K., Ruiter, A. J., & Pakmor, R. 2013, *A&A*, 559, L5
- Sestito, F., Arden-Arentsen, A., Vitali, S., et al. 2024a, *A&A*, 690, A333
- Sestito, F., Venn, K. A., Arentsen, A., et al. 2023, *MNRAS*, 518, 4557
- Sestito, F., Vitali, S., Jofre, P., et al. 2024b, *A&A*, 689, A201
- Sharp, R., Saunders, W., Smith, G., et al. 2006, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 6269, *Ground-based and Airborne Instrumentation for Astronomy*, ed. I. S. McLean & M. Iye, 62690G
- Shipp, N., Erkal, D., Drlica-Wagner, A., et al. 2021, *ApJ*, 923, 149
- Siegel, M. H., Dotter, A., Majewski, S. R., et al. 2007, *ApJ*, 667, L57
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Skúladóttir, Á., Hansen, C. J., Salvadori, S., & Choplin, A. 2019, *A&A*, 631, A171
- Skúladóttir, Á., Puls, A. A., Amarsi, A. M., et al. 2023, *The Messenger*, 190, 19
- Skúladóttir, Á. & Salvadori, S. 2020, *A&A*, 634, L2
- Skúladóttir, Á., Vanni, I., Salvadori, S., & Lucchesi, R. 2024, *A&A*, 681, A44
- Smecker-Hane, T. A. & McWilliam, A. 2002, arXiv e-prints, astro
- Sneden, C., Cowan, J. J., & Gallino, R. 2008, *ARA&A*, 46, 241
- Starkenburg, E., Martin, N., Youakim, K., et al. 2017, *MNRAS*, 471, 2587
- Strolger, L.-G., Riess, A. G., Dahlen, T., et al. 2004, *ApJ*, 613, 200
- Suda, T., Katsuta, Y., Yamada, S., et al. 2008, *PASJ*, 60, 1159
- Sullivan, M., Le Borgne, D., Pritchett, C. J., et al. 2006, *ApJ*, 648, 868
- Tinsley, B. M. 1968, *Astrophysical Journal*, vol. 151, p. 547, 151, 547
- Tinsley, B. M. 1979, *ApJ*, 229, 1046
- Tolstoy, E., Hill, V., & Tosi, M. 2009, *ARA&A*, 47, 371
- Travaglio, C., Gallino, R., Arnone, E., et al. 2004, *ApJ*, 601, 864
- Van der Swaelmen, M., Hill, V., Primas, F., & Cole, A. A. 2013, *A&A*, 560, A44
- Vasiliev, E. & Belokurov, V. 2020, *MNRAS*, 497, 4162
- Venn, K. A., Irwin, M., Shetrone, M. D., et al. 2004, *AJ*, 128, 1177
- Venn, K. A., Starkenburg, E., Malo, L., Martin, N., & Laevens, B. P. M. 2017, *MNRAS*, 466, 3741
- Villemoes, P., Arnesen, A., Heijkenskjold, F., & Wannstrom, A. 1993, *Journal of Physics B Atomic Molecular Physics*, 26, 4289
- Vitali, S., Arentsen, A., Starkenburg, E., et al. 2022, *MNRAS*, 517, 6121

- White, S. D. M. & Frenk, C. S. 1991, *ApJ*, 379, 52
White, S. D. M. & Rees, M. J. 1978, *MNRAS*, 183, 341
Wiseman, P., Sullivan, M., Smith, M., et al. 2021, *MNRAS*, 506, 3330
Wosley, S. E. & Weaver, T. A. 1995, *ApJS*, 101, 181
Yang, Y., Caffau, E., Bonifacio, P., et al. 2024, arXiv e-prints,
arXiv:2409.15414

Appendix A: Radial velocities

In Fig. A.1, we present the RV values obtained using the method described in Sect. 3.1 for the three setups. Notably, the different sets of values show good agreement with the exception of two stars where the redder setup H665 yields RV values $\approx 5 \text{ km s}^{-1}$ higher than the other setups. Upon inspecting the spectra of these two stars, we found that the spectra quality for the redder setup was compromised due to sky emission lines in this wavelength region. Consequently, we decided to compute the final RV for these two targets excluding the H665-RV values.




Fig. A.1. Radial velocity values with their uncertainties for the entire FLAMES sample inferred using the three setups. A diagonal line is shown as a guide to better illustrate the agreement between the results of the different setups, which almost all fall on a 1:1 relation.

Appendix B: Metallicities

The metallicity results from the three different setups are illustrated in Fig. B.1. Each metallicity sample was inferred following the methodology outlined in Sect. 3.3. The left panel displays the $[\text{Fe}/\text{H}]$ values selected for averaging as the final measurements for this study. The other two panels show that the metallicities obtained from HR15 are systematically higher compared to those from the other two setups. Therefore, the HR15 data were excluded from the final averaged iron abundances.




Fig. B.1. Comparison between the different sets of spectroscopic iron abundances obtained with the three different GIRAFFE filters.