A PIECEWISE-LINEAR FORMULA FOR RIGID REPRESENTATIONS OF DYNKIN QUIVERS

DENIZ KUS, MARKUS REINEKE

ABSTRACT. We derive a piecewise-linear formula for the rigid representation of a Dynkin quiver of a given dimension vector, and illustrate the formula in several examples.

1. INTRODUCTION

It is a fundamental fact in the representation theory of Dynkin quivers that any dimension vector admits a unique (up to isomorphism) rigid representation, that is, one with vanishing group of self-extensions. This fact already plays a decisive role in proofs of Gabriel's theorem stating that the isomorphism classes of representations of a Dynkin quiver are parametrized by non-negative integer valued functions on the set of positive roots of the corresponding simply-laced root system (we recall basic notation, facts and techniques from the representation theory of quivers in Sections 2.1, 2.2).

It is thus natural to ask how the rigid representation of a given dimension vector is given in this parametrization. Our main result is a closed piecewise-linear formula for this (Theorem 1), which is proved using standard methods of Auslander-Reiten theory, and of the theory of general subrepresentations, in Section 2.3. We illustrate the efficiency of this formula by making it explicit, and comparing it to natural rank conditions, for representations of type A quivers in Section 3.

Acknowledgments: This work originated in a research seminar on flat families of quiver Grassmannians at Ruhr-University Bochum in 2022. The authors would like to thank all participants, in particular M. Bertozzi, M. Boos, V. Genz, W. Gnedin and V. Rappel, for their contributions.

2. Piecewise-linear formula: the main result

2.1. For all techniques and more details on the representation theory of quivers, we refer to [4]. Let Q be a finite quiver with set of vertices Q_0 and set of arrows Q_1 , arrows being denoted by $\alpha : i \to j$ for $i, j \in Q_0$. Let $t, h : Q_1 \to Q_0$ be the maps associating to an arrow α its tail and head, respectively. The associated Euler form is given by

$$\langle \mathbf{d}, \mathbf{e} \rangle = \sum_{i \in Q_0} d_i e_i - \sum_{\alpha: i \to j} d_i e_j, \quad \mathbf{d} = (d_i)_i, \ \mathbf{e} = (e_i)_i \in \mathbb{Z}Q_0.$$

For an algebraically closed field \mathbb{K} , we consider finite-dimensional \mathbb{K} -representations V of Q, consisting of \mathbb{K} -vector spaces V_i for $i \in Q_0$ and \mathbb{K} -linear maps $V_{\alpha} : V_i \to V_j$ for all arrows $\alpha : i \to j$. They naturally form a \mathbb{K} -linear abelian category rep_{\mathbb{K}}Q, which is equivalent to the category of finite dimensional left modules over the path algebra $\mathbb{K}Q$ of Q. For a path β in the path algebra we denote by V_{β} the composition of the

linear maps V_{α} , $\alpha \in Q_1$ along the path β . The tuple $\dim(V) = (\dim_{\mathbb{K}} V_i)_{i \in Q_0} \in \mathbb{N}Q_0$ is called the dimension vector of V. We have

$$\dim_{\mathbb{K}} \operatorname{Hom}_{Q}(V, W) - \dim_{\mathbb{K}} \operatorname{Ext}_{Q}^{1}(V, W) = \langle \operatorname{dim}(V), \operatorname{dim}(W) \rangle$$

for all representations V, W of Q. The set

$$R_{\mathbf{d}}(Q) = \bigoplus_{\alpha: i \to j} \operatorname{Hom}_{\mathbb{K}}(V_i, V_j), \ \mathbf{d} \in \mathbb{N}Q_0$$

of all representations of Q on the vector spaces V_i of dimension d_i is an affine space, on which the algebraic group

$$G_{\mathbf{d}} = \prod_{i \in Q_0} \operatorname{GL}(V_i)$$

acts via base change

$$(g_i)_i \cdot (V_\alpha)_\alpha = (g_j V_\alpha g_i^{-1})_{\alpha:i \to j}$$

so that the orbits \mathcal{O}_V of $G_{\mathbf{d}}$ in $R_{\mathbf{d}}(Q)$ correspond bijectively to the isomorphism classes [V] of \mathbb{K} -representations of Q of dimension vector \mathbf{d} . We have

(2.1)
$$\dim R_{\mathbf{d}}(Q) - \dim \mathcal{O}_{V} = \dim_{\mathbb{K}} \operatorname{Ext}_{Q}^{1}(V, V)$$

$$\dim G_{\mathbf{d}} - \dim R_{\mathbf{d}}(Q) = \langle \mathbf{d}, \mathbf{d} \rangle.$$

2.2. We assume in the rest of the paper that Q is a Dynkin quiver (that is, the underlying unoriented graph is a disjoint union of Dynkin diagrams of type A_n , D_n , E_6 , E_7 , E_8). Then the quadratic form associated to the Euler form $\langle -, - \rangle$ is positive definite. We denote by

$$\Phi^+ = \{ \alpha \in \mathbb{N}Q_0 \, | \, \langle \alpha, \alpha \rangle = 1 \}$$

the set of positive roots of this form, or, equivalently, of the root system for the Dynkin type of Q. By the classical theorem of Gabriel, the isomorphism classes of indecomposable K-representations of Q are in bijection to Φ^+ . More precisely, for every $\alpha \in \Phi^+$ there exists a unique (up to isomorphism) indecomposable representations U_{α} such that $\dim(U_{\alpha}) = \alpha$. By the theorem of Krull-Remak-Schmidt, every finite-dimensional representation V of Q is then isomorphic to a direct sum

(2.2)
$$V \cong \bigoplus_{\alpha \in \Phi^+} U^{\mathbf{m}_V(\alpha)}_{\alpha}$$

for a unique function $\mathbf{m}_V : \Phi^+ \to \mathbb{N}$. In particular, for a given dimension vector \mathbf{d} , there are only finitely many isomorphism classes of \mathbb{K} -representations of Q of dimension vector \mathbf{d} . Equivalently, there are only finitely many orbits of the connected algebraic group $G_{\mathbf{d}}$ in the irreducible algebraic variety $R_{\mathbf{d}}(Q)$, thus there exists a unique open orbit $\mathcal{O}_{\mathbf{d},\mathrm{ri}} \subseteq R_{\mathbf{d}}(Q)$, corresponding to a so-called rigid representation $V_{\mathbf{d},\mathrm{ri}}$ uniquely defined up to isomorphism. It is already uniquely determined by the properties $\dim(V_{\mathbf{d},\mathrm{ri}}) = \mathbf{d}$ and $\operatorname{Ext}^1_Q(V_{\mathbf{d},\mathrm{ri}}, V_{\mathbf{d},\mathrm{ri}}) = 0$ (see equation (2.1)). However, a formula for the decomposition (2.2) of $V_{\mathbf{d},\mathrm{ri}}$ into indecomposables is still not present in the literature. In special cases, the description is straightforward; for example, we have $V_{\alpha,\mathrm{ri}} \simeq U_{\alpha}$ for all positive roots $\alpha \in \Phi^+$.

Our aim is to derive a closed, piecewise-linear, formula for the decomposition of $V_{\mathbf{d},\mathrm{ri}}$ and illustrate the formula in type A leading also to maximizing certain rank tuples. We write $\mathbf{e} \hookrightarrow \mathbf{d}$ (resp. $\mathbf{d} \twoheadrightarrow \mathbf{e}$) if every \mathbb{K} -representation of Q of dimension vector \mathbf{d} admits a subrepresentation (resp. factor representation) of dimension vector \mathbf{e} . In the following, we use the convenient notation $[x]_{+} = \max\{x, 0\}$ for $x \in \mathbb{R}$. The main result is the following.

Theorem 1. Let Q be a Dynkin quiver. For the rigid representation $V_{\mathbf{d},\mathrm{ri}}$ of Q of dimension vector \mathbf{d} , we have

 $\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}(\alpha) = [\min\{\langle \mathbf{e}, \mathbf{d} \rangle, \langle \mathbf{d}, \mathbf{e}' \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha, \ \alpha \twoheadrightarrow \mathbf{e}' \neq 0)\}]_+ \text{ for all } \alpha \in \Phi^+.$

Before proving the theorem, we recall a few concepts from the representation theory of quivers, namely almost split sequences, and generic subrepresentations/extensions.

By Auslander-Reiten theory, for every indecomposable representation U_{α} , there exists a so-called almost split sequence

$$U_{\alpha} \stackrel{f}{\longrightarrow} X \stackrel{g}{\longrightarrow} V$$

inducing the following exact sequence of $\operatorname{mod}_{\mathbb{K}}$ -valued \mathbb{K} -linear functors on $\operatorname{rep}_{\mathbb{K}}Q$:

 $0 \longrightarrow \operatorname{Hom}_Q(V, -) \longrightarrow \operatorname{Hom}_Q(X, -) \longrightarrow \operatorname{Hom}_Q(U_{\alpha}, -) \longrightarrow S_{U_{\alpha}} \longrightarrow 0,$

where $S_{U_{\alpha}}$ is the simple functor whose value at any representation Y is the vector space $\mathbb{K}^{\mathbf{m}_{Y}(\alpha)}$ (the above sequence thus providing a projective resolution of this functor). In particular, for every representation Y, we have

$$\mathbf{m}_{Y}(\alpha) = \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{\alpha}, Y) - \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(X, Y) + \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(V, Y).$$

Moreover, the representation X fulfills $\operatorname{Ext}_Q^1(X, X) = 0$ and is therefore rigid of dimension vector $\alpha + \operatorname{dim}(V)$. If U_{α} is an injective indecomposable, then $X = U_{\alpha}/\operatorname{soc}(U_{\alpha})$ and V = 0. Otherwise, the almost split sequence is short exact, the representation V is again indecomposable, and from the above exact sequence of functors, we see immediately that every non-split mono starting in U_{α} factors over f; dually, every non-split epi to V factors over g.

The association of V to U_{α} is functorial, in fact given by the inverse Auslander-Reiten translation functor which, in the Dynkin case, is given as $\tau^{-1} = \text{Ext}_Q^1((\mathbb{K}Q)^*, -)$; the Auslander-Reiten formulas then imply $\text{Hom}_Q(V, U_{\alpha}) = 0$. The functor τ^{-1} induces a bijection from isomorphism classes of non-injective indecomposables to isomorphism classes of non-projective indecomposables, and it induces a linear map, again denoted by $\tau^{-1} \in \text{GL}(\mathbb{Z}Q_0)$, on the level of dimension vectors. This map satisfies

(2.3)
$$\langle \tau^{-1} \mathbf{d}, \mathbf{e} \rangle = -\langle \mathbf{e}, \mathbf{d} \rangle.$$

2.3. Given two K-representations V_1 and V_2 of Q, there exists a unique (up to isomorphism) representation Y where the function $\dim_{\mathbb{K}} \operatorname{End}(Y)$ obtains its minimum among all extensions of V_1 by V_2 ; we call it the generic extension of V_1 by V_2 and denote it by $V_1 * V_2$. We say that a representation W degenerates to a representation W' if the orbit $\mathcal{O}_{W'}$ is contained in the Zariski-closure of \mathcal{O}_W . In particular, $V_{d,ri}$ degenerates to W for any representation W of dimension vector \mathbf{d} .

We then have the following result from [2, Proposition 2.4]: if V_1 degenerates to V'_1 and V_2 degenerates to V'_2 , and Y is an arbitrary extension of V'_1 by V'_2 , then $V_1 * V_2$ degenerates to Y. From this we conclude the following.

(2.4) If $V_{\mathbf{d},\mathrm{ri}}$ admits a factor of dimension vector \mathbf{e} , then $\mathbf{d} \rightarrow \mathbf{e}$.

Indeed, if $V_{\mathbf{d},\mathrm{ri}}/W' \cong W$ for some representation W of dimension vector \mathbf{e} , then $V_{\mathbf{e},\mathrm{ri}}$ degenerates to W. Thus, by the above discussion, $V_{\mathbf{e},\mathrm{ri}} * W'$ degenerates to $V_{\mathbf{d},\mathrm{ri}}$. But

this implies $V_{\mathbf{d},\mathrm{ri}} \simeq V_{\mathbf{e},\mathrm{ri}} * W'$ and hence $V_{\mathbf{e},\mathrm{ri}}$ is a factor of $V_{\mathbf{d},\mathrm{ri}}$. Similarly, we obtain the statement for subrepresentations.

(2.5) If $V_{\mathbf{d},\mathrm{ri}}$ admits a subrepresentation of dimension vector \mathbf{e} , then $\mathbf{e} \hookrightarrow \mathbf{d}$.

We will use these two statements in the rest of this paper. We can now prove the theorem.

Proof of Theorem 1: The function $(V, W) \mapsto \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(V, W)$ is upper semicontinuous on the product $R_{\mathbf{e}}(Q) \times R_{\mathbf{d}}(Q)$ and its minimal value, denoted by $\operatorname{hom}(\mathbf{e}, \mathbf{d})$, is obtained on the product of open orbits, that is, $\operatorname{hom}(\mathbf{e}, \mathbf{d}) = \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(V_{\mathbf{e}, \mathrm{ri}}, V_{\mathbf{d}, \mathrm{ri}})$. In particular, for every $\alpha \in \Phi^+$, we have

 $\dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{\alpha}, V_{\mathbf{d}, \operatorname{ri}}) = \operatorname{hom}(\alpha, \mathbf{d}).$

Moreover, using the Euler form and a formula of Schofield [5, Theorem 5.4], we get

$$\hom(\mathbf{e}, \mathbf{d}) = \max\{\langle \mathbf{f}, \mathbf{d} \rangle : \mathbf{e} \twoheadrightarrow \mathbf{f}\}.$$

Let $U_{\alpha} \xrightarrow{f} X \xrightarrow{g} V$ the almost split sequence starting in U_{α} for some $\alpha \in \Phi^+$ and let γ and β be the dimension vectors of X and V, respectively. From the discussion in Subsection 2.2 we have

$$\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}(\alpha) = \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{\alpha}, V_{\mathbf{d},\mathrm{ri}}) - \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(X, V_{\mathbf{d},\mathrm{ri}}) + \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(V, V_{\mathbf{d},\mathrm{ri}})$$
$$= \operatorname{hom}(\alpha, \mathbf{d}) - \operatorname{hom}(\gamma, \mathbf{d}) + \operatorname{hom}(\beta, \mathbf{d})$$
$$= \max\{\langle \mathbf{e}, \mathbf{d} \rangle : \alpha \twoheadrightarrow \mathbf{e}\} - \max\{\langle \mathbf{e}, \mathbf{d} \rangle : \gamma \twoheadrightarrow \mathbf{e}\} + \max\{\langle \mathbf{e}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{e}\}.$$

For $\mathbf{e} \in \mathbb{N}Q_0$, we denote by $Q_{\mathbf{e}}$ the set of all dimension vectors \mathbf{f} such that $\mathbf{e} \to \mathbf{f}$. We claim that

(2.6)
$$Q_{\gamma} = (Q_{\alpha} + Q_{\beta}) \setminus \{\alpha\}.$$

In case of U_{α} beeing injective, we have $\beta = 0$ and $\gamma = \dim(U_{\alpha}/\operatorname{soc}(U_{\alpha}))$. So identity (2.6) follows from the fact that $\operatorname{soc}(U_{\alpha})$ is simple, and thus the proper factors of U_{α} are precisely the factors of $U_{\alpha}/\operatorname{soc}(U_{\alpha})$. So we assume that U_{α} is non-injective, and prove identity (2.6) in three steps.

Step 1: We first prove $Q_{\gamma} \subseteq (Q_{\alpha} + Q_{\beta}) \setminus \{\alpha\}$. Suppose that $\mathbf{f} \in Q_{\gamma}$ and let Y = X/C be a factor of X of dimension vector \mathbf{f} . Then $U_{\alpha}/(U_{\alpha} \cap C)$ is a factor of U_{α} , say of dimension vector \mathbf{f}_1 , and

$$X/(U_{\alpha}+C) \simeq (X/U_{\alpha})/((U_{\alpha}+C)/U_{\alpha}) \simeq V/((U_{\alpha}+C)/U_{\alpha})$$

is a factor of V, say of dimension vector \mathbf{f}_2 , so that $\mathbf{f} = \mathbf{f}_1 + \mathbf{f}_2$. Hence the inclusion is obtained from (2.4) if we can exclude the case $\mathbf{f} = \alpha$. However, if $\mathbf{f} = \alpha$ we have that U_{α} is already a factor of X, which is impossible by the above exact sequence of functors characterizing almost split sequences, namely $\operatorname{Hom}_Q(X, U_{\alpha}) = \operatorname{Hom}_Q(V, U_{\alpha}) = 0$ since $\operatorname{End}_Q(U_{\alpha}) \simeq \mathbb{K}$.

Step 2: Next, we prove $(Q_{\alpha} \setminus \{\alpha\}) + Q_{\beta} \subseteq Q_{\gamma}$. So let $p: U_{\alpha} \to Y$ and $q: V \to Z$ be surjective maps, where p is not an isomorphism. Hence, p factors through $f: U_{\alpha} \to X$ to a map $\hat{p}: X \to Y$ which is still surjective. We can also compose q with the map $g: X \to V$. It is then easily verified that the map $(\hat{p}, q \circ g): X \to Y \oplus Z$ is again surjective.

Step 3: Finally, we prove $\alpha + (Q_{\beta} \setminus \{0\}) \subseteq Q_{\gamma}$. So let $Z \neq 0$ and $V \to Z$ be surjective with kernel $K \to V$ being non-split epi, thus factoring to an embedding $K \to X$, and $X \to X/K$ is the desired factor of dimension vector $\alpha + \dim(Z)$.

So identity (2.6) is proven, and we rewrite it in the following ways:

(2.7)
$$Q_{\alpha} + Q_{\beta} = Q_{\gamma} \cup \{\alpha\}$$

(2.8)
$$Q_{\gamma} = ((Q_{\alpha} \setminus \{\alpha\}) + Q_{\beta}) \cup (\alpha + (Q_{\beta} \setminus \{0\}),$$

and we will make use of the following simple identity:

(2.9)
$$[\min\{x, y, x+y\}]_{+} = [\min\{x, y\}]_{+}$$

Given a subset $B \subseteq \mathbb{N}Q_0$, we abbreviate

$$\max\langle B, \mathbf{d} \rangle = \max\{\langle \mathbf{f}, \mathbf{d} \rangle : f \in B\}$$

and calculate:

$$\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}(\alpha) = \max \langle Q_{\alpha} + Q_{\beta}, \mathbf{d} \rangle - \max \langle Q_{\gamma}, \mathbf{d} \rangle$$

$$\stackrel{(2.7)}{=} \max \{\max \langle Q_{\gamma}, \mathbf{d} \rangle, \langle \alpha, \mathbf{d} \rangle\} - \max \langle Q_{\gamma}, \mathbf{d} \rangle = [\min \{ \langle \alpha - \mathbf{e}, \mathbf{d} \rangle : \gamma \twoheadrightarrow \mathbf{e} \}]_{+}$$

$$\stackrel{(2.8)}{=} [\min \{ \min \{ \langle \alpha - \mathbf{e} - \mathbf{f}, \mathbf{d} \rangle : \mathbf{e} \in Q_{\alpha} \setminus \{\alpha\}, \mathbf{f} \in Q_{\beta} \},$$

$$\min \{ \langle -\mathbf{f}, \mathbf{d} \rangle : \mathbf{f} \in Q_{\beta} \setminus \{0\} \} \}]_{+}$$

$$= [\min \{ \min \{ \langle \mathbf{e} - \mathbf{f}, \mathbf{d} \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha, \beta \twoheadrightarrow \mathbf{f} \}, \min \{ \langle -\mathbf{f}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{f} \neq 0 \} \}]_{+}$$

$$= [\min \{ \min \{ \langle \mathbf{e} - \mathbf{f}, \mathbf{d} \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha, \beta \twoheadrightarrow \mathbf{f} \neq 0 \}, \min \{ \langle \mathbf{e}, \mathbf{d} \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha \},$$

$$\min \{ \langle -\mathbf{f}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{f} \neq 0 \} \}]_{+}$$

$$\stackrel{(2.9)}{=} [\min \{ \min \{ \langle \mathbf{e}, \mathbf{d} \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha \}, \min \{ -\mathbf{f}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{f} \neq 0 \} \}]_{+}$$

$$= \left[\min \left\{ \min \left\{ \langle \mathbf{e}, \mathbf{d} \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha \right\}, \min \left\{ -\langle \mathbf{f}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{f} \neq 0 \right\} \right]_{+}.$$

Denoting by Q_{α}^{inj} the set of dimension vectors of injective factor representations of U_{α} , the properties of the Auslander-Reiten translation imply that

$$Q_{\beta} \setminus \{0\} = \tau^{-1}(Q_{\alpha} \setminus Q_{\alpha}^{\mathrm{inj}}).$$

Together with the formula (2.3), this yields

$$\min\{-\langle \mathbf{f}, \mathbf{d} \rangle : \beta \twoheadrightarrow \mathbf{f} \neq 0\} = \min\{\langle \mathbf{d}, \mathbf{e}' \rangle : \mathbf{e}' \in Q_{\alpha} \setminus Q_{\alpha}^{\operatorname{inj}}\},\$$

and thus the calculations above give

$$\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}(\alpha) = \left[\min\{\langle \mathbf{e}, \mathbf{d} \rangle, \ \langle \mathbf{d}, \mathbf{e}' \rangle : \ 0 \neq \mathbf{e} \hookrightarrow \alpha, \ \mathbf{e}' \in Q_{\alpha} \setminus Q_{\alpha}^{\mathrm{inj}}\}\right]_{+}.$$

We claim that this already equals

$$[\min\{\langle \mathbf{e}, \mathbf{d} \rangle, \ \langle \mathbf{d}, \mathbf{e}' \rangle : 0 \neq \mathbf{e} \hookrightarrow \alpha, \ \alpha \twoheadrightarrow \mathbf{e}' \neq 0\}]_+,$$

which then proves the theorem.

Indeed, suppose that $\alpha \to \mathbf{e}' \neq 0$ for \mathbf{e}' the dimension vector of an injective representation I, thus in particular $\operatorname{Hom}_Q(U_\alpha, I) \neq 0$. The inverse Nakayama functor $\nu^{-1} = \operatorname{Hom}_Q((\mathbb{K}Q)^*, -)$ then yields a projective representation $\nu^{-1}I$ such that

$$\operatorname{Hom}_Q(U_\alpha, I) \simeq \operatorname{Hom}_Q(\nu^{-1}I, U_\alpha)^*,$$

thus we can choose a non-zero map $0 \neq \varphi : \nu^{-1}I \to U_{\alpha}$. We denote its (projective) kernel by P, its image by U', and by \mathbf{f} the dimension vector of U' which satisfies $0 \neq \mathbf{f} \hookrightarrow \alpha$ by (2.5). Then

$$0 \to P \to \nu^{-1}I \to U' \to 0$$

is a projective resolution, and we find for any dimension vector \mathbf{d} that

$$\begin{split} \langle \mathbf{f}, \mathbf{d} \rangle &= \langle \mathbf{dim}(\nu^{-1}I), \mathbf{d} \rangle - \langle \mathbf{dim}(P), \mathbf{d} \rangle = \langle \mathbf{d}, \mathbf{dim}(I) \rangle - \langle \mathbf{dim}(P), \mathbf{d} \rangle = \\ &= \langle \mathbf{d}, \mathbf{e}' \rangle - \langle \mathbf{dim}(P), \mathbf{d} \rangle \leq \langle \mathbf{d}, \mathbf{e}' \rangle, \end{split}$$

which shows that the above minimum does not change when adding the terms $\langle \mathbf{d}, \mathbf{e}' \rangle$ for $0 \neq \mathbf{e}' \in Q_{\alpha}^{\text{inj}}$.

2.4. To make our main result more efficient, we have to determine, for a given positive root $\alpha \in \Phi^+$, the dimension vectors $\mathbf{e} \neq 0$ satisfying $\mathbf{e} \hookrightarrow \alpha$ and $\alpha \twoheadrightarrow \mathbf{e}$ respectively.

Lemma 2. Let $\alpha \in \Phi^+$ and $\mathbf{e} \neq 0$ be a non-zero element of $\mathbb{N}Q_0$. We have $\mathbf{e} \hookrightarrow \alpha$ if and only if

$$[\langle \beta, \alpha \rangle]_+ \ge \langle \beta, \mathbf{e} \rangle, \quad \forall \beta \in \Phi^+$$

Dually, we have $\alpha \twoheadrightarrow \mathbf{e}$ if and only if

$$[\langle \alpha, \beta \rangle]_+ \ge \langle \mathbf{e}, \beta \rangle \quad \forall \beta \in \Phi^+.$$

Proof. By [1, Theorem 4.2], we have $\mathbf{e} \hookrightarrow \alpha$ if and only if $\dim_{\mathbb{K}} \operatorname{Hom}_Q(U_{\beta}, U_{\alpha}) \geq \langle \beta, \mathbf{e} \rangle$ for all $\beta \in \Phi^+$. Now Dynkin quivers are representation-directed, that is, there exists a partial ordering \preceq on Φ^+ such that $\operatorname{Hom}_Q(U_{\beta}, U_{\alpha}) \neq 0$ already implies $\beta \preceq \alpha$, and $\operatorname{Ext}^1_Q(U_{\beta}, U_{\alpha}) \neq 0$ implies $\alpha \prec \beta$. This immediately gives $\dim_{\mathbb{K}} \operatorname{Hom}_Q(U_{\beta}, U_{\alpha}) = [\langle \beta, \alpha \rangle]_+$. The dual statement is similar and the lemma is proved.

3. Applications to type A: rank tuples

Let Q be an arbitrary type A_n quiver. The positive roots are parametrized by intervals in this quiver,

$$\Phi^{+} = \{ \alpha_{ij} : 1 \le i \le j \le n \},\$$

where $(\alpha_{ij})_k = 1$ precisely if $i \leq k \leq j$. We denote by $U_{i,j}$ the corresponding indecomposable representations. For a subset $I \subseteq \{1, \ldots, n\}$, we denote by \mathbf{e}_I the dimension vector such that $(\mathbf{e}_I)_i = 1$ when $i \in I$, and zero otherwise. Then $0 \neq \mathbf{e}_I \hookrightarrow \alpha_{ij}$ if and only if $\emptyset \neq I \subseteq \{i, \ldots, j\}$ is closed under successors, that is, if $k \in I$ and there is an arrow $k \to l$ in Q, then $l \in I$, for $i \leq k, l \leq j$. Dually, $\alpha_{ij} \twoheadrightarrow \mathbf{e}_I \neq 0$ if and only if $\emptyset \neq I \subseteq \{i, \ldots, j\}$ is closed under predecessors.

3.1. We consider the equioriented type A_n quiver

$$1 \to 2 \to \cdots \to n.$$

From the discussion above, the only non-zero dimension vectors of subrepresentations (resp. quotient representations) of the corresponding indecomposable $U_{i,j}$ are the α_{kj} (resp. α_{ik}) for $i \leq k \leq j$. We thus find with our main result that

$$\mathbf{m}_{V_{\mathbf{d},ri}}(\alpha_{ij}) = \left[\min\left\{d_k - d_{i-1}, d_k - d_{j+1} : i \le k \le j\right\}\right]_+, \quad \forall 1 \le i \le j \le n$$

where we understand $d_0 = 0 = d_{n+1}$. The representation $V = V_{\mathbf{d},\mathrm{ri}}$, given as

$$V_1 \stackrel{f_1}{\to} V_2 \stackrel{f_2}{\to} \cdots \stackrel{f_{n-1}}{\to} V_n$$

is determined up to isomorphism by the ranks

$$r_{ij}(V) = \operatorname{rank}(f_{j-1} \circ \cdots \circ f_i : V_i \to V_j)$$

for $1 \leq i \leq j \leq n$ (where we view the empty composition of maps as the identity map, that is, $r_{ii}(V) = \dim_{\mathbb{K}} V_i$). We have $r_{ij}(V) = \min\{d_i, \ldots, d_j\}$. To see this, let i < j (the case i = j is true by convention) and observe

$$r_{ij}(V) = \dim_{\mathbb{K}} V_i - \dim_{\mathbb{K}} \operatorname{Ker}(f_{j-1} \circ \ldots \circ f_i) = d_i - \dim_{\mathbb{K}} \operatorname{Hom}_Q(U_{i,j-1}, V)$$
$$= d_i - \operatorname{hom}(\alpha_{i,j-1}, \mathbf{d}) = d_i - \max\{d_i - d_k : i \le k \le j\} = \min\{d_i, \ldots, d_j\}.$$

The above formula thus constructs a representation V where all the ranks $r_{ij}(V)$ reach their theoretical maximum min $\{d_i, \ldots, d_j\}$.

3.2. Now we consider the case when Q has a unique sink, that is,

 $1 \to 2 \to \dots \to s \leftarrow \dots \leftarrow n - 1 \leftarrow n.$

We thus find with our main result that for all $1 \le i \le j \le n$ we have (ignoring the redundant elements not contributing to the minimum)

$$\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}(\alpha_{ij}) = \begin{cases} \left[\min\left\{d_k + d_{k'} - d_s, d_s - d_{i-1} - d_{j+1}, \\ d_{k'} - d_{j+1}, d_k - d_{i-1} : i \le k \le s \le k' \le j\right\}\right]_+, & i < s \text{ and } j > s \\ \left[\min\left\{d_k - d_{j+1}, d_k - d_{i-1} : i \le k \le j\right\}\right]_+, & j < s \text{ or } i > s \\ \left[\min\left\{d_k - d_{i-1} : i \le k \le j\right\}\right]_+, & j = s \\ \left[\min\left\{d_k - d_{j+1} : i \le k \le j\right\}\right]_+, & i = s \end{cases}$$

where we understand again $d_0 = d_{n+1} = 0$. The representation $V = V_{\mathbf{d},\mathrm{ri}}$, given as

$$V_1 \xrightarrow{f_1} V_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{s-1}} V_s \xleftarrow{f_s} \cdots \xleftarrow{f_{n-2}} V_{n-1} \xleftarrow{f_{n-1}} V_n$$

is again determined up to isomorphism by the ranks $r_{ij}(V)$, $1 \le i \le j \le n$, where $r_{ii}(V) = \dim_{\mathbb{K}}(V_i)$ as above and otherwise

$$r_{ij}(V) = \begin{cases} \operatorname{rank}(f_{j-1} \circ \cdots \circ f_i : V_i \to V_j), & \text{if } j \leq s \\ \operatorname{rank}(f_i \circ \cdots \circ f_{j-1} : V_j \to V_i), & \text{if } i \geq s \\ \operatorname{rank}((f_{s-1} \circ \cdots \circ f_i) \oplus (f_s \circ \cdots \circ f_{j-1}) : V_i \oplus V_j \to V_r), & \text{if } j > s, i < s \end{cases}$$

We have

 $r_{ij}(V) = \min\{d_i, \dots, d_j\}, \text{ if } j \le s \text{ or } i \ge s$

as in the equioriented case, and for i < s < j we get

$$r_{ij}(V) = d_i + d_j - \dim_{\mathbb{K}} \operatorname{Hom}_Q(U_{i,j}, V) = d_i + d_j - \hom(\alpha_{i,j}, \mathbf{d})$$

= $d_i + d_j - \max\{d_i + d_j - d_k - d_{k'}, d_i + d_j - d_s : i \le k \le s, s \le k' \le j\}$
= $\min\{d_k + d_{k'}, d_s : i \le k \le s, s \le k' \le j\}$
= $\min\{\min\{d_i, \dots, d_s\} + \min\{d_s, \dots, d_j\}, d_s\}.$

So in both cases, the rigid representation is the unique representation maximizing certain ranks. This will be generalized to arbitrary orientations in the next subsection and can be interpreted as a coherent linear algebra description of the rigid representation. **3.3.** Consider an arbitrary orientation of the type A quiver Q and let Q(i, j) for $i \leq j$ be the full subquiver of Q with vertices $\{i, \ldots, j\}$. Moreover, we introduce the following notation.

- Let $Q^{si}(i, j)$ (resp. $Q^{so}(i, j)$) be the set of sinks in Q(i 1, j + 1) different from the nodes *i* and *j* (resp. the set of sources in Q(i, j)).
- For $k \in Q^{si}(i, j)$ we denote by w_k^{ij} (resp. u_k^{ij}) the unique longest path in Q(i, k) (resp. Q(k, j)) ending in k.

Then we have a projective resolution of $U_{i,j}$:

$$0 \longrightarrow P_1 := \bigoplus_{k \in Q^{\mathrm{si}}(i,j)} P(k) \xrightarrow{\varphi} P_0 := \bigoplus_{\ell \in Q^{\mathrm{so}}(i,j)} P(\ell) \longrightarrow U_{i,j} \longrightarrow 0$$

where P(k) denotes as usual the indecomposable projective module corresponding to the node k. The map φ restricted to P(k) is given by $\varphi(e_k) = w_k^{ij} - u_k^{ij}$ (where e_k is the constant path at node k). Now for any $V \in R_{\mathbf{d}}(Q)$ the functor $\operatorname{Hom}_Q(-, V)$ yields:

$$0 \to \operatorname{Hom}_Q(U_{i,j}, V) \to \operatorname{Hom}_Q(P_0, V) \xrightarrow{\Phi_{i,j}^V} \operatorname{Hom}_Q(P_1, V)$$

and hence

(3.1)
$$\dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{i,j}, V) = \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(P_{0}, V) - \operatorname{rank}(\Phi_{i,j}^{V}).$$

Recall the elements w_k^{ij} and u_k^{ij} from above and set $r_{ij}(V) := \operatorname{rank}(A_{ij}(V))$ for all $1 \le i \le j \le n$ where

(3.2)
$$A_{ij}(V) := \begin{bmatrix} V_{w_{k_1}^{ij}} & V_{u_{k_1}^{ij}} & 0 & \cdots & 0 & 0 \\ 0 & V_{w_{k_2}^{ij}} & V_{u_{k_2}^{ij}} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & V_{w_{k_r}^{ij}} & V_{u_{k_r}^{ij}} \end{bmatrix}$$

and $Q^{\rm si}(i,j) = \{k_1 < \dots < k_r\}.$

Proposition 3. Let $V \in R_{\mathbf{d}}(Q)$. Then $V \cong V_{\mathbf{d},\mathrm{ri}}$ if and only if

(3.3)
$$r_{ij}(V) = \sum_{\ell \in Q^{\text{so}}(i,j)} d_{\ell} - \hom(\alpha_{i,j}, \mathbf{d}), \text{ for all } 1 \le i \le j \le n.$$

Proof. We claim that $r_{ij}(V) = \operatorname{rank}(\Phi_{i,j}^V)$ for all $1 \le i \le j \le n$. Before we prove the claim, let us see how this proves the proposition. From (3.1) we have

$$\operatorname{rank}(\Phi_{i,j}^{V_{\mathbf{d},\mathrm{ri}}}) = \sum_{\ell \in Q^{\mathrm{so}}(i,j)} d_{\ell} - \hom(\alpha_{i,j}, \mathbf{d}).$$

Conversely, if V satisfies (3.3), we get again with (3.1) that $\operatorname{rank}(\Phi_{i,j}^V) = \operatorname{rank}(\Phi_{i,j}^{V_{\mathbf{d},\mathrm{ri}}})$ for $1 \leq i \leq j \leq n$, thus

$$\dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{i,j}, V) = \dim_{\mathbb{K}} \operatorname{Hom}_{Q}(U_{i,j}, V_{\mathbf{d}, \mathrm{ri}}), 1 \le i \le j \le n.$$

Hence $V \cong V_{\mathbf{d},\mathrm{ri}}$ since representations are determined up to isomorphism by the dimensions of homomorphism spaces from indecomposables. So it remains to show $r_{ij}(V) = \mathrm{rank}(\Phi_{i,j}^V)$. The rank of $\Phi_{i,j}^V$ is obviously given by the dimension of the space $\{g \circ \varphi : g \in \mathrm{Hom}_Q(P_0, V)\}$. Note that any $g \in \mathrm{Hom}_Q(P(k), V)$ is determined

by an element $V_k^g \in V_k$ via $g(e_k) = V_k^g$. Let $Q^{so}(i, j) = \{\ell_1 < \cdots < \ell_p\}$ and define a map

$$\{g \circ \varphi : g \in \operatorname{Hom}_Q(P_0, V)\} \to \operatorname{Im}\left(\sum_{m=1}^r V_{w_{k_m}^{ij}} \oplus -V_{u_{k_m}^{ij}} : V_{\ell_1} \oplus \cdots \oplus V_{\ell_p} \to V_{k_1} \oplus \cdots \oplus V_{k_r}\right)$$

as follows. Recall that

$$V_{w_{k_m}^{ij}} \oplus -V_{u_{k_m}^{ij}} : V_{h(w_{k_m}^{ij})} \oplus V_{h(u_{k_m}^{ij})} \to V_{k_m}$$

and define the assignment

$$g \circ \varphi \mapsto \sum_{m=1}^{r} V_{w_{k_m}^{ij}}(V_{h(w_{k_m}^{ij})}^g) - V_{u_{k_m}^{ij}}(V_{h(u_{k_m}^{ij})}^g) = \sum_{m=1}^{r} g \circ \varphi(e_{k_m}).$$

Recall that the starting vertices of w_k^{ij} and u_k^{ij} , respectively, correspond to sources, that is, the projective module satisfies $P(h(w_k^{ij})) \subseteq P_0$. This map is in fact a vector space isomorphism. To see this let $\beta \in P(k) \subseteq P_1$ be a path starting at k. Then

$$(g \circ \varphi)(\beta) = V_{\beta}(g \circ \varphi(e_k)) = V_{\beta}(g(w_k^{ij}) - g(u_k^{ij})) = V_{\beta}(V_{w_k^{ij}}(V_{h(w_k^{ij})}^g) - V_{u_k^{ij}}(V_{h(u_k^{ij})}^g))$$

and the injectivity is established. The surjectivity is clear, since given an element

$$\sum_{m=1}' V_{w_{k_m}^{ij}}(z_{h(w_{k_m}^{ij})}) - V_{u_{k_m}^{ij}}(z_{h(u_{k_m}^{ij})})$$

in the image we can simply define $g \in \operatorname{Hom}_Q(P_0, V)$ by $g(e_{h(w_k^{ij})}) = z_{h(w_k^{ij})}$ and $g(e_{h(u_k^{ij})}) = z_{h(u_k^{ij})}$ respectively. The map $\sum_{m=1}^r V_{w_{k_m}^{ij}} \oplus -V_{u_{k_m}^{ij}}$ is represented by the matrix $A_{i,j}(V)$ up to some minus signs, which are negligible for the rank. \Box

So the rigid representation $V_{\mathbf{d},\mathrm{ri}}$ is the unique representation up to isomorphism in $R_{\mathbf{d}}(Q)$ maximizing the rank tuple $(r_{ij}(V))_{1 \leq i \leq j \leq n}$ coordinatewise.

We conclude this section with a few remarks and further questions:

- The piecewise-linear nature of our formula is natural a priori, from the fact that the functions $\mathbf{m}_{V_{\mathbf{d},\mathrm{ri}}}$ have to be supported on one of finitely many maximal subset of roots with vanishing Ext_Q^1 -groups between the corresponding indecomposable representations, which feature on the Catalan combinatorial aspects of quivers [3].
- The minimization in Theorem 1 includes many redundancies, and it would be interesting to find an optimal form of the formula.
- Due to the Auslander-Reiten theoretic nature of the proof of Theorem 1, generalizations to the case of preprojective or preinjective dimension vectors for arbitrary acyclic quivers can be expected.

References

- Kathrin Möllenhoff and Markus Reineke. Embeddings of representations. Algebr. Represent. Theory, 18(4):977–987, 2015.
- [2] Markus Reineke. Generic extensions and multiplicative bases of quantum groups at q = 0. Represent. Theory, 5:147–163, 2001.
- [3] Claus Michael Ringel. The Catalan combinatorics of the hereditary Artin algebras. In Recent developments in representation theory, volume 673 of Contemp. Math., pages 51–177. Amer. Math. Soc., Providence, RI, 2016.

DENIZ KUS, MARKUS REINEKE

- [4] Ralf Schiffler. *Quiver representations*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer, Cham, 2014.
- [5] Aidan Schofield. General representations of quivers. Proc. London Math. Soc. (3), 65(1):46-64, 1992.

Ruhr-University Bochum, Faculty of Mathematics, Universitätsstrasse 150, 44780 Bochum (Germany)

Email address: deniz.kus@rub.de, markus.reineke@rub.de

10